



PHILIPPINE COVID-19 LIVING CLINICAL PRACTICE GUIDELINES

As of January 2022

Disclaimer

As a living guideline, the recommendations will be updated, and new recommendations will be added as the evidence evolves. The living recommendations are based on the best evidence available in scientific literature at the time of its formulation. However, this living CPG is not a comprehensive guide to all practice questions and management options on COVID-19. This is not meant to restrict the practitioner in using sound clinical judgement and sharing the decision with the patient, and from considering other management options according to the patient's particular needs and preferences. This CPG can serve to inform policy, but it is not meant to serve as a basis for approving or denying financial coverage or insurance claims merely because of nonconformance with recommendations. Neither are the recommendations supposed to be considered as legal rules for dictating certain modes of action to the exclusion of others.

Acknowledgements

The Philippine COVID-19 Living CPG funded by the Department of Health's (DOH) AHEAD Program through the Philippine Council for Health Research and Development (PCHRD) and the Disease Prevention and Control Bureau (DPCB).

This project was implemented under the Institute of Clinical Epidemiology, National Institutes of Health (NIH), University of the Philippines Manila (UPM). It was completed with the valuable contribution of 190 people representing the different stakeholders

The Philippine COVID-19 Living CPG team dedicates this work to the patients braving their journey with this disease; to all Filipinos who are equally affected physically, emotionally, socially, economically, among others, and to all healthcare professionals contributing to this fight against COVID-19 through patient care and research.

The content of this CPG is the intellectual property of the Department of Health (DOH). We request for proper use of citations when any part of this document is used for presentation to the public.

Contact Us

Send us an email at covidcpg.ph@gmail.com for any questions or clarifications on the outputs and process of this Living CPG. You may also suggest a clinical question for the consideration of the Living Clinical Practice Guidelines COVID-19 Taskforce.

Participating Professional Societies and Institutions



ASIA PACIFIC CENTER FOR
EVIDENCE BASED HEALTHCARE

Table of Contents

Disclaimer	i
Acknowledgements.....	ii
Participating Professional Societies and Institutions.....	iii
Table of Contents	iv
List of Tables	vi
List of Figures	vi
Executive Summary	i
INTRODUCTION.....	1
Objectives	1
Target Population	1
Intended Users	2
CPG DEVELOPMENT METHODOLOGY	3
Overview of Philippine COVID-19 Living CPG Development Process	3
Guideline Preparation.....	5
Evidence Synthesis	8
Evidence to Decision: Formulating Recommendations	10
Guideline Dissemination.....	14
Guideline Monitoring and Evaluation.....	16
External Review	16
Updating of Guidelines	17
Editorial Independence.....	17
Evidence and Recommendations for the Screening and Diagnosis of COVID-19	19
Evidence and Recommendations for the Treatment of COVID-19	47
Evidence and Recommendations for the Critical Care and Respiratory Management of COVID-19	72
Evidence and Recommendations for the Non-Pharmacologic Interventions for Prevention and Control of COVID-19.....	89
Evidence and Recommendations for the Vaccines and Prophylactic Interventions for COVID-19	103
Evidence and Recommendations for the Adjunct Interventions for Prevention and Control of COVID-19	131
DISCUSSION.....	222
Outputs of the Philippine COVID-19 Living CPG Project	222
Applicability Issues.....	222

Implementation Tools	223
Process Evaluation	223
Research Implications	225
Conclusions and Recommendations	225
REFERENCES.....	226
APPENDICES	227
Appendix A. Members of the Philippine COVID-19 Living CPG Task Force ...	227
Appendix B. Decisions of the Oversight Committee –.....	239
Review of Conflict of Interest.....	239
Appendix C. General Search Strategy for COVID-19.....	244
Appendix D. Breakdown of Consensus Meetings for Phase 2 of the Philippine COVID-19 Living Clinical Practice Guidelines	245
Appendix E. AGREE Reporting Checklist	246

List of Tables

Table 1. Topics covered in the Philippine COVID-19 Living CPG.....	6
Table 2. Factors influencing certainty of evidence.....	9
Table 3. Definitions and Implications of each GRADE Certainty of Evidence.	10
Table 4. Implications of the Strength of Recommendation to Patients, Clinicians, and Policymakers	11
Table 5. Outcome ratings for each CPG panel.....	12
Table 6. Top locations of website visitors of the Philippine COVID-19 Living CPG..	224
Table 7. Top evidence summaries downloaded from the Philippine COVID-19 Living CPG.....	224

List of Figures

Figure 1. Philippine COVID-19 Living CPG Development Process.	4
Figure 2. Initial PSMID Webpage for the Philippine COVID-19 LCPG in March 2021.	14
Figure 3. Latest PSMID Webpage for the Philippine COVID-19 LCPG.	15
Figure 4. Living Recommendations Document for the Philippine COVID-19 Living CPG.	15
Figure 5. Contact details in the Webpage for the Philippine COVID-19 Living CPG.	16
Figure 7. Geographical location of website visitors of the Philippine COVID-19 Living CPG.	224

Executive Summary

Coronavirus disease 2019 (COVID-19) has grown into a pandemic and global crisis affecting multiple sectors of society. Over 180 million confirmed COVID-19 cases have been reported globally, with 2.8 million of these cases from the Philippines as of December 2021. Despite the national strategies implemented to curtail the health and economic impact of COVID-19 in the country, epidemiologic projections have yet to point to a foreseeable end to the pandemic, especially with the recent rise of variants with increased transmissibility. Thus, the Philippine COVID-19 Living CPG aimed to provide up-to-date, evidence-based recommendations on the management of COVID-19 among adults with or at risk for COVID-19. Thematic areas included in this CPG were screening and diagnosis, treatment, critical care and respiratory management, non-pharmacologic interventions, vaccines and prophylactic interventions, and adjunct interventions for COVID-19. This document also serves as an update of the PSMID-PCP-PCCP-PCHTM-PRA July 2020 Interim Guidance.

Following the standard CPG development process outlined in the DOH Manual for CPG Development and the GRADE methodology, 90 evidence summaries and 136 recommendations were generated by 47 consensus panelists representing 20 health organizations and institutions. 11 evidence summaries were updated within the first 6 months. The second phase of the project was conducted in the last quarter of 2021 wherein 73 evidence summaries were generated (32 new topics and 41 updated from the first phase). A summary of the latest recommendations is presented here.

The CPG recommendations were used in constructing management algorithms for COVID-19. Process evaluation using website analytics revealed that the CPG recommendations were mostly accessed in regions with the greatest number of new cases and active cases. Furthermore, a list of top evidence summaries accessed reflected topics that CPG users needed the most guidance on, or that remain to be contentious as of the present date.

Severity Classification of COVID-19

The Philippine COVID-19 Living CPG used the following definitions for the spectrum of severity of COVID-19 (as of October 28, 2021):

Mild COVID-19 – no pneumonia or desaturation, acute onset of fever and cough or any three (3) or more of the following: fever, cough, coryza, sore throat, diarrhea, anorexia/nausea/vomiting, loss of sense of smell or taste, general weakness/body malaise/fatigue, headache, myalgia

Moderate COVID-19 – with pneumonia*, BUT no difficulty of breathing or shortness of breath, RR < 30 breaths/min, oxygen saturation[#] \geq 94% at room air; **OR** without pneumonia but with risk factors for progression: elderly (\geq 60 years old) and/or with comorbidities

Severe COVID-19 – with pneumonia and ANY one of the following: signs of respiratory distress, oxygen saturation < 94% at room air, RR \geq 30 breaths/minute, requiring oxygen supplementation

Critical COVID-19 – with pneumonia and ANY one of the following: impending respiratory failure requiring high flow oxygen, non-invasive or invasive ventilation, acute respiratory distress syndrome, sepsis or shock, deteriorating sensorium, multi-organ failure, acute thrombosis

*Pneumonia – evidence of lower respiratory disease during clinical assessment (e.g. cough, fever plus crackles) and/or imaging (CXR, ultrasound, CT scan)

[#]Proper recording of the O₂ saturation: finger should be inserted in the oximeter for about 10-20 seconds, patient should be still and not talking



Summary of Recommendations on Screening and Diagnosis

Recommendation	Strength of Recommendation	Certainty of Evidence
<p>We suggest doing an initial screening for ANY influenza-like illness, typical and atypical COVID-19 symptoms* within the past 14 days in apparently healthy adults and children, especially for individuals with known exposure to a laboratory-confirmed case of COVID-19</p> <p><i>Symptoms include but not limited to: fever/chills, cough, shortness of breath/dyspnea, sore throat, runny nose, myalgia, headache, fatigue/malaise, diarrhea, nausea/vomiting, abdominal pain, anosmia, ageusia, wheezing, chest pain, altered mental status, seizures, rash, pink eye</i></p>	Weak	Very Low
<p>We suggest pulse oximetry with close clinical monitoring by qualified medical personnel in suspected and confirmed COVID-19 patients especially those who are at high risk for deterioration.</p>	Weak	Very Low
<p>We recommend the use of the following specimens as alternative specimens to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19 among symptomatic and asymptomatic patients suspected of COVID-19 in hospital and outpatient settings:</p> <ul style="list-style-type: none"> • oropharyngeal swab • saliva drool/spit and oral saliva • nasal swab/wash • throat swab <p><i>*SARS COV-2 RT-PCR of nasopharyngeal swabs remains the diagnostic test of choice to confirm the diagnosis of COVID-19 among suspected individuals.</i></p>	Strong	Moderate Moderate Moderate Low
<p>We suggest the use of saliva swab and posterior oropharyngeal saliva specimens as an alternative specimen to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19 among symptomatic and asymptomatic patients with suspected COVID-19 in hospital and community/outpatient settings.</p>	Weak	Low
<p>We recommend against the use of sputum as an alternative specimen to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19.</p>	Strong	Very Low
<p>There is no evidence to recommend the use of bronchoalveolar lavage as an alternative specimen to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19.</p>	-	-

Recommendation	Strength of Recommendation	Certainty of Evidence
We suggest the use of rapid antigen test for the diagnosis of symptomatic individuals suspected of COVID-19 as an alternative to RT-PCR if all the following conditions are met: <ul style="list-style-type: none"> Individuals are in the early phase of illness (less than or equal to 7 days from onset of symptoms) Testing kits demonstrated sensitivity of more than or equal to 80% AND have very high specificity of more than or equal to 97% 	Weak	Low
We suggest against the use of rapid antigen test for screening purposes	Weak	Low
We suggest against the use of saliva as specimen for rapid antigen test in patients suspected of COVID-19 infection.	Weak	Low
We suggest against the use of rapid antigen tests alone in asymptomatic patients suspected of COVID-19 infection	Weak	Low
We suggest the use of rapid antigen tests for the diagnosis of individuals suspected of COVID-19 during the setting of an outbreak provided that all the following conditions are met: <ul style="list-style-type: none"> Individuals are in the early phase of illness (less than or equal to 7 days from onset of symptoms); AND Testing kits demonstrated sensitivity of more than or equal to 80% AND have very high specificity of more than or equal to 97%. 	Weak	Very Low
There is insufficient evidence to recommend for or against the use of repeat antigen testing for screening or diagnosis of COVID-19.	-	Very Low
We suggest the use of self-administered rapid antigen test for the diagnosis of COVID-19 in symptomatic individuals, provided that ALL OF THE FOLLOWING conditions are met: <ol style="list-style-type: none"> Ease of collecting samples is ensured; Ease of interpretation is ensured; Test kits have passed flex studies; AND Individuals present with symptoms for less than 7 days. 	Weak	Low
We suggest against the use of self-administered rapid antigen test for routine screening of COVID-19	Weak	Low
There is insufficient evidence to recommend the use of breath test in detecting COVID-19 infection.	-	Low
We suggest the use of pooled RT-PCR testing in targeted* low-risk and low-prevalence populations using a pool size of 5 in individuals suspected of COVID-19 infection. *Target population refer to the list of PSP and DOH	Weak	Moderate

Recommendation	Strength of Recommendation	Certainty of Evidence
We suggest repeating RT-PCR testing when the initial RT-PCR test is negative among symptomatic patients highly suspected to have COVID-19 infection.	Weak	Low
There is insufficient evidence to recommend an RT-PCR cycle threshold cut-off value* to determine infectivity among COVID-19 confirmed patients <i>* Interpretation of RT-PCR cycle threshold values may vary and is dependent on the PCR assay used, gene target, sample type, and timing of sample collection</i>	-	Very Low
We suggest using antibody tests that accurately measure IgG or total antibodies to determine COVID-19 seroprevalence among adults when needed for public health purposes.	Weak	Very Low
We suggest against using antibody tests detecting IgM to determine COVID-19 seroprevalence among adults	Weak	Very Low
We suggest against using lateral flow immunoassay (LFIA) tests to determine COVID-19 seroprevalence among adults	Weak	Very Low
We recommend against routine measurement of SARS-CoV-2 antibody titers after vaccination.	Strong	-
We recommend against the use of SARS-CoV-2 antibody testing to diagnose presumptive COVID-19 reinfection among symptomatic patients previously diagnosed with COVID-19.	Strong	Very Low
We recommend the use of both clinical risk assessment and RT-PCR* to screen for COVID-19 among asymptomatic individuals scheduled for non-emergency surgery. <i>*Use high-risk PPE regardless of RT-PCR or Ag-RDT test results in areas with prevalence of 1% or higher.</i>	Strong	Very Low
We recommend the use of both clinical risk assessment and Antigen-Rapid Diagnostic Test (Ag-RDT)** to screen for COVID-19 among asymptomatic individuals scheduled for non-emergency surgery when RT-PCR testing is not available or when turnaround time of results is prolonged. <i>**Ag-RDT should have a Sn of 80% and Sp of 97%</i>	Strong	Very Low
For asymptomatic, not severely immunocompromised fully vaccinated adults, we suggest the use of the following symptom-based criteria for return-to-work clearance: a. At least 8 days have passed since the first positive COVID-19 RT-PCR test; AND a. No symptoms have developed during this period.	Weak	Very Low
For asymptomatic, not severely immunocompromised not fully vaccinated adults, we suggest the use of the following symptom-based criteria for return to work clearance:	Weak	Very Low

Recommendation	Strength of Recommendation	Certainty of Evidence
<p>a. At least 10 days have passed since the first positive COVID-19 RT-PCR test; AND</p> <p>No symptoms have developed during this period.</p>		
<p>For symptomatic, not severely immunocompromised adults with mild-to-moderate COVID-19 diagnosis and any vaccination status, we suggest the use of the following symptom-based criteria for return to work clearance:</p> <p>a. At least 10 days have passed since the onset of symptoms; AND</p> <p>b. No fever during the previous 24 hours; AND</p> <p>c. There has been substantial improvement in respiratory symptoms of the acute illness.</p>	Weak	Very Low
<p>For symptomatic, not severely immunocompromised adults with severe-to-critical COVID-19 diagnosis and any vaccination status, we suggest the use of the following symptom-based criteria for return to work clearance: (<i>Very low certainty of evidence; Weak recommendation</i>)</p> <p>a. At least 21 days have passed since the onset of symptoms; AND</p> <p>b. No fever during the previous 24 hours; AND</p> <p>c. There has been substantial improvement in respiratory symptoms of the acute illness.</p>	Weak	Very Low
<p>For symptomatic, severely immunocompromised adults* with any vaccination status, we suggest the use of the following for return to work clearance:</p> <p>b. At least 22 days have passed since the onset of symptoms; AND</p> <p>c. No fever during the previous 24 hours; AND</p> <p>d. There has been substantial improvement in respiratory symptoms of the acute illness; AND</p> <p>a. PCR test results are negative on at least 1 respiratory specimen</p>	Weak	Very Low
<p>We suggest against the use of chest x-ray to diagnose COVID 19 infection among asymptomatic individuals.</p>	Weak	Very Low
<p>We suggest chest x-ray to facilitate rapid triage, infection control and clinical management among any of the following:</p> <p>a. patients with mild features of COVID 19 at risk for progression</p> <p>b. patients with moderate to severe features of COVID 19 patients with symptoms of at least 5 days duration</p>	Weak	Very Low
<p>We suggest against the routine use of CT scan for diagnosing COVID-19 among suspected patients with COVID-19 presenting at the emergency department if RT-PCR testing is readily available with timely results.</p>	Weak	Very Low

Recommendation	Strength of Recommendation	Certainty of Evidence
<p>If RT-PCR test is not available, we suggest using non-contrast chest CT scan for symptomatic patients suspected of having COVID-19 to guide early triage and management under the following conditions:</p> <ul style="list-style-type: none"> • Patients with mild COVID-19 who are at risk for progression (elderly, with comorbidities) • Patients with moderate to severe COVID-19 	Weak	Very Low
We suggest against the use of lung ultrasound alone in diagnosing patients with suspected COVID-19 infection.	Weak	Low
<p><u>To guide the decision to admit patients with COVID-19 to the hospital:</u></p> <p>We suggest the use of the following scoring systems:</p> <ol style="list-style-type: none"> a. Age, BUN, number of Comorbidities, CRP, SpO₂/FiO₂ ratio, Platelet count, Heart rate (ABC2-SPH) risk score, b. Confusion Urea Respiration Blood Pressure (CURB-65) severity score, c. Risk Stratification in the Emergency Department in Acutely Ill Older Patients (RISE-UP) score, and d. Rapid Emergency Medicine Score (REMS) <p>There is insufficient evidence to recommend the use of the 4C Mortality Score, COVID Outcome Prediction in the Emergency Department (COPE) model, and Quick Sepsis-related Organ Failure Assessment (qSOFA) score. (<i>Very low certainty of evidence</i>)</p>	Weak	Low
<p><u>To guide in the expectant monitoring of hospitalized patients:</u></p> <p>We suggest the use of the 4C Deterioration model.</p> <p>There is insufficient evidence to recommend the use of the Modified Early Warning Score (MEWS) and National Early Warning Score 2 (NEWS2) scoring systems.</p>	-	Very Low
There is insufficient evidence to recommend the use of specific cut-off values of CRP, LDH and Ferritin to guide immunotherapy in COVID-19.	-	Very Low
We suggest the use of D-dimer to guide anticoagulation of patients with COVID-19, because of its significant association with mortality, thromboembolism, and worsening severity of disease.	Weak	Low
<p>There is insufficient evidence in using symptoms*, biologic factors or severity of acute COVID-19 in predicting the development of long COVID-19 symptoms.</p> <p><i>*The most common symptoms of long COVID-19 identified were fatigue, dyspnea, sleep disturbance, anxiety or depression, and memory impairment</i></p>	-	Very Low

Recommendation	Strength of Recommendation	Certainty of Evidence
We suggest against the use of procalcitonin alone as a basis for initiating antibiotic therapy among COVID-19 confirmed patients	Weak	Very Low
If available, we recommend using a procalcitonin level of less than or equal to 0.25ng/ml for discontinuing antibiotic therapy among COVID-19 confirmed patients.	Strong	Very Low
We suggest against the use of PF4 antibody ELISA Heparin Induced Thrombocytopenia (HIT) test kits and non-ELISA rapid HIT test kits for COVID-19 Vaccine Induced Thrombosis and Thrombocytopenia (VITT).	Weak	Low
We suggest against using serum tryptase for patients who had anaphylaxis after receiving COVID-19 vaccine.	Weak	Very Low



Summary of Recommendations on Treatment

Recommendation	Strength of Recommendation	Certainty of Evidence
We recommend against the use of hydroxychloroquine/chloroquine, with or without azithromycin among patients with COVID-19 infection.	Strong	Moderate
We recommend against the use of azithromycin among patients with COVID-19 infection.	Strong	Moderate
There is insufficient evidence to recommend the use of favipiravir among patients with COVID-19.	-	Low
We suggest against the use of remdesivir in patients with COVID-19 infection who have O ₂ saturation $\geq 94\%$ and do not require oxygen supplementation.	Weak	Low
We suggest the addition of remdesivir to dexamethasone in patients with COVID-19 infection who have O ₂ saturation $< 94\%$ and/or requiring oxygen supplementation.	Weak	Low
We suggest against the use of remdesivir in patients with COVID-19 infection who are already on invasive mechanical ventilation or ECMO. <i>*For patients who progress to invasive mechanical ventilation while on remdesivir, the drug can be continued.</i>	Weak	Low
We suggest the use of molnupiravir within 5 days of symptom onset among non-hospitalized adult patients (18 years old and older) with mild to moderate COVID-19 infection with at least one risk factor* for progression.	Weak	Low
We suggest against the use of baloxavir as treatment for patients with COVID-19 infection.	Weak	Very low
We recommend against the use of oseltamivir as treatment for patients with COVID-19 infection.	Strong	Very low
We recommend against the use of lopinavir/ritonavir as treatment for COVID-19 infection.	Strong	Moderate
We recommend the addition of tocilizumab to systemic steroids in patients showing rapid respiratory deterioration and/or requiring high doses of oxygen (high-flow nasal cannula, noninvasive or invasive mechanical ventilation) and with elevated biomarkers of inflammation (CRP).	Strong	Moderate
We recommend against the use of tocilizumab among patients with COVID-19 infection who do not require oxygen.	Strong	Very low

Recommendation	Strength of Recommendation	Certainty of Evidence
We suggest the use of baricitinib in addition to dexamethasone and remdesivir as treatment for hospitalized COVID-19 patients who require low-flow oxygen, high-flow oxygen, and non-invasive ventilation.	Weak	Low
There is insufficient evidence to recommend baricitinib as an alternative to tocilizumab as treatment for hospitalized COVID-19 patients.	-	Very low
There is insufficient evidence to recommend the use of imatinib among patients with COVID-19 infection	-	Low
We suggest against the use of tofacitinib among hospitalized COVID-19 patients	Weak	Low
We suggest against the use of leronlimab as treatment for COVID-19	Weak	Very low
We suggest against the use of infliximab among patients with COVID-19 infection	Weak	Very Low
We suggest against the use of bevacizumab as treatment for patients with COVID-19 infection	Weak	Very Low
We recommend against the use of ivermectin for the treatment of patients of any severity	Strong	Very low
We suggest against the use of ivermectin combined with doxycycline for the treatment of patients with COVID-19.	Weak	Very low
We suggest against the use of artesunate, artemisinin or pyronaridine tetraphosphate + artesunate in the treatment of COVID-19	Weak	Very Low
We suggest against the use of colchicine in the treatment of COVID-19.	Weak	Low
We suggest against the use of interferon in the treatment of hospitalized patients with moderate to critical COVID-19.	Weak	Very low
There is insufficient evidence to recommend the use of fluvoxamine among COVID-19 patients	-	Low
We suggest the use of bamlanivimab and etesevimab combination therapy as treatment for mild to moderate, non-hospitalized COVID-19 patients with at least 1 risk factor for progression to severe disease.	Weak	Low
We suggest casirivimab + imdevimab as treatment for symptomatic, non-hospitalized patients with at least 1 risk factor for severe COVID-19.	Weak	Moderate
We recommend against casirivimab + imdevimab as treatment for hospitalized COVID-19 patients.	Strong	Low
There is insufficient evidence to recommend casirivimab-imdevimab as treatment for asymptomatic COVID-19 patients	-	Low
We suggest against the use of regdanvimab for the treatment of mild to moderate COVID-19	Weak	Very Low

Recommendation	Strength of Recommendation	Certainty of Evidence
We recommend against the use of convalescent plasma among patients with COVID-19 infection.	Strong	Moderate
We suggest against the use of intravenous immunoglobulin as treatment for moderate to severe COVID-19.	Weak	Very low
There is insufficient evidence to recommend using umbilical cord-derived mesenchymal stem cell therapy among adults with severe COVID-19.	-	Very low
There is insufficient evidence to recommend the use of inhaled corticosteroids as treatment for non-hospitalized patients	-	Very low
We recommend against the use of steam inhalation alone in the treatment of COVID-19.	Strong	Very low
There is no evidence to recommend the use of VCO as treatment among patients with COVID-19 infection.	-	-
There is insufficient evidence to recommend the use of Lianhua in treatment of patients with non-severe COVID-19	-	Very low
We suggest against the use of famotidine in the treatment of COVID-19.	Weak	Very low
We recommend against the use of ibuprofen as treatment among patients with COVID-19 infection.	Strong	Very low



Summary of Recommendations on Critical Care and Respiratory Management

Recommendation	Strength of Recommendation	Certainty of Evidence
We recommend the use of dexamethasone for up to 10 days among patients with severe and critical COVID-19.	Strong	High
We recommend the use of 6 mg to 12 mg per day of dexamethasone among patients with severe and critical COVID-19.	Strong	Moderate
We recommend against the use of corticosteroids among mild and moderate (non-oxygen requiring) COVID-19 patients	Strong	Moderate
We suggest that steroid therapy be initiated as soon as diagnosed or categorized as severe or critical COVID-19.	Weak	Very Low
We recommend the use of prophylactic over therapeutic dose anticoagulation among hospitalized patients with moderate, severe or critical COVID-19 disease unless there are any contraindications.	Strong	Low
We recommend the use of standard dose prophylactic anticoagulation over intermediate dose prophylactic anticoagulation among hospitalized patients with COVID-19 disease unless there are any contraindications.	Strong	Moderate
We recommend against the routine use of antibiotics in patients with severe and critical COVID-19 infection, unless with suspicion of secondary bacterial co-infection. For patients on empiric antibiotics, they should be assessed daily for the need for discontinuation, continuation or de-escalation based on clinical and laboratory parameters.	Strong	Very low
There is insufficient evidence on the use of hemoperfusion among patients with COVID-19 infection.	-	Very low
We suggest the use of conservative fluid management rather than liberal fluid management strategy in mechanically ventilated adult COVID-19 patients with acute respiratory distress syndrome who have been adequately resuscitated*. <i>*without tissue hypoperfusion and fluid responsiveness</i>	Weak	Low
We suggest self-proning in non-intubated patients with severe and critical COVID-19	Weak	Very low
There is insufficient evidence to recommend the use of side lying in non-intubated patients with severe and critical COVID-19	-	Very low
We suggest the use of high-flow nasal cannula oxygenation rather than non-invasive ventilation (e.g., helmet CPAP, mask NIV) in patients with COVID-19 infection and acute	Weak	Very low

Recommendation	Strength of Recommendation	Certainty of Evidence
hypoxemic respiratory failure who do not respond to conventional oxygen therapy.		
We suggest the use of a lung protective ventilation strategy (tidal volume 4-8 mL/kg predicted body weight and plateau pressure less than 30 cmH ₂ O) in patients with COVID-19 infection and ARDS.	Weak	Very low
There is insufficient evidence to recommend the use of a higher PEEP strategy. We suggest individualizing PEEP or employ a PEEP strategy based on respiratory mechanics (i.e., compliance) in patients with COVID-19 infection.	Weak	Low
There is insufficient evidence to recommend a driving pressure limited strategy in patients with COVID-19 infection. We suggest keeping the driving pressure \leq 14 cmH ₂ O.	Weak	Low
We suggest the use of rapid sequence intubation for COVID-19 patients to reduce infection among healthcare workers performing the procedure.	Weak	Very low
We suggest the use of Venovenous Extracorporeal Membrane Oxygenation (ECMO) for judiciously selected COVID-19 patients with severe ARDS based on the Extracorporeal Life Support Organization (ELSO) criteria.	Weak	Very low
There is insufficient evidence to recommend the use of hyperbaric oxygen therapy for the management of COVID-19 patients.	-	Very low
We suggest light over deep sedation in COVID-19 patients who are mechanically ventilated and who are anxious or agitated.	Weak	Very low
We recommend against the use of nitric oxide among patients with COVID-19	Strong	Low
We recommend against the use of etoposide among patients with COVID-19 pneumonia in cytokine storm.	Strong	Very low
We recommend individualized pulmonary rehabilitation with pre-intervention medical clearance for long COVID patients who show residual respiratory symptoms.	Strong	Moderate
There is insufficient evidence to recommend the use of pirfenidone or nintedanib among patients with post COVID-19 pulmonary fibrosis	-	Very low




Summary of Recommendations on Non-Pharmacologic Interventions

Recommendation	Strength of Recommendation	Certainty of Evidence
We recommend that healthcare workers not directly taking care of COVID-19 patients, and other persons with high risk of exposure to COVID-19 should use properly fitted surgical masks instead of cloth masks.	Strong	Moderate
We suggest using a cloth mask that fits snugly on the face and made of at least two layers of cotton (e.g., t-shirt fabric) or non-woven nylon with aluminum nose bridge for the general public with low risk of exposure to COVID-19 in outdoor or indoor areas to prevent COVID-19 infections.	Weak	Low
We suggest against requiring the use of face shields in addition to face masks among the general public in non-healthcare settings.	Weak	Very low
We recommend the addition of face shields to face masks among the general public in areas with sustained community transmission of SARS-CoV-2.	Strong	Very low
We recommend using medical face mask plus face shield and standard personal protective equipment among health care workers not directly involved in the care of COVID-19 patients in areas with sustained community transmission of SARS-COV2.	Strong	Very low
There is no evidence to recommend the use of copper-containing over non-copper-containing masks to decrease COVID-19 transmission	-	-
We recommend the use of appropriate PPE to include mask (N95 or higher), fluid repellent sealed well-fitting long gown, double gloves, apron, full face shield or goggles or visor, scrub hat, and disposable shoe covers or dedicated closed footwear among surgeons engaged in aerosol generating procedures of suspected or confirmed COVID-19 patients.	Strong	Very low
We recommend the use of at least a surgical face mask and face shield for protection against COVID-19 infection among healthcare workers in the outpatient setting not performing aerosol generating procedures. Additional PPEs such as medical gowns and gloves should be worn as part of standard precautions during the performance of other procedures.	Strong	Very low
We recommend the use of the following PPE: disposable hat, medical protective mask (N95 or higher standard), goggles or face shield (anti-fog), medical protective clothing, disposable gloves and disposable shoe covers or dedicated closed footwear as an effective intervention	Strong	Moderate

Recommendation	Strength of Recommendation	Certainty of Evidence
in the prevention of COVID-19 among health care workers in areas with possible direct patient care of confirmed or probable COVID-19 patients and possible performance of aerosol generating procedures.		
We recommend against the use of ionizing air purifier to reduce COVID-19 transmission in the community.	Strong	Low
We recommend against the use of footbaths for the prevention and control of COVID-19 transmission.	Strong	Very low
We recommend against the use of misting tents or disinfection chambers for preventing and controlling COVID-19 transmission.	Strong	Very low
We recommend against the use of UV lamps or other UV devices in any place outside of a controlled clinic or hospital setting to prevent and control COVID-19 transmission.	Strong	Low
We suggest the use of HEPA filter as an option to improve air quality in indoor spaces with inadequate ventilation.	Weak	Low
We recommend the use of carbon dioxide (CO ₂) monitors in enclosed spaces to guide actions to improve ventilation and reduce transmission of SARS-COV-2	Strong	Moderate
In situations where there is shortage of filtering facepiece respirators (FFR), we suggest the use of Hydrogen Peroxide Vapor (HPV), Ultraviolet Germicidal Irradiation (UVGI), moist heat and peracetic acid dry fogging system (PAF) as options for N95 mask decontamination as recommended by the manufacturer based on their ability to reduce SARS-COV-2 load while still maintaining N95 mask integrity.	Weak	Low
We recommend against the use of autoclave and alcohol as these methods alter filtering facepiece respirator's (N95) integrity and degrade filtration efficacy.	Strong	Very low
We suggest against the use of protective physical barrier enclosures (ex. aerosol box) for the prevention of COVID-19 among health care providers who perform aerosol generating medical procedures*. <i>*Proper PPEs should be used by health care providers when performing aerosol-generating procedures.</i>	Weak	Very low
We suggest using protective physical barriers in areas where physical distancing cannot be adhered to (e.g., offices, reception desk)**. <i>**Adequate ventilation, physical distancing, use of facemasks and personal hygiene should still be maintained to prevent COVID-19 infections. Regular cleaning and disinfection of physical barriers should be practiced.</i>	Weak	Very low
We recommend cleaning and disinfecting surfaces using the appropriate disinfecting chemical agents such as	Strong	Low

Recommendation	Strength of Recommendation	Certainty of Evidence
<p>0.5% sodium hypochlorite solution (bleach) or 70% alcohol to prevent COVID-19 infection.</p> <p>For high touch surfaces and high traffic areas, such as in the workplace, disinfection should be done before shift, intermittently during shift and after the shift.</p> <p>For household disinfection, once daily disinfection of high touch surfaces is recommended.</p>		



Summary of Recommendations on Vaccines and Prophylactic Interventions

Recommendation	Strength of Recommendation	Certainty of Evidence
<p>We recommend the use of the following vaccines to prevent symptomatic SARS-CoV-2 infection among adults:</p> <ul style="list-style-type: none"> a. BNT162b2 (Pfizer/BioNTech) (given as 0.3ml (30ug) intramuscular injections, in 2 doses, 21 days apart) b. mRNA-1273 (Moderna) (given as 0.5ml (100ug) intramuscular injections, in 2 doses, 28 days apart) c. ChAdOx1 (AstraZeneca) (given as 0.5 ml (5 x 10⁶ vp) intramuscular injections, in 2 doses, at least 12 weeks apart) d. Gam-COVID-Vac (Gamaleya) (given as rAd-26 0.5ml intramuscular injection, then rAd-5S 0.5 ml intramuscular injection 21 days after) e. Ad26.COVS.2.S (Janssen/Johnson&Johnson) (given as 0.5ml single dose intramuscular injection) 	Strong	Moderate
<p>We recommend the use of CoronaVac (Sinovac) (given as 0.5ml (600SU) intramuscular injection, in 2 doses, at 28 days apart) to prevent symptomatic SARS-CoV-2 infection among healthy adults.</p>	Strong	Low
<p>We recommend the use of BNT162b2 (Pfizer/BioNTech), mRNA-1273 (Moderna), ChAdOx1 (Astrazeneca), Gam-COVID-Vac (Gamaleya) and Ad26.COVS.2.S (Janssen/Johnson&Johnson) vaccines to prevent symptomatic SARS-CoV-2 infection in older adults (>64-year-old).</p>	Strong	Low
<p>We suggest the use of CoronaVac (Sinovac) to prevent SARS-COV-2 infection in older adults (>60 years old)</p>	Weak	Low
<p>We recommend the use of BNT162b2 (Pfizer/BioNTech), mRNA-1273 (Moderna), ChAdOx1 (Astrazeneca), Gam-COVID-Vac (Gamaleya), CoronaVac (Sinovac) and Ad26.COVS.2.S (Janssen/Johnson&Johnson) vaccines in pregnant and lactating women after consultation with a physician.</p>	Strong	Very low
<p>We recommend the use of BNT162b2 (Pfizer/BioNTech), mRNA-1273 (Moderna), ChAdOx1 (Astrazeneca), Gam-COVID-Vac (Gamaleya) and Ad26.COVS.2.S (Janssen/Johnson&Johnson) vaccines to prevent SARS-CoV-2 infection in adults who have stable medical comorbidities and are at risk for severe infection.</p>	Strong	Moderate
<p>We suggest the use of CoronaVac (Sinovac) to prevent SARS-CoV-2 infection in adults who have stable medical comorbidities and are at risk for severe infection.</p>	Weak	Very low

Recommendation	Strength of Recommendation	Certainty of Evidence
We recommend the use of BNT162b2 (Pfizer/BioNTech), mRNA-1273 (Moderna), ChAdOx1 (Astrazeneca), Gam-COVID-Vac (Gamaleya), CoronaVac (Sinovac) and Ad26.COV2.S (Janssen/ Johnson&Johnson) vaccines to prevent SARS-CoV-2 infection in immunocompromised patients (i.e., diagnosed with HIV, hepatitis B and C, those with cancer undergoing chemotherapy, transplant patients receiving immune-suppression) after medical clearance from a physician.	Strong	Low
We recommend against the use of these vaccines for those who have known allergies to the contents / excipients of the vaccine, such as polysorbate: (ChAdOx1 (Astrazeneca), Gam-COVID-Vac (Gamaleya) and Ad26.COV2.S (Janssen/ Johnson&Johnson) and polyethylene glycol or PEG200 DMG (BNT162b2 (Pfizer/BioNTech) and mRNA-1273 (Moderna).	Strong	Moderate to high
We recommend the use of BBIBP-CorV (Sinopharm), given as 200U (WIV04) or 4ug (HBO2) in 0.5 ml in 2 doses, 21 days apart, to prevent symptomatic and asymptomatic COVID-19 infection among healthy adults (18 to 59 years old)	Strong	Moderate
We suggest the use of BBIBP-CorV to prevent severe COVID-19 infection among healthy adults (18 to 59 years old)	Weak	Low
We suggest the use of BBIBP-CorV to prevent symptomatic COVID-19 infection in the following: a. Adults with co-morbidities b. Older persons	Weak	Very Low
There is insufficient evidence to recommend for or against the use of BBIBP-CorV to prevent COVID-19 infection among the following: a. Children (3-17 years old) b. Immunocompromised population c. Pregnant and lactating women	-	Very Low
In areas where the SARS-CoV-2 variants of concern are prevalent, there is insufficient evidence to recommend for or against the use of BBIBP-CorV to prevent COVID	-	Very Low
We recommend the use of the CoronaVac (Sinovac), [given as 0.5 mL (600SU)] to prevent symptomatic SARS-CoV-2 infection in: <ul style="list-style-type: none"> • Healthy Adults • Pregnant women in their first trimester after consultation with a physician • Pregnant women in their 2nd and 3rd trimester and lactating women • Adults who have medical comorbidities • Immunocompromised patients after medical clearance from a physician (the immunocompromised include those diagnosed with HIV, hepatitis B and C, those with cancer 	Strong Strong Strong Strong Strong	Low Very Low Very Low Low Low

Recommendation	Strength of Recommendation	Certainty of Evidence
undergoing chemotherapy, transplant patients receiving immunosuppression)		
We suggest the use of CoronaVac (Sinovac) to prevent SARS-CoV-2 infection in older adults (>60 years old)	Weak	Low
We suggest against the use of CoronaVac (Sinovac) to prevent SARS-CoV-2 infection in children (3 to 17 years old)	Weak	Very Low
In areas where Delta is the predominant variant of concern, we recommend the use of CoronaVac (Sinovac)	Strong	Very Low
Under the current context of low vaccine coverage and inadequate vaccine supply, we recommend against booster vaccination using CoronaVac (Sinovac) in the healthy, adult population (18 years old and above)	Strong	Low
For immunocompromised patients who received primary CoronaVac (Sinovac) vaccination, we recommend for heterologous booster vaccination	Strong	Very Low
We recommend the use of BBV152 (Covaxin/Bharat), 0.5 mL/dose, in a two-dose regimen, 28 days apart for the prevention of symptomatic COVID-19 infection in healthy adults	Strong	Moderate
We suggest the use of BBV152 (Covaxin/Bharat), 0.5 mL/dose, in a two-dose regimen, 28 days apart for the prevention of symptomatic COVID-19 infection:		
a. Adults who have stable medical co-morbidities and are at high risk for severe infection	Weak	Low
b. Healthy, older adults (>60 years old)	Weak	Low
c. Pregnant and lactating women, after discussing with a physician	Weak	-
d. Immunocompromised patients, after discussing with a physician	Weak	-
We suggest against the use of BBV152 (Covaxin/Bharat) for the prevention of COVID-19 in children and adolescents	Weak	-
We recommend against the use of BBV152 (Covaxin/Bharat) in individuals who have known allergies to its contents/excipients	-	-
We suggest the use of the rAd26 (Sputnik Light), given as 10 ¹¹ vp per 0.5ml, single dose, intramuscularly to prevent symptomatic SARS-CoV-2 infection in:	Weak	Low
a. Healthy adults		
b. Older adults (60 years and older)		
c. Adults with comorbidities		
We suggest against the use of rAd26 (Sputnik Light) to prevent symptomatic SARS-CoV-2 infection in:		
a. Children (3-17 years)	Weak	-
b. Pregnant and lactating women		
c. Immunocompromised		

Recommendation	Strength of Recommendation	Certainty of Evidence
In areas where Alpha, Beta or Delta is the predominant variant of concern, we suggest the use of rAd26 (Sputnik Light) to prevent COVID-19 infection.	Weak	Low
We suggest the following homologous booster vaccination regimen for the general adult population:		
a. BNT162b2	Weak	Low
b. mRNA-1273	Weak	Low
c. ChAdOx1	Weak	Very Low
d. Ad26.Cov2.	Weak	Very Low
e. CoronaVac	Weak	Very Low
f. BBIBP-CorV	Weak	Very Low
There is insufficient evidence to recommend the following homologous booster vaccination in the general population:	-	-
a. Gam-COVID-Vac		
b. BBV152		
We suggest the following heterologous booster vaccination regimen for the general adult population:		
a. BNT162b2 primary, mRNA-1273 booster		
b. BNT162b2 primary, Ad26.CoV2.S booster		
c. mRNA-1273 primary, BNT162b2 booster		
d. mRNA-1273 primary, Ad26.CoV2.S booster		
e. ChAdOx1 primary, BNT162b2 booster	Weak	Very Low
f. Ad26.COVID.S primary, BNT162b2 booster		
g. Ad26.COVID.S primary, mRNA-1273 booster		
h. CoronaVac primary, BNT162b2 booster		
i. CoronaVac primary, ChAdOx1 booster		
j. BBIBP-CorV primary, BNT162b2 booster		
We suggest the following homologous booster vaccination for the immunocompromised population:		
a. BNT162b2	Weak	Very Low
b. mRNA-1273	Weak	Low
There is insufficient evidence to recommend the following homologous booster vaccination for the immunocompromised population:		
a. ChAdOx1		
b. Ad26.CoV2.S	-	-
c. CoronaVac		
d. Gam-COVID-Vac		
e. BBV152		
f. BBIBP-CorV		
We suggest the following heterologous booster vaccination regimen for the immunocompromised population:		
a. an mRNA vaccine primary, another mRNA vaccine booster	Weak	Very Low
b. an mRNA vaccine primary, ChAdOx1 booster	Weak	Low
c. BNT162b2 primary, mRNA-1273 booster	Weak	Very Low
d. BNT162b2 primary, Ad26.CoV2.S booster	Weak	Very Low
e. mRNA-1273 primary, Ad26.CoV2.S booster	Weak	Very Low

Recommendation	Strength of Recommendation	Certainty of Evidence
We recommend the use of heterologous COVID-19 vaccination for those with serious adverse event to the first dose.	Strong	Very Low
We suggest the use of heterologous COVID-19 vaccination in the event of the unavailability of the second dose in the recommended schedule	Weak	Very Low
In areas where the Delta variant is the predominant circulating variant, we recommend the use of the following vaccine to prevent symptomatic and severe COVID-19:		
a. 2 doses of BBV152 (Covaxin/Bharat)	Strong	Moderate
b. 2 doses of BNT162b2 (Pfizer)	Strong	Low
c. 2 doses of mRNA-1273 (Moderna)	Strong	Low
d. 2 doses of ChAdOx1 (Astra Zeneca)	Strong	Low
2 doses of CoronaVac (Sinovac)	Strong	Very Low
In areas where the Delta variant is the predominant circulating variant, we suggest the use of the following vaccines to prevent symptomatic and severe COVID-19:		
a. Ad26.CoV2 (Janssen)	Weak	Low
b. Gam-COVID-Vac (Sputnik V)	Weak	Low
We recommend the use of the BNT162b2 (Pfizer/BioNTech) vaccine, [given as 0.3 mL (30 ug) intramuscular injections, in 2 doses, 21 days apart] for children 12-15 years old to prevent symptomatic SARS-CoV-2 infection.	Strong	Moderate
We suggest the use of the mRNA-1273 (Moderna) vaccine, [given as 0.5 mL (100 ug) intramuscular injections, in 2 doses, 28 days apart] for children 12-17 years old to prevent symptomatic SARS-CoV-2 infection.	Weak	Low
We suggest against the use of Coronavac (Sinovac), [given as 0.5 mL (600 SU) intramuscular injection, in 2 doses, 28 days apart] for children 3-17 years old to prevent symptomatic SARS-CoV-2 infection.	Weak	-
There is no evidence on the use of ChAdOx1 (Astrazeneca), Gam-COVID-Vac (Gamaleya), Ad26.COVS.2.S (Janssen/ Johnson&Johnson) among children <18 years old to prevent SARS-CoV-2 infection.	-	-

Recommendation	Strength of Recommendation	Certainty of Evidence
We suggest the use of following vaccines, after the first trimester, for the prevention of COVID-19 infection in pregnant and lactating women.		
a. BNT162b2 (Pfizer)	Weak	Low
b. mRNA-1273 (Moderna)	Weak	Low
c. ChAdOx1 (AstraZeneca)	Weak	-
d. Ad26.CoV2.S (Janssen/Johnson&Johnson)	Weak	-
e. CoronaVac (Sinovac)	Weak	-
f. BBIBP-CorV (Sinopharm)	Weak	-
g. BBV152 (Covaxin)	Weak	-
We suggest <u>against</u> the use of the following vaccines for the prevention of COVID-19 infection in pregnant and lactating women:		
a. Gam-CoV-Vac (Sputnik V)	Weak	-
b. NVX-2373 (Novavax)	Weak	-
We suggest against the use of BCG vaccine for the prevention of COVID-19 infection.	Weak	Very low
We suggest the use of casirivimab-imdevimab as day 4 post-exposure prophylaxis for COVID-19 close contacts, ages 12 years and above weighing at least 40kg, who are at risk for severe disease or hospitalization	Weak	Moderate
We recommend against the use of melatonin as prevention for COVID-19 infection.	Strong	Very low
We recommend against the use of Vitamin D supplementation to prevent COVID-19 infection.	Strong	Very low
We recommend against the use of zinc supplementation to prevent COVID-19 infection.	Strong	Very low
We recommend against the use of HCQ for pre-exposure prophylaxis in adults who are at high risk of exposure to COVID-19 cases.	Strong	Moderate
We recommend against the use of HCQ for post-exposure prophylaxis in adults who are exposed to COVID-19 cases.	Strong	Low
We recommend against the use of lopinavir/ritonavir for chemoprophylaxis in individuals exposed to COVID-19 patients.	Strong	Very low
We recommend against the use of ivermectin as COVID-19 prophylaxis for the general population.	Strong	Very low
We recommend against the use of ivermectin for COVID-19 as post-exposure prophylaxis for household contacts of confirmed COVID-19 patients.	Strong	Very low
We recommend against the use of ivermectin for COVID-19 as prophylaxis for healthcare workers.	Strong	Very low

Recommendation	Strength of Recommendation	Certainty of Evidence
There is insufficient evidence to recommend the use of saline nasal irrigation (SNI) to prevent COVID-19 in healthy individuals.	-	Very low
We recommend against the use of steam inhalation in the prevention of COVID-19	Strong	Very low
There is insufficient evidence on the use of aspirin as prophylaxis against COVID-19-induced coagulopathy among patients with COVID-19.	-	Very low
There is insufficient evidence to recommend the use of antiseptic mouthwash or gargle to prevent COVID-19 in healthy individuals.	-	Very low



Summary of Recommendations on Adjunct Interventions

Recommendation	Strength of Recommendation	Certainty of Evidence
There is insufficient evidence to recommend the use of zinc as adjunct treatment for patients with COVID-19 infection.	-	Low
We suggest against the use of B vitamins as adjunct in the treatment of patients with COVID-19.	Weak	Very low
There is insufficient evidence to recommend the use of Vitamin C as adjunct treatment for patients with COVID-19 infection.	-	Low
There is insufficient evidence to recommend the use of Vitamin D supplementation as adjunct treatment for patients with COVID-19 infection.	-	Very low
There is insufficient evidence to recommend the use of melatonin as adjunct treatment for patients with COVID-19 infection.	-	Very low
There is no evidence to recommend the use of virgin coconut oil as adjunct treatment for patients with COVID-19 infection.	-	-
There is no evidence to recommend Lagundi (<i>Vitex negundo</i>) as adjunctive treatment for patients with COVID-19 infection	-	-
There is no evidence to recommend Tawa-tawa (<i>Euphorbia hirta</i>) as adjunctive treatment for patients with COVID-19 infection	-	-
There is insufficient evidence to recommend the use of fatty acid supplements as adjunctive treatment for patients with COVID-19.	-	Low
We recommend against the use of intravenous N-acetylcysteine as adjunct treatment for patients with COVID-19 infection.	Strong	Moderate
We recommend continuing maintenance RAAS blockers for hypertension among patients with COVID-19 infection.	Strong	Moderate
There is insufficient evidence to recommend statins as adjunctive treatment in patients with COVID-19	-	Very low
We suggest that ibuprofen may still be used as symptomatic treatment of patients with COVID-19 infection if clinically warranted. Concurrent use of ibuprofen is not associated with worsening of COVID-19 outcomes.	Weak	Very low
There is insufficient evidence to recommend discontinuation of aspirin as maintenance therapy for underlying medical conditions in patients with COVID-19.	-	Very low

Recommendation	Strength of Recommendation	Certainty of Evidence
We recommend against the use of any antiseptic mouthwash as an adjunctive therapy for patients with COVID-19	Strong	Very Low
We recommend against the use of any antiseptic mouthwash to prevent COVID-19 in healthy individuals	Strong	Very Low
We suggest against the use of nasal spray as an adjunct to treatment of COVID-19 infection	Weak	Low

INTRODUCTION

The coronavirus disease 2019 (COVID-19) has grown into a pandemic and global crisis infecting more than 572 million people worldwide and causing more than six million deaths [1]. As of August 27, 2022, the number of cases in the Philippines has reached more than 3.8 million with 61, 613 COVID-19 related deaths [2]. The national strategy towards the new normal is prevention, detection, isolation, treatment, and reintegration. Since the launch of the national vaccination campaign against COVID-19 in March 2021, the Philippines had 72 million fully vaccinated individuals as of August 26, 2022. Notwithstanding these strategies, none of the epidemiologic projections on COVID-19 in the Philippines point to a foreseeable end of the pandemic, especially with the rise of variants with increased transmissibility.

Given the magnitude of the impact of COVID-19 in the country, in addition to the concurrent infodemic potentially causing misinformation and disinformation among clinicians, public health officials, and policy makers, there is a need for evidence-based guidelines for the effective management and control of the spread of this disease. Existing international guidelines and living systematics reviews on COVID-19 need to be contextualized for the recommendations to be applicable to local end-users and other stakeholders.

Objectives

The Philippine COVID-19 Living CPG aimed to provide up-to-date, evidence-based recommendations on the treatment, diagnosis, infection prevention, and control of COVID-19 among adults with or at risk for COVID-19 using the GRADE methodology. Specifically, this project:

1. Identified priority questions related to COVID-19 management, infection prevention and control
2. Summarized available literature on each priority question related to COVID-19 management, infection prevention and control
3. Formulated recommendations on COVID-19 management, infection prevention, and control based on the evidence summaries presented
4. Updated selected recommendations on COVID-19 management, infection prevention and control based on predefined parameters

Target Population

This CPG was intended to apply primarily for adult Filipinos aged 18 years old and above diagnosed with, or at risk of COVID-19. The severity of COVID-19 was indicated in several recommendations if it is severity-specific. Other clinical characteristics, such as comorbidities, that would affect the recommendations were indicated clearly in the wording, as appropriate.

Intended Users

The following groups are the expected target users of this Living CPG:

1. Public health professionals, such as provincial/city/municipal health officers, program managers, public health nurses, etc., to inform their localized decisions in implementing national policies on COVID-19, such as on public health standards, management, and preventive interventions.
2. Clinicians in the hospitals, quarantine centers, and other treatment facilities handling COVID-19 patients, such as generalist physicians, internists, infectious disease specialists, pulmonologists, other specialist physicians, staff nurses, hospital administrators, etc., to inform their individual clinical decisions from diagnosis to treatment and prevention.
3. Academicians and researchers, especially those working on related COVID-19 topics, to guide their research initiatives in addressing the identified gaps during the evidence synthesis of this CPG
4. Policymakers and local government officials, such as the Department of Health, Philippine Health Insurance Corporation, Inter-agency Task Force for the Management of Emerging Infectious Diseases, Food and Drug Administration, Health Technology Assessment Council, etc., to inform their national policies on COVID-19, including standards of care in outpatient and in-patient settings

CPG DEVELOPMENT METHODOLOGY

The development process of the Philippine COVID-19 Living CPG followed the Philippine Department of Health's Manual for Clinical Practice Guideline Development [5] and the Grading of Recommendations, Assessment, Development and Evaluation or GRADE Approach [6]. The reporting of this CPG manuscript was based on the AGREE Reporting Checklist [7]. Some of the questions in the base CPG were updated following the living CPG methodology [8].

Overview of Philippine COVID-19 Living CPG Development Process

The following development process was undertaken by the Philippine COVID-19 Living CPG. Further details were presented in succeeding sections.

1. **Guideline Preparation** – The Steering Committee (SC) identified and convened members of the Living CPG task force: Evidence Review Experts (ERE) or Technical Working Group (TWG) and the Consensus Panel. A total of 24 specialty societies and stakeholders were represented in the second phase of the Philippine COVID-19 Living CPG. The SC reviewed and determined the scope of the guideline and also identified new and existing interventions that needed to be updated. Key questions were judged according to the criteria enumerated in the DOH Manual for CPG development

Several orientation sessions were conducted for the technical reviewers and consensus panel members on the COVID CPG development process. Technical reviewers were re-trained on evidence synthesis and the GRADE methodology. Consensus panel members were oriented on how to interpret the evidence summaries and apply the GRADE evidence-to-decision framework.

2. **Evidence Synthesis** - Evidence Review Experts (ERE) reviewed and appraised existing CPGs, systematic reviews, preprints and published literature, prepared evidence summaries, and drafted evidence-based recommendations. During the second phase of the CPG, five technical coordinators with expertise in CPG Development and Evidence-Based Medicine oversaw the retrieval and appraisal of evidence and the creation of the draft recommendations. Technical assistants were also assigned to each task force as search specialists to perform continuous surveillance for new evidence on existing guideline questions. Four technical writers ensured that the evidence summaries are uniform, concise, and clear. The Steering Committee organized several practice sessions for the ERE to finalize their presentations, and discuss them with other EREs, Steering Committee, and technical experts. Evidence summaries were collated, formatted, and prepared for presentation to the consensus panel.

3. **Evidence to Decision** – Upon completion of the evidence summaries by the ERE, several *en banc* meetings with the multidisciplinary Consensus Panel were conducted to present and discuss the evidence summaries and draft recommendations. The consensus panel voted on the strength and direction of the recommendations.
4. **Living CPG Process** – From the base CPG created in the above standard guideline development process, some of the questions were prioritized to a *living* status and updated depending on the following: (1) current priority for decision making, or (2) new evidence available likely to change existing recommendation [8]. The EREs working on living recommendations performed continued surveillance of literature to update the living systematic review with new evidence and updated the Evidence Summary tables and draft recommendations for the panel discussion. The Steering Committee reviewed the updated evidence summary document and revised draft recommendation. As needed, the Consensus Panel concerned was convened again in an online meeting to discuss the new evidence and any changes in the living recommendation.

The Living CPG Development Process is summarized in Figure 1 below:




Figure 1. Philippine COVID-19 Living CPG Development Process.

Guideline Preparation

COMPOSITION OF THE GUIDELINE TASK FORCE

The Steering Committee were composed of members representing one or more of the following expertise: CPG methodology, clinical epidemiology, family medicine, internal medicine, infectious diseases, pulmonology and critical care, infection control, and public health. Aside from clinicians, there was also a representative from the DOH. All members have technical knowledge and expertise on clinical management and policy development related to COVID-19.

The Evidence Review Experts (ERE) were composed of members with one or more of the following expertise: methodologists, clinical epidemiologists, evidence-based medical practitioners. They preferably had previous training and experience in CPG development and evidence synthesis.

The Consensus Panel was composed of multi-sectoral representatives such as health practitioners, both specialists and non-specialists, and patient advocates. Aside from clinicians, there were also representatives from the DOH. All panel members were the designated representatives of the relevant professional groups and stakeholder organizations and were selected based on their content expertise and experience, and potential conflicts of interest. The panelists, being involved directly in COVID-19 patient care and some having been infected themselves, acted also as patient advocates to reflect patients' and public's views and preferences.

Refer to Appendix A for the full composition of the Philippine COVID-19 Living CPG Task Force, including their professional and institutional affiliations. Their declarations of conflicts of interest are presented in Appendix B.

KEY CLINICAL ISSUES AND QUESTIONS

The Philippine COVID-19 Living CPG tackled six central themes in COVID-19, and each theme was represented by a task force headed by a technical coordinator:

- Screening and diagnosis
- Treatment
- Critical care and respiratory management
- Non-pharmacologic interventions
- Vaccines and prophylactic interventions
- Adjunct interventions

Table 1 below summarizes the topics covered per panel. The Steering Committee, together with the TWG and other key stakeholders, finalized the health questions to be addressed in the CPG. The detailed population, interventions/ tests, and outcomes were presented in the appropriate sections for each theme.

Table 1. Topics covered in the Philippine COVID-19 Living CPG.

Screening and Diagnosis	Treatment
<ul style="list-style-type: none"> • 14-day COVID-19 symptom-based test • Pulse oximeter monitoring • Choice of sample for RT-PCR (including saliva testing) • Rapid antigen tests • Self administered rapid antigen tests • Breath test • Pooled sample testing using RT-PCR • Repeat testing using RT-PCR • Antibody testing for reinfection • Antibody tests (LFIA, ECLIA, and ELISA) for seroprevalence • Clinical risk assessment and RT-PCR testing for patients undergoing surgery • Return to work after COVID-19 (non HCW and HCWs) • Thoracic Imaging in the diagnosis of COVID-19 (Chest X-ray, Chest CT Scan and Lung Ultrasound) • Prognostic factors of severe disease and mortality • LDH, CRP, and Ferritin for immunotherapy • D-dimer for anticoagulation • Heparin Induced Thrombocytopenia (HIT) Test kits • Risk Factors for Long COVID • Serum Trypsin 	<ul style="list-style-type: none"> • Hydroxychloroquine (HCQ) or Chloroquine (CQ) with or without Azithromycin • Azithromycin • Favipiravir • Remdesivir • Molnupiravir • Baloxavir • Oseltamivir • Lopinavir/ritonavir • Tocilizumab • Baricitinib • Imatinib • Tofacitinib • Leronlimab • Infliximab • Bevacizumab • Ivermectin • Artesunate • Colchicine • Interferon • Fluvoxamine • Bamlanivimab-etesevimab • Casirivimab-imdevimab • Regdanvimab • Convalescent Plasma • Intravenous immunoglobulin • Mesenchymal stem cell therapy • Inhalational steroids • Steam inhalation • Virgin coconut oil • Traditional Chinese Medicine (Lianhua capsules) • Famotidine • Ibuprofen
Critical Care and Respiratory Management	Non-Pharmacologic Interventions
<ul style="list-style-type: none"> • Systemic corticosteroids • Anticoagulation • Empiric antimicrobials • Hemoperfusion • Fluid management • Proning and side-lying in non-intubated COVID-19 patients • High-Flow Nasal Cannula (versus Non-invasive ventilation or Conventional oxygen therapy) for acute hypoxemic respiratory failure • Non-invasive ventilation 	<ul style="list-style-type: none"> • Cloth masks • Face masks and protective eyewear (face shield or goggles) • Copper mask • Ionizing air filters • Foot bath • Misting tents • UV lamps • HEPA filter • Carbon dioxide monitors • Methods of decontaminating N95 mask for reuse

- | | |
|--|--|
| <ul style="list-style-type: none"> • Lung protective ventilation, PEEP, and driving pressure for ARDS patients • Rapid sequence intubation versus delayed intubation • ECMO • Hyperbaric Oxygen Therapy • Sedation, neuromuscular blockade and paralysis • Inhaled Nitric Oxide • Etoposide for cytokine storm • Pulmonary rehabilitation for patients with long COVID-19 • Pirfenidone vs. Nintedanib for Long COVID | <ul style="list-style-type: none"> • Minimum PPE during surgeries • PPEs for HCWs in OPDs in areas with sustained community transmission • PPEs for HCWs in the hospital setting (wards, ICUs, and emergency rooms) • Acrylic physical barriers • Methods of surface disinfection |
|--|--|

Vaccines and Prophylactic Interventions

- COVID-19 Vaccines
- Vaccination for children
- Vaccination for Pregnant and Lactating Women
- Booster Dosing
- Heterologous vaccination
- Vaccines against the Delta variant
- Hydroxychloroquine (HCQ) or Chloroquine (CQ)
- Lopinavir/ Ritonavir
- Melatonin
- Nasal saline irrigation
- Antiseptic mouthwash/ gargle
- Steam inhalation
- Vitamin D
- Zinc supplements
- BCG vaccine
- Ivermectin
- Aspirin as prophylaxis against COVID-19 induced coagulopathy
- Casirivimab-imdevimab

Adjunct Interventions

- Zinc
- Vitamin B
- Vitamin C
- Vitamin D
- Melatonin
- Virgin Coconut Oil
- Lagundi
- Tawa-tawa
- Oral fatty acid supplements
- N-Acetylcysteine
- RAAS antagonists as maintenance therapy
- Statins
- Ibuprofen and worse COVID-19 symptoms
- Aspirin as maintenance therapy
- Antiseptic mouthwash or gargle
- Nasal sprays

Evidence Synthesis

The general approach for the evidence reviews for this CPG was the identification of existing systematic reviews and CPGs on COVID-19. Reference lists were checked vis-a-vis the search yield of the evidence reviewers. If there were none found, or the systematic reviews and CPGs were not high-quality nor updated, a *de novo* systematic review was done. Otherwise, high-quality and up-to-date review CPG evidence summaries were used for generating recommendations.

Each clinical question was reviewed by at least two reviewers, with the oversight of an expert technical coordinator. This was done to ensure reproducibility of the following study assessments: Inclusion/ exclusion of studies, study quality appraisal, and data extraction.

SEARCH METHODS

Primary studies and systematic reviews were searched from December 2020 to December 2021, using the following sources:

- Electronic databases: MEDLINE through PubMed and Cochrane CENTRAL Database
- Pre-print databases: ChinaXiv.org, MedRxiv.org, and BioRxiv.org
- Trial registries: USA ClinicalTrials.gov, China ChiCtr.org, and WHO ICTRP
- Living COVID-19 databases: COVID-19 Open Living Evidence Synthesis (<https://covid-nma.com/>), COAP Living Evidence on COVID-19 (https://zika.ispm.unibe.ch/assets/data/pub/search_beta/), and L-OVE Database (<https://iloveevidence.com>)
- COVID-19 Living CPGs: Australia (<https://covid19evidence.net.au/>), US NIH (<https://www.covid19treatmentguidelines.nih.gov/>), and WHO (<https://www.who.int/publications/i/item/therapeutics-and-covid-19-living-guideline>)

A final check of the comprehensiveness and completeness of the search was done by checking references used in relevant articles on the UpToDate Clinical Decision Support System (<http://uptodate.com/>).

Detailed search strategies for each clinical question were presented in the respective full-text evidence summaries. Refer to Appendix C for the search terms used for COVID-19 and the study design filters.

INCLUSION AND EXCLUSION CRITERIA

As a rule, questions on clinical efficacy and safety of interventions were answered using randomized controlled trials. If there were limited or no RCTs available, observational studies were included. For questions on diagnostic tests, appropriately designed diagnostic accuracy studies were sought.

The target population depended on the clinical question, whether it was on patients with COVID-19, individuals at high risk of COVID-19, or the general population. Due to the limited resources available, only those articles in the English language were

included. Specific details on inclusion and exclusion criteria were presented in the respective full-text evidence summaries.

STUDY QUALITY ASSESSMENT

Quality appraisal of primary studies and systematic reviews was done by at least two independent reviewers. The Painless EBM questions on validity [8] were prescribed to be used for quality appraisal of therapy, diagnosis, harm, and systematic review questions. Risk of bias assessments were summarized in evidence tables within the respective full-text evidence summaries.

Certainty of evidence for each outcome was determined using the GRADE approach [5]. The overall certainty of evidence was determined by the ERE by considering the lowest certainty across all critical and important outcomes. There were different factors considered by the reviewers in determining the certainty of evidence, as summarized in Table 2.

DATA SYNTHESIS

Meta-analysis was done to pool the treatment effects or the diagnostic performance indices, as appropriate to the clinical question. When studies and results cannot be combined, a narrative synthesis was done, and relevant information was summarized in a table.

Table 2. Factors influencing certainty of evidence [6].

Certainty of Evidence	Study Design – Intervention Questions	Study Design – Diagnosis Questions	Factors that Decrease COE (by 1 to 2 levels)	Factors that Increase COE (by 1 to 2 levels)
High	Randomized controlled trial	Appropriate cross-sectional or cohort studies in patients with diagnostic uncertainty	<ul style="list-style-type: none"> • Risk of Bias • Inconsistency • Indirectness • Imprecision • Publication Bias 	<ul style="list-style-type: none"> • Large magnitude of effect • Plausible confounding • Dose-response gradient
Moderate				
Low	Observational study			
Very Low				

Evidence to Decision: Formulating Recommendations

The Consensus Panel evaluated the direction and strength of recommendation using the GRADE approach and the Evidence to Decision Framework, based on the (1) overall quality of evidence for each question, (2) balance between benefits and harms, (3) values, preferences, and burden on patients, (4) cost and resource use, and (5) other considerations such as feasibility, equity and acceptability.

CERTAINTY OF EVIDENCE AND STRENGTH OF RECOMMENDATIONS

The certainty of evidence was one of the bases of the Consensus Panel in making the final recommendation. Table 3 shows the definition and implication of each:

Table 3. Definitions and Implications of each GRADE Certainty of Evidence [6].

GRADE Certainty of Evidence	Definition	Implication
High	We are very confident that the true effect lies close to that of the estimate of the effect.	Further research is very unlikely to change confidence in the estimate of effect
Moderate	We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different	Further research is likely to have an important impact on confidence in the estimate of effect and may change the estimate
Low	Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect.	Further research is very likely to have an important impact on confidence in the estimate of effect and is likely to change the estimate
Very Low	We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect	Any estimate of effect is very uncertain

The strength of recommendation could either be strong or weak. However, there were three reasons where the consensus panels were unable to make a recommendation [6]:

1. Confidence in effect estimates is so low that the panels feel a recommendation is too speculative.
2. Trade-offs are so closely balanced, and the values and preferences, and resource implications are not known or too variable.
3. Management options have very different undesirable consequences, and individual patients' reactions to these consequences are likely to be variable

A strong recommendation was stated as "We recommend/ We recommend against...", while a weak recommendation was worded "We suggest/ We suggest against..."

Finally, when no recommendation can be made, the sentence starts with “There is no/insufficient evidence to recommend...”

The implications of strong and conditional recommendations are enumerated in Table 4 [5].

PATIENT VIEWS AND PREFERENCES

Patient views and preferences were represented by the nurses who had direct patient care encounters and consensus panel members who were directly involved in various aspects of COVID-19 care: clinician, administrator, researcher. Some of the panelists were COVID-19 patients themselves or had relatives and friends afflicted with COVID-19. This strategy ensured that patient views and preferences are still considered in the formulation of recommendations.

Table 4. Implications of the Strength of Recommendation to Patients, Clinicians, and Policymakers [6].

	Strong Recommendation	Weak Recommendation
Patients	Most individuals in this situation would want the recommended course of action and only a small proportion would not.	Most individuals in this situation would want the suggested course of action, but many would not.
	Most individuals should receive the recommended course of action.	Recognize that different choices will be appropriate for different patients.
Clinicians	Adherence to this recommendation according to the guideline could be used as a quality criterion or performance indicator.	Clinicians must help each patient arrive at a management decision consistent with her or his values and preferences.
Policy makers	The recommendation can be adopted as policy in most situations including for the use as performance indicators.	Policy making will require substantial debates and the involvement of many stakeholders. Policies are also more likely to vary between regions.

RESOURCE IMPLICATIONS

Since COVID-19 is a relatively new disease that is being studied internationally, and most COVID-19 diagnostics and interventions are still investigational, there were limited economic evaluations available. In the absence of this information, consensus panelists considered the cost and other local resources needed for the recommendations. This discussion could be found in the *Consensus issues* subsection of each evidence summary, when appropriate.

RATING OF OUTCOMES

The Consensus Panel rated outcomes for each set of clinical questions according to whether they were critical, important but not critical, or of low importance for decision making. Critical outcomes were primary factors that should influence a recommendation, while those with lower importance did not bear on these recommendations. On a scale of 1-9, those rated 7-9 were critical outcomes, 4-6 were important but not critical outcomes, and 1-3 were outcomes of limited importance. Table 5 below shows the result of the ranking of outcomes per CPG panel:

Table 5. Outcome ratings for each CPG panel.

CPG Panel	Critical Outcomes	Important but not critical outcomes
Screening and Diagnosis	<ul style="list-style-type: none"> • Sensitivity and specificity • Positive and negative predictive values • Development of COVID-19 • Mortality • Positive and negative likelihood ratios • False positive and false negative rates • Resource savings • Number of RT-PCR positive samples 	<ul style="list-style-type: none"> • Psychological effects of testing • Physical harm of testing
Treatment – Outpatients and Mild to Moderate COVID Patients	<ul style="list-style-type: none"> • All-cause mortality • Clinical improvement/ Time to clinical cure • Respiratory distress/ Need for mechanical ventilation or oxygen support therapy • Progression to severe COVID-19 • Need for hospitalization or ICU admission (including duration) • Adverse events • Serious adverse events 	<ul style="list-style-type: none"> • Duration of mechanical ventilation • Improvement in Chest x-ray or CT scan • Viral negative conversion/ viral clearance • Time to negative conversion (PCR)
Treatment – Severe to Critical COVID Patients	<ul style="list-style-type: none"> • All-cause mortality • Clinical improvement/ Time to clinical cure • Respiratory distress/ Need for mechanical ventilation • Hospitalization • ICU admission • Need for supportive oxygen therapy • Adverse events • Serious adverse events 	<ul style="list-style-type: none"> • Duration of mechanical ventilation • Duration of hospitalization • Viral negative conversion/ Viral clearance • Improvement in Chest X-ray/ CT Scan • Time to negative conversion (PCR)
Critical Care and Respiratory Management	<ul style="list-style-type: none"> • COVID-19 related deaths • Need for mechanical ventilation • Clinical improvement • Length of hospital and ICU stay • Thromboembolic events • Bleeding • Serious adverse events 	<ul style="list-style-type: none"> • Duration of Mechanical ventilation • Adverse events • Improvement in oxygenation • Length of hospital and ICU stay
Non-Pharmacologic Interventions	<ul style="list-style-type: none"> • Incidence of COVID-19 infection (any severity) • Filtration efficacy • Prevention of influenza-like illness • Safety of use 	
Vaccines and Prophylactic Interventions	<ul style="list-style-type: none"> • COVID-19-related deaths • Incidence of COVID-19 • Incidence of hospitalization or ER visit • Incidence of mechanical ventilation • Adverse events • Serious adverse events 	

Vaccines and Prophylactic Interventions - Pregnant and Lactating Women	<ul style="list-style-type: none"> • Newborn antibody levels • Pregnancy outcomes: <ul style="list-style-type: none"> ○ Gestational Hypertension ○ Gestational Diabetes ○ Premature Rupture of membranes ○ IUGR • Delivery Outcomes <ul style="list-style-type: none"> ○ Spontaneous Abortion ○ Fetal death in utero or still birth ○ Preterm delivery ○ APGAR Score <7 • Neonatal outcomes <ul style="list-style-type: none"> ○ NICU Admission ○ Presence of congenital anomalies • Breastmilk antibodies • Changes in breastmilk 	<ul style="list-style-type: none"> • Fetal, placental or cord blood antibody levels
Adjunct Interventions	<ul style="list-style-type: none"> • Clinical recovery/improvement • All-cause mortality • Need for/ Duration of mechanical ventilation • Need for hospitalization • Adverse events • Serious adverse events • Clinical deterioration/ progression of respiratory distress/progression to severe COVID-19 • Need for ICU admission • Need for supportive oxygen therapy/mechanical ventilation 	<ul style="list-style-type: none"> • Time to negative conversion • Duration of hospital stay • Improvement in Chest x-ray/CT Scan • Duration of mechanical ventilation

CONSENSUS PROCESS

A skilled facilitator moderated the discussions during the consensus meetings. Each member voted on the draft recommendation as follows: yes, no, or abstain. The consensus was defined as at least 75% agreement among the members for both the direction and strength of recommendation. If consensus was not reached, members discussed the reasons in support of their votes for or against the recommendation. The voting was repeated, up to three rounds, until a consensus was reached. Any issues left unsettled after the *en banc* meeting were finalized through a modified Delphi activity.

There were two recommendations that did not reach a consensus after voting for three rounds and a repeat meeting after a modified Delphi activity. These were on the recommendations for heterologous booster vaccination (October 2021) and the use of nasal sprays as adjunctive therapy (December 2021). The recommendation on heterologous vaccination has been resolved as of December 27, 2021 following an update on the evidence available.

Guideline Dissemination

Three methods were used in the dissemination of the Philippine COVID-19 Living CPG: (1) online webpage, (2) Living Recommendations document, and (3) full-text CPG manuscript.

The online webpage of the Philippine COVID-19 Living CPG (Figures 2 and 3) was hosted on the PSMID website. This was launched on March 21, 2021, and has undergone improvements from the feedback of CPG users and members of the Living CPG task force.

The short *Living Recommendations document* (Figure 4) contained the content in the PSMID website, including the introduction, CPG methodology, members of the living CPG task force, and the actual recommendation statements. The evidence summaries were not included in this document. This shorter format allowed for an easily accessible document for use by practitioners and selected laypersons.

This full-text CPG manuscript, as well as the complete evidence base, will be submitted to the DOH National Clearinghouse for national promotion regarding use and uptake of the recommendations, including activities such as releasing a department memorandum to notify stakeholders, publicizing the CPG through the DOH newsletter and to other appropriate agencies, and issuing press releases, news articles, and social media posts. The final manuscript will be made available as electronic copies through the websites of DOH and PSMID.

Furthermore, several dissemination fora have already been conducted during relevant meetings of professional societies, where several members of the Steering Committee and Consensus Panels presented. More avenues for dissemination will be undertaken to promote the use and value of this CPG's recommendations.

Real-time updates of living recommendations were published on the CPG webpage and disseminated to various stakeholders. Further updates will be announced during the DOH daily updates on COVID-19, promoted on various social media platforms, and published on the PSMID website

Figure 2. Initial PSMID Webpage for the Philippine COVID-19 LCPG in March 2021.





Figure 3. Latest PSMID Webpage for the Philippine COVID-19 LCPG.



Philippine COVID-19 Living Recommendations
Updated May 28, 2021

Home About Disclaimer Contact Us Download

The Philippine COVID-19 Living Recommendations document is brought to you by the Institute of Clinical Epidemiology, National Institutes of Health, UP Manila in cooperation with the Philippine Society of Microbiology and Infectious Diseases (PSMID). This was funded by the DOH Ahead Program through the DOST-Philippine Council for Health Research and Development (PCHRD).

INTRODUCTION


Given the magnitude of the impact of COVID-19 in the for clinical practice guidelines to optimize health care the rapid pace of scientific developments on COVID-19 providers to guide their diagnosis and treatment decisions inform policy and provide timely guidance on effective

While there are existing international guidelines and by local experts, end-users and other relevant stakeholders switched to a living status based on the likelihood of n

Screening and Diagnosis

- Should the 14-day symptom-based test be used in screening for COVID-19 infection in apparently healthy adults?
- Which clinical specimens can be used as an alternative to nasopharyngeal swab RT-PCR* for the diagnosis of COVID-19?
- Should rapid antigen tests be used in the diagnosis of COVID-19 in clinically suspected patients?
- Should pooled testing using RT-PCR for SARS-CoV-2, versus individual testing using RT-PCR, be used for screening and surveillance for SARS-CoV-2 in patients with suspected COVID-19 infection?
- Should repeat RT-PCR testing after an initial negative RT-PCR versus single RT-PCR testing be done to diagnose COVID-19 in symptomatic patients with high index of suspicion?
- Among asymptomatic individuals scheduled for non-urgent, non-emergency surgery, should RT-PCR and clinical risk assessment vs clinical risk assessment alone be done to screen for COVID-19?
- Should antibody tests be used for COVID-19 seroprevalence studies among adult populations?

Figure 4. Living Recommendations Document for the Philippine COVID-19 Living CPG.



Philippine COVID-19 Living Recommendations
As of 04 June 2021

By:
Institute of Clinical Epidemiology,
National Institutes of Health, UP Manila

In cooperation with:
Philippine Society for Microbiology
and Infectious Diseases

Funded by:
DOH AHEAD Program
through the DOST-PCHRD

IMPORTANT NOTICE: These living recommendations will be updated as new evidence are published in the medical literature. It is critical that you take note of the date when the evidence was last reviewed. Additional recommendations may have to be developed as needed.

1 | Philippine COVID-19 Living Recommendations As of 04 June 2021

Guideline Monitoring and Evaluation

Guideline implementation would be assessed through process and impact evaluation. Only a process evaluation was feasible during the project implementation using webpage analytics. Refer to the subsection on *Process Evaluation* in the *Discussion* section of this manuscript.

Impact evaluation for the Philippine COVID-19 Living CPG would include bi-annual surveys of the following (1) clinicians managing COVID-19 patients, (2) public health practitioners coordinating local PDITR+ strategies in the community, and (3) the public regarding their compliance to non-pharmacologic interventions and any preventive measures.


The quality of care rendered to COVID-19 patients can be assessed by measuring adherence of healthcare providers and institutions to the recommendations of the Philippine COVID-19 Living CPG. Strong recommendations would be included in a quality-of-care checklist on COVID-19 care, while weak recommendations would be relevant if the identified conditions are satisfied.

Finally, a scheduled bi-annual review would be conducted to evaluate the process efficiency and scientific quality of the Philippine COVID-19 Living CPG.

External Review

The CPG webpage served the dual purpose of a dissemination method and a way to collect the external reviews of the CPG processes, evidence summaries, and recommendations. This website (Figure 5) also allowed health professionals and key stakeholders to suggest additional clinical questions that could be included in the scope of this CPG.

Figure 5. Contact details in the Webpage for the Philippine COVID-19 Living CPG.



Over the weeks and months, we have gathered feedback from users and members of the Living CPG Taskforce to improve the readability of the webpage, such as toggling of topics, recommendations, and evidence summaries, changing from topics to questions in the listing, rearranging various sections into headers (such as CPG methodology, task force members, contact details, etc.), and other formatting changes. As an illustration, one feedback received on Lianhua led to the update of that evidence summary, and the inclusion of the health professional who gave the feedback in the evidence review team on Lianhua.

Furthermore, the *en banc* project review meeting with the Philippine Council for Health Research and Development and the screening appraisal of the Department of Health-Disease Prevention and Control Bureau will provide important inputs on the improvement of the living CPG development.

Updating of Guidelines

Due to the rapidly evolving science of COVID-19 treatment and diagnosis, the Philippine COVID-19 Living CPG was updated continuously. See the *Living CPG Process* under the *Overview of Philippine COVID-19 Living CPG Development Process* section for specific details on the process of updating this Living CPG.

After the initial DOH-PCHRD funding for six months, the DOH Disease Prevention and Control Bureau has provided funding support for another two and a half months to continue the surveillance search for the “living recommendations”. Further funding will be sought from professional societies and other government agencies to ensure the sustainability of the living CPG throughout the COVID-19 pandemic.

Editorial Independence

FUNDING SOURCE

This CPG project was funded by the Department of Health AHEAD Program through the Philippine Council for Health Research and Development for the first 6 months and subsequently funded by the DOH Disease Prevention and Control Bureau during the last quarter of 2021. Though the DOH was part of the Steering Committee and the Consensus Panels, their influence on the guideline content was limited to the identification of key clinical questions and the discussion of the recommendations. The funding agency did not have any undue influence on the evidence review conducted, as well as on the interpretation of the research data available.

MANAGEMENT OF CONFLICTS OF INTEREST

All members involved in the creation of this Clinical Practice Guideline, including the Steering Committee, Technical Working Group, and Consensus Panel, declared any potential conflicts of interest within the last 4 years, using a uniform Declaration of Conflict of Interest (DCOI) form as recommended in the DOH Manual [5]. These were reviewed by an independent Oversight Committee (OC) and the Steering Committee, to screen and manage the COIs declared. The Oversight Committee is responsible for recommending the extent of participation that can be allowed. The decisions of the

Oversight Committee will be reported and published in the final manuscript of the Living CPG.

The Oversight Committee has come up with the following guide as bases for their decisions:

- a. **Manageable A** - if there are intellectual conflicts of interest only. They can vote but they need to declare their intellectual conflicts (e.g., affiliation with institutions, positions in an organization, authorship in paper or CPG)
- b. **Manageable B** - if there are some intellectual and financial conflicts of interest. They cannot vote but they can share their expertise with the group. Examples include panelists from government agencies directly involved in the implementation of the program and panelists from the agency funding the guidelines. The specific terms of management shall be set forth by the OC and shall relate to specific clinical questions.
- c. **Disqualifying** - the scope or nature of some relationships negates management and will sometimes outweigh the content expertise an individual may bring by serving as a full panelist, resulting in disqualification.

Evidence and Recommendations for the Screening and Diagnosis of COVID-19

Should the 14-day symptom-based test be used in screening for COVID-19 infection in apparently healthy adults and children?

As of 29 November 2021

RECOMMENDATION

We suggest doing an initial screening for ANY influenza-like illness, typical and atypical COVID-19 symptoms* within the past 14 days in apparently healthy adults and children, especially for individuals with known exposure to a laboratory-confirmed case of COVID-19. (*Very low certainty of evidence; Weak recommendation*)

**Symptoms include but not limited to: fever/chills, cough, shortness of breath/dyspnea, sore throat, runny nose, myalgia, headache, fatigue/malaise, diarrhea, nausea/vomiting, abdominal pain, anosmia, ageusia, wheezing, chest pain, altered mental status, seizures, rash, pink eye*

KEY FINDINGS

- Evidence on the accuracy of 14-day symptom-based testing was found from three observational studies (n=8,475).[1-3] This symptom-based test had a wide range of sensitivity (2.2-100%) and specificity (29-99%) in detecting COVID-19 among adults and children. Variability in accuracy appeared to be associated with heterogeneity in the populations tested and the characteristics of the index test.
- The diagnostic accuracy of individual symptoms differed depending on the exposure history of the populations where the test was applied. For those with unclear exposure, individuals had excellent specificity (>90%) but poor sensitivity (<50%). For close contacts of confirmed COVID-19 cases, nasal congestion/rhinorrhea, headache, cough, sore throat, were the individual symptoms that had at least 70% sensitivity. For children, accuracy was highest for headache, nasal congestion or rhinorrhea, fever or chills, and sore throat.
- Combining individual symptoms can increase the specificity to as high as 99.9% but with no substantial improvements in sensitivity. The Centers for Disease Control (CDC) symptom list showed the highest sensitivity (100%) but lowest specificity (21-45%) in both adults and children population while influenza-like illness showed the highest specificity (86-96%) but lowest sensitivity (43-54%).

CONSENSUS ISSUES

A weak recommendation was made based on evidence including studies that were conducted prior to the new variant of concern, Omicron, which was noted to present with symptoms not typical of previous variants. Additionally, majority of the studies were on adults while one study was on the pediatric population. The panelists emphasized that the list of symptoms is not exhaustive and that presence of any of the symptoms would warrant further investigation through a follow-up confirmatory diagnostic test.

Should pulse oximetry be used for at-home monitoring of COVID-19 patients?

As of 22 November 2021

RECOMMENDATION

We suggest pulse oximetry with close clinical monitoring by qualified medical personnel in suspected and confirmed COVID-19 patients especially those who are at high risk for deterioration. (*Very low certainty of evidence; Weak recommendation*)

KEY FINDINGS

- A total of 20 observational studies [4-23] on pulse oximetry monitoring for suspected and confirmed COVID-19 patients were included in this review.
- Effect estimates could not be pooled due to serious heterogeneity across studies. Lowered percentages of admissions, readmissions, and mortalities with pulse oximetry monitoring compared to no monitoring suggest feasibility and safety of remote monitoring using pulse oximetry but the certainty of evidence was very low.
- A single study [22] on device accuracy showed that medical and consumer-grade pulse oximeters, particularly Oxywatch (Sn 92.2%, 95% CI 87.3-97.1; Sp 60%, 95% CI 49.59-70.41), SM (Sn 90.7%, 95% CI 85.7-95.7; Sp 67.6%, 95% CI 56.7-78.5), and Onyx (Sn 92.1%, 95% CI 87.3-96.8; Sp 67.6%, 95% CI 56.9-78.2), were comparable to standard emergency department (ED) monitor units.
- Another study [23] reported that certain levels of oxygen saturation (SpO₂) have high sensitivity and specificity for risk of mortality and intensive care unit (ICU) admission. However, the overall certainty of evidence for this outcome was very low because of non-blinding, imprecision, and significant heterogeneity.

CONSENSUS ISSUES

A weak recommendation was made due to the limitations and risk of inaccuracy of pulse oximeters. The panel emphasized (1) ensuring the quality of the device by purchasing from reliable sources and (2) taking measures to get the best reading (e.g., following manufacturer instructions and ensuring adequate battery supply).

Considering the risk of inaccurate measurements that may result in unrecognized low oxygen saturation levels, the role of qualified medical personnel in pulse oximetry monitoring was highlighted by the panelists. Medical personnel should be available to provide instructions, to respond to caregiver and/or patient concerns, and to monitor signs and symptoms of deterioration

Which clinical specimens can be used as an alternative to nasopharyngeal swab RT-PCR* for the diagnosis of COVID-19?

As of 20 February 2021

RECOMMENDATION

We recommend the use of the following specimens as alternative specimens to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19 among symptomatic and asymptomatic patients suspected of COVID-19 in hospital and outpatient settings:

- Oropharyngeal swab (*Moderate certainty of evidence; Strong recommendation*)
- Saliva drool/spit and oral saliva (*Moderate certainty of evidence; Strong recommendation*)
- Nasal swab/wash (*Moderate certainty of evidence; Strong recommendation*)
- Throat swab (*Low certainty of evidence; Strong recommendation*)

We suggest the use of saliva swab and posterior oropharyngeal saliva specimens as an alternative specimen to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19 among symptomatic and asymptomatic patients with suspected COVID-19 in hospital and community/outpatient settings. (*Low certainty of evidence; Weak recommendation*)

RECOMMENDATION

We recommend against the use of sputum as an alternative specimen to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19. (*Very low certainty of evidence; Strong recommendation*)

RECOMMENDATION

There is no evidence to recommend the use of bronchoalveolar lavage as an alternative specimen to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19.

*SARS COV-2 RT-PCR of nasopharyngeal swabs remains the diagnostic test of choice to confirm the diagnosis of COVID-19 among suspected individuals.

KEY FINDINGS

- One cross-sectional study [24] on the use of oropharyngeal swab RT-PCR as an alternative clinical specimen to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19 showed that oropharyngeal swab had comparable sensitivity and specificity to nasopharyngeal swab RT-PCR.
- A meta-analysis [25] of 19 observational studies on the use of saliva as an alternative clinical specimen to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19 concluded that saliva had comparable sensitivity and specificity to nasopharyngeal swab RT-PCR.
- Two cross-sectional studies [24,26] on the use of nasal swab/wash RT-PCR as an alternative specimen to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19 showed that nasal swab/wash had comparable sensitivity and specificity to nasopharyngeal swab RT-PCR.

- A cross-sectional study [26] on the use of throat swab RT-PCR as an alternative specimen to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19 also showed that throat swab had comparable sensitivity and specificity to nasopharyngeal swab RT-PCR.
- A cross-sectional study from a meta-analysis [25] on the use of sputum RT-PCR as an alternative specimen to nasopharyngeal swab RT-PCR for the diagnosis of COVID-19 showed that sputum had lower sensitivity and specificity compared to nasopharyngeal swab RT-PCR.
- All the above studies were assessed to have low risk of bias.
- No studies compared the sensitivity and specificity of bronchoalveolar lavage RT-PCR to nasopharyngeal swab RT-PCR. Hence, no conclusion and recommendation can be made for this clinical specimen.

CONSENSUS ISSUES

Currently, oropharyngeal, and nasopharyngeal specimens are collected simultaneously; however, in resource-limited settings, the panel recognized the need for alternative clinical specimens to nasopharyngeal swab RT-PCR, along with the positive implication of single specimens on resource use. As a result, the use of oropharyngeal swab, oral saliva specimens, nasal swab/wash and throat swab were recommended as alternative clinical specimens to nasopharyngeal swab RT-PCR.

The differences between oropharyngeal swab and throat swab samples were clarified. Although the two specimens are collected in the same area, they were considered by the panel as dissimilar specimens due to the differences in sample collection technique. The panel made a strong recommendation for using throat swab samples due to its relatively high sensitivity.

The panel opted to strongly recommend against the use of sputum specimens as an alternative to nasopharyngeal swab samples. This was primarily due to the risk of viral transmission when obtaining sputum samples, coupled with the very low certainty of the evidence for its use.

Among patients suspected to have COVID-19, how accurate are rapid antigen tests compared to RT-PCR for the diagnosis of COVID-19?

As of 22 November 2021

RECOMMENDATION

We suggest the use of rapid antigen test for the diagnosis of symptomatic individuals suspected of COVID-19 as an alternative to RT-PCR if all the following conditions are met: *(Low certainty of evidence; Weak recommendation)*

- a. Individuals are in the early phase of illness (less than or equal to 7 days from onset of symptoms); AND
- b. Testing kits demonstrated sensitivity of more than or equal to 80% AND have very high specificity of more than or equal to 97%.

We suggest the use of rapid antigen tests for the diagnosis of individuals suspected of COVID-19 during the setting of an outbreak provided that all the following conditions are met: *(Very low certainty of evidence; Weak recommendation)*

- a. Individuals are in the early phase of illness (less than or equal to 7 days from onset of symptoms); AND
- b. Testing kits demonstrated sensitivity of more than or equal to 80% AND have very high specificity of more than or equal to 97%.

RECOMMENDATION

We suggest against the use of rapid antigen test for screening purposes. *(Low certainty of evidence; Weak recommendation)*

We suggest against the use of saliva as specimen for rapid antigen test in patients suspected of COVID-19 infection. *(Low certainty of evidence; Weak recommendation)*

We suggest against the use of rapid antigen tests alone in asymptomatic patients suspected of COVID-19 infection. *(Low certainty of evidence; Weak recommendation)*

RECOMMENDATION

There is insufficient evidence to recommend for or against the use of repeat antigen testing for screening or diagnosis of COVID-19. *(Very low certainty of evidence)*

A negative rapid antigen test should be confirmed with an RT-PCR in settings or situations wherein COVID-19 is highly suspected (e.g., symptomatic or asymptomatic close contacts of probable or confirmed COVID-19 individuals).

KEY FINDINGS

- A total of 164 observational studies [27-190] assessed the diagnostic accuracy of rapid antigen tests (RAgTs) against reverse transcriptase-polymerase chain reaction (RT-PCR) as the reference standard. Studies included different test brands, specimen types and timing of collection, symptom status, cycle threshold

(CT) values, and populations, namely inpatients, children, and healthcare workers among others.

- The overall sensitivity of RAgTs was moderate at 0.71 (95% CI 0.68-0.73) while specificity was excellent at 0.995 (95% CI 0.993-0.996). This was comparable to the data of the previous evidence summary with a pooled sensitivity of 0.72 (95% CI 0.64-0.78) and specificity of 0.99 (95% CI 0.99-1.0).
- On subgroup analysis, RAgT showed higher sensitivity when used in symptomatic individuals (Sn 0.74, 95% CI 0.71-0.78), when conducted during the early phase or first week of illness (Sn 0.79, 95% CI 0.75-0.82), in positive specimens with Ct value <25 (Sn 0.94, 95% CI 0.92-0.96), and in other Ct thresholds considered as “high” viral load (Sn 0.89, 95% CI 0.85-0.92). Pooled sensitivity of commonly used specimen types falls between 65% to 79%. FDA-approved RAgT brands have pooled sensitivities ranging from 0-90% with improved performance of commonly used RAgT brands when used in symptomatic individuals.
- In outbreak settings, RAgT use remained to have excellent specificity (Sp 0.966, 95% CI 0.997-0.999) with similar sensitivity (Sn 0.68, 95% CI 0.45-0.84) but with less precise estimates.
- The overall certainty of evidence was low because of serious risk of bias in all domains (high and unclear risk in patient selection, conduct of index test and reference standard, and flow and timing) and serious inconsistency. Despite performing pre-specified subgroup analyses, significant heterogeneity was still observed. In certain subgroup analyses such as the use of RAgT in outbreak settings and saliva specimens, certainty of evidence was further downgraded to very low due to imprecision attributed to wide interval estimates.

CONSENSUS ISSUES

The panel was unanimous against (1) the use of rapid antigen test for screening purposes, (2) the use of saliva as specimen for rapid antigen tests, and (3) the use of rapid antigen test alone in asymptomatic patients suspected of COVID-19 infection due to the observed lower sensitivity of these tests under such conditions. A unanimous decision on the insufficiency of evidence to recommend for or against the use of repeat antigen testing was also made.

Majority of the panelists agreed that the following conditions should be met when using rapid antigen tests:

- a. Individuals are in the early phase of illness, because antigen tests perform best during this period; and
- b. Testing kits have a sensitivity of more than or equal to 80% and specificity of more than or equal to 97%, because the quality of the test kit should be ensured.

One of eleven panelists raised a concern on the specified sensitivity and specificity of the testing kits, as these are based on the Health Technology Assessment Council (HTAC) of the local Department of Health (DOH).

A weak recommendation on the use of rapid antigen tests for diagnosing COVID-19 suspects during outbreaks was made based on nine observational studies with unclear patient selection, conduct of reference standard, and patient flow and timing. The risk of exposure was an important consideration for the panel, citing that it is not cost-effective to test everyone during an outbreak. However, the risk stratification of participants was not specified in any of the studies.

Among patients suspected to have COVID-19, how accurate are self-administered rapid antigen tests alone compared to RT-PCR for the diagnosis of COVID-19?

As of 11 November 2021

RECOMMENDATION

We suggest the use of self-administered rapid antigen test for the diagnosis of COVID-19 in symptomatic individuals, provided that ALL OF THE FOLLOWING conditions are met: (*Low certainty of evidence; Weak recommendation*)

1. Ease of collecting samples is ensured;
2. Ease of interpretation is ensured;
3. Test kits have passed flex studies; AND
4. Individuals present with symptoms for less than 7 days.

RECOMMENDATION

We suggest against the use of self-administered rapid antigen test for routine screening of COVID-19. (*Low certainty of evidence; Weak recommendation*)

KEY FINDINGS

- Seven observational studies [191-197] assessed the diagnostic accuracy of self-administered rapid antigen tests against RT-PCR as the reference standard. The studies included varied test brands (n=6), specimen types, and symptom status.
- The pooled sensitivity of self-administered rapid antigen test was moderate at 0.77 (95% CI 0.62-0.87) while the pooled specificity was high at 0.996 (95% CI 0.99-1.00). Pooled sensitivity estimates must be interpreted with caution due to the substantial heterogeneity ($I^2=94\%$) across studies.
- On subgroup analysis, self-administered rapid antigen test showed higher sensitivity when used in the following conditions:
 - Symptomatic individuals (Sn 0.81, 95% CI 0.69-0.89);
 - Specimens taken from exhaled breath (Sn 0.92, 95% CI 0.64-1.0) or nasal mid-turbinate (Sn 0.81, 95% CI 0.73-0.87);
 - Specimens with high viral loads at RT-PCR cycle threshold <25 (Sn 0.87, 95% CI 0.68-0.88);
 - Specific brands of rapid antigen test, namely Inflammacheck device (Sn 0.92, 95% CI 0.64-1.0), Drager antigen test (Sn 0.89, 95% CI 0.79-0.95), and Abbott Panbio (Sn 0.84, 95% CI 0.71-0.94); and
 - Studies with high methodological quality or low risk of bias (Sn 0.79, 95% CI 0.68-0.86).
- The overall certainty of evidence for test sensitivity was low because of serious inconsistency (high heterogeneity) and risk of bias issues (patient selection, conduct of index test, and reference standard).

CONSENSUS ISSUES

The Panel unanimously agreed that all the following four conditions should be met when using self-administered antigen tests for the diagnosis of COVID-19:

1. Ease of sample collection, because incorrect performance of self-administered tests largely affects diagnostic accuracy;

2. Ease of interpretation, because proper interpretation is important for the accurate clinical management of patients;
3. The test kits should have passed the flex studies, because the quality of the self-administered test kit should be ensured; and
4. Individuals present with symptoms for less than 7 days, because antigen tests perform best during this period; and beyond this, the use of the test is not cost-effective, thereby incurring costs without added benefit.

The panel decided on a weak recommendation based on evidence including seven observational studies where performance of self-testing by participants was supervised by trained personnel either onsite or via telehealth. None were conducted in a home setting. Additionally, the studies did not specify if the participants were close contacts of COVID-19 patients or if they have a high- or low-risk of contracting COVID-19.

Other issues raised include (1) the lack of locally FDA-approved self-administered antigen test kits, (2) differentiating antigen tests manufactured for self-administration versus those that are not (i.e., for trained personnel), (3) the method of reporting the test results, and (4) the subsequent management and/or protocols after a positive test result (e.g., contact tracing).

Should breath tests be used to diagnose COVID-19 infection?

As of 29 November 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of breath test in detecting COVID-19 infection. (*Low certainty of evidence*)

KEY FINDINGS

This review included three cross-sectional [198-200] and three prospective studies [201-203] on the use of breath tests in the diagnosis of COVID-19 infection. The overall accuracy of breath tests was high, with sensitivity of 97% (95% CI 0.90-0.99) and specificity of 85% (95% CI 0.72-0.92). However, the overall certainty of evidence was low due to significant heterogeneity. This heterogeneity may be attributed to the different mechanisms of the devices despite using the same idea of breath testing.

CONSENSUS ISSUES

Despite the addition of five new studies since the previous recommendation, insufficient evidence remains to recommend for or against breath tests. The diagnostic accuracy of breath tests cannot be ascertained due to the heterogeneity across studies. The panel also raised concerns on the availability and accessibility of the test, its cost, and ease of use.

Should pooled testing using RT-PCR for SARS-CoV-2, versus individual testing using RT-PCR, be used for screening and surveillance for SARS-CoV-2 in patients with suspected COVID-19 infection?

As of 6 March 2021

RECOMMENDATION

We suggest the use of pooled RT-PCR testing in targeted* low-risk and low-prevalence populations using a pool size of 5 in individuals suspected of COVID-19 infection. (*Moderate certainty of evidence; Weak recommendation*)

**For targeted populations refer to the list of Philippine Society of Pathologists and Department of Health*

KEY FINDINGS

- Twenty-one cross-sectional studies [204-224] (N=220,253) were found that used pooled RT-PCR testing for SARS-CoV2. Six were diagnostic accuracy studies that compared pooled testing with individual testing, and 15 were pragmatic clinical validation studies of pooled testing that did individual testing of positive pools. Studies had varying study populations, use case, index test kits, and pool sizes (5 to 16).
- Among the six diagnostic accuracy studies with a total of 5,987 participants, there was moderately high pooled sensitivity (Sn 81%, 95% CI 72-88%; $I^2=73.6\%$; moderate certainty of evidence) and high pooled specificity (Sp 99%, 95% CI 98-100%; $I^2=1.84\%$; high certainty of evidence), given the positivity rate of 2.7 to 15% within the study populations.
- The positive predictive value based on 21 studies ranged from 67 to 100%.
- Resource savings in the number of test kits used ranged from 49 to 89%.
- Identified harms of pooled testing were delayed turnaround time for positive samples and laboratory errors.
- The overall risk of bias was low in seven studies, among which were six diagnostic accuracy studies that contributed to pooled sensitivity and specificity, and low in 14 studies, mainly due to the lack of independent assessment between the index and reference tests.

CONSENSUS ISSUES

The set recommendation included positivity rate and pool size; however, there were no studies that investigated the specificity and sensitivity of pooled testing for different pool sizes across different prevalence settings. Since the data presented did not clearly define the risk or the prevalence settings, pooled testing was suggested to be used only at a specific target population despite the moderate quality of evidence.

Should repeat RT-PCR testing after an initial negative RT-PCR (versus single RT-PCR testing) be done to diagnose COVID-19 in symptomatic patients?

As of 6 March 2021

RECOMMENDATION

We suggest repeating RT-PCR testing when the initial RT-PCR test is negative in symptomatic patients with high index of suspicion for COVID-19 infection. (*Low certainty of evidence; Weak recommendation*)

KEY FINDINGS

- Two cohort studies [225-226] involving 368 patients were found on the accuracy of repeat RT-PCR testing after an initial negative test to diagnose COVID-19 in symptomatic patients with a high index of suspicion. The evidence was assigned a low-quality rating due to the serious risk of bias and serious imprecision.
- The sensitivity of repeat RT-PCR testing ranged from 0.83 (95% CI 0.75-0.90) to 0.85 (95% CI 0.62-0.97), approximately 15% higher compared to the sensitivity of a single RT-PCR test, which ranged from 0.68 (95% CI 0.58-0.76) to 0.70 (95% CI 0.46-0.88).
- The specificity of repeat testing was consistently very high (Sp 1.00, 95% CI 0.89-1.00).

CONSENSUS ISSUES

The recommendation applies only to symptomatic patients with high index of suspicion for COVID-19. Since the disease severity and the level of suspicion were not clearly defined in the studies, the level of suspicion may vary. Moreover, no specific recommendation was made regarding the specific time interval between the initial and the repeat test as well as the frequency of repeat PCR tests.

Among COVID-19 confirmed patients, should certain RT-PCR cycle threshold values be used to determine infectivity?

As of 15 December 2021

RECOMMENDATION

There is insufficient evidence to recommend an RT-PCR cycle threshold cut-off value to determine infectivity among COVID-19 confirmed patients. Interpretation of RT-PCR cycle threshold values may vary and is dependent on the PCR assay used, gene target, sample type, and timing of sample collection. (*Very low certainty of evidence*)

KEY FINDINGS

- Twelve observational studies [227-238] were included in this review on the use of cycle threshold (Ct) as a surrogate marker of infectivity as evidenced by viral isolation in culture. One systematic review and meta-analysis [239] was included on the association of Ct value with patient clinical outcomes.

- Very low certainty evidence showed that among COVID-19 cases, Ct values of 24 to 35.6 were associated with isolation of SARS-CoV-2 virus in culture. Lower Ct values (Ct <25-30) were associated with increased disease severity and mortality.
- Among convalescent and clinically recovered patients with persistent positive RT-PCR test, Ct values ranged from 30 to 41.7. Samples from these cases had a low yield of virus isolation in culture and had degraded viral fragments on genome sequencing.
- Interpretation of Ct values must be done with caution due to variations in PCR assay methods, target gene, sample type, and timing of sample collection.

CONSENSUS ISSUES

RT-PCR cycle threshold values may have utility specifically for patients with previously documented COVID-19 infection. However, evidence remains insufficient on the cut-off cycle threshold value that differentiates infectious virus from viral remnants.

Should antibody tests be used for COVID-19 seroprevalence studies and monitoring vaccine response among adults?

As of 22 November 2021

RECOMMENDATION

We suggest using antibody tests that accurately measure IgG or total antibodies to determine COVID-19 seroprevalence among adults when needed for public health purposes. *(Very low certainty of evidence; Weak recommendation)*

RECOMMENDATION

We suggest against using antibody tests detecting IgM to determine COVID-19 seroprevalence among adults. *(Very low certainty of evidence; Weak recommendation)*

We suggest against using lateral flow immunoassay (LFIA) tests to determine COVID-19 seroprevalence among adults. *(Very low certainty of evidence; Weak recommendation)*

We recommend against routine measurement of SARS-CoV-2 antibody titers after vaccination. *(No evidence; Strong recommendation)*

KEY FINDINGS

- This review included 19 observational studies [243-261] with a total of 28,566 samples that evaluated the diagnostic accuracy of antibody tests compared with reverse transcription polymerase chain reaction (RT-PCR) in seroprevalence studies.
- The studies were of moderate to high methodologic quality. The overall certainty of evidence was rated very low due to serious risk of bias (recall bias) and inconsistency.
- Heterogeneity across studies was substantial ($I^2 > 90\%$). The sensitivity of antibody tests ranged from 14.4 to 100% while specificity ranged from 54.9 to 99.6%.
- No studies evaluating the accuracy of antibody tests in determining vaccine response compared to RT-PCR-diagnosed breakthrough infections were found.

CONSENSUS ISSUES

Recommendations on antibody testing were made in the context of public health purposes (i.e., to identify the percentage of people in a population who may have been previously infected). The panel was unanimous against the use of antibody tests detecting IgM and LFIA tests while the majority voted for the use of antibody tests detecting IgG or total antibodies with a weak recommendation. Concern was raised on the cross-reactivity of the tests with other coronaviruses. Thus, emphasis was made on ensuring the accuracy of the antibody test kits used.

The panel was initially divided about antibody testing post-vaccination. Six of eleven panelists voted that there is insufficient evidence to recommend for or against it, citing that the public should be given the choice given the lack of evidence. On the other hand, five panelists voted against routine antibody testing post-vaccination despite the lack of evidence due to: (1) the unavailability of a real neutralizing antibody test in the market and (2) the unclear cutoff of antibody level that is predictive of COVID-19 protection. The panelists eventually reached consensus in the second round of voting, with the majority choosing for a strong recommendation because antibody tests post-vaccination lack clinical utility at this point.

Among symptomatic individuals previously diagnosed with COVID-19, should antibody testing be done to diagnose presumptive COVID-19 reinfection?

As of 09 April 2021

RECOMMENDATION

We recommend against the use of SARS-CoV-2 Ab testing to diagnose presumptive COVID-19 reinfection among symptomatic patients previously diagnosed with COVID-19* (*Very low certainty of evidence; Strong recommendation*).

*NAAT (RT-PCR) and Genomic sequencing are the recommended diagnostic tests to confirm COVID-19 reinfection.

KEY FINDINGS

- No studies directly assessed the accuracy of SARS-CoV-2 Ab testing in diagnosing presumptive COVID-19 reinfection compared to RT-PCR as the reference standard. Only three retrospective observational studies [262-264] reported on the accuracy of SARS-CoV-2 IgG/IgM Ab in diagnosing COVID-19. There is very low certainty regarding these estimates due to very serious risk of bias concerns, imprecision, indirectness, and inconsistency.
- The sensitivity of SARS-CoV-2 IgG/IgM ranged from 0.19 (95% CI 0.4-0.46) to 0.89 (95% CI 0.71-0.98) and specificity of 0.50 (95% CI 0.01-0.99) to 1.00 (95% CI 0.89-1.00).
- Subgroup analysis suggested that the sensitivity of Ab testing was low under the following conditions: (a) when used within 0-15 days from symptom onset, (b) Ab tests that assess IgM, (c) using lateral flow qualitative immunoassay (LFIA) technique.
- Specificity was consistently high (>89%) regardless of the type of antibody detected or if either LFIA or chemiluminescence immunoassay (CLIA) techniques were done. However, specificity was high only if the test was performed more than 16 days from symptom onset.

CONSENSUS ISSUES

The studies reviewed did not perform subgroup analysis according to 4-fold titer rise at a given interval. Due to the very low quality of evidence, the use of SARS-CoV-2 Ab testing to diagnose presumptive COVID-19 reinfection was not recommended.

Among asymptomatic individuals scheduled for non-urgent, non-emergency surgery, should RT-PCR and clinical risk assessment vs clinical risk assessment alone be done to screen for COVID-19?

As of 09 April 2021

RECOMMENDATION

We recommend using both clinical risk assessment and RT-PCR* to screen for COVID-19 among asymptomatic individuals scheduled for non-emergency surgery. *(Very low certainty of evidence; Strong recommendation)*

We recommend the use of both clinical risk assessment and Antigen-Rapid Diagnostic Test (Ag-RDT)** to screen for COVID-19 among asymptomatic individuals scheduled for non-emergency surgery when RT-PCR testing is not available or turnaround time of results is prolonged. *(Very low certainty of evidence; Strong recommendation)*

**Always use high-risk PPE regardless of RT-PCR or Ag-RDT test results in areas with prevalence of 1% or higher*

***Ag-RDT should have a Sn of 80% and Sp of 97%*

KEY FINDINGS

- Based on one cohort study [240] with very low quality, the diagnostic accuracy of clinical risk assessment alone in detecting COVID-19 compared to RT-PCR was found to be poor, with a sensitivity of 0.42 (95% CI 0.15-0.72) and a specificity of 0.85 (95% CI 0.76-0.92). Clinical risk assessment also resulted in more false negative and false positive results.
- Very low certainty evidence from one economic modeling study [241] suggested that universal pre-endoscopy virus testing using Ag-RDT, standard RT-PCR, or rapid PCR combined with high-risk PPE use in all patients irrespective of test results was more cost-effective compared to no pre-endoscopy testing and no high risk PPE use, at an assumed prevalence rate of 1% or higher among asymptomatic individuals.
- Patients for elective surgery who tested positive on any pre-operative COVID-19 tests or clinical assessment were at least three times more at risk of experiencing pulmonary complications or death compared to those who tested negative based on one cohort study [242] with very low quality. Delaying surgery to at least seven weeks from a COVID-19 diagnosis also showed benefit. Given this data on the risks and benefits associated with a COVID-19 diagnosis as well as the high false negative rates of clinical risk assessment alone, clinical risk assessment would appear to cause more harm compared to more objective tests.

CONSENSUS ISSUES

Despite the very low quality of evidence, the majority voted to strongly recommend the use of both RT-PCR testing and clinical risk assessment to screen for COVID-19 among asymptomatic individuals scheduled for non-emergency surgery primarily due to the potential impact of a false negative result on the safety of the patient and health care staff involved as well as on the infection control processes of hospitals. RT-PCR

was also recommended as it is now readily available in most hospitals. However, a panelist suggested that RT-PCR and PPE should only be conditionally recommended in areas with prevalence rates of 1% or higher.

The specification of the sensitivity and specificity for the Ag-RDT was the reason for the strong recommendation on the use of clinical risk assessment and Ag-RDT to screen for COVID-19 among asymptomatic individuals scheduled for non-emergency surgery when RT-PCR testing is not available. However, other panelists were concerned about the availability of antigen tests that would meet the set specification in terms of sensitivity and specificity.

What criteria should be used for allowing workers who were previously infected with COVID-19 to return to work?

As of 17 December 2021

RECOMMENDATION

For asymptomatic, not severely immunocompromised fully vaccinated adults, we suggest the use of the following symptom-based criteria for return-to-work clearance:

(Very low certainty of evidence; Weak recommendation)

- a. At least 8 days have passed since the first positive COVID-19 RT-PCR test; AND
- b. No symptoms have developed during this period.

For asymptomatic, not severely immunocompromised not fully vaccinated adults, we suggest the use of the following symptom-based criteria for return-to-work clearance:

(Very low certainty of evidence; Weak recommendation)

- a. At least 10 days have passed since the first positive COVID-19 RT-PCR test; AND
- b. No symptoms have developed during this period.

For symptomatic, not severely immunocompromised adults with mild-to-moderate COVID-19 diagnosis and any vaccination status, we suggest the use of the following symptom-based criteria for return-to-work clearance:

(Very low certainty of evidence; Weak recommendation)

- a. At least 10 days have passed since the onset of symptoms; AND
- b. No fever during the previous 24 hours; AND
- c. There has been substantial improvement in respiratory symptoms of the acute illness.

For symptomatic, not severely immunocompromised adults with severe-to-critical COVID-19 diagnosis and any vaccination status, we suggest the use of the following symptom-based criteria for return-to-work clearance:

(Very low certainty of evidence; Weak recommendation)

- a. At least 21 days have passed since the onset of symptoms; AND
- b. No fever during the previous 24 hours; AND
- c. There has been substantial improvement in respiratory symptoms of the acute illness.

For symptomatic, severely immunocompromised adults with any vaccination status, we suggest the use of the following for return-to-work clearance:

(Very low certainty of evidence; Weak recommendation)

- a. At least 22 days have passed since the onset of symptoms; AND
- b. No fever during the previous 24 hours; AND
- c. There has been substantial improvement in respiratory symptoms of the acute illness; AND
- d. PCR test results are negative on at least 1 respiratory specimen.

Note:

Severely immunocompromised individuals include the following:

- Individuals receiving active chemotherapy for cancer
- Being within one year out from receiving a hematopoietic stem cell or solid organ transplant
- Untreated HIV infection with CD4 <200
- Primary immunodeficiency
- Taking immunosuppressive medications (e.g., drugs to suppress rejection of transplanted organs or to treat rheumatologic conditions such as mycophenolate and rituximab)
- Taking more than 20mg a day of prednisone for more than 14 days

KEY FINDINGS

- This review included six observational studies [265-270] that explored the relationship of vaccination status on various outcomes related to infectivity. From this data, the review provides additional recommendations related to duration of isolation for return-to-work clearance.
- One study [268] found no significant difference ($p=0.16$) in the proportion of culture-positive results between fully vaccinated and not fully vaccinated individuals (OR 0.67, 95% CI 0.38-1.18). The duration of viral culture positivity between the two groups were comparable (median of 5 days for both) and suggested no significant difference in terms of duration of infectivity (mean difference=0 days). Another study [267] concluded that vaccinated individuals had faster viral clearance by 2 days (5.5 days, 95% credible intervals [CrI] 4.6-6.5) vs. 7.5 days for unvaccinated (95% CrI 6.8-8.2 days) and shorter infection duration by 2.3 days (8.7 days, 95% CrI 7.6-9.9) vs 11.0 days (95% CrI 10.3-11.8) for unvaccinated).
- Based on three observational studies, [265,266,269] the secondary attack rate was lower by 8.43% (95% CI -18.03%, 1.17%) for fully vaccinated compared to not fully vaccinated individuals, but this difference was not statistically significant ($P=0.09$).
- Compared to immunocompetent individuals, time to PCR clearance was not significantly different for severely (adjusted hazards ratios [aHR] 0.98, 95% CI 0.84-1.15) and moderately immunocompromised patients (aHR 0.86; 95% CI 0.71-1.05). Delayed time to PCR clearance was seen for specific subgroups of patients: solid organ transplant (aHR 0.64, 95% CI 0.42-0.97), diabetes (aHR 0.82, 95% CI 0.73-0.93), obesity (aHR 0.90, 95% CI 0.83-0.98), rheumatologic disease (aHR 0.90, 95% CI 0.83-0.98), ≥ 3 comorbidities (aHR 0.73, 95% CI 0.60-0.88), older age (aHR 0.996, 95% CI 0.993-0.999).
- The overall certainty of evidence for each of the outcomes was rated very low. Downgrading occurred due to risk of bias issues across the included studies, imprecision related to wide confidence intervals and small sample sizes, and inconsistency.

CONSENSUS ISSUES

The evidence base included studies that investigated the clearance of COVID-19 infection in relation to vaccination status, symptom presentation (i.e., asymptomatic versus symptomatic), and immunocompromised state (i.e., non-, moderately, or severely immunocompromised). These factors were taken into consideration in recommending the criteria for allowing workers who were previously infected with COVID-19 to return to work.

Despite studies including non-, moderately, and severely immunocompromised individuals, the panelists decided to make recommendations that only distinguished severely versus non-severely immunocompromised individuals due to insufficiency of evidence among moderately immunocompromised individuals.

Among asymptomatic, not severely immunocompromised individuals, separate recommendations were made depending on vaccination status. Evidence base including one study showed faster viral clearance and shorter infection duration by two days among fully vaccinated compared to not fully vaccinated individuals. The panelists unanimously voted for a weak recommendation due to indirectness of the evidence, wherein the reference standard used was time to RT-PCR clearance instead of viral culture positivity.

Among severely immunocompromised individuals, the evidence on prolonged viral shedding was an important consideration for the panelists. However, data remains lacking on the duration of infectivity of the virus among this subgroup. The evidence base included only one study on this subgroup and the reference standard for infectivity was RT-PCR instead of viral culture. For this reason, a panelist opined that a test-based criteria for clearance to work would be a more prudent strategy compared to a symptom-based criteria. However, concern on the impracticality of a test-based approach was raised as this may lead to prolonged isolation of patients who continue to shed detectable SARS-CoV-2 RNA on RT-PCR but are no longer infectious. A weak recommendation was made mainly due to the very low certainty of evidence.

Among individuals suspected of COVID-19, how accurate are thoracic imaging modalities compared to RT-PCR alone in diagnosing COVID-19?

As of 13 December 2021

RECOMMENDATION

We suggest against the use of chest x-ray to diagnose COVID-19 infection among asymptomatic individuals. (*Very low certainty of evidence; Weak recommendation*)

We suggest against the use of lung ultrasound alone in diagnosing patients with suspected COVID-19 infection. (*Very low certainty of evidence; Weak recommendation*)

We suggest against the routine use of CT scan for diagnosing COVID-19 among suspected patients with COVID-19 presenting at the emergency department if RT-PCR testing is readily available with timely results. (*Very low certainty of evidence; Weak recommendation*)

RECOMMENDATION

We suggest chest x-ray to facilitate rapid triage, infection control, and clinical management among any of the following: (*Very low certainty of evidence; Weak recommendation*)

- a. Patients with mild features of COVID-19 at risk for progression
- b. Patients with moderate to severe features of COVID 19
- c. Patients with symptoms of at least 5 days duration

If RT-PCR is not available, we suggest using non-contrast chest CT scan for symptomatic patients suspected of having COVID-19 to guide early triage and management under the following conditions: (*Very low certainty of evidence; Weak recommendation*)

- a. Mild COVID-19 patients who are at risk for progression
- b. Moderate to severe COVID-19 patients

KEY FINDINGS

- A total of 81 observational studies [271-351] were assessed for the diagnostic accuracy of chest x-ray, lung ultrasound, and chest CT scan against reverse transcriptase-polymerase chain reaction (RT-PCR) on individuals suspected of COVID-19.
- **Chest x-ray:** Overall sensitivity was 72% (62-81%) and overall specificity was 76% (67-86%). Results were comparable to the findings of the previous review that showed a sensitivity of 74% and specificity of 76%. Considerable heterogeneity is still seen ($I^2=95\%$). Sensitivity was higher for studies that involved experienced readers, used standardized chest x-ray scoring systems, and when testing was done late in the disease course. Studies with high risk of bias tended to produce similar but less precise estimates. Overall certainty of evidence remained very low due to very serious risk of bias and inconsistency, and serious imprecision.
- **Lung ultrasound:** Overall sensitivity was 93% (86-97%) and specificity was 52% (33-71%). The previous review showed a lower sensitivity of 88% but a higher

specificity of 63%. Sensitivity appeared higher when the test is used for symptomatic patients and when reader impression is used instead of a scoring system. The highest accuracy estimates were produced when only high quality studies were included in the analysis (Sn 97%, 95% CI 89, 100%; Sp 73%, 95% CI 45, 92%). Overall certainty of evidence was downgraded from low to very low due to very serious inconsistency, and seriousness on risk of bias and imprecision.

- **Chest CT scan:** Overall sensitivity was 85% (81-88%) and overall specificity was 78% (71-84%). Heterogeneity was very high at $I^2=100\%$. Findings were almost similar to the estimates (Sn 88% and Sp 80%) of the previous review. Accuracy estimates were higher under the following situations: (1) test is used among symptomatic patients, (2) contrast-enhanced CT scan machines, (3) results are interpreted by experienced readers, (4) standardized scoring systems are used, (5) when only high-quality studies were considered. Overall certainty of evidence remained very low due to the seriousness of risk of bias and impression, and the very serious issues on inconsistency.

CONSENSUS ISSUES

The panel was unanimous against the use of chest x-ray and lung ultrasound in diagnosing COVID-19 due to the widespread availability of alternatives, such as antigen testing, even in the absence of RT-PCR. Additionally, the evidence base showed that detecting COVID-19 through these imaging modalities are dependent on the experience of the reader. However, a weak recommendation was decided since the evidence did not compare these imaging modalities with rapid antigen tests.

The panel was also unanimous in recommending chest x-ray to facilitate triage, infection control, and clinical management, especially in the absence of RT-PCR and rapid antigen tests, due to the widespread availability and rapid results of chest x-ray. A weak recommendation was made, however, since the accuracy of this modality varies depending on the experience of the reader.

Similarly, the panel was for the use of non-contrast chest CT scan in guiding early triage and clinical management if RT-PCR is not available. A weak recommendation for this was made due to concerns on its risk of radiation and cost, and the evidence base including studies conducted abroad.

Among adult patients diagnosed with COVID-19, should prognostic models be used to predict the likelihood of severe disease and mortality?

As of 17 December 2021

RECOMMENDATION

To guide the decision to admit adult patients with COVID-19 to the hospital:

We suggest the use of age, BUN, number of comorbidities, CRP, SpO₂/FiO₂ ratio, platelet count, Heart rate (ABC2-SPH) risk score, Confusion Urea Respiration Blood Pressure (CURB-65) severity score, Risk Stratification in the Emergency Department in Acutely Ill Older Patients (RISE-UP) score, and Rapid Emergency Medicine Score (REMS). *(Low certainty of evidence; Weak recommendation)*

To guide in the expectant monitoring of hospitalized adult patients, we suggest the use of the 4C Deterioration model. *(Low certainty of evidence; Weak recommendation)*

RECOMMENDATION

To guide the decision to admit adult patients with COVID-19 to the hospital, there is insufficient evidence to recommend the use of 4C Mortality Score, COVID Outcome Prediction in the Emergency Department (COPE) model, and Quick Sepsis-related Organ Failure Assessment (qSOFA) score. *(Very low certainty of evidence)*

To guide in the expectant monitoring of hospitalized adult patients, there is insufficient evidence to recommend the use of Modified Early Warning Score (MEWS) and National Early Warning Score 2 (NEWS2), Clinical Frailty Scale (CFS), and the COVID-GRAM model. *(Very low certainty of evidence)*

KEY FINDINGS

- In this version, two new prognostic models were reviewed and 13 new studies [352-364] were added. One study is a binational prospective cohort which validated 4C mortality score and COVID-GRAM on a larger population in South America and Europe [352]. Three more studies [353-355] validated COVID-GRAM but have an overall unclear risk of bias and low quality of evidence. Nine studies [356-364] validated the use of Clinical Frailty Scale in prognosticating elderly COVID-19 patients, but the overall assessment for risk of bias was rated high and the level of evidence was rated very low.
- In total, 46 cohort studies [352-397] on prognostic models for clinical deterioration and mortality of individuals with COVID-19 were found. Most of the studies (n=36) were assessed to have high risk of bias due to issues in participant selection and analysis. There were eight studies [352, 357-358, 360, 364-367] with unclear, and two studies [368-369] with low risk of bias.
- For predicting mortality, the following models demonstrated fair-to-good predictive ability: 4C mortality score, ABC2-SPH, CURB-65, REMS, RISE UP and COVID-GRAM models. Poor to fair prediction was noted for the qSOFA model, with one new study yielding lower AUC estimates compared to the previously included studies. Only one model, the QCOVID model for mortality validated for the UK setting, demonstrated high predictive ability.

- For predicting clinical deterioration, available prognostic models showed varied performance. The 4C deterioration which has been investigated in only one study with low risk of bias, showed fair predictive ability. The MEWS model has poor prediction of clinical deterioration while NEWS2 has inconsistent prediction (poor to good). The Clinical Frailty Scale (CFS), increases the risk of mortality among elderly COVID-19 patients.
- None of these models have been validated in the Philippine population. Thus, validation studies are needed before these models can be used to inform practice.

CONSENSUS ISSUES

The recommendations on prognostic models are limited to adult patients since the evidence base included studies only on the adult population. No studies were conducted among pediatric patients. The panelists were unanimous in all recommendations on prognostic models.

Should LDH, CRP, and Ferritin be used to guide immunotherapy in patients with COVID-19?

As of 13 December 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of specific cut-off values of CRP, LDH and Ferritin to guide the initiation of immunotherapy in patients with COVID-19 (*Very low certainty of evidence*)

KEY FINDINGS

- Moderate quality of evidence showed significant correlation of CRP >10 mg/L with mortality (four studies [398]) and poor outcome (ten studies [399]).
- High quality of evidence from seven studies [398] showed significant correlation between elevated LDH and progression to severe disease.
- Moderate quality of evidence from 18 studies [399] showed significant correlation between LDH >250 U/L and poor outcome.
- Low quality of evidence from four studies [398,400-402] did not find significant correlation between elevated LDH and mortality.
- Very low quality of evidence showed significant correlation between elevated CRP and progression to severe disease (one study [403]) and ICU admission (one study [403]) in pediatric patients.

CONSENSUS ISSUES

The evidence base included studies that investigated the levels of LDH, CRP, and ferritin that are correlated with progression to severe disease, poor outcomes, or mortality, rather than improved survival with immunotherapy. Thus, the panel unanimously decided that insufficient evidence remains to use specific cut-off values of these markers to guide immunotherapy. Additionally, the panel cited the availability of other parameters that may be used to guide initiation of immunotherapy besides these markers.

Should D-dimer be used to guide anticoagulation among adult patients with COVID-19?

As of 26 May 2021

RECOMMENDATION

We suggest the use of D-dimer to guide anticoagulation of adult patients with COVID-19, because of its significant association with mortality, thromboembolism, and worsening of disease (*Low certainty of evidence; Weak recommendation*)

KEY FINDINGS

- We found a total of 25 observational studies [404-428] on the association between D-dimer and the outcomes of mortality, worsening severity, or thromboembolism. In general, the included studies showed increased odds of in-hospital mortality (OR 5.57, 95% CI 2.74-11.31), worsening severity (critical illness OR 1.91-2.58); disease progression (HR 2.84, 95% CI 2.10-3.85), or need for mechanical ventilation (HR 3.28, 95% CI 1.071-0.10), and thromboembolism (OR 5.61, 95% CI 3.97-7.94), with higher D-dimer levels across different COVID-19 severities.
- Most studies yielded imprecise effect measures, due to the small number of event outcomes. Most of the studies were found to have serious risk of bias, with issues on data censoring, incomplete laboratory data, and unclear adequacy of follow-up rates. Differences in D-dimer cut-offs, definitions of critical illness and disease progression, and severities of COVID-19 in the study population contributed to the heterogeneity across studies.
- While the predictive ability of D-dimer for mortality appeared to be fair to good, prediction of worsening severity or progression of disease is inconsistent.

CONSENSUS ISSUES

Due to the varied cut-off values used in the included studies for this review, the recommendation did not include a specific cut-off value of D-dimer to predict mortality, thromboembolism, and worsening severity of disease. Further, the laboratories in the Philippines also make use of varying cut-offs due to the different assays and machines used, hence, it is difficult to define a specific cut-off value.

Should procalcitonin be used to guide the initiation of antibiotic therapy in patients diagnosed with COVID-19?

As of 13 December 2021

RECOMMENDATION

For initiating antibiotic therapy, we suggest against the use of procalcitonin alone as a basis for initiating antibiotic therapy among COVID-19 confirmed patients. (*Very low certainty of evidence; Weak recommendation*)

RECOMMENDATION

For discontinuing antibiotic therapy:

If available, we recommend using a procalcitonin level of less than or equal to 0.25 ng/mL for discontinuing antibiotic therapy among COVID-19 confirmed patients. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

- Four retrospective cohort studies [429-432] were included in this review. Three [429-431] provided data for calculation of diagnostic accuracy estimates, while one [432] reported on the benefits and safety of using an antimicrobial therapy discontinuation protocol based on procalcitonin measurements. Studies used varying cut-off levels for procalcitonin and used it for guiding discontinuation of antibiotics.
- Sensitivity ranged from 0.40 to 0.87, while specificity ranged from 0.43 to 0.88. Across the three studies, a total of 42/816 (5.1%) had positive results on microbiological testing. No stratification as to whether patients with higher procalcitonin levels were associated with more severe or less severe COVID-19 cases.
- Using procalcitonin-based algorithms was found to be associated with lower antimicrobial exposure (mean 5.4 vs. 4.4 days) and consumption (mean DDD 8.4 vs. 6.8). This translates to a 30% lower risk for both mean duration of antibiotic therapy (adjusted ratios of means [ROM] 0.70; 95% CI 0.6, 0.9) as well as mean DDD (ROM 0.70, 95% CI 0.6, 0.8). A trend towards lower 30-day mortality rates was noted for participants with procalcitonin measurements (adjusted prevalence ratio [PR] 0.6, 95% CI 0.4, 1.1). Mean patient antimicrobial consumption also showed a decreasing trend over time in the procalcitonin group (β slope -0.07 , 95% CI -0.11 , -0.03) compared to those without procalcitonin measurements.
- Overall certainty of evidence was rated very low due to serious risk of bias, imprecision, and inconsistency.

CONSENSUS ISSUES

Antibiotic misuse and abuse among COVID-19 patients is a common practice that needs to be addressed. Thus, the panel recognized that procalcitonin may have utility in guiding antibiotic use among COVID-19 patients. However, the panel was against the use of procalcitonin alone in initiating antibiotic therapy, citing that clinical evaluation and other laboratory parameters should be considered as well since procalcitonin levels can be elevated in certain conditions even in the absence of bacterial infection. Additionally, the evidence base included studies that investigated

absolute cut-offs of procalcitonin rather than the comparative decrease or increase in its levels.

In contrast, the panel was unanimous in recommending the use of procalcitonin levels to discontinue antibiotic therapy among COVID-19 patients. This was to provide guidance to practitioners due to the common practice of continuing antibiotics indefinitely among critically ill patients.

Cited reasons for a weak recommendation on the use of procalcitonin for discontinuing antibiotic therapy include:

1. concerns on the cost and availability of procalcitonin;
2. very low certainty of evidence; and
3. use of indirect evidence to identify the cut-off level for discontinuation (i.e., the evidence base aimed to identify procalcitonin levels that differentiate the presence or absence of co-infection rather than the continuation or discontinuation of antibiotic therapy).

Cited reasons for a strong recommendation on the use of procalcitonin for discontinuing antibiotic therapy include:

1. the need to prevent antibiotic resistance, considering that multidrug resistance is a common cause of death among critically ill or severe COVID-19 patients; and
2. to prevent the higher costs associated with unnecessary antibiotic use (i.e., While procalcitonin is expensive, even an extra day of unnecessary antibiotic therapy would equate to higher costs for the patient).

Should certain risk factors be used to predict the development of long COVID?

As of 29 November 2021

RECOMMENDATION

There is insufficient evidence in using symptoms*, biologic factors or severity of acute COVID-19 in predicting the development of long covid symptoms. (*Very low certainty of evidence*)

**The most common symptoms of long COVID identified were fatigue, dyspnea, sleep disturbance, anxiety or depression, and memory impairment.*

KEY FINDINGS

- Evidence for this review was obtained from 37 observational studies included in two systematic reviews [433-434] as well as three additional observational studies [435-437] describing possible risk factors for the development of long COVID.
- Majority of hospitalized patients diagnosed as having long COVID presented with the following symptoms: weakness, fatigue, dyspnea, cognitive/memory impairment, sleep disorder, and anxiety/psychosocial symptoms.
- For patients in the community setting, dyspnea, reduced quality of life, weakness, chest pain, palpitations, arthralgia, and myalgia were highly common. Older age, female sex, presence of comorbidities, and more severe status during the initial infection were likely to be associated with the development of long COVID during follow-up, although studies showed inconsistent results.
- Very low certainty evidence showed that some of these risk factors were associated with individual symptoms characteristic of long COVID. Our analysis was limited by the significant heterogeneity among the studies included, the different time frames of follow-up, and the studies' inclusion criteria.

CONSENSUS ISSUES

Predicting the development of long COVID-19 using risk factors can help identify which patients should be followed up more closely. However, there is insufficient evidence on the predictive risk factors, precluding any recommendation to be made.

Should heparin induced thrombocytopenia (HIT) test kits be used for COVID-19 vaccine induced thrombosis with thrombocytopenia (VITT)?

As of 29 November 2021

RECOMMENDATION

We suggest against the use of PF4 antibody ELISA Heparin Induced Thrombocytopenia (HIT) test kits and non-ELISA rapid HIT test kits for COVID-19 Vaccine Induced Thrombosis and Thrombocytopenia (VITT). (*Very low certainty of evidence; Weak recommendation*)

KEY FINDINGS

- One cross-sectional observational study [438] was included on the use of anti-platelet factor 4 (anti-PF4) assay test kits for the diagnosis of COVID-19 vaccine-induced thrombosis and thrombocytopenia (VITT).
- The overall certainty of evidence was very low because of serious concerns of risk of bias, inconsistency, and imprecision.
- The study analyzed samples from 43 patients with suspected VITT using six brands of anti-PF4 ELISA test kits and four brands of anti-PF4 non-ELISA test kits. Sensitivity ranged from 0.71-0.97 and specificity ranged from 0.56-1.00 for the ELISA test kits, while sensitivity ranged from 0.00-0.45 and specificity ranged from 0.67-1.00 for the non-ELISA test kits.

CONSENSUS ISSUES

The panel unanimously decided against the use of HIT test kits for COVID-19 VITT, citing that the test has a slow turnaround time and immediate clinical management remains the same regardless of the test result. Concerns were also raised on its cost-effectiveness as well as availability in the local setting. Test kits employing ELISA demonstrate relatively high sensitivity and specificity but have limited availability locally.

Should serum tryptase be used to test individuals who had anaphylaxis after receiving COVID-19 vaccine?

As of 29 November 2021

RECOMMENDATION

We suggest against using serum tryptase for patients who had anaphylaxis after receiving COVID-19 vaccine. (*Very low certainty of evidence; Weak recommendation*)

KEY FINDINGS

- Eleven observational studies (five case reports [439-443], two case series [444-445], three retrospective chart reviews [446-448] and one prospective survey [449]) with 33 total patients showed conflicting results and uncertainty with regards to the use of serum tryptase in the diagnosis of COVID-19 vaccine induced anaphylaxis.
- Nine patients presented with a history of allergy but none of them showed elevated tryptase levels.
- Four patients presented with elevated serum tryptase levels but have no allergic history elicited/provided.
- There is uncertainty with regards to the accuracy of serum tryptase in the diagnosis of COVID-19 vaccine-induced anaphylaxis due to the lack of study samples.
- The overall certainty of evidence was very low because of heterogeneity in duration of tryptase determination, definition of elevated values, variable clinical classification criteria, and lack of an objective reference standard for diagnosis of anaphylaxis.

CONSENSUS ISSUES

The panel unanimously decided against the use of serum tryptase to diagnose COVID-19 vaccine-induced anaphylaxis, citing that the test is not cost-effective, and anaphylaxis remains to be a clinical diagnosis. While serum tryptase may have a role in patients with atypical presentation, it would still be prudent to treat the patient especially if there is high clinical suspicion regardless of the test result. Additionally, the use of tryptase test in settings of mass vaccination poses feasibility issues because (1) the sample should be taken within 30 minutes up to two hours after the patient presents with a reaction, and (2) there is a paucity of laboratories offering the test. However, a weak recommendation was made since the test may still benefit a small subset of the population and may have more utility in the emergency department setting.

Evidence and Recommendations for the Treatment of COVID-19

Should hydroxychloroquine/ chloroquine, with or without azithromycin be used in the treatment of patients with COVID-19 infection?

As of 19 February 2021

RECOMMENDATION

We recommend against the use of hydroxychloroquine/chloroquine, with or without azithromycin among patients with COVID-19 infection. (*Moderate certainty of evidence, Strong recommendation*)

KEY FINDINGS

- Twenty-two trials [1-22] showed that hydroxychloroquine (HCQ)-containing treatment regimens did not significantly improve the outcomes of patients with COVID-19 disease compared with placebo or standard of care. Eleven (11) of these trials provided evidence of low certainty that the use of HCQ for the treatment of COVID-19 infection was significantly associated with a higher risk of adverse events (i.e., diarrhea, headache, rashes, and fatigue).
- There was limited evidence with low to very low certainty which showed that treatment with HCQ combined with azithromycin does not show any significant difference from placebo in any of the efficacy outcomes. Adverse events were more frequent with the hydroxychloroquine plus azithromycin group compared to placebo.

Should azithromycin be used in the treatment of patients with COVID-19 infection?

As of 1 December 2021

RECOMMENDATION

We recommend against the use of azithromycin among patients with COVID-19 infection. (*Moderate certainty of evidence, Strong recommendation*)

KEY FINDINGS

Based on 11 RCTs and one additional RCT [23-30] for asymptomatic to mild COVID - 19 patients, there was no significant difference in clinical outcomes with the use of azithromycin compared to placebo among patients with COVID-19 across all disease severity. Likewise, there was no significant difference in serious adverse events and cardiac arrhythmias among all patients but there was a noted significantly higher risk among those given azithromycin for non-serious adverse events for the asymptomatic to mild group with diarrhea being the most common adverse event. This was not noted among the moderate to severe group.

Among patients with COVID-19, should favipiravir be used for treatment?

As of 8 November 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of favipiravir among patients diagnosed with COVID-19. (*Very low certainty of evidence*)

KEY FINDINGS

- Seven (7) randomized controlled trials (RCTs) [31-37] found on the use of favipiravir among patients with COVID-19. Pooled results showed a modest benefit in clinical improvement on day 7 favoring favipiravir compared to standard of care; however, clinical improvement on day 28 showed no significant benefit. There was also significant benefit in time to clinical improvement and time to negative conversion. Incidence of viral negative conversion was not significantly different between favipiravir and standard of care.
- There was no significant difference on the incidence of adverse events and serious adverse events. Report on adverse events, although an important outcome, was not rated as a critical outcome to be included in the decision making. The overall certainty of evidence was rated low due to serious risk of bias, inconsistency, and very serious imprecision in several critical outcomes.

CONSENSUS ISSUES

Results from the study are mostly inconclusive and there are still no recommendations on the use of favipiravir outside clinical trials. There are ongoing clinical trials, including one local study currently recruiting participants. Results from these ongoing studies will help further evaluate the use of favipiravir in the treatment of COVID-19.

Should remdesivir be used in the treatment of patients with COVID-19 infection?

As of 19 February 2021

RECOMMENDATION

We suggest the addition of remdesivir to dexamethasone in patients with COVID-19 infection who have O₂ saturation < 94% and/or requiring oxygen supplementation and are high risk for progression. (*Low certainty of evidence, Weak recommendation*)

**For patients who progress to invasive mechanical ventilation while on remdesivir, the drug may be continued.*

RECOMMENDATION

We suggest against the use of remdesivir in patients with COVID-19 infection who have O₂ saturation ≥94% and do not require oxygen supplementation. (*Low certainty of evidence, Weak recommendation*)

RECOMMENDATION

We suggest against the use of remdesivir in patients with COVID-19 infection who are already on invasive mechanical ventilation or ECMO. (*Low certainty of evidence, Weak recommendation*)

KEY FINDINGS

- Four randomized controlled trials on the use of remdesivir as treatment for COVID-19 were found [38-41]. Low certainty of evidence shows that remdesivir has limited effect on all cause-mortality, clinical improvement, and initiation of mechanical ventilation among confirmed COVID-19 patients. However, remdesivir appears to be beneficial in the time to clinical improvement especially among cases needing supplemental oxygen but not high flow oxygen/ mechanical ventilation.
- Remdesivir did not show increased risk for serious adverse events. Availability and cost of intervention should be considered before making recommendations regarding its use locally.

CONSENSUS ISSUES

Early introduction of remdesivir in the treatment of COVID-19 is preferred because of its action on the polymerase resulting in less viral replication. Remdesivir is a relatively safe drug, but its cost should be considered. Hence, routine use of the drug is not recommended. There are 26 ongoing trials pertaining to the efficacy and safety of remdesivir for the treatment of COVID-19.

Among patients with COVID-19, should molnupiravir be used for treatment?

As of 10 January 2022

RECOMMENDATION

We suggest the use of molnupiravir within 5 days of symptom onset among non-hospitalized adult patients (18 years old and older) with mild to moderate COVID-19 infection with at least one risk factor* for progression. (*Low certainty of evidence, Weak recommendation*)

**Risk factors for progression include:*

age >60 years, active cancer, chronic kidney disease, chronic obstructive pulmonary disease, obesity, serious heart conditions or diabetes mellitus

KEY FINDINGS

- Five (5) randomized controlled trials (RCTs) [42-46] studied the effect of molnupiravir on the treatment of COVID-19 compared to standard of care and/or placebo. Molnupiravir significantly decreased the need for hospitalization at day 29. Subgroup analysis based on the severity showed significant reduction in mortality in non-hospitalized mild to moderate patients but did not show significant benefit among hospitalized mild to severe patients. For other critical outcomes, molnupiravir had no apparent effect on clinical improvement based on patient reported outcome measures (FLU-PRO), WHO COVID-19 ordinal scale, National Early Warning Score (NEWS2) and sustained recovery. Adverse events and serious adverse events were similar between molnupiravir and standard of care and/or placebo.
- The phase 2a study had randomization issues and reporting bias. The serious risk of bias and imprecision led to the downgrading of evidence to low certainty.

CONSENSUS ISSUES

Molnupiravir showed general net benefit, specifically for mortality. The drug should be given specifically within the given time period (within 5 days of symptom onset). There are still issues to consider about the availability and use since it is still limited (currently available under compassionate special permit issued by FDA).

Should baloxavir be used in the treatment of patients with COVID-19 infection?

As of 20 May 2021

RECOMMENDATION

We suggest against the use of baloxavir as treatment for patients with COVID-19 infection. (Very low certainty of evidence, Weak recommendation)

KEY FINDINGS

There was only one randomized controlled trial [47] included in this review. Treatment with baloxavir did not lead to significant reduction in the need for invasive mechanical ventilation, admission to intensive care unit, hospitalization and clinical improvement (very low quality of evidence). There was no report of mortality in any of treatment arms. Currently, there are two ongoing clinical trials on the efficacy of baloxavir as treatment for COVID-19.

CONSENSUS ISSUES

Only one small study was included in the review, which showed consistent result of no significant benefit across all the outcomes measured (i.e., need for mechanical ventilator/ ECMO, admission to ICU, hospitalization, clinical improvement at 14 days, adverse events and time to clinical improvement). It was noted that baloxavir is a repurposed drug for COVID-19 and is originally indicated for influenza. It costs Php 450 per tablet and is usually given within 48 hours from the onset of symptoms for the treatment of influenza. In terms of health equity, it was raised that since its benefit is yet to be established, resources should be allocated to more known and established drugs where the benefits are certain.

Should oseltamivir be used for the treatment of COVID-19?

As of 22 May 2021

RECOMMENDATION

We recommend against the use of oseltamivir as treatment for patients with COVID-19 infection. (*Very low certainty of evidence, Strong recommendation*)

KEY FINDINGS

Based on the five retrospective cohort studies [48-52] included in this review, oseltamivir was associated with increased risk of mortality (Odds Ratio (OR), 4.20; [95%CI 4.03, 4.38], very low certainty of evidence). Moreover, it was associated with risk of disease progression (OR 5.22; [95% CI, 2.00, 13.02], low quality of evidence) as well as longer time to viral clearance (standard mean difference (SMD) of 1.65 days longer [95% CI 1.27, 2.03], low quality of evidence). Currently, there are five ongoing clinical trials on the efficacy of oseltamivir as treatment for COVID-19.

CONSENSUS ISSUES

The panel made a strong recommendation against the use of oseltamivir noting that the mortality rate among patients given the drug is higher compared to those given standard of care (i.e., 27% vs. 6%). Likewise, results showed that the progression to severe disease is five times more likely among patients taking oseltamivir.

Should lopinavir/ritonavir be used in the treatment of COVID-19?

As of 07 April 2021

RECOMMENDATION

We recommend against the use of lopinavir/ritonavir as treatment for COVID-19 infection. (*Moderate certainty of evidence, Strong recommendation*)

KEY FINDINGS

Based on 5 randomized controlled trials [53-57] enrolling both suspected and confirmed COVID-19 patients, lopinavir/ritonavir combination did not significantly affect clinical improvement, all-cause mortality, viral negative conversion, need for mechanical ventilation, or WHO progression score level 7 or higher, when compared to standard of care. No difference in adverse events were also seen between lopinavir/ritonavir and control groups. The overall quality of evidence ranged from low to high across different outcomes.

Among patients with COVID-19, should tocilizumab be used for treatment?

As of 28 October 2021

RECOMMENDATION

We recommend the addition of tocilizumab to systemic steroids in patients showing rapid respiratory deterioration and/or requiring high doses of oxygen (high-flow nasal cannula, noninvasive or invasive mechanical ventilation) and with elevated biomarkers of inflammation (CRP). (*Moderate certainty of evidence, Strong recommendation*)

RECOMMENDATION

We recommend against the use of tocilizumab among patients with COVID-19 infection who do not require oxygen. (*Very low certainty of evidence, Strong recommendation*)

KEY FINDINGS

Fifteen (15) randomized controlled clinical trials (RCTs) [58-72] (N = 8,937) that investigated the effectiveness of tocilizumab among confirmed hospitalized COVID-19 patients compared to placebo and/or standard of care was found. Tocilizumab significantly reduced all-cause mortality, time to clinical improvement, and the need for mechanical ventilation, with no significant increase in adverse events among hospitalized patients.

CONSENSUS ISSUES

Report on adverse events, although an important outcome, was not rated as a critical outcome to be included in the decision making. The over-all assessment of the certainty of evidence is based only on critical outcomes identified by the consensus panel, hence, the quality of evidence was retained as moderate.

There is no new evidence on the use of tocilizumab among patients with COVID-19 infection who do not require oxygen. Given the lack of evidence, potential adverse effects and risks, this recommendation also considers indiscriminate use to avoid misuse or overuse of tocilizumab among patients who do not require oxygen.

Among patients with COVID-19, should baricitinib be used for treatment?

As of 21 October 2021

RECOMMENDATION

We suggest the use of baricitinib in addition to dexamethasone and remdesivir as treatment for hospitalized COVID-19 patients who require low-flow oxygen, high-flow oxygen, and non-invasive ventilation. (*Low certainty of evidence; Weak recommendation*)

RECOMMENDATION

There is insufficient evidence to recommend baricitinib as an alternative to tocilizumab as treatment for hospitalized COVID-19 patients. (*Very low quality of evidence*)

KEY FINDINGS

The evidence on the use of baricitinib comes from 2 RCTs [73-75] among hospitalized patients with moderate to severe COVID-19. Both studies found that there was a reduction in the all-cause mortalities, progression of oxygen use, and serious adverse events in the baricitinib group compared to the control group. However, there was no significant difference in the duration of hospitalization between the two groups.

CONSENSUS ISSUES

The recommendation to give baricitinib in addition to dexamethasone and remdesivir was made based on 2 randomized controlled trials wherein baricitinib was given as an add-on therapy to dexamethasone and remdesivir among hospitalized patients with moderate to severe COVID-19. At present, dexamethasone, a systemic corticosteroid, is considered standard of care for patients requiring oxygen supplementation (High quality of evidence; Strong recommendation), while remdesivir may be considered as an additional therapy for patients who require oxygen supplementation but not on invasive mechanical ventilation (Low quality of evidence, Conditional recommendation). Although remdesivir is not currently considered standard care, there are studies that show possible synergistic effect in concurrently giving an anti-viral to shorten viral clearance plus an immunomodulator to address impending cytokine storm. Addition of baricitinib is recommended only for non-intubated patients (i.e., on low-flow oxygen, high-flow oxygen, and non-invasive ventilation).

As of writing, the available evidence on baricitinib is not as robust as tocilizumab, and there is only 1 retrospective observational study, which directly compared baricitinib versus tocilizumab. As such, there is insufficient evidence to make recommendations on whether baricitinib may be used as an alternative to tocilizumab. Strategies for conservation of tocilizumab is beyond the scope of the consensus panel, and may be discussed by the local expert group working on COVID-19 treatment algorithms.

Among patients with COVID-19, should imatinib be used for treatment?

As of 8 November 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of imatinib among patients with COVID-19 infection. (*Low certainty of evidence*)

KEY FINDINGS

There is one (1) randomized controlled trial (RCT) [76] that investigated the effect of imatinib compared to placebo as treatment for hospitalized patients with COVID-19 requiring supplemental oxygen. Imatinib was associated with significant reduction in the duration of mechanical ventilation and duration of intensive care unit (ICU) stay. However, there was no significant benefit for several outcomes such as mortality at day 28, need for ICU admission, need for mechanical ventilation, discontinuation of supplemental oxygen and mechanical ventilation, discontinuation of ventilation and supplemental oxygen in 48 hours, and duration of hospital admission. There was also no significant difference in the number of adverse events. The overall certainty of evidence was rated low; evidence was downgraded for serious risk of bias and imprecision due to wide confidence intervals and small number of events.

CONSENSUS ISSUES

Further trials are needed to recommend the use of imatinib for the treatment of COVID-19.

Among patients with COVID-19, should tofacitinib be used for treatment?

As of 21 October 2021

RECOMMENDATION

We suggest against the use of tofacitinib among hospitalized COVID-19 patients. (*Low certainty of evidence, Weak recommendation*)

KEY FINDINGS

- There is one (1) randomized controlled trial (RCT) [77] that investigated the effect of tofacitinib compared to placebo as treatment for patients with COVID-19. Patients treated with tofacitinib had a significant reduction in death or respiratory failure. Tofacitinib did not show significant effect in all-cause mortality, need for mechanical ventilation or extracorporeal membrane oxygenation (ECMO), cure (defined as resolution of fever, cough, or need for ventilatory/oxygen support), length of hospitalization, length of ICU stay, and duration of mechanical ventilation.
- There was no significant increase in serious adverse events and adverse events between the tofacitinib and placebo group. However, there was a significant increase in adverse events leading to treatment discontinuation for patients given tofacitinib compared to placebo, with increase in transaminase levels and lymphopenia being the most commonly reported adverse events. The very serious

imprecision due to the limited number of events contributed to the downgrading of evidence to a low certainty of evidence.

CONSENSUS ISSUES

The recommendation against the use of tofacitinib among hospitalized patients with COVID-19 was mainly due to the drug's safety issues. Although the study showed that there was benefit in reducing death or respiratory failure, particularly among patients on low-flow supplemental oxygen, patients who received tofacitinib were three times more likely to experience adverse events leading to treatment discontinuation. Furthermore, the US FDA and Health Canada issued safety alert on the use of tofacitinib due to increased risk of serious cardiovascular-related events (heart attack, stroke), cancer (lymphoma, lung cancer), thrombosis and death.

Among patients with COVID-19, should leronlimab be used for treatment?

As of 28 October 2021

RECOMMENDATION

We suggest against the use of leronlimab as treatment for COVID-19. (*Very low certainty of evidence, Weak recommendation*)

KEY FINDINGS

Two (2) unpublished randomized controlled trial (RCTs) [78-82] investigated the effect of leronlimab compared to placebo as treatment for patients with COVID-19. There was no significant benefit in the use of leronlimab in reducing mortality, length of hospital stay, resolution of clinical symptoms, and time to symptom resolution. There was no significant difference in serious adverse events and adverse events. The overall quality of evidence was rated very low because of serious risk of bias and very serious imprecision.

CONSENSUS ISSUES

The evidences of the review are now based on unpublished randomized controlled trials (previously from case series and a case study), however, similar to the previous consensus issue, the panel agreed that further trials are needed to recommend the use of leronlimab for the treatment of COVID-19.

Among patients with COVID-19, should infliximab be used for treatment?

As of 12 October 2021

RECOMMENDATION

We suggest against the use of infliximab among patients with COVID-19 infection (*Very low certainty of evidence, Weak recommendation*)

KEY FINDINGS

There is one (1) pre-print randomized controlled trial (RCT) [83] that reported the effect of infliximab compared to standard of care as treatment for patients with COVID-19. Results of the RCT showed no significant difference in mortality, hospital discharge in 28 days, and time to improvement in the WHO clinical progression scale. The RCT suffered from lack of allocation concealment, blinding, and poor comparability of treatment groups. It was also terminated prematurely due to futility. The very serious risk of bias and imprecision contributed to the downgrading of evidence to a very low certainty of evidence.

CONSENSUS ISSUES

The recommendation against the use of infliximab among patients with COVID-19 was made based on a small, pre-print randomized controlled trial (n = 63), which did not show any benefit on mortality, hospital discharge within 28 days, and time to clinical improvement. The panel also emphasized that the infliximab arm of the study was prematurely terminated due to futility on interim analysis, explaining its small sample size.

Among patients with COVID-19, should bevacizumab be used for treatment?

As of 12 October 2021

RECOMMENDATION

We suggest against the use of bevacizumab as treatment for COVID-19. (*Very low certainty of evidence, Weak recommendation*)

KEY FINDINGS

One (1) small non-randomized clinical trial [84] investigated on the potential use of bevacizumab for treatment of COVID-19. The study showed that bevacizumab led to net potential harm with minimal potential benefit only in terms of duration of oxygen support. Evidence was inconclusive for hospital discharge. Adverse events in the bevacizumab group were elevation of liver enzymes (30%), reduced hemoglobin (19%), decreased platelet counts (15%), elevation of blood pressure (11%), elevated blood urea nitrogen (7%), and sepsis (7%).

CONSENSUS ISSUES

Currently available evidence came from one small non-randomized clinical trial with only 27 patients in the treatment group and 26 patients in the external control cohort.

Administration of bevacizumab led to net potential harm with unclear potential benefit in only 1 non-critical outcome (i.e., duration of oxygen support). Its cost and potential negative effect on equity, since the drug is also being used as treatment for lung cancer, were also considered by the consensus panel. At present, bevacizumab is not readily accessible due to its limited availability and large cost. Use of bevacizumab among COVID-19 patients when its benefit is still unclear, may lead to even greater difficulty for lung cancer patients to gain access to this drug.

Among patients with COVID-19, should ivermectin be used for treatment?

As of 6 December 2021

RECOMMENDATION

We recommend against the use of ivermectin for the treatment of patients with COVID-19 of any severity. (*Very low certainty of evidence, Strong recommendation*)

We suggest against the use of ivermectin combined with doxycycline for the treatment of patients with COVID-19. (*Very low certainty of evidence, Weak recommendation*)

We suggest against the use of ivermectin combined with doxycycline for the treatment of patients with COVID-19. (*Very low certainty of evidence, Weak recommendation*)

KEY FINDINGS

- There are 16 randomized controlled trials (RCTs) [85-113] that investigated the effects of ivermectin as treatment for patients with COVID-19. No significant overall mortality benefit was found. Subgroup analysis by disease severity also showed no significant benefit for low dose ivermectin. We cannot estimate the risk of benefit or harm associated with high dose ivermectin because no deaths were reported among the three included RCTs. Sensitivity analysis revealed that publication status and study quality did not influence our estimates.
- Treatment with ivermectin was not significantly associated with clinical deterioration, need for mechanical ventilation, clinical improvement, reduction in hospital length of stay, time to symptom resolution, and virologic clearance. The risk for serious and non-serious adverse events was not significantly different among patients who received ivermectin. Our results agree with a recent Cochrane systematic review done in May 2021.
- These results must be interpreted in the context of very low certainty of evidence. The certainty of evidence was downgraded due to varying degrees of risk of bias in most studies, inconsistency, and imprecision in several critical outcomes.

CONSENSUS ISSUES

The review showed that ivermectin has no clear benefit for mortality and all other outcomes for patients with different disease severity, hence the panel made a general recommendation for all COVID-19 patients regardless of severity (mild, moderate, severe or critical).

This update provided additional evidence ivermectin did not differ significantly from placebo in terms of critical outcomes in the treatment of COVID-19. Hence, given the ongoing misuse and abuse of the drug, the panel unanimously voted for a strong recommendation against the use of ivermectin. Other considerations included issues on the pharmacologic property of the drug, given that the drug is registered for veterinary use, the need for higher doses, and concerns regarding adverse events. The panel also considered the issue on health equity wherein other medications for COVID-19 are available, hence resources should be allocated to these more effective

and efficacious treatment with clear benefits. There are still a number of ongoing trials, including a local one, which will be considered once data is available.

There is no new evidence for ivermectin combined with doxycycline available, hence, no update was done and the previous recommendations were retained.

Among patients with COVID-19, should artesunate (artemisinin) be used for treatment?

As of 18 November 2021

RECOMMENDATION

We suggest against the use of artesunate, artemisinin or pyronaridine tetraphosphate + artesunate in the treatment of COVID-19. (*Very low certainty of evidence, Weak recommendation*)

KEY FINDINGS

There are three (3) randomized controlled trials (RCTs) [114-116] that investigate the effect of artesunate (artemisinin) compared to varying standards of care or placebo as treatment for patients with COVID-19. Artesunate showed no significant difference in mortality, clinical deterioration, improvement in chest CT scan or X-ray, time to virologic clearance, and mild and moderate-to-severe adverse effects. Two (2) studies reported shorter time to clinical improvement for artesunate. The overall certainty of evidence was rated very low due to very serious imprecision, serious risk of bias, and in some outcomes, inconsistency and indirectness.

CONSENSUS ISSUES

There may be evidence on the in vitro effect of the drug. Despite being a new drug, we have a similar drug being used for malaria (Artemether-lumefantrine). However, given that the evidences available are very limited and the effect of drug is not yet clear, the panel unanimously suggests against the use of it. We need to wait for ongoing studies and more evidence for its use and effect on COVID 19.

Among patients with COVID-19, should colchicine be used for treatment?

As of 8 November 2021

RECOMMENDATION

We suggest against the use of colchicine in the treatment of COVID-19 patients.
(*Low certainty of evidence, Weak recommendation*)

KEY FINDINGS

- Five (5) randomized controlled trials (RCTs) [117-122] investigated the effect of colchicine compared to standard of care as treatment for patients with COVID-19. Colchicine showed net potential harm (significant increase in adverse events) with no significant benefit in all-cause mortality, need for mechanical ventilation, clinical improvement (defined as hospital discharge) within 28 days, need for hospitalization, and need for hemodialysis or hemofiltration.
- One large RCT reported that colchicine did not significantly shorten duration of hospitalization, while 2 smaller RCTs reported significantly shorter duration of hospitalization (data could not be pooled due to inadequate data provided). All studies had risk of bias issues as there were concerns in allocation concealment, blinding, attrition and selective reporting of outcome. The serious risk of bias and issues with inconsistency and imprecision in one critical outcome contributed to the downgrading of evidence to very low certainty of evidence.

CONSENSUS ISSUES

The addition of a large multicenter trial (RECOVERY trial) still showed that colchicine led to net potential harm (significant increase in adverse events) with no significant benefit in terms of all-cause mortality, clinical improvement (defined as hospital discharge) within 28 days, need for hospitalization, need for mechanical ventilation, and need for hemodialysis or hemofiltration. Hence, the consensus panel decided to maintain the previous recommendation against the use of colchicine in patients with COVID-19, regardless of hospitalization status.

Among patients with COVID-19, should interferon be used for treatment?

As of 6 December 2021

RECOMMENDATION

We recommend against the use of interferon in the treatment of COVID-19 patients. *(Very low certainty of evidence, Strong recommendation)*

KEY FINDINGS

Nine (9) published randomized controlled clinical trials (RCTs) [123-131] (N = 5,957) investigated the efficacy and safety of interferon in the treatment of COVID-19 compared to standard of care and/or placebo. Results showed significant benefit on viral negative conversion among patients given interferon, however, there was inconclusive evidence in terms of the critical outcomes such as all-cause mortality, clinical improvement, need for mechanical ventilation, progression to severe disease, ICU admission, adverse events, and serious adverse events.

CONSENSUS ISSUES

Recent review included more participants (close to 6,000) yet, interferon still did not show any clear benefit for all-cause mortality and other critical outcomes. There is a need to emphasize the potential harm and side effects of the drug as well as its high cost.

Among patients with COVID-19, should fluvoxamine be used for treatment?

As of 8 November 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of fluvoxamine among COVID-19. *(Low certainty of evidence)*

KEY FINDINGS

Two (2) published randomized controlled trials (RCTs) [187-188] (N = 1,649) investigated on the effectiveness of fluvoxamine compared to placebo among confirmed symptomatic non-hospitalized COVID-19 patients compared. Results showed significant reduction in emergency room visits and the need for hospitalization. There was inconclusive evidence in terms of other critical outcomes such as all-cause mortality, clinical deterioration, viral negative conversion, adverse events, and serious adverse events.

CONSENSUS ISSUES

Current evidence showed that although fluvoxamine appeared to reduce the need for emergency room visit or hospitalization, there was inconclusive evidence in terms of other critical outcomes such as all-cause mortality, clinical deterioration, adverse events, and serious adverse events. The sample size of the two randomized controlled trials may still be too small to reach a level of significance, precluding any

recommendation to be made. As of writing, there are 9 ongoing clinical trials, results of which may further elucidate on fluvoxamine's effectiveness in the treatment of COVID-19.

Among patients with COVID-19, should bamlanivimab in combination with etesevimab be used for treatment?

As of 12 October 2021

RECOMMENDATION

We suggest the use of bamlanivimab and etesevimab combination therapy as treatment for mild to moderate, non-hospitalized COVID-19 patients with at least 1 risk factor* for progression to severe disease. (*Low certainty of evidence, Weak recommendation*)

**Risk factors for severe COVID-19: age ≥ 65 years, body-mass index ≥ 35 kg/m², cardiovascular disease (including hypertension), chronic lung disease (including asthma), chronic metabolic disease (including diabetes), chronic kidney disease (including receipt of dialysis), chronic liver disease, and immunocompromised conditions*

KEY FINDINGS

The evidence on the use of bamlanivimab + etesevimab combination comes from two (2) randomized controlled trials (RCTs) [154-155] among non-hospitalized patients with COVID-19. Both studies found that there was a significant reduction in COVID-19 related hospitalizations, all-cause mortalities, symptom resolution, and reduction in viral load among the participants who received the bamlanivimab + etesevimab cocktail compared to placebo. There was no significant difference in adverse events between the two groups.

CONSENSUS ISSUES

The panel favoured the use of bamlanivimab and etesevimab among non-hospitalized COVID-19 patients who are at risk for severe disease, based on the results of 2 randomized controlled trials that showed net potential benefit in terms of COVID-19 related hospitalizations and all-cause mortality, reduction in total symptom score, and number of days to symptom resolution, with no significant difference in terms of adverse events. Concern regarding the drug's effectivity against variants was raised by one of the panelists. As of writing, the drug has no emergency use authorization from the Philippine FDA, thus may only be used in the context of clinical trials.

Among patients with COVID-19, should casirivimab-imdevimab be used for treatment?

As of 20 December 2021

RECOMMENDATION

We suggest the use of casirivimab-imdevimab as treatment for symptomatic, non-hospitalized patients with at least 1 risk factor* for severe COVID-19. (*Moderate certainty of evidence, Weak recommendation*)

*Risk factors: age >50 years, obesity, cardiovascular disease (including hypertension), chronic lung disease (including asthma), chronic metabolic disease (including diabetes), chronic kidney disease (including receipt of dialysis), chronic liver disease, and immunocompromised conditions

RECOMMENDATION

We recommend against casirivimab-imdevimab as treatment for hospitalized COVID-19 patients. (*Low certainty of evidence, Strong recommendation*)

RECOMMENDATION

There is insufficient evidence to recommend casirivimab-imdevimab as treatment for asymptomatic COVID-19 patients. (*Low certainty of evidence*)

KEY FINDINGS

Six (6) RCTs [156-161] evaluated the efficacy of casirivimab-imdevimab cocktail as treatment for patients with COVID-19. Among non-hospitalized patients given casirivimab-imdevimab, there was a significant reduction in combined end-point of need for invasive mechanical ventilation or death, COVID-19-related medically-assisted visits, duration of symptoms, and serious adverse events. However, evidence was largely inconclusive for hospitalized patients and asymptomatic, non-hospitalized patients. There was trend towards benefit in clinical recovery among hospitalized seronegative patients (negative for serum SARS-CoV-2 antibodies at baseline), but not for seropositive patients (positive for serum SARS-CoV-2 antibodies at baseline). There was no benefit in all-cause mortality regardless of hospitalization status.

CONSENSUS ISSUES

Addition of 2 new pre-print randomized controlled trials (RCTs) still showed that casirivimab and imdevimab cocktail is beneficial for symptomatic, non-hospitalized COVID-19 patients with at least 1 risk factor for progression to severe disease. Although there was moderate certainty of evidence, the recommendation was weak due to inconclusive evidence on all-cause mortality, large cost, need for emergency room visit for drug administration and close monitoring and equity considerations. At present, the drug is only available in private tertiary hospitals.

The evidence remained inconclusive for hospitalized COVID-19 patients. The previous update included 1 pre-print RCT (RECOVERY trial), which on subgroup analysis showed that casirivimab-imdevimab appeared to have benefit in terms of all-cause mortality, need for mechanical ventilation, and clinical recovery among hospitalized patients who were seronegative at baseline (negative for serum SARS-CoV-2

antibodies). However, addition of 1 (one) new pre-print RCT showed trend towards benefit in clinical recovery only. The panel maintained the previous recommendation against the use of casirivimab-imdevimab cocktail among hospitalized COVID-19 patients because of uncertain balance of effects in this specific subset of patients and both RCTs are still pre-print. As of writing, there are 8 ongoing clinical trials on casirivimab-imdevimab, 2 of which are among hospitalized patients. Results of these trials may further elucidate on the cocktail's effectiveness in the treatment of hospitalized COVID-19 patients.

Lastly, the only available study for asymptomatic, non-hospitalized COVID-19 patients is a small pre-print RCT with low certainty of evidence. The panel deemed that current evidence is still insufficient to make any recommendations for this subset of patients.

Among patients with COVID-19, should regdanvimab be used for treatment?

As of 20 December 2021

RECOMMENDATION

We suggest against the use of regdanvimab for the treatment of mild to moderate COVID-19. (*Very low certainty of evidence, Weak recommendation*)

KEY FINDINGS

Two (2) randomized controlled trials (RCTs) [162-163] evaluated the efficacy of regdanvimab as treatment for patients with COVID-19. The overall certainty of evidence was very low because of serious risk of bias, imprecision, and high probability of publication bias. Among patients with mild to moderate COVID-19 infection, regdanvimab showed minimal potential benefit in terms of clinical recovery at Day 14 and Day 28. There was inconclusive evidence in terms of other critical outcomes such as all-cause mortality, need for hospitalization, and oxygen therapy requirement. There was no significant difference in adverse events between patients treated with regdanvimab compared to placebo.

CONSENSUS ISSUES

Very limited evidence (2 small randomized controlled trials, one of which is still a pre-print) support the use of regdanvimab in treatment of mild to moderate COVID-19. Although regdanvimab showed potential benefit in terms of clinical recovery at Day 14 and Day 28, the evidence was inconclusive for other critical outcomes such as all-cause mortality, need for hospitalization, or clinical deterioration. Clinical recovery among patients with moderate to severe symptoms at baseline was defined as improvement of symptoms and not necessarily complete resolution of symptoms. Concern regarding the effect of possible publication bias on subjective outcomes such as clinical recovery was raised, since currently available studies were both funded by regdanvimab's manufacturing company. Hence, the panel suggested against use of regdanvimab considering its limited evidence, potentially large cost, and availability of treatment alternatives with more robust evidence that has shown benefit on more objective outcomes (e.g., all-cause mortality and need for hospitalization) among patients with mild to moderate COVID-19.

Among patients with COVID-19, should convalescent plasma be used for treatment?

As of 18 November 2021

RECOMMENDATION

We recommend against the use of convalescent plasma in patients with COVID-19 infection. (*Moderate certainty of evidence, Strong recommendation*)

KEY FINDINGS

- There are 22 randomized controlled trials (RCTs) [132-153] that compared the effect of convalescent plasma therapy against placebo and/or standard of care among confirmed COVID-19 patients. Pooled estimates of critical patient outcomes (i.e., all-cause mortality) on the use of convalescent plasma were not statistically significant. Exploratory subgroup analysis done for all-cause mortality by age, severity of disease, and timing of administration did not show any statistically significant benefit except for the subgroup of early administration (defined as within 3 days of hospitalization) of high-level titers of convalescent plasma (RR 0.42, 95% CI 0.21, 0.86; $I^2 = 0\%$; 4 RCTS, $n = 376$).
- The incidence of adverse and serious adverse events (e.g., transfusion-related events) were not significantly different between the convalescent plasma group and those given standard care or placebo. The over-all certainty of evidence was retained as moderate.

CONSENSUS ISSUES

Serious adverse events and progression to respiratory distress/respiratory failure were re-rated and still considered as critical outcomes. However, the panel unanimously voted to give more value to the effect of convalescent plasma on all-cause mortality, clinical improvement, and need for invasive ventilation and ICU admission, hence, the over-all certainty of evidence was retained as moderate.

Recommendation remains strong given that convalescent plasma is no different from placebo in terms of efficacy, clinical outcomes, and harm; yet there is a lot to consider in terms of cost, value preferences, equity, and feasibility.

Should intravenous immunoglobulin (IVIg) be used for the treatment of COVID-19?

As of 18 May 2021

RECOMMENDATION

We suggest against the use of intravenous immunoglobulin as treatment for moderate to severe COVID-19. (*Very low certainty of evidence, Weak recommendation*)

KEY FINDINGS

Four (4) RCTs [164-167] that were of low to very low quality found that the use of intravenous immunoglobulin (IVIg) did not significantly reduce mortality or risk of mechanical ventilation among patients with moderate to severe COVID-19. However, one very low quality RCT found higher incidence of virologic clearance as assessed by negative RT PCR with the use of IVIg. There was also no significant increase risk of adverse events with its use in COVID-19.

CONSENSUS ISSUES

A conditional recommendation was made while waiting for the results of the 31 ongoing trials.

Should mesenchymal stem cell therapy be used for the treatment of COVID-19?

As of 29 May 2021

RECOMMENDATION

There is insufficient evidence to recommend using umbilical cord-derived mesenchymal stem cell therapy among adults with severe COVID-19 (PaO₂/FiO₂ ratio ≤ 300 mmHg). (*Very low certainty of evidence*)

KEY FINDINGS

We found 3 randomized controlled trials [168-170] evaluating the effectiveness and safety of mesenchymal stem cell (MSC) therapy in COVID-19 treatment. One RCT was at high risk of bias while the remaining 2 RCTs were of moderate quality. Validity issues included unclear allocation concealment, lack of blinding and incomplete outcome reporting. Based on very low certainty of evidence, umbilical cord-derived MSC (UC-MSC) therapy reduces mortality and hastens clinical improvement in adults with severe COVID-19. Limited by small sample sizes, adverse events were not significantly different between MSC and control groups.

CONSENSUS ISSUES

Although the current available evidence shows benefit in terms of mortality and time to clinical improvement, no recommendation was made due to the small sample size of the included studies. Further, more patients in the mesenchymal stem cell (MSC) therapy arm received co-interventions (i.e., remdesivir and corticosteroids) compared to the control arm which may have overestimated the true effect of MSC.

Among patients with COVID-19, should inhaled corticosteroids be used for treatment?

As of 18 November 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of inhaled corticosteroids in treatment of non-hospitalized COVID-19 patients. *(Very low certainty of evidence)*

KEY FINDINGS

Four (4) randomized controlled trials (RCTs) [171-174] investigated the effectiveness of inhaled corticosteroids for the treatment of COVID-19. Use of inhaled corticosteroids (e.g., budesonide and ciclesonide) among non-hospitalized patients with mild to moderate COVID-19 demonstrated benefit only for subjective outcomes such as self-reported clinical recovery or improvement at day 14 and day 28/30 but not for objective outcomes such as mortality, need for hospitalization, emergency room visit, and duration of hospitalization. There was no significant difference in serious adverse events and adverse events.

CONSENSUS ISSUES

Current evidence shows that inhaled corticosteroids (i.e., budesonide and ciclesonide) had benefit for subjective outcomes such as self-reported recovery on day 14 and day 28/30. However, evidence was inconclusive in terms of objective outcomes such as mortality, emergency room visit, need for hospitalization, duration of hospitalization, and clinical improvement (as evaluated by an assessor). Self-reported recovery may be biased since participants are not blinded, hence a more objective marker of clinical status such as oxygen saturation is more reliable. Only 1 study explicitly excluded patients with asthma or chronic obstructive pulmonary disease (COPD), and the perceived positive effect may have been due to improvement of asthma or COPD and not COVID infection. As of writing, there are 13 ongoing clinical trials, results of which may further elucidate on the effectiveness of inhaled corticosteroids in the treatment of COVID-19.

Should steam inhalation be used for the treatment of COVID-19?

As of 12 March 2021

RECOMMENDATION

We recommend against the use of steam inhalation alone in the treatment of COVID-19. *(Very low certainty of evidence, Strong recommendation)*

KEY FINDINGS

Based on a single arm observational study [175] with high risk of bias, there is currently only very low certainty of evidence showing the possible benefit of steam inhalation in the prevention of developing symptomatic COVID-19 among exposed healthy individuals and reducing symptoms and number of days to negative SARS-COV-2 RT-PCR test of COVID-19 confirmed individuals. Meanwhile, there is indirect evidence

highlighting the significant adverse effects of steam inhalation among individuals using it for symptomatic relief from the colds.

CONSENSUS ISSUES

The panel strongly recommended against the use of steam inhalation as treatment for COVID-19, despite the very low quality of evidence, because it was recognized that the potential for harm outweighs the benefit.

Should virgin coconut oil (VCO) be used in the treatment of patients with COVID-19 infection?

As of 20 February 2021

RECOMMENDATION

There is no evidence to recommend the use of VCO as treatment among patients with COVID-19 infection.

KEY FINDINGS

To date, there is no available completed clinical trial directly investigating effectiveness or safety of virgin coconut oil as an adjunct treatment for COVID-19 patients of any age or disease severity. Indirect evidence was noted from two studies of VCO on other viruses such as human immunodeficiency viruses. [178-179]

CONSENSUS ISSUES

It is important to note that this recommendation is strictly for the use of VCO in the treatment of COVID-19 infection and should not be confused with another living recommendation pertaining to the use of VCO in the adjunctive treatment of COVID-19.

The ongoing clinical trial for VCO is not yet registered with the local Food and Drug Administration, hence, the evidence that will be generated cannot be used as an additional indication for the marketing purpose of VCO.

Among patients with COVID-19, should Lianhua be used for treatment?

As of 06 December 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of Lianhua in the treatment of patients with non-severe COVID-19. (Very low certainty of evidence)

KEY FINDINGS

Five (5) randomized controlled trials investigated the effect of Lianhua compared to standard of care as treatment for patients with COVID-19. Lianhua showed significant benefit in preventing clinical deterioration or progression to severe disease among patients with non-severe COVID 19. There was no significant benefit in mortality, and day-14 improvement in fever, cough and fatigue. There was no significant difference

in adverse events or serious adverse events between the Lianhua and control group. The overall certainty of evidence was rated very low due to very serious risk of bias and serious imprecision in some critical outcomes.

CONSENSUS ISSUES

Although the recent review showed some benefit on the symptomatic treatment (clinical deterioration), the panel considered that the uncertainties on the quality of the evidence outweigh the trend in benefit. First, the subset of patients (percentage of mild and moderate cases) were not clearly stated in the included studies. This may actually affect the trend towards benefit since patients with mild COVID are expected to improve and have shorter time to recovery. Second, the definition of outcomes, particularly total symptom recovery may be too lax as it accounted only for at least one of the major symptoms. The panel also considered that since this is a regulated drug, there are uncertainties about the reported harm (expected adverse effects from ephedra such as hypertension and tachycardia was not assessed in the study procedures or reported in the results) and serious adverse events (unclear if the reported events were transient nor how severe the cases were). There are also inconsistencies in the direction of the clinical outcomes. While there is some benefit seen in reducing clinical deterioration, no definite benefit was seen in terms of clinical improvement of individual symptoms.

Should famotidine be used for the treatment of COVID-19?

As of 30 May 2021

RECOMMENDATION

We suggest against the use of famotidine in the treatment of COVID-19. (*Very low certainty of evidence, Weak recommendation*)

KEY FINDINGS

Very low-quality evidence from seven retrospective cohort studies [180-186] was found on the use of famotidine for the treatment of COVID-19. There was no significant reduction in the risk of mortality, mechanical ventilation, and composite outcome of mortality or mechanical ventilation. However, there was a significant increase in the risk of the composite outcome of mortality, mechanical ventilation, and intensive care unit admission. Adverse events were not examined in these retrospective cohort studies.

Should ibuprofen be used in the treatment of patients with COVID-19 infection?

As of 5 March 2021

RECOMMENDATION

We recommend against the use of ibuprofen as treatment among patients with COVID-19 infection. (*Very low certainty of evidence, Strong recommendation*)

KEY FINDINGS

Currently, there are no randomized controlled trials that assessed the efficacy of ibuprofen as a treatment for COVID-19. Based on the two observational studies (one retrospective cohort and one case series) [176-177] that have not been peer-reviewed, there is limited evidence that ibuprofen treatment could improve COVID-19 symptoms and outcomes.

CONSENSUS ISSUES

It is important to note that this recommendation is strictly for the use of ibuprofen in the treatment of COVID-19 infection and should not be confused with another living recommendation pertaining to the effect of the concurrent use of ibuprofen with other COVID-19 outcomes.

Although direct evidence of very low quality suggests a trend towards benefit, there was indirect evidence of possible harm. The benefit that was seen from the first study might not be solely because of ibuprofen given that other patients were given other medications. The severity of COVID-19 infection was not also specified in the first study and it was assumed that it included patients of all severity (i.e., mild, moderate, severe) since the information were taken from electronic health records of different hospitals.

Evidence and Recommendations for the Critical Care and Respiratory Management of COVID-19

Should intravenous corticosteroids be used in COVID-19?

As of 03 January 2022

RECOMMENDATION

We recommend the use of dexamethasone for up to 10 days among patients with severe and critical COVID-19. *(Moderate certainty of evidence; Strong recommendation)*

We recommend the use of 6mg to 12mg per day of dexamethasone among patients with severe and critical COVID-19. *(Moderate certainty of evidence; Strong recommendation)*

We suggest that steroid therapy be initiated as soon as diagnosed or categorized as severe and critical COVID-19. *(Very low certainty of evidence; Weak recommendation)*

RECOMMENDATION

We recommend against the use of corticosteroids among mild and moderate (non-oxygen requiring) COVID-19 patients. *(Moderate certainty of evidence; Strong recommendation)*

KEY FINDINGS

- Fourteen randomized controlled trials (RCTs) [1-14] provided data on the effect of different intravenous (IV) corticosteroids versus placebo or standard of care on all-cause mortality in severe and critical COVID-19 patients. Compared to placebo, only dexamethasone showed a statistically significant reduction in the risk of mortality. However, patients in this group had significantly longer duration of hospital stay. In terms of adverse events, no significant difference was found between the IV corticosteroids and control groups. No significant benefit on 28-day mortality and a tendency towards harm were observed when dexamethasone was given to COVID-19 patients who did not require oxygen therapy.
- In comparing 12mg and 6mg dosing of dexamethasone, no significant differences were found in terms of all-cause mortality and life support-free days while marginal benefits in ventilator-free days, cardiovascular support-free days, and renal replacement therapy-free days were observed in the 12mg group. No significant difference in adverse events were observed between the two dosing regimens.[15]
- Seven retrospective cohort studies [16-22] evaluated the effect in the timing of administration of different corticosteroids among severe and critical COVID-19 patients. Significant benefit was found only when systemic corticosteroids were started early within 24 hours of admission compared to non-early initiation beyond 24 hours. Mechanical ventilation was likewise significantly reduced when systemic corticosteroids were initiated within 24 hours of admission.

CONSENSUS ISSUES

Dexamethasone has a better pharmacokinetic profile (i.e., longer acting than hydrocortisone and methylprednisolone) and better anti-inflammatory effect as compared to other steroids with less corticoid effects (e.g., less water retention). Low-dose steroids (i.e., 6mg of dexamethasone) are preferred by physicians. As for other corticosteroids such as methylprednisolone and hydrocortisone, there is insufficient evidence to recommend its use in patients with COVID-19 infection who require supplemental oxygenation.

Should anticoagulation be used in treating patients diagnosed with COVID-19?

As of 26 October 2021

RECOMMENDATION

We recommend the use of prophylactic over therapeutic dose anticoagulation among hospitalized patients with moderate, severe or critical COVID-19 disease unless there are any contraindications. (*Low certainty of evidence; Strong recommendation*)

We recommend the use of standard dose prophylactic anticoagulation over intermediate dose prophylactic anticoagulation among hospitalized patients with COVID-19 disease unless there are any contraindications. (*Moderate certainty of evidence; Strong recommendation*)

KEY FINDINGS

- Eight randomized controlled trials (RCTs) [23-30] were included in this review to compare therapeutic versus prophylactic anticoagulation and to compare intermediate dose versus standard dose prophylactic anticoagulation in COVID-19 patients. Quality of evidence ranged from very low to moderate due to issues regarding risk of bias, and imprecision due to underpowered population, and wide confidence intervals.
- Comparing therapeutic and prophylactic anticoagulation (AC) [23-28], there was no significant difference over-all in terms of mortality and organ support-free days among critically ill and stable patients. There was a significantly lower over-all incidence of venous thromboembolism (VTE) for therapeutic anticoagulation but there was significantly higher risk for major and minor bleeding for therapeutic anticoagulation considering all populations. However, no significant difference was seen in the incidence of VTE among critically ill patients. There was no significant difference in major bleeding among subgroups of critically ill and stable patients but the trend pointed to a higher incidence among those undergoing therapeutic anticoagulation. For minor bleeding, there was a significantly higher observed incidence among critically ill patients.
- For the comparison of intermediate versus standard prophylactic anticoagulation dose among severely ill patients [29,30], there was no statistically significant difference in terms of mortality and incidence of venous thromboembolism. There was also no significant difference in terms of major and minor bleeding, but the trend showed higher incidence when given intermediate prophylactic dose anticoagulation.

CONSENSUS ISSUES

A strong recommendation in favor of prophylactic dose anticoagulation was unanimously given despite the low certainty of evidence to give emphasis on preventing harm, as the risk of bleeding was significantly higher with therapeutic dose anticoagulation. It was further emphasized that the duration of anticoagulation, which was not directly addressed by the studies included in this review, should be individualized based on the patient's thromboembolic and bleeding risks.

Should empiric antimicrobial coverage be given to patients with severe and critical COVID-19?

As of 15 April 2021

RECOMMENDATION

We recommend against the use of antibiotics in patients with severe and critical COVID-19 infection, unless with suspicion of secondary bacterial co-infection. For patients on empiric antibiotics, they should be assessed daily for the need for discontinuation, continuation or de-escalation based on clinical and laboratory parameters. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

Very low quality of evidence from one retrospective observational report was found on the use of empiric antimicrobials for COVID-19.[31] Early administered antibiotics did not significantly impact mortality or delayed hospital acquired infections in critically ill patients with COVID-19.

CONSENSUS ISSUES

Despite the low quality of evidence available, the panel voted for a strong recommendation against the routine use of antibiotics due to concerns on antimicrobial resistance.

Should hemoperfusion be used in patients diagnosed with COVID-19?

As of 01 December 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of hemoperfusion among patients diagnosed with COVID-19. (*Very low certainty of evidence*)

KEY FINDINGS

Presently, no randomized clinical trials have been published to provide data on the use of hemoperfusion in patients with COVID-19. Two retrospective cohort studies [32,33] showed statistically significant benefit in terms of mortality. Improvement in some clinical parameters such as respiratory rate, heart rate, and peripheral oxygen saturation, and selected markers such as C-reactive protein, erythrocyte sedimentation rate, and serum ferritin levels were also significantly observed among COVID-19 patients who received hemoperfusion. However, the benefits in clinical and laboratory parameters failed to translate to more clinically important outcomes such as decrease in the length of hospital and intensive care unit stay. Moreover, the studies were of very low certainty of evidence being non-clinical trials.

CONSENSUS ISSUES

No randomized controlled trials were available for review and the observational studies included did not adjust for confounders. With the available sparse evidence, the use of hemoperfusion can be suggested in COVID-19 patients with clinical deterioration despite standard medical therapy (including Tocilizumab). However, there is no consensus among experts in the panel on the use of hemoperfusion in COVID-19. Clinical trials are needed to be able to identify and evaluate the balance of benefits, harm, and cost for an invasive mode of treatment such as hemoperfusion especially since immunotherapy has been made available for the management of COVID-19.

Should a conservative fluid management strategy be used in mechanically ventilated adult COVID-19 patients?

As of 05 March 2021

RECOMMENDATION

We suggest the use of conservative fluid management rather than liberal fluid management strategy in mechanically ventilated adult COVID-19 patients with acute respiratory distress syndrome who have been adequately resuscitated*. (*Low quality of evidence; Weak recommendation*)

**without tissue hypoperfusion and fluid responsiveness*

KEY FINDINGS

Analysis of four randomized controlled trials [34-37], which included 1,106 hemodynamically stable mechanically ventilated patients with acute respiratory distress showed a trend towards decreased risk of mortality in the conservative fluid management strategy group. Furthermore, the use of conservative fluid management strategy resulted in significantly increased ventilator-free days, significant decrease in the length of intensive care unit stay and duration of mechanical ventilation. In terms of adverse events, the use of conservative fluid management strategy showed no significant difference in renal failure-free days and need for renal replacement therapy. The overall quality of studies included in this analysis was found to have moderate risk of bias with common issues on selection, performance, and reporting bias.

CONSENSUS ISSUES

The recommendation of initiating a conservative fluid management strategy should be employed in adequately resuscitated and hemodynamically stable mechanically ventilated patients with acute respiratory distress syndrome. Studies included in the evidence review for this clinical question excluded patients with hemodynamic instability or those with more than minimal requirement for vasopressors for septic shock. It should be emphasized that fluid resuscitation for septic shock should be prioritized in order to optimize adequate perfusion and hemodynamic stability. Fluid responsiveness should also be adequately assessed as conservative fluid management strategy must not be initiated until patient is deemed fluid or volume unresponsive.

Should side lying position be used in patients with severe to critical COVID-19?

Should self-proning be used in non-intubated patients with severe COVID-19?

As of 26 October 2021

RECOMMENDATION

We suggest self-proning position in non-intubated patients with severe and critical COVID-19. (*Very low certainty of evidence; Weak recommendation*)

RECOMMENDATION

There is insufficient evidence to recommend the use of side lying in non-intubated patients with severe and critical COVID-19. (*Very low certainty of evidence*)

KEY FINDINGS

- Among non-intubated severe patients with COVID 19 with oxygen saturation of at least 90% and oxygen requirement of less than 6 liters per minute, pooled results of 4 randomized controlled trials [38-41] showed no difference in the duration of proning and outcomes such as mortality, need for intubation and the need for intensive care.
- A case series [42] of five patients with COVID-19 associated ARDS showed that side lying accompanied with Positive End Expiratory Pressure (PEEP) titration provided a statistically significant benefit by decreasing the incidence of overdistension and lung collapse. Adverse events were not reported.
- The certainty of evidence for both side lying and proning are very low due to serious risk of bias, substantial heterogeneity and imprecision.

CONSENSUS ISSUES

Self-proning in non-intubated patients with severe and critical COVID-19 was suggested despite the lack of significant benefit in terms of mortality, need for endotracheal intubation, and need for intensive care unit stay on the basis that self-proning may still offer some benefit on improving oxygenation citing theoretical effect and personal experience, and taking into consideration the existing recommendations made by various international guidelines as well.

There was very limited evidence to recommend side lying for the same subset of patients, although it was recognized that there may be some benefit. This intervention will depend on the physician's prerogative in situations where self-proning is not possible. Potential harms such as patient discomfort and risk of accidental removal of peripheral lines and endotracheal tubes and the need for additional healthcare workers to perform proning in sedated and mechanically ventilated patients should be considered for both patient and health care worker in attempting this intervention.

Should high flow nasal cannula be used for patients with COVID-19 and acute respiratory failure?

As of 01 December 2021

RECOMMENDATION

We suggest the use of high flow nasal cannula for patients with severe to critical COVID-19 who do not respond to conventional oxygen therapy (low flow nasal cannula/face mask). (*Low certainty of evidence; Weak recommendation*)

KEY FINDINGS

Two randomized controlled trials [43,44] comparing the use of high flow nasal cannula (HFNC) versus conventional oxygen therapy (COT) in COVID-19 patients with acute respiratory failure showed significant improvement of $\text{PaO}_2/\text{FiO}_2$ ratio among patients who received HFNC. However, no significant benefit was found in terms of 30-day mortality, length of hospital stay, length of intensive care unit stay, and eventual tracheal intubation and intensive care unit admission. Certainty of evidence was low because of unclear to high-risk of selection and detection bias, and imprecision in most of the critical outcomes.

CONSENSUS ISSUES

The use of high flow nasal cannula should only be considered when patients fail to respond to low flow nasal cannula or face mask. It is not intended to be the immediate first line respiratory support for COVID-19 patients. It was initially promoted due to its capability to deliver high oxygen concentration, particularly when coupled with the potential harm or risk of viral aerosolization with non-invasive ventilation. The comparison of the efficacy of high flow nasal cannula and non-invasive ventilation is discussed in a separate review.

Should non-invasive ventilation be used over high flow nasal cannula for patients with severe and critical COVID-19?

As of 03 January 2022

RECOMMENDATION

We suggest the use of either high flow nasal cannula or non-invasive positive pressure ventilation in COVID-19 patients with hypoxemic respiratory failure in the absence of any indication for emergent invasive mechanical ventilation. (*Low certainty of evidence; Weak recommendation*)

KEY FINDINGS

Two randomized controlled trials [45,46] were evaluated to compare the effect of non-invasive ventilation (NIV) and high flow nasal cannula (HFNC) oxygenation in improving clinical outcomes in COVID-19 patients with respiratory failure. Direct comparison of NIV in the form of helmet and face mask CPAP with HFNC in 218 COVID-19 patients with hypoxemia showed that reduction in mortality and need for endotracheal intubation were inconclusive. Certainty of evidence was low due to serious risk of bias and serious imprecision. Indirect mixed treatment comparison of NIV in the form of helmet CPAP and HFNC among COVID-19 patients with hypoxemia also showed no significant difference in terms of in-hospital mortality, need for mechanical ventilation, intensive care unit admission, and length of hospital stay.

CONSENSUS ISSUES

The risk of aerosolization using non-invasive ventilation was not discussed in the identified studies, but case series and reports have suggested minimal risk for health care workers. Standard operating procedure includes the use of filters in the expiratory limb tubing for non-invasive ventilation and use of face masks for patients on high flow nasal cannula. Physicians must be cognizant of the indications for intubation such as continued and progressive deterioration, and signs of respiratory failure.

Should lung protective ventilation, high PEEP and driving pressure-limited strategies be used in the management of adult patients with COVID-19-associated acute respiratory distress syndrome?

As of 19 February 2021

RECOMMENDATION

We suggest the use of a lung protective ventilation strategy (tidal volume 4-8 mL/kg predicted body weight and plateau pressure less than 30 cmH₂O) in patients with COVID-19 infection and ARDS. (*Very low certainty of evidence; Weak recommendation*)

RECOMMENDATION

There is insufficient evidence to recommend the use of a higher PEEP strategy. We suggest to individualize PEEP or employ a PEEP strategy based on respiratory mechanics (i.e., compliance) in patients with COVID-19 infection. (*Low certainty of evidence*)

There is insufficient evidence to recommend a driving pressure limited strategy in patients with COVID-19 infection. We suggest to keep the driving pressure ≤ 14 cmH₂O. (*Low certainty of evidence*)

KEY FINDINGS

There were no clinical trials evaluating the (1) effectiveness and harms of lung protective ventilation strategy and high PEEP strategies in patients with COVID-19-associated ARDS; (2) effectiveness and harms of higher PEEP versus lower PEEP strategies in conjunction with LTVV in patients with COVID-19-associated ARDS; and, (3) effectiveness and harms of a driving pressure-limited strategy in patients with COVID-19 and ARDS.

Lung Protective Ventilation

- Higher tidal volume was associated with the same or higher risk of 28-day mortality but there was no significant difference in ventilator-free days in multivariable models.[47] A smaller case study [48] also showed that barotrauma cases were observed to have significantly lower V_T than the controls. These studies provided very low quality of evidence for this component of the review question.

Higher PEEP and Low Tidal Volume Ventilation

- Higher PEEP was not associated with differences in 28-day mortality nor ventilator-free days in adults with COVID-19-related ARDS.[47] A case-control study showed that PEEP was not statistically significantly different between patients with and without barotrauma.[48] These studies provided low quality of evidence for this component of the review question.

Driving Pressure

- There was no significant difference in 28-day mortality, in-hospital mortality, mechanical ventilation-free days, hospital stay and barotrauma were observed.[49]

These studies provided low quality of evidence for this component of the review question.

CONSENSUS ISSUES

During the early stages of the disease, COVID-19 ARDS may not be the same as the usual ARDS. In COVID-19-related ARDS, compliance can be high or normal even in patients with very low PaO₂/FiO₂ ratios. In these cases, the lungs are not recruitable and the use of high levels of PEEP will not lead to better oxygenation.

As the respiratory mechanics (compliance) in COVID-19 ARDS are not uniformly correlated with the severity of the hypoxemia, it is best to individualize PEEP based on compliance and driving pressure rather than titrating PEEP based on the severity of the PaO₂/FiO₂ ratio. Thus, a higher PEEP strategy (using the high PEEP/ FiO₂ table) for moderate to severe ARDS is not recommended for all patients with COVID 19 infections due to possible complications such as barotrauma.

We suggest titrating lung volumes (tidal volume 4-8 ml/kg predicted body weight) and PEEP to maintain the plateau pressure less than 30 and to have the lowest driving pressure. A driving pressure of <14cm H₂O is recommended. The driving pressure is the difference between the plateau pressure and the PEEP, or tidal volume over the respiratory system compliance.

Should rapid sequence intubation or delayed sequence intubation be used for the management of COVID-19?

As of 30 June 2021

RECOMMENDATION

We suggest the use of rapid sequence intubation for COVID-19 patients to reduce infection among healthcare workers performing the procedure. (*Very low certainty of evidence; Weak recommendation*)

KEY FINDINGS

- There were no studies directly comparing rapid sequence intubation (RSI) and delayed sequence intubation (DSI). For each modality, there were very low quality evidence, which had limitations in study design and indirectness.
- Both RSI and DSI improved the post-intubation oxygen saturations of the patients. The study on RSI [50] reported that 10.4% of the patients died within 24 hours of intubation.
- On the other hand, the study on DSI [51] reported no deaths within an unspecified time period. The study on RSI reported cardiac arrest in 2% of the patients and pneumothorax in 5.9% of the patients post intubation. Conversely, the study on DSI reported that none of the patients had cardiac arrest post intubation within an unspecified time period. The study on RSI reported that there was no evidence of cross infection in the anesthesiologists who intubated the COVID-19 patients.

CONSENSUS ISSUES

There is no direct comparison between rapid sequence intubation (RSI) and delayed sequence intubation (DSI). RSI does not pertain to the timing of intubation but rather it pertains to the sequence of events.

RSI helps prevent the aerosolization of COVID-19 particles and is safer for the healthcare workers performing RSI. RSI is done rapidly and prevents further deterioration of patients who are in need of oxygen supplementation. However, there is a potential to develop hypoxemia among patients who have undergone RSI. RSI is commonly done locally except in patients with collapsed airways. Depending on the condition, RSI may be modified to suit the patients' needs. On the other hand, delayed sequence intubation (DSI) is generally much safer for patients because it ensures that the procedure is not conducted haphazardly (patients are pre-oxygenated and sedatives and neuromuscular blockers are administered). In addition, RSI is also recommended by the other societies except for agitated patients. Additional costs for anesthesia must be considered in performing RSI.

Should extracorporeal membrane oxygenation (ECMO) be used in the management of Acute Respiratory Distress Syndrome (ARDS) among COVID-19 patients?

As of 03 January 2022

RECOMMENDATION

We suggest the use of Extracorporeal Membrane Oxygenation (ECMO) for judiciously selected COVID-19 patients with severe Acute Respiratory Distress Syndrome (ARDS) based on the ELSO criteria. (*Very low certainty of evidence; Weak recommendation*)

KEY FINDINGS

Four cohort studies on the use of ECMO in adult patients with critical COVID-19 or COVID-19 ARDS were reviewed in this evidence update.[52-55] Pooled results from four studies showed inconclusive mortality benefit with the use ECMO versus standard of care. In one study [52], occurrence of procedure-related infection reported as a serious adverse event was significantly associated with ECMO therapy. The certainty of evidence is very low because of the risk of bias attributed to lack of randomized studies, inconsistency due to significant heterogeneity, and imprecision due to small study sample sizes.

CONSENSUS ISSUES

VV-ECMO is not indicated for all severe COVID-19 patients with ARDS. The general guidelines for the use of ECMO among non-COVID patients must be followed given that there are no guidelines for the use of ECMO among patients with COVID-19. Likewise, standard selection criteria, indications and contraindications should be followed in recommending the use of ECMO. The protocol for the treatment of severe ARDS must be followed. VV-ECMO is used once other modalities (e.g., proning, mechanical ventilation and lung protective strategies) did not work and the patients are qualified for ECMO.

There is a limited number of ECMO machines that are locally available. In addition, manipulation of this machine requires manpower, training and additional medical equipment. The use of ECMO must be judicious due to the considerable concern on the high cost and local availability of ECMO. ECMO is also used for the treatment of ARDS caused by leptospirosis and cytokine adsorption among COVID patients. The use of ECMO is included in the COVID fund and it costs around Php 2,000,000.00 per day, however, the costs are not fully covered. The results of ongoing trials are needed to show that the use of ECMO is effective and safe among patients with COVID-19 infection.

Should hyperbaric oxygen therapy be used in COVID-19 patients with hypoxemia?

As of 01 December 2021

RECOMMENDATION

We suggest against the use of hyperbaric oxygen therapy for the management of COVID-19 patients with hypoxemia due to insufficient evidence. (*Very low certainty of evidence; Weak recommendation*)

KEY FINDINGS

- Two interventional studies (non-randomized and randomized controlled trials) [56,57] were included in this review update on the use of hyperbaric oxygen therapy (HBOT) among COVID-19 patients with hypoxemia. HBOT appears to have a tendency towards benefit in reducing need for mechanical ventilation (HR 0.26, 95% CI 0.07-0.98) and in increasing the proportion of patients successfully weaned from oxygen support (RR 6.33, 95% CI 2.15-18.62).
- The effects of HBOT on mortality, oxygen saturation, and clinical improvement scores were inconclusive given the wide confidence intervals. The intervention appears generally safe with only a few minor adverse events including claustrophobia (RR 1.60, 95% CI of 0.07-38.20) and ear pain (RR 4.81, 95% CI of 0.27-86.47) reported. The overall certainty of evidence was low because of non-randomization, imprecision, and presence of possible confounders.

CONSENSUS ISSUES

The potential for the use of hyperbaric oxygen therapy in COVID-19 is recognized based on some reported benefit and minimal harm. However, its large cost and limited availability are important factors to consider especially since the noted benefits are based on a singular trial with very low certainty of evidence. More clinical trials are thus needed before recommending its use in patients with COVID-19.

Should sedation and neuromuscular blockade be done in mechanically ventilated patients with COVID-19-associated acute respiratory distress syndrome?

As of 03 January 2022

RECOMMENDATION

We suggest light over deep sedation in COVID-19 patients who are mechanically ventilated and who are anxious or agitated. (*Very low certainty of evidence; Weak recommendation*)

RECOMMENDATION

We suggest against the routine use of neuromuscular blockade in mechanically ventilated patients with COVID-19 associated respiratory distress syndrome. (*Low certainty of evidence; Weak recommendation*)

KEY FINDINGS

- There are currently no available randomized clinical trials testing for the effect of sedation and neuromuscular blockade in COVID-19 patients with acute respiratory distress syndrome (ARDS). Only indirect evidence from eight studies (two randomized clinical trials (RCTs) [58,59] on the use of sedation versus no sedation in mechanically ventilated patients and six RCTs [60-65] on the use of neuromuscular blockade agent (NMBA) compared to light sedation alone or placebo with or deep sedation in mechanically ventilated patients with moderate to severe ARDS) showed no significant benefit in 90-day mortality, length of hospital stay, and ventilator free-days.
- In terms of adverse events, major thromboembolic complications were significantly observed among mechanically ventilated patients on sedation while the incidence of delirium, accidental extubation, and ventilator-associated pneumonia did not significantly differ. The use of NMBA in moderate-to-severe ARDS showed significant reduction in 28- and 90-day mortality, improvement in PaO₂/FiO₂ ratio at 72 hours when compared to a deep sedation strategy without NMBA. Likewise, the use of NMBA reduced the risk of barotrauma and pneumothorax compared to no NMBA.

CONSENSUS ISSUES

Light sedation for mechanically ventilated COVID-19 patients has been suggested to help in managing agitation and anxiety. Certain patients, however, who are on paralytics and prone position, as well as those who exhibit ventilator asynchrony must be considered for deep sedation. On the other hand, routine neuromuscular blockade is not recommended unless there are indications for paralysis: as supportive management to facilitate lung protective strategies or prone ventilation.

Should inhaled nitric oxide be used in patients with COVID-19?

As of 26 October 2021

RECOMMENDATION

We recommend against the use of nitric oxide among patients with COVID-19. (*Low certainty of evidence; Strong recommendation*)

KEY FINDINGS

- One open-label randomized controlled trial [66] showed that the administration of inhaled nitric oxide as an adjunct to standard-of-care among hospitalized patients with moderate COVID-19 infection had no significant difference in 28-day mortality, need for mechanical ventilation, intensive care unit length of stay, hospital length of stay, viral clearance, and two-point WHO Ordinal Scale improvement when compared to standard-of-care alone.
- The incidence of methemoglobinemia was also not significantly different among patients who were given inhaled nitric oxide compared to those who received standard-of-care. The certainty of evidence was low due to lack of allocation concealment and serious imprecision of effect estimates.

CONSENSUS ISSUES

A strong recommendation against the use of nitric oxide was unanimously given on the basis of low certainty of currently available evidence (non-peer reviewed pre-print article with small population and imprecision of effect estimates) to justify its use and the high cost of the intervention.

Should etoposide be given among patients with severe COVID-19 pneumonia in cytokine storm?

As of 15 April 2021

RECOMMENDATION

We recommend against the use of etoposide among patients with COVID-19 pneumonia in cytokine storm. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

Very low quality evidence was found on the use of etoposide as treatment for severe COVID-19: one case series [67] and one case report.[68] The case series showed an improvement in PaO₂-FiO₂ ratio (PFR) post-treatment of etoposide. Among the 14 patients studied in both studies, three patients expired while ten patients improved and were subsequently discharged. Adverse effects of etoposide include alopecia, gastrointestinal symptoms, acute hypersensitivity reactions, myelosuppression, and rarely, secondary malignancies such as acute myelocytic leukemia.

CONSENSUS ISSUES

There is not enough high-quality evidence to show that etoposide is useful for the management of COVID-19 patients with pneumonia in cytokine storm. There are significant adverse events with the use of etoposide, namely: alopecia, nausea,

vomiting, myelosuppression, acute hypersensitivity reactions, hepatotoxicity, hypotension from rapid infusion, and secondary malignancies.

Should pulmonary rehabilitation be done among long COVID patients with residual pulmonary symptoms to improve pulmonary function and quality of life?

As of 30 June 2021

RECOMMENDATION

We recommend individualized pulmonary rehabilitation with pre-intervention medical clearance for long COVID patients who show residual respiratory symptoms. (*Moderate certainty of evidence; Strong recommendation*)

KEY FINDINGS

- We found one randomized controlled trial that investigated the effect of pulmonary rehabilitation (PR) among long COVID patients with residual respiratory symptoms.[69] The study was found to have a moderate risk of bias due to absence of blinding for both patients and outcome assessors and unclear allocation process. They also enrolled only participants above 65 years of age.
- Based on moderate quality of evidence, pulmonary function tests of those who received pulmonary rehabilitation significantly improved. Quality of life and anxiety scales also significantly differed between groups, with improvement noted in the PR group compared to no PR. There was no significant difference for depression and activities of daily living between groups and within groups before and after the intervention. The RCT did not report any adverse events; indirect evidence from PR done on COPD patients [70] likewise showed no to low number of adverse events which did not hinder the feasibility and safety of the procedure of pulmonary rehabilitation.

CONSENSUS ISSUES

The panel recommends that the start and duration of pulmonary rehabilitation of each patient should be individualized depending on the assessment of a pulmonologist. Studies showed that the assessment of pulmonary rehabilitation among long COVID patients should start at least 6 months after their hospital admission and last for as long as 6 weeks. However, recommendations on when the assessment for pulmonary rehabilitation should start differ across professional medical societies. An international task force with representation from the European Respiratory Society and the American Thoracic Society recommends that the assessment should be done 6 to 8 weeks after hospital discharge in order to identify patients who will have residual symptoms. In addition, pulmonary rehabilitation for other disease conditions lasts for 6 to 9 weeks.

Should pirfenidone versus nintedanib be used as therapy for post-COVID-19 pulmonary fibrosis?

As of 26 October 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of pirfenidone or nintedanib among patients with post-COVID-19 pulmonary fibrosis. (*Very low certainty of evidence*)

KEY FINDINGS

Limited evidence is available on the use of pirfenidone or nintedanib among patients with post-COVID-19 fibrosis, most of which are from published case reports. There are no clinical trials or observational studies among this group of patients which directly compare both anti-fibrotic agents, nor are there studies comparing each drug individually to placebo. Indirect evidence from prospective and retrospective cohort studies on the use of pirfenidone versus nintedanib among non-COVID patients with idiopathic pulmonary fibrosis [71-77] showed no difference in mortality rate, respiratory-related hospitalizations, acute exacerbation of pulmonary fibrosis and percent predicted forced vital capacity (FVC%) at 6 and 12 months of treatment; higher frequency of diarrhea and transaminitis with nintedanib; and a higher rate of discontinuation due to adverse events with pirfenidone.

CONSENSUS ISSUES

Evidence to support this recommendation is indirect as the studies included do not involve patients with post-COVID-19 pulmonary fibrosis. This is due to the paucity of available data on the use of pirfenidone and nintedanib in this population. There is no strong evidence to favor one anti-fibrotic agent over the other except for less side effects with nintedanib. Both drugs are costly, with annual expenses amounting to approximately Php 4.3M and Php 2.4M for pirfenidone and nintedanib, respectively. This may increase the inequity of treatment among those who cannot afford these drugs.

Evidence and Recommendations for the Non-Pharmacologic Interventions for Prevention and Control of COVID-19

Should cloth masks be used to prevent COVID-19 infection caused by Variants of Concern (VoC)?

As of 03 December 2021

RECOMMENDATION

We recommend the proper use of either a well-fitted cloth mask or a medical mask in the community setting. If a cloth mask will be used, we suggest that it should be made of at least two layers of cotton (e.g., t-shirt fabric) or non-woven nylon with aluminum nose bridge. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

At the time of writing, there were no published nor preprint studies that directly investigated the efficacy or effectiveness of cloth masks in preventing COVID-19 infection caused by specific SARS-CoV-2 variants of concern (VoC). There was indirect evidence from two studies that investigated cloth masks for the prevention of COVID-19 in general: one was a case-control study [1] that showed no significant difference between cloth mask and medical mask, and the other was a cluster-randomized trial [2] which showed that both cloth and surgical mask reduced the proportion of people with COVID-19-like symptoms in the community. The indirect evidence from these two studies suggested that cloth masks can reduce symptomatic COVID-19 cases at the community level, albeit at a lesser degree compared to medical masks. It also suggested that community use of both surgical and cloth masks can reduce symptomatic COVID-19 cases at the community level.

CONSENSUS ISSUES

The panel unanimously recommended the proper use of either a well-fitted cloth or a medical mask in the community setting. Whether to add qualifiers or not (regarding, for instance, the type of fabric and the particular setting) to this updated recommendation, as was done in the previous one, weighed heavily on the consensus panel and was discussed extensively. Eventually, it was highlighted that regardless of the community area or situation (e.g., crowded places, close-contact settings, confined and enclosed spaces, indoors or outdoors), proper wearing of a well-fitted mask must be emphasized and is still strongly recommended.

Is a facemask with face shield more effective than facemask alone in reducing SARS COV2 transmission?

As of 05 November 2021

RECOMMENDATION

We suggest against requiring the use of face shields in addition to face masks among the general public in non-healthcare settings. (*Very low certainty of evidence; Weak recommendation*)

RECOMMENDATION

We recommend the addition of face shields to face masks among the general public in areas with sustained community transmission of SARS-CoV-2. (*Very low certainty of evidence; Strong recommendation*)

We recommend the use of face shield plus medical face mask and standard personal protective equipment among health care workers not directly involved in the care of COVID-19 patients in areas with sustained community transmission of SARS-COV2. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

- There was no available direct evidence for face shield plus face mask versus face mask alone in the general public. In the initial review done in May 2021, there were three studies that were included; for this updated review (until September 2021) we found two additional case-control studies. All these five studies were conducted among health care workers: four were done in the health care setting and one in the community.
- Three case-control studies [3-5] showed a trend toward benefit with the use of face shields but this was inconclusive (OR 0.85, 95% CI 0.68-1.08). Two pre- and post-face shield use surveillance studies [6-7] showed significant benefit (OR 0.28, 95% CI 0.22-0.37 and OR 0.04, 95% CI 0.00-0.69), respectively. The overall certainty of evidence was very low as the studies were non-randomized, unadjusted for confounders and with high risk of bias as well as indirectness.

CONSENSUS ISSUES

For the first statement, which suggests against obligatory public use of face shields, the panel's primary consideration was the insufficient evidence from existing literature that were assessed to be of very low certainty. This assessment stemmed from the indirectness of the studies in which they enrolled healthcare workers – rather than the general public – in the community, and included laboratory experiments not directly involving humans, as well as studies that investigated different viral infections with varied routes of transmission. Other factors affecting the panel's decision were public preference and observed poor compliance, environmental pollution, safety issues particularly in workplaces, additional costs, increasing vaccination rates, declining COVID-19 incidence, and the individuals' capacity to assess their own risk in the

community. Despite this suggestion to rescind the mandatory use of face shields, continued proper use of well-fitted face masks is still recommended.

Due to the perceived benefit of providing an additional mechanical barrier against viral infections, the panel maintained a strong recommendation for wearing both face masks and face shields in high-risk environments such as crowded public transit, poorly ventilated spaces, and areas with high incidence of COVID-19 cases. In support of this, the panel considered the indirect evidence taken from a high-quality meta-analysis that showed a 78% reduction of risk in contracting respiratory infections from SARS-CoV-2, SARS, or MERS when an eye protector, such as face shield or goggles, was used compared with none.

There was a unanimous decision on adopting the previous recommendation on the use of face shield plus medical face mask for the healthcare workers and no concerns were mentioned.

Should copper-containing masks be used to decrease SARS-CoV-2 transmission?

As of 03 December 2021

RECOMMENDATION

There is no evidence to recommend the use of copper-containing over non-copper-containing masks to decrease SARS-CoV-2 transmission.

KEY FINDINGS

Currently, there are no clinical trials directly comparing copper-containing masks with non-copper-containing masks in terms of SARS-CoV-2 transmission except in two in-vitro studies [8-9]. One study demonstrated that the metal leaching potential of copper-containing textiles could potentially lead to adverse events attributed to inhalation or ingestion of copper. In terms of cost, copper-containing masks are more expensive than regular masks.

CONSENSUS ISSUES

Due to the absence of clinical trials on the efficacy of copper-containing masks in reducing COVID-19 transmission, the panel members opted to have no recommendation. One panelist voted against the use of copper masks because of the higher costs compared with using surgical or cloth masks and the potential harm of metal leaching from copper-containing textiles.

What is the appropriate PPE to be used use during surgeries to reduce the risk of virus transmission?

As of 30 June 2021

RECOMMENDATION

We recommend the use of appropriate PPE to include mask (N95 or higher standard), fluid repellent sealed well-fitting long gown, double gloves, apron, full face shield or goggles or visor, scrub hat, and disposable shoe covers or dedicated closed footwear among surgeons engaged in aerosol generating procedures of suspected or confirmed COVID-19 patients. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

- Five observational studies (2 cohorts, 2 cross-sectional, and 1 case series) investigated the effectiveness of PPE use in reducing SARS-CoV-2 transmission among healthcare workers involved in surgical aerosol-generating procedures (AGP).[10-14]
- Very low evidence suggests the protective effect of an appropriate PPE on surgeons engaged in AGP procedures of suspected or confirmed COVID-19 patients. Consistent N95 mask use reduced the odds of SARS-CoV-2 infections significantly (OR 0.37 [95% CI 0.21, 0.67], 1 study, n=195 participants) than inconsistent N95 use among healthcare workers involved AGP. Consistent gown use significantly reduced the odds of SARS-CoV-2 infections (OR 0.59 [95% CI 0.46, 0.77] I²= 0%, 2 studies, n= 941 participants) than inconsistent gown use amongst healthcare workers performing AGP.
- Consistent glove use reduced the odds of SARS-CoV-2 significantly (OR 0.42 [95% CI 0.43, 0.55] I²=34%, 3 studies, n=978) than inconsistent glove use among healthcare workers performing AGP. Very low evidence suggests the protective effects of N95 mask, gown, gloves, face shield/goggles, apron, and scrub hat in reducing SARS-CoV-2 transmission among healthcare workers performing AGP procedures.

CONSENSUS ISSUES

Although shoe cover was not mentioned in the assessed studies and in the recommendations from other groups, the panel agreed to include this in the minimum PPE required in surgery as it is part of the standard precaution. A strong recommendation was given despite the very low quality of evidence since the enumerated PPE is the existing minimum standard protection recommended for healthcare workers directly caring for COVID-19 patients. The panel also emphasized strict adherence to protocols and the appropriate use of this minimum PPE to prevent COVID-19 infection.

What is the appropriate PPE for healthcare workers in the outpatient setting to reduce the risk of virus transmission?

As of 30 June 2021

RECOMMENDATION

We recommend the use of at least surgical face mask and face shield for protection against COVID-19 infection among healthcare workers in the outpatient setting not performing aerosol generating procedures. Additional PPEs such as medical gowns and gloves should be worn as part of standard precautions during the performance of other procedures. *(Very low certainty of evidence; Strong recommendation)*

KEY FINDINGS

- There are no available direct evidence comparing the effectiveness of N95 versus surgical mask in COVID-19 infection among healthcare workers in the outpatient setting. Meta-analysis comparing the two among healthcare workers in general showed no significant difference in their effectiveness in preventing clinical and laboratory viral infection. One RCT [15] investigated the difference of N95 and surgical mask in protecting healthcare workers in different outpatient setting from viral respiratory infection and noted no significant difference between the two.
- Indirect evidence also shows more adverse skin reactions for those wearing N95 respirators as compared to surgical masks. The use of face shield on the other hand in addition to face mask provided added protection from acquiring COVID-19 among community healthcare workers in India based on a before and after study.[7] The use of gowns and gloves are standards of care in medicine whenever handling patient's body fluids and this recommendation is still applicable in the current setting.

CONSENSUS ISSUES

The superiority of N95 over medical face masks cannot be established based on the imprecise effects noted from the evidence. The high likelihood of dermatosis or skin infections while using N95 respirators was also noted. Considering these issues, the panel deemed that medical face masks would be more cost-effective as long as only non-aerosol generating procedures are done in the outpatient setting. The use of additional PPEs may be required depending on the procedure that will be performed, consistent with the recommendation of the US Centers for Disease Control (CDC). Face shields are preferred over goggles as it offers a greater level of protection from droplets.

Despite the very low quality of evidence, a strong recommendation was formulated as the listed PPEs are already considered the minimum standard protection needed by the healthcare workers. Strict adherence to appropriate use of these PPEs is emphasized.'

What is the appropriate PPE for health care workers in the wards, ICU and emergency room to reduce the risk of virus transmission?

As of 30 June 2021

RECOMMENDATION

We recommend the use of the following PPE: disposable hat, medical protective mask (N95 or higher standard), goggles or face shield (anti-fog), medical protective clothing, disposable gloves and disposable shoe covers or dedicated closed footwear as an effective intervention in the prevention of COVID-19 among health care workers in areas with possible direct patient care of COVID-19 positive patients and aerosol generating procedures. (*Moderate certainty of evidence; Strong recommendation*)

KEY FINDINGS

Four studies and a case report were found on the use of PPE among health care workers to prevent COVID infection.[16-20] Moderate certainty evidence from three of the studies showed that the use of Level 2 PPE (disposable hat, medical protective mask (N95 or higher standard), goggles (anti-fog) or protective mask (anti-fog), medical gown clothing or white coats covered by medical protective clothing, disposable gloves and disposable shoe covers), N95 respirators and face shields protected health care workers in hospital settings from COVID-19 infections. On the other hand, very low certainty evidence showed no significant protective effect from the use of face/surgical masks, gowns, and/or disposable gloves if used individually.

CONSENSUS ISSUES

Direct patient care is defined as hands on, face-to-face contact with patients for the purpose of diagnosis, treatment and monitoring. This recommendation was made by the panel as it prioritized giving the best available protection to the healthcare workers. Whenever possible, hospital administrators should invest in these PPEs. Strict adherence to the appropriate use of PPEs must be observed even if healthcare workers have already been vaccinated against COVID-19.

Should ionizing air filter be used in the prevention and control of COVID-19 infection in public spaces with sustained community transmission?

As of 30 June 2021

RECOMMENDATION

We recommend against the use of ionizing air purifier to reduce COVID-19 transmission in the community. (*Low certainty of evidence; Strong recommendation*)

KEY FINDINGS

- No direct evidence was found assessing the effectiveness of ionizing air filters in reducing SARS-CoV-2 infections. Five experimental studies [21-24] reported using an ionizing air purifier in reducing airborne particles, mostly in uninhabited laboratory settings. Ionizing air purifiers can efficiently remove the fine and ultrafine particles. However, its effectiveness in eliminating airborne organisms for infection control is lacking. Ozone, a dangerous respiratory irritant produced by some ionizing air purifiers, is a health risk to users.
- Overall quality: Most of the studies were at high risk of bias, with common issues on selecting tested ionizing air purifiers and the assessor's blinding.

CONSENSUS ISSUES

One of the studies noted that when an area is inhabited, reducing the particulate matter becomes insignificant once people move within the household, which consequently makes the ionizing air purifier ineffective. The panel also recognized that the harm caused by this intervention outweighs its benefit because one of the apparent disadvantages of ionizers is the emission of ozone, a powerful oxidant that may inflict health hazards through long-term or high-dose exposure.

Should foot baths be used in the prevention and control of COVID-19 infection?

As of 30 June 2021

RECOMMENDATION

We recommend against the use of foot baths for prevention and control of COVID-19 transmission. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

No evidence was found evaluating the effectiveness of footbaths in preventing or controlling COVID-19 infections.

Should misting tents or disinfection chambers be used in preventing and controlling COVID-19 transmission?

As of 30 June 2021

RECOMMENDATION

We recommend against the use of misting tents or disinfection chambers for preventing and controlling COVID-19 transmission. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

There are no existing or ongoing studies investigating the effectiveness and safety of sanitation tents for preventing and/or controlling COVID-19 infections. Indirect evidence from a rapid systematic review [26] of 6 laboratory studies and 1 in-vitro case control study showed effective viral inactivation after application of various cleaning products. However, disinfecting substances pose significant health risks, especially when used outside their manufacturer's recommendation or with repeated, prolonged exposure.

CONSENSUS ISSUES

The strong recommendation was based on the intrinsic irritant properties of the chemicals used in disinfection. Although some health facilities have implemented misting tents, evidence was not established to show their effectiveness to prevent and control COVID-19 transmission. Likewise, it was noted that aerosols are not actually recommended because the contact time is not enough to kill the microorganism.

Should ultraviolet (UV) lamps be used in the prevention and control of COVID-19 infection in public spaces in locations with sustained community transmission?

As of 30 June 2021

RECOMMENDATION

We recommend against the use of UV lamps or other UV devices in any place outside of a controlled clinic or hospital setting to prevent and control COVID-19 transmission. (*Low certainty of evidence; Strong recommendation*)

KEY FINDINGS

- No direct evidence was found evaluating the effectiveness of ultraviolet lamps in the prevention and control of COVID-19 infections in public spaces in locations with sustained community transmission.
- Indirect evidence of low quality showed some benefit in reducing the incidence of viral infection in a hospital ward.[25-31] However, the evidence for its potential harm such as skin erythema, ocular itching, blurring and conjunctival injections, was more significant.

CONSENSUS ISSUES

A strong recommendation was made based on the potential adverse reactions and the risks associated with UV lamps. Although the panel recognizes the germicidal effect of UV light in clinical settings, emphasis was made to limiting its use only in controlled environments (i.e., without the presence of human beings) with trained staff to minimize its potential health hazards. Since the use of personal UV lamps and devices in households has also become widespread following advertisements, the public must be appropriately informed that misuse of these devices may cause harmful, long-term effects on health.

Should high efficiency particulate air (HEPA) filters be used in the prevention and control of COVID-19 infection in public spaces and locations with sustained community transmission?

As of 30 June 2021

RECOMMENDATION

We suggest the use of HEPA filter as an option to improve air quality for COVID-19 prevention and control in indoor spaces with inadequate ventilation. (*Low certainty of evidence; Weak recommendation*)

KEY FINDINGS

No direct evidence was found assessing the effectiveness of HEPA filters in preventing and controlling COVID-19 infection in public spaces and locations with sustained community transmission. Low quality evidence from 3 laboratory experiments and 1 case series [32-35] demonstrated that HEPA filters appear to significantly improve air quality.

CONSENSUS ISSUES

A weak recommendation was made because of indirect evidence showing the benefit of HEPA filter in improving air quality. HEPA filters may be useless in public spaces with uncontrolled airflow and should be used only in areas where air exchange is compromised. To ensure that HEPA filters serve their purpose, the amount of air that can be filtered per hour by the machine must be matched with the size of the room. Proper installation and regular maintenance are likewise important to avoid contaminated air from recirculating back to the room and to maximize the machine's lifespan. In spite of the use of HEPA filters, minimum health standards should still be observed.

Should carbon dioxide (CO₂) monitors be used to reduce transmission of COVID-19?

As of 05 November 2021

RECOMMENDATION

We recommend the use of carbon dioxide (CO₂) monitors in enclosed spaces to guide actions to improve ventilation and reduce transmission of SARS-CoV-2. *(Moderate certainty of evidence; Strong recommendation)*

KEY FINDINGS

- We found no studies that directly answered the question on whether carbon dioxide level determination can be used to prevent SARS-CoV-2 infection. There were observational studies [36-38] that reported no infection when there was increased air ventilation and the CO₂ levels were low. In these studies, many factors affecting CO₂ levels were considered simultaneously real-time, such as use of personal protective equipment (PPEs), activity of persons in the room, duration of stay in the room, and size of the room, among others. The CO₂ levels, as detected by carbon dioxide sensors, were correlated with air ventilation and in most studies, the CO₂ level was designated to be a surrogate or proxy to COVID-19 risk of transmission. Indirect evidence showed that people exposed to a room with more than or equal to 1000 parts per million (ppm) of carbon dioxide in the air, as measured by a CO₂ monitor, had as high as 16x higher risk of contracting tuberculosis, compared to those in a room with lower CO₂ levels.[39]
- The other studies [40-42] investigated the presence of the virus in air (aerosol and droplets), and as fomites. Although there was a disagreement existed among the studies, as two studies were able to isolate the virus in the air, while the other was able to isolate the virus only as fomites, it may be possible that the negative growth of the virus may not mean that it is absent.

CONSENSUS ISSUES

The panel made a strong recommendation for the use of carbon dioxide (CO₂) monitors because of the moderate certainty of evidence based on an indirect study that showed much higher risk of contracting tuberculosis when exposed to a room whose air reached 1000 parts per million (ppm) of carbon dioxide. The panel believed CO₂ monitors could serve as a real-time guide to initiate activities that improve air ventilation (such as promoting distancing, opening windows, or turning on electric fans). However, two panelists still voted for a weak recommendation due to (1) unknown accuracy of various commercial monitors in detecting CO₂ levels and (2) concerns regarding the actual use of industrial-grade monitors (training of personnel, number of monitors needed, calibration, preventive maintenance) in different institutions. Even with the use of CO₂ monitors, the public must continue to observe the precautionary measures of handwashing, wearing face masks, and observing physical distancing to avoid infection with COVID-19.

What are effective decontamination techniques for N95 reuse?

As of 30 June 2021

RECOMMENDATION

In situations where there is shortage of filtering facepiece respirators (FFR), we suggest the use of Hydrogen Peroxide Vapor (HPV), Ultraviolet Germicidal Irradiation (UVGI), moist heat and peracetic acid dry fogging system (PAF) as options for N95 mask decontamination as recommended by the manufacturer based on their ability to reduce SARS-CoV-2 load and infectivity while still maintaining N95 mask integrity. (*Low certainty of evidence; Weak recommendation*)

RECOMMENDATION

We recommend against the use of autoclave and alcohol as these methods alter filtering facepiece respirator's (N95) integrity and degrade filtration efficacy. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

Eight quasi-experimental studies reported the effects of six decontamination techniques (i.e., alcohol, autoclave, moist heat, HPV, PAF, UVGI) on SARS-CoV-2 laden N95 masks.[43-50] There is low evidence on use of Hydrogen Peroxide Vapor (HPV), Ultraviolet Germicidal Irradiation (UVGI), moist heat and peracetic acid dry fogging system (PAF) as options for N95 mask decontamination. These decontamination methods resulted in $\geq 4.79 \log_{10}$ reductions in SARS-CoV-2. Except for autoclave, they preserved the quality fit of N95 masks after 10 decontamination cycles. There is very low evidence against the use of autoclave and alcohol for N95 mask decontamination as these methods alter N95 integrity and degrade filtration efficacy.

CONSENSUS ISSUES

Reuse of N95 masks should be considered only in cases of shortages and not during times of normal supply. It is important to specify the maximum number of cycles by which the decontamination techniques can still preserve the functional and physical integrity of the N95 mask: (1) autoclave- up to 2 cycles; and (2) moist heat, HPV and PAF- up to 10 cycles.

Regarding local availability, some noted that HPV is usually not affordable in resource-limited settings, making UVGI the preferred option due to its lower cost. In practice, however, HPV is considered more effective in decontamination. Its mist can reach all the spaces once halo fogging is done, which is beneficial to N95 mask since it has many pores. In contrast, UVGI can only disinfect the area reached by the UV light. Although HPV is available in the country, its use will entail the need for a special decontamination room and a machine. Some noted that HPV is usually not affordable in resource-limited settings. The panel also stressed the importance of checking the condition of the N95 mask before decontamination and not just solely rely on the

manufacturer's recommendation. Likewise, the proper procedure before decontamination should also be emphasized.

On the recommendation against the use of autoclave, it was clarified that although it can destroy the virus, the physical and functional integrity of the N95 mask are affected. In terms of the other decontamination techniques, no study was found on the effectiveness of air dry alone, exposure to sunlight, soap and water, 0.5% chlorine solution, and for other disinfecting agents such as benzalkonium chloride.

Should protective physical barriers be used to prevent COVID-19?

As of 30 June 2021

RECOMMENDATION

We suggest against the use of protective physical barrier enclosures (ex. aerosol box) for the prevention of COVID-19 among health care providers who perform aerosol generating medical procedures*. (*Very low certainty of evidence; Weak recommendation*)

**Proper PPEs should be used by health care providers when performing aerosol-generating procedures.*

RECOMMENDATION

We suggest the use of protective physical barriers in the prevention of COVID-19 in areas where social distancing cannot be adhered to (e.g., offices, reception desk). (*Very low certainty of evidence; Weak recommendation*)

***Adequate ventilation, physical distancing, use of facemasks and personal hygiene should still be maintained to prevent COVID-19 infections. Regular cleaning and disinfection of physical barriers should be practiced.*

KEY FINDINGS

- Two case series reported that there were no COVID-19 infections among health care providers who used protective barrier enclosures during aerosol generating procedures on COVID-19 patients.[51,52] One systematic review reported that barrier devices were effective at either preventing or reducing the number of particles escaping the system only when negative pressure suction was applied.[53]
- As for physical barriers (i.e., sneeze guards), we did not find any studies on its efficacy in preventing COVID-19 infections. Two computational fluid-particle dynamics (CFPD) simulation studies, however, showed that physical barriers could potentially reduce aerosol transmission within shared spaces such as classrooms or airplane cabins.[54,55]

CONSENSUS ISSUES

Protective physical barrier enclosures refer to passive protective barriers that were used to protect the health care workers during the early stages of the pandemic when PPE supplies were limited. The physical barrier enclosure can be used as an

additional layer of protection in situations where the HCP does not have adequate PPE for aerosol generating procedures. However, these barriers should be used with suction or negative pressure devices to reduce the risk of transmission from secondary aerosols that are trapped underneath the plastic barrier. Previous studies showed that aerosols can still escape a physical barrier enclosure. In addition, concerns related to efficiency and usability of such barriers have also been reported, such as: (1) longer intubation time; (2) difficulty in maneuvering the arms because of the rigidity of the box; (3) view of the anesthesiologist or ER doctor is blocked by the plastic or acrylic box). The Philippine Society of Anesthesiologists (PSA) also no longer recommends the use of physical barrier enclosures.

The use of physical barriers is not mandatory, but an option in areas where social distancing cannot be maintained. The proper disinfection of these barriers is emphasized. It was noted that the World Health Organization and Centers for Disease Control and Prevention have specifications and instruction for disinfection of barriers.

Should surfaces be disinfected to prevent COVID-19 infection?

As of 30 June 2021

RECOMMENDATION

We recommend the practice of cleaning and disinfecting surfaces using the appropriate disinfecting chemical agents such as 0.5% sodium hypochlorite solution (bleach) or 70% alcohol to prevent COVID-19 infection. (*Low certainty of evidence; Strong recommendation*)

For high touch surfaces and high traffic areas, such as in the workplace, disinfection should be done before shift, intermittently during, and after the shift.

For household disinfection, once daily disinfection on high touch surfaces is recommended.

KEY FINDINGS

No randomized controlled studies were found directly addressing this question. Indirect evidence from a rapid systematic review [26] of 7 studies, 5 laboratories, 1 case control, and 1 experimental provided data on the effects of cleaning products on SARS-CoV-2 inactivation. Indirect evidence from laboratory studies showed that SARS-CoV-2 is inactivated by common disinfectants within 1 minute when used according to their manufacturer's specifications and precautionary measures. However, disinfectants such as 0.1% sodium hypochlorite and 70% alcohol are potential irritants to the skin, ocular tissue, and respiratory tract and should be handled with proper protection and precaution.

CONSENSUS ISSUES

The panel noted that alcohol is actually recommended for use on skin but is now also being used in the environment because of its accessibility. As for sodium hypochlorite solution, the DOH Department Memorandum 2020-0157 (“Guidelines on Cleaning and Disinfection in Various Settings as an Infection Prevention and Control Measure Against COVID-19”) provides instructions on its dilution and appropriate use.

Evidence and Recommendations for the Vaccines and Prophylactic Interventions for COVID-19

Among persons at risk, what is the clinical efficacy, effectiveness and safety of BBIBP-CorV (Sinopharm) in the prevention of SARS-CoV-2 infection?

As of 02 December 2021

RECOMMENDATION

We recommend the use of BBIBP-CorV (Sinopharm), given as 200U (WIV04) or 4ug (HBO2) in 0.5 ml in 2 doses, 21 days apart, to prevent symptomatic and asymptomatic COVID-19 infection among healthy adults (18 to 59 years old). *(Moderate certainty of evidence; Strong recommendation)*

We suggest the use of BBIBP-CorV to prevent severe COVID-19 infection among healthy adults (18 to 59 years old). *(Low certainty of evidence; Weak recommendation)*

We suggest the use of BBIBP-CorV to prevent symptomatic COVID-19 infection in the following:

- a. Adults with comorbidities *(Very low certainty of evidence; Weak recommendation)*
- b. Older persons (60 years and older) *(Very low certainty of evidence; Weak recommendation)*

RECOMMENDATION

There is insufficient evidence to recommend for or against the use of BBIBP-CorV to prevent COVID-19 infection among the following:

1. Children (3-17 years old) *(Very low certainty of evidence)*
2. Immunocompromised population *(Very low certainty of evidence)*
3. Pregnant and lactating women *(Very low certainty of evidence)*

In areas where the SARS-CoV-2 variants of concern are prevalent, there is insufficient evidence to recommend for or against the use of BBIBP-CorV to prevent COVID. *(Very low certainty of evidence)*

KEY FINDINGS

The evidence base for the clinical efficacy, effectiveness, and safety of BBIBP-CorV (Sinopharm), as of October 29, 2021 included one network meta-analysis[3], four randomized controlled trials[4-7], 14 observational studies and two regulatory authority reports[8-24]. Trial results showed that BBIBP-CorV, particularly the HBO2 formulation, provided sufficient protection against symptomatic and asymptomatic COVID-19 infection, within a follow up period of 77days. Limited information was available regarding effectiveness against severe infection and on protection for the older population. Immunologic studies show that the vaccine is immunogenic even

among children[7]. Limited information from observational studies on the effectiveness of the vaccine for the immunocompromised suggested some protection and immunogenicity. Available safety data shows acceptable safety profile with no adverse event of interest reported.

CONSENSUS ISSUES

The decision of the Panel to defer any recommendation for the use of BBIBP-CorV on children and the immunocompromised was based on the limited evidence of efficacy. As no correlate of protection has been established for COVID-19 vaccines, the Panel did not feel that immunogenic response was sufficient to make a recommendation, especially from studies with a small number of participants. The same issue was considered in withholding a recommendation with the use of BBIBP-CorV against the SARS-CoV-2 variants of concern.

Is CoronaVac (Sinovac) effective and safe in the prevention of COVID-19-infections?: A Rapid Review (Update)

As of 28 October 2021

RECOMMENDATION

We recommend the use of the CoronaVac (Sinovac), given as (given as 0.5 mL (600SU) to prevent symptomatic SARS-CoV-2 infection in:

- Healthy Adults (*Low certainty of evidence; Strong recommendation*)
- Pregnant women in their first trimester after consultation with a physician (*Very Low certainty of evidence; Strong recommendation*)
- Pregnant women in their 2nd and 3rd trimester and lactating women (*Very Low certainty of evidence; Strong recommendation*)
- Adults who have medical comorbidities (*including chronic respiratory disease and infection, cardiovascular disease, chronic kidney disease, cerebrovascular disease, diabetes mellitus, obesity, neurologic disorder, chronic liver disease and others like sickle cell disease, thalassemia, or Down's syndrome, as per DOH guidelines dated April 5, 2021 on the A3 Priority Group*) (*Low certainty of evidence; Strong recommendation*)
- Immunocompromised patients after medical clearance from a physician (*the immunocompromised include those diagnosed with HIV, hepatitis B and C, those with cancer undergoing chemotherapy, transplant patients receiving immunosuppression*) (*Low certainty of evidence; Strong recommendation*)

In areas where Delta is the predominant variant of concern, we recommend the use of CoronaVac (Sinovac) (*Very Low certainty of evidence; Strong recommendation*)

For immunocompromised patients who received primary CoronaVac (Sinovac) vaccination, we recommend for heterologous booster vaccination (*Very Low certainty of evidence; Strong recommendation*)

RECOMMENDATION

We suggest the use of CoronaVac (Sinovac) to prevent SARS-CoV-2 infection in older adults (>60 years old). (*Low certainty of evidence; Weak recommendation*)

RECOMMENDATION

We suggest against the use of CoronaVac (Sinovac) to prevent SARS-CoV-2 infection in children (3 to 17 years old) (*Very Low certainty of evidence; Weak recommendation*)

Under the current context of low vaccine coverage and inadequate vaccine supply, we recommend against booster vaccination using CoronaVac (Sinovac) in the healthy, adult population (18 years old and above) (*Low certainty of evidence; Strong recommendation*)

KEY FINDINGS

The updated search on September 7, 2021 provided additional real-world evidence on the clinical effectiveness of CoronaVac in the general, as well as the immunocompromised population, in whom immune response to the vaccine is diminished. One trial showed that CoronaVac was immunogenic and safe among children aged 3 to 17 years old[30]. Studies among the immunocompromised population showed reduced immunologic responses after CoronaVac vaccination[42, 44, 60]. Additional evidence showed sufficient protection against COVID-19 infection by the Delta variant despite decreased immunologic response elicited by CoronaVac against the variant of concern [61,62]. However, available evidence suggests escape from immunity against the Gamma variant after CoronaVac vaccination[63]. It is also uncertain whether a booster results in additional protection against infection in terms of clinical outcomes[28, 69-71]. Studies suggest declining immunologic marker levels over time with concomitant decreased vaccine effectiveness in the older population[34]. The effect of booster vaccination in providing additional protection is limited[28].

CONSENSUS ISSUES

In the light of new evidence since the last version of the guidelines, the panel revised the wordings and updated the strengths of the recommendations accordingly.

The panel members were divided and could not reach consensus regarding homologous booster vaccination for CoronaVac. Despite three rounds of voting and additional evidence search followed by a Delphi and a fourth round of voting, no consensus was reached. The arguments for and against homologous boosting of these 4 vaccines in the immunocompromised are as follows:

FOR:

1. The immunocompromised are vulnerable and at risk of severe COVID-19 infection and should be given the necessary protection from effective vaccination.

2. Primary vaccination has been found to result in poor immunogenic response in this population. Without a third/additional dose, these patients remain relatively unprotected and would likely have breakthrough infections.
3. Majority of the local population, and this likely includes the immunocompromised, received either ChAdOx1 or CoronaVac as their primary vaccination. Hence, despite the lack of a strong evidence of efficacy/effectiveness, giving them a booster using the same vaccine is better than not giving one.

AGAINST:

1. Homologous boosting using these vaccines for this population may turn out to be a waste of precious resources, given the lack of evidence that demonstrates clinical or even immunologic effectiveness and safety.
2. Ongoing trials and continued evidence generation may soon provide the necessary answer to the question and it may be prudent to wait for their results.
3. Heterologous boosting may be a better option, considering the current evidence of a satisfactory benefit/harm ratio.

The panel gave a strong recommendation for heterologous booster vaccination despite the low certainty of the evidence because of its significantly better immunogenic response compared to homologous boosting combined with the acceptable safety profile. More importantly, with the inconsistent supply of vaccines, and the need for the provide an effective vaccination regimen to the immunocompromised, the heterologous booster vaccination may be the only available option at this time

Is vaccination with BBV152 (Covaxin/Bharat) effective and safe in the prevention of COVID-19 infections?: A Rapid Review

As of 21 October 2021

RECOMMENDATION

We recommend the use of BBV152 (Covaxin/Bharat), 0.5 mL/dose, in a two-dose regimen, 28 days apart for the prevention of symptomatic COVID-19 infection in healthy adults. (*Moderate certainty of evidence; Strong recommendation*)

RECOMMENDATION

We suggest the use of BBV152 (Covaxin/Bharat), 0.5 mL/dose, in a two-dose regimen, 28 days apart for the prevention of symptomatic COVID-19 infection:

- e. Adults who have stable medical co-morbidities and are at high risk for severe infection (*Low certainty of evidence; Weak recommendation*)
- f. Healthy, older adults (>60 years old) (*Low certainty of evidence; Weak recommendation*)
- g. Pregnant and lactating women, after discussing with a physician (*No direct evidence; Weak recommendation*)
- h. Immunocompromised patients, after discussing with a physician (*No direct evidence; Weak recommendation*)

RECOMMENDATION

We suggest against the use of BBV152 (Covaxin/Bharat) for the prevention of COVID-19 in children and adolescents. *(No evidence; Weak recommendation)*

We recommend against the use of BBV152 (Covaxin/Bharat) in individuals who have known allergies to its contents/excipients. *(Best practice statement)*

KEY FINDINGS

The search performed on September 20, 2021 included one (1) randomized controlled clinical trial (RCT) (n=25,798)[77] and seven (7) observational studies[78-84], which provided evidence on the efficacy, effectiveness, and safety of BBV152 (Covaxin/Bharat) in the prevention of SARS-CoV-2 infection, including those due to the variants of concern.

Is NVX-Cov2373 (Novavax) effective and safe in the prevention of COVID-19 infection?

As of 27 December 2021

RECOMMENDATIONS

We suggest the use of NVX-CoV2373 (Novavax), given as 5ug (with 50ug Matrix M1 adjuvant) two doses, intramuscular, 21 days apart, for the prevention of symptomatic and severe SARS-CoV-2 infection in healthy adults. *(Low certainty of evidence; Weak recommendation)*

We suggest the use of NVX-CoV2373 (Novavax), given as 5ug (with 50ug Matrix M1 adjuvant) two doses, intramuscular, 21 days apart, for the prevention of symptomatic SARS-CoV-2 infection in older adults (>65 years old). *(Low certainty of evidence; Weak recommendation)*

We suggest the use of NVX-CoV2373 (Novavax), given as 5ug (with 50ug Matrix M1 adjuvant) two doses, intramuscular, 21 days apart, for the prevention of symptomatic SARS-CoV-2 infection in adults with comorbidities. *(Moderate certainty of evidence; Weak recommendation)*

In areas where the Alpha variant is predominant, we suggest the use of the NVX-CoV2373 (Novavax) given as 5ug (with 50ug Matrix-M1 adjuvant), two doses, intramuscular, 21 days apart, to prevent symptomatic SARS-CoV-2 infection. *(Low certainty of evidence; Weak recommendation)*

RECOMMENDATIONS

We suggest against the use of NVX-CoV2373 (Novavax), for the prevention of symptomatic SARS-CoV-2 infection in the immunocompromised population (specifically HIV positive individuals). (*Very low certainty of evidence; Weak recommendation*)

We suggest against the use of NVX-CoV2373 for the prevention of symptomatic SARS-CoV-2 infection among pregnant and lactating women. (*No direct evidence; Weak recommendation*)

In areas where the Beta variant is predominant, we suggest against the use of the NVX-CoV2373 (Novavax) to prevent symptomatic SARS-CoV-2 infection. (*Low certainty of evidence; Weak recommendation*)

We recommend against the use of the NVX-CoV2373 (Novavax) in individuals who have known allergies to its contents/excipients, such as Matrix-M1. (*Best practice statement*)

RECOMMENDATION

There is insufficient evidence to recommend for or against the use of NVX-2373 for the prevention of symptomatic SARS-CoV-2 infection among children.

KEY FINDINGS

The evidence base on the efficacy and safety of NVX-Cov2373, as of November 12, 2021, includes five RCTs and one comparative cohort study[86, 88-92]. Results show that the vaccine provides excellent protection against symptomatic COVID-19 among healthy adults, among older persons above 65 years of age, among high-risk groups and those with comorbidities. The vaccine is highly immunogenic. Effectiveness is preserved against the Alpha variant, but one study showed potential loss of effect against the Beta variant. The vaccine is reactogenic.

CONSENSUS ISSUES

The Panel highly considered equity issues in the recommendations for the use of NVX-2373 in the prevention of SARS-CoV-2 infection. However, in the case of populations who were at high risk of severe infection and are immunocompromised such as the HIV positive individuals and in the pregnant and lactating women, the Panel took into consideration the lower efficacy estimates and the wide confidence interval with the lower border breaching the 30% threshold combined with the signal of additional harm with the higher adverse reaction risk so that a recommendation against the use of the vaccine was given.

Are vaccines effective and safe in the prevention of COVID-19 infections?

As of April 23, 2021

RECOMMENDATIONS

We recommend the use of the following vaccines to prevent symptomatic SARS-CoV-2 infection in adults: (*Moderate certainty of evidence; Strong recommendation*)

- a. **BNT162b2 (Pfizer/BioNTech)** given as 0.3ml (30ug) intramuscular injections, in 2 doses, 21 days apart
- b. **mRNA-1273 (Moderna)** given as 0.5ml (100ug) intramuscular injections, in 2 doses, 28 days apart
- c. **ChAdOx1 (AstraZeneca)** given as 0.5 ml (5×10^6 vp) intramuscular injections, in 2 doses, **at least 12 weeks** apart
- d. **Gam-COVID-Vac (Gamaleya)** given as rAd-26 0.5ml intramuscular injection, then rAd-5S 0.5 ml intramuscular injection 21 days after
- e. **Ad26.COVS.2.S (Janssen/Johnson&Johnson)** given as 0.5ml single dose intramuscular injection

We recommend the use of CoronaVac (Sinovac) (given as 0.5ml (600SU) intramuscular injection, in 2 doses, at 28 days apart) to prevent symptomatic SARS-CoV-2 infection among adults: (*Low certainty of evidence; Strong recommendation*)

CONSENSUS ISSUES

It was noted that ChAdOx1 was originally designed for a 21-day dosing interval, but because of some problems in logistics during the trial, different dosing intervals were implemented and the vaccine efficacy per dosing interval was recorded. The above recommendation, i.e., at least 12 weeks, reflects the dosing interval with the highest observed vaccine efficacy of 81.3% (95% CI 60.3-91.2).

With regard to Coronavac, a strong recommendation was made despite the low certainty of evidence because the panel considered its availability and the very high vaccine efficacy in preventing severe COVID-19. Although a 14-day dosing interval was utilized in the trial, it was noted that the recommendation on a 28-day dosing interval was based on the submission of the manufacturer to the Hongkong Food and Health Bureau. It was assumed that a longer duration may have better vaccine efficacy based on the immunogenicity data (seroconversion on 14-days versus 28-days) and the result of the subgroup analysis (<21 days versus >21 days). It was likewise explained that in general, increasing the interval between the doses of vaccines will provide better immunogenicity because it will give ample time for the population of lymphocytes in the lymph nodes to be replenished, thus resulting in a more robust immune response.

RECOMMENDATIONS

We recommend the use of BNT162b2 (Pfizer/BioNTech), mRNA-1273 (Moderna), ChAdOx1 (Astrazeneca), Gam-COVID-Vac (Gamaleya) and Ad26.COV2.S (Janssen/ Johnson&Johnson) vaccines to prevent symptomatic SARS-CoV-2 infection in older adults (>64 year old). (*Low certainty of evidence; Strong recommendation*)

We recommend the use of these vaccines in pregnant and lactating women after consultation with a physician. (*Very low certainty of evidence; Weak recommendation*)

RECOMMENDATION

There is insufficient evidence to recommend the use of CoronaVac to prevent symptomatic SARS-CoV-2 Infection in older adults (>60-year-old). (*Very low certainty of evidence*)

CONSENSUS ISSUES

The strength of recommendation was changed from weak to strong because, although the certainty of evidence is low, the benefits of vaccinating the elderly who are at risk of severe disease outweigh the harm as reported in the evidence presented, which showed lower adverse event rates in the said population compared to the younger group.

The panel agreed not to recommend CoronaVac due to the very low certainty of evidence wherein the interim analysis showed imprecision because of the very wide confidence interval for symptomatic COVID-19. Moreover, there was no disaggregation of data into mild, moderate or severe COVID-19 in older adults ≥ 60 y/o. There was also no data on harm.

RECOMMENDATION

We recommend the use of these vaccines in pregnant and lactating women after consultation with a physician. (*Very low certainty of evidence; Weak recommendation*)

CONSENSUS ISSUES

The certainty of evidence was changed from low to very low given that there was no evidence on either efficacy or safety in pregnant and lactating women because they were excluded from the trials. Regarding the risk of COVID-19 infection in the fetus, there is no evidence to date of vertical transmission, but there is increased incidence of premature birth and other complications arising from the pregnancy itself. There is also lack of evidence on transmitting COVID-19 infection through breastmilk. The risk of horizontal transmission in the household versus from the mother is the same provided that infection prevention and control (IPC) measures are observed. It was emphasized that the discussion with a physician should involve informing these women of the benefits and risks of vaccination, specific to the timing of its administration during the pregnancy. Physicians should be educated on these risks and benefits for the delivery of proper advice.

There was a discussion on the registry involving Pfizer and Moderna vaccines, where pregnant women who get vaccinated can volunteer to join and report their outcomes. This real world analysis showed that there was no difference in the incidence of early trimester complications. Moreover, it was mentioned that the WHO report already included eight (8) pregnancies after the J&J vaccination, which noted the following results: one (1) spontaneous abortion, one (1) elective abortion, while the rest had no reported congenital anomalies.

RECOMMENDATIONS

We recommend the use of BNT162b2 (Pfizer/BioNTech), mRNA-1273 (Moderna), ChAdOx1 (Astrazeneca), Gam-COVID-Vac (Gamaleya) and Ad26.COV2.S (Janssen/ Johnson&Johnson) vaccines to prevent SARS-CoV-2 infection in adults who have stable medical comorbidities and those who are at risk for severe infection. (*Moderate certainty of evidence; Strong recommendation*)

We suggest the use of CoronaVac to prevent SARS-CoV-2 infection in adults who have stable medical comorbidities and those who are at risk for severe infection. (*Very low certainty of evidence; weak recommendation*)

CONSENSUS ISSUES

A weak recommendation was made for Coronavac because the panel took into account the absence of any estimate of vaccine efficacy specific for this subgroup. Although the Brazilian trial included healthcare workers with stable medical comorbidities, the proportion of this subgroup as well as the vaccine efficacy for this specific population are unknown. As such, the panel considered the estimates of vaccine efficacy for the entire trial population as indirect evidence to support its use on those with comorbidities.

RECOMMENDATIONS

We recommend the use of these vaccines to prevent SARS-CoV-2 infections in immunocompromised patients (i.e., diagnosed with HIV, hepatitis B and C, those with cancer undergoing chemotherapy, transplant patients receiving immune-suppression) after medical clearance from a physician. (*Low certainty of evidence; Strong recommendation*)

CONSENSUS ISSUES

Despite the low certainty of evidence, a strong recommendation was made because the benefits of vaccination outweigh any potential harm. It was noted that there are no specific subgroup results for the immunocompromised patients and the expected vaccine efficacy would be lower, but then the vaccine will still give them protection against COVID-19. The panel also emphasized that a medical clearance from any physician should be sufficient to facilitate the vaccination.

RECOMMENDATIONS

We recommend against the use of these vaccines in children to prevent SARS-CoV-2 infection: (*Weak recommendation*)

- BNT162b2: <16 years old
- ChAdOx1: <18 years old

There is no evidence on the use of mRNA-1273, GamCOVID-Vac, Ad26.COV2.S and CoronaVac in children to prevent SARS-CoV-2 infection.

CONSENSUS ISSUES

A weak recommendation was made for BNT162b2 and ChAdOx1 because these are the only vaccines that included children in the trial protocol, the results of which are still pending. It was clarified that the recommendation against its use was not because of safety issues.

Based on the existing protocols, children are excluded in the clinical trials of mRNA-1273, GamCOVID-Vac, Ad26.COV2.S and CoronaVac. There is no statement of planned recruitment for the younger population.

RECOMMENDATIONS

We recommend against the use of particular vaccines in individuals who have known allergies to the contents/excipients of that vaccine, such as polysorbate (ChAdOx1, Gam-COVID-Vac and Ad26.COV2.S) and polyethylene glycol or PEG200 DMG (BNT162b2 and mRNA-1273). (*Moderate to high certainty of evidence; Strong recommendation*)

CONSENSUS ISSUES

It was clarified that the recommendation was specific for polysorbate and PEG because these two (2) excipients are notorious for hypersensitivity reactions based on different regulatory authorities. CoronaVac does not contain any of these, but the Philippine Society of Allergy, Asthma and Immunology (PSAAI) has been receiving reports on allergic reactions to CoronaVac and this is currently being investigated.

OTHER CONSENSUS ISSUES

The panel agreed to place in a separate document (i.e., guidance or standard operating procedure) the recommendations on (1) advising the recipients regarding adverse reaction and adverse events as well as the (2) implementation of a pharmacovigilance program and regular evidence review upon vaccine use.

KEY FINDINGS

As of April 23, 2021, interim trial data on the safety and efficacy of six COVID-19 vaccines have been made publicly available [95-196]. The BNT162b2 (Pfizer/BioNTech), the mRNA-1273 (Moderna), the ChAdOx1 (AstraZeneca), the

Gam-COVID-Vac (Gamaleya), and the Ad26.Cov2.S (Janssen/Johnson&Johnson) vaccines demonstrated satisfactory vaccine efficacy against symptomatic COVID-19 infection among adults in the short term with moderate certainty. Limited available information provides low certainty evidence that the CoronaVac (Sinovac) vaccine also provides satisfactory protection against symptomatic COVID-19 infection among adults. Data on the efficacy against severe COVID-19 infection and asymptomatic COVID-19 infection are still inconclusive, except for Ad26.CoV2.S, which demonstrated, with moderate certainty, good efficacy in preventing moderate and/or severe COVID-19 infection and acceptable protection against asymptomatic COVID-19 infection 28 days after vaccination. Efficacy data on preventing death from COVID-19 infection are still inconclusive. Very limited Phase 3 trial data are available to inform vaccine efficacy against the different variants of SARS-CoV-2.

Administration of these vaccines was associated with higher proportions of adverse reactions compared with the control, although serious adverse event rates were comparable. These adverse events, mostly from reactions to the vaccines, were mild to moderate and of short duration. Long term efficacy and safety data are still lacking.

Real world evidence supports clinical trial findings on the efficacy of BNT162b2, mRNA-1273, ChAdOx1, Gam-COVID-Vac and CoronaVac. A rare phenomenon, the vaccine-induced thrombotic thrombocytopenia has been found to have a possible link with the use of ChAdOx1 and Ad26.COV2.S. However, regulatory authorities have maintained a positive benefit-ratio for its continued use. No safety signals have been identified with the mRNA vaccines. Cases of anaphylaxis have been reported with the mRNA vaccines and with ChAdOx1 but they remain in very low numbers. Deaths after vaccination are also rare and often assessed as not related.

Is rAd26 (Sputnik Light) effective and safe in the prevention of COVID-19 infections?: A Rapid Review

As of 04 November 2021

RECOMMENDATION

We suggest the use of the rAd26 (Sputnik Light), given as 10^{11} vp per 0.5ml, single dose, intramuscularly to prevent symptomatic SARS-CoV-2 infection in:

- d. Healthy adults (*Low certainty, Weak recommendation*)
- e. Older adults (60 years and older) (*Low certainty, Weak recommendation*)
- f. Adults with comorbidities (*Low certainty, Weak recommendation*)

In areas where Alpha, Beta or Delta is the predominant variant of concern, we suggest the use of rAd26 (Sputnik Light) to prevent COVID-19 infection. (*Very Low certainty, Weak recommendation*)

RECOMMENDATION

We suggest against the use of rAd26 (Sputnik Light) to prevent symptomatic SARS-CoV-2 infection in:

- d. Children (3-17 years) (*No evidence, Weak recommendation*)
- e. Pregnant and lactating women (*No evidence, Weak recommendation*)
- f. Immunocompromised (*No evidence, Weak recommendation*)

KEY FINDINGS

Five observational studies [209-213] provided low to very low certainty evidence on the effectiveness and safety of rAd26 (Sputnik Light) against COVID-19 infection. Immunogenicity studies showed that rAd26 induced significant humoral and cellular response against SARS-COV-2 and its variants. Observational studies found rAd26 to provide sufficient protection against COVID-19 infection, hospitalization and death, particularly for the older person and those with comorbidities. Associated adverse events were mostly due to reactogenicity and were mild and transient. No evidence was found on the effect of rAd26 on children, pregnant women and the immunocompromised.

Among adults who received the standard full doses of any COVID-19 vaccine, what is the clinical and immunologic efficacy and effectiveness and safety of a booster?

As of 27 December 2021

RECOMMENDATION

We suggest the following homologous booster vaccination regimen for the general adult population:

- g. BNT162b2 (*Low certainty of evidence; Weak recommendation*)
- h. mRNA-1273 (*Low certainty of evidence; Weak recommendation*)
- i. ChAdOx1 (*Very low certainty of evidence; Weak recommendation*)
- j. Ad26.Cov2.S (*Very low certainty of evidence; Weak recommendation*)
- k. CoronaVac (*Very low certainty of evidence; Weak recommendation*)
- l. BBIBP-CorV (*Very low certainty of evidence; Weak recommendation*)

We suggest the following heterologous booster vaccination regimen for the general adult population:

- k. BNT162b2 primary, mRNA-1273 booster (*Very low certainty of evidence; Weak recommendation*)
- l. BNT162b2 primary, Ad26.CoV2.S booster (*Very low certainty of evidence; Weak recommendation*)
- m. mRNA-1273 primary, BNT162b2 booster (*Very low certainty of evidence; Weak recommendation*)
- n. mRNA-1273 primary, Ad26.CoV2.S booster (*Very low certainty of evidence; Weak recommendation*)
- o. ChAdOx1 primary, BNT162b2 booster (*Very low certainty of evidence; Weak recommendation*)
- p. Ad26.COv2.S primary, BNT162b2 booster (*Very low certainty of evidence; Weak recommendation*)
- q. Ad26.COv2.S primary, mRNA-1273 booster (*Very low certainty of evidence; Weak recommendation*)
- r. CoronaVac primary, BNT162b2 booster (*Very low certainty of evidence; Weak recommendation*)
- s. CoronaVac primary, ChAdOx1 booster (*Very low certainty of evidence; Weak recommendation*)
- t. BBIBP-CorV primary, BNT162b2 booster (*Very low certainty of evidence; Weak recommendation*)

We suggest the following homologous booster vaccination for the immunocompromised population:

- a. BNT162b2 (*Very low certainty of evidence; Weak recommendation*)
- b. mRNA-1273 (*Low certainty of evidence; Weak recommendation*)

We suggest the following heterologous booster vaccination regimen for the immunocompromised population:

- f. an mRNA vaccine primary, another mRNA vaccine *booster* (*Very low certainty of evidence; Weak recommendation*)
- g. an mRNA vaccine primary, ChAdOx1 booster (*Low certainty of evidence; Weak recommendation*)
- h. BNT162b2 primary, mRNA-1273 booster (*Very low certainty of evidence; Weak recommendation*)
- i. BNT162b2 primary, Ad26.CoV2.S booster (*Very low certainty of evidence; Weak recommendation*)
- j. mRNA-1273 primary, Ad26.CoV2.S booster (*Very low certainty of evidence; Weak recommendation*)

RECOMMENDATION

There is insufficient evidence to recommend the following homologous booster vaccination in the general population:

- a. Gam-COVID-Vac
- b. BBV152

There is insufficient evidence to recommend the use of the heterologous booster vaccination regimens other than the combinations included above in the general adult population.

There is insufficient evidence to recommend the following homologous booster vaccination for the immunocompromised population:

- g. ChAdOx1
- h. Ad26.CoV2.S
- i. CoronaVac
- j. Gam-COVID-Vac
- k. BBV152
- l. BBIBP-CorV

There is insufficient evidence to recommend the use of the heterologous booster vaccination regimen other than the combinations included above in the immunocompromised population.

KEY FINDINGS

Fifty-one studies [216-265] are now included in the review, including: 8 randomized trials; 16 studies involving immunocompromised population; and 20 studies on heterologous booster vaccination. Clinical outcomes of booster vaccination remain to be available only for BNT162b2 homologous vaccination from three observational studies and from on study which included BBIBP-CorV. A new study provides vaccine effectiveness estimates for BNT162b2 homologous vaccination. Immunogenicity studies among the general population comparing pre- and post-boost humoral response consistently show significant increases titers and seropositivity rates regardless of the booster regimen. Available data on cellular response after boosters

among the general population is limited and suggests increased response. Immunogenic response after boosters among the immunocompromised showed inconsistency. Low certainty evidence suggests booster vaccination to be safe, with heterologous booster vaccination associated with higher adverse reaction rates compared with homologous booster vaccination.

CONSENSUS ISSUES

The main considerations in the recommendations of the Panel were the positive benefit/harm ratio of the administration of a booster compared with no boosters and increasing vaccine equity, providing flexibility, and optimizing available vaccines by recommending the use of vaccine booster regimen despite low certainty of evidence and for those with comparatively lower efficacy (compared to other combinations).

Among adults, what is the clinical and immunologic efficacy and effectiveness and safety of heterologous COVID-19 vaccination compared to standard homologous COVID-19 vaccination in preventing COVID-19 infection?

As of 22 October 2021

RECOMMENDATION

We recommend the use of heterologous COVID-19 vaccination for those with serious adverse event to the first dose. (*Very low certainty of evidence; Strong recommendation*)

We suggest the use of heterologous COVID-19 vaccination in the event of the unavailability of the second dose in the recommended schedule. (*Very low certainty of evidence; Weak recommendation*)

KEY FINDINGS

Three RCTs, 18 observational studies, and 1 case report[275-295] investigated the use of heterologous vaccination using the following vaccines: BNT162b2, ChAdOx1, mRNA1273, CoronaVac, BBV152, Ad5-nCoV. No study was found using Gam-COVID-Vac, NVX-CoV2373, and BBIBP-CorV as a component of the vaccination regimen.

The overall certainty of evidence for efficacy/effectiveness is very low due to the study design (i.e. observational studies with a lack of control for confounding factors), the use of surrogate outcomes (immunogenicity), missing outcomes, and short follow up. The overall certainty of evidence for safety is low because of the study design (i.e. observational studies), unclear to lack of blinding, and the short follow-up.

Only one retrospective population-based cohort study[283] with ChAdOx1/BNT162b2 or mRNA-1273 as the heterologous regimen showed clinical efficacy against SARS-CoV-2 infection, hospitalization, and death.

Current evidence shows that heterologous primary vaccination is immunogenic and results in acceptable adverse reaction rates. Different combinations and regimen perform differently.

CONSENSUS ISSUES

The panel decided to give a strong recommendation, despite the very low certainty of evidence, on the use of heterologous vaccination among those with serious adverse event to the first dose. This is because of the need to provide the sufficient protection afforded by the second dose, especially for the vulnerable and at-risk populations. The importance of informed consent was emphasized by the panel, particularly on the benefits and harms of the administration of heterologous vaccination versus delayed administration of the second dose in the context of vaccine unavailability.

Are COVID-19 vaccines efficacious in preventing COVID-19 infections caused by the B.1.617.2 (Delta) Variant?

As of 28 October 2021

RECOMMENDATION

In areas where the Delta variant is the predominant circulating variant, we recommend for the use of the following vaccine to prevent symptomatic and severe COVID-19:

- e. 2 doses of BBV152 (Covaxin/Bharat)
(*Moderate certainty of evidence; Strong recommendation*)
- f. 2 doses of BNT162b2 (Pfizer)
(*Low certainty of evidence; Strong recommendation*)
- g. 2 doses of mRNA-1273 (Moderna)
(*Low certainty of evidence; Strong recommendation*)
- h. 2 doses of ChAdOx1 (Astra Zeneca)
(*Low certainty of evidence; Strong recommendation*)
- i. 2 doses of CoronaVac (Sinovac)
(*Very low certainty of evidence; Strong recommendation*)

In areas where the Delta variant is the predominant circulating variant, we suggest the use of the following vaccines to prevent symptomatic and severe COVID-19:

- c. Ad26.CoV2 (Janssen)
(*Low certainty of evidence; Weak recommendation*)
- d. Gam-COVID-Vac (Sputnik V)
(*Low certainty of evidence; Weak recommendation*)

CONSENSUS ISSUES

The panel agreed that all vaccines may be used in for protection against the infections caused by the Delta variant, despite some reduction in vaccine effectiveness. The panel's decisions on the strength of recommendation were based mainly on the certainty of evidence supporting the effectiveness. Vaccines with clinical evidence of effectiveness were given strong recommendations for use, whereas vaccines with only immunologic evidence of effectiveness were given weak recommendations.

Among children <18 years old, what is the efficacy/effectiveness and safety of COVID-19 vaccines compared to placebo in preventing COVID-19?

As of 21 October 2021

RECOMMENDATION

We recommend the use of the BNT162b2 (Pfizer/BioNTech) vaccine, [given as 0.3 mL (30 ug) intramuscular injections, in 2 doses, 21 days apart] for children 12-15 years old to prevent symptomatic SARS-CoV-2 infection. (*Moderate certainty of evidence; Strong recommendation*)

We suggest the use of the mRNA-1273 (Moderna) vaccine, [given as 0.5 mL (100 ug) intramuscular injections, in 2 doses, 28 days apart] for children 12-17 years old to prevent symptomatic SARS-CoV-2 infection. (*Low certainty of evidence; Weak recommendation*)

RECOMMENDATION

We suggest against the use of Coronavac (Sinovac), [given as 0.5 mL (600 SU) intramuscular injection, in 2 doses, 28 days apart] for children 3-17 years old to prevent symptomatic SARS-CoV-2 infection. (*No evidence; Weak recommendation*)

KEY FINDINGS

This review provides the evidence on the use of COVID-19 vaccines in children based on one (1) systematic review, three (3) randomized controlled trials (RCTs), two (2) real world effectiveness studies, five (5) case series/reports, and six (6) regulatory agency reports[330-345]. Findings show that BNT162b2 (Pfizer/BioNTech) and mRNA-1273 (Moderna) provide sufficient protection against COVID-19 infection among adolescents who are at least 12 years old. More reactogenicity events were noted but adverse events were generally non-severe. Higher immunogenicity responses were also seen with children compared to adults. CoronaVac (Sinovac) was found to be immunogenic among children aged 3-17 years old.

Real world evidence on the use of mRNA vaccines in children supported the findings in the clinical trials. Myocarditis was detected among children after mRNA vaccination, although rare.

CONSENSUS ISSUES

The issue of myocarditis associated with the mRNA vaccines in children was a major consideration in the recommendation, particularly with the mRNA-1273 (Moderna) vaccine. Despite the low incidence of post-vaccination myocarditis, some Panel members considered it significant enough, especially when normal children are concerned, to weigh heavily on the benefit risk ratio. The suspension of the use of mRNA-1273 among children in some countries due to the myocarditis issue was also raised, resulting in the cautious recommendation given for this vaccine. The Panel emphasized that warnings should be given to the parents and caregivers of the

vaccine recipients to consult immediately in case of symptoms occurring after vaccination, which may suggest the development of myocarditis.

Is COVID-19 vaccination effective and safe among pregnant and lactating individuals and their infants in the prevention of COVID-19 infections?

As of 27 December 2021

RECOMMENDATION

We suggest the use of following vaccines, after the first trimester, for the prevention of COVID-19 infection in pregnant and lactating women.

- h. BNT162b2 (Pfizer) (*Low certainty of evidence; Weak recommendation*)
- i. mRNA-1273 (Moderna) (*Low certainty of evidence; Weak recommendation*)
- j. ChAdOx1 (AstraZeneca) (*No direct evidence; Weak recommendation*)
- k. Ad26.CoV2.S (Janssen/Johnson&Johnson) (*No direct evidence; Weak recommendation*)
- l. CoronaVac (Sinovac) (*No direct evidence; Weak recommendation*)
- m. BBIBP-CorV (Sinopharm) (*No direct evidence; Weak recommendation*)
- n. BBV152 (Covaxin) (*No direct evidence; Weak recommendation*)

RECOMMENDATION

We suggest against the use of the following vaccines for the prevention of COVID-19 infection in pregnant and lactating women:

- c. Gam-CoV-Vac (Sputnik V) (*No direct evidence; Weak recommendation*)
- d. NVX-2373 (Novavax) (*No direct evidence; Weak recommendation*)

KEY FINDINGS

The current evidence base includes forty-two reports [357-394] that investigated the effectiveness, immunogenicity, and safety of COVID-19 vaccination among pregnant and lactating women and their infants. Three were systematic reviews and 39 were observational studies. Available evidence demonstrates that COVID-19 vaccination provides sufficient protection against COVID-19 disease for pregnant and lactating women, with transfer of antibody protection to the newborn, without significant increase in the risk of maternal and neonatal adverse events nor adverse pregnancy and delivery outcomes.

CONSENSUS ISSUES

The Panel took into consideration that pregnant and lactating women are at a higher risk for severe COVID infection and are a particularly vulnerable population to recommend COVID-19 vaccination for them. Despite the absence of direct evidence of the efficacy of some vaccines for this population, the Panel still recommended their use using indirect evidence of efficacy and safety from the other vaccines and from the general adult population. The decision to recommend against the use of Gam-COVID-Vac (Sputnik V) was made based on the manufacturer's restriction of their

product among pregnant women, in the absence of direct evidence of its effect. On the other hand, the decision to recommend against the use of NVX-2373 for pregnant women was mainly due to its low efficacy seen in the immunocompromised population and a higher risk of adverse reaction.

Is BCG vaccination effective and safe in the prevention of COVID-19 infections?

As of 09 April 2021

RECOMMENDATION

We suggest against the use of BCG vaccine for the prevention of COVID-19 infection. (*Very low certainty of evidence; weak recommendation*)

KEY FINDINGS

A systematic review of the literature on April 2, 2021 did not yield any completed randomized trial on the efficacy and safety of BCG vaccination in the prevention of COVID-19 infection. Five retrospective cohort studies and one case control study[403-407] were included in this review and all showed high risk of bias and conflicting results. As of March 6, 2021, 22 studies that address this issue have been registered, 21 of which are randomized controlled trials.

CONSENSUS ISSUES

A weak recommendation against the use of BCG vaccination for COVID-19 prevention was made by the Panel in consideration of the following issues. Most of the evidence included in the review are retrospective cohorts and one case control. Among the articles, the timing of BCG administration is variable or unclear and the outcomes were self-reported and not necessarily RT-PCR confirmed. Also, although an RCT was recently completed in Greece on the potential utility of BCG vaccine as prophylaxis for COVID-19, it is still a preprint and not yet included in the review. There is also clinical trial evidence in the elderly that it improves outcomes for other respiratory viral infections. Furthermore, it is relatively inexpensive and there is no evidence of serious harm. It was discussed that BCG vaccine is usually given among children to prevent disseminated or miliary tuberculosis.

It was also discussed in the panel meeting that the Philippines has a policy on birth doses for BCG within 24-hours of delivery; however, this is not fully implemented in health centers since it is available as a multi-dose vial, they do not administer this if there is only one infant. On the other hand, it was noted that there is no national policy on BCG vaccination among adults.

Among close contacts of COVID-19 patients, should casirivimab + imdevimab cocktail be used as post-exposure prophylaxis?

As of 04 November 2021

RECOMMENDATION

We suggest the subcutaneous use of casirivimab + imdevimab as day 4 post-exposure prophylaxis for COVID-19 close contacts*, ages 12 years and above weighing at least 40 kilograms, who are at risk for severe disease or hospitalization**. (Moderate certainty of evidence; weak recommendation)

*The definition of close contacts is the same as in the Living COVID CPG guidelines.

**This includes the following people: elderly; BMI >25; those with chronic diseases such as hypertension, diabetes, and chronic kidney disease; those who are not expected to mount an adequate immune response to the vaccine due to immunosuppressive therapy or those in an immunocompromised state.

KEY FINDINGS

There was 1 randomized controlled trial[414] that investigated casirivimab + imdevimab cocktail as post-exposure prophylaxis for RT-PCR SARS-CoV-2 negative close contacts of COVID-19 patients. The results showed a significant decrease in symptomatic and asymptomatic COVID-19 infection. Among those who developed infection, casirivimab + imdevimab resulted in significant decrease in duration of infection. There was no significant difference in serious adverse events between those given casirivimab + imdevimab and placebo. The overall certainty of evidence was rated moderate due to imprecision in 1 critical outcome.

CONSENSUS ISSUES

Despite moderate certainty of evidence, the panel decided on a weak recommendation for prophylactic use of casirivimab + imdevimab cocktail given the following factors related to equity: (1) prohibitive cost (PHP 25,000-30,000), (2) potential problems with accessibility, (3) limited supply, (4) EUA mandate last October 2021 specifically allowing its use for treatment only and (5) issues on applicability.

The recommendation is based on a single multi-center, randomized controlled trial that was done in the United States. There is a very limited window to administer the drug therefore, the poor contact tracing and the delayed release of test results are issues in our setting that compromises the applicability of the results of this study. The vaccination status of the participants as well as the prevalent viral strains during the time of the trial were considerations of the panel. While neither of the two were discussed in the study, the study was implemented from June 2020 to March 2021.

Should melatonin be used in the prevention of COVID-19 infection?

As of 26 February 2021

RECOMMENDATION

We recommend against the use of melatonin as prevention for COVID-19 infection.
(*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

One observational study [426] reported lower odds of getting a positive test result for COVID-19 among those taking melatonin compared to those who were not. The study was found to be with serious risk of bias because of absence of randomization and allocation concealment and the lack of details which precluded assessment of blinding and follow-up. For safety, one systematic review[427] reported that overall, melatonin supplements have a good safety profile. However, there are some adverse effects that are dose- and timing-related, and one study stated that caution should be exercised in the use of melatonin in patients with hypertension [428].

CONSENSUS ISSUES

It was discussed that among the limitations of the efficacy study was the lack of information on the dose administered and the duration of intake; hence, the panel could not compare it against the usual dose used for sleep disorder, which is one (1) 3-mg capsule a day for at least a month. The cost per capsule ranges from Php 16.00 to 20.00. Regarding melatonin's adverse effects, it was noted that indirect studies were evaluated (i.e., use of 2 mg to 10 mg melatonin for different indications); hence, the observation of excessive sleepiness may be dose-related. The panel raised a concern on the potential for its misuse or overuse considering that it may be marketed as prophylaxis for COVID-19, which is an off-label indication. Likewise, its potential adverse effects from long-term use are unknown.

Considering that the certainty of evidence to support the use of melatonin is very low, and in the context of potential adverse events as well as cost considerations, the panel decided to strongly recommend against its use for COVID-19 infection.

Should Vitamin D supplementation be used in the prevention of COVID-19 infection?

As of 18 March 2021

RECOMMENDATION

We recommend against the use of Vitamin D supplementation to prevent COVID-19 infection. (*Very low certainty of evidence; Strong recommendation*).

KEY FINDINGS

A retrospective cohort study on frail, elderly nursing home residents (N=66)[437] reported that fewer patients died among those who received 80,000 IU of Vitamin D during the week following the diagnosis of COVID-19, or within one month prior to diagnosis, when compared to patients who received other medications. There was no direct evidence on the safety of Vitamin D supplementation as prevention for COVID-19.

CONSENSUS ISSUES

It was pointed out that although the study by Jolliffe et al. found that Vitamin D supplementation reduced the risk of acute respiratory infection (ARI) overall, this systematic review was done prior to the COVID-19 pandemic, hence the population assessed did not include COVID-19 patients. In relation to the preventive effect of Vitamin D against ARI, it was noted that the duration of the trials in the systematic review ranged from 7 weeks to 12 months, and such a duration for effect to be seen is not deemed beneficial since vaccines will soon be available. Overall, the panel recognized that all the studies evaluated on its efficacy and safety were indirect evidence.

Should zinc supplementation be used in the prevention of COVID-19 infection?

As of 18 March 2021

RECOMMENDATION

We recommend against the use of zinc supplementation to prevent COVID-19 infection. (*Very low certainty of evidence; Strong recommendation*).

KEY FINDINGS

No direct published evidence was found on the use of zinc supplements in the prevention of COVID-19. Among non-COVID adult participants, zinc acetate doses of 100 to 150 mg/day taken for months had few adverse effects. Excessive intake of zinc may lead to gastrointestinal disturbances. When taken together with other supplements, zinc may reduce copper and iron levels in the body.

CONSENSUS ISSUES

A concern was raised that the recommendation may be misconstrued that zinc supplementation is not needed, but it was clarified that this will be specific only for COVID-19 infection.

Should hydroxychloroquine/ chloroquine be used in the prevention of COVID-19?

As of 12 March 2021

RECOMMENDATION

We recommend against the use of HCQ for pre-exposure prophylaxis in adults who are at high risk of exposure to COVID-19 cases. (*Moderate certainty of evidence; Strong recommendation*)

We recommend against the use of HCQ for post-exposure prophylaxis in adults who are exposed to COVID-19 cases. (*Low certainty of evidence; Strong recommendation*).

KEY FINDINGS

Two randomized controlled clinical trials on the use of hydroxychloroquine (HCQ) as pre-exposure prophylaxis for COVID-19 showed tendency for reduction in the risk of developing COVID-19 infection. However, there was significant increase in the risk of adverse events. Three RCTs were found on the use of HCQ for post-exposure prophylaxis for COVID-19 no significant reduction in the risk of developing COVID-19 infection and a trend toward increase in the risk of adverse events. [474-478] Most of the studies were found to be at risk for bias with common issues in ascertainment bias and one study without blinding.

CONSENSUS ISSUES

For pre-exposure prophylaxis, HCQ will only have 2% benefit considering the worst-case scenario. With regard to its safety profile, there is clear harm in terms of gastrointestinal (e.g., diarrhea, abdominal cramps, nausea and vomiting) and neurological (e.g., dizziness) adverse events, which are incapacitating to health workers if they are to use this as prophylaxis.

For post exposure prophylaxis, the low-quality evidence showed no significant benefit of HCQ, and a trend toward increase in the risk of adverse events was likewise observed.

Should lopinavir/ritonavir be used as prophylaxis for the prevention of COVID-19?

As of 12 March 2021

RECOMMENDATION

We recommend against the use of lopinavir/ritonavir for chemoprophylaxis in individuals exposed to COVID-19 patients. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

There is currently no available direct evidence on the use of lopinavir/ritonavir in the prevention of COVID-19 among healthy individuals, particularly those exposed to confirmed COVID-19 individuals. There is very low-quality indirect evidence of lopinavir/ritonavir as post-exposure prophylaxis for an outbreak of MERS-CoV [481].

CONSENSUS ISSUES

The panel strongly recommends against the use of lopinavir/ritonavir for chemoprophylaxis because of the very low certainty of evidence. The drug combination is not commercially available and can only be availed of through the HIV program. In addition to the limited accessibility, there might also be competition with HIV patients who need it as a regular regimen. Given the limited supply in the country, it is not deemed to be a feasible intervention for prophylaxis since a large quantity of lopinavir/ritonavir will be required in such cases.

Should ivermectin be used as COVID-19 prophylaxis for the general population?

As of 17 April 2021

RECOMMENDATION

We recommend against the use of ivermectin as COVID-19 prophylaxis for the general population. (*Very low certainty of evidence; Strong recommendation*)

We recommend against the use of ivermectin for COVID-19 as post-exposure prophylaxis for household contacts of confirmed COVID-19 patients. (*Very low certainty of evidence; Strong recommendation*)

We recommend against the use of ivermectin for COVID-19 as prophylaxis for healthcare workers. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

Four very low quality randomized controlled trials[493-496] were found on the use of ivermectin as COVID-19 prophylaxis. All RCTs were found to have very serious risk of bias in terms of lack of blinding, analysis of incomplete outcome data, and selective reporting. One RCT showed lower rates of developing COVID-19 related symptoms. Moreover, two RCTs showed lower RT-PCR-confirmed COVID-19 infection rates in

the ivermectin group compared to non-intervention group. However, one of these two RCTs administered a co-intervention in the ivermectin group, which poses serious validity issues. Lastly, one cluster RCT showed inconclusive RT-PCR-confirmed results for ivermectin. Mild adverse events were reported such as gastrointestinal upset, fatigue, sleepiness, pruritus, numbness, and burning sensation, all of which did not necessitate discontinuation of therapy.

CONSENSUS ISSUES

The studies included in the review had very serious or high risk of bias. In particular, the study by Elgazzar et al. (2021) had a very low overall certainty of evidence due to the risk of bias and serious imprecision from the wide 95% confidence interval (CI). Using the WHO Considerations for Evaluation of COVID-19 Vaccines as reference for efficacy of prophylaxis or preventive therapy, the upper end of the computed CI exceeded the threshold for primary efficacy endpoint estimate 0.70 (30% efficacy). The Shoumann et al. (2021) study also has a serious validity issue due to the premature termination of the control group, and lack of pretermination protocol, thus leading to selective reporting. Lastly, the results of the Chahla et al. (2021) study also have validity issues due to the presence of a co-intervention in the treatment arm.

The panel recognized the great potential for its misuse or overuse. The panel also stressed that there is a need to have concrete evidence on safety, as well as on the appropriate dose and dosing frequency, which the current very low quality evidence did not provide. Another issue raised was that only a compassionate special permit (CSP) has been granted to two specific hospitals that applied for the permit, despite the current registration of ivermectin products as veterinary treatment for internal and external animal parasites. Hence, there may be legal implications when a positive recommendation to use it as a prophylaxis is issued. The human-grade ivermectin, on the other hand, is still applying for emergency use authorization (EUA) from the Philippine Food and Drug Administration. Considering the vaccine hesitancy of the public, a concern was raised by the panel that if a recommendation to an alternative to the vaccine as prophylactic agent will be made, then people will most likely opt not to get vaccinated, undermining the national vaccination program of the government.

Should saline nasal irrigation be used for the prevention of COVID-19?

As of 12 March 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of saline nasal irrigation (SNI) to prevent COVID-19 in healthy individuals. (*Very low certainty of evidence*)

KEY FINDINGS

There are no direct studies assessing the benefit or harm of saline nasal irrigation use among healthy individuals as a preventive strategy for COVID-19. Only indirect evidence from an observational study[508] using saline spray compared to no nasal spray in the prevention of common colds showed shorter duration of symptoms of nasal blockage and discharge but no difference in number of respiratory infections.

CONSENSUS ISSUES

The panel did not provide a recommendation on SNI because of the very low certainty of evidence on the use of nasal spray to prevent the common cold.

Should steam inhalation be used for the prevention of COVID-19?

As of 12 March 2021

RECOMMENDATION

We recommend against the use of steam inhalation in the prevention of COVID-19. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

Based on two single arm observational studies[516-517] with high risk of bias, there is currently only very low certainty evidence showing the possible benefit of steam inhalation in the prevention of developing symptomatic COVID-19 among exposed healthy individuals and reducing symptoms and number of days to negative SARS-COV-2 RT-PCR test of COVID-19 confirmed individuals. Meanwhile, there is indirect evidence highlighting the significant adverse effects of steam inhalation among individuals using it for symptomatic relief from colds.

CONSENSUS ISSUES

The panel strongly recommended against the use of steam inhalation as prevention for COVID-19, despite the very low certainty of evidence, because it was recognized that the potential for harm outweighs the benefit.

Should aspirin be used for prophylaxis against COVID-19-induced coagulopathy in patients with COVID-19?

As of 02 June 2021

RECOMMENDATION

There is insufficient evidence on the use of aspirin as prophylaxis against COVID-19-induced coagulopathy among patients with COVID-19. (*Very low certainty of evidence*)

KEY FINDINGS

Very low quality evidence from one retrospective cohort study [526] was found on the use of aspirin prophylaxis for COVID-19 induced coagulopathy among COVID-19 patients. There was a significant reduction in the risk of mortality. COVID-19 induced coagulopathy and adverse events were not examined in this study.

Evidence and Recommendations for the Adjunct Interventions for Prevention and Control of COVID-19

Should zinc be used as adjunctive treatment for COVID-19 infection?

As of 21 December 2021

RECOMMENDATION

There is insufficient evidence to recommend zinc as adjunctive treatment for COVID-19 infection. (*Low certainty of evidence*)

KEY FINDINGS

This updated review [1-6] showed inconclusive results on the efficacy of zinc as adjunctive treatment for COVID-19 for the outcomes of in-hospital mortality and hospitalization of outpatients. There was a significantly higher number of adverse events in the group that received zinc compared to control.

CONSENSUS ISSUES

Despite the risk of adverse events (i.e., infusion site irritation and gastrointestinal side effects), the panelists agreed that the evidence of benefit and harm of zinc as an adjunct therapy for COVID-19 still poses uncertainty. From their experience with the use of oral zinc for prevention of other illnesses, there had been no significant events reported to identify oral zinc as a cause for concern. Its potential for benefit was still given high consideration, particularly pending the results of ongoing studies. One voted against its use due to the signal for harm (i.e., hospitalization) among ambulatory patients and the risk of adverse events in the zinc group.

Should B Vitamins be used as an adjunct in the treatment of COVID-19?

As of 30 June 2021

RECOMMENDATION

We suggest against the use of B vitamins as adjunct in the treatment of patients with COVID-19. (*Very low certainty of evidence; Weak recommendation*)

KEY FINDINGS

There was one cohort study on the use of vitamin D/magnesium/vitamin B12 (DMB) supplementation on patients with COVID-19 infection.[7] Likelihood of subsequent oxygen therapy including ICU support was significantly less for those taking DMB. However, there was no significant difference when looking at oxygen therapy alone or ICU support alone. There were no adverse events directly attributed to DMB use in the cohort study. There is very low certainty of evidence that suggests an association between excessive levels of Vitamin B12 and poorer outcomes in COVID-19 patients.[8] Overall, there is very low certainty of evidence due to risk of bias based on the observational design, as well as issues with directness (DMB was given in combination with vitamin D and magnesium) and imprecision (due to limited sample size).

CONSENSUS ISSUES

Vitamin B plays an important role in cell functioning and boosting the immune system. Therefore, there is a need to assess the potential of Vitamin B as an adjunct treatment for COVID-19.

There were no studies that assessed the use of Vitamin B alone as an adjunct treatment for COVID-19 but there are still ongoing studies that can provide a clear picture on the use of Vitamin B as adjunct treatment. There were also no studies found that assessed the correlation of Vitamin B deficiency and cytokine storm. However, a prospective study that investigated the levels of Vitamin B12 and folate plasma level among patients admitted for COVID-19 pneumonia until they were transferred to ICU or death ensued suggested that there was a potential association between high plasma levels of Vitamin B12 and increased risk of mortality.

Should vitamin C be used in the adjunctive treatment of COVID-19?

As of 21 December 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of vitamin C as adjunctive treatment for patients with COVID-19. (*Low certainty of evidence*)

KEY FINDINGS

There are eight RCTs included in this review but only four, that used intravenous vitamin C for hospitalized patients with moderate to severe COVID-19, were pooled.[9-12] For the outcome of mortality, there was a trend towards benefit with small negligible harm. There was no significant benefit and inconclusive results for length of hospital stay, length of ICU stay and need for mechanical ventilation. One RCT using oral vitamin C in the outpatient setting [13] and three RCTs using Vitamin C in addition to other adjunctive treatments [14-16] showed inconclusive results. Adverse events reported were flushing, headache, nausea, vomiting, stomach pain, and diarrhea.

CONSENSUS ISSUES

The evidence still posed low certainty in terms of efficacy (i.e., reduction in length of hospital and ICU stay, need for mechanical ventilation for hospitalized patients, and mortality among both hospitalized and non-hospitalized patients) and safety. The panel could not suggest against the use of vitamin C since only one study for non-hospitalized patients showed inconclusive effects on mortality and adverse events.

Among patients with COVID-19, should Vitamin D be used as adjunctive treatment?

As of 03 December 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of Vitamin D supplementation as an adjunct treatment for patients with COVID-19 infection. (*Very low certainty of evidence*)

KEY FINDINGS

We found eight randomized controlled trials of COVID-19 patients who were given vitamin D as adjunctive treatment.[17-24] We found no significant benefit for the following outcomes: mortality, ICU admission, need for mechanical ventilation, hospital length of stay, clinical improvement, and virologic clearance. Subgroup analysis revealed that vitamin D status did not significantly change our estimates. These results must be interpreted in the context of very low certainty of evidence due to serious risk of bias, serious inconsistency, and serious imprecision among the included studies.

CONSENSUS ISSUES

The panel members perceived that the quality of evidence, showing negligible benefit, is still very low. A panel member voted against using Vitamin D as an adjunct due to (1) the possibility of it being prescribed despite the inconclusive evidence of benefit, (2) additional expenses that would be incurred if prescribed, and (3) probable drug interactions with other medications.

Should melatonin be used in the adjunctive treatment of COVID-19?

As of 30 June 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of melatonin as adjunct treatment for patients with COVID-19 infection. (*Very low certainty of evidence*)

KEY FINDINGS

There was only one very low quality RCT that compared melatonin with standard treatment [25], and it showed significant reduction in cough, dyspnea, and fatigue, and significantly shorter hospital stay and return to baseline health in those given melatonin. There was no significant difference in the proportion of patients discharged on day 14. No adverse events were observed in the use of melatonin among COVID-19 patients.

Should virgin coconut oil be used in the adjunctive treatment of COVID-19?

As of 30 June 2021

RECOMMENDATION

There is no evidence to recommend the use of VCO as treatment among patients with COVID-19 infection.

KEY FINDINGS

Virgin coconut oil (VCO) is rich in lauric acid and pharmacologically active metabolite monolaurin. In vitro studies have found that VCO has anti-inflammatory, antioxidant, antibacterial, antifungal, and antiviral properties. In a clinical trial involving HIV and AIDS patients, VCO treatment led to an increase in CD4+ and lymphocyte counts as well as reduction in viral load. Currently, there are no published studies assessing the efficacy and safety of virgin coconut oil as an adjunctive treatment for COVID-19.

Should Lagundi (*Vitex negundo*) be used as adjunctive treatment for COVID-19 infection?

As of 29 October 2021

RECOMMENDATION

There is no evidence to recommend Lagundi (*Vitex negundo*) as adjunctive treatment for patients with COVID-19 infection.

KEY FINDINGS

Lagundi (*Vitex negundo*), a medicinal herbal cough remedy widely used in the Philippines, was considered a potential adjunctive treatment for COVID-19. We found no published studies on *Vitex negundo* on patients with COVID-19 but there is one completed local clinical trial [26] whose full results are not yet available.

CONSENSUS ISSUES

Considerations of the panel were the lack of completed studies to date and the possibility of misinterpretation, stemming from anecdotal accounts of relief of symptoms, which could lead to its use even without definite benefits.

Should Tawa-tawa (*Euphorbia hirta*) be used as adjunctive treatment for COVID-19 infection?

As of 29 October 2021

RECOMMENDATION

There is no evidence to recommend Tawa-tawa (*Euphorbia hirta*) as adjunctive treatment for patients with COVID-19 infection.

KEY FINDINGS

Tawa-tawa (*Euphorbia hirta*) is a medicinal herb widely used for febrile illness in the Philippines; it is considered as a potential adjunctive treatment for COVID-19. We found no published studies on *Euphorbia hirta* in patients with COVID-19. There is one ongoing local clinical trial.

CONSENSUS ISSUES

Due to the lack of evidence, the panel decided not to make a recommendation for the use of Tawa-tawa as adjunctive treatment for COVID-19. One panelist opted to abstain from the vote because her inclination was to suggest against the use of Tawa-tawa. While it has been used in the treatment of dengue, it has no proven benefits for COVID-19 infection and may cause harm.

Should oral fatty acid supplements be used as adjunct treatment for patients with COVID-19?

As of 30 June 2021

RECOMMENDATION

There is insufficient evidence to recommend the use of fatty acid supplements as adjunct treatment for patients with COVID-19. (*Low certainty of evidence*)

KEY FINDINGS

There is low certainty of evidence based on one RCT (N=128) that compared omega-3-fatty acid supplementation of high-protein enteral feeding versus high-protein enteral feeding alone in hospitalized patients with COVID-19 [27], which showed significant reduction in mortality (available case analysis, RR 0.85, [0.73, 0.99]). The RCT did not report adverse events. Indirect evidence on adverse events from two meta-analyses of RCTs on critically-ill ICU patients (non-COVID), mostly with ARDS, given fatty acid supplementation versus control, showed that gastrointestinal adverse events were common but did not differ significantly between groups (RR 1.04, 95% CI 0.96-1.13; moderate certainty of evidence).

CONSENSUS ISSUES

There were no reported adverse events in the study that assessed the effect of addition of 1000mg omega-3-fatty acid supplementation to enteral feeding of hospitalized patients with COVID-19. There was no significant difference in reported adverse events between groups given fatty acid supplementation or placebo in critically-ill ICU patients (non-COVID).

In terms of cost, a capsule of 1000mg omega-3-fatty acid may cost less than Php 20.00 in the market. It was also remarked that patient preference may be a factor because of the fishy taste.

Should N-acetylcysteine be used as an adjunct treatment for patients diagnosed with COVID-19?

As of 30 June 2021

RECOMMENDATION

We recommend against the use of intravenous N-acetylcysteine as adjunct treatment for patients with COVID-19 infection. (*Moderate certainty of evidence; Strong recommendation*)

KEY FINDINGS

One randomized controlled trial comparing NAC to placebo with 135 participants [28] was found to have no significant difference compared to control for mortality, invasive mechanical ventilation, ICU admission, length of hospital and ICU stay. A case series of ten respirator-dependent patients who responded to IV NAC showed reduction in inflammatory markers wherein six patients rebounded once NAC was discontinued and eight were discharged while two remained hospitalized.[29] Indirect evidence in a low-quality systematic review of NAC vs placebo in patients with Acute Respiratory Distress Syndrome failed to show any difference in the mortality, but significantly reduced ICU stay. [30]

CONSENSUS ISSUES

The panel distinguished between the oral and intravenous (IV) preparations of N-acetylcysteine (NAC), noting in part the cost of the IV agent.

A study included in this review which compared NAC and placebo among suspected or confirmed COVID-19 patients found no significant difference on its primary and secondary outcomes (i.e., mortality, invasive mechanical ventilation and ICU admission). However, NAC may essentially still be used for its other clinical indications (i.e., as mucolytic) on patients with COVID-19, but not necessarily for the treatment of COVID-19.

There were also a few studies included in the review that compared IV-NAC and placebo among ARDS patients, and although it was also noted that NAC has no ancillary role on the treatment of ARDS, it is used for intubated patients for its mucolytic and immunomodulating properties.

Should RAAS blockers be continued in patients with COVID-19?

As of 30 June 2021

RECOMMENDATION

We recommend continuing maintenance RAAS blockers for hypertension among patients with COVID-19 infection. (*Moderate certainty of evidence; Strong recommendation*)

KEY FINDINGS

Based on two RCTs (N=811) with moderate certainty of evidence [31,32], there is probably little or no significant reduction in the risk of deaths and severe disease for patients with hypertension and COVID-19 who continued RAAS blockers compared to those who discontinued RAAS blockers.

Should statins be used as adjunctive treatment in patients with COVID-19?

As of 29 October 2021

RECOMMENDATION

There is insufficient evidence to recommend statins as adjunctive treatment in patients with COVID-19. (*Very low certainty of evidence*)

KEY FINDINGS

Statins are lipid-lowering drugs known to have pleiotropic effects against inflammation and thrombosis. Very low quality of evidence from two randomized clinical trials [33-35] showed that among COVID-19 patients admitted to the ICU, the only significant finding was a decrease in hospital stay. There was no significant benefit for the following outcomes: composite risk for adjudicated venous or arterial thrombosis, the risk for ECMO treatment, the risk for all-cause mortality, or the individual components of adjudicated venous thromboembolism, all-cause mortality, or need for mechanical ventilation.

CONSENSUS ISSUES

Due to perceived risk of harm, two panelists were inclined to suggest against the use of statins as adjunctive treatment in patients with COVID-19.

Does the concurrent use of Ibuprofen worsen COVID-19 outcomes?

As of 30 June 2021

RECOMMENDATION

We suggest that Ibuprofen may still be used as symptomatic treatment of patients with COVID-19 infection if clinically warranted. Concurrent use of ibuprofen is not associated with worsening of COVID-19 outcomes. (*Very low certainty of evidence; Weak recommendation*)

KEY FINDINGS

There has been concern on the use of ibuprofen in COVID-19. Fang et al. reported that the point of entry for SARS-COV-2 is through angiotensin- converting enzyme 2 receptor, which is believed to be upregulated by ibuprofen. Based on six observational studies assessed [36-41], there was no significant association between ibuprofen use and worse COVID-19 outcomes: composite outcome (death, acute respiratory distress syndrome, ICU admission, shock) OR 1.06 (95% CI 0.81-1.40), Death OR 1.07 (95% CI 0.35-3.24), and progression of symptoms after propensity score matching. Five clinical trials on Ibuprofen use in COVID-19 are still on-going.

Should aspirin, taken as maintenance therapy for underlying medical conditions, be discontinued in patients with COVID-19?

As of 30 June 2021

RECOMMENDATION

There is insufficient evidence to recommend discontinuation of aspirin as maintenance therapy for underlying medical conditions in patients with COVID-19. (*Very low certainty of evidence*)

KEY FINDINGS

Very low-quality evidence from three retrospective cohort studies [42-44] was found on the continuation of aspirin as maintenance for underlying medical conditions among patients with COVID-19. There was no significant decrease in the odds of mortality in the use of aspirin. However, there was a significant increase in the odds of the composite outcome of mortality or discharge to hospice with continuation of aspirin. Thromboembolic and adverse events were not examined in these studies.

CONSENSUS ISSUES

Further studies are needed for the panel to make a recommendation on the discontinuation of aspirin as maintenance therapy in patients with COVID-19. There is certainty of benefit for continuing aspirin for underlying medical conditions. However, the benefit of its use among patients with COVID-19 is still uncertain.

Should antiseptic mouthwashes/gargles be used as adjunctive treatment for COVID-19 infection?

As of 21 December 2021

RECOMMENDATION

We recommend against the use of any antiseptic mouthwash as an adjunctive therapy for patients with COVID-19. (*Very low certainty of evidence; Strong recommendation*)

RECOMMENDATION

We recommend against the use of any antiseptic mouthwash to prevent COVID-19 in healthy individuals. (*Very low certainty of evidence; Strong recommendation*)

KEY FINDINGS

There were no studies that directly investigated antiseptic mouthwash for the prevention of COVID-19 in people without the infection. Seven randomized controlled trials (RCTs) on the use of antiseptic mouthwashes as adjunctive therapy in patients with COVID-19 were found.[45-51] Of these, six RCTs studied the effect of antiseptic mouthwashes on the SARS-CoV-2 viral load or cycle threshold. The studies, which used different agents, showed no clear clinical benefit. Only one RCT, which used hydrogen peroxide mouthwash, investigated symptom relief and found no significant benefit. Increased risk of thyroid dysfunction with povidone iodine mouthwash and nasal powder exposure was reported in one RCT. Another RCT reported increased risk of burning throat sensation with hydrogen peroxide mouthwash. For both adjunctive treatment and prevention of COVID-19, there was evidence for harm and no clear benefit with the use of antiseptic mouthwash.

CONSENSUS ISSUES

The panelists voted unanimously to recommend against the use of antiseptic mouthwash as an adjunctive treatment mainly due to the unclear clinical benefit based on the surrogate outcomes and the risk of harm among patients with COVID-19 from the very low certainty evidence. The evidence of harm on using mouthwash, which came from the studies used for adjunctive treatment, led the panel to also formulate a strong recommendation against its use for the prevention of COVID-19 infection in healthy individuals. They considered the probable risk of harm from long-term exposure to particular mouthwash components if used as a preventive strategy. Their strong recommendations also took into account individuals with possible or suspected thyroid problem. A panelist expressed that mouthwashes are unnecessary because people breathe in virus-containing particles that are likely to settle in their nasal passages. Furthermore, these antiseptic gargles could result in wasteful spending.

Should nasal sprays be used in the prevention and treatment of COVID-19 infection?

As of 03 December 2021

RECOMMENDATION

We suggest against the use of nasal spray as an adjunct to treatment of COVID-19 infection. (*Low certainty of evidence; Weak recommendation*)

RECOMMENDATION

There was no consensus on the use of nasal spray in addition to other preventive interventions such as vaccination, proper use of personal protective equipment, and adherence to quarantine and isolation protocols to prevent COVID-19 infection.

KEY FINDINGS

As an adjunct to prevention, two randomized controlled trials on healthy asymptomatic healthcare workers providing direct care to COVID-19 patients showed that the use of iota-carrageenan or dimethylsulfoxide ethanol nasal spray provided significant benefit in reducing the risk of COVID-19 infection compared to routine care with or without saline placebo.[52,53] The risk of adverse events and discontinuation of the intervention due to intolerance, comparing treatment and control groups, was inconclusive. Nasal sprays in these trials were used as an adjunct to proper personal protective equipment and universal precautions. Overall certainty of evidence was low.

As an adjunct to treatment, nine randomized controlled trials on patients with mild and moderate COVID-19 [54-62] showed that the use of nasal sprays containing different active agents (one study each for glycerol, hydrogen peroxide, nitric oxide, ivermectin, mometasone and triamcinolone; povidone iodine in three studies) did not result in significant benefit in terms of viral clearance and showed inconsistent benefit in terms of symptomatic relief. Furthermore, the risk for adverse events (mostly minor, such as sensation of nasal and throat burning, frequent sneezing, altered taste sensation, and minor epistaxis) was significantly higher in the nasal spray arm compared to placebo. Overall certainty of evidence was low.

CONSENSUS ISSUES

The statement suggesting against the use of nasal spray as an adjunct to treatment was unanimously approved by the panel members due to the high certainty of the risk of adverse events and the uncertain benefits from the use of glycerol- and hydrogen-peroxide-based nasal sprays. For the panel, the statement warranted a weak recommendation because (1) it is not a primary intervention and (2) despite the risk of harm, it may not necessarily result in overuse, unlike other drugs such as antibiotics.

The decision regarding the statement on the use of nasal spray as an adjunctive to other preventive interventions reached an impasse. The panel members who were inclined towards its use despite possible added costs based their arguments on the risk reduction of developing COVID-19, its availability in the local market, and its

authorization from the Philippine Food and Drug Administration as a medical device intended for short-term use. They also claimed that it could be a weak recommendation and that it would not be a replacement for any existing recommended intervention. However, other panel members who voted against its use placed a greater value on the benefit of vaccinations. A panel member also voiced opposition because the word “suggest” implied a strong recommendation for use despite the fact that only two small trials demonstrated benefit and several clinical trials are still underway. Another posited that glycerol-based sprays, which can be oily, might obstruct breathing when used with masks.

Evidence and Recommendations for the Screening and Diagnosis of COVID-19

1. Dixon BE, Wools-Kaloustian K, Fadel WF, Duszynski TJ, Yiannoutsos C, Halverson PK, et al. Symptoms and symptom clusters associated with SARS-CoV-2 infection in community-based populations: Results from a statewide epidemiological study. *medRxiv* [Internet]. 2020;e00146. Available from: <https://doi.org/10.1016/j.sciaf.2019.e00146>
2. Kimball A, Hatfield KM, Arons M, James A, Taylor J, Spicer K, et al. Asymptomatic and Presymptomatic SARS-CoV-2 Infections in Residents of a Long-Term Care Skilled Nursing Facility —. *Morb Mortal Wkly Rep Summ CDC*. 2020;69(13):377–81.
3. Reses, H.E., Fajans, M., Lee, S.H. *et al.* Performance of existing and novel surveillance case definitions for COVID-19 in household contacts of PCR-confirmed COVID-19. *BMC Public Health* 21, 1747 (2021). <https://doi.org/10.1186/s12889-021-11683-y>
4. Gootenberg DB, Kurtzman N, O'Mara T, et al, Developing a pulse oximetry home monitoring protocol for patients suspected with COVID-19 after emergency department discharge. *BMJ Health & Care Informatics* 2021;28:e100330. doi: 10.1136/bmjhci-2021- 100330
5. Galbraith M, Kelso P, Levine M, Wasserman RC, Sikka J, Read JS. Addressing silent hypoxemia with COVID-19: Implementation of an outpatient pulse oximetry program in Vermont. *Public Health Pract (Oxf)*. 2021 Nov;2:100186. doi: 10.1016/j.puhip.2021.100186. Epub 2021 Aug 27. PMID: 34467257; PMCID: PMC8390119.
6. Shah S, Majmudar K, Stein A, Gupta N, Suppes S, Karamanis M, Capannari J, Sethi S, Patte C. Novel Use of Home Pulse Oximetry Monitoring in COVID-19 Patients Discharged From the Emergency Department Identifies Need for Hospitalization. *Acad Emerg Med*. 2020 Aug;27(8):681-692. doi: 10.1111/acem.14053. Epub 2020 Jul 23. PMID: 32779828; PMCID: PMC7323027.
7. Kyriakides J, Khani A, Coleman R, Kelly C. Analysis of an Ambulatory Care Pathway for Patients With COVID-19 Utilising Remote Pulse Oximetry at a London District General Hospital. *Cureus*. 2021 Jan 29;13(1):e12979. doi: 10.7759/cureus.12979. PMID: 33659119; PMCID: PMC7920215.
8. Banzi R, Sala L, Colmi A, Gerardi C, Nattino G, Occhipinti F, Stura R, Targetti E, Bertolini G. Fattibilità ed efficacia di un modello di monitoraggio domiciliare avanzato dei pazienti affetti da CoViD-19 o sospetti [Feasibility and efficacy of home monitoring for patients with suspected or confirmed COVID-19.]. *Recenti Prog Med*. 2020 Oct;111(10):584-592. Italian. doi: 10.1701/3453.34418. PMID: 33078008.
9. Hutchings O, Dearing C, Jagers D, et al. Virtual health care for community management of patients with COVID-19. *MedRxiv* [preprint] 2020:2020.05.11.20082396. doi: 10.1101/2020.05.11.20082396
10. Vindrola-Padros C, Sidhu MS, Georghiou T, Sherlaw-Johnson C, Singh KE, Tomini SM, Ellins J, Morris S, Fulop NJ. The implementation of remote home monitoring models during the COVID-19 pandemic in England. *EClinicalMedicine*. 2021 Apr;34:100799. doi: 10.1016/j.eclinm.2021.100799. Epub 2021 Mar 30. PMID: 33817610; PMCID: PMC8008987.
11. O'Carroll O, MacCann R, O'Reilly A, Dunican EM, Feeney ER, Ryan S, Cotter A, Mallon PW, Keane MP, Butler MW, McCarthy C. Remote monitoring of oxygen saturation in individuals with COVID-19 pneumonia. *Eur Respir J*. 2020 Aug 13;56(2):2001492. doi: 10.1183/13993003.01492-2020. PMID: 32616588; PMCID: PMC7331654.
12. Gaeta T, Chiricolo G, Mendoza C, Vaccari N, Melville L, Melniker L, Bove J. 124 Impact of a Novel Telehealth Follow-Up Protocol for At-Risk Emergency Department Patients Discharged With Presumptive or Confirmed COVID-19. *Ann Emerg Med*. 2020 Oct;76(4):S49. doi: 10.1016/j.annemergmed.2020.09.135. Epub 2020 Oct 29. PMCID: PMC7598558.
13. Clarke J, Flott K, Fernandez Crespo R, Ashrafian H, Fontana G, Bengier J, Darzi A, Elkin S. Assessing the safety of home oximetry for COVID-19: a multisite retrospective observational study. *BMJ Open*. 2021 Sep 14;11(9):e049235. doi: 10.1136/bmjopen- 2021-049235. PMID: 34521666; PMCID: PMC8441226.
14. Lancet EA, Gonzalez D, Alexandrou NA, Zabbar B, Lai PH, Hall CB, Braun J, Zeig-Owens R, Isaacs D, Ben-Eli D, Reisman N, Kaufman B, Asaeda G, Weiden MD, Nolan A, Teo H, Wei E, Natsui S, Philippou C, Prezant DJ. Prehospital hypoxemia, measured by pulse oximetry, predicts hospital outcomes during the New York City COVID-19 pandemic. *J Am Coll Emerg Physicians Open*. 2021 Mar 17;2(2):e12407. doi: 10.1002/emp2.12407. PMID: 33748809; PMCID: PMC7967703.

15. Grutters LA, Majoor KI, Mattern ESK, et al. Home telemonitoring makes early hospital discharge of COVID-19 patients possible. *J Am Med Inform Assoc* 2020;ocaa168. doi: 10.1093/jamia/ocaa168.
16. Gordon WJ, Henderson D, DeSharone A, Fisher HN, Judge J, Levine DM, MacLean L, Sousa D, Su MY, Boxer R. Remote Patient Monitoring Program for Hospital Discharged COVID-19 Patients. *Applied clinical informatics*. 2020 Oct;11(05):792- 801.
17. Kodama R, Arora S, Anand S, Choudhary A, Weingarten J, Francesco N, Chiricolo G, Silber S, Mehta PH. Reengineering the Discharge Transition Process of COVID-19 Patients Using Telemedicine, Remote Patient Monitoring, and Around-the-Clock Remote Patient Monitoring from the Emergency Department and Inpatient Units. *Telemed J E Health*. 2021 Oct;27(10):1188-1193. doi: 10.1089/tmj.2020.0459. Epub 2020 Dec 14. PMID: 33320031.
18. Silven AV, Petrus AH, Villalobos-Quesada M, Dirikgil E, Oerlemans CR, Landstra CP, Boosman H, van Os HJ, Blanker MH, Treskes RW, Bonten TN. Telemonitoring for patients with COVID-19: recommendations for design and implementation. *Journal of Medical Internet Research*. 2020 Sep 2;22(9):e20953.
19. Francis NA, Stuart B, Knight M, Vancheeswaran R, Oliver C, Willcox M, Barlow A, Moore M. Predictors of adverse outcome in patients with suspected COVID-19 managed in a virtual hospital setting: a cohort study. *medRxiv*. 2020 Jan 1.
20. Wilcock J, Grafton-Clarke C, Coulson T. What is the value of community oximetry monitoring in people with SARS-CoV-2? A prospective, open-label clinical study. *medRxiv*. 2021 Jan 1.
21. Dirikgil E, Roos R, Groeneveld GH, Heringhaus C, Silven AV, Petrus AHJ, Villalobos- Quesada M, Tsonaka R, van der Boog PJM, Rabelink TJ, Bos WJW, Chavannes NH, Atsma DE, Teng YKO. Home monitoring reduced short stay admissions in suspected COVID-19 patients: COVID-box project. *Eur Respir J*. 2021 Aug 5;58(2):2100636. doi: 10.1183/13993003.00636-2021. PMID: 33795321; PMCID: PMC8015644.
22. Schradling WA, McCafferty B, Grove J, Page DB. Portable, consumer-grade pulse oximeters are accurate for home and medical use: Implications for use in the COVID-19 pandemic and other resource-limited environments. *J Am Coll Emerg Physicians Open*. 2020 Oct 20;1(6):1450-1458. doi: 10.1002/emp2.12292. PMID: 33392548; PMCID: PMC7771801.
23. Inada-Kim M, Chmiel FP, Boniface MJ, Pocock H, Black JJM, Deakin CD. Validation of home oxygen saturations as a marker of clinical deterioration in patients with suspected COVID-19. *MedRxiv [preprint]* 2020:2020.11.06.20225938. doi:10.1101/2020.11.06.20225938
24. Agulló V, Fernández-González M, Ortiz de la Tabla V, et al. Evaluation of the rapid antigen test Panbio COVID-19 in saliva and nasal swabs in a population-based point-of-care study. *J Infect*. 2020 Jan 1. Available from: <https://doi.org/10.1016/j.jinf.2020.12.007>
25. Albert E, Torres I, Bueno F, et al. Field evaluation of a rapid antigen test (Panbio™ COVID-19 Ag Rapid Test Device) for COVID-19 diagnosis in primary healthcare centres. *Clin Microbiol Infect*. 2020.
26. Cerutti F, Burdino E, Milia M, et al. Urgent need of rapid tests for SARS CoV-2 antigen detection: Evaluation of the SD-Biosensor antigen test for SARS-CoV-2. *J Clin Virol*. 2020;132:104654.
27. Abdelrazik AM, Elshafie SM, Abdelaziz HM. Potential Use of Antigen-Based Rapid Test for SARS-CoV-2 in Respiratory Specimens in Low-Resource Settings in Egypt for Symptomatic Patients and High-Risk Contacts. *Lab Med*. 2021;52(2):e46-e49. doi:10.1093/labmed/lmaa104
28. Abusrewil Z, Alhudiri IM, Kaal HH, El Meshri SE, Ebrahim FO, Dalyoum T, et al. Time scale performance of rapid antigen testing for SARS-CoV-2: Evaluation of 10 rapid antigen assays. *J Med Virol*. 2021;(June):4–10. doi: 10.1002/jmv.27186
29. Adnan N, Khandker S, Haq A, Chaity M, Khalek A, Nazim A et al. Detection of SARS-CoV-2 by antigen ELISA test is highly swayed by viral load and sample storage condition. *Expert Review of Anti-infective Therapy*. 2021;:1-9.
30. Agarwal J, Das A, Pandey P, Sen M, Garg J. "David vs. Goliath": A simple antigen detection test with potential to change diagnostic strategy for SARS-CoV-2. *J Infect Dev Ctries*. 2021 Jul 31;15(7):904-909. doi: 10.3855/jidc.13925. PMID: 34343113.
31. Albert E, Torres I, Bueno F, et al. Field evaluation of a rapid antigen test (Panbio™ COVID-19 Ag Rapid Test Device) for COVID-19 diagnosis in primary healthcare centres. *Clin Microbiol Infect*. 2021;27(3):472.e7-472.e10. doi:10.1016/j.cmi.2020.11.004
32. Alemany A, Baró B, Ouchi D, et al. Analytical and clinical performance of the panbio COVID-19 antigen-detecting rapid diagnostic test. *J Infect*. 2021;82(5):186-230. doi:10.1016/j.jinf.2020.12.033

33. Allan-Blitz LT, Klausner JD. A Real-World Comparison of SARS-CoV-2 Rapid Antigen Testing versus PCR Testing in Florida. *J Clin Microbiol.* 2021 Sep 20;59(10):e0110721. doi: 10.1128/JCM.01107-21. Epub 2021 Aug 4. PMID: 34346715; PMCID: PMC8451433.
34. Amer R, Samir M, Gaber O, EL-Deeb N, Abdelmoaty A, Ahmed A et al. Diagnostic performance of rapid antigen test for COVID-19 and the effect of viral load, sampling time, subject's clinical and laboratory parameters on test accuracy. *Journal of Infection and Public Health.* 2021;14(10):1446-1453.
35. Andreani J, Lupo J, Germi R, Laugier C, Roccon M, Larrat S et al. Evaluation of six commercial SARS-CoV-2 rapid antigen tests in nasopharyngeal swabs: Better knowledge for better patient management?. *Journal of Clinical Virology.* 2021;143:104947.
36. Asai N, Sakanashi D, Ohashi W, et al. Efficacy and validity of automated quantitative chemiluminescent enzyme immunoassay for SARS-CoV-2 antigen test from saliva specimen in the diagnosis of COVID-19. *J Infect Chemother.* 2021;27(7):1039-1042. doi:10.1016/j.jiac.2021.03.021
37. Baccani I, Morecchiato F, Chilleri C, Cervini C, Gori E, Matarrese D et al. Evaluation of Three Immunoassays for the Rapid Detection of SARS-CoV-2 antigens. *Diagnostic Microbiology and Infectious Disease.* 2021;101(2):115434.
38. Bachman C, Grant B, Anderson C, Alonzo L, Garing S, Byrnes S et al. Clinical validation of an open-access SARS-COV-2 antigen detection lateral flow assay, compared to commercially available assays. *PLOS ONE.* 2021;16(8):e0256352.
39. Baro B, Rodo P, Ouchi D, Bordoy AE, Saya Amaro EN, Salsench SV, et al. Performance characteristics of five antigen-detecting rapid diagnostic test (Ag-RDT) for SARS-CoV-2 asymptomatic infection: a head-to-head benchmark comparison. *J Infect.* 2021;82(6):269–75. doi: 10.1016/j.jinf.2021.04.009.
40. Bello-Chavolla O, Antonio-Villa N, Fernández-Chirino L, Guerra E, Fermín-Martínez C, Márquez-Salinas A et al. Diagnostic performance and clinical implications of rapid SARS-CoV-2 antigen testing in Mexico using real-world nationwide COVID-19 registry data. *PLOS ONE.* 2021;16(8):e0256447.
41. Berger A, Nsoga M, Perez-Rodriguez F, Aad Y, Sattonnet-Roche P, Gayet-Ageron A et al. Diagnostic accuracy of two commercial SARS-CoV-2 antigen-detecting rapid tests at the point of care in community-based testing centers. *PLOS ONE.* 2021;16(3):e0248921.
42. Bianco G, Boattini M, Barbui A, Scozzari G, Riccardini F, Coggiola M et al. Evaluation of an antigen-based test for hospital point-of-care diagnosis of SARS-CoV-2 infection. *Journal of Clinical Virology.* 2021;139:104838.
43. Billaud G, Gaymard A, Lina B, Laboratoire de Virologie des HCL CNR des virus des infections respiratoires. Evaluation du Test Antigénique ABBOTT SARS-COV2 ABBOT. Lyon, France: SFM (French Society of Microbiology), 2020.
44. Blairon L, Wilmet A, Beukinga I, Tré-Hardy M. Implementation of rapid SARS-CoV-2 antigenic testing in a laboratory without access to molecular methods: Experiences of a general hospital. *Journal of Clinical Virology.* 2020;129:104472.
45. Bornemann L, Kaup O, Kleideiter J, Panning M, Ruprecht B, Wehmeier M. Real-life evaluation of the Sofia SARS-CoV-2 antigen assay in a large tertiary care hospital. *Journal of Clinical Virology.* 2021;140:104854.
46. Mboumba Bouassa R, Veyer D, Péré H, Bélec L. Analytical performances of the point-of-care SIENNA™ COVID-19 Antigen Rapid Test for the detection of SARS-CoV-2 nucleocapsid protein in nasopharyngeal swabs: A prospective evaluation during the COVID-19 second wave in France. *International Journal of Infectious Diseases.* 2021;106:8-12.
47. Boum Y, Fai KN, Nikolay B, Mboringong AB, Bebell LM, Ndifon M, et al. Performance and operational feasibility of antigen and antibody rapid diagnostic tests for COVID-19 in symptomatic and asymptomatic patients in Cameroon: a clinical, prospective, diagnostic accuracy study. *Lancet Infect Dis.* 2021;21(8):1089–96. [https://doi.org/10.1016/S1473-3099\(21\)00132-8](https://doi.org/10.1016/S1473-3099(21)00132-8)
48. Brihn A, Chang J, OYong K, Balter S, Terashita D, Rubin Z et al. Diagnostic Performance of an Antigen Test with RT-PCR for the Detection of SARS-CoV-2 in a Hospital Setting — Los Angeles County, California, June–August 2020. *MMWR Morbidity and Mortality Weekly Report.* 2021;70(19):702-706.
49. Bruzzzone B, De Pace V, Caligiuri P, Ricucci V, Guarona G, Pennati B et al. Comparative diagnostic performance of rapid antigen detection tests for COVID-19 in a hospital setting. *International Journal of Infectious Diseases.* 2021;107:215-218.

50. Bulilete O, Lorente P, Leiva A, et al. Panbio™ rapid antigen test for SARS-CoV-2 has acceptable accuracy in symptomatic patients in primary health care. *J Infect.* 2021;82(3):391-398. doi:10.1016/j.jinf.2021.02.014
51. Caputo V, Bax C, Colantoni L, et al. Comparative analysis of antigen and molecular tests for the detection of Sars-CoV-2 and related variants: A study on 4266 samples. *Int J Infect Dis.* 2021;108:187-189.
52. Caruana G, Croxatto A, Kampouri E, Kritikos A, Opota O, Foerster M, et al. Implementing SARS-CoV-2 rapid antigen testing in the emergency ward of a Swiss university hospital: The INCREASE study. *Microorganisms.* 2021;9(4):798.. doi: 10.3390/microorganisms9040798.
53. Caruana G, Lebrun L, Aebischer O, Opota O, Urbano L, de Rham M et al. The dark side of SARS-CoV-2 rapid antigen testing: screening asymptomatic patients. *New Microbes and New Infections.* 2021;42:100899.
54. Cassuto NG, Gravier A, Colin M, Theillay A, Pires-Roteira D, Pallay S, et al. Evaluation of a SARS-CoV-2 antigen-detecting rapid diagnostic test as a self-test: Diagnostic performance and usability. *J Med Virol.* 2021;93(12):6686–92. doi: 10.1002/jmv.27249.
55. Cento V, Renica S, Matarazzo E, Antonello M, Colagrossi L, Di Ruscio F et al. Frontline Screening for SARS-CoV-2 Infection at Emergency Department Admission by Third Generation Rapid Antigen Test: Can We Spare RT-qPCR?. *Viruses.* 2021;13(5):818.
56. Cerutti F, Burdino E, Milia MG, Alice T, Gregori G, Bruzzzone B, et al. Urgent need of rapid tests for SARS CoV-2 antigen detection: Evaluation of the SD-Biosensor antigen test for SARS-CoV-2. *J Clin Virol.* 2020;132(104654):104654.
57. Chaimayo, C., Kaewnaphan, B., Tanlieng, N. et al. Rapid SARS-CoV-2 antigen detection assay in comparison with real-time RT-PCR assay for laboratory diagnosis of COVID-19 in Thailand. *Virol J* 17, 177 (2020). <https://doi.org/10.1186/s12985-020-01452-5>
58. Chiu R, Kojima N, Mosley G, Cheng K, Pereira D, Brobeck M et al. Evaluation of the INDICAID COVID-19 Rapid Antigen Test in Symptomatic Populations and Asymptomatic Community Testing. *Microbiology Spectrum.* 2021;9(1).
59. Ciotti M, Maurici M, Pieri M, Andreoni M, Bernardini S. Performance of a rapid antigen test in the diagnosis of SARS-CoV-2 infection. *J Med Virol.* 2021;93(5):2988–91. doi: 10.1002/jmv.26830. Epub 2021 Feb 9. PMID: 33527409; PMCID: PMC8014551.
60. Courtellemont L, Guinard J, Guillaume C, Giaché S, Rzepecki V, Seve A, et al. High performance of a novel antigen detection test on nasopharyngeal specimens for diagnosing SARS-CoV-2 infection. *J Med Virol.* 2021;93(5):3152–7. doi: 10.1002/jmv.26896.
61. Denina M, Giannone V, Curtioni A, Zanotto E, Garazzino S, Urbino A et al. Can we trust in Sars-CoV-2 rapid antigen testing? Preliminary results from a paediatric cohort in the emergency department. *Irish Journal of Medical Science (1971 -).* 2021;.
62. Diao B, Wen K, Zhang J, Chen J, Han C, Chen Y, et al. Accuracy of a nucleocapsid protein antigen rapid test in the diagnosis of SARS-CoV-2 infection. *Clin Microbiol Infect.* 2021;27(2):289.e1-289.e4. doi: 10.1016/j.cmi.2020.09.057. Epub 2020 Oct 5. PMID: 33031947; PMCID: PMC7534827.
63. Dierks S, Bader O, Schwanbeck J, Groß U, Weig M, Mese K et al. Diagnosing SARS-CoV-2 with Antigen Testing, Transcription-Mediated Amplification and Real-Time PCR. *Journal of Clinical Medicine.* 2021;10(11):2404.
64. González-Donapetry P, García-Clemente P, Bloise I, García-Sánchez C, Sánchez Castellano M, Romero M et al. Think of the Children. *Pediatric Infectious Disease Journal.* 2021;40(5):385-388.
65. Eleftheriou I, Dasoula F, Dimopoulou D, Lebessi E, Serafi E, Spyridis N, et al. Real-life evaluation of a COVID-19 rapid antigen detection test in hospitalized children. *J Med Virol.* 2021;93(10):6040–4. 10.1002/jmv.27149
66. Escrivá B, Mochón M, González R, García C, Pla A, Ricart A et al. “The effectiveness of rapid antigen test-based for SARS-CoV-2 detection in nursing homes in Valencia, Spain”. *Journal of Clinical Virology.* 2021;143:104941.
67. Favresse J, Gillot C, Oliveira M, Cadrobbi J, Elsen M, Euchet C, Laffineur K, Rosseels C, Van Eeckhoudt S, Nicolas JB, Morimont L, Dogné JM, Dourx J. Head-to-Head Comparison of Rapid and Automated Antigen Detection Tests for the Diagnosis of SARS-CoV-2 Infection. *J Clin Med.* 2021 Jan 13;10(2):265. doi: 10.3390/jcm10020265. PMID: 33450853; PMCID: PMC7828347.
68. Fenollar F, Bouam A, Ballouche M, Fuster L, Prudent E, Colson P et al. Evaluation of the Panbio COVID-19 Rapid Antigen Detection Test Device for the Screening of Patients with COVID-19. *Journal of Clinical Microbiology.* 2021;59(2).

69. Ferté T, Ramel V, Cazanave C, Lafon M, Bébéar C, Malvy D et al. Accuracy of COVID-19 rapid antigenic tests compared to RT-PCR in a student population: The StudyCov study. *Journal of Clinical Virology*. 2021;141:104878.
70. García-Fiñana M, Hughes DM, Cheyne CP, Burnside G, Stockbridge M, Fowler TA, et al. Performance of the Innova SARS-CoV-2 antigen rapid lateral flow test in the Liverpool asymptomatic testing pilot: Population based cohort study. *BMJ*. 2021;374:n1367. doi: 10.1136/bmj.n1637
71. Ford L, Lee C, Pray I, Cole D, Bigouette J, Abedi G et al. Epidemiologic Characteristics Associated With Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Antigen-Based Test Results, Real-Time Reverse Transcription Polymerase Chain Reaction (rRT-PCR) Cycle Threshold Values, Subgenomic RNA, and Viral Culture Results From University Testing. *Clinical Infectious Diseases*. 2021;73(6):e1348-e1355.
72. Fourati S, Audureau E, Chevaliez S, Pawlotsky JM. Évaluation de la performance diagnostique des tests rapides d'orientation diagnostique antigéniques COVID-19. France: AP-HP Hopitaux universitaires Henri-Mondor, 2020.
73. Fourati S, Langendorf C, Audureau E, et al. Performance of six rapid diagnostic tests for SARS-CoV-2 antigen detection and implications for practical use. *J Clin Virol*. 2021;142:104930. doi:10.1016/j.jcv.2021.104930
74. Frediani JK, Levy JM, Rao A, et al. Multidisciplinary assessment of the Abbott BinaxNOW SARS-CoV-2 point-of-care antigen test in the context of emerging viral variants and self-administration. *Sci Rep*. 2021;11(1):14604. Published 2021 Jul 16. doi:10.1038/s41598-021-94055-1
75. Pérez-García F, Romanyk J, Moya Gutiérrez H, Labrador Ballester A, Pérez Ranz I, González Arroyo J et al. Comparative evaluation of Panbio and SD Biosensor antigen rapid diagnostic tests for COVID-19 diagnosis. *Journal of Medical Virology*. 2021;93(9):5650-5654.
76. Gili A, Paggi R, Russo C, Cenci E, Pietrella D, Graziani A, et al. Evaluation of Lumipulse® G SARS-CoV-2 antigen assay automated test for detecting SARS-CoV-2 nucleocapsid protein (NP) in nasopharyngeal swabs for community and population screening. *Int J Infect Dis*. 2021;105:391–6. doi: 10.1016/j.ijid.2021.02.098.
77. Gremmels H, Winkel BMF, Schuurman R, Rosingh A, Rigter NAM, Rodriguez O, et al. Real-life validation of the Panbio™ COVID-19 antigen rapid test (Abbott) in community-dwelling subjects with symptoms of potential SARS-CoV-2 infection. *EClinicalMedicine*. 2021;31(100677):100677.DOI: doi.org/10.1016/j.eclinm.2020.100677.
78. Gupta A, Khurana S, Das R, Srigrayan D, Singh A, Mittal A, et al. Rapid chromatographic immunoassay-based evaluation of COVID-19: A cross-sectional, diagnostic test accuracy study & its implications for COVID-19 management in India. *Indian J Med Res*. 2021;153(1 & 2):126–31. DOI: 10.4103/ijmr.IJMR_3305_20.
79. Halfon P, Penaranda G, Khiri H, Garcia V, Drouet H, Philibert P et al. An optimized stepwise algorithm combining rapid antigen and RT-qPCR for screening of COVID-19 patients. *PLOS ONE*. 2021;16(9):e0257817.
80. Hartard C, Berger S, Josse T, Schvoerer E, Jeulin H. Performance evaluation of an automated SARS-CoV-2 Ag test for the diagnosis of COVID-19 infection on nasopharyngeal swabs. *Clin Chem Lab Med*. 2021;59(12):2003–9.. doi: 10.1515/ccbm-2021-0569.
81. Häuser, F., Sprinzi, M.F., Dreis, K.J. et al. Evaluation of a laboratory-based high-throughput SARS-CoV-2 antigen assay for non-COVID-19 patient screening at hospital admission. *Med Microbiol Immunol* 210, 165–171 (2021). <https://doi.org/10.1007/s00430-021-00706-5>
82. Herrera V, Hsu V, Adewale A, Johnson L, Hendrix T, Kuhlman J et al. Testing Healthcare Workers Exposed to COVID19 using Rapid Antigen Detection. 2020;.
83. Hirotsu Y, Maejima M, Shibusawa M, Nagakubo Y, Hosaka K, Amemiya K, et al. Comparison of automated SARS-CoV-2 antigen test for COVID-19 infection with quantitative RT-PCR using 313 nasopharyngeal swabs, including from seven serially followed patients. *Int J Infect Dis*. 2020;99:397–402. doi: 10.1016/j.ijid.2020.08.029. Epub 2020 Aug 12. PMID: 32800855; PMCID: PMC7422837.
84. Holzner C, Pabst D, Anastasiou O, Dittmer U, Manegold R, Risse J et al. SARS-CoV-2 rapid antigen test: Fast-safe or dangerous? An analysis in the emergency department of an university hospital. *Journal of Medical Virology*. 2021;93(9):5323-5327.
85. Homza M, Zelena H, Janosek J, Tomaskova H, Jezo E, Kloudova A et al. Performance of Seven SARS-CoV-2 Self-Tests Based on Saliva, Anterior Nasal and Nasopharyngeal Swabs Corrected for Infectiousness in Real-Life Conditions: A Cross-Sectional Test Accuracy Study. *Diagnostics*. 2021;11(9):1567.

86. Ifko M, Skvarc M. Use of immunochromatographic sars-cov-2 antigen testing in eight long-term care facilities for the elderly. *Healthc.* 2021;9(7):2–7. doi: 10.3390/healthcare9070868
87. Igloi Z, Velzing J, van Beek J, van de Vijver D, Aron G, Ensing R, et al. Clinical evaluation of Roche SD Biosensor rapid antigen test for SARS-CoV-2 in municipal health service testing site, the Netherlands. *Emerg Infect Dis.* 2021;27(5):1323–9. doi: 10.3201/eid2705.204688.
88. Iqbal B, Khan M, Shah N, Dawood M, Jehanzeb V, Shafi M. Comparison of SARS-CoV-2 antigen electrochemiluminescence immunoassay to RT-PCR assay for laboratory diagnosis of COVID-19 in Peshawar. *Diagnosis.* 2021;0(0).
89. Ishii T, Sasaki M, Yamada K, Kato D, Osuka H, Aoki K, et al. Immunochromatography and chemiluminescent enzyme immunoassay for COVID-19 diagnosis. *J Infect Chemother.* 2021;27(6):915–8. doi: 10.1016/j.jiac.2021.02.025.
90. Jääskeläinen AE, Ahava MJ, Jokela P, Szirovicza L, Pohjala S, Vapalahti O, et al. Evaluation of three rapid lateral flow antigen detection tests for the diagnosis of SARS-CoV-2 infection. *J Clin Virol.* 2021;137(104785):104785 doi: 10.1016/j.jcv.2021.104785.
91. James AE, Gulley T, Kothari A, Holder K, Garner K, Patil N. Performance of the BinaxNOW coronavirus disease 2019 (COVID-19) Antigen Card test relative to the severe acute respiratory coronavirus virus 2 (SARS-CoV-2) real-time reverse transcriptase polymerase chain reaction (rRT-PCR) assay among symptomatic and asymptomatic healthcare employees. *Infect Control Hosp Epidemiol.* 2021;1–3. doi: 10.1017/ice.2021.20.
92. Jegerlehner S, Suter-Riniker F, Jent P, Bittel P, Nagler M. Diagnostic accuracy of a SARS-CoV-2 rapid antigen test in real-life clinical settings: Antigen tests in real-life clinical settings. *Int J Infect Dis.* 2021;109(January):118–22. doi: <https://doi.org/10.1016/j.ijid.2021.07.01>
93. Jung C, Levy C, Varon E, et al. Diagnostic Accuracy of SARS-CoV-2 Antigen Detection Test in Children: A Real-Life Study. *Front Pediatr.* 2021;9:647274. Published 2021 Jul 15. doi:10.3389/fped.2021.647274
94. Kahn M, Schuierer L, Bartenschlager C, Zellmer S, Frey R, Freitag M et al. Performance of antigen testing for diagnosis of COVID-19: a direct comparison of a lateral flow device to nucleic acid amplification based tests. *BMC Infectious Diseases.* 2021;21(1).
95. Kanaujia R, Ghosh A, Mohindra R, et al. Rapid antigen detection kit for the diagnosis of SARS-CoV-2 - are we missing asymptomatic patients? [published online ahead of print, 2021 Jul 19]. *Indian J Med Microbiol.* 2021;S0255-0857(21)04168-2. doi:10.1016/j.ijmmb.2021.07.003
96. Karon BS, Donato LJ, Bridgeman AR, Blommel JH, Kipp B, Maus A, et al. Analytical Sensitivity and Specificity of Four Point of Care Rapid Antigen Diagnostic Tests for SARS-CoV-2 Using Real-Time Quantitative PCR, Quantitative Droplet Digital PCR, and a Mass Spectrometric Antigen Assay as Comparator Methods. *Clin Chem.* 2021;9:1–9. doi: 10.1093/clinchem/hvab138
97. Kim D, Lee J, Bal J, Seo S, Chong C, Lee J et al. Development and Clinical Evaluation of an Immunochromatography-Based Rapid Antigen Test (GenBody™ COVAG025) for COVID-19 Diagnosis. *Viruses.* 2021;13(5):796.
98. Mathur P, Kiro V, Gupta A, Singh P, Sharad N, Khurana S et al. Evaluation of COVID-19 antigen fluorescence immunoassay test for rapid detection of SARS-CoV-2. *Journal of Global Infectious Diseases.* 2021;13(2):91.
99. Klein JAF, Krüger LJ, Tobian F, Gaeddert M, Lainati F, Schnitzler P, et al. Head-to-head performance comparison of self-collected nasal versus professional-collected nasopharyngeal swab for a WHO-listed SARS-CoV-2 antigen-detecting rapid diagnostic test. *Med Microbiol Immunol [Internet].* 2021;210(4):181–6. Available from: <https://doi.org/10.1007/s00430-021-00710-9> doi: 10.1007/s00430-021-00710-9
100. Kobayashi R, Murai R, Moriai M, Nirasawa S, Yonezawa H, Kondoh T, et al. Evaluation of false positives in the SARS-CoV-2 quantitative antigen test. *J Infect Chemother.* 2021;27(10):1477–81. doi: 10.1016/j.jiac.2021.06.019.
101. Koeleman J, Brand H, de Man S, Ong D. Clinical evaluation of rapid point-of-care antigen tests for diagnosis of SARS-CoV-2 infection. *European Journal of Clinical Microbiology & Infectious Diseases.* 2021;40(9):1975-1981.
102. Kolwijck E, Brouwers-Boers M, Broertjes J, van Heeswijk K, Runderkamp N, Meijer A et al. Validation and implementation of the Panbio COVID-19 Ag rapid test for the diagnosis of SARS-CoV-2 infection in symptomatic hospital Trained personnels. *Infection Prevention in Practice.* 2021;3(2):100142.
103. Korenkov M, Poopalasingam N, Madler M, Vanshylla K, Eggeling R, Wirtz M et al. Evaluation of a Rapid Antigen Test To Detect SARS-CoV-2 Infection and Identify Potentially Infectious Individuals. *Journal of Clinical Microbiology.* 2021;59(9).

104. Krüger L, Gaeddert M, Köppel L, Brümmer L, Gottschalk C, Miranda I et al. Evaluation of the accuracy, ease of use and limit of detection of novel, rapid, antigen-detecting point-of-care diagnostics for SARS-CoV-2. 2020;.
105. Krüger L, Gaeddert M, Tobian F, Lainati F, Gottschalk C, Klein J et al. The Abbott PanBio WHO emergency use listed, rapid, antigen-detecting point-of-care diagnostic test for SARS-CoV-2—Evaluation of the accuracy and ease-of-use. *PLOS ONE*. 2021;16(5):e0247918.
106. Krüger L, Klein J, Tobian F, Gaeddert M, Lainati F, Klemm S et al. Evaluation of accuracy, exclusivity, limit-of-detection and ease-of-use of LumiraDx™: An antigen-detecting point-of-care device for SARS-CoV-2. *Infection*. 2021;.
107. Krüttgen A, Cornelissen C, Dreher M, Hornef M, Imöhl M, Kleines M. Comparison of the SARS-CoV-2 Rapid antigen test to the real star Sars-CoV-2 RT PCR kit. *Journal of Virological Methods*. 2021;288:114024.
108. Kurihara Y, Kiyasu Y, Akashi Y, Takeuchi Y, Narahara K, Mori S, et al. The evaluation of a novel digital immunochromatographic assay with silver amplification to detect SARS-CoV-2. *J Infect Chemother*. 2021;27(10):1493–7. doi: 10.1016/j.jiac.2021.07.006
109. Kweon O, Lim Y, Kim H, Choi Y, Kim M, Choi S et al. Evaluation of rapid SARS-CoV-2 antigen tests, AFIAS COVID-19 Ag and ichroma COVID-19 Ag, with serial nasopharyngeal specimens from COVID-19 patients. *PLOS ONE*. 2021;16(4):e0249972.
110. Kyritsi M, Vontas A, Voulgaridi I, Matziri A, Komnos A, Babalis D et al. Rapid Test Ag 2019-nCoV (PROGNOSIS, BIOTECH, Larissa, Greece); Performance Evaluation in Hospital Setting with Real Time RT-PCR. *International Journal of Environmental Research and Public Health*. 2021;18(17):9151.
111. Lambert-Niclot S, Cuffel A, Le Pape S, Vauloup-Fellous C, Morand-Joubert L, Roque-Afonso A-M, et al. Evaluation of a rapid diagnostic assay for detection of SARS-CoV-2 antigen in nasopharyngeal swabs. *J Clin Microbiol* [Internet]. 2020;58(8). Available from: <http://dx.doi.org/10.1128/JCM.00977-20>.
112. Landaas ET, Storm ML, Tollånes MC, Barlinn R, Kran A-MB, Bragstad K, et al. Diagnostic performance of a SARS-CoV-2 rapid antigen test in a large, Norwegian cohort. *J Clin Virol*. 2021;137(104789):104789. doi: 10.1016/j.jcv.2021.104789.
113. Leixner G, Voill-Glaninger A, Bonner E, Kreil A, Zadnikar R, Viveiros A. Evaluation of the AMP SARS-CoV-2 rapid antigen test in a hospital setting. *International Journal of Infectious Diseases*. 2021;108:353-356.
114. Leli C, Di Matteo L, Gotta F, et al. Performance of a SARS-CoV-2 antigen rapid immunoassay in patients admitted to the emergency department. *Int J Infect Dis*. 2021;110:135-140. doi:10.1016/j.ijid.2021.07.043
115. L'Huillier A, Lacour M, Sadiku D, Gadiri M, De Siebenthal L, Schibler M et al. Diagnostic Accuracy of SARS-CoV-2 Rapid Antigen Detection Testing in Symptomatic and Asymptomatic Children in the Clinical Setting. *Journal of Clinical Microbiology*. 2021;59(9).
116. Linares M, Pérez-Tanoira R, Carrero A, Romanyk J, Pérez-García F, Gómez-Herruz P et al. Panbio antigen rapid test is reliable to diagnose SARS-CoV-2 infection in the first 7 days after the onset of symptoms. *Journal of Clinical Virology*. 2020;133:104659.
117. Lindner AK, Nikolai O, Rohardt C, Burock S, Hülso C, Bölke A, et al. Head-to-head comparison of SARS-CoV-2 antigen-detecting rapid test with professional-collected nasal versus nasopharyngeal swab. *Eur Respir J*. 2021 May 6;57(5):2004430. doi: 10.1183/13993003.04430-2020.
118. Lindner AK, Nikolai O, Rohardt C, Kausch F, Wintel M, Gertler M, et al.. Diagnostic accuracy and feasibility of patient self-testing with a SARS-CoV-2 antigen-detecting rapid test. *J Clin Virol*. 2021 Aug;141:104874. doi: 10.1016/j.jcv.2021.104874.
119. Liotti F, Menchinelli G, Lalle E, Palucci I, Marchetti S, Colavita F et al. Performance of a novel diagnostic assay for rapid SARS-CoV-2 antigen detection in nasopharynx samples. *Clinical Microbiology and Infection*. 2021;27(3):487-488.
120. Mak G, Cheng P, Lau S, Wong K, Lau C, Lam E et al. Evaluation of rapid antigen test for detection of SARS-CoV-2 virus. *Journal of Clinical Virology*. 2020;129:104500.
121. Maniscalco M, Ambrosino P, Ciullo A, Fuschillo S, Valente V, Gaudiosi C et al. A Rapid Antigen Detection Test to Diagnose SARS-CoV-2 Infection Using Exhaled Breath Condensate by A Modified Inflammacheck® Device. *Sensors*. 2021;21(17):5710.
122. Martín-Sánchez V, Fernández-Villa T, Carvajal Urueña A, Rivero Rodríguez A, Reguero Celada S, Sánchez Antolín G et al. Role of Rapid Antigen Testing in Population-Based SARS-CoV-2 Screening. *Journal of Clinical Medicine*. 2021;10(17):3854.

123. Matsuda EM, de Campos IB, de Oliveira IP, Colpas DR, Carmo AMDS, Brígido LFM. Field evaluation of COVID-19 antigen tests versus RNA based detection: Potential lower sensitivity compensated by immediate results, technical simplicity, and low cost. *J Med Virol.* 2021 Jul;93(7):4405-4410. doi: 10.1002/jmv.26985. Epub 2021 Apr 8. PMID: 33788270; PMCID: PMC8250877.
124. Mboma O, Rieke E, Ahmad-Nejad P, Wirth S, Aydin M. Diagnostic Performance of SARS-CoV-2 Rapid Antigen Test in a Large, German Cohort. *Children (Basel).* 2021 Aug 8;8(8):682. doi: 10.3390/children8080682. PMID: 34438573; PMCID: PMC8394520.
125. McKay S, Tobolowsky F, Moritz E, Hatfield K, Bhatnagar A, LaVoie S et al. Performance Evaluation of Serial SARS-CoV-2 Rapid Antigen Testing During a Nursing Home Outbreak. *Annals of Internal Medicine.* 2021;174(7):945-951.
126. Merino-Amador P, González-Donapetry P, Domínguez-Fernández M, González-Romo F, Sánchez-Castellano M, Seoane-Estevez A et al. Clinitest rapid COVID-19 antigen test for the diagnosis of SARS-CoV-2 infection: A multicenter evaluation study. *Journal of Clinical Virology.* 2021;143:104961.
127. Mertens P, De Vos N, Martiny D, Jassoy C, Mirazimi A, Cuypers L, et al. Development and potential usefulness of the COVID-19 Ag Respi-Strip diagnostic assay in a pandemic context. *Front Med (Lausanne).* 2020;7:225.. doi: 10.3389/fmed.2020.00225. PMID: 32574326; PMCID: PMC7227790.
128. Möckel M, Corman V, Stegemann M, Hofmann J, Stein A, Jones T et al. SARS-CoV-2 antigen rapid immunoassay for diagnosis of COVID-19 in the emergency department. *Biomarkers.* 2021;26(3):213-220.
129. Fernandez-Montero A, Argemi J, Rodríguez J, Ariño A, Moreno-Galarraga L. Validation of a Rapid Antigen Test as a Screening Tool for SARS-CoV-2 Infection in Asymptomatic Populations. Sensitivity, Specificity and Predictive Values. *SSRN Electronic Journal.* 2021;.
130. Muhi S, Tayler N, Hoang T, et al. Multi-site assessment of rapid, point-of-care antigen testing for the diagnosis of SARS-CoV-2 infection in a low-prevalence setting: A validation and implementation study. *Lancet Reg Health West Pac.* 2021;9:100115. doi:10.1016/j.lanwpc.2021.100115
131. Nagura-Ikeda M, Imai K, Tabata S, Miyoshi K, Murahara N, Mizuno T, et al. Clinical Evaluation of Self-Collected Saliva by Quantitative Reverse Transcription-PCR (RT-qPCR), Direct RT-qPCR, Reverse Transcription-Loop-Mediated Isothermal Amplification, and a Rapid Antigen Test To Diagnose COVID-19. *J Clin Microbiol.* 2020 Aug 24;58(9):e01438-20. doi: 10.1128/JCM.01438-20. PMID: 32636214; PMCID: PMC7448663.
132. Nalumansi A, Lutalo T, Kayiwa J, Watera C, Balinandi S, Kiconco J, et al. Field evaluation of the performance of a SARS-CoV-2 antigen rapid diagnostic test in Uganda using nasopharyngeal samples. *Int J Infect Dis.* 2021;104:282–6. doi: 10.1016/j.ijid.2020.10.073. Epub 2020 Oct 30. PMID: 33130198; PMCID: PMC7836828.
133. Nash B, Badea A, Reddy A, Bosch M, Salcedo N, Gomez A et al. Validating and modeling the impact of high-frequency rapid antigen screening on COVID-19 spread and outcomes. 2020;.
134. Nikolai O, Rohardt C, Tobian F, Junge A, Corman V, Jones T et al. Anterior nasal versus nasal mid-turbinate sampling for a SARS-CoV-2 antigen-detecting rapid test: does localisation or professional collection matter?. *Infectious Diseases.* 2021;:1-6.
135. Nomoto H, Yamamoto K, Yamada G, Suzuki M, Kinoshita N, Takasaki J et al. Time-course evaluation of the quantitative antigen test for severe acute respiratory syndrome coronavirus 2: The potential contribution to alleviating isolation of COVID-19 patients. *Journal of Infection and Chemotherapy.* 2021;27(11):1669-1673.
136. Nordgren J, Sharma S, Olsson H, Jämtberg M, Falkeborn T, Svensson L et al. SARS-CoV-2 rapid antigen test: High sensitivity to detect infectious virus. *Journal of Clinical Virology.* 2021;140:104846.
137. Nörz D, Olearo F, Perisic S, Bauer M, Riester E, Schneider T et al. Multicenter Evaluation of a Fully Automated High-Throughput SARS-CoV-2 Antigen Immunoassay. *Infectious Diseases and Therapy.* 2021;.
138. Ngo Nsoga MT, Kronig I, Perez Rodriguez FJ, et al. Diagnostic accuracy of Panbio rapid antigen tests on oropharyngeal swabs for detection of SARS-CoV-2. *PLoS One.* 2021;16(6):e0253321. Published 2021 Jun 24. doi:10.1371/journal.pone.0253321
139. Oh S-M, Jeong H, Chang E, Choe PG, Kang CK, Park WB, et al. Clinical application of the standard Q COVID-19 Ag test for the detection of SARS-CoV-2 infection. *J Korean Med Sci.* 2021;36(14). doi: 10.3346/jkms.2021.36.e101.

140. Okoye NC, Barker AP, Curtis K, Orlandi RR, Snavely EA, Wright C, et al. Performance Characteristics of BinaxNOW COVID-19 Antigen Card for Screening Asymptomatic Individuals in a University Setting. *J Clin Microbiol*. 2021 Mar 19;59(4):e03282-20. doi: 10.1128/JCM.03282-20.
141. Olearo F, Nörz D, Heinrich F, Sutter JP, Roedl K, Schultze A, et al.. Handling and accuracy of four rapid antigen tests for the diagnosis of SARS-CoV-2 compared to RT-qPCR. *J Clin Virol*. 2021 Apr;137:104782. doi: 10.1016/j.jcv.2021.104782.
142. Orsi A, Pennati BM, Bruzzone B, Ricucci V, Ferone D, Barbera P, et al. On-field evaluation of a ultra-rapid fluorescence immunoassay as a frontline test for SARS-CoV-2 diagnostic. *J Virol Methods*. 2021;295(114201):114201.. doi: 10.1016/j.jviromet.2021.114201. Epub 2021 May 28. PMID: 34058287; PMCID: PMC8161776.
143. Osmanodja B, Budde K, Zickler D, Naik M, Hofmann J, Gertler M et al. Accuracy of a Novel SARS-CoV-2 Antigen-Detecting Rapid Diagnostic Test from Standardized Self-Collected Anterior Nasal Swabs. *Journal of Clinical Medicine*. 2021;10(10):2099.
144. Osterman A, Baldauf HM, Eleteby M, et al. Evaluation of two rapid antigen tests to detect SARS-CoV-2 in a hospital setting. *Med Microbiol Immunol*. 2021;210(1):65-72. doi:10.1007/s00430-020-00698-8
145. Osterman A, Iglhaut M, Lehner A, Späth P, Stern M, Autenrieth H et al. Comparison of four commercial, automated antigen tests to detect SARS-CoV-2 variants of concern. *Medical Microbiology and Immunology*. 2021;.
146. Paul D, Gupta A, Rooge S, Gupta E. Performance evaluation of automated chemiluminescence immunoassay based antigen detection – Moving towards more reliable ways to predict SARS-CoV-2 infection. *Journal of Virological Methods*. 2021;298:114299.
147. Peña M, Ampuero M, Garcés C, Gaggero A, García P, Velasquez M et al. Performance of SARS-CoV-2 rapid antigen test compared with real-time RT-PCR in asymptomatic individuals. *International Journal of Infectious Diseases*. 2021;107:201-204.
148. Peña-Rodríguez M, Viera-Segura O, García-Chagollán M, Zepeda-Nuño JS, Muñoz-Valle JF, Mora-Mora J, et al. Performance evaluation of a lateral flow assay for nasopharyngeal antigen detection for SARS-CoV-2 diagnosis. *J Clin Lab Anal*. 2021;35(5):e23745. <https://doi.org/10.1002/jcla.23745>
149. Public Health England. Preliminary report from the Joint PHE Porton Down & University of Oxford SARS-CoV-2 test development and validation cell : Rapid evaluation of Lateral Flow Viral Antigen detection devices (LFDs) for mass community testing : 2020;(November):2–7.
150. Pickering S, Batra R, Merrick B, Snell L, Nebbia G, Douthwaite S et al. Comparative performance of SARS-CoV-2 lateral flow antigen tests and association with detection of infectious virus in clinical specimens: a single-centre laboratory evaluation study. *The Lancet Microbe*. 2021;2(9):e461-e471.
151. Pilarowski G, Marquez C, Rubio L, Peng J, Martinez J, Black D, et al. Field performance and public health response using the BinaxNOW TM Rapid SARS-CoV-2 antigen detection assay during community-based testing. *Clin Infect Dis*. 2020;73(9):e3098–101.
152. Pollock NR, Jacobs JR, Tran K, Cranston AE, Smith S, O’Kane CY, et al. Performance and implementation evaluation of the Abbott BinaxNOW rapid antigen test in a high-throughput drive-through community testing site in Massachusetts. *J Clin Microbiol*. 2021;59(5). Available from: <http://dx.doi.org/10.1128/JCM.00083-21>
153. Pollock NR, Tran K, Jacobs JR, Cranston AE, Smith S, O’Kane CY, et al. Performance and operational evaluation of the Access Bio CareStart rapid Antigen test in a high-throughput drive-through community testing site in Massachusetts. *Open Forum Infect Dis*. 2021;8(7):ofab243. <https://doi.org/10.1093/ofid/ofab243>
154. Porte L, Legarraga P, Vollrath V, Aguilera X, Munita J, Araos R et al. Evaluation of a novel antigen-based rapid detection test for the diagnosis of SARS-CoV-2 in respiratory samples. *International Journal of Infectious Diseases*. 2020;99:328-333.
155. Porte L, Legarraga P, Iruretagoyena M, Vollrath V, Pizarro G, Munita J et al. Evaluation of two fluorescence immunoassays for the rapid detection of SARS-CoV-2 antigen—new tool to detect infective COVID-19 patients. *PeerJ*. 2021;9:e10801.
156. Pray IW, Ford L, Cole D, Lee C, Bigouette JP, Abedi GR, et al. Performance of an antigen-based test for asymptomatic and symptomatic SARS-CoV-2 testing at two university campuses - Wisconsin, September-October 2020. *MMWR Morb Mortal Wkly Rep*. 2021;69(5152):1642–7 doi: <http://dx.doi.org/10.15585/mmwr.mm695152a3>
157. Prince-Guerra JL, Almendares O, Nolen LD, Gunn JKL, Dale AP, Buono SA, et al. Evaluation of Abbott BinaxNOW rapid antigen test for SARS-CoV-2 infection at two community-based testing

- sites - Pima County, Arizona, November 3-17, 2020. *MMWR Morb Mortal Wkly Rep*. 2021;70(3):100–5. doi: <http://dx.doi.org/10.15585/mmwr.mm7003e3>
158. Ristić M, Nikolić N, Čabarkapa V, Turkulov V, Petrović V. Validation of the STANDARD Q COVID-19 antigen test in Vojvodina, Serbia. *PLoS One*. 2021;16(2 February):1–13. doi: 10.1371/journal.pone.0247606
 159. Rottenstreich A, Zarbiv G, Kabiri D, Porat S, Sompolinsky Y, Reubinoff B, et al. Rapid antigen detection testing for universal screening for severe acute respiratory syndrome coronavirus 2 in women admitted for delivery. *Am J Obstet Gynecol*. 2021;224(5):539–40. doi: 10.1016/j.ajog.2021.01.002
 160. Carbonell-Sahuquillo S, Lázaro-Carreño MI, Camacho J, Barrés-Fernández A, Albert E, Torres I, et al. Evaluation of a rapid antigen detection test (Panbio™ COVID-19 Ag Rapid Test Device) as a point-of-care diagnostic tool for COVID-19 in a pediatric emergency department. *J Med Virol*. 2021;93(12):6803–7. doi: 10.1002/jmv.27220
 161. Salvagno G, Gianfilippi G, Fiorio G, Pighi L, De Nitto S, Henry B et al. Performance of Fujirebio Espline SARS-CoV-2 Rapid Antigen Test for Identifying Potentially Infectious Individuals. *SSRN Electronic Journal*. 2021;.
 162. Schildgen V, Demuth S, Lüsebrink J, Schildgen O. Limits and opportunities of SARS-CoV-2 antigen rapid tests: An experienced-based perspective. *Pathogens*. 2021;10(1):38. doi: 10.3390/pathogens10010038
 163. Schuit E, Veldhuijzen IK, Venekamp RP, et al. Diagnostic accuracy of rapid antigen tests in asymptomatic and presymptomatic close contacts of individuals with confirmed SARS-CoV-2 infection: cross sectional study. *BMJ*. 2021;374:n1676. Published 2021 Jul 27. doi:10.1136/bmj.n1676
 164. Scohy A, Anantharajah A, Bodéus M, Kabamba-Mukadi B, Verroken A, Rodriguez-Villalobos H. Low performance of rapid antigen detection test as frontline testing for COVID-19 diagnosis. *J Clin Virol*. 2020;129(104455):104455. doi: doi.org/10.1016/j.jcv.2020.104455
 165. Seitz T, Schindler S, Winkelmeyer P, Zach B, Wenisch C, Zoufaly A, et al. Evaluation of rapid antigen tests based on saliva for the detection of SARS-CoV-2. *J Med Virol*. 2021;93(7):4161–2. doi: 10.1002/jmv.26983
 166. Seynaeve Y, Heylen J, Fontaine C, et al. Evaluation of Two Rapid Antigenic Tests for the Detection of SARS-CoV-2 in Nasopharyngeal Swabs. *J Clin Med*. 2021;10(13):2774. Published 2021 Jun 24. doi:10.3390/jcm10132774
 167. Shah M, Salvatore P, Ford L, Kamitani E, Whaley M, Mitchell K et al. Performance of Repeat BinaxNOW Severe Acute Respiratory Syndrome Coronavirus 2 Antigen Testing in a Community Setting, Wisconsin, November 2020–December 2020. *Clinical Infectious Diseases*. 2021;73(Supplement_1):S54–S57.
 168. Shaikh N, Friedlander E, Tate P, Liu H, Chang C, Wells A et al. Performance of a Rapid SARS-CoV-2 Antigen Detection Assay in Symptomatic Children. *Pediatrics*. 2021;148(3):e2021050832.
 169. Shrestha B, Neupane AK, Pant S, Shrestha A, Bastola A, Rajbhandari B, et al. Sensitivity and specificity of lateral flow antigen test kits for COVID-19 in asymptomatic population of Quarantine centre of Province 3. *Kathmandu Univ Med J (KUMJ)*. 2020;18(2):36–9.
 170. Soleimani R, Deckers C, Huang T, Bogaerts P, Evrard S, Wallemme I et al. Rapid COVID-19 antigenic tests: Usefulness of a modified method for diagnosis. *Journal of Medical Virology*. 2021;93(9):5655–5659.
 171. Sood N, Shetgiri R, Rodriguez A, Jimenez D, Treminino S, Daflos A, et al. Evaluation of the Abbott BinaxNOW rapid antigen test for SARS-CoV-2 infection in children: Implications for screening in a school setting. *PLoS One*. 2021;16(4):e0249710. doi: 10.1371/journal.pone.0249710.
 172. Stohr JJM, Zwart VF, Goderski G, Meijer A, Nagel-Imming CRS, Kluytmans-van den Bergh MFQ, et al. Self-testing for the detection of SARS-CoV-2 infection with rapid antigen tests for people with suspected COVID-19 in the community. *Clin Microbiol Infect [Internet]*. 2021; Available from: <http://dx.doi.org/10.1016/j.cmi.2021.07.039>.
 173. Stokes NL, Reed KA, Berbari EF, Vetter S, Binnicker MJ. Evaluation of the BinaxNOW COVID-19 rapid antigen test in an asymptomatic patient population undergoing pre-procedural screening. *J Clin Microbiol*. 2021;JCM0165021.
 174. Stokes W, Berenger BM, Portnoy D, Scott B, Szelewicki J, Singh T, et al. Clinical performance of the Abbott Panbio with nasopharyngeal, throat, and saliva swabs among symptomatic individuals with COVID-19. *Eur J Clin Microbiol Infect Dis*. 2021;40(8):1721–6. doi: 10.1007/s10096-021-04202-9.

175. Takeda Y, Mori M, Omi K. SARS-CoV-2 qRT-PCR Ct value distribution in Japan and possible utility of rapid antigen testing kit [Internet]. bioRxiv. 2020. Available from: <http://dx.doi.org/10.1101/2020.06.16.20131243>
176. Thakur P, Saxena S, Manchanda V, Rana N, Goel R, Arora R. Utility of Antigen-Based Rapid Diagnostic Test for Detection of SARS-CoV-2 Virus in Routine Hospital Settings. *Laboratory Medicine*. 2021;.
177. Torres I, Poujois S, Albert E, Colomina J, Navarro D. Evaluation of a rapid antigen test (Panbio™ COVID-19 Ag rapid test device) for SARS-CoV-2 detection in asymptomatic close contacts of COVID-19 patients. *Clin Microbiol Infect*. 2021 Apr;27(4):636.e1-636.e4. doi: 10.1016/j.cmi.2020.12.022.
178. Tsai S, Lee W, Chen P, Hung S. Real world clinical performance of SARS-CoV-2 rapid antigen tests in suspected COVID-19 cases in Taiwan. *Journal of the Formosan Medical Association*. 2021;120(11):2042-2043.
179. Turcato G, Zaboli A, Pfeifer N, Ciccariello L, Sibilio S, Tezza G, et al. Clinical application of a rapid antigen test for the detection of SARS-CoV-2 infection in symptomatic and asymptomatic patients evaluated in the emergency department: A preliminary report. *J Infect*. 2021;82(3):e14–6.
180. Uwamino Y, Nagata M, Aoki W, Nakagawa T, Inose R, Yokota H, et al. Accuracy of rapid antigen detection test for nasopharyngeal swab specimens and saliva samples in comparison with RT-PCR and viral culture for SARS-CoV-2 detection. *J Infect Chemother*. 2021;27(7):1058–62.
181. Van der Moeren N, Zwart VF, Lodder EB, Van den Bijllaardt W, Van Esch HRJM, Stohr JJJM, et al. Evaluation of the test accuracy of a SARS-CoV-2 rapid antigen test in symptomatic community dwelling individuals in the Netherlands. *PLoS One* [Internet]. 2021;16(5 May):1–11. Available from: <http://dx.doi.org/10.1371/journal.pone.0250886>
182. Van Honacker E, Van Vaerenbergh K, Boel A, De Beenhouwer H, Leroux-Roels I, Cattoir L. Comparison of five SARS-CoV-2 rapid antigen detection tests in a hospital setting and performance of one antigen assay in routine practice: a useful tool to guide isolation precautions?. *Journal of Hospital Infection*. 2021;114:144-152.
183. Veyrenche N, Bolloré K, Pisoni A, Bedin A-S, Mondain A-M, Ducos J, et al. Diagnosis value of SARS-CoV-2 antigen/antibody combined testing using rapid diagnostic tests at hospital admission [Internet]. bioRxiv. 2020. Available from: <http://dx.doi.org/10.1101/2020.09.19.20197855>
184. Von Ahnen T, von Ahnen M, Wirth U, Schardey H, Herdtle S. Evaluation of a rapid-antigen test for COVID-19 in an asymptomatic collective. *Wiener Medizinische Wochenschrift*. 2021;.
185. Wachinger J, Olaru ID, Horner S, Schnitzler P, Heeg K, Denking CM. The potential of SARS-CoV-2 antigen-detection tests in the screening of asymptomatic persons [published online ahead of print, 2021 Jul 26]. *Clin Microbiol Infect*. 2021;S1198-743X(21)00412-2. doi:10.1016/j.cmi.2021.07.020
186. Wagenhäuser I, Knies K, Rauschenberger V, Eisenmann M, McDonogh M, Petri N, et al. Clinical performance evaluation of SARS-CoV-2 rapid antigen testing in point of care usage in comparison to RT-qPCR. *EBioMedicine*. 2021;69(103455):103455. *EBioMedicine*. 2021 Jul;69:103455. doi: 10.1016/j.ebiom.2021.103455. Epub 2021 Jun 26. PMID: 34186490; PMCID: PMC8234263.
187. Weitzel T, Legarraga P, Iruretagoyena M, Pizarro G, Vollrath V, Araos R, et al. Head-to-head comparison of four antigen-based rapid detection tests for the diagnosis of SARS-CoV-2 in respiratory samples [Internet]. bioRxiv. 2020. Available from: <http://dx.doi.org/10.1101/2020.05.27.119255>
188. Yokota I, Sakurazawa T, Sugita J, Iwasaki S, Yasuda K, Yamashita N, Fujisawa S, Nishida M, Konno S, Teshima T. Performance of Qualitative and Quantitative Antigen Tests for SARS-CoV-2 Using Saliva. *Infect Dis Rep*. 2021 Aug 24;13(3):742-747. doi: 10.3390/idr13030069.
189. Young S, Taylor SN, Cammarata CL, Roger-Dalbert C, Montano A, Griego-Fullbright C, et al. Clinical evaluation of BD Veritor™ SARS-CoV-2 point-of-care test performance compared to PCR-based testing and versus the Sofia® 2 SARS Antigen point-of-care test [Internet]. bioRxiv. 2020. Available from: <http://dx.doi.org/10.1101/2020.09.01.20185777>.
190. Study to Evaluate the Performance of the Thermo COVID-19 Rapid Antigen Test for Detection of SARS-CoV-2 [Internet]. *Clinicaltrials.gov*. [cited 2021 Nov 4]. Available from: <https://clinicaltrials.gov/ct2/show/NCT04878068?term=rapid+antigen+tests&cond=COVID-19&intr=rapid+antigen+test&draw=2&rank=1>
191. Klein JAF, Krüger LJ, Tobian F, Gaeddert M, Lainati F, Schnitzler P, et al. Head-to-head performance comparison of self-collected nasal versus professional-collected nasopharyngeal

- swab for a WHO-listed SARS-CoV-2 antigen-detecting rapid diagnostic test. *Med Microbiol Immunol* [Internet]. 2021;210(4):181–6. Available from: <https://doi.org/10.1007/s00430-021-00710-9> doi: 10.1007/s00430-021-00710-9
192. Lindner AK, Nikolai O, Rohardt C, Burock S, Hülso C, Bölke A, Gertler M, Krüger LJ, Gaeddert M, Tobian F, Lainati F, Seybold J, Jones TC, Hofmann J, Sacks JA, Mockenhaupt FP, Denkinge CM. Head-to-head comparison of SARS-CoV-2 antigen-detecting rapid test with professional-collected nasal versus nasopharyngeal swab. *Eur Respir J*. 2021 May 6;57(5):2004430. doi: 10.1183/13993003.04430-2020.
 193. Lindner AK, Nikolai O, Rohardt C, Kausch F, Wintel M, Gertler M, Burock S, Horig M, Bernhard J, Tobian F, Gaeddert M, Lainati F, Corman VM, Jones TC, Sacks JA, Seybold J, Denkinge CM, Mockenhaupt FP. Diagnostic accuracy and feasibility of patient self-testing with a SARS-CoV-2 antigen-detecting rapid test. *J Clin Virol*. 2021 Aug;141:104874. doi: 10.1016/j.jcv.2021.104874.
 194. Osmanodja B, Budde K, Zickler D, Naik M, Hofmann J, Gertler M et al. Accuracy of a Novel SARS-CoV-2 Antigen-Detecting Rapid Diagnostic Test from Standardized Self-Collected Anterior Nasal Swabs. *Journal of Clinical Medicine*. 2021;10(10):2099.
 195. García-Fiñana M, Hughes DM, Cheyne CP, Burnside G, Stockbridge M, Fowler TA, et al. Performance of the Innova SARS-CoV-2 antigen rapid lateral flow test in the Liverpool asymptomatic testing pilot: Population based cohort study. *BMJ*. 2021;374:n1367. doi: 10.1136/bmj.n1367
 196. Frediani JK, Levy JM, Rao A, et al. Multidisciplinary assessment of the Abbott BinaxNOW SARS-CoV-2 point-of-care antigen test in the context of emerging viral variants and self-administration. *Sci Rep*. 2021;11(1):14604. Published 2021 Jul 16. doi:10.1038/s41598-021-94055-1
 197. Maniscalco M, Ambrosino P, Ciullo A, Fuschillo S, Valente V, Gaudiosi C et al. A Rapid Antigen Detection Test to Diagnose SARS-CoV-2 Infection Using Exhaled Breath Condensate by A Modified Inflammacheck® Device. *Sensors*. 2021;21(17):5710.
 198. Ruszkiewicz D, Sanders D, O'Brien R, Hempel F, Reed M, Riepe A et al. Diagnosis of COVID-19 by analysis of breath with gas chromatography-ion mobility spectrometry - a feasibility study. *EClinicalMedicine*. 2020;29-30:100609. doi: 10.1016/j.eclim.2020.100609
 199. Maniscalco M, Ambrosino P, Ciullo A, Fuschillo S, Valente V, Gaudiosi C et al. A Rapid Antigen Detection Test to Diagnose SARS-CoV-2 Infection Using Exhaled Breath Condensate by A Modified Inflammacheck® Device. *Sensors*. 2021;21(17):5710. doi: 10.3390/s21175710
 200. Steppert C, Steppert I, Sterlacci W, Bollinger T. Rapid detection of SARS-CoV-2 infection by multicapillary column coupled ion mobility spectrometry (MCC-IMS) of breath. A proof of concept study. 2020;. doi: 10.1101/2020.06.30.20143347
 201. Wintjens A, Hintzen K, Engelen S, Lubbers T, Savelkoul P, Wesseling G et al. Applying the electronic nose for pre-operative SARS-CoV-2 screening. *Surgical Endoscopy*. 2020;. doi: 10.1007/s00464-020-08169-0
 202. Grassin-Delyle S, Roquencourt C, Moine P, Saffroy G, Carn S, Heming N et al. Metabolomics of exhaled breath in critically ill COVID-19 patients: A pilot study. *EBioMedicine*. 2021;63:103154. doi: 10.1016/j.ebiom.2020.103154
 203. de Vries R, Vigeveno R, Mulder S, Farzan N, Vintges D, Goeman J et al. Ruling out SARS-CoV-2 infection using exhaled breath analysis by electronic nose in a public health setting. 2021;. doi: 10.1101/2021.02.14.21251712
 204. Abdalhamid B, Bilder CR, McCutchen EL, Hinrichs SH, Koepsell SA, Iwen PC. Assessment of Specimen Pooling to Conserve SARS CoV-2 Testing Resources. *Am J Clin Pathol*. 2020;153(6):715–8.
 205. Alcoba-Florez J, Gil-Campesino H, García-Martínez de Artola D, Díez-Gil O, Valenzuela-Fernández A, González-Montelongo R, Ciuffreda L, Flores C. Increasing SARS-CoV-2 RT-qPCR testing capacity by sample pooling. *Int J Infect Dis*. 2021 Feb;103:19-22. doi: 10.1016/j.ijid.2020.11.155. Epub 2020 Nov 19. PMID: 33220439; PMCID: PMC7674967.
 206. Alizad-Rahvar AR, Vafadar S, Totonchi M, Sadeghi M. False-negative mitigation in group testing for COVID-19 screening. *medRxiv*. 2020;1–18.
 207. Barak N, Ben-Ami R, Sido T, Perri A, Shtoyer A, Rivkin M, Licht T, Peretz A, Magenheimer J, Fogel I, Livneh A, Daich Y, Oiknine-Djian E, Benedek G, Dor Y, Wolf DG, Yassour M; Hebrew University-Hadassah COVID-19 Diagnosis Team. Lessons from applied large-scale pooling of 133,816 SARS-CoV-2 RT-PCR tests. *Sci Transl Med*. 2021 Apr 14;13(589):eabf2823. doi: 10.1126/scitranslmed.abf2823. Epub 2021 Feb 22. PMID: 33619081; PMCID: PMC8099176.
 208. Cesselli D, Bulfoni M, Marzinotto S, et al. Implementation and validation of a pooling strategy for a sustainable screening campaign for the presence of SARS-CoV-2. *medRxiv* [Internet]. 2020;2020.08.28.20174946. Available from: <https://doi.org/10.1101/2020.08.28.20174946>

209. Chhikara K, Kanta P, Ghosh A, Prakash RC, Goyal K, Singh MP. Validation of SARS CoV-2 detection by real-time PCR in matched pooled and deconvoluted clinical samples before and after nucleic acid extraction: a study in tertiary care hospital of North India. *Diagn Microbiol Infect Dis*. 2020;99(January):115206.
210. Chong BSW, Tran T, Druce J, Ballard SA, Simpson JA, Catton M. Sample pooling is a viable strategy for SARS-CoV-2 detection in low-prevalence settings. *Pathology*. 2020;52(7):796–800.
211. Christoff AP, Cruz GNF, Sereia AFR, et al. Swab Pooling for Large-Scale RT-qPCR Screening of SARS-CoV-2. *SSRN Electron J*. 2020.
212. Das S, Lau AF, Youn JH, Khil PP, Zelazny AM, Frank KM. Pooled Testing for Surveillance of SARS-CoV-2 in Asymptomatic Individuals. *J Clin Virol*. 2020 Nov;132:104619. doi: 10.1016/j.jcv.2020.104619. Epub 2020 Sep 3. PMID: 32920431; PMCID: PMC7470739.
213. de Salazar A, Aguilera A, Trastoy R, et al. Sample pooling for SARS-CoV-2 RT-PCR screening. *Clin Microbiol Infect*. 2020;26.
214. Garg A, Ghoshal U, Patel SS, Singh DV, Arya AK, Vasanth S, Pandey A, Srivastava N. Evaluation of seven commercial RT-PCR kits for COVID-19 testing in pooled clinical specimens. *J Med Virol*. 2021 Apr;93(4) 2281–2286. doi:10.1002/jmv.26691. PMID: 33230819; PMCID: PMC7753435.
215. Gavars D, Perminovs D, Stasulans J, et al. Saliva as testing sample for SARS-CoV-2 detection by RT-PCR in low prevalence community setting [Preprint]. *Medrxiv* [Internet]. 2020;2020.10.20.20216127. Available from: <http://medrxiv.org/content/early/2020/10/21/2020.10.20.20216127.abstract>
216. Hogan CA, Sahoo MK, Pinsky BA. Sample Pooling as a Strategy to Detect Community Transmission of SARS-CoV-2: Letter. *JAMA*. 323(1967–9).
217. Li H, Sun K, Persing DH, Tang Y-W, Shen D. Real-time Screening of Specimen Pools for Coronavirus Disease 2019 (COVID-19) Infection at Sanya Airport, Hainan Island, China. *Clin Infect Dis*. 2020;2019(Figure 1):2019–21.
218. Lo R, Barrientos A, Espiritu B, et al. An Evaluation of Pooling Strategies for qRT-PCR Testing for SARS-CoV-2 Infection: A pragmatic multi-site parallel operational study by The Philippine Society of Pathologists, Inc. *Philipp J Pathol*. 2020;5(2):1–22.
219. Lüsebrink J, Schildgen V, Schildgen O. Pooling is an insufficient strategy to avoid health care staff to patient transmission of SARS-CoV-2. *Infect Control Hosp Epidemiol*. 2020;1–3.
220. Mastrianni D, Falivena R, Brooks T, et al. Pooled Testing for SARS-CoV-2 in Hospitalized Patients. *J Hosp Med*. 2020;15(9):538–9.
221. Pasomsub E, Watcharananan SP, Watthanachockchai T, et al. Saliva sample pooling for the detection of SARS-CoV-2. *J Med Virol* [Internet]. 2020;0–3. Available from: <http://dx.doi.org/10.1002/jmv.26460>
222. Singh AK, Nema RK, Joshi A, et al. Evaluation of pooled sample analysis strategy in expediting case detection in areas with emerging outbreaks of COVID-19: A pilot study. *PLoS One* [Internet]. 2020;15(9 September 2020):1–8. Available from: <http://dx.doi.org/10.1371/journal.pone.0239492>
223. Wang H, Hogan CA, Miller JA, et al. Performance of Nucleic Acid Amplification Tests for Detection of Severe Acute Respiratory Syndrome Coronavirus 2 in Prospectively Pooled Specimens. *Emerg Infect Dis* [Internet]. 2020;27(1). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/33183494>
224. Mohanty S, Ravindra A, Gupta K, et al. Intricacies in characterizing positivity in pooled sample testing for SARS-CoV-2. *J Med Virol*. 2020;0–2.
225. Ai J, Gong J, Xing L, et al. 2020. “Analysis of Factors Associated Early Diagnosis in Coronavirus Disease 2019 (COVID-19).” *MedRxiv*. medRxiv. doi:10.1101/2020.04.09.20059352.
226. Ai J, Zhang H, Xu T, et al. 2020. “Optimizing Diagnostic Strategy for Novel Coronavirus Pneumonia, a Multi-Center Study in Eastern China.” *MedRxiv*. medRxiv.
227. Singanayagam A. et al. Duration of infectiousness and correlation with RT-PCR cycle threshold values in cases of COVID-19, England, January to May 2020. *Euro Surveill*. 2020;25(32):pii=2001483. Available from <https://doi.org/10.2807/1560-7917.ES.2020.25.32.2001483> [Accessed 24 September 2021]
228. La Scola, B. et al. Viral RNA load as determined by cell culture as a management tool for discharge of SARS-CoV-2 patients from infectious disease (Brief Report). *Eur J Clin Microbiol Infect Dis* (2020) 39:1059–1061. Available at <https://doi.org/10.1007/s10096-020-03913-9> [Accessed 24 September 2021]
229. Laferl, H. et al. An approach to lifting self-isolation for health care workers with prolonged shedding of SARS-CoV-2 RNA. *Infection* (2021) 49:95–101. Available at <https://doi.org/10.1007/s15010-020-01530-4>. [Accessed 24 September 2021]
230. Piralla, A. et al. Residual SARS-CoV-2 RNA in nasal swabs of convalescent COVID-19:

231. Is prolonged quarantine always justified? *International Journal of Infectious Diseases* 2021 102:299-302. Available at <https://doi.org/10.1016/j.ijid.2020.10.072> [Accessed 01 November 2021]
232. Ladhani, et al. Investigation of SARS-CoV-2 outbreaks in six care homes in London, April 2020. *EClinicalMedicine* (26)2020 100533. Available at <https://doi.org/10.1016/j.eclinm.2020.100533>. [Accessed 31 October 2021].
233. Folquiera, et al. Persistent SARS-CoV-2 replication in severe COVID-19. *medRxiv preprint*. Available at i: <https://doi.org/10.1101/2020.06.10.20127837.t>. [Accessed 31 October 2021]
234. Brown et al. Snapshot PCR surveillance for SARS-CoV-2 in hospital staff in England. *Journal of Infection* 81 (2020): 427-434. Available at <https://doi.org/10.1016/j.jinf.2020.06.069>. [Accessed 1 November 2021].
235. Bullard, J. et al. Predicting infectious SARS-CoV-2 from diagnostic samples. *Clin Infect Dis*. 2020;71(10):2663-2666. Available from doi:10.1093/cid/ciaa638. [Accessed 1 November 2021]
236. Gniazdowski, V. et al. Repeat COVID-19 Molecular Testing: Correlation with Recovery of Infectious Virus, Molecular Assay Cycle Thresholds, and Analytical Sensitivity. *medRxiv* 2020.08.05.20168963. Available from doi: <https://doi.org/10.1101/2020.08.05.20168963> [Accessed 1 November 2021]
237. Basile, K., et al. Cell-based culture of SARS-CoV-2 informs infectivity and safe de-isolation assessments during COVID-19. *Clinical Infectious Diseases*. Available from doi:10.1093/cid/ciaa1579 [Accessed 31 October 2021]
238. Huang, C-G. et al., Culture-based virus isolation to evaluate potential infectivity of clinical specimens tested for COVID-19. *J Clin Microbiol* 58:e01068-20. <https://doi.org/10.1128/JCM.01068-20>. [Accessed 24 September 2021]
239. Young et al. Viral Dynamics and Immune Correlates of Coronavirus Disease 2019 (COVID-19) Severity. *Clinical Infectious diseases* 2020. Available at DOI: 10.1093/cid/ciaa1280. [Accessed 1 November 2021].
240. Gruskay JA, Dvorzhinskiy A, Konnaris MA, et al. Universal Testing for COVID-19 in Essential Orthopaedic Surgery Reveals a High Percentage of Asymptomatic Infections. *J Bone Joint Surg Am*. 2020 Aug 19;102(16):1379-1388. doi: 10.2106/JBJS.20.01053. PMID: 32516279.
241. The Infectious Diseases Society of America Guidelines on the Diagnosis of COVID-19: Molecular Diagnostic Testing. 23 Dec 2020. Available from: <https://www.idsociety.org/practice-guideline/covid-19-guideline-diagnostics/>. 11
242. Timing of surgery following SARS-CoV-2 infection: an international prospective cohort study. *COVIDSurg Collaborative. Anaesthesia*. Published 09 Mar 2021. Available from: <https://associationofanaesthetists-publications.onlinelibrary.wiley.com/action/doSearch?ContribAuthorStored=COVIDSurg+Collaborative>). <https://doi.org/10.1111/anae.15458>
243. Mortgat L, Barbezange C, Fischer N, Heyndrickx L, Hutse V, Thomas I, et al. SARS-CoV-2 Prevalence and Seroprevalence among Healthcare Workers in Belgian Hospitals: Baseline Results of a Prospective Cohort Study. *medRxiv*. 2020.
244. Mulchandani R, Taylor-Philips S, Jones HE, Ades T, Borrow R, Linley E, et al. Self assessment overestimates historical COVID-19 disease relative to sensitive serological assays: Cross sectional study in UK key workers. *medRxiv*. 2020.
245. Shields AM, Faustini SE, Perez-Toledo M, Jossi S, Allen JD, Al-Taei S, et al. Serological responses to SARS-CoV-2 following non-hospitalised infection: Clinical and ethnodemographic features associated with the magnitude of the antibody response. *medRxiv*. 2020.
246. Ige O, Buru AS, Lamido TZ, Mohammed T, Dogara L, Patience II, et al. Evaluation of SARS-CoV2 antibody Rapid Diagnostic Test kits (RDTs) and Real Time-Polymerase Chain Reaction (Rt-PCR) for COVID-19 Diagnosis in Kaduna, Nigeria. 2020;2196482:1–24.
247. Gonzalez JM, Santos-Barbosa JC, Jaller C, Otalora G, Hernandez LJ, Guevara-Suarez M, et al. Seroreactivity to SARS-CoV-2 in individuals attending a university campus in Bogotá Colombia. *medRxiv*. 2020;1–13.
248. Silva VO, de Oliveira EL, Castejon MJ, Yamashiro R, Ahagon CM, López-Lopes GI, et al. Prevalence of antibodies against SARS-CoV-2 in professionals of a public health laboratory at São Paulo, SP, Brazil. *medRxiv*. 2020.
249. Lorent D, Nowak R, Roxo C, Lenartowicz E, Makarewicz A, Zaremba B, et al. 2021.
250. Fong D, San Nicolò KO, Alber M, Mitterer M. Evaluating the longitudinal effectiveness of preventive measures against COVID-19 and seroprevalence of IgG antibodies to SARS-CoV-2 in cancer outpatients and healthcare workers. *Wien Klin Wochenschr*. 2021.

251. Khan SMS, Qurieshi MA, Haq I, Majid S, Ahmad J, Ayub T, et al. Seroprevalence of SARS-CoV-2-specific IgG antibodies in Kashmir, India, 7 months after the first reported local COVID-19 case: results of a population-based seroprevalence survey from October to November 2020. *BMJ Open*. 2021;11(9):e053791.
252. Jespersen S, Mikkelsen S, Greve T, Kaspersen KA, Tolstrup M, Boldsen JK, et al. Severe Acute Respiratory Syndrome Coronavirus 2 Seroprevalence Survey Among 17 971 Healthcare and Administrative Personnel at Hospitals, Prehospital Services, and Specialist Practitioners in the Central Denmark Region. *Clin Infect Dis*. 2020;(Xx Xxx):1–8.
253. Flannery DD, Gouma S, Dhudasia MB, Mukhopadhyay S, Pfeifer MR, Woodford EC, et al. SARS-CoV-2 seroprevalence among parturient women. *medRxiv*. 2020;5709(July):1–9.
254. Carrat F, de Lamballerie X, Rahib D, Blanche H, Lapidus N, Artaud F, et al. Antibody status and cumulative incidence of SARS-CoV-2 infection among adults in three regions of France following the first lockdown and associated risk factors: a multicohort study. *International journal of epidemiology*. 2021.
255. Robinson AA V, Weaving AG, Philips BBJ, Eziefula CAC, E DKES, F TC. Real-world experience of SARS-CoV-2 antibody assays in UK healthcare workers. 2021;21(3):1–6.
256. Nasrallah GK, Dargham SR, Shurrah F, Al-Sadeq DW, Al-Jighefee H, Chemaitelly H, et al. Analytic comparison between three high-throughput commercial SARS-CoV-2 antibody assays reveals minor discrepancies in a high-incidence population. *Sci Rep*. 2021;11(1):11837.
257. Afzal N, Tariq N, Raza S, Shakeel D. Diagnostic Accuracy of Electro-Chemiluminescence Immunoassay Anti-SARS-CoV-2 Serological Test. *Cureus*. 2021;13(1):1–6.
258. Santarelli A, Lalitsasivimol D, Bartholomew N, Reid S, Reid J, Lyon C, et al. The seroprevalence of SARS-CoV-2 in a rural southwest community. *J Am Osteopath Assoc [Internet]*. 2021;121(2):199–210. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/33567087>
259. Papasavas P, Olugbile S, Wu U, Robinson K, Roberts AL, O'Sullivan DM, et al. Seroprevalence of SARS-CoV-2 antibodies, associated epidemiological factors and antibody kinetics among healthcare workers in Connecticut. *J Hosp Infect*. 2021;114:117–25.
260. Percivalle E, Cambiè G, Cassaniti I, Nepita EV, Maserati R, Ferrari A, et al. Prevalence of SARS-CoV-2 specific neutralising antibodies in blood donors from the Lodi Red Zone in Lombardy, Italy, as at 06 April 2020. *Eurosurveillance [Internet]*. 2020;25(24):1–5. Available from: <http://dx.doi.org/10.2807/1560-7917.ES.2020.25.24.2001031>
261. Caillard S, Benotmane I, Meidinger C, Jegou V, Ludwiller S, Rihon A, et al. SARS-Cov-2 Seroprevalence in a French Kidney Transplant Center Located Within a "High-risk" Zone. *Transplantation*. 2021;105(10):2165–9.
262. Jin Y, Wang M, Zuo Z, et al. Diagnostic value and dynamic variance of serum antibody in coronavirus disease 2019. *Int J Infect Dis*. 2020 May; 94:49–52. doi: 10.1016/j.ijid.2020.03.065. Epub 2020 Apr 3. PMID: 32251798; PMCID: PMC7194885.
263. Liu Ying, Liu Yue-ping, Diao Bo, et al. Diagnostic Indexes of a Rapid IgG/IgM Combined Antibody Test for SARS-CoV-2 *medRxiv* 2020.03.26.20044883; doi: <https://doi.org/10.1101/2020.03.26.20044883>
264. Paradiso A, De Summa S, Loconsole D, et al. Clinical meanings of rapid serological assay in patients tested for SARS-Co2 RT-PCR *medRxiv* 2020.04.03.20052183; doi: <https://doi.org/10.1101/2020.04.03.20052183>
265. Samuel Clifford et al.. "Effectiveness of BNT162b2 and ChAdOx1 against SARS-CoV-2 household transmission - a prospective cohort study in England." *medRxiv* , no. (2021): 2021.11.24.21266401. Accessed November 27, 2021. doi: 10.1101/2021.11.24.21266401.
266. Brechje de Gier et al.. "Vaccine effectiveness against SARS-COV-2 transmission and infections among households and other close contacts of confirmed cases, the Netherlands, February to May 2021." www.eurosurveillance.org Accessed November 27, 2021. doi: 10.2807/1560-7917.
267. Kissler SM, Fauver JR, Mack C, et al. Viral dynamics of SARS-CoV-2 variants in vaccinated and unvaccinated persons. *N Engl J Med*. 1 Dec 2021. DOI: 10.1056/NEJMc2102507.
268. Phillip P. Salvatore et al.. "Transmission potential of vaccinated and unvaccinated persons infected with the SARS-CoV-2 Delta variant in a federal prison, July–August 2021." *medRxiv* , no. (2021): 2021.11.12.21265796. Accessed November 27, 2021. doi: 10.1101/2021.11.12.21265796.
269. Anika Singanayagam et al.. "Community transmission and viral load kinetics of the SARS-COV-2 delta (B.1.617.2) variant in vaccinated and unvaccinated individuals in the UK: a prospective longitudinal, cohort study." 2021: *Lancet Infectious Disease*. doi: [doi.org/10.1016/S1473-3099\(21\)00648-4](https://doi.org/10.1016/S1473-3099(21)00648-4).

270. Rachel Epstein et al.. "SARS-CoV-2 PCR Clearance in Immunocompromising Conditions: Is Test-Based Removal From Isolation Necessary in Severely Immunocompromised Individuals?" *openforum infectious diseases*. Accessed December 9, 2021. Doi 10.1093/ofid/ofab164.
271. Borakati A, Perera A, Johnson J, Sood T. Diagnostic accuracy of X-ray versus CT in COVID-19: A propensity-matched database study. *BMJ Open*. 2020;10(11):1–14. doi: 10.1136/bmjopen-2020-042946
272. Roy Choudhury SH, Shahi PK, Sharma S, Dhar R. Utility of chest radiography on admission for initial triaging of COVID-19 in symptomatic patients. *ERJ Open Res [Internet]*. 2020;6(3):00357–2020. Available from: <http://dx.doi.org/10.1183/23120541.00357-2020>
273. Cozzi A, Schiaffino S, Arpaia F, Della Pepa G, Tritella S, Bertolotti P, et al. Chest x-ray in the COVID-19 pandemic: Radiologists' real-world reader performance. *Eur J Radiol [Internet]*. 2020 Nov;132(January):109272. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0720048X20304617>
274. Du R, Tsougenis ED, Ho JWK, Chan JKY, Chiu KWH, Fang BXH, et al. Machine learning application for the prediction of SARS-CoV-2 infection using blood tests and chest radiograph. *Sci Rep [Internet]*. 2021;11(1):1–13. Available from: <https://doi.org/10.1038/s41598-021-93719-2>
275. Gibbons RC, Magee M, Goett H, Murrett J, Genninger J, Mendez K, et al. Lung Ultrasound vs. Chest X-Ray Study for the Radiographic Diagnosis of COVID-19 Pneumonia in a High-Prevalence Population. In: *Journal of Emergency Medicine*. 2021. p. 615–25. doi: 10.1016/j.jemermed.2021.01.041
276. Gichoya JW, Sinha P, Davis M, Dunkle JW, Hamlin SA, Herr KD, et al. Multireader evaluation of radiologist performance for COVID-19 detection on emergency department chest radiographs. *Clin Imaging [Internet]*. 2021;82(May 2021):77–82. Available from: <https://doi.org/10.1016/j.clinimag.2021.10.018>
277. Hwang EJ, Kim H, Yoon SH, Goo JM, Park CM. Implementation of a deep learning-based computeraided detection system for the interpretation of chest radiographs in patients suspected for covid-19. *Korean J Radiol*. 2020;21(10):1150–60.
278. Hwang EJ, Kim KB, Kim JY, Lim JK, Nam JG, Choi H, et al. COVID-19 pneumonia on chest X-rays: Performance of a deep learning-based computer-aided detection system. *PLoS One [Internet]*. 2021;16(6 June):1–16. Available from: <http://dx.doi.org/10.1371/journal.pone.0252440>
279. Ippolito D, Pecorelli A, Maino C, Capodaglio C, Mariani I, Giandola T, et al. Diagnostic impact of bedside chest X-ray features of 2019 novel coronavirus in the routine admission at the emergency department: case series from Lombardy region. *Eur J Radiol [Internet]*. 2020;129(May):109092. Available from: <https://doi.org/10.1016/j.ejrad.2020.109092>
280. Khan SA, Manohar M, Khan M, Asad S, Adil SO. Radiological profile of patients undergoing chest x-ray and computed tomography scans during covid-19 outbreak. *Pakistan J Med Sci*. 2021;37(5):1288–94. doi: 10.12669/pjms.37.5.4290
281. Majeed AI, Raza F, Riaz SK, Kanwal J. The Essential Role Of Conventional Radiography In Covid-19; Perspective Of A Developing Country. *J Ayub Med Coll Abbottabad*. 2021;33(2):267–73.
282. Murphy K, Smits H, Knoop AJG, Korst MBJM, Samson T, Scholten ET, et al. COVID-19 on chest radiographs: A multireader evaluation of an artificial intelligence system. *Radiology*. 2020;296(3):E166–72. doi: 10.1148/radiol.2020201874
283. Pare JR, Camelo I, Mayo KC, Leo MM, Dugas JN, Nelson KP, et al. Point-of-care lung ultrasound is more sensitive than chest radiograph for evaluation of COVID-19. *West J Emerg Med*. 2020;21(4):771–8. doi: 10.5811/westjem.2020.5.47743
284. Gaia C, Maria Chiara C, Silvia L, Chiara A, Maria Luisa DC, Giulia B, et al. Chest CT for early detection and management of coronavirus disease (COVID-19): a report of 314 patients admitted to Emergency Department with suspected pneumonia. *Radiol Medica [Internet]*. 2020;125(10):931–42. Available from: <https://doi.org/10.1007/s11547-020-01256-1>
285. Peyrony O, Marbeuf-Gueye C, Truong V, Giroud M, Rivière C, Khenissi K, et al. Accuracy of Emergency Department Clinical Findings for Diagnosis of Coronavirus Disease 2019. *Ann Emerg Med [Internet]*. 2020 Oct;76(4):405–12. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7254017/pdf/main.pdf>
286. Schiaffino S, Tritella S, Cozzi A, Carriero S, Blandi L, Ferraris L, et al. Diagnostic Performance of Chest X-Ray for COVID-19 Pneumonia during the SARS-CoV-2 Pandemic in Lombardy, Italy. *J Thorac Imaging*. 2020;35(4):W105–6. doi: 10.1097/RTI.0000000000000533
287. Stephanie S, Shum T, Cleveland H, Challa SR, Herring A, Jacobson FL, et al. Determinants of chest radiography sensitivity for covid-19: A multi-institutional study in the United States. *Radiol Cardiothorac Imaging*. 2020;2(5). doi: 10.1148/ryct.2020200337

288. Stevens BJ. Reporting radiographers' interpretation and use of the British Society of Thoracic Imaging's coding system when reporting COVID-19 chest x-rays. *Radiography*. 2021;27(1):90–4. doi: 10.1016/j.radi.2020.06.010
289. Tung-Chen Y, Algora-Martín A, Llamas-Fuentes R, Rodríguez-Fuertes P, Martínez Virto AM, Sanz-Rodríguez E, et al. Point-of-care ultrasonography in the initial characterization of patients with COVID-19. *Med Clin (Barc)*. 2021;156(10):477–84. doi: 10.1016/j.medcli.2020.12.007
290. Bar S, Lecourtois A, Diouf M, Goldberg E, Bourbon C, Arnaud E, et al. The association of lung ultrasound images with COVID-19 infection in an emergency room cohort. *Anaesthesia*. 2020;75(12):1620–5. doi: 10.1111/anae.15175
291. Bitar ZI, Shamsah M, Maadarani OS, Bamasood OM, Al-foudari H. Appropriateness of lung ultrasound for the diagnosis of COVID -19 pneumonia . *Heal Sci Reports*. 2021;4(2):1–7. doi: 10.1002/hsr2.302
292. Dini FL, Bergamini C, Allegrini A, Scopelliti M, Secco G, Miccoli M, et al. Bedside wireless lung ultrasound for the evaluation of COVID-19 lung injury in senior nursing home residents. *Monaldi Arch Chest Dis*. 2020;90(3):523–7. doi: 10.4081/monaldi.2020.1446
293. Fonsi GB, Sapienza P, Brachini G, Andreoli C, De Cicco ML, Cirillo B, et al. Is Lung Ultrasound Imaging a Worthwhile Procedure for Severe Acute Respiratory Syndrome Coronavirus 2 Pneumonia Detection? *J Ultrasound Med*. 2021;40(6):1113–23. doi: 10.1002/jum.15487
294. Haak SL, Renken IJE, Jager LC, Lameijer H, Van Der Kolk BBYM. Diagnostic accuracy of point-of-care lung ultrasound in COVID-19. *Emerg Med J*. 2021;38(2):94–9. doi: 10.1136/emmermed-2020-210125
295. Lieveeld AWE, Kok B, Schuit FH, Azijli K, Heijmans J, van Laarhoven A, et al. Diagnosing COVID-19 pneumonia in a pandemic setting: Lung Ultrasound versus CT (LUVCT) – a multicentre, prospective, observational study . *ERJ Open Res [Internet]*. 2020;6(4):00539–2020. Available from: <http://dx.doi.org/10.1183/23120541.00539-2020>
296. Narinx N, Smismans A, Symons R, Frans J, Demeyere A, Gillis M. Feasibility of using point-of-care lung ultrasound for early triage of COVID-19 patients in the emergency room. *Emerg Radiol*. 2020;27(6):663–70. doi: 10.1007/s10140-020-01849-3
297. Ai T, Yang Z, Hou H, et al. Correlation of chest CT and RT-PCR testing for coronavirus disease 2019 (COVID-19) in China: a report of 1014 cases. *Radiology* 2020;296:E32–E40. doi: 10.1148/radiol.2020200642
298. Aslan S, Becki T, Cakir I, Ekiz M, Yavuz I, Sahin A. Diagnostic performance of low-dose chest CT to detect COVID-19: A Turkish population study. *Diagn Interv Radiol* 2021;27:181-7. doi: 10.5152/dir.2020.20350
299. Bağcı Z, Keçeli AM. Comparison of computed tomography (CT) findings with RT-PCR in the diagnosis of COVID-19 in children. *Pediatr Pulmonol*. 2021;56(7):1985–91. doi: 10.1002/ppul.25426
300. Barbosa P, Bitencourt A, Miranda Gd, Almeida M, Chonjniak R. Chest CT accuracy in the diagnosis of SARS-CoV-2 infection: initial experience in a cancer center. *Radiol Bras* 2020;53:211-5. doi: 10.1590/0100-3984.2020.0040
301. Bellini D, Panvini N, Rengo M, Vicini S, Lichtner M, Tieghi T, et al. Diagnostic accuracy and interobserver variability of CO-RADS in patients with suspected coronavirus disease-2019: a multireader validation study. *Eur Radiol*. 2021;31(4):1932–40.
302. Berkel B Van, Vandevenne J, Coursier K, Alberts V, Offenwert J Van, Verduyck J, et al. Chest CT Diagnosis of COVID-19: Accuracy using CO-RADS and CT-Involvement Scoring. *J Belgian Soc Radiol*. 2021;105(1):1–7. doi: 10.5334/JBSR.2342
303. Besutti G, Giorgi Rossi P, Iotti V, Spaggiari L, Bonacini R, Nitrosi A, et al. Accuracy of CT in a cohort of symptomatic patients with suspected COVID-19 pneumonia during the outbreak peak in Italy. *Eur Radiol*. 2020;30(12):6818–27. doi: 10.1007/s00330-020-07050-x
304. Caruso D, Zerunian M, Polici M, Pucciarelli F, Polidori T, Rucci C, et al. Chest CT Features of COVID-19 in Rome, Italy. *Radiology*. 2020;296(2):E79–85. doi: 10.1148/radiol.2020201237
305. de Smet K, de Smet D, Ryckaert T, Laridon E, Heremans B, Vandenbulcke R, et al. Diagnostic performance of chest CT for SARS-CoV-2 infection in individuals with or without COVID-19 symptoms. *Radiology*. 2020;298(1):E30–7. doi: 10.1148/RADIOL.2020202708
306. Debray MP, Tarabay H, Males L, Chalhoub N, Mahdjoub E, Pavlovsky T, et al. Observer agreement and clinical significance of chest CT reporting in patients suspected of COVID-19. *Eur Radiol*. 2021;31(2):1081–9. doi: 10.1007/s00330-020-07126-8
307. Deng Z, Zhange X, Li Y, Xu H, Gang Y, Wang H. Value of chest CT screening in the early COVID-19 outbreak. *Chinese Journal of Radiology* 2020;54:430-4. doi: 10.3760/cma.j.cn112149-20200218-00187

308. Dofferhoff A, Swinkels A, Sprong T, Berk Y, Spanbroek M, Nabuurs-Franssen M. Diagnostic algorithm for COVID-19 at the ER. *Nederlandsche Tijdschrift voor Geneeskunde* 2020;164:D5042.
309. Ducray V, Vlachomitrou AS, Bouscambert-Duchamp M, Si-Mohamed S, Gouttard S, Mansuy A, et al. Chest CT for rapid triage of patients in multiple emergency departments during COVID-19 epidemic: experience report from a large French university hospital. *Eur Radiol*. 2021;31(2):795–803. doi: 10.1007/s00330-020-07154-4
310. Falaschi Z, Danna PSC, Arioli R, Pasché A, Zagaria D, Percivale I, et al. Chest CT accuracy in diagnosing COVID-19 during the peak of the Italian epidemic: A retrospective correlation with RT-PCR testing and analysis of discordant cases. *Eur J Radiol* [Internet]. 2020 Sep;130(January):109192. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0720048X20303818>
311. Fujioka T, Takahashi M, Mori M, Tsuchiya J, Yamaga E, Horii T, et al. Evaluation of the Usefulness of CO-RADS for Chest CT in Patients Suspected of Having COVID-19. *Diagnostics*. 2020;10(9). doi: 10.3390/diagnostics10090608
312. Gezer NS, Ergun B, Barış MM, Appak Ö, Sayiner AA, Balci P, et al. Covid-19 s: A new proposal for diagnosis and structured reporting of covid-19 on computed tomography imaging. *Diagnostic Interv Radiol*. 2020;26(4):315–22. doi: 10.5152/dir.2020.20351
313. Giannitto C, Sposta FM, Repici A, Vatteroni G, Casiraghi E, Casari E, et al. Chest CT in patients with a moderate or high pretest probability of COVID-19 and negative swab. *Radiol Medica* [Internet]. 2020;125(12):1260–70. Available from: <https://doi.org/10.1007/s11547-020-01269-w>
314. Gietema HA, Zelis N, Nobel JM, Lambriksi LJG, Alphen LBV, Lashof AMLO, et al. CT in relation to rt-PCR in diagnosing covid-19 in the netherlands: A prospective study. *PLoS One* [Internet]. 2020;15(7 July):1–10. Available from: <http://dx.doi.org/10.1371/journal.pone.0235844>
315. Grando RD, Brentano VB, Zanardo AP, Hertz FT, Júnior LCA, Prietto dos Santos JF, et al. Clinical usefulness of tomographic standards for COVID-19 pneumonia diagnosis: Experience from a Brazilian reference center. *Brazilian J Infect Dis* [Internet]. 2020 Nov;24(6):524–33. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1413867020301471>
316. Guillo E, Bedmar Gomez I, Dangeard S, Bennani S, Saab I, Tordjman M, et al. COVID-19 pneumonia: Diagnostic and prognostic role of CT based on a retrospective analysis of 214 consecutive patients from Paris, France. *Eur J Radiol* [Internet]. 2020 Oct;131(January):109209. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0720048X20303983>
317. He J, Luo L, Luo Z, Lyu J, Ng M, Shen X, et al. Diagnostic performance between CT and initial real-time RT-PCR for clinically suspected 2019 coronavirus disease (COVID-19) patients outside Wuhan, China. *Respir Med* [Internet]. 2020 Jul;168(January):105980. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0954611120301207>
318. Hermans JJR, Groen J, Zwets E, Boxma-De Klerk BM, Van Werkhoven JM, Ong DSY, et al. Chest CT for triage during COVID-19 on the emergency department: myth or truth? *Emerg Radiol*. 2020;27(6):641–51.
319. Hernigou J, Cornil F, Poignard A, El Bouchaïbi S, Mani J, Naouri JF, et al. Thoracic computerised tomography scans in one hundred eighteen orthopaedic patients during the COVID-19 pandemic: identification of chest lesions; added values; help in managing patients; burden on the computerised tomography scan department. *Int Orthop*. 2020;44(8):1571–80. doi: 10.1007/s00264-020-04651-5
320. Herpe G, Lederlin M, Naudin M, Ohana M, Chaumoitre K, Gregory J, et al. Efficacy of Chest CT for COVID-19 Pneumonia Diagnosis in France. *Radiology* [Internet]. 2021 Feb;298(2):E81–7. Available from: <http://pubs.rsna.org/doi/10.1148/radiol.2020202568>
321. Kalin S. Evaluation of Radiological Findings in Pediatric Patients with COVID-19 in Turkey. *North Clin Istanbul*. 2021;8(4):332–9. doi: 10.14744/nci.2020.97947
322. Kavak S, Duymus R. RSNA and BSTI grading systems of COVID-19 pneumonia: comparison of the diagnostic performance and interobserver agreement. *BMC Med Imaging* [Internet]. 2021;21(1). Available from: <https://doi.org/10.1186/s12880-021-00668-3>
323. Kiziloglu I, Sener A, Siliv N. Comparison of rapid antibody test and thorax computed tomography results in patients who underwent RT-PCR with the pre-diagnosis of COVID-19. *Int J Clin Pract*. 2021;(February):1–7. doi: 10.1111/ijcp.14524
324. Korevaar DA, Kootte RS, Smits LP, van den Aardweg JG, Bonta PI, Schinkel J, et al. Added value of chest computed tomography in suspected COVID-19: An analysis of 239 patients. *Eur Respir J*. 2020;56(2):38–41. doi: 10.1183/13993003.01377-2020
325. Krdzalic J, de Jaegere TMH, Kwee RM. Diagnostic performance of chest CT in screening patients with suspected COVID-19 infection in a Western population. *Br J Radiol* [Internet]. 2020 Sep 1;93(1113):20200643. Available from: <https://www.birpublications.org/doi/10.1259/bjr.20200643>

326. Kuzan TY, Murzoğlu Altıntoprak K, Çiftçi HÖ, Ergül U, Ünal Özdemir NB, Bulut M, et al. A comparison of clinical, laboratory and chest CT findings of laboratory-confirmed and clinically diagnosed COVID-19 patients at first admission. *Diagn Interv Radiol*. 2021;27(3):336–43. doi: 10.5152/dir.2020.20270
327. Li Z, Zeng B, Lei P, Liu J, Fan B, Shen Q, et al. Differentiating pneumonia with and without COVID-19 using chest CT images: from qualitative to quantitative. *J Xray Sci Technol*. 2020;28(4):583–9. doi: 10.3233/xst-200689
328. Luo L, Luo Z, Jia Y, Zhou C, He J, Lyu J, et al. CT differential diagnosis of COVID-19 and non-COVID-19 in symptomatic suspects: A practical scoring method. *BMC Pulm Med*. 2020;20(1):1–9. doi: 10.1186/s12890-020-1170-6
329. Luo N, Zhang H, Zhou Y, Kong ZX, Sun WH, Huang N, et al. Utility of chest CT in diagnosis of COVID-19 pneumonia. *Diagnostic Interv Radiol*. 2020;26(5):437–42. doi: 10.5152/dir.2020.20144
330. Mei X, Lee HC, Diao K yue, Huang M, Lin B, Liu C, et al. Artificial intelligence-enabled rapid diagnosis of patients with COVID-19. *Nat Med* [Internet]. 2020;26(8):1224–8. Available from: <http://dx.doi.org/10.1038/s41591-020-0931-3>
331. Mirahmadizadeh A, Pourmontaseri Z, Afrashteh S, Hosseinzadeh M, Karimi J, Sharafi M. Sensitivity and specificity of chest computed tomography scan based on RT-PCR in COVID-19 diagnosis. *Polish J Radiol*. 2021;86:e74–7. doi: 10.5114/pjr.2021.103858
332. Miranda Magalhães Santos JM, Paula Alves Fonseca A, Pinheiro Zarattini Anastacio E, Formagio Minenelli F, Furtado De Albuquerque Cavalcanti C, Borges Da Silva Teles G. Initial results of the use of a standardized diagnostic criteria for chest computed tomography findings in coronavirus disease 2019. *J Comput Assist Tomogr*. 2020;44(5):647–51. doi: 10.1097/RCT.0000000000001054
333. Nair AV, McInnes M, Jacob B, Kumar D, Soman DK, Subair HSV, et al. Diagnostic accuracy and inter-observer agreement with the CO-RADS lexicon for CT chest reporting in COVID-19. *Emerg Radiol*. 2021;28(6):1045–54. doi: 10.1007/s10140-021-01967-6
334. Ohana M, Muller J, Severac F, Bilbault P, Behr M, Oberlin M, et al. Temporal variations in the diagnostic performance of chest CT for Covid-19 depending on disease prevalence: Experience from North-Eastern France. *Eur J Radiol* [Internet]. 2021 Jan;134(January):109425. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0720048X2030615X>
335. Özer H, Kılınçer A, Uysal E, Yormaz B, Cebeci H, Durmaz MS, et al. Diagnostic performance of Radiological Society of North America structured reporting language for chest computed tomography findings in patients with COVID-19. *Jpn J Radiol* [Internet]. 2021;39(9):877–88. Available from: <https://doi.org/10.1007/s11604-021-01128-2>
336. Özkaraçaklı MA, Özkurt H, Bardakçı Mİ, Akıllı İK, Yanç U, Altuntaş Y, et al. Comparison of chest computed tomography findings of RT-PCR negative and RT-PCR positive cases in COVID-19 patients. *Clin Imaging*. 2022;82(October 2021):7–12. doi: 10.1016/j.clinimag.2021.10.013
337. Patel M, Chowdhury J, Zheng M, Abramian O, Verga S, Zhao H. High resolution CHEST CT (HRCT) evaluation in patients hospitalized with COVID-19 infection. [Preprint] 2020. Available from: 10.1101/2020.05.26.20114082
338. Peng D, Zhang J, Xu Y, Liu Z, Wu P. The role of procalcitonin in early differential diagnosis of suspected children with COVID-19. . [Preprint] 2020. Available from: <https://doi.org/10.1101/2020.04.07.20057315>
339. Prokop M, Van Everdingen W, Van Rees Vellinga T, Van Ufford HQ, Stöger L, Beenen L, et al. CO-RADS: A Categorical CT Assessment Scheme for Patients Suspected of Having COVID-19- Definition and Evaluation. *Radiology*. 2020;296(2):E97–104. doi: 10.1148/radiol.2020201473
340. Ravikanth R. Diagnostic accuracy of chest computed tomography in improving the false negative rate as compared to reverse transcriptase polymerase chain reaction in coronavirus disease 2019 pneumonia: A cross sectional analysis of 348 cases from India. *Lung India* [Internet]. 2021;38(7):11. Available from: <http://www.lungindia.com/text.asp?2021/38/7/11/310900>
341. Revel M, Boussouar S, de Margerie-Mellon C, Saab I, Lapotre T, Mompont D, et al. Study of Thoracic CT in COVID-19: The STOIC Project. *Radiology* [Internet]. 2021 Oct;301(1):E361–70. Available from: <http://pubs.rsna.org/doi/10.1148/radiol.2021210384>
342. Saeed GA, Helali AA Al, Shah A, Almazrouei S, Ahmed LA. Chest CT performance and features of COVID-19 in the region of Abu Dhabi, UAE: a single institute study. *Chinese J Acad Radiol* [Internet]. 2021; Available from: <https://doi.org/10.1007/s42058-021-00075-1>
343. Schulze-Hagen M, Yüksel C, Sähn M, Isfort P, Kuhl C, Hübel C, et al. Low-dose chest CT for the diagnosis of COVID-19 a systematic, prospective comparison with PCR. *Dtsch Arztebl Int*. 2020;117(22–23):389–95. doi: 10.3238/arztebl.2020.0389

344. Shirota G, Sato Y, Itoh D, Gonoi W, Hayashi TY, Sugita Y, et al. Pitfalls in chest CT findings of COVID-19 patients infected during hospitalisation. *Clin Imaging*. 2021;78(January):146–53. doi: 10.1016/j.clinimag.2021.03.017
345. Simsek Yurt N, Ocak M, Yurt YC. Epidemiological and clinical characteristics of 3334 cases with prediagnosis coronavirus disease-2019 (COVID-19) in Turkey. *Int J Clin Pract*. 2021;75(10):1–8. doi: 10.1111/ijcp.14496
346. Song S, Wu F, Liu Y, Jiang H, Xiong F, Guo X, et al. Correlation between Chest CT Findings and Clinical Features of 211 COVID-19 Suspected Patients in Wuhan, China. *Open Forum Infect Dis*. 2020;7(6):1–8. doi: 10.1093/ofid/ofaa171
347. Steuwe A, Rademacher C, Valentin B, Köhler MH, Appel E, Keitel V, et al. Dose-optimised chest computed tomography for diagnosis of Coronavirus Disease 2019 (COVID-19) - Evaluation of image quality and diagnostic impact. *J Radiol Prot*. 2020;40(3):877–91. doi: 10.1088/1361-6498/aba16a
348. Thomas C, Naudin M, Tasu JP, Leclerc C, Depaire L, Suberville M, et al. Efficacy of chest CT scan for COVID-19 diagnosis in a low prevalence and incidence region. *Eur Radiol*. 2021;31(11):8141–6. doi: 10.1007/s00330-021-07863-4
349. Vicini S, Panvini N, Bellini D, Rengo M, Ciotola M, De Vivo M, et al. Radiographers and COVID-19 pneumonia: Diagnostic performance using CO-RADS. *Radiography [Internet]*. 2021 Nov;27(4):1078–84. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1078817421000493>
350. Wang T, Xiong Z, Zhou H, Luo W, Tang H, Liu J. Design, validation, and clinical practice of standardized imaging diagnostic report for COVID-19. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* 2020;45:229-35. Doi: 10.11817/j.issn.1672-7347.2020.200152
351. Xiong Z, Fu L, Zhou H, Liu J, Wang A, Huang Y. Construction and evaluation of a novel diagnosis pathway for 2019-coronavirus disease. *Zhonghua Yi Xue za Zhi* 2020;100:1223-9. Doi: 10.3760/cma.j.cn112137-20200228-00499
352. Lazar Neto, F., Marino, L. O., Torres, A., Cilloniz, C., Meirelles Marchini, J. F., Garcia de Alencar, J. C., Palomeque, A., Albacar, N., Brandão Neto, R. A., Souza, H. P., Ranzani, O. T., Bortolotto, A. L., Müller Veiga, A. D., Bellintani, A. P., Fantinatti, B. L., Nicolao, B. R., Caldeira, B. T., Umehara Juck, C. E., Bueno, C. G., ... Gomez Gomez, L. M. (2021). Community-acquired pneumonia severity assessment tools in patients hospitalized with COVID-19: A validation and clinical applicability study. *Clinical Microbiology and Infection*, 27(7). <https://doi.org/10.1016/j.cmi.2021.03.002>
353. Armiñanzas, C., Arnaiz de las Revillas, F., Gutiérrez Cuadra, M., Arnaiz, A., Fernández Sampedro, M., González-Rico, C., Ferrer, D., Mora, V., Suberviola, B., Latorre, M., Calvo, J., Olmos, J. M., Cifrián, J. M., & Fariñas, M. C. (2021). Usefulness of the covid-gram and curb-65 scores for predicting severity in patients with covid-19. *International Journal of Infectious Diseases*, 108, 282–288. <https://doi.org/10.1016/j.ijid.2021.05.048>
354. Liang, W., Liang, H., Ou, L., Chen, B., Chen, A., Li, C., Li, Y., Guan, W., Sang, L., Lu, J., Xu, Y., Chen, G., Guo, H., Guo, J., Chen, Z., Zhao, Y., Li, S., Zhang, N., Zhong, N., & He, J. (2020). Development and validation of a clinical risk score to predict the occurrence of critical illness in hospitalized patients with covid-19. *JAMA Internal Medicine*, 180(8), 1081. <https://doi.org/10.1001/jamainternmed.2020.2033>
355. Doğanay, F., & Ak, R. (2021). Performance of the curb-65, Isaric-4C and covid-gram scores in terms of severity for Covid-19 patients. *International Journal of Clinical Practice*, 75(10). <https://doi.org/10.1111/ijcp.14759>
356. Apea, V. J., Wan, Y. I., Dhairyawan, R., Puthucheary, Z. A., Pearse, R. M., Orkin, C. M., & Prowle, J. R. (2020). Ethnicity and outcomes in patients hospitalised with COVID-19 infection in East London: An observational cohort study. *Medrxiv*. <https://doi.org/10.1101/2020.06.10.20127621>
357. Cobos-Siles, M., Cubero-Morais, P., Arroyo-Jiménez, I., Rey-Hernández, M., Hernández-Gómez, L. Vargas-Parra, D. J., González-Fernández, M., Cazorla-González, M., Gabella-Martín, M., Ruíz-Albi, T., Berezo-García, J. A., García-Cruces-Méndez, J. F., Miramontes-González, J. P., & Corral-Gudino, L. (2020). Cause-specific death in hospitalized individuals infected with SARS-COV-2: More than just acute respiratory failure or thromboembolic events. *Internal and Emergency Medicine*, 15(8), 1533–1544. <https://doi.org/10.1007/s11739-020-02485-y>
358. Dres, M., Hajage, D., Lebbah, S., Kimmoun, A., Pham, T., Béduneau, G., Combes, A., Mercat, A., Guidet, B., Demoule, A., & Schmidt, M. (2021). Characteristics, management, and prognosis of elderly patients with covid-19 admitted in the ICU during the first wave: Insights from the Covid-ICU Study. *Annals of Intensive Care*, 11(1). <https://doi.org/10.1186/s13613-021-00861-1>

359. Miles, A., Webb, T. E., Mcloughlin, B. C., Mannan, I., Rather, A., Knopp, P., & Davis, D. (2020). Outcomes from covid-19 across the range of frailty: Excess mortality in fitter older people. *European Geriatric Medicine*, 11(5), 851–855. <https://doi.org/10.1007/s41999-020-00354-7>
360. Aw, D., Woodrow, L., Ogliari, G., & Harwood, R. (2020). Association of frailty with mortality in older inpatients with COVID-19: A cohort study. *Age and Ageing*, 49(6), 915–922. <https://doi.org/10.1093/ageing/afaa184>
361. Jung, C., Flaatten, H., Fjølner, J., Bruno, R. R., Wernly, B., Artigas, A., Bollen Pinto, B., Schefold, J. C., Wolff, G., Kelm, M., Beil, M., Sviri, S., van Heerden, P. V., Szczeklik, W., Czuczwar, M., Elhadi, M., Joannidis, M., Oeyen, S., Zafeiridis, T., ... Zegers, M. (2021). The impact of frailty on survival in elderly intensive care patients with covid-19: The COVIP study. *Critical Care*, 25(1). <https://doi.org/10.1186/s13054-021-03551-3>
362. Aliberti, M. J., Szlejf, C., Avelino-Silva, V. I., Suemoto, C. K., Apolinario, D., Dias, M. B., Garcez, F. B., Trindade, C. B., Amaral, J. R., Melo, L. R., Aguiar, R. C., Coelho, P. H., Hojaij, N. H., Saraiva, M. D., Silva, N. O., Jacob-Filho, W., & Avelino-Silva, T. J. (2021). Covid -19 is not over and age is not enough: Using frailty for prognostication in hospitalized patients. *Journal of the American Geriatrics Society*, 69(5), 1116–1127. <https://doi.org/10.1111/jgs.17146>
363. Hewitt, J., Carter, B., Vilches-Moraga, A., Quinn, T. J., Braude, P., Verduri, A., Pearce, L., Stechman, M., Short, R., Price, A., Collins, J. T., Bruce, E., Einarsson, A., Rickard, F., Mitchell, E., Holloway, M., Hesford, J., Barlow-Pay, F., Clini, E., ... Guaraldi, G. (2020). The effect of frailty on survival in patients with covid-19 (COPE): A multicentre, European, observational cohort study. *The Lancet Public Health*, 5(8). [https://doi.org/10.1016/s2468-2667\(20\)30146-8](https://doi.org/10.1016/s2468-2667(20)30146-8)
364. Thompson, J. V., Meghani, N. J., Powell, B. M., Newell, I., Craven, R., Skilton, G., Bagg, L. J., Yaqoob, I., Dixon, M. J., Evans, E. J., Kambele, B., Rehman, A., & Ng Man Kwong, G. (2020). Patient characteristics and predictors of mortality in 470 adults admitted to a District General Hospital in England with covid-19. *Epidemiology and Infection*, 148. <https://doi.org/10.1017/s0950268820002873>
365. Knight SR, Ho A, Pius R, Buchan I, Carson G, Drake TM, et al. Risk stratification of patients admitted to hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: development and validation of the 4C Mortality Score. *BMJ*. 2020;370:m3339.
366. Marcolino M, Pires M, Ramos L, Silva R, Oliveira L, Carvalho R, et al. ABC2-SPH risk score for in-hospital mortality in COVID-19 patients: development, external validation and comparison with other available scores. medRxiv preprint doi: <https://doi.org/10.1101/2021.02.01.21250306>; this version posted February 3, 2021
367. Jimenez-Solem E et al. Developing And Validating Covid-19 Adverse Outcome Risk Prediction Models From A Bi-National European Cohort Of 5594 Patients. doi: <https://doi.org/10.1101/2020.10.06.20207209>
368. Clift AK, Coupland CAC, Keogh RH, Diaz-Ordaz K, Williamson E, Harrison EM et al. Living risk prediction algorithm (QCOVID) for risk of hospital admission and mortality from coronavirus 19 in adults: national derivation and validation cohort study. *BMJ* 2020;371:m3731 <http://dx.doi.org/10.1136/bmj.m3731>
369. Nafilyan V, Humberstone B, Mehta N, Diamond I, Coupland C, Lorenzi L et al. An external validation of the QCovid risk prediction algorithm for risk of mortality from COVID-19 in adults: national validation cohort study in England. medRxiv preprint <https://doi.org/10.1101/2021.01.22.21249968> this version posted January 25, 2021
370. Gupta R, Harrison E, Ho A, Docherty A, Knight S, Smeden M et al. Development and validation of the ISARIC 4C Deterioration model for adults hospitalised with COVID-19: a prospective cohort study. *Lancet Respir Med* 2021. [https://doi.org/10.1016/S2213-2600\(20\)30559-2](https://doi.org/10.1016/S2213-2600(20)30559-2)
371. Adderley N, Taverner T, Price M, Sainsbury C, Greenwood D, Chandan J et al. Development and external validation of prognostic models for COVID-19 to support risk stratification in secondary care. medRxiv preprint doi: <https://doi.org/10.1101/2021.01.25.21249942>; this version posted January 30, 2021.
372. Ageno W, Cogliati C, Perego M, Girelli D, Crisaffuli E, Pizzolo F et al. Clinical risk scores for the early prediction of severe outcomes in patients hospitalized for COVID-19. *Internal and Emergency Medicine* 2020. <https://doi.org/10.1007/s11739-020-02617-4>
373. Artero A, Madrazo M, Fernandez-Garces M, Miguez A, Garcia A, Vieitez A Severity Scores in COVID-19 Pneumonia: a Multicenter, Retrospective, Cohort Study. *J Gen Intern Med* DOI: 10.1007/s11606-021-06626-7.
374. Baker KF, Hanrath A, van der Loeff I, Kay L, Back J, Duncan C. National Early Warning Score 2 (NEWS2) to identify inpatient COVID-19 deterioration: a retrospective analysis. *Clinical Medicine* 2021, Vol 21 No 2 March 2021.

375. Bartoletti M, Giannella M, Scudeller L, Tedeschi S, Rinaldi M, Bussini L et al. Development and validation of a prediction model for severe respiratory failure in hospitalized patients with SARS-CoV-2 infection: a multicentre cohort study (PREDI-CO study). *Clinical Microbiology and Infection*. 2020 <https://doi.org/10.1016/j.cmi.2020.08.003>.
376. Berenguer J, Borobia A, Ryan P, Rodriguez-Bano J, Bellon J, Harrin I et al. Development and validation of a prediction model for 30-day mortality in hospitalised patients with COVID-19: the COVID-19 SEIMC score. *Thorax* 2020. <http://dx.doi.org/10.1136/thoraxjnl-2020-216001>.
377. Bradley P, Frost F, Tharmaratnam K, et al. Utility of established prognostic scores in COVID-19 hospital admissions: multicenter prospective evaluation of CURB-65, NEWS2 and qSOFA. *BMJ Open Res* 2020;7:e000729. doi:10.1136.bmjresp-2020-000729.
378. Carr E, Bendayan R, Bean D, Stammers M, Wang W, Zhang H, Searle T et al. Evaluation and improvement of the NEWS2 for COVID-19: a multi-hospital study. *BMC Medicine* 19:23. <https://doi.org/10.1186/s12916-020-01893-3>.
379. Castro V, McCoy T, Perlis R. Consistency of performance of adverse outcome prediction models for hospitalized COVID-19 patients. medRxiv preprint doi: <https://doi.org/10.1101/2021.02.25.21252410>; this version posted March 1, 2021.
380. Chua F, Vancheeswaran R, Draper A, Vaghela T, Knight M, Mogal R et al. Early prognostication of COVID-19 to guide hospitalisation versus outpatient monitoring using a point-of-test risk prediction score *Thorax* Published Online First: 10 March 2021. doi: 10.1136/thoraxjnl-2020-216425
381. Codon J, Lip G, Rey H, Iniesta A, Rosillo S, Castrejon S et al. Prediction of thromboembolic events and mortality by the CHADS₂ and the CHA₂DS₂-VASc in COVID-19. *Europace* (2021) 00, 1–11 doi:10.1093/europace/euab015.
382. El-Solh AA, Lawson Y, Carter M, El-Solh DA, Mergenhagen KA (2020) Comparison of in-hospital mortality risk prediction models from COVID-19. *PLoS ONE* 15(12): e0244629. <https://doi.org/10.1371/journal.pone.0244629>
383. Gupta RK, Marks M, Samuels THA, et al. Systematic evaluation and external validation of 22 prognostic models among hospitalised adults with COVID-19: An observational cohort study. *Eur Respir J* 2020; in press (<https://doi.org/10.1183/13993003.03498-2020>).
384. King JT, Jr., Yoon JS, Rentsch CT, Tate JP, Park LS, Kidwai-Khan F, et al. (2020) Development and validation of a 30-day mortality index based on pre-existing medical administrative data from 13,323 COVID-19 patients: The Veterans Health Administration COVID-19 (VACO) Index. *PLoS ONE* 15(11): e0241825. <https://doi.org/10.1371/journal.pone.0241825>
385. Li, J., Chen, Y., Chen, S. et al. Derivation and validation of a prognostic model for predicting in-hospital mortality in patients admitted with COVID-19 in Wuhan, China: the PLANS (platelet lymphocyte age neutrophil sex) model. *BMC Infect Dis* 20, 959 (2020). <https://doi.org/10.1186/s12879-020-05688-y>
386. Fang-Yan Liu, Xue-Lian Sun, Yong Zhang, Lin Ge, Jing Wang, Xiao Liang, Jun-Fen Li, Chang-Liang Wang, Zheng-Tao Xing, Jagadish K. Chhetri, Peng Sun, Piu Chan. Evaluation of the Risk Prediction Tools for Patients With Coronavirus Disease 2019 in Wuhan, China: A Single-Centered, Retrospective, Observational Study. *Critical Care Medicine* DOI: 10.1097/CCM.0000000000004549
387. Liu H, Chen J, Yang Q, Wang Y, Cai J, Li H. Development and validation of a risk score using complete blood count to predict in-hospital mortality in COVID-19 patients, *Med* (2020), <https://doi.org/10.1016/j.medj.2020.12.013>
388. Mancilla-Galindo J, Vera- Zertuche JM, Navarro-Cruz AR, Segura-Badilla O, Reyes-Velázquez G, Tepepa-López FJ, Aguilar-Alonso P, Vidal-Mayo JdeJ, Kammar- García A (2020). Development and validation of the patient history COVID-19 (PH-Covid19) scoring system: a multivariable prediction model of death in Mexican patients with COVID-19. *Epidemiology and Infection* 148, e286, 1–8. <https://doi.org/10.1017/S095026882000290>
389. Guillermo Rodriguez-Nava, Maria Adriana Yanez-Belloa, Daniela Patricia Trelles-Garciaa, Chul Won Chunga, Harvey J. Friedmanb,c, David W. Hinesd. Performance of the quick COVID-19 severity index and the Brescia-COVID respiratory severity scale in hospitalized patients with COVID-19 in a community hospital setting. *International Journal of Infectious Diseases* 102 (2021) 571–576
390. Nicholson C, Wooster L, Sigurslide H, Li R, Jiang W, Tian W, Cardenas C, Malhotra R. Estimating risk of mechanical ventilation and in-hospital mortality among adult COVID-19 patients admitted to Mass General Brigham: The VICE and DICE scores. *EClinicalMedicine* 33 (2021) 100765
391. Paranjape N, Staples L, Stradwick C, Ray H, Saldanha I. Development and validation of a predictive model for critical illness in adult patients requiring hospitalization for COVID-19

- medRxiv preprint doi: <https://doi.org/10.1101/2021.01.22.21250289>; this version posted January 22, 2021.
392. Richardson D, Faisal M, Fiori M, *et al.* Use of the first National Early Warning Score recorded within 24 hours of admission to estimate the risk of in-hospital mortality in unplanned COVID-19 patients: a retrospective cohort study. *BMJ Open* 2021;11:e043721. doi:10.1136/bmjopen-2020-043721
 393. Schöning V, Liakoni E, Baumgartner C, Exadaktylos A, Hautz W, Atkinson A, Hammann F Development and validation of a prognostic COVID-19 severity assessment (COSA) score and machine learning models for patient triage at a tertiary hospital. *J Transl Med* (2021) 19:56 <https://doi.org/10.1186/s12967-021-02720-w>
 394. Tanboga IH, Canpolat U, Çetin E, Kundi H, Çelik O, Çağlayan M, Ata N, Özeke O, Çay S, Kaymaz C, Topaloğlu S. Development and validation of clinical prediction model to estimate the probability of death in hospitalized patients with COVID-19: Insights from a nationwide database. *J Medical Virology* 2021;1–8 DOI: 10.1002/jmv.26844
 395. van Dam P, Zelis N, van Kuijk S, Linkens A, Bruggemann R, Spaetgens B *et al.* Performance of prediction models for short term outcome in COVID-19 patients in the emergency department: a retrospective study. *Ann Med* 2021 Dec;53(1):402-409. doi: 10.1080/07853890.2021.1891453
 396. van Dam PMEL, Zelis N, Stassen P, *et al.* Validating the RISE UP score for predicting prognosis in patients with COVID-19 in the emergency department: a retrospective study. *BMJ Open* 2021;11:e045141. doi:10.1136/bmjopen-2020-045141
 397. van Klaveren D, Rekkas A, Alisma J, Verdonschot R, Koning D, Kamps M *et al.* COVID Outcome Prediction in the Emergency Department (COPE): Development and validation of a model for predicting death and need for intensive care in COVID-19 patients. medRxiv preprint doi: <https://doi.org/10.1101/2020.12.30.20249023>; this version posted February 8, 2021
 398. Zhang L, Hou J, Ma FZ, Li J, Xue S, Xu ZG. The common risk factors for progression and mortality in COVID-19 patients: a meta-analysis. *Arch Virol.* 2021;166(8):2071-2087. doi:10.1007/s00705-021-05012-2
 399. Malik P, Patel U, Mehta D, *et al.* Biomarkers and outcomes of COVID-19 hospitalisations: systematic review and meta-analysis. *BMJ Evid Based Med.* 2021;26(3):107-108. doi:10.1136/bmjebm-2020-111536
 400. Mahat RK, Panda S, Rathore V, Swain S, Yadav L, Sah SP. The dynamics of inflammatory markers in coronavirus disease-2019 (COVID-19) patients: A systematic review and meta-analysis. *Clin Epidemiol Glob Health.* 2021;11:100727. doi:10.1016/j.cegh.2021.100727
 401. Melo AKG, Milby KM, Caparroz ALMA *et al.* Biomarkers of cytokine storm as red flags for severe and fatal COVID-19 cases: A living systematic review and meta-analysis. *PLoS One.* 2021;16(6):e0253894. doi:10.1371/journal.pone.0253894
 402. Wang L, Yang LM, Pei SF, *et al.* CRP, SAA, LDH, and DD predict poor prognosis of coronavirus disease (COVID-19): a meta-analysis from 7739 patients. *Scand J Clin Lab Invest.* 2021;1-8. doi:10.1080/00365513.2021.2000635
 403. Shi Q, Wang Z, Liu J, *et al.* Risk factors for poor prognosis in children and adolescents with COVID-19: A systematic review and meta-analysis. *EClinicalMedicine.* 2021;41:101155. doi:10.1016/j.eclim.2021.101155
 404. Benito N, Filella D, Mateo J, Fortuna AM, Gutierrez-Alliende JE, Hernandez N, Gimenez AM, Pomar V, Castellvi I, Corominas H, Casademont J and Domingo P (2020) Pulmonary Thrombosis or Embolism in a Large Cohort of Hospitalized Patients With Covid-19. *Front. Med.* 7:557. doi: 10.3389/fmed.2020.005574.
 405. Zhang L, Yan X, Fan Q, Liu H, Liu X, Liu Z, Zhang Z. D-dimer levels on admission to predict in-hospital mortality in patients with Covid-19. *J Thromb Haemost.* 2020 Jun;18(6):1324-1329. doi: 10.1111/jth.14859.
 406. Duan J, Wang X, Chi J, Chen H, Bai L, Hu Q, Han X, Hu W, Zhu L, Wang X, Li Y, Zhou C, Mou H, Yan X, Guo S. Correlation between the variables collected at admission and progression to severe cases during hospitalization among patients with COVID-19 in Chongqing. *J Med Virol.* 2020 Nov;92(11):2616-2622. doi: 10.1002/jmv.26082. Epub 2020 Jun 9.
 407. Zhang W, Sang L, Shi J, Zhong M, Jiang L, Song B, Kang L, Zhang Y, Zhang D, Yu Y, Zheng X. Association of D-dimer elevation with inflammation and organ dysfunction in ICU patients with COVID-19 in Wuhan, China: a retrospective observational study. *Aging (Albany NY).* 2021 Feb 11;13(4):4794-4810. doi: 10.18632/aging.202496. Epub 2021 Feb 11.
 408. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, Xiang J, Wang Y, Song B, Gu X, Guan L, Wei Y, Li H, Wu X, Xu J, Tu S, Zhang Y, Chen H, Cao B. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet.* 2020 Mar

- 28;395(10229):1054-1062. doi: 10.1016/S0140-6736(20)30566-3. Epub 2020 Mar 11. Erratum in: *Lancet*. 2020 Mar 28;395(10229):1038. Erratum in: *Lancet*. 2020 Mar 28;395(10229):1038.
409. Al-Samkari H, Karp Leaf RS, Dzik WH, Carlson JCT, Fogerty AE, Waheed A, Goodarzi K, Bendapudi PK, Bornikova L, Gupta S, Leaf DE, Kuter DJ, Rosovsky RP. COVID-19 and coagulation: bleeding and thrombotic manifestations of SARS-CoV-2 infection. *Blood*. 2020 Jul 23;136(4):489-500. doi: 10.1182/blood.2020006520.
 410. Peiró ÓM, Carrasquer A, Sánchez-Gimenez R, Lal-Trehan N, Del-Moral-Ronda V, Bonet G, Fort-Gallifa I, Picó-Plana E, Bastón-Paz N, Gutiérrez C, Bardaji A. Biomarkers and short-term prognosis in COVID-19. *Biomarkers*. 2021 Mar;26(2):119-126. doi: 10.1080/1354750X.2021.1874052. Epub 2021 Jan 18.
 411. Yang C, Liu F, Liu W, Cao G, Liu J, Huang S, Zhu M, Tu C, Wang J, Xiong B. Myocardial injury and risk factors for mortality in patients with COVID-19 pneumonia. *Int J Cardiol*. 2021 Mar 1;326:230-236. doi: 10.1016/j.ijcard.2020.09.048. Epub 2020 Sep 23.
 412. Gil-Rodrigo A, Miró Ó, Piñera P, Burillo-Putze G, Jiménez S, Martín A, Martín-Sánchez FJ, Jacob J, Guardiola JM, García-Lamberechts EJ, Espinosa B, Martín Mojarro E, González Tejera M, Serrano L, Agüera C, Soy E, Llauger L, Juan MÁ, Palau A, Del Arco C, Rodríguez Miranda B, Maza Vera MT, Martín Quirós A, Tejada de Los Santos L, Ruiz de Lobera N, Iglesias Vela M, Torres Garate R, Alquézar-Arbé A, González Del Castillo J, Llorens P; en representación de la red de investigación SIESTA. Analysis of clinical characteristics and outcomes in patients with COVID-19 based on a series of 1000 patients treated in Spanish emergency departments. *Emergencias*. 2020 Ago;32(4):233-241.
 413. Yao Y, Cao J, Wang Q, Shi Q, Liu K, Luo Z, Chen X, Chen S, Yu K, Huang Z, Hu B. D-dimer as a biomarker for disease severity and mortality in COVID-19 patients: a case control study. *J Intensive Care*. 2020 Jul 10;8:49. doi: 10.1186/s40560-020-00466-z.
 414. Cen Y, Chen X, Shen Y, Zhang XH, Lei Y, Xu C, Jiang WR, Xu HT, Chen Y, Zhu J, Zhang LL, Liu YH. Risk factors for disease progression in patients with mild to moderate coronavirus disease 2019-a multi-centre observational study. *Clin Microbiol Infect*. 2020 Sep;26(9):1242-1247. doi: 10.1016/j.cmi.2020.05.041. Epub 2020 Jun 9.
 415. Yue T, Zhou W, He J, Wang H, Liu Y, Wang B, Zhu Q, Xia H, Hu H. Combined clinical and imaging features better predict the critical outcomes of patients with SARS-COV-2. *Medicine (Baltimore)*. 2021 Mar 26;100(12):e25083. doi: 10.1097/MD.00000000000025083.
 416. García-Ortega A, Oscullo G, Calvillo P, López-Reyes R, Méndez R, Gómez-Olivas JD, Bekki A, Fonfría C, Trilles-Olaso L, Zaldívar E, Ferrando A, Anguera G, Briones-Gómez A, Reig-Mezquida JP, Feced L, González-Jiménez P, Reyes S, Muñoz-Núñez CF, Carreres A, Gil R, Morata C, Toledo-Pons N, Martí-Bonmati L, Menéndez R, Martínez-García MÁ. Incidence, risk factors, and thrombotic load of pulmonary embolism in patients hospitalized for COVID-19 infection. *J Infect*. 2021 Feb;82(2):261-269. doi: 10.1016/j.jinf.2021.01.003. Epub 2021 Jan 10.
 417. Mestre-Gómez B, Lorente-Ramos RM, Rogado J, Franco-Moreno A, Obispo B, Salazar-Chiriboga D, Saez-Vaquero T, Torres-Macho J, Abad-Motos A, Cortina-Camarero C, Such-Díaz A, Ruiz-Velasco E, Churruca-Sarasqueta J, Muñoz-Rivas N; Infanta Leonor Thrombosis Research Group. Incidence of pulmonary embolism in non-critically ill COVID-19 patients. Predicting factors for a challenging diagnosis. *J Thromb Thrombolysis*. 2021 Jan;51(1):40-46. doi: 10.1007/s11239-020-02190-9.
 418. Mouhat B, Besutti M, Bouiller K, Grillet F, Monnin C, Ecartot F, Behr J, Capellier G, Soumagne T, Pili-Floury S, Besch G, Mourey G, Lepiller Q, Chirouze C, Schiele F, Chopard R, Meneveau N. Elevated D-dimers and lack of anticoagulation predict PE in severe COVID-19 patients. *Eur Respir J*. 2020 Oct 22;56(4):2001811. doi: 10.1183/13993003.01811-2020.
 419. Cho ES, McClelland PH, Cheng O, Kim Y, Hu J, Zenilman M, D'Ayala M. Utility of D-dimer for diagnosis of deep vein thrombosis in coronavirus disease-19 infection. *Journal of Vascular Surgery* January 2021. <https://doi.org/10.1016/j.jvsv.2020.07.009>
 420. Demelo-Rodríguez P, Cervilla-Muñoz E, Ordieres-Ortega L, Parra-Virto A, Toledano-Macías M, Toledo-Samaniego N, García-García A et al. Incidence of asymptomatic deep vein thrombosis in patients with COVID-19 pneumonia and elevated D-dimer levels. *Thrombosis Research* 2020: 192, 23-26. <https://doi.org/10.1016/j.thromres.2020.05.018>
 421. Dujardin RWG, Hilderink BN, Haksteen WE, Middeldorp S, Vlaar APJ, Thachil J et al. Biomarkers for the prediction of venous thromboembolism in critically ill COVID-19 patients. *Thrombosis Research* 2020. 196: 308-312. <https://doi.org/10.1016/j.thromres.2020.09.01>
 422. Kampouri E, Filippidis P, Viala B, Méan M, Pantet O, Desgranges F, Tschopp J, Regina J, Karachalias E, Bianchi C, Zermatten MG, Jatton K, Qanadli SD, Bart PA, Pagani JL, Guery B, Alberio L, Papadimitriou-Olivgeris M, RegCOVID Research Group. Predicting Venous

- Thromboembolic Events in Patients with Coronavirus Disease 2019 Requiring Hospitalization: an Observational Retrospective Study by the COVIDIC Initiative in a Swiss University Hospital. *Biomed Res Int*. 2020 Nov 6;2020:9126148. doi: 10.1155/2020/9126148.
423. Creel-Bulos C, Liu M, Auld SC, Gaddh M, Kempton CL, Sharifpour M, Snieciniski RM, Maier CL, Nahab FB, Rangaraju S. Trends and diagnostic value of D-dimer levels in patients hospitalized with coronavirus disease 2019. *Medicine* 2020;99:46(e23186). <http://dx.doi.org/10.1097/MD.00000000000023186>
 424. Nauka PC, Baronb SW, Assab A, Mohrmannb L, Jindala S, Oranb E et al. Utility of D-dimer in predicting venous thromboembolism in non-mechanically ventilated COVID-19 survivors *Thrombosis Research* 199 (2021) 82–84. <https://doi.org/10.1016/j.thromres.2020.12.023>
 425. Zhang L, Feng X, Zhang D, Jiang C, Mei H, Wang J, Zhang C, Li H, Xia X, Kong S, Liao J, Jia H, Pang X, Song Y, Tian Y, Wang B, Wu C, Yuan H, Zhang Y, Li Y, Sun W, Zhang Y, Zhu S, Wang S, Xie Y, Ge S, Zhang L, Hu Y, Xie M. Deep Vein Thrombosis in Hospitalized Patients With COVID-19 in Wuhan, China: Prevalence, Risk Factors, and Outcome. *Circulation*. 2020 Jul 14;142(2):114-128. doi: 10.1161/CIRCULATIONAHA.120.046702. Epub 2020 May 18. Erratum in: *Circulation*. 2020 Jul 14;142(2):e33.
 426. Alabyad D, Rangaraju S, Liu M, Imran R, Kempton CL, Sharifpour M, et al. (2021) Validation of an admission coagulation panel for risk stratification of COVID-19 patients. *PLoS ONE* 16(3): e0248230. <https://doi.org/10.1371/journal.pone.0248230>
 427. Rali P, O'Corragain O, Oresanya L, Yu D, Sheriff O, Weiss R, Myers C, Desai P, Ali N, Stack A, Bromberg M, Lubitz AL, Panaro J, Bashir R, Lakhter V, Caricchio R, Gupta R, Dass C, Maruti K, Lu X, Rao AK, Cohen G, Criner GJ, Choi ET; Temple University COVID-19 Research Group. Incidence of venous thromboembolism in coronavirus disease 2019: An experience from a single large academic center. *J Vasc Surg Venous Lymphat Disord*. 2021 May;9(3):585-591.e2. doi: 10.1016/j.jvsv.2020.09.006. Epub 2020 Oct 5.
 428. Wang T, Tang C, Chen R, et al. Clinical Features of Coronavirus Disease 2019 Patients With Mechanical Ventilation: A Nationwide Study in China. *Crit Care Med*. 2020;48(9):e809-e812. doi:10.1097/CCM.0000000000004473
 429. Chen, S., Zhu, Q., Xiao, Y., Wu, C., Jiang, Z., Liu, L., & Qu, J. (2021). Clinical and etiological analysis of co-infections and secondary infections in COVID-19 patients: An observational study. *The clinical respiratory journal*, 15(7), 815–825. <https://doi.org/10.1111/crj.13369>
 430. Hughes, S., Mughal, N., & Moore, L. (2021). Procalcitonin to Guide Antibacterial Prescribing in Patients Hospitalised with COVID-19. *Antibiotics (Basel, Switzerland)*, 10(9), 1119. <https://doi.org/10.3390/antibiotics10091119>
 431. Cheng, K., He, M., Shu, Q., Wu, M., Chen, C., & Xue, Y. (2020). Analysis of the Risk Factors for Nosocomial Bacterial Infection in Patients with COVID-19 in a Tertiary Hospital. *Risk management and healthcare policy*, 13, 2593–2599. <https://doi.org/10.2147/RMHP.S277963>
 432. Calderon, M., Li, A., Bazo-Alvarez, J. C., Dennis, J., Baker, K. F., Schim van der Loeff, I., Hanrath, A. T., Capstick, R., Payne, B., Weiland, D., Hunter, E., & Schwab, U. (2021). Evaluation of procalcitonin-guided antimicrobial stewardship in patients admitted to hospital with COVID-19 pneumonia. *JAC-antimicrobial resistance*, 3(3), dlab133. <https://doi.org/10.1093/jacamr/dlab133>
 433. Martimbianco ALC, Pacheco RL, Bagattini AM, Rlera R. Frequency, signs and symptoms, and criteria adopted for long COVID-19: A systematic review. *Int J Clin Pract*. 2021; 75:e14357. doi:10.1111/jcp.14357
 434. Michelen M,, Manoharan L, Elkheir N, et al. Characterising long COVID: a living systematic review. *BMJ Global Health* 2021;6:e005427. doi:10.1136/bmjgh-2021-005427
 435. Liu M, Lv F, Zheng Y, Xiao K. A prospective cohort study on radiological and physiological outcomes of recovered COVID-19 patients 6 months after discharge. *Quant Imaging Med Surg* 2021;11(9):4181-4192. doi: 10.21037/qims-20-1294
 436. Pilotto, ., Cristillo, V., Cotti Piccinelli, S. et al. Long-term neurological manifestations of COVID-19: prevalence and predictive factors. *Neurol Sci* (2021). doi: 10.1007/s10072-021-05586-4
 437. Wallis T, Heiden E, Welham B, et al. Risk factors for persistent abnormality on chest radiographs at 12-weeks post hospitalization with PCR confirmed COVID-19. *Respir Res* 2021; 22:157. doi:10.1186/s12931-021-01750-8
 438. Platten S, Bartlett A, MacCallum P, Makris M, McDonald V, Singh D, Scully M, Pavord S. Evaluation of laboratory assays for anti-platelet factor 4 antibodies after ChAdOx1 nCoV-19 vaccination. *J Thromb Haemost*. 2021 Aug;19(8):2007-2013. doi: 10.1111/jth.15362. Epub 2021 Jul 5. PMID: 33973336; PMCID: PMC8236994.
 439. Sellaturay P, Nasser S, Islam S, et al. Polyethylene glycol (PEG) is a cause of anaphylaxis to the Pfizer/BioNTech mRNA COVID-19 vaccine. *Clinical & Experimental Allergy*; 51: 861–863.

440. Restivo, Vincenzo, et al. "Allergy to Polyethylenglicole of Anti-SARS CoV2 Vaccine Recipient: A Case Report of Young Adult Recipient and the Management of Future Exposure to SARS-CoV2." *Vaccines*, vol. 9, no. 5, 2021, p. 412., <https://doi.org/10.3390/vaccines9050412>.
441. Jiang SY, Smith EM, Vo V, Akdis C, Nadeau KC. Non-immunoglobulin E-mediated allergy associated with Pfizer-BioNTech coronavirus disease 2019 vaccine excipient polyethylene glycol. *Annals of Allergy, Asthma & Immunology* 2021. doi:10.1016/j.anai.2021.09.012.
442. Park HJ, Montgomery JR, Boggs NA. Anaphylaxis After the Covid-19 Vaccine in a Patient With Cholinergic Urticaria. *Military Medicine* 2021. doi:10.1093/milmed/usab138.
443. Pitlick MM, Park MA, Gonzalez-Estrada A, et al. Biphasic anaphylaxis after first dose of messenger RNA coronavirus disease 2019 vaccine with positive polysorbate 80 skin testing result. *Annals of Allergy, Asthma & Immunology*; 127: 498–499.
444. Lim, Xin Rong, et al. "Pseudo-Anaphylactic Reactions to Pfizer BNT162b2 Vaccine: Report of 3 Cases of Anaphylaxis Post Pfizer BNT162b2 Vaccination." *Vaccines*, vol. 9, no. 9, 2021, p. 974., <https://doi.org/10.3390/vaccines9090974>.
445. Warren CM, Snow TT, Lee AS, et al. Assessment of Allergic and Anaphylactic Reactions to mRNA COVID-19 Vaccines With Confirmatory Testing in a US Regional Health System. *JAMA Network Open* 2021; 4. Epub ahead of print 2021. DOI: 10.1001/jamanetworkopen.2021.25524.
446. Laisuan, Wannada, et al. "CoronaVac COVID-19 Vaccine-Induced Anaphylaxis: Clinical Characteristics and Revaccination Outcomes." *Journal of Asthma and Allergy*, Volume 14, 2021, pp. 1209–1215., <https://doi.org/10.2147/jaa.s333098>.
447. Rasmussen TH, Mortz CG, Georgsen TK, Rasmussen HM, Kjaer HF, Bindselev-Jensen C. Patients with suspected allergic reactions to COVID-19 vaccines can be safely revaccinated after diagnostic work-up. *Clinical and Translational Allergy* 2021;11. doi:10.1002/ct2.12044.
448. Kohli-Pamnani A, Zapata K, Gibson T, et al. Coronavirus disease 2019 vaccine hypersensitivity evaluated with vaccine and excipient allergy skin testing. *Annals of Allergy, Asthma & Immunology* 2021. Epub ahead of print 2021. DOI: 10.1016/j.anai.2021.08.417.
449. Blumenthal KG, Robinson LB, Camargo CA, et al. Acute Allergic Reactions to mRNA COVID-19 Vaccines. *Jama*; 325: 1562.

Evidence and Recommendations for the Treatment of COVID-19

1. Hernandez-Cardenas C, Thirion-Romero I, Rivera-Martinez N, Meza-Meneses P, Remigio-Luna A, Perez-Padilla R. Hydrochloroquine for the treatment of severe respiratory infection by COVID-19: A randomized controlled trial. *MedRxiv*. 2021;
2. Johnston C, Brown E, Stewart J, Karita HC, Patricia K, John D, et al. Hydroxychloroquine with or without azithromycin for treatment of early SARS-CoV-2 infection among high-risk outpatient adults: A randomized clinical trial. *EClinicalMedicine* [Internet]. 2021; Available from: [https://www.thelancet.com/journals/eclinm/article/PIIS2589-5370\(21\)00053-5/fulltext](https://www.thelancet.com/journals/eclinm/article/PIIS2589-5370(21)00053-5/fulltext)
3. Beltran Gonzales JL, Gamez MG, Enciso EAM, Maldonado RJE, Palacios DH. Efficacy and safety of Ivermectin and Hydroxychloroquine in patients with severe COVID-19. A randomized controlled trial [Internet]. *MedRxiv*. 2021 [cited 2021 Mar 9]. Available from: <https://www.medrxiv.org/content/10.1101/2021.02.18.21252037v1>
4. Abd-El Salam S, Esmail ES, Khalaf M, Abdo EF, Medhat MA, El Ghafar MSA, et al. Hydroxychloroquine in the treatment of COVID-19: A multicenter randomized controlled study. *Am J Trop Med Hyg*. 2020;103(4):1635–9.
5. Ader F, Peiffer-Smadja N, Poissy J, Bouscambert-Duchamp M, Belhadi D, Diallo A, et al. Antiviral drugs in hospitalized patients with COVID-19 - the DisCoVeRy trial. *MedRxiv* [Internet]. 2021; Available from: <https://www.medrxiv.org/content/10.1101/2021.01.08.20248149v1>
6. Amaravadi R. The PATCH Trial (Prevention And Treatment of COVID-19 With Hydroxychloroquine) (PATCH) [Internet]. [cited 2020 Dec 28]. Available from: <https://clinicaltrials.gov/ct2/show/results/NCT04329923>
7. Cavalcanti A, Zampieri F, Rosa R, Azvedo L, Veiga V, Avezum A, et al. Hydroxychloroquine with or without Azithromycin in Mild to Moderate Covid 19. *NEJM*. 2020;383:2041–52.
8. Chen C-P, Lin Y-C, Chen T-C, Tseng T-Y, Wong H-L, Kuo C-Y, et al. A multicenter, randomized, open-label, controlled trial to evaluate the efficacy and tolerability of hydroxychloroquine and a retrospective study in adult patients with mild to moderate coronavirus disease 2019 (COVID-19). *PLoS One* [Internet]. 2020;15(12):e0242763. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/33264337>
9. Chen J, Liu D, Liu L, Liu P, Xu Q, Xia L, et al. A pilot study of hydroxychloroquine in treatment of patients with moderate COVID-19. *Zhejiang Da Xue Xue Bao Yi Xue Ban* [Internet].

- 2020;49(2):215–9. Available from: <http://www.zjujournals.com/med/EN/10.3785/j.issn.1008-9292.2020.03.03>
10. Chen L, Zhang Z, Fu J, Feng Z, Zhang S, Han Q, et al. Efficacy and safety of chloroquine or hydroxychloroquine in moderate type of COVID 19: a prospective open-label randomized controlled study. *MedRxiv* [Internet]. Available from: <https://doi.org/10.1101/2020.06.19.20136093>
11. Chen Z, Hu J, Zhang Z, Jiang S, Han S, Yan D, et al. Efficacy of hydroxychloroquine in patients with COVID-19: results of a randomized clinical trial. 2020.
12. Dubee V, Roy P, Vielle B, Parot-Schinkel E, Blanchet O, Darsonval A, et al. A placebo-controlled double-blind trial of hydroxychloroquine in mild-to moderate COVID-19. *Medrxiv*.
13. Horby P, Mafham M, Staplin N, Emberson J, Wiselka M, Utiasnowski A. Effect of Hydroxychloroquine in Hospitalized Patients with Covid-19. *N Engl J Med*. 2020;383(21):2030–40.
14. Kamran S, Mirza Z, Azam R, Ullah N, Saeed F, Alamgir W. Clearing the fog: Is hydroxychloroquine effective in reducing Corona virus disease-2019 progression : A randomized controlled trial. *MedRxiv* [Internet]. Available from: <https://doi.org/10.1101/2020.07.30.20165365>
15. Lyngbakken M, Berdal J, Eskesen A, Kvale D, Olsen I, Rueegg C. A pragmatic randomized controlled trial reports lack of efficacy of hydroxychloroquine on coronavirus disease 2019 viral kinetics. *Nat Commun* [Internet]. 2020;11:5284. Available from: <https://doi.org/10.1038/s41467-020-19056-6>
16. Mitja O, Carbacho-Monne M, Ubals M, Tebe C, Penafiel J, Tobias A. Hydrochloroquine for early treatment of adults with mild coronavirus disease 2019: A randomized controlled trial. *Clin Infect Dis*.:ciaa1009.
17. Omrani A, Pathan S, Thomas S, Harris T, Coyle P, Thomas C, et al. Randomized double-blinded placebo-controlled trial of hydroxychloroquine with or without azithromycin for virologic cure of non-severe Covid-19. *EClinical Med*. 2020;29:100645.
18. Pan H, Peto R, Henao-Restrepo A, Preziosi M, Sathiyamoorthy V, Karim Q. Repurposed antiviral drugs for Covid-19 – Interim WHO Solidarity trial results. *NEJM* [Internet]. 2021;384(6):497–511. Available from: <https://www.nejm.org/doi/pdf/10.1056/NEJMoa2023184?articleTools=true>
19. Self W, Semler M, Leither L, Casey J, Angus D, Brower R, et al. Effect of hydroxychloroquine on clinical status at 14 days in hospitalized patients with COVID-19. A randomized clinical trial. *JAMA*. 2020;324(21):2165–76.
20. Skipper CP, Pastick KA, Engen NW, Bangdiwala AS, Abassi M, Lofgren SM, et al. Hydroxychloroquine in Nonhospitalized Adults With Early COVID-19: A Randomized Trial. *Ann Intern Med*. 2020;173(8):623–31.
21. Tang W, Cao Z, Han M, Wang Z, Chen J, Sun W, et al. Hydrochloroquine in patients with mainly mild to moderate coronavirus disease 2019: open label, randomized controlled trial. *BMJ* [Internet]. 2020;369. Available from: <http://dx.doi.org/10.1136/bmj.m1849>
22. Ulrich RJ, Troxel AB, Carmody E, Eapen J, Bäcker M, DeHovitz JA, et al. Treating COVID-19 with hydroxychloroquine (TEACH): A multicenter, double-blind randomized controlled trial in hospitalized patients. *Open Forum Infect Dis*. 2020;7(10):ofaa446.
23. Sekhavati E, Jafari F, SeyedAlinaghi S, Jamalimoghadamsiahkali S, Sadr S, Tabarestani M et al. Safety and effectiveness of azithromycin in patients with COVID-19: An open-label randomised trial. *International Journal of Antimicrobial Agents*. 2020;56(4):106143.
24. Furtado R, Berwanger O, Fonseca H, Corrêa T, Ferraz L, Lapa M et al. Azithromycin in addition to standard of care versus standard of care alone in the treatment of patients admitted to the hospital with severe COVID-19 in Brazil (COALITION II): a randomised clinical trial. *The Lancet*. 2020;396(10256):959-967
25. RECOVERY Collaborative Group. Azithromycin in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *Lancet*. 2021;397(10274):605-612.
26. Kamel AM, Monem MSA, Sharaf NA, Magdy N, Farid SF. Efficacy and safety of azithromycin in Covid-19 patients: A systematic review and meta-analysis of randomized clinical trials. *Rev Med Virol* [Internet]. 2021 [cited 2021 Nov 5];e2258. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1002/rmv.2258>
27. Sharma S, Singh A, Banerjee T. Antibacterial agents used in COVID-19: A systematic review and meta-analysis. *Environmental Sustainability*. 2021; 4:503-513. Available from: <https://doi.org/10.1007/s42398-021-00194-6>

28. Popp M, Stegemann M, Riemer M, Metzendorf M-I, Romero CS, Mikolajewska A, et al. Antibiotics for the treatment of COVID-19. *Cochrane Database of Systematic Reviews* 2021; 10:CD015025. doi: 10.1002/14651858.CD015025.
29. Oldenburg CE, Pinsky BA, Brogdon J, et al. Effect of Oral Azithromycin vs Placebo on COVID-19 Symptoms in Outpatients With SARS-CoV-2 Infection: A Randomized Clinical Trial. *JAMA*. 2021;326(6):490–498. doi:10.1001/jama.2021.11517
30. Echeverría-Esnal D, Martín-Ontiyuelo C, Navarrete-Rouco ME, Cuscó MD, Ferrández O, Horcajada JP, Grau S. Azithromycin in the treatment of COVID-19: a review. *Expert Review of Anti-infective Therapy*. 2021;19(2):147-163. doi: 10.1080/14787210.2020.1813024
31. Lou Y, Liu L, Yao H, et al. Clinical Outcomes and Plasma Concentrations of Baloxavir Marboxil and Favipiravir in COVID-19 Patients: An Exploratory Randomized, Controlled Trial. *Eur J Pharm Sci*. 2021;157:105631. doi:10.1016/j.ejps.2020.105631
32. Ivashchenko AA, Dmitriev KA, Vostokova NV, et al. AVIFAVIR for Treatment of Patients With Moderate Coronavirus Disease 2019 (COVID-19): Interim Results of a Phase II/III Multicenter Randomized Clinical Trial. *Clin Infect Dis*. 2021;73(3):531-534. doi:10.1093/cid/ciaa1176
33. Dabbous HM, El-Sayed MH, El Assal G, et al. Safety and efficacy of favipiravir versus hydroxychloroquine in management of COVID-19: A randomised controlled trial. *Sci Rep*. 2021;11(1):7282. Published 2021 Mar 31. doi:10.1038/s41598-021-85227-0
34. Balykova L.A., Granovskaya M.V., Zaslavskaya K.Ya., Simakina E.N., Agafina A.S., Ivanova A.Yu., Kolontarev K.B., Pushkar D.Yu. New possibilities for targeted antiviral therapy for COVID-19. Results of a multicenter clinical study of the efficacy and safety of using the drug Areplivir. *Infektsionnye bolezni: novosti, mneniya, obuchenie* [Infectious Diseases: News, Opinions, Training]. 2020; 9 (3): 16–29. DOI: <https://doi.org/10.33029/2305-3496-2020-9-3-16-29> (in Russian)
35. Ruzhentsova TA, Chukhlaev PV, Khavkina DA, Garbuzov AA, Osheshnyuk RA, Soluyanov TN, et. al. Phase 3 Trial of Coronavir (Favipiravir) in patients with mild to moderate COVID-19, <https://ssrn.com/abstract=3696907>
36. Udawadia ZR, Singh P, Barkate H, Patil S, Rangwala S, Pendse A, et. al. International Journal of Infections Diseases. 2021; 103:62-71 <https://doi.org/10.1016/j.ijid.2020.11.142>
37. Zhao H, Zhang C, Zhu Q, et al. Favipiravir in the treatment of patients with SARS-CoV-2 RNA recurrent positive after discharge: A multicenter, open-label, randomized trial. *Int Immunopharmacol*. 2021;97:107702. doi:10.1016/j.intimp.2021.107702
38. W. H. O. Solidarity Trial Consortium, Pan H, Peto R, et al. Repurposed Antiviral Drugs for Covid-19 - Interim WHO Solidarity Trial Results. *N Engl J Med* 2020.
39. Spinner CD, Gottlieb RL, Criner GJ, et al. Effect of Remdesivir vs Standard Care on Clinical Status at 11 Days in Patients With Moderate COVID-19: A Randomized Clinical Trial. *JAMA* 2020; 324(11): 1048-57.
40. Wang Y, Zhang D, Du G, et al. Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial. *The Lancet* 2020; 395(10236): 1569-78.
41. Beigel JH, Tomashek KM, Dodd LE, et al. Remdesivir for the Treatment of Covid-19 - Final Report. *N Engl J Med* [9] Salvarani C, Dolci G, Massari M, et al. Effect of Tocilizumab vs Standard Care on Clinical
42. Khoo S, et al. Optimal dose and safety of molnupiravir in patients with early SARS-CoV-2: a Phase I, open-label, dose-escalating, randomized controlled study. *J Antimicrob Chemother*. 2021 Aug 27;dkab318. doi: 10.1093/jac/dkab318. Epub ahead of print. PMID: 34450619.
43. Fischer W, et al. A Phase 2a clinical trial of Molnupiravir in patients with COVID-19 shows accelerated SARS-CoV-2 RNA clearance and elimination of infectious virus. *Sci Transl. Med*. 2021 December 23; DOI: 10.1126/scitranslmed.abl7430
44. Caraco Y, et al. Phase 2/3 Trial of Molnupiravir for Treatment of Covid-19 in Nonhospitalized Adults. *NEJM*. 2021 December 16. DOI: 10.1056/EVIDoa2100043
45. Arribas J. et al. Randomized Trial of Molnupiravir or Placebo in Patients Hospitalized with Covid-19. *NEJM*. 2021 December 16. DOI: 10.1056/EVIDoa2100044
46. Bernal A, da Silva, M.M, Masungie DB, Kovalchuk E, Gonzalez A, Delos Reyes V, Martín-Quiros A, et al. Molnupiravir for Oral Treatment of Covid-19 in Nonhospitalized Patients. *NEJM*. 2021 December 16. DOI: 10.1056/NEJMoa2116044
47. Lou Y, Liu L, Yao H. et al. Clinical outcomes and plasma concentrations of baloxavir marboxil and favipiravir in COVID-19 patients: an exploratory randomized, controlled trial. 2020. *Eur J of Pharm Sci* 157 105631.
48. Macilla-Galindo J, Garcia-Mendez J, Marquez-Sanchez J et a. All-cause mortality among patients treated with repurposed antivirals and antibiotics for COVID-19 in Mexico City: A real-

- world observational study. medRxiv 2020 preprint doi:<https://doi.org/10.1101/2020.10.13.20211797>.
49. Lee S, Park G, Moon Y, Sung K. Clinical characteristics and risk factors for fatality and severity in patients with coronavirus disease in Korea: A nationwide population-based retrospective study using the Korean Health Insurance Review and Assessment Service (HIRA) Database. *Int J Environ Res Public Health* 2020; 17, 8559; doi:10.3390/ijerph1728559.
 50. Liu Q, Fang X, Tian L et al. Arbidol treatment with reduced mortality of adult patients with COVID-19 Wuhan, China: a retrospective cohort study. medRxiv preprint doi:<https://doi.org/10.1101/2020.04.11.20056523>.
 51. Chen X, Zhang Y, Zhu B, et al. Association of clinical characteristics and antiviral drugs with viral RNA clearance in patients with COVID-19 in Guangzhou, China: a retrospective cohort study. medRxiv preprint doi: <https://doi.org/10.1101/2020.04.02.200589941>.
 52. Hu F, Yin G, Chen Y. et al. Corticosteroids, oseltamivir and delayed admission are independent risk factors for prolonged viral shedding in patients with coronavirus disease 2019. *Clin Respir J* 2020; 14:1067-1075. Doi:10.1111/crj.13243
 53. Cao B., Wang, Y., Wen, D., Liu, W., Wang, J., Fan, G., . . . Wang, C. (2020). A trial of lopinavir–ritonavir in adults hospitalized with severe covid-19. *New England Journal of Medicine*, 382(19), 1787-1799. doi:10.1056/nejmoa2001282
 54. Ader, F., Peiffer-Smadja, N., Poissy, J., Bouscambert-Duchamp, M., Belhadi, D., Diallo, A., . . . Mentre, F. (2021). Antiviral drugs in hospitalized patients With covid-19 - the DisCoVeRy trial. doi:10.1101/2021.01.08.20248149 [pre-print]
 55. Li, Y., Xie, Z., Lin, W., Cai, W., Wen, C., Guan, Y., . . . Li, L. (2020 dec 18). Efficacy and safety of LOPINAVIR/RITONAVIR Or ARBIDOL in adult patients with Mild/Moderate COVID-19: An Exploratory randomized controlled trial. *Med*, 1(1), 105-113. doi:10.1016/j.medj.2020.04.001
 56. Horby, P. W., Mafham, M., Bell, J. L., Linsell, L., Staplin, N., Emberson, J., . . . Landray, M. J. (2020 oct 5). Lopinavir–ritonavir in patients admitted to hospital With Covid-19 (RECOVERY): A RANDOMISED, CONTROLLED, OPEN-LABEL, Platform trial. *The Lancet*, 396(10259), 1345-1352. doi:10.1016/s0140-6736(20)32013-4
 57. Pan, H., Peto, R., Karim, Q. A., Alejandria, M., Henao-Restrepo, A. M., García, C. H., . . . Swaminathan, S. (2021). Repurposed antiviral drugs FOR covid-19 –interim Who SOLIDARITY trial results. *New England Journal of Medicine*, 384th ser., 497-511. doi:10.1056/NEJMoa2023184
 58. Gordon AC, Mouncey PR, Al-Beidh F, Rowan KM, Nichol AD, Arabi YM, et al. Interleukin-6 Receptor Antagonists in Critically Ill Patients with Covid-19. *N Engl J Med*. 2021;384(16):1491-502.
 59. Rosas IO, Bräu N, Waters M, Go RC, Hunter BD, Bhagani S, et al. Tocilizumab in Hospitalized Patients with Severe Covid-19 Pneumonia. *N Engl J Med*. 2021;384(16):1503-16.
 60. Salama C, Han J, Yau L, Reiss WG, Kramer B, Neidhart JD, et al. Tocilizumab in Patients Hospitalized with Covid-19 Pneumonia. *N Engl J Med*. 2021;384(1):20-30.
 61. Derde LP, Gordon AC, Mouncey PR, Al-Beidh F, Rowan KM, Nichol AD, et al. Pre-print: Effectiveness of Tocilizumab, Sarilumab, and Anakinra for critically ill patients with COVID-19 The REMAP-CAP COVID-19 Immune Modulation Therapy Domain Randomized Clinical Trial. 2021
 62. Horby PWRG. Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *Lancet*. 2021;397(10285):1637-45.
 63. Hermine O, Mariette X, Tharaux PL, Resche-Rigon M, Porcher R, Ravaud P, et al. Effect of Tocilizumab vs Usual Care in Adults Hospitalized With COVID-19 and Moderate or Severe Pneumonia: A Randomized Clinical Trial. *JAMA Intern Med*. 2021;181(1):32-40.
 64. Hermine O, Mariette X, Tharaux PL, Resche-Rigon M, Simon TM, Porcher R, et al . Pre-print: Tocilizumab Plus Dexamethasone in Patients with Moderate-to-Severe COVID-19 Pneumonia: a Randomized Clinical Trial of the CORIMUNO-19 Study Group. 2021
 65. Salvarani C, Dolci G, Massari M, Merlo DF, Cavuto S, Savoldi L, et al. Effect of Tocilizumab vs Standard Care on Clinical Worsening in Patients Hospitalized With COVID-19 Pneumonia: A Randomized Clinical Trial. *JAMA Intern Med*. 2021;181(1):24-31.
 66. Stone JH, Frigault MJ, Serling-Boyd NJ, Fernandes AD, Harvey L, Foulkes AS, et al. Efficacy of Tocilizumab in Patients Hospitalized with Covid-19. *N Engl J Med*. 2020;383(24):2333-44.
 67. Wang D, Fu B, Peng Z, Yang D, Han M, Li M, et al. Tocilizumab in patients with moderate or severe COVID-19: a randomized, controlled, open-label, multicenter trial. *Front Med*. 2021:1-9.

68. Veiga VC, Prats J, Farias DLC, Rosa RG, Dourado LK, Zampieri FG, et al. Effect of tocilizumab on clinical outcomes at 15 days in patients with severe or critical coronavirus disease 2019: randomised controlled trial. *BMJ*. 2021;372:n84
69. Soin AS, Kumar K, Choudhary NS, Sharma P, Mehta Y, Kataria S, et al. Tocilizumab plus standard care versus standard care in patients in India with moderate to severe COVID-19-associated cytokine release syndrome (COVINTOC): an open-label, multicentre, randomised, controlled, phase 3 trial. *Lancet Respir Med*. 2021;9(5):511-521
70. Rutgers A, Westerweel P, van der Holt B, Simone Postma, van Vonderen MGA, Djura P. Piersma, et al. Preprint: Timely administration of tocilizumab improves survival of hospitalized COVID-19 patients. 2021.
71. Talaschian M, Akhtari M, Mahmoudi M, Mostafaei S, Jafary M, Hussein A, et al. Preprint: Tocilizumab Failed to Reduce Mortality in Severe COVID-19 Patients: Results From a Randomized Controlled Clinical Trial. *Research Square*. 2021.
72. Hamed DM, Belhoul KM, Al Maazmi NA, Ghayoor F, Moin M, Al Suwaidi M, et al. Intravenous methylprednisolone with or without tocilizumab in patients with severe COVID-19 pneumonia requiring oxygen support: A prospective comparison. *Journal of Infection and Public Health*. 2021; 9(8): 985-989
73. Kalil AC, Patterson TF, Mehta AK, Tomashek KM, Wolfe CR, Ghazaryan V, et al. Baricitinib plus Remdesivir for Hospitalized Adults with Covid-19. *N Engl J Med*. 2021;384(9):795-807.
74. Marconi VC, Ramanan AV, de Bono S, et al. Efficacy and safety of baricitinib for the treatment of hospitalised adults with COVID-19 (COV-BARRIER): a randomised, double-blind, parallel-group, placebo-controlled phase 3 trial. *Lancet Respir Med* 2021. doi:10.1016/S2213-2600(21)00331-3.
75. Rosas J, Liano FP, Canto ML et al. Experience with the use of baricitinib and tocilizumab monotherapy or combined, in patients with interstitial pneumonia secondary to Coronavirus COVID19: A real-world study. *Reumatol Clin*. 2020. <https://doi.org/10.1016/j.reuma.2020.10.009>
76. Aman J, Duijvelaar E, Botros L, Kianzad A, Schippers J, Smeele P et al. Imatinib in patients with severe COVID-19: a randomised, double-blind, placebo-controlled, clinical trial. *The Lancet Respiratory Medicine*. 2021;9(9):957-968.
77. Guimaraes PO, Quirk D, Furtado RH, Maia LN, Saraiva JF, Antunes MO, et al. Tofacitinib in Patients Hospitalized with Covid-19 Pneumonia. *N Engl J Med*. 2021;385:406-15.
78. CytoDyn. A Phase 2 Study of Leronlimab for Mild to Moderate Coronavirus Disease 2019 (COVID-19). Available from https://mod.isirv.org/repository/avg_2020/Harish_Seetamraju.pdf. Accessed September 20, 2021.
79. CytoDyn. CytoDyn to Submit Newly Completed Topline Report of CD12 Trial Results to Regulatory Agencies in Multiple Countries including India and Philippines. Available form <https://www.cytodyn.com/newsroom/press-releases/detail/533/cytodyn-to-submit-newly-completed-topline-report-of-cd12>. Accessed September 20, 2021.
80. ClinicalTrials.gov. Study to evaluate the efficacy and safety of leronlimab for mild to moderate COVID-19. Available from <https://clinicaltrials.gov/ct2/show/NCT04343651>. Accessed September 20, 2021.
81. ClinicalTrials.gov. Study to Evaluate the Efficacy and Safety of Leronlimab for Patients With Severe or Critical Coronavirus Disease 2019 (COVID-19). Available from <https://clinicaltrials.gov/ct2/show/NCT04347239>. Accessed September 20, 2021.
82. EU Clinical Trials Register. A Phase 2b/3, Randomized, Double Blind, Placebo Controlled, Adaptive Design Study to Evaluate the Efficacy and Safety of Leronlimab for Patients with Severe or Critical Coronavirus Disease 2019 (COVID-19). Available from <https://www.clinicaltrialsregister.eu/ctr-search/trial/chinaxiv.org2020-001996-33/GB>. Accessed September 20, 2021.
83. Fisher B, et al. Namilumab or infliximab compared to standard of care in hospitalized patients with COVID-19 (CATALYST): a phase 2 randomized adaptive trial. Preprint. Available from: doi: <https://doi.org/10.1101/2021.06.02.21258204>
84. Pang, Jiaojaio, et al. Efficacy and tolerability of bevacizumab in patients with severe COVID-19. *Nature Communications* (2021)12:814
85. Castañeda-Sabogal A, Chambergó-Michilot D, Toro-Huamanchumo CJ, Silva-Rengifo C, Gonzales-Zamora J, Barboza JJ. Outcomes of ivermectin in the treatment of COVID-19: a systematic review and meta-analysis. *medRxiv* 2021; 2021.01.26.21250420.
86. Bryant A, Lawrie TA, Dowswell T, Fordham EJ, Mitchell S, Hill SR, et al. Ivermectin for Prevention and Treatment of COVID-19 Infection: A Systematic Review, Meta-analysis, and Trial Sequential Analysis to Inform Clinical Guidelines. *Am J Ther* 2021; 28: e434–e460.
87. Popp M, Stegemann M, Metzendorf M-I, Gould S, Kranke P, Meybohm P, et al. Ivermectin for

- preventing and treating COVID-19. *Cochrane database Syst Rev* 2021; 7: CD015017.
88. Roman YM, Burela PA, Pasupuleti V, Piscocya A, Vidal JE, Hernandez A V. Ivermectin for the treatment of COVID-19: A systematic review and meta-analysis of randomized controlled trials. *medRxiv* 2021; 2021.05.21.21257595.
89. Zein AFMZ, Sulistiyana CS, Raffaello WM, Pranata R. Ivermectin and mortality in patients with COVID-19: A systematic review, meta-analysis, and meta-regression of randomized controlled trials. *Diabetes Metab Syndr* 2021; 15: 102186.
90. Karale S, Bansal V, Makadia J, Khan H, Spandana Ghanta S, Singh R, et al. A meta-analysis of mortality, need for ICU admission, use of mechanical ventilation and adverse effects with ivermectin use in COVID-19 patients. *medRxiv* 2021; 2021.04.30.21256415.
91. Heidary F, Gharebaghi R. Ivermectin: a systematic review from antiviral effects to COVID-19 complementary regimen. *J Antibiot (Tokyo)* 2020; 73: 593–602.
92. Hill A, Abdulamir AS, Ahmed S, Asghar A, Babalola OE, Basri R, et al. *Preliminary meta-analysis of randomized trials of ivermectin to treat SARS-CoV-2 infection Authors: Andrew Hill on behalf of the International Ivermectin Project Team* International Ivermectin Project Team.*
93. Padhy BM, Mohanty RR, Das S, Meher BR. Therapeutic potential of ivermectin as add on treatment in COVID 19: A systematic review and meta-analysis. *J Pharm Pharm Sci* 2020; 23: 462–469.
94. Rakedzon S, Neuberger A, Domb AJ, Petersiel N, Schwartz E. From hydroxychloroquine to ivermectin: what are the anti-viral properties of anti-parasitic drugs to combat SARS-CoV-2? *J Travel Med* 2021; 28: 1–9.
95. Kalfas S, Visvanathan K, Chan K, Drago J. The therapeutic potential of ivermectin for COVID-19: a systematic review of mechanisms and evidence. *medRxiv* 2020; 2020.11.30.20236570.
96. Kow CS, Merchant HA, Mustafa ZU, Hasan SS. The association between the use of ivermectin and mortality in patients with COVID-19: a meta-analysis. *Pharmacol Rep* 2021; 1–7.
97. Bhowmick S, Dang A, Vallish BN, Dang S. Safety and Efficacy of Ivermectin and Doxycycline Monotherapy and in Combination in the Treatment of COVID-19: A Scoping Review. *Drug safety* 2021; 44: 635–644.
98. Ahmed S, Karim MM, Ross AG, Hossain MS, Clemens JD, Sumiya MK, et al. A five-day course of ivermectin for the treatment of COVID-19 may reduce the duration of illness. *Int J Infect Dis* 2021; 103: 214–216.
99. Podder C, Chowdhury N, Sina M, Haque W. Outcome of ivermectin treated mild to moderate COVID-19 cases: a single-centre, open-label, randomised controlled study. *IMC J Med Sci* 2020; 14: 2.
100. Chaccour C, Casellas A, Blanco-Di Matteo A, Pineda I, Fernandez-Montero A, Ruiz-Castillo P, et al. The effect of early treatment with ivermectin on viral load, symptoms and humoral response in patients with non-severe COVID-19: A pilot, double-blind, placebo-controlled, randomized clinical trial. *EClinicalMedicine*; 32. Epub ahead of print February 2021. DOI: 10.1016/j.eclinm.2020.100720.
101. Bukhari K, Asghar A, Ahmad Mangat S, Abdullah M, Fatima T, Mustafa A. Efficacy of ivermectin in COVID-19 patients with mild to moderate disease. *medRxiv* 2021; 2021.02.02.21250840.
102. Chachar AZK, Khan KA, Asif M, Tanveer K, Khaqan A, Basri R, et al. Effectiveness of Ivermectin in SARS-CoV-2/COVID-19 Patients. *Int J Sci* 2020; 9: 31–35.
103. Abd-Elsalam S, RA N, Badawi R, Khalaf M, ES E, Soliman S, et al. Clinical Study Evaluating the Efficacy of Ivermectin in COVID-19 Treatment: A Randomized Controlled Study. *J Med Virol*. Epub ahead of print 2021. DOI: 10.1002/jmv.27122.
104. Vallejos J, Zoni R, Bangher M, Villamandos S, Bobadilla A, Plano F, et al. Ivermectin to prevent hospitalizations in patients with COVID-19 (IVERCOR-COVID19) a randomized, double-blind, placebo-controlled trial. *BMC Infect Dis* 2021; 21: 635.
105. Krolewiecki A, Lifschitz A, Moragas M, Travacio M, Valentini R, Alonso DF, et al. Antiviral effect of high-dose ivermectin in adults with COVID-19: A proof-of-concept randomized trial. *EClinicalMedicine* 2021; 37: 100959.
106. Mohan A, Tiwari P, TM S, Mittal S, Patel A, Jain A, et al. Single-dose oral ivermectin in mild and moderate COVID-19 (RIVET-COV): A single-centre randomized, placebo-controlled trial. *J Infect Chemother*. Epub ahead of print 2021. DOI: 10.1016/j.jiac.2021.08.021.
107. Kishoria N, Mathur SL, Parmar V, Kaur RJ, Agarwal H, Parihar BS, et al. Ivermectin As Adjuvant To Hydroxychloroquine In Patients Resistant To Standard Treatment For Sars-Cov-2: Results Of An Open-Label Randomized Clinical Study. *Paripex Indian J Res*. Epub ahead of print 2020. DOI: 10.36106/PARIPEX/4801859.
108. Ravikirti R, Pattadar C, Raj R, Agarwal N, Biswas B, PK M, et al. Evaluation of Ivermectin as a

- Potential Treatment for Mild to Moderate COVID-19: A Double-Blind Randomized Placebo Controlled Trial in Eastern India. *J Pharm Pharm Sci* 2021; 24: 343–350.
109. Okumuş N, Demirtürk N, Çetinkaya RA, Sultan H, Han A, Güner R, et al. Evaluation of the effectiveness and safety of adding ivermectin to treatment in severe COVID-19 patients. *ResearchSquare*. Epub ahead of print February 2021. DOI: 10.21203/rs.3.rs-224203/v1.
 110. Gonzalez B, Lenin J, González Gámez M, Antonio E, Enciso M, Josue R, et al. Efficacy and safety of ivermectin and hydroxychloroquine in patients with severe COVID-19: a randomized controlled trial. *medRxiv* 2021; 2021.02.18.21252037.
 111. Pott-Junior H, Bastos Paoliello MM, Miguel A de QC, da Cunha AF, de Melo Freire CC, Neves FF, et al. Use of ivermectin in the treatment of COVID-19: A pilot trial. *Toxicol Reports* 2021; 8: 505–510.
 112. López-Medina E, López P, Hurtado IC, Dávalos DM, Ramirez O, Martínez E, et al. Effect of ivermectin on time to resolution of symptoms among adults with mild COVID-19: a randomized clinical trial. *JAMA - J Am Med Assoc* 2021; 325: 1426–1435.
 113. Biber A, Mandelboim M, Harmelin G, Lev D, Ram L, Shaham A, et al. Favorable outcome on viral load and culture viability using Ivermectin in early treatment of non-hospitalized patients with mild COVID-19 – A double-blind, randomized placebo-controlled trial. *medRxiv* 2021; 2021.05.31.21258081.
 114. Trieu V, Saund S, Rahate P, Barge V, Nalk K, Windlass H et al. Targeting TGF- β pathway with COVID-19 Drug Candidate ARTIVeda/PulmoHeal Accelerates Recovery from Mild-Moderate COVID-19. 2021;.
 115. Lin, Wu, Xie et al. Clinical trial on Artesunate in the treatment of COVID-19. *Chinese Critical Care Medicine*. 2020;.
 116. Shin Poong Pharm. Co., Ltd. A Multi-center, Randomized, Double-blind, Parallel, Placebo-Controlled, Phase II Clinical Trial to Evaluate Efficacy and Safety of Pyramax in Mild to Moderate COVID-19 Patients. South Korea; 2021.
 117. Tardif JC, Bouabdallaoui N, L'Allier PL, Gaudet D, Shah B, Pillinger MH, et al. Efficacy of colchicine in non-hospitalized patients with COVID-19. *Lancet Respir Med*. 2021;9(8):924-932.
 118. Stewart S, Yang KCK, Atkins K, Dalbeth N, Robinson PC. Adverse events during oral colchicine use: a systematic review and meta-analysis of randomised controlled trials. *Arthritis Res Ther* 22, 28 (2020). <https://doi.org/10.1186/s13075-020-2120-7>
 119. Lopes MIF, Bonjorno LP, Giannini MC, Amaral NB, Menezes PI, Dib SM, et al. Beneficial effects of colchicine for moderate to severe COVID-19: a randomised, double-blinded, placebo-controlled clinical trial. *RMD Open*. 2021;7(1):e001455. doi:10.1101/2020.08.06.20169573
 120. Deftereos SG, Giannopoulos G, Vrachatis DA, Siasos GD, Giotaki SG, Gargalianos P, et al. Effect of colchicine vs standard care on cardiac and inflammatory biomarkers and clinical outcomes in patients hospitalized with coronavirus disease 2019: the GRECCO-19 randomized clinical trial. *JAMA Netw Open* 2020;3:e2013136. doi:10.1001/jamanetworkopen.2020.13136
 121. Salehzadeh F, Pourfarzi F, Ataei S. The Impact of Colchicine on The COVID-19 Patients; A Clinical Trial Study. 2020. preprint. 10.21203/rs.3.rs-69374/v1.
 122. RECOVERY Collaborative Group, Horby PW, Campbell M, Spata E, Emberson JR, Staplin N, et al. Colchicine in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *Lancet Respir Med*. 2021;SS2213-2600(21):00435-5. doi:10.1101/2021.05.18.21257267;
 123. Pan H, Peto R, Henao-Restrepo AM, Preziosi MP, Sathiyamoorthy V, Abdool Karim Q, et al. Repurposed Antiviral Drugs for Covid-19 - Interim WHO Solidarity Trial Results. *N Engl J Med*. 2021 Feb 11;384(6):497-511. doi: 10.1056/NEJMoa2023184. Epub 2020 Dec 2. PMID: 33264556; PMCID: PMC7727327.
 124. Davoudi-Monfared E, Rahmani H, Khalili H, Hajiabdolbaghi M, Salehi M, Abbasian L, Kazemzadeh H, Yekaninejad MS. A Randomized Clinical Trial of the Efficacy and Safety of Interferon β -1a in Treatment of Severe COVID-19. *Antimicrob Agents Chemother*. 2020 Aug 20;64(9):e01061-20. doi: 10.1128/AAC.01061-20. PMID: 32661006; PMCID: PMC7449227.
 125. Monk PD, Marsden RJ, Tear VJ, Brookes J, Batten TN, Mankowski M, et al. Inhaled Interferon Beta COVID-19 Study Group. Safety and efficacy of inhaled nebulised interferon beta-1a (SNG001) for treatment of SARS-CoV-2 infection: a randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet Respir Med*. 2021 Feb;9(2):196-206. doi: 10.1016/S2213-2600(20)30511-7. Epub 2020 Nov 12. PMID: 33189161; PMCID: PMC7836724. (Synairgen SG016 Phase 2 trial SNG001)
 126. Ader F, Peiffer-Smadja N, Poissy J, Bouscambert-Duchamp M, Belhadi D, Diallo A, et al. An open-label randomized controlled trial of the effect of lopinavir/ritonavir, lopinavir/ritonavir plus

- IFN- β -1a and hydroxychloroquine in hospitalized patients with COVID-19. *Clinical Microbiology and Infection*. 2021 May 26;S1198-743X(21)00259-7. doi: 10.1016/j.cmi.2021.05.020.
127. Kalil AC, Mehta AK, Patterson TF, Erdmann N, Gomez CA, Jain MK, et al. Efficacy of interferon beta-1a plus remdesivir compared with remdesivir alone in hospitalized adults with COVID-19: a double-blind, randomized, placebo-controlled, phase 3 trial. *Lancet Respir Med*. 2021:S2213-2600(21)00384-2
 128. Darazam IA, Shokouhi S, Pourhoseingholi MA, Irvani SSN, Mokhtari M, Shabani M, et. al. Role of interferon therapy in severe COVID-19: the COVIFERON randomized controlled trial. *Nature Portfolio*. 2021;11:8059
 129. Rahmani H, Davoudi-Monfared ED, Nourian A, Hossein K, Hajizadeh N, Jalalabadi NZ, et. Al. Interferon beta-1b in treatment of severe COVID-19: A randomized clinical trial. *International Immunopharmacology* 2020. 88; 106903
 130. Bhushan S, Wanve S, Koradia P, Bhomia V, Soni P, Chakraborty S, et al. Efficacy and Safety of pegylated interferon- α 2b in moderate COVID-19: a phase 3, randomized, comparator-controlled, open-label study. *International Journal of Infectious Disease*. 2021;111; 281-287
 131. Pandit A, Bhalani N, Bhushan S, Koradia P, Gargiya S, Bhomia V, Kansagra K. Efficacy and safety of pegylated interferon alfa-2b in moderate COVID-19: A phase II, randomized controlled open-label study. *International Journal of Infectious Disease*. 2021;105; 516-521
 132. Simonovich V, Burgos Pratx L, Scibona P, Beruto M, Vallone M, Vázquez C et al. 2021. A Randomized Trial of Convalescent Plasma in Covid-19 Severe Pneumonia. *New England Journal of Medicine*. 2021; 384:619-29. DOI: 10.1056/NEJMoa2031304
 133. Agarwal A, Mukherjee A, Kumar G, Chatterjee P, Bhatnagar T, Malhotra P. 2020. Convalescent plasma in the management of moderate covid-19 in adults in India: open label phase II multicentre-randomised controlled trial (PLACID Trial). *BMJ*. 2020; 371:m3939 <http://dx.doi.org/10.1136/bmj.m393>
 134. Ray Y, Paul S, Bandopadhyay P, D'Rozario R, Sarif J, Lahiri A et al. 2020. Clinical and immunological benefits of convalescent plasma therapy in severe COVID-19: insights from a single center open label randomized control trial. 2020. Preprint. doi: <https://doi.org/10.1101/2020.11.25.20237883>
 135. AlQahtani M, Abdulrahman A, Almadani A, Alali S, Al Zamrooni A, Hejab A et al. 2021. Randomized controlled trial of convalescent plasma therapy against standard therapy in patients with severe COVID-19 disease. *Nature*. 2021; 11:9927 | <https://doi.org/10.1038/s41598-021-89444-5>
 136. Li L, Zhang W, Hu Y, Tong X, Zheng S, Yang J et al. 2020. Effect of Convalescent Plasma Therapy on Time to Clinical Improvement in Patients With Severe and Life-threatening COVID-19. *JAMA*. 2020; 324(5):460.
 137. Gharbharan A, Jordans C, Geurtsvankessel C, den Hollander J, Karim F, Mollema F et al. 2021. Convalescent Plasma for COVID-19. A randomized clinical trial. *Nature*. 2021; 12:3189 | <https://doi.org/10.1038/s41467-021-23469-2>
 138. Avendano-Sola C, Ramos-Martinez A, Munez-Rubio E, Ruiz-Antoran B, Malo de Molina R, Torres F et al. 2021. Convalescent Plasma for COVID-19: A multicenter randomized clinical trial. *J Clin Invest*. 2021; in press <https://doi.org/10.1172/JCI152740>.
 139. Bajpai M, Kumar S, Maheshwari A, Chhabra K, kale P, Gupta A et al. 2020. Efficacy of Convalescent Plasma Therapy compared to Fresh Frozen Plasma in Severely ill COVID-19 Patients: A Pilot Randomized Controlled Trial. 2020. Preprint, doi: <https://doi.org/10.1101/2020.10.25.20219337>
 140. Balcells ME, Rojas L, Le Corre N, Martí'nez-Valdebenito C, Ceballos ME, Ferre's M, et al. 2021. Early versus deferred anti-SARS-CoV-2 convalescent plasma in patients admitted for COVID-19: A randomized phase II clinical trial. *PLoS Med*. 2021; 18(3): e1003415. <https://doi.org/10.1371/journal.pmed.1003415>
 141. Horby, Peter & Estcourt, Lise & Peto, Leon & Emberson, Jonathan & Staplin, Natalie & Spata, Enti & Pessoa-Amorim, Guilherme & Campbell, Mark & Roddick, Alistair & Brunskill, Nigel & George, Tina & Zehnder, Daniel & Tiberi, Simon & Aung, Ni & Uriel, Alison & Widdrington, John & Koshy, George & Brown, Thomas & Scott, Steven & Landray, Martin. 2021. Convalescent plasma in patients admitted to hospital with COVID-19 (RECOVERY): a randomized, controlled, open-label, platform trial. *Lancet*. 2021; 397: 2049–59 [https://doi.org/10.1016/S0140-6736\(21\)00897-7](https://doi.org/10.1016/S0140-6736(21)00897-7) 10.1101/2021.03.09.21252736.
 142. O'Donnell, Max & Grinsztejn, Beatriz & Cummings, Matthew & Justman, Jessica & Lamb, Matt & Eckhardt, Christina & Phillip, Neena & Cheung, Kenneth & Abrams, Darryl & Gupta, Vinay & Diniz, Maria & Cardoso, Sandra & Rajagopalan, Kartik & Borden, Sarah & Wolf, Allison

- &Bitan, Zachary & Meyer, Benjamin & Jacobson, Samuel & Kantor, Alex & Lipin, W. 2021. A randomized, double-blind, controlled trial of convalescent plasma in adults with severe COVID-19. *J Clin Invest.* 2021; 131(13):e150646. <https://doi.org/10.1172/JCI150646>.
143. Pouladzadeh M, Safdarian M, Eshghi P, Abolghasemi H, Bavani AG, Sheibani B, Moradi Choghakabodi P, Fegghi A, GhafourianBoroujerdnia M, Forouzan A, Jalali Far MA, Kaydani GA, Rajaei E, Amin M, Torabizadeh M, Yousefi F, Hadaddezfuli R. 2021. A randomized clinical trial evaluating the immunomodulatory effect of convalescent plasma on COVID-19-related cytokine storm. *Intern Emerg Med.* 2021 Apr 10:1–11. doi: 10.1007/s11739-021-02734-8. Epub ahead of print. PMID: 33837906; PMCID: PMC8035885.
 144. Salman OH, Mohamed HSA. 2020. Efficacy and safety of transfusing plasma from COVID-19 survivors to COVID-19 victims with severe illness. A double-blinded controlled preliminary study. *Egypt J Anaesth.* 2020; 36:264–72. <https://doi.org/10.1080/11101849.2020.18420>
 145. Bennett-Guerrero E, Romeiser JL, Talbot LR, Ahmed T, Mamone LJ, Singh SM, Hearing JC, Salman H, Holiprosad DD, Freedenberg AT, Carter JA, Browne NJ, Cosgrove ME, Shevik ME, Generale LM, Andrew MA, Nachman S, Fries BC. 2021. Stony Brook Medicine COVID Plasma Trial Group. Severe Acute Respiratory Syndrome Coronavirus 2 Convalescent Plasma Versus Standard Plasma in Coronavirus Disease 2019 Infected Hospitalized Patients in New York: A Double-Blind Randomized Trial. *Crit Care Med.* 2021 Apr 16. doi: 10.1097/CCM.0000000000005066. Epub ahead of print. PMID: 33870923.
 146. Körper S, Weiss M, Zickler D, Wiesmann T, Zacharowski K, M.Corman V, Grüner B, Ernst L, Spieth P, Lepper PM, Bentz M, Zinn S, Paul G, Kalbhenn J, Dollinger M, Rosenberger P, Kirschning T, Thiele T, Appl T, Mayer B, Schmidt M, Drosten C, Wulf H, Kruse JM, Jungwirth B, Seifried E, Schrezenmeier H. 2021. CAPSID Clinical Trial Group. High Dose Convalescent Plasma in COVID-19: Results from the Randomized Trial CAPSID. *J Clin Invest.* 2021; in press <https://doi.org/10.1172/JCI152264>.
 147. Bégin P, Callum J, Jamula E, Cook R, Heddle N, Tinmouth A, et. al. 2021. Convalescent plasma for hospitalized patients with COVID-19 and the effect of plasma antibodies: a randomized controlled, open-label trial The CONCOR-1 Study Group. Preprint. doi: <https://doi.org/10.1101/2021.06.29.21259427>
 148. Estcourt LJ. 2021. Convalescent Plasma in Critically ill Patients with Covid-19 The REMAP-CAP Investigators. Preprint. doi: <https://doi.org/10.1101/2021.06.11.21258760>
 149. Balcells ME, Rojas L, Le Corre N, Martí nez-Valdebenito C, Ceballos ME, Ferre´s M, et al. 2021. Early versus deferred anti-SARS-CoV-2 convalescent plasma in patients admitted for COVID-19: A randomized phase II clinical trial. *PLoS Med* 18(3): e1003415. <https://doi.org/10.1371/journal.pmed.1003415>
 150. Kirenga B, ByakikaKibwika P, Muttamba W, et al. 2021. Efficacy of convalescent plasma for treatment of COVID-19 in Uganda. *BMJ Open Res* 2021; 8:e001017. doi: 10.1136/bmjresp-2021-001017
 151. Korley, F.K., V. Durkalski-Mauldin, S.D. Yeatts, K. Schulman, R.D. Davenport, L.J. Dumont, N. El Kassar, L.D. Foster, J.M. Hah, S. Jaiswal, A. Kaplan, E. Lowell, J.F. McDyer, J. Quinn, D.J. Triulzi, C. Van Huysen, V.L.W. Stevenson, K. Yadav, C.W. Jones, B. Kea, A. Burnett, J.C. Reynolds, C.F. Greineder, N.L. Haas, D.G. Beiser, R. Silbergleit, W. Barsan, and C.W. Callaway. 2021. Early Convalescent Plasma for High-Risk Outpatients with Covid-19. *NEJM.* 2021; in press. DOI: 10.1056/NEJMoa2103784
 152. Rasheed A, Fatak D, Hashim H, Maulood M, Kabah K, Almusawi Y, Abdulamir A. 2020. The therapeutic potential of convalescent plasma therapy on treating critically-ill COVID-19 patients residing in respiratory care units in hospitals in Baghdad, Iraq. *Le Infezioni in Medicina.* 2020; 3: 357-366.
 153. Sekine L, Arns B, Fabro BR, et al.2021. Convalescent plasma for COVID19 in hospitalized patients: an open-label randomized clinical trial. *Eur Respir J* 2021; in press (<https://doi.org/10.1183/13993003.01471-2021>).
 154. Gottlieb RL, Nirula A, Chen P, et al. Effect of bamlanivimab as monotherapy or in combination with etesevimab on viral load in patients with mild to moderate COVID-19: A randomized clinical trial. *JAMA - J Am Med Assoc.* [Internet]. 2021;325(7):632-644. Available from: doi:10.1001/jama.2021.0202
 155. Dougan M, Nirula, Azizad M et al. Bamlanivimab plus Etesevimab in Mild or Moderate COVID-19. *N Engl J Med.* 2021. [Internet]. Available from: doi:10.1056/NEJMoa2102685.
 156. Weinreich DM, Sivapalasingam S, Norton T, Ali S, Gao H, Bhore R, et al. REGEN-COV Antibody Combination and Outcomes in Outpatients with Covid-19. *N Engl J Med* [Internet]. 2021 Sep 29 [cited 2021 Oct 10]; Available from: <https://doi.org/10.1056/NEJMoa2108163>

157. Horby PW & Landray MJ. Casirivimab and imdevimab in patients admitted to hospital with COVID-19 (RECOVERY): a randomized, controlled, open-label, platform trial. 2021. Preprint. 10.1101/2021.6.15.21258542.
158. Somersan-Karakaya S, Mylonakis E, Menon VP, Wells JC, Ali S, Sivapalasingam S, et al. REGEN-COV for Treatment of Hospitalized Patients with Covid-19. 2021. Preprint. 10.1101/2021.11.05.21265656.
159. O'Brien MP, Forleo-Neto E, Sarkar N, Isa F, Hou P, et al. Subcutaneous REGEN-COV Antibody Combination in Early Asymptomatic SARS-CoV-2 Infection: A Randomized Controlled Trial. 2021. Preprint. 10.1101/2021.06.14.21258569.
160. Portal-Celhay C, Forleo-Neto E, Eagan W, Musser BJ, Davis JD, Turner KC, Norton T, et al. Phase 2 dose-ranging study of the virologic efficacy and safety of the combination COVID-19 antibodies casirivimab and imdevimab in the outpatient setting. 2021. Pre-print. 10.1101/2021.11.09/21265912.
161. Weinrich DM, Sivapalasingam S, Norton T, Ali S, Gao H, Bhore R, et al. REGEN-COV Antibody Cocktail in Outpatients with Covid-19. 2021. Preprint. 10.1101/2021.06.09.21257915.
162. Kim JY, Jang YR, Hong JH, et al. Safety, Virologic Efficacy, and Pharmacokinetics of CT-P59, a Neutralizing Monoclonal Antibody Against SARS-CoV-2 Spike Receptor-Binding Protein: Two Randomized, Placebo-Controlled, Phase I Studies in Healthy Individuals and Patients With Mild SARS-CoV-2 Infection. 18 Aug 2021. Available from <https://doi.org/10.1016/j.clinthera.2021.08.009>
163. Eom, JS, Ison M, Streinu-Cercel A, et al. Efficacy and safety of CT-P59 plus standard of care: a phase 2/3 randomized, double-blind, placebo controlled trial in outpatients with mild-to-moderate SARS-CoV-2 infection. 15 March 2021. Available from <https://www.researchsquare.com/article/rs-296518/v1>
164. Gharebaghi N, Nejadrahim R, Mousavi SJ, Sadat-Ebrahimi S-R, Hajizadeh R. The use of intravenous immunoglobulin gamma for the treatment of severe coronavirus disease 2019: a randomized placebo-controlled double-blind clinical trial. *BMC Infect Dis.* 2020;20(1):786. doi:10.1186/s12879-020-05507-4
165. Raman RS, Bhagwan Barge V, Anil Kumar D, et al. A Phase II Safety and Efficacy Study on Prognosis of Moderate Pneumonia in Coronavirus Disease 2019 Patients With Regular Intravenous Immunoglobulin Therapy. *J Infect Dis.* February 2021. doi:10.1093/infdis/jiab098
166. Sakoulas G, Geriak M, Kullar R, et al. Intravenous Immunoglobulin (IVIG) Significantly Reduces Respiratory Morbidity in COVID-19 Pneumonia: A Prospective Randomized Trial. *medRxiv.* January 2020:2020.07.20.20157891. doi:10.1101/2020.07.20.20157891
167. Tabarsi P, Barati S, Jamaati H, et al. Evaluating the effects of Intravenous Immunoglobulin (IVlg) on the management of severe COVID-19 cases: A randomized controlled trial. *Int Immunopharmacol.* 2021;90(January):107205. doi:10.1016/j.intimp.2020.107205
168. Shu L, Niu C, Li R, Huang T, Wang Y, Huang M, et al. Treatment of severe COVID-19 with human umbilical cord mesenchymal stem cells. *Stem Cell Res Ther.* 2020;11(1):1–11.
169. Lanzoni G, Linetsky E, Correa D, Messinger Cayetano S, Alvarez RA, Kouroupis D, et al. Umbilical cord mesenchymal stem cells for COVID-19 acute respiratory distress syndrome: A double-blind, phase 1/2a, randomized controlled trial. *Stem Cells Transl Med.* 2021;10(5):660–73.
170. Shi L, Huang H, Lu X, Yan X, Jiang X, Xu R, et al. Effect of human umbilical cord-derived mesenchymal stem cells on lung damage in severe COVID-19 patients: a randomized, double-blind, placebo-controlled phase 2 trial. *Signal Transduct Target Ther.* 2021;6(1).
171. Yu LM, Bafadhel M, Dorward J, Hayward G, Saville BR, Gbinigie O, Van Hecke O, Ogburn E, Evans PH, Thomas NPB, Patel MG, Richards D, Berry N, Detry MA, Saunders C, Fitzgerald M, Harris V, Shanyinde M, de Lusignan S, Andersson MI, Barnes PJ, Russell REK, Nicolau DV, Ramakrishnan S, Hobbs FDR, Butler CC, PRINCIPLE Trial Collaborative Group. 2021. Inhaled budesonide for COVID-19 in people at high risk of complications in the community in the UK (PRINCIPLE): a randomized, controlled, open-label, adaptive platform trial. *Lancet* 2021; 398: 843–55. Published Online August 10, 2021. [https://doi.org/10.1016/S0140-6736\(21\)01744-X](https://doi.org/10.1016/S0140-6736(21)01744-X)
172. Ramakrishnan S, Nicolau D, Langford B, Mahdi M, Jeffers H, Mwasuku C, Krassowska K, Fox R, Binnian I, Glover V, Bright S, Butler C, Cane JL, Halner A, Matthews PC, Donnelly LE, Simpson JL, Baker JR, Fadai NT, Peterson S, Bengtsson T, Barnes PJ, Russell REK, Bafadhel M. 2021. Inhaled budesonide in the treatment of early COVID-19 (STOIC): a phase 2, open-label, randomized controlled trial. *Lancet Respir Med* 2021; 9: 763–72. Published Online April 9, 2021 [https://doi.org/10.1016/S2213-2600\(21\)00160-0](https://doi.org/10.1016/S2213-2600(21)00160-0)
173. Song, J.-Y.; Yoon, J.-G.; Seo, Y.-B.; Lee, J.; Eom, J.-S.; Lee, J.-S.; Choi, W.-S.; Lee, E.-Y.;

- Choi, Y.-A.; Hyun, H.-J.; et al. 2021. Ciclesonide Inhaler Treatment for Mild-to-Moderate COVID-19: A Randomized, Open-Label, Phase 2 Trial. *J. Clin. Med.* 2021, 10, 3545. <https://doi.org/10.3390/jcm10163545>
174. Clemency B, Varughese R, Gonzalez-Rojas Y, Morse CG, Phipatanakul W, Koster DJ, Blaiss MS. 2021. A randomized controlled trial of inhaled ciclesonide for outpatient treatment of symptomatic COVID-19 infections. *medRxiv preprint*. September 12, 2021. doi: <https://doi.org/10.1101/2021.09.07.21261811>.
 175. la Marca G, Barp J, Frenos S, Mugelli A, Galli L, Calistri E, et al. Thermal inactivation of SARS COVID-2 virus: Are steam inhalations a potential treatment? *Life Sci [Internet]*. 2021 Jan 15 [cited 2021 Feb 22];265. Available from: <https://pubmed.ncbi.nlm.nih.gov/33232690/>
 176. Wang D, Fu B, Peng Z, et al. Preprint. Tocilizumab ameliorates the hypoxia in COVID-19 moderate patients with bilateral pulmonary lesions: a randomized, controlled, open-label, multicenter trial. SSRN 2020.
 177. Salama C, Han J, Yau L, et al. Tocilizumab in Patients Hospitalized with Covid-19 Pneumonia. *N Engl J Med* 2021; 384(1): 20-30
 178. Dayrit C. Coconut oil in health and disease: its and monolaurin's potential as cure for HIV/AIDS. 2000. Accessed from: <http://coconutresearchcenter.org/wp-content/uploads/2015/11/article10526.pdf>
 179. Widhiarta KD. Virgin coconut oil for HIV-positive people. *Cord.* 2016; 3(1):50-57.
 180. Freedberg DE, Conigliaro J, Wang TC, Tracey KJ, Callahan MV, Abrams JA; Famotidine Research Group. Famotidine use is associated with improved clinical outcomes in hospitalized COVID-19 Patients: a propensity score matched retrospective cohort study. *Gastroenterology*. 2020 Sep;159(3):1129-1131.e3. doi: 10.1053/j.gastro.2020.05.053.
 181. Mura C, Preissner S, Nahles S, Heiland M, Bourne PE, Preissner R. Clinical evidence for improved outcomes with histamine antagonists and aspirin in 22,560 COVID-19 patients. *medRxiv*. 2021 Apr 5. doi: 10.1101/2021.03.29.21253914
 182. Cheung KS, Hung IFN, Leung WK. Association between famotidine use and COVID-19 severity in hong kong: a territory-wide study. *Gastroenterology*. 2021 Apr;160(5):1898-1899. doi: 10.1053/j.gastro.2020.05.098.
 183. Mather JF, Seip RL, McKay RG. Impact of famotidine use on clinical outcomes of hospitalized patients with COVID-19. *Am J Gastroenterol*. 2020 Oct;115(10):1617-1623. doi: 10.14309/ajg.0000000000000832.
 184. Shoaibi A, Fortin SP, Weinstein R, Berlin JA, Ryan P. Comparative effectiveness of famotidine in hospitalized COVID-19 patients. *Am J Gastroenterol*. 2021 Apr;116(4):692-699. doi: 10.14309/ajg.0000000000001153
 185. Yeramneni S, Doshi P, Sands K, Cooper M, Kurbegov D, Fromell G. Famotidine use is not associated with 30-day mortality: a coarsened exact match study in 7158 hospitalized patients with coronavirus disease 2019 from a large healthcare system. *Gastroenterology*. 2021 Feb;160(3):919-921.e3. doi: 10.1053/j.gastro.2020.10.011.
 186. Zhou J, Wang X, Lee S, Wu WKK, Cheung BMY, Zhang Q, Tse G. Proton pump inhibitor or famotidine use and severe COVID-19 disease: a propensity score-matched territory-wide study. *Gut*. Epub ahead of print: 04 December 2020. doi: 10.1136/gutjnl-2020-323668.
 187. Lenze EJ, Mattar C, Zorumski CF, Stevens A, Schweiger J, Nicol GE, et al. Fluvoxamine vs Placebo and Clinical Deterioration in Outpatients With Symptomatic COVID-19. *JAMA Intern Med*. 2020;324(22):2292-2300
 188. Reis G, dos Santos Moreira-Silva EA, Silva DCM, Thabane L, Milagres AC, Ferreira TS, et al. Effect of early treatment with fluvoxamine on risk of emergency care and hospitalisation among patients with COVID-19: the TOGETHER randomised, platform clinical trial. *Lancet Glob Health*. 2021 Oct;S2214109X21004484.

Evidence and Recommendations for the Critical Care and Respiratory Management of COVID-19

1. Dequin P-F, Heming N, Meziani F, Plantefève G, Voiriot G, Badié J, et al. Effect of hydrocortisone on 21-day mortality or respiratory support among critically ill patients with COVID-19. *JAMA*. 2020Sep2;324(13).
2. Tomazini BM, Maia IS, Cavalcanti AB, Berwanger O, Rosa RG, Veiga VC, et al. Effect of dexamethasone on days alive and ventilator-free in patients with moderate or severe acute respiratory distress syndrome and COVID-19. *JAMA*. 2020Oct6;324(13):1307–16.

3. Munch MW, Meyhoff TS, Helleberg M, Kjær MBN, Granholm A, Hjortsø CJ, et al. Low-dose hydrocortisone in patients with COVID-19 and severe hypoxia: The COVID steroid randomised, placebo-controlled trial. *Acta Anaesthesiologica Scandinavica*. 2021Jun17;65(10):1421–30.
4. Sterne JA, Murthy S, Diaz JV, Slutsky AS, Villar J, Angus DC, et al. Association between administration of systemic corticosteroids and mortality among critically ill patients with COVID-19. *JAMA*. 2020Oct6;324(13):1330–241.
5. Edalatfard M, Akhtari M, Salehi M, Naderi Z, Jamshidi A, Mostafaei S, et al. Intravenous methylprednisolone pulse as a treatment for hospitalised severe COVID-19 patients: Results from a randomised controlled clinical trial. *European Respiratory Journal*. 2020;56(6).
6. Ghanei M, Solaymani-Dodaran M, Qazvini A, Ghazale AH, Setarehdan SA, Saadat SH, et al. The efficacy of corticosteroids therapy in patients with moderate to severe SARS-COV-2 infection: A Multicenter, randomized, open-label trial. *Respiratory Research*. 2021Sep15;22(245).
7. Corral-Gudino L, Bahamonde A, Arnaiz-Revillas F, Gómez-Barquero J, Abadía-Otero J, García-Ibarbia C, et al. Methylprednisolone in adults hospitalized with COVID-19 pneumonia. *Wiener klinische Wochenschrift*. 2021Oct9;133(7-8):303–11.
8. Jamaati H, Hashemian SMR, Farzanegan B, Malekmohammad M, Tabarsi P, Marjani M, et al. No clinical benefit of high dose corticosteroid administration in patients with COVID-19: A preliminary report of a randomized clinical trial. *European Journal of Pharmacology*. 2021Feb16;897:173947.
9. Jeronimo CM, Farias ME, Val FF, Sampaio VS, Alexandre MA, Melo GC, et al. Methylprednisolone as adjunctive therapy for patients hospitalized with coronavirus disease 2019 (COVID-19; MetCOVID): A randomized, double-blind, phase iib, placebo-controlled trial. *Clinical Infectious Diseases*. 2020May1;72(9).
10. Horby P, Lim WS, Emberson JR, Mafham M, Bell JL, Linsell L, et al. Dexamethasone in hospitalized patients with covid-19. *New England Journal of Medicine*. 2021Feb25;384(8):693–704.
11. Angus DC, Derde L, Al-Beidh F, Annane D, Arabi Y, Beane A, et al. Effect of hydrocortisone on mortality and organ support in patients with severe COVID-19. *JAMA*. 2020Oct6;324(13).
12. Solanich X, Antolí A, Rocamora-Blanch G, Padullés N, Fanlo-Maresma M, Iriarte A, et al. Methylprednisolone pulses plus tacrolimus in addition to standard of care vs. standard of care alone in patients with severe COVID-19. A randomized controlled trial. *Frontiers in Medicine*. 2021Jun14;8.
13. Tang X, Feng Y-M, Ni J-X, Zhang J-Y, Liu L-M, Hu K, et al. Early use of corticosteroid may prolong SARS-COV-2 shedding in non-intensive care unit patients with COVID-19 pneumonia: A Multicenter, single-blind, randomized control trial. *Respiration*. 2021Jan22;100(2):116–26.
14. Farahani RH, Mosaed R, Nezami-Asl A, chamanara M, Soleiman-Meigooni S, Kalantar S, et al. Evaluation of the efficacy of methylprednisolone pulse therapy in treatment of COVID-19 adult patients with severe respiratory failure: Randomized, clinical trial. *Research Square [Internet]*. 2020Sep9 [cited 2021Nov20]; Available from: <https://www.researchsquare.com/article/rs-66909/v1>
15. Munch MW, Russell L, Uhre KR, Lindgaard AL, Degn JF, Wetterslev M, et al. Effect of 12 mg vs 6 mg of dexamethasone on the number of days alive without life support in adults with COVID-19 and severe hypoxemia. *JAMA*. 2021Nov9;326(18).
16. Bahl A, Johnson S, Chen NW. Timing of corticosteroids impacts mortality in hospitalized COVID-19 patients. *Intern Emerg Med*. 2021 Sep;16(6):1593-1603. doi: 10.1007/s11739-021-02655-6. Epub 2021 Feb 5. PMID: 33547620; PMCID: PMC7864133.
17. Sulaiman K, Al Juhani O, Korayen GB, Eljaaly K, Alhubaishi A, Al Harbi O, Badreldin HA, Al Yousif GA, Vishwakarma R, Altebainawi A, Albelwi S, Almutairi R, Almousa M, Alghamdi R, Alharbi A, Algami R, Akhani N, Al Hartin A, Alissa A, Al Homaid S, Al Qahtani K, Al Atassi A, Al Ghamdi G. Early Versus Late Use of Dexamethasone in Critically Ill Patients with COVID-19: A Multicenter, Cohort Study. *ResearchSquare [Preprint]*. 2021 Jul 26 [cited 2021 Nov 27]. Available from <https://www.researchsquare.com/article/rs-349677/v1>
18. Monedero P, Gea A, Castro P, Candela-Toha AM, Hernández-Sanz ML, Arruti E, Villar J, Ferrando C; COVID-19 Spanish ICU Network. Early corticosteroids are associated with lower mortality in critically ill patients with COVID-19: a cohort study. *Crit Care*. 2021 Jan 4;25(1):2. doi: 10.1186/s13054-020-03422-3. PMID: 33397463; PMCID: PMC7780210.
19. Akhtar H, Khalid S, Rahman FU, Ali S, Afridi M, Khader YS, Hassan F, Akhtar N, Khan MM, Ikram A. Delayed admissions and efficacy of steroid use in patients with critical and severe COVID-19: an apprehensive approach. *J Public Health (Oxf)*. 2021 Sep 27;fdab239. doi: 10.1093/pubmed/fdab239.

20. Dupuis C, de Montmollin E, Buetti N, Goldgran-Toledano D, Reignier J, Schwebel C, Domitile J, Neuville M, Ursino M, Siami S, Ruckly S, Alberti C, Mourvillier B, Bailly S, Laurent V, Gainnier M, Souweine B, Timsit JF; OutcomeReaTM research network. Impact of early corticosteroids on 60-day mortality in critically ill patients with COVID-19: A multicenter cohort study of the OUTCOMEREA network. *PLoS One*. 2021 Aug 4;16(8):e0255644. doi: 10.1371/journal.pone.0255644. PMID: 34347836; PMCID: PMC8336847.
21. Moreno A, Vargas C, Azocar F, Villarroel F, Cofré M, Oppliger H, Ríos F, Raijmakers M, Silva-Ayarza I, Beltrán C, Zamora F. Steroids and mortality in non-critically ill COVID-19 patients: a propensity score-weighted study in a Chilean cohort. *Int J Infect Dis*. 2021 Sep 20;112:124-129. doi: 10.1016/j.ijid.2021.09.038. Epub ahead of print. PMID: 34547488; PMCID: PMC8450146.
22. Li Y, Zhou X, Li T, Chan S, Yiqi Y, Ai JW, Zhang H, Sun F, Zhang Q, Zhu L, Shao L, Xu B, Zhang W. Corticosteroid prevents COVID-19 progression within its therapeutic window: a multicentre, proof-of-concept, observational study. *Emerg. Microbes Infect.*, 9:1, 1869-1877, DOI: 10.1080/22221751.2020.1807885
23. Lopes RD, Silva PGM de B e, Furtado RHM, Macedo AVS, Bronhara B, Damiani LP, et al. Therapeutic versus prophylactic anticoagulation for patients admitted to hospital with COVID-19 and elevated D-dimer concentration (ACTION): an open-label, multicentre, randomised, controlled trial. *Lancet [Internet]*. 2021 Jun 12 [cited 2021 Sep 11];397(10291):2253–63. Available from: <http://www.thelancet.com/article/S0140673621012034/fulltext>
24. The ATTACC A-4a, and R-CI. Therapeutic Anticoagulation with Heparin in Noncritically Ill Patients with Covid-19. <https://doi.org/10.1056/NEJMoa2105911> [Internet]. 2021 Aug 4 [cited 2021 Sep 11];385(9):790–802. Available from: <https://www.nejm.org/doi/10.1056/NEJMoa2105911>
25. Sholzberg M, Tang GH, Rahhal H, AlHamzah M, Kreuziger LB, Áinle FN, et al. Heparin for Moderately Ill Patients with Covid-19. *medRxiv [Internet]*. 2021 Jul 9 [cited 2021 Sep 25];2021.07.08.21259351. Available from: <https://www.medrxiv.org/content/10.1101/2021.07.08.21259351v2>
26. The REMAP-CAP A-4a, and AI. Therapeutic Anticoagulation with Heparin in Critically Ill Patients with Covid-19. <https://doi.org/10.1056/NEJMoa2103417> [Internet]. 2021 Aug 4 [cited 2021 Sep 11];385(9):777–89. Available from: <https://www.nejm.org/doi/10.1056/NEJMoa2103417>
27. ACB L, DA do ES, MC S, RN G, LB A, A P-F, et al. Therapeutic versus prophylactic anticoagulation for severe COVID-19: A randomized phase II clinical trial (HESACOVID). *Thromb Res [Internet]*. 2020 Dec 1 [cited 2021 Sep 11];196:359–66. Available from: <https://pubmed.ncbi.nlm.nih.gov/32977137/>
28. Spyropoulos AC, Goldin M, Giannis D, Diab W, Wang J, Khanijo S, et al. Efficacy and Safety of Therapeutic-Dose Heparin vs Standard Prophylactic or Intermediate-Dose Heparins for Thromboprophylaxis in High-risk Hospitalized Patients With COVID-19: The HEP-COVID Randomized Clinical Trial. *JAMA Intern Med [Internet]*. 2021 Oct 7 [cited 2021 Oct 12]; Available from: <https://jamanetwork.com/journals/jamainternalmedicine/fullarticle/2785004>
29. US P, I C, A W, P TE, C W, S D, et al. Standard prophylactic versus intermediate dose enoxaparin in adults with severe COVID-19: A multi-center, open-label, randomized controlled trial. *J Thromb Haemost [Internet]*. 2021 Sep 1 [cited 2021 Sep 11];19(9):2225–34. Available from: <https://pubmed.ncbi.nlm.nih.gov/34236768/>
30. Mazloomzadeh S, Khaleghparast S, Ghadrdoost B, Mousavizadeh M, Baay MR, et al. Effect of Intermediate-Dose vs Standard-Dose Prophylactic Anticoagulation on Thrombotic Events, Extracorporeal Membrane Oxygenation Treatment, or Mortality Among Patients With COVID-19 Admitted to the Intensive Care Unit: The INSPIRATION Randomized Clinical Trial. *JAMA [Internet]*. 2021 Apr 27 [cited 2021 Sep 11];325(16):1620–30. Available from: <https://jamanetwork.com/journals/jama/fullarticle/2777829>
31. Buetti N., Mazzuchelli, T., Lo Priore, E., Balmelli, C., Llamas, M., Pallanza, M., Elzi, L., Consonni, V., Trimboli, P., Forni-Ogna, V. and Bernasconi, E., 2020. Early administered antibiotics do not impact mortality in critically ill patients with COVID-19. *Journal of Infection*, 81(2), pp.e148-e149. Available at: <www.sciencedirect.com/.../journal-of-infection/vol/81/issue/2> [Accessed 5 April 2021].
32. Soleimani A, Moeini Taba SM, Hasibi Taheri S, Loghman AH, Shayestehpour M. The effect of Hemoperfusion on Outcome, Clinical and Laboratory findings of Patients with Severe COVID-19: A Retrospective Study [published online ahead of print, 2021 Sep 2]. *New Microbes New Infect*. 2021;100937. doi:10.1016/j.nmni.2021.100937

33. Darazam I, Kazempour M, Pourhoseingholi M, Hatami F, Rabiei M, Gharehbagh F, et al. Efficacy of Hemoperfusion in Severe and Critical Cases of COVID-19. Preprint version Research square. Available from: <https://doi.org/10.21203/rs.3.rs-555429/v1>
34. Hu W, Lin CW, Liu BW, Hu WH, Zhu Y. Extravascular lung water and pulmonary arterial wedge pressure for fluid management in patients with acute respiratory distress syndrome. *Multidiscip Respir Med*. 2014 Jan 16;9(1):3.
35. Wiedemann HP, Wheeler AP, Bernard GR, Thompson BT, Hayden D, deBoisblanc B, Connors AF Jr, Hite RD, Harabin AL, National Heart, Lung, and Blood Institute Acute Respiratory Distress Syndrome (ARDS) Clinical Trials Network. Comparison of two fluid-management strategies in acute lung injury. *N Engl J Med*. 2006 Jun 15;354(24):2564-75.
36. Martin GS, Mangialardi RJ, Wheeler AP, Dupont WD, Morris JA, Bernard GR. Albumin and furosemide therapy in hypoproteinemic patients with acute lung injury. *Crit Care Med*. 2002 Oct;30(10):2175-82.
37. Martin GS, Moss M, Wheeler AP, Mealer M, Morris JA, Bernard GR. A randomized, controlled trial of furosemide with or without albumin in hypoproteinemic patients with acute lung injury. *Crit Care Med*. 2005 Aug;33(8):1681-7.
38. Rosén J, von Oelreich E, Fors D, Jonsson Fagerlund M, Taxbro K, Skorup P, et al. Awake prone positioning in patients with hypoxemic respiratory failure due to COVID-19: the PROFLO multicenter randomized clinical trial. *Ann Am Thorac Soc* [Internet]. 2021;7(1):00692–2020. Available from: <http://dx.doi.org/10.1183/23120541.00692-2020>
39. Kharat A, Dupuis-Lozeron E, Cantero C, Marti C, Groscurin O, Lolachi S, et al. Self-proning in COVID-19 patients on low-flow oxygen therapy: a cluster randomised controlled trial. *ERJ Open Res* [Internet]. 2021;7(1):00692–2020. Available from: <http://dx.doi.org/10.1183/23120541.00692-2020>
40. Jayakumar D, Ramachandran, DNB P, Rabindrarajan, DNB E, Vijayaraghavan, MD BKT, Ramakrishnan, AB N, Venkataraman, AB R. Standard Care Versus Awake Prone Position in Adult Nonintubated Patients With Acute Hypoxemic Respiratory Failure Secondary to COVID-19 Infection—A Multicenter Feasibility Randomized Controlled Trial. *J Intensive Care Med*. 2021;36(8):918–24.
41. Taylor SP, Bundy H, Smith WM, Skavroneck S, Taylor B, Kowalkowski MA. Awake prone positioning strategy for nonintubated hypoxic patients with covid-19 a pilot trial with embedded implementation evaluation. *Ann Am Thorac Soc*. 2021;18(8):1360–8.
42. Miček M, Otáhal M, Borges JB, Alcala GC, Hladík D, Kuriščák E, et al. Targeted lateral positioning decreases lung collapse and overdistension in COVID-19-associated ARDS. *BMC Pulm Med* [Internet]. 2021;21(1):1–7. Available from: <https://doi.org/10.1186/s12890-021-01501-x>
43. Teng X bao, Shen Y, Han M feng, Yang G, Zha L, Shi J feng. The value of high-flow nasal cannula oxygen therapy in treating novel coronavirus pneumonia. *Eur J Clin Invest*. 2021;51(3):0–1.
44. Perkins GD, Ji C, Connolly BA, Couper K, Lall R, Baillie JK, et al. An adaptive randomized controlled trial of non-invasive respiratory strategies in acute respiratory failure patients with COVID-19. *medRxiv* [Internet]. 2021;2021.08.02.21261379. Available from: <https://www.medrxiv.org/content/10.1101/2021.08.02.21261379v1%0Ahttps://www.medrxiv.org/content/10.1101/2021.08.02.21261379v1.abstract>
45. Grieco DL, Menga LS, Cesarano M, Rosà T, Spadaro S, Bitondo MM, et al. Effect of Helmet Noninvasive Ventilation vs High-Flow Nasal Oxygen on Days Free of Respiratory Support in Patients with COVID-19 and Moderate to Severe Hypoxemic Respiratory Failure: The HENIVOT Randomized Clinical Trial. *JAMA - J Am Med Assoc*. 2021 May 4;325(17):1731–43.
46. Nair PR, Haritha D, Behera S, Kayina CA, Maitra S, Anand RK, et al. Comparison of High-Flow Nasal Cannula and Noninvasive Ventilation in Acute Hypoxemic Respiratory Failure Due to Severe COVID-19 Pneumonia. *Respir Care*. 2021;66(12):1824–30.
47. Botta M, Tsonas AM, Pillay J, Boers LS, Algera AG, Bos LDJ, et al. Ventilation management and clinical outcomes in invasively ventilated patients with COVID-19 (PRoVENT-COVID): a national, multicentre, observational cohort study. *Lancet Respir Med*. 2020;19(20):1–10.
48. Udi J, Lang CN, Zotzmann V, Krueger K, Fluegle A, Bamberg F, et al. Incidence of Barotrauma in Patients With COVID-19 Pneumonia During Prolonged Invasive Mechanical Ventilation – A Case-Control Study. *J Intensive Care Med*. 2020;
49. Romano MLP, Maia IS, Laranjeira LN, Damiani LP, Paisani DDM, Borges MDC, et al. Driving Pressure-limited Strategy for Patients with Acute Respiratory Distress Syndrome A Pilot Randomized Clinical Trial. *Ann Am Thorac Soc*. 2020;17(5):596–604.

50. Yao W, Wang T, Jiang B, Gao F, Wang L, Zheng H et al. Emergency tracheal intubation in 202 patients with COVID-19 in Wuhan, China: lessons learnt and international expert recommendations. *British Journal of Anaesthesia*. 2020;125(1):e28-e37.
51. Weingart S, Trueger N, Wong N, Scofi J, Singh N, Rudolph S. Delayed Sequence Intubation: A Prospective Observational Study. *Annals of Emergency Medicine*. 2015;65(4):349-355.
52. Cheng W, Ma XD, Su LX, Long Y, Liu DW, Du B, et al. Retrospective Study of Critically Ill COVID-19 Patients With and Without Extracorporeal Membrane Oxygenation Support in Wuhan, China. *Front Med (Lausanne)*. 2021 Oct 12;8:659793.
53. Li S, Xiong J, Du Z, Lai W, Ma X, et al. Extracorporeal membrane oxygenation (ECMO) for critically ill patients with coronavirus disease 2019 (COVID-19): A retrospective cohort study. *J Card Surg*. 2021 Oct;36(10):3554-3560.
54. Mustafa AK, Joshi DJ, Alexander PJ, Tabachnick DR, Cross CA, et al. Comparative Propensity Matched Outcomes in Severe COVID-19 Respiratory Failure-Extracorporeal Membrane Oxygenation or Maximum Ventilation Alone. *Ann Surg*. 2021 Nov 1;274(5):e388-e394.
55. Nguyen NT, Sullivan B, Sagebin F, Hohmann SF, Amin A, Nahmias J. Analysis of COVID-19 Patients With Acute Respiratory Distress Syndrome Managed With Extracorporeal Membrane Oxygenation at US Academic Centers. *Ann Surg*. 2021 Jul 1;274(1):40-44.
56. Gorenstein SA, Castellano ML, Slone ES, Gillette B, Liu H, Alsamarraie C, et al. Hyperbaric oxygen therapy for COVID-19 patients with respiratory distress: treated cases versus propensity-matched controls. *Undersea Hyperb Med*. 2020;47(3):405-13.
57. Petrikov SS, Evseev AK, Levina OA, Shabanov AK, Kulabukhov V v., Kutrovskaya NY, et al. Hyperbaric oxygen therapy in patients with COVID-19. *Obshchaya Reanimatologiya*. 2020;16(6):4-18.
58. Olsen HT, Nedergaard HK, Strøm T, Oxlund J, Wian KA, Ytrebø LM, et al. Nonsedation or Light Sedation in Critically Ill, Mechanically Ventilated Patients. *N Engl J Med*. 2020 Mar 19;382(12):1103-1111. doi: 10.1056/NEJMoa1906759. Epub 2020 Feb 16. PMID: 32068366.
59. Strøm T, Martinussen T, Toft P. A protocol of no sedation for critically ill patients receiving mechanical ventilation: a randomised trial. *Lancet*. 2010 Feb 6;375(9713):475-80. doi: 10.1016/S0140-6736(09)62072-9. Epub 2010 Jan 29. PMID: 20116842.
60. Forel JM, Roch A, Marin V, Michelet P, Demory D, Blache JL, et al. Neuromuscular blocking agents decrease inflammatory response in patients presenting with acute respiratory distress syndrome. *Crit Care Med* 2006;34:2749-57.
61. Gainnier M, Roch A, Forel JM, Thirion X, Arnal JM, Donati S, et al. Effect of neuromuscular blocking agents on gas exchange in patients presenting with acute respiratory distress syndrome. *Crit Care Med* 2004;32:113-9.
62. Guervilly C, Bisbal M, Forel JM, Mechat M, Lehingue S, Bourenne J, et al. Effects of neuromuscular blockers on transpulmonary pressures in moderate to severe acute respiratory distress syndrome. *Intensive Care Med* 2017;43:408-18.
63. Lyu G, Wang X, Jiang W, Cai T, Zhang Y. [Clinical study of early use of neuromuscular blocking agents in patients with severe sepsis and acute respiratory distress syndrome]. *Zhonghua Wei Zhong Bing Ji Jiu Yi Xue* 2014;26:325-9.
64. National Heart, Lung, and Blood Institute PETAL Clinical Trials Network, Moss M, Huang DT, Brower RG, Ferguson ND, Ginde AA, Gong MN, et al. Early Neuromuscular Blockade in the Acute Respiratory Distress Syndrome. *N Engl J Med*. 2019 May 23;380(21):1997-2008. doi: 10.1056/NEJMoa1901686. Epub 2019 May 19. PMID: 31112383; PMCID: PMC6741345.
65. Papazian L, Forel JM, Gacouin A, Penot-Ragon C, Perrin G, Loundou A, et al.; ACURASYS Study Investigators. Neuromuscular blockers in early acute respiratory distress syndrome. *N Engl J Med* 2010;363:1107-16.
66. Moni M, Madathil T, Sathyapalan D, Menon V, Gutjahr G, Edathadathil F, et al. A feasibility trial to evaluate the composite efficacy of inhaled Nitric Oxide in the treatment of Covid 19 pneumonia : Impact on viral load and clinical outcomes [Internet]. *bioRxiv*. 2021. Available from: <http://dx.doi.org/10.1101/2021.04.15.21255300>.
67. Montero-Baladia, M., et al. "Etoposide treatment adjunctive to immunosuppressants for critically ill COVID-19 patients." *The Journal of Infection*, 81(3) (2020), 452-482. <https://doi.org/10.1016/j.jinf.2020.06.006>
68. Patel, M., Dominguez, E., Sacher, D., Desai, P., Chandar, A., Bromberg, M., Caricchio, R., Criner, G. J., & Temple University COVID-19 Research Group (2021). Etoposide as Salvage Therapy for Cytokine Storm Due to Coronavirus Disease 2019. *Chest*, 159(1), e7-e11. <https://doi.org/10.1016/j.chest.2020.09.077>

69. Liu K, Zhang W, Yang Y, Zhang J, Li Y, Chen Y. Respiratory rehabilitation in elderly patients with COVID-19: A randomized controlled study. *Complement Ther Clin Pract* [Internet]. 2020;39:101166. Available from: <https://doi.org/10.1016/j.ctcp.2020.101166> <https://doi.org/10.1016/j.ctcp.2020.101166>
70. He M, Yu S, Wang L, Lv H, Qiu Z. Efficiency and safety of pulmonary rehabilitation in acute exacerbation of chronic obstructive pulmonary disease. *Med Sci Monit*. 2015 Mar 18;21:806-12. doi: 10.12659/MSM.892769. PMID: 25783889; PMCID: PMC4374486.
71. Bargagli E, Piccioli C, Rosi E, Torricelli E, Turi L, Piccioli E, et al. Pirfenidone and nintedanib in idiopathic pulmonary fibrosis: Real-life experience in an Italian referral centre. *Pulmonology*. 2019;25(3): 149-153. Available from: <https://doi.org/10.1016/j.pulmoe.2018.06.003>
72. Barratt SL, Mulholland S, Jbour KA, Steer H, Gutsche M, Foley N, et al. South-west of England's experience of the safety and tolerability pirfenidone and nintedanib for the treatment of idiopathic pulmonary fibrosis (IPF). *Frontiers in Pharmacology*. 2018;9(1480): 1-10. doi: 10.3389/fphar.2018.01480.
73. Belhassen M, Dalon F, Nolin M, Van Ganse E. Comparative outcomes in patients receiving pirfenidone or nintedanib for idiopathic pulmonary fibrosis. *Respiratory Research*. 2021;22(135): 1-11. Available from: <https://doi.org/10.1186/s12931-021-01714-y>
74. Cerri S, Monari M, Guerrieri A, Donatelli P, Bassi I, Garuti M, et al. Real-life comparison of pirfenidone and nintedanib in patients with idiopathic pulmonary fibrosis: a 24-month assessment. *Respir Med*. 2019;159: 1-6. Available from: <https://doi.org/10.1016/j.rmed.2019.105803>
75. Galli JA, Pandya A, Vega-Olivo M, Dass C, Zhao H, Criner G. Pirfenidone and nintedanib for pulmonary fibrosis in clinical practice: tolerability and adverse drug reactions. *Respirology*. 2017;22: 1171-78. doi: 10.1111/resp.13024
76. Isshiki T, Sakamoto S, Yamasaki A, Shimizu H, Miyoshi S, Nakamura Y, et al. Incidence of acute exacerbation of idiopathic pulmonary fibrosis in patients receiving antifibrotic agents: real-world experience. *Respir Med*. 2021;187: 1-6. Available from: <https://doi.org/10.1016/j.rmed.2021.106551>.
77. Santoleri F, Auriemma L, Spacone A, Marinari S, Esposito F, De Vita F, et al. Adherence, persistence and effectiveness in real life. Multicenter long-term study on the use of pirfenidone and nintedanib in the treatment of idiopathic pulmonary fibrosis. *J Pharm Pract*. 2021;XX(X): 1-6. doi: 10.1177/08971900211008625.

Evidence and Recommendations for the Non-Pharmacologic Interventions for COVID-19

1. Doung-Ngern P, Suphanchaimat R, Panjangampatthana A, et al. Case-Control Study of Use of Personal Protective Measures and Risk for SARS-CoV 2 Infection, Thailand. *Emerg Infect Dis*. 2020; 26: 2607–2616.
2. Abaluck J, Kwnong L, Styczynski A, et al. The Impact of Community Masking on COVID-19: A Cluster-Randomized Trial in Bangladesh [Internet]. 2021 [cited 2021 Sept 17]. Available from: https://www.poverty-action.org/sites/default/files/publications/Mask_RCT____Symptomatic_Seropositivity_083121.pdf
3. Khalil M, Alam M, Arefin M, Chowdhury M, Huq M, Chowdhury J. Role of personal protective measures in prevention of COVID-19 spread among physicians in Bangladesh: a multicenter cross-sectional comparative study. *SN Compr Clin Med*.2(10):1733-9.
4. Chatterjee P, Anand T, Singh K, Rasaily R, Singh R, Das S, et al. Healthcare workers & SARS-CoV-2 infection in India: A case-control investigation in the time of COVID-19. *Indian J Med Res*. 2020;151(5):459-67.
5. Rodriguez-Lopez M, Parra B, Vergara E, Rey L, Salcedo M, Arturo G, et al. A case-control study of factors associated with SARS-CoV-2 infection among healthcare workers in Colombia. *BMC Infect Dis*. 2021;21(1):878.
6. Mohajer M, Panthagani K, Lasco T, Lembecke B, Hemmige V. Association between universal face shield in a quaternary care center and reduction of SARS-COV-2 infections among healthcare personnel and hospitalized patients. *Int J Infect Dis*. 2021;105:252-5.
7. Bhaskar M, Arun S. SARS-CoV-2 infection among community health workers in India before and after use of face shields. *JAMA*. 2020;324(13):1348.

8. Jung S, Yang J-Y, Byeon E-Y, Kim D-G, Lee D-G, Ryoo S, et al. Copper-coated polypropylene filter face mask with SARS-CoV-2 antiviral ability. *Polymers*. 2021;13(9):1367.
9. Hewawaduge C, Senevirathne A, Jawalagatti V, Kim JW, Lee JH. Copper-impregnated three-layer mask efficiently inactivates SARS-CoV-2. *Environmental Research*. 2021;196:110947.
10. Chatterjee R, Bajwa S, Dwivedi D, Kanji R, Ahammed M, Shaw R. COVID-19 Risk Assessment Tool: Dual application of risk communication and risk governance. *Progress in Disaster Science*. 2020 Oct;7:100109.
11. Heinzerling A, Stuckey MJ, Scheuer T, Xu K, Perkins KM, Resseger H, et al. Transmission of COVID-19 to Health Care Personnel During Exposures to a Hospitalized Patient — Solano County, California, February 2020. *MMWR Morb Mortal Wkly Rep*. 2020 Apr 17;69(15):472–6.
12. Khalil MM, Alam MM, Arefin MK, Chowdhury MR, Huq MR, Chowdhury JA, et al. Role of Personal Protective Measures in Prevention of COVID-19 Spread Among Physicians in Bangladesh: a Multicenter Cross-Sectional Comparative Study. *SN Compr Clin Med*. 2020 Aug 28;1–7.
13. Liu M, Cheng S-Z, Xu K-W, Yang Y, Zhu Q-T, Zhang H, et al. Use of personal protective equipment against coronavirus disease 2019 by healthcare professionals in Wuhan, China: cross sectional study. *BMJ*. 2020 Jun 10;369:m2195.
14. Periyasamy P, Ng BH, Ali UK, Rashid ZZ, Kori N. Aerosolized SARS-CoV-2 transmission risk: Surgical or N95 masks? *Infect Control Hosp Epidemiol*. 2020 Sep 15;1–2.
15. Radonovich LJ, Simberkoff MS, Bessesen MT, Brown AC, Cummings DAT, Gaydos CA, et al. N95 respirators vs medical masks for preventing influenza among health care personnel: A randomized clinical trial. *JAMA - J Am Med Assoc*. 2019;322(9):824–33.
16. Wang Q, Huang X, Bai Y, et al. Epidemiological characteristics of COVID-19 in medical staff members of neurosurgery departments in Hubei province: a multicentre descriptive study. *medRxiv* 2020; published online April 24. DOI:10.1101/2020.04.20.20064899 (preprint).
17. Al Mohajer M, Panthagani KM, Lasco T, Lembcke B, Hemmige V. Association between universal face shield in a quaternary care center and reduction of SARS-COV2 infections among healthcare personnel and hospitalized patients. *Int J Infect Dis*. 2021 Apr; 105:252-255. doi: 10.1016/j.ijid.2021.02.060. Epub 2021 Feb 18. PMID: 33610788; PMCID: PMC7891045.
18. Wang X, Pan Z, Cheng Z. Association between 2019-nCoV transmission and N95 respirator use. *J Hosp Infect* 2020; May;105(1):104-105. doi: 10.1016/j.jhin.2020.02.021. Epub 2020 Mar 3. PMID: 32142885; PMCID: PMC7134426.
19. Khalil MM, Alam MM, Arefin MK, Chowdhury MR, Huq MR, Chowdhury JA, Khan AM. Role of Personal Protective Measures in Prevention of COVID-19 Spread Among Physicians in Bangladesh: a Multicenter Cross-Sectional Comparative Study. *SN Compr Clin Med*. 2020 Aug 28;1-7. doi: 10.1007/s42399-020-00471-1. Epub ahead of print. PMID: 32904377; PMCID: PMC7454131.
20. Heinzerling A, Stuckey MJ, Scheuer T, Xu K, Perkins KM, Resseger H, Magill S, Verani JR, Jain S, Acosta M, Epton E. Transmission of COVID-19 to Health Care Personnel During Exposures to a Hospitalized Patient - Solano County, California, February 2020. *MMWR Morb Mortal Wkly Rep*. 2020 Apr 17;69(15):472-476. doi: 10.15585/mmwr.mm6915e5. PMID: 32298249; PMCID: PMC7755059.
21. Berry D, Mainelis G, Fennell D. Effect of an Ionic Air Cleaner on Indoor/Outdoor Particle Ratios in a Residential Environment. *Aerosol Science and Technology*. 2007 Apr;41(3):315–28.
22. Grinshpun SA, Mainelis G, Trunov M, Adhikari A, Reponen T, Willeke K. Evaluation of ionic air purifiers for reducing aerosol exposure in confined indoor spaces. *Indoor Air*. 2005 Aug;15(4):235–45.
23. Hyun J, Lee S-G, Hwang J. Application of corona discharge-generated air ions for filtration of aerosolized virus and inactivation of filtered virus. *Journal of Aerosol Science*. 2017 May;107:31–40.
24. Shi S, Zhu S, Lee ES, Zhao B, Zhu Y. Performance of wearable ionization air cleaners: Ozone emission and particle removal. *Aerosol Science and Technology*. 2016 Mar 3;50(3):211–21.
25. Huang R, Agranovski I, Pyankov O, Grinshpun S. Removal of viable bioaerosol particles with a low-efficiency HVAC filter enhanced by continuous emission of unipolar air ions. *Indoor Air*. 2008 Apr;18(2):106–12.
26. Shimabukuro PMS, Duarte ML, Imoto AM, Atallah ÁN, Franco ESB, Peccin MS, Taminato M. Environmental cleaning to prevent COVID-19 infection. A rapid systematic review. *Sao Paulo Med J*. 2020 Nov-Dec;138(6):505-514. doi: 10.1590/1516-3180.2020.0417.09092020. PMID: 33206913.

27. Hickerson RP, Conneely MJ, Tsutsumi SKH, Wood K, Jackson DN, Ibbotson SH, Eadie E. Minimal, superficial DNA damage in human skin from filtered far-ultraviolet-C (UV-C). *Br J Dermatol*. 2021 Jan 16. doi: 10.1111/bjd.19816. Epub ahead of print. PMID: 33452809.
28. Ramos CCR, Roque JA, Sarmiento DB, Suarez LEG, Sunio JTP, Hilario AL. Use of ultraviolet-C in environmental sterilization in hospitals: A systematic review on efficacy and safety. *International Journal of Health sciences*. 2020 Nov;14(6):52–65.
29. Derraik JGB, Anderson WA, Connelly EA, Anderson YC. Rapid Review of SARS-CoV-1 and SARS-CoV-2 Viability, Susceptibility to Treatment, and the Disinfection and Reuse of PPE, Particularly Filtering Facepiece Respirators. *Int J Environ Res Public Health*. 2020 Aug 22;17(17):6117. doi: 10.3390/ijerph17176117. PMID: 32842655; PMCID: PMC7504573.
30. Buonanno M, Ponnaiya B, Welch D, Stanislauskas M, Randers-Pehrson G, Smilenov L, et al. Germicidal efficacy and mammalian skin safety of 222-nm UV light. *Radiat Res* 2017;187:483–91.
31. Pavia M, Simpser E, Becker M, Mainquist WK, Velez KA. The effect of ultraviolet-C technology on viral infection incidence in a pediatric long-term care facility. *Am J Infect Control* 2018;46:720–2.
32. Qian H, Li Y, Sun H, Nielsen PV, Huang X, Zheng X (2010). Particle removal efficiency of the portable HEPA air cleaner in a simulated hospital ward. *Building Simulation*. 3:215–224.
33. Gedic IE and Alar T (2020). Protective measures undertaken during chest tube thoracostomy in COVID-19 outbreak. *Indian Journal of Thoracic and Cardiovascular Surgery*. <https://doi.org/10.1007/s12055-020-01090-0>
34. J.P. Gibbons, J. Hayes, C.J. Skeritt, J.M. O'Byrne, C.J. Green. Custom solution for personal protective equipment (PPE) in the orthopaedic setting: retrofitting Stryker Flyte T5 PPE system. *Journal of Hospital Infection* 108 (2021) 55e63.
35. Perry JL, Agui JH, Vijayakumar R. Submicron and Nanoparticulate Matter Removal by HEPA-Rated Media Filters and Packed Beds of Granular Materials. USA: National Aeronautics and Space Administration. NASA/TM-2016-218224. May 2016. Downloaded from <https://ntrs.nasa.gov/search.jsp?R=20170005166>.
36. Di Gilio A, Palmisani J, Pulimeno M, Cerino F, Cacace M, Miani A, de Gennaro G. CO2 concentration monitoring inside educational buildings as a strategic tool to reduce the risk of Sars-CoV-2 airborne transmission. *Environ Res* 2021; 202:111560. Available from: <https://doi.org/10.1016/j.envres.2021.111560>.
37. Vassella CC, Koch J, Henzi A, Jordan A, Waeber R, Ianniccone R, Charriere R. From spontaneous to strategic natural window ventilation: Improving indoor air quality in Swiss Schools. *Int J Hyg Environ Health*. 2021. 113736.
38. Lu Y, Li Y, Zhou H, Lin J, Zheng Z, Xu H, et al. Affordable measures to monitor and alarm nosocomial SARS-CoV-2 infection due to poor ventilation. *Indoor Air*. 2021; 00:1–10. doi:10.1111/ina.12899.
39. Du CR, Wang SC, Yu MC, Chiu TF, Wang JY, Chuang PC, et al. Effect of ventilation improvement during a tuberculosis outbreak in under ventilated university buildings. *Indoor Air*. 2020; 30:422–432.
40. Ma J, Qi X, Chen H, Li X, Zhang Z, Wang H, et al. Coronavirus Disease 2019 Patients in Earlier Stages Exhaled Millions of Severe Acute Respiratory Syndrome Coronavirus 2 Per Hour. *Clin Infect Dis*. 2020. doi: 10.1093/cid/ciaa1283.
41. Lednicky, J. A.; Lauzardo, M.; Fan, Z. H.; Jutla, A.; Tilly, T. B.; Gangwar, M.; Usmani, M.; Shankar, S. N.; Mohamed, K.; Eiguren-Fernandez, A.; et al. Viable SARS-CoV-2 in the Air of a Hospital Room with COVID-19 Patients. *Int. J. Infect. Dis*. 2020, 100, 476–482.
42. Cheng VCC, Wong SC, Chan VWM, So SYC, Chen JHK, Yip CCY, et al. Air and environmental sampling for SARS-CoV-2 around hospitalized patients with corona virus disease 2019 (COVID-19). *Infect Control Hosp Epidemiol*. 2020; 41 (11): 1258–1265. doi: 10.1017/ice.2020.282
43. Fischer RJ, Morris DH, van Doremalen N, Sarchette S, Matson MJ, Bushmaker T, et al. Assessment of N95 respirator decontamination and re-use for SARS-CoV-2 [Internet]. *Infectious Diseases (except HIV/AIDS)*; 2020 Apr [cited 2021 Jan 29]. Available from: <http://medrxiv.org/lookup/doi/10.1101/2020.04.11.20062018>
44. Smith JS, Hanseler H, Welle J, Rattray R, Campbell M, Brotherton T, et al. Effect of various decontamination procedures on disposable N95 mask integrity and SARS-CoV-2 infectivity. *J Clin Trans Sci*. 2020 Jun 11;1–5.
45. Kumar A, Kasloff SB, Leung A, Cutts T, Strong JE, Hills K, et al. N95 Mask Decontamination using Standard Hospital Sterilization Technologies [Internet]. *Infectious Diseases (except*

- HIV/AIDS); 2020 Apr [cited 2021 Jan 31]. Available from: <http://medrxiv.org/lookup/doi/10.1101/2020.04.05.20049346>
46. Daeschler SC, Manson N, Joachim K, Chin AWH, Chan K, Chen PZ, et al. Effect of moist heat reprocessing of N95 respirators on SARS-CoV-2 inactivation and respirator function. *CMAJ*. 2020 Oct 13;192(41):E1189–97.
 47. Ibáñez-Cervantes G, Bravata-Alcántara JC, Nájera-Cortés AS, Meneses-Cruz S, Delgado-Balbuena L, Cruz-Cruz C, et al. Disinfection of N95 masks artificially contaminated with SARS-CoV-2 and ESKAPE bacteria using hydrogen peroxide plasma: Impact on the reutilization of disposable devices. *Am J Infect Control*. 2020 Sep;48(9):1037–41.
 48. Simmons SE, Carrion R, Alfson KJ, Staples HM, Jinadatha C, Jarvis WR, et al. Deactivation of SARS-CoV-2 with pulsed-xenon ultraviolet light: Implications for environmental COVID-19 control. *Infect Control Hosp Epidemiol*. 2020 Aug 3;1–4.
 49. Ozog DM, Sexton JZ, Narla S, Pretto-Kernahan CD, Mirabelli C, Lim HW, et al. The effect of ultraviolet C radiation against different N95 respirators inoculated with SARS-CoV-2. *Int J Infect Dis*. 2020 Nov;100:224–9.
 50. Czubyrt MP, Stecy T, Popke E, Aitken R, Jabusch K, Pound R, et al. N95 mask reuse in a major urban hospital: COVID-19 response process and procedure. *Journal of Hospital Infection*. 2020 Oct;106(2):277–82.
 51. Giampalmo M, Pasquesi R, Solinas E. Easy and Accessible Protection against Aerosol Contagion during Airway Management. *Anesthesiology*. 2020;133(3):686–7.
 52. Convissar D, Chang CY, Choi WE, Chang MG, Bittner EA. The Vacuum Assisted Negative Pressure Isolation Hood (VANISH) System: Novel Application of the Stryker Neptune™ Suction Machine to Create COVID-19 Negative Pressure Isolation Environments. *Cureus*. 2020;12(5):e8126.
 53. Price C, Ben-Yakov M, Choi J, Orchanian-Cheff A, Tawadrous D. Barrier enclosure use during aerosol-generating medical procedures: A scoping review. *Am J Emerg Med*. 2021;41:209–18.
 54. Abuhegazy M, Talaat K, Anderoglu O, Poroseva SV. Numerical investigation of aerosol transport in a classroom with relevance to COVID-19. *Phys Fluids (1994)*. 2020;32(10):103311. doi:10.1063/5.0029118
 55. Talaat K, Abuhegazy M, Mahfoze OA, Anderoglu O, Poroseva SV. Simulation of aerosol transmission on a Boeing 737 airplane with intervention measures for COVID-19 mitigation. *Phys Fluids (1994)*. 2021 Mar 1;33(3):033312. doi: 10.1063/5.0044720. Epub 2021 Mar 16. PMID: 33897238; PMCID: PMC8060968.

Evidence and Recommendations for the Vaccines and Prophylactic Interventions for COVID-19

1. PFDA. Emergency Use Authorization (EUA) for COVID-19 Vaccine (Vero Cell), Inactivated manufactured by Sinopharm/China National Pharmaceutical Group/ Beijing Institute of Biological Products (BIBP) Co., Ltd. [Internet]. [cited 2021 Oct 25]. Available from: <https://www.fda.gov.ph/wp-content/uploads/2021/09/EUA-DOH-procured-Sinopharm-Website.pdf>
2. PFDA. Reports of Suspected Adverse Reaction to COVID-19 Vaccines (01 March to 24 October 2021) [Internet]. [cited 2021 Oct 30]. Available from: <https://www.fda.gov.ph/wp-content/uploads/2021/10/Reports-of-suspected-adverse-reaction-to-COVID-19-vaccines-as-of-24-October-2021.pdf>
3. Rogliani P, Chetta A, Cazzola M, Calzetta L. SARS-CoV-2 Neutralizing Antibodies: A Network Meta-Analysis across Vaccines. *Vaccines*. 2021.
4. Al Kaabi N, Zhang Y, Xia S, Yang Y, Al Qahtani MM, Abdulrazzaq N, et al. Effect of 2 Inactivated SARS-CoV-2 Vaccines on Symptomatic COVID-19 Infection in Adults: A Randomized Clinical Trial. *JAMA* [Internet]. 2021;326(1):35–45. Available from: <https://dx.doi.org/10.1001/jama.2021.8565>
5. Feng Y, Chen J, Yao T, Chang Y, Li X, Xing R. Safety and Immunogenicity of Inactivated SARS-CoV-2 Vaccine in High-Risk Occupational Population: a randomized, parallel, controlled clinical trial [Internet]. *medRxiv*. 2021 [cited 2021 Oct 25]. Available from: <https://doi.org/10.1101/2021.08.06.21261696>
6. Xia S, Zhang Y, Wang Y, Wang H, Tang Y, Gao G, et al. Safety and immunogenicity of an inactivated SARS-CoV-2 vaccine, BBIBP-CorV: a randomised, double-blind, placebo-controlled, phase 1/2 trial. *Lancet Infect Dis*. 2021;21:39–51.

7. Xia S, Zhang Y, Wang Y, Wang H, Yang Y, Gao G. Safety and immunogenicity of an inactivated COVID-19 vaccine, BBIBP-CorV, in people younger than 18 years: a randomised, double-blind, controlled, phase 1/2 trial. *Lancet Infect Dis*. 2021.
8. Huang B, Dai L, Wang H, Hu Z, X Y, Tan W. Serum sample neutralisation of BBIBP-CorV and ZF2001 vaccines to SARS-CoV-2 501Y.V2 [Internet]. *The Lancet. Microbe*. 2021 [cited 2021 Oct 25]. Available from: <https://doi.org/10.1016/j.lanmic.2021.100082>
9. Jeewandara C, Aberathna I, Pushpakumara P, Kamaladasa A, Guruge D. Antibody and T cell responses to Sinopharm/BBIBP-CorV in naïve and previously infected individuals in Sri Lanka. *medRxiv*. 2021.
10. Liu T, Dai J, Yang Z, Yu X, Xu Y, Shi X. Inactivated SARS-CoV-2 vaccine does not influence the profile of prothrombotic antibody nor increase the risk of thrombosis in a prospective Chinese cohort [Internet]. *Science Bulletin*. 2021 [cited 2021 Oct 25]. Available from: <https://doi.org/10.1016/j.scib.2021.07.033>
11. Badano M, Sabbione F, Keitelman I, Pereson M, Aloisi N, Colado A. Humoral response to the BBIBP-CorV vaccine over time in healthcare workers with or without exposure to SARS-CoV-2 [Internet]. *medRxiv*. 2021 [cited 2021 Oct 7]. Available from: <https://doi.org/10.1101/2021.10.02.21264432>
12. Ferenci T, Sarkadi B. Virus neutralizing antibody responses after two doses of BBIBP-CorV (Sinopharm, Beijing CNBG) vaccine. *medRxiv*. 2021.
13. Jahromi M, Al Sheikh M. Partial protection of Sinopharm vaccine against SARS COV2 during recent outbreak in Bahrain. *Microbial Pathogenesis*. 2021.
14. WHO. Background document on the inactivated COVID-19 vaccine BIBP developed by China National Biotech Group (CNBG), Sinopharm.
15. Etemadifar M, Abhari A, Nouri H, Sigari A, Daliyeh M, Maracy M. Self-reported safety of the BBIBP-1 CorV (Sinopharm) COVID-19 vaccine among Iranian people with multiple sclerosis [Internet]. *medRxiv*. 2021 [cited 2021 Oct 26]. Available from: <https://doi.org/10.1101/2021.10.17.21265114>
16. Sahraian M, Ghadiri F, Azimi A, Moghadasi A. Adverse events reported by Iranian patients with multiple sclerosis after the first dose of Sinopharm BBIBP-CorV. *Vaccine*. 2021;39:6347–50.
17. Drulovic J, Ivanovic J, Marinovic V, Tamas O, Veselinovic N, Cujic D. Humoral response to SARS-CoV-2 COVID-19 vaccines in patients with multiple sclerosis treated with immune reconstitution therapies [Internet]. Vol. 51, *Multiple Sclerosis and Related Disorders*. 2021 [cited 2021 Oct 28]. Available from: <https://doi.org/10.1016/j.msard.2021.103150>
18. Holt S, Mahmoud S, Ahmed W, Acuna J, Al Madani A, Eltantawy W. An analysis of antibody responses and clinical sequelae of the Sinopharm HB02 COVID19 vaccine in dialysis patients in the United Arab Emirates [Internet]. *Nephrology*. 2021 [cited 2021 Oct 28]. Available from: <https://doi.org/10.1111/nep.13980>
19. Ariamanesh M, Parouhan P, Peyroshabany B, Fazilat-Panah D, Dehghani M, Nabavifard M. Immunogenicity and Safety of the inactivated SARS-CoV-2 vaccine (BBIBP-CorV) in patients with malignancy [Internet]. *medRxiv*. 2021 [cited 2021 Oct 7]. Available from: <https://doi.org/10.1101/2021.09.02.21262760>
20. Abu-Halaweh S, Alqassieh R, Suleiman A, Al-Sabbagh M, AbuHalaweh M, AlKhader D. Qualitative Assessment of Early Adverse Effects of Pfizer–BioNTech and Sinopharm COVID-19 Vaccines by Telephone Interviews. *Vaccines* [Internet]. 2021;9:950. Available from: <https://doi.org/10.3390/vaccines9090950>
21. Saeed B, Al-Shahrabi R, Alhaj S, Alhokhardi Z, Andres A. Side Effects and Perceptions Following Sinopharm COVID-19 Vaccination. *medRxiv*. 2021.
22. Liu Y, Zeng Q, Deng C, Li M, Li L, Liu D. Robust induction of B cell and T cell responses by a third dose of inactivated SARS-CoV-2 vaccine. *medRxiv*. 2021.
23. PFDA. COVID-19 Vaccine (Vero Cell), Inactivated [COVID-19 Vaccine Sinopharm] [Internet]. [cited 2021 Oct 31]. Available from: <https://www.fda.gov.ph/covid-19-vaccine-vero-cell-inactivated-covid-19-vaccine-sinopharm/>
24. WHO-SAGE. Interim recommendations for use of the inactivated COVID-19 vaccine BIBP developed by China National Biotech Group (CNBG), Sinopharm. Interim Guidance. May 7 2021 [Internet]. [cited 2021 Oct 31]. Available from: https://www.who.int/publications/i/item/WHO-2019-nCoV-vaccines-SAGE_recommendation-BIBP-2021
25. Palacios R, Batista A, Albuquerque CS, Patiño E, Santos J, Conde M. Efficacy and Safety of a COVID-19 Inactivated Vaccine in Healthcare Professionals in Brazil: The PROFISCOV Study [Internet]. *SSRN*. 2021 [cited 2021 Apr 12]. Available from: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3822780

26. Tanriover MD, Doganay HL, Akova M, Güner HR, Azap A, Akhan S, et al. Efficacy and safety of an inactivated whole-virion SARS-CoV-2 vaccine (CoronaVac): interim results of a double-blind, randomised, placebo-controlled, phase 3 trial in Turkey. *Lancet* [Internet]. 2021;398(10296):213–22. Available from: [https://dx.doi.org/10.1016/S0140-6736\(21\)01429-X](https://dx.doi.org/10.1016/S0140-6736(21)01429-X)
27. Badan Pengawas Obat dan Makanan (Indonesia Food and Drug Administration). Fact sheet for health care providers emergency use authorization (EUA) of Coronavac [Internet]. [cited 2021 Apr 10]. Available from: <http://pionas.pom.go.id/obat-baru/coronavac-suspensi-injeksi-3-mcg05-ml>
28. Wu Z, Hu Y, Xu M, Z C, Yang W, Jiang Z, et al. Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine (CoronaVac) in healthy adults aged 60 years and older: a randomised, double-blind, placebo-controlled, phase 1/2 clinical trial. *Lancet*. 2021;21:803–12.
29. Bueno S, Abarca K, Gonzales P, Galvez N, Soto G, Duarte L. Interim report: Safety and immunogenicity of an inactivated vaccine against SARS-CoV-2 in healthy chilean adults in a Phase 3 clinical trial [Internet]. *MedRxiv*. April 2021.
30. Han B, Song Y, Li C, Yang W, Ma Q, Jiang Z, et al. Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine (CoronaVac) in healthy children and adolescents: a double-blind, randomised, controlled, phase 1/2 clinical trial. *Lancet Infect Dis* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/b469151f8e2d8574e3f680501b44ed50c82fafdf>
31. Banho C, Sacchetto L, Campos G, Bittar C, Possebon F, Ullmann L. Effects of SARS-CoV-2 P.1 introduction and the impact of COVID-19 vaccination on the epidemiological landscape of São José Do Rio Preto, Brazil. *medRxiv*. 2021.
32. Victora C, Castro M, Gurgenda S, Medeiros A, Franca G, Barros A. Estimating the early impact of vaccination against COVID-19 on deaths among elderly people in Brazil: Analyses of routinely-collected data on vaccine coverage and mortality. *EClinicalMedicine* [Internet]. 2021;38:101036. Available from: <https://doi.org/10.1016/j.eclinm.2021.101036>
33. Jara A, Undurraga E, Gonzalez C, Paredes F, Fontecilla T, Jara G. Effectiveness of an Inactivated SARS-CoV-2 Vaccine in Chile. *N Engl J Med*. 2021;385(10):875–84.
34. Cequeira-Silva T, Oliveira V, Pescarini J, Junior J, Machado T, Florez-Ortiz R, et al. Influence of age on the effectiveness and duration of protection in Vaxzevria and CoronaVac vaccines. *medRxiv*. 2021.
35. Alencar C, Cavalcanti L, de Almeida M, Barbosa P, Cavalcante K, de Melo D, et al. High Effectiveness of SARS-CoV-2 Vaccines in Reducing COVID-19-Related Deaths in over 75-Year-Olds, Ceará State, Brazil. *Trop Med Int Heal*. 2021;6:129.
36. de Faria E, Guedes A, Oliveira M, Moreira M, Maia F, Barboza A, et al. Performance of vaccination with CoronaVac in a cohort of healthcare workers (HCW) - preliminary report. *medRxiv*. 2021.
37. Hitchings M, Ranzani O, Torres M, de Oliveira S, Said R, Borg R. Effectiveness of CoronaVac among healthcare workers in the setting of high SARS-CoV-2 Gamma variant transmission in Manaus, Brazil: A test-negative case-control study [Internet]. *medRxiv*. 2021 [cited 2021 Sep 3]. Available from: <https://www.medrxiv.org/content/10.1101/2021.04.07.21255081v4.full.pdf>
38. Toniasso S, Fernandes F, Jovelevith D, Filho F, Takahasi A, Baldin C, et al. Reduction in COVID-19 prevalence in healthcare workers in a university hospital in southern Brazil after the start of vaccination. *Int J Infect Dis*. 2021;109:283–5.
39. Bayram A, Demirbakan H, Günel Karadeniz P, Erdogan M, Koçer I. Quantitation of antibodies against SARS-CoV-2 spike protein after two doses of CoronaVac in healthcare workers. *J Med Virol* [Internet]. 2021;93(9):5560–7. Available from: <https://dx.doi.org/10.1002/jmv.27098>
40. Muenia N, Garcia-Salum T, Pardo-Roa C, Serrano E, Levican J, Avendano M, et al. Long-lasting neutralizing antibody responses in SARS-CoV-2 seropositive individuals are robustly boosted by immunization with the CoronaVac and BNT162b2 vaccines. *medRxiv*. 2021.
41. Ozdemir H, Tosun S, Coskuner S, Demir S. Assessment of Factors Affecting Inactivated COVID-19 (CORONAVAC) Vaccine Response and Antibody Response in Healthcare Professionals. *Research Square*. 2021.
42. Seyahi E, Bakhdiyarli G, Oztas M, Kuskucu MA, Tok Y, Sut N, et al. Antibody response to inactivated COVID-19 vaccine (CoronaVac) in immune-mediated diseases: a controlled study among hospital workers and elderly. *Rheumatol Int* [Internet]. 2021;41(8):1429–40. Available from: <https://dx.doi.org/10.1007/s00296-021-04910-7>
43. Bichara C, Queiroz M, Amoras E, Vaz G, Vallinoto I, Bichara C. Assessment of anti-SARS-CoV-2 antibodies post-Coronavac vaccination in the Amazon region of Brazil. *Research Square*. 2021.
44. Karacin C, Eren T, Zeynelgil E, Imamoglu G, Altinbas M, Karadag I. Immunogenicity and safety of the CoronaVac vaccine in patients with cancer receiving active systemic therapy [Internet].

- Future Oncology. 2021 [cited 2021 Sep 3]. Available from: <https://www.futuremedicine.com/doi/pdf/10.2217/fon-2021-0597>
45. Yigit M, Ozkaya-Parlakay A, Cosgun Y, Ince Y, Bulut Y, Senel E. Should a third booster dose be scheduled after two doses of CoronaVac? A single-center experience. *J Med Virol.* 2021;(1–4).
 46. Karamese M, Tutuncu E. The effectiveness of inactivated SARS-CoV-2 vaccine (CoronaVac) on antibody response in participants aged 65 years and older. *J Med Virol.* 2021;(1–5).
 47. Organizacion Panamericana de la Salud. Consolidated regional and global information on adverse events following immunization (AEFI) against COVID-19 and other updates. 12 July 2021 [Internet]. [cited 2021 Sep 7]. Available from: <https://covid-19pharmacovigilance.paho.org/img/recursos/60f9ac887614123b2aaa2f56d.pdf>
 48. Organizacion Panamericana de la Salud. Consolidated regional and global information on adverse events following immunization (AEFI) against COVID-19 and other updates. 29 July 2021 [Internet]. [cited 2021 Sep 7]. Available from: <https://covid-19pharmacovigilance.paho.org/img/recursos/611ea8b0a08be304063c51820.pdf>
 49. Food and Health Bureau. The Government of Hong Kong Special Autonomous. Report on Evaluation of Safety, Efficacy and Quality of CoronaVac COVID-19 Vaccine (Vero Cell) Inactivated [Internet]. 2021. [cited 2021 Apr 2]. Available from: https://www.fhb.gov.hk/download/our_work/health/201200/e_evaluation_report_CoronaVac.pdf
 50. Food and Drug Administration Philippines. Reports of Suspected Adverse Reaction to COVID-19 Vaccines (01 March to 5 September).
 51. Durmaz K, Temiz A, Suhal T, Dursun R, Abdelmaksoud A. Allergic and Cutaneous reactions following Inactivated SARS-CoV-2 vaccine (CoronaVac®) in Healthcare workers [Internet]. *Clinical and experimental dermatology.* 2021 [cited 2021 Sep 3]. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/ced.14896>
 52. Akdas E, Ogut B, Erdem O, Oztas M, Ilter N. Cutaneous reactions following CoronaVac COVID-19 vaccination: a case series of six healthcare workers from a single centre [Internet]. *J Eur Acad Dermatol Venereol.* 2021 [cited 2021 Sep 3]. Available from: <https://onlinelibrary.wiley.com/doi/epdf/10.1111/jdv.17592>
 53. Orenay O, Balta I, Yigit D, Eksioğlu M. Systemic drug-related intertriginous and flexural exanthema like eruption after CoronaVac vaccine. *J Eur Acad Dermatol Venereol.* 2021.
 54. Lopes N, Pinilla C, Gerbase A. Erythema multiforme after CoronaVac vaccination. *J Eur Acad Dermatol Venereol.* 2021.
 55. Onsun N, Kaya G, Isik B, Gunes B. A generalized pustular psoriasis flare after CoronaVac COVID-19 vaccination: Case report. *Heal Promot Pr.* 2021;11(2):261–2.
 56. Dirim A, Safak S, Andac B, Garayea N, Demir E, Artan A. Minimal change disease following vaccination with CoronaVac [Internet]. *Clinical Kidney Journal.* 2021 [cited 2021 Sep 3]. Available from: <https://doi.org/10.1093/ckj/sfab123>
 57. Unver S, Hahalu A, Yildirim S. Nephrotic syndrome and acute kidney injury following CoronaVac anti-SARS-CoV-2 vaccine. *Clinical Kidney Journal.* 2021.
 58. Leber H, Sant'Ana L, da Silva N, Raio M, Mazzezo T, Endo C. Acute Thyroiditis and Bilateral Optic Neuritis following SARS-CoV-2 Vaccination with CoronaVac: A Case Report. *Ocular Immunology and inflammation.* 2021.
 59. Wan E, Chui C, Lai F, E C, X L, V Y. Bell's palsy following vaccination with mRNA (BNT162b2) and inactivated (CoronaVac) SARS-CoV-2 vaccines: a case series and nested case-control study. *Lancet Infect Dis.* 2021.
 60. Medeiros-Ribeiro A, Aikawa N, Saad C, Yuki E, Pedrosa T, Gusco S, et al. Immunogenicity and safety of the CoronaVac inactivated vaccine in patients with autoimmune rheumatic diseases: a phase 4 trial [Internet]. *Nature medicine.* 2021 [cited 2021 Sep 3]. Available from: <https://www.nature.com/articles/s41591-021-01469-5.pdf>
 61. Kang M, Yi Y, Li Y, Sun L, Deng A, Hu T, et al. Effectiveness of Inactivated COVID-19 Vaccines Against COVID-19 Pneumonia and Severe Illness Caused by the B.1.617.2 (Delta) Variant: Evidence from an Outbreak in Guangdong, China [Internet]. SSRN. 2021 [cited 1BC Sep 6]. Available from: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3895639
 62. Li X, Huang Y, Jing Q, Zhang C, Qin P, Guan W. Efficacy of inactivated SARS-CoV-2 vaccines against the Delta variant infection in Guangzhou: A test negative case-control real-world study [Internet]. *Emerg Microbes Infect.* 2021 [cited 2021 Sep 3]. Available from: <https://doi.org/10.1080/22221751.2021.1969291>
 63. Ranzani O, Hitchings M, Dorion M, D'Agostini T, de Paula R, de Paula O, et al. Effectiveness of the CoronaVac vaccine in the elderly population during a Gamma variant-associated epidemic of COVID-19 in Brazil: A test-negative case-control study. *medRxiv.* 2021.

64. Hu J, Wei X, Xiang J, Peng P, Xu F, Wu K. Reduced neutralization of SARS-CoV-2 B.1.617 variant by inactivated and RBD-subunit vaccine. *bioRxiv*. 2021.
65. Vacharathit V, Aeiwsakun P, Manopwisedjaroen S, Crisaowakarn C, Laopanupong T. SARS-CoV-2 variants of concern exhibit reduced sensitivity to live-virus neutralization in sera from CoronaVac vaccinees and naturally infected COVID-19 patients [Internet]. *MedRxiv*. 2021 [cited 2021 Aug 18]. Available from: <https://www.medrxiv.org/content/10.1101/2021.07.10.21260232v2.full.pdf>
66. Kara E, Tanir F, Demirhindi H, Mete B, Kibar F, Cetiner S. Humoral immune response in inactivated SARS-CoV-2 vaccine: When should a booster dose be administered? *medRxiv*. 2021.
67. Liao Y, Zhang Y, Zhao H, Pu J, Zhao Z, Li D. Intensified antibody response elicited by boost suggests immune memory in individuals administered two doses of SARS-CoV-2 inactivated vaccine. *Emerg Microbes Infect*. 2021;10(1):1112–5.
68. Jantarabenjakul W, Chantasrisawad N, Puthanakit T, Wacharapluesadee S, Hirankarn N. Short-Term Immune Response After Inactivated 1 SARS-CoV-2 (CoronaVac®, Sinovac) And ChAdOx1 nCoV-19 (Vaxzevria®, Oxford-AstraZeneca) Vaccinations in Thai Health Care Workers. *medRxiv*. 2021.
69. Pan H, Wu Q, Zeng Q, Zeng G, Yang J, Jiang D. Immunogenicity and safety of a third dose, and immune persistence of CoronaVac vaccine in healthy adults aged 18-59 years: interim results from a double-blind, randomized, placebo-controlled phase 2 clinical trial. *MedRxiv*. 2021.
70. Liao Y, Zhang Y, Zhao H, Pu J, Zhao Z, Li D, et al. Intensified antibody response elicited by boost suggests immune memory in individuals administered two doses of SARS-CoV-2 inactivated vaccine. *Emerg Microbes Infect* [Internet]. 2021;10(1):1112–5. Available from: <https://dx.doi.org/10.1080/22221751.2021.1937328>
71. Wang K, Cao Y, Zhou Y, Wu J, Jia Z, Hu Y, et al. A third dose of inactivated vaccine augments the potency, breadth, and duration of anamnestic responses against SARS-CoV-2. *medRxiv*. 2021.
72. Li J, Hou L, Guo X, Jin P, Wu S, Zhu J. Heterologous prime-boost immunization with CoronaVac and Convidecia. *medRxiv*.
73. WHO. Interim recommendations for use of the inactivated COVID-19 vaccine, CoronaVac, developed by Sinovac. 24 May 2021 [Internet]. [cited 2021 Jun 1]. Available from: https://www.who.int/publications/i/item/WHO-2019-nCoV-vaccines-SAGE_recommendation-Sinovac-CoronaVac-2021.1
74. Food and Drug Administration Philippines. Emergency Use Authorization (EUA) for SARS-CoV-2 Vaccine (Vero Cell) Inactivated [Coronavac] Sinovac Life Sciences Co., Ltd [Internet]. [cited 2021 Mar 24]. Available from: <https://www.fda.gov.ph/wp-content/uploads/2021/03/EUA-SINOVAC-WEBSITE-3-1.pdf>
75. Darbar S, Agarwal S, Saha S. COVID19 Vaccine: COVAXIN - India's first indigenous fight against coronavirus (A Review). [Internet]. *Parana Journal of Science and Education*. 2021 [cited 2021 Sep 20]. Available from: <https://www.researchgate.net/publication/352639479>
76. Shresta Y, Venkataraman R, Moktan J, Chitti R, Yadav S. Covid-19 Vaccine Authorized in India- A Mini Review. *SSRN*. 2021.
77. Ella R, Reddy S, Blackwelder W, Potdar V, Yadav P, Sarangi V. Efficacy, safety, and lot to lot immunogenicity of an inactivated SARS-CoV-2 vaccine (BBV152): a, double-blind, randomised, controlled phase 3 trial. *medRxiv*. 2021
78. Singh A. Antibody Response after Second-dose of ChAdOx1 nCov-19 and BBV-152 among Health Care Workers in India: Final Results of Cross-Sectional Coronavirus Vaccine-induced Antibody Titre (COVAT) study. *medRxiv*. 2021.
79. Rajpurohit P, Suva M, Rajpurohit H, Singh Y, Boda P. Retrospective observational survey of adverse events following immunization comparing tolerability of Covishield and Covaxin vaccines in the real world. *J Pharmacol Vigil drug Res*. 2021;2(3):21–6.
80. Arora P, Sardana K, Mathachan S, Malhotra P. Herpes zoster after inactivated COVID-19 vaccine: A cutaneous adverse effect of the vaccine [Internet]. *J Cosmet Dermatol*. 2021 [cited 2021 Sep 10]. Available from: <https://doi.org/10.1111/jocd.14268>
81. Soltanpoor P, Norouzi P. Subacute Thyroiditis Following COVID-19 Vaccination [Internet]. [cited 2021 Sep 20]. Available from: <https://doi.org/10.22541/au.162714686.61648474/v1>
82. Kant R, Dwivedi G, Zaman K, Sahay RR, Sapkal G, Kaushal H, et al. Serendipitous COVID-19 Vaccine-Mix in Uttar Pradesh, India: Safety and Immunogenicity Assessment of a Heterologous Regime. *medRxiv* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/b3f2dc4bbd31ad9d07ff4475cd7257fe571e71c8>
83. Sapkal GN, Yadav PD, Ella R, Deshpande GR, Sahay RR, Gupta N, et al. Inactivated COVID-19

- vaccine BBV152/COVAXIN effectively neutralizes recently emerged B.1.1.7 variant of SARS-CoV-2. *J Travel med* [Internet]. 2021;28(4). Available from: <https://dx.doi.org/10.1093/jtm/taab051>
84. Yadav P, Sapkal G, Ella R, RR S, DA N, DY P, et al. Neutralization of Beta and Delta variant with sera of COVID-19 recovered cases and vaccinees of inactivated COVID-19 vaccine BBV152/Covaxin. *J Travel Med* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/63199af438567149c18bbd9816d71484590ebe62>
 85. Philippine Food and Drug Administration. Emergency Use Authorization (EUA) for Whole Virion, Inactivated Corona Virus Vaccine [Covaxin] [Internet]. [cited 2021 Sep 20]. Available from: <https://www.fda.gov.ph/wp-content/uploads/2021/06/EUA-Bharat-website.pdf>
 86. Heath P, Galiza E, Baxter D, Boffito M, Browne D, Burns F. Safety and Efficacy of NVX-CoV2373 Covid-19 Vaccine. *N Engl J Med* [Internet]. 2021;385:1172–83. Available from: <https://doi.org/10.1056/NEJMoa2107659>
 87. Novavax and Serum Institute of India Receive Emergency Use Authorization for COVID-19 Vaccine in Indonesia [Internet]. [cited 2021 Nov 13]. Available from: <https://ir.novavax.com/2021-11-01-Novavax-and-Serum-Institute-of-India-Receive-Emergency-Use-Authorization-for-COVID-19-Vaccine-in-Indonesia>
 88. Dunkle L, Kotloff K, Gay C, Anez G, Adelglass J, Barrat Hernandez A. Efficacy and Safety of NVX-CoV2373 in Adults in the United States and Mexico [Internet]. *MedRxiv*. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1101/2021.10.05.21264567>
 89. Shinde V, Bhikha S, Hoosain Z, Archary M, Bhorat Q, Fairlie L. Efficacy of NVX-CoV2373 Covid-19 Vaccine against the B.1.351 Variant. *NEJM* [Internet]. 2021;384:1899–909. Available from: <https://doi.org/10.1056/NEJMoa2103055>
 90. Formica N, Mallory R, Albert G, Robinson M, Plested J, Cho I. Evaluation of a SARS-CoV-2 Vaccine NVX-CoV2373 in Younger and Older Adults [Internet]. *MedRxiv*. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1101/2021.02.26.21252482>
 91. Keech C, Albert G, Cho I, Robertson A, Reed P, Neal S, et al. Phase 1–2 Trial of a SARS-CoV-2 Recombinant Spike Protein Nanoparticle Vaccine. *N Engl J Med* [Internet]. 2020;383:2320–32. Available from: <https://doi.org/10.1056/NEJMoa2026920>
 92. Shen X, Tang H, Pahon R, Smith G, Glenn G. Neutralization of SARS-CoV-2 Variants B.1.429 and B.1.351 [Internet]. *NEJM*. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1056/NEJMc2103740>
 93. Higgins J, Green S. *Cochrane handbook for systematic reviews*. Higgins J, Green S, editors. The Cochrane Collaboration; 2011.
 94. Guyatt GH. GRADE: an emerging consensus on rating certainty of evidence and strength of recommendations. *BMJ*. 2008;336:924–6.
 95. Polack F, Thomas S, Kitchin N, Absalon J, Gurtman A, Lockhart S. Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med*. 2020;383(27):2603–15.
 96. Baden L, El Sahly H, Essink B, Kotloff K, Frey S, Novak R. Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N Engl J Med*. 2020.
 97. Voysey M, Clemens S, Madhi S, Weckx L, Folegatti P, Aley P. Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *Lancet*. 2021;397(10269):99–111.
 98. Ramasamy M, Minassian A, Ewer K, Flaxman A, Folegatti P, Owens D. Safety and immunogenicity of ChAdOx1 nCoV-19 vaccine administered in a prime-boost regimen in young and old adults (COV002): a single-blind, randomised, controlled, phase 2/3 trial. *Lancet*. 2020;396(10267):1979–93.
 99. Voysey M, Clemens S, Madhi S, Weckx L. Single-dose administration and the influence of the timing of the booster dose on immunogenicity and efficacy of ChAdOx1 nCoV-19 (AZD1222) vaccine: a pooled analysis of four randomised trials. *Lancet* [Internet]. 2021. Available from: [https://doi.org/10.1016/S0140-6736\(21\)00432-3](https://doi.org/10.1016/S0140-6736(21)00432-3)
 100. Madhi S, Baillie V, Cutland C, Voysey A, Koen L, Fairlie K. Efficacy of the ChAdOx1 nCoV-19 Covid-19 Vaccine against the B.1.351 Variant [Internet]. *New England Journal of Medicine*. 2021. Available from: <https://www.nejm.org/doi/pdf/10.1056/NEJMoa2102214?articleTools=true>
 101. Emary KR, Golubchick T, Aley PK, et al. Efficacy of ChAdOx1 nCoV-19 (AZD1222) vaccine against SARS-CoV-2 variant of concern 202012/01 (B.1.1.7): an exploratory analysis of a randomised controlled trial. *Lancet* [Internet]. 2021;397:1351–62. Available from: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(21\)00628-0/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(21)00628-0/fulltext)

102. Logunov D, Dolzhikova I, Shcheblakov D, Tukhvatulin A, Zubkhova O. Safety and efficacy of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine: an interim analysis of a randomised controlled phase 3 trial in Russia. *Lancet* [Internet]. 2021. Available from: [https://doi.org/10.1016/S0140-6736\(21\)00234-8](https://doi.org/10.1016/S0140-6736(21)00234-8)
103. Sadoff J, Gray G, Vandebosch A, Cardenas V, Shukarev G, Grinsztejn B. Safety and Efficacy of Single-Dose Ad26.COV2.S Vaccine against Covid-19 [Internet]. *N Engl J Med*. 2021 [cited 2021 Apr 23]. Available from: <https://www.nejm.org/doi/pdf/10.1056/NEJMoa2101544?articleTools=true>
104. Palacios R, Batista A, Albuquerque CS, Patiño E, Santos J, Conde M. Efficacy and Safety of a COVID-19 Inactivated Vaccine in Healthcare Professionals in Brazil: The PROFISCOV Study [Internet]. SSRN. 2021 [cited 2021 Apr 12]. Available from: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3822780
105. Bueno S, Abarca K, Gonzales P, Galvez N, Soto G, Duarte L. Interim report: Safety and immunogenicity of an inactivated vaccine against SARS-CoV-2 in healthy chilean adults in a Phase 3 clinical trial [Internet]. *MedRxiv*. 2021. Available from: April 1, 2021
106. Vaccines and Related Biological Products Advisory Committee Meeting December 10, 2020 Meeting FDA Briefing Document Pfizer BioNTech COVID 19 Vaccine. In [cited 2020 Dec 31]. Available from: <https://www.fda.gov/media/144245/download>
107. Vaccines and Related Biological Products Advisory Committee Meeting December 17, 2020 Meeting Presentation - FDA Review of Efficacy and Safety of Moderna COVID-19 Vaccine Emergency Use Authorization Request [Internet]. [cited 2020 Dec 21]. Available from: <https://www.fda.gov/media/144585/download>
108. Vaccines and Related Biological Products Advisory Committee Meeting February 26, 2021. FDA Briefing Document Janssen Ad26.COV2.S Vaccine for the Prevention of COVID-19 [Internet]. 2021. [cited 2021 Mar 5]. Available from: <https://www.fda.gov/media/146217/download>
109. Vaccines and Related Biological Products Advisory Committee Meeting December 10, 2020 Meeting Sponsor Briefing Document Pfizer BioNTech COVID 19 Vaccine [Internet]. [cited 2021 Apr 23]. Available from: <https://www.fda.gov/media/144246/download>
110. European Medicines Agency (EMA). Comirnaty - Assessment Report. 21 December 2020 [Internet]. [cited 2020 Dec 21]. Available from: www.ema.europa.eu/en/medicines/human/EPAR/comirnaty#assessment-history-section
111. European Medicines Agency (EMA) . Comirnaty - Product Information [Internet]. [cited 2020 Dec 21]. Available from: https://www.ema.europa.eu/en/documents/product-information/comirnaty-epar-product-information_en.pdf
112. UK Medicines and Healthcare Products Regulatory Agency (MHRA). Summary of the Public Assessment Report for the Pfizer/BioNTech COVID-19 vaccine [Internet]. [cited 2020 Dec 30]. Available from: <https://www.gov.uk/government/publications/regulatory-approval-of-pfizer-biontech-vaccine-for-covid-19/summary-public-assessment-report-for-pfizerbiontech-covid-19-vaccine>
113. UK Medicines and Healthcare Products Regulatory Authority (MHRA). Conditions of Authorisation for Pfizer/BioNTech COVID-19 vaccine [Internet]. [cited 2020 Dec 30]. Available from: <https://www.gov.uk/government/publications/regulatory-approval-of-pfizer-biontech-vaccine-for-covid-19/conditions-of-authorisation-for-pfizerbiontech-covid-19-vaccine>
114. World Health Organization (WHO). Background document on the mRNA vaccine BNT162b2 (Pfizer-BioNTech) against COVID-19. 14 January 2021 [Internet]. [cited 2021 Apr 23]. Available from: <https://www.who.int/publications/i?healthtopics=b6bd35a3-cf4f-4851-8e80-85cb0068335b>
115. World Health Organization. (2020). mRNA vaccines against COVID-19: Pfizer-BioNTech COVID-19 vaccine BNT162b2: prepared by the Strategic Advisory Group of Experts (SAGE) on immunization working group on COVID-19 vaccines, 22 December 2020 [Internet]. [cited 2021 Jan 4]. Available from: <https://apps.who.int/iris/handle/10665/338096>
116. Vaccines and Related Biological Products Advisory Committee. US Food and Drugs Administration (USFDA). mRNA-1273 Sponsor Briefing Document Vaccines and Related Biological Products Advisory Committee Meeting December 17, 2020. In [cited 2020 Dec 21]. Available from: <https://www.fda.gov/media/144452/download>
117. European Medicines Agency (EMA). Public Assessment Report. COVID-19 Vaccine Moderna. [Internet]. [cited 2021 Feb 9]. Available from: https://www.ema.europa.eu/en/documents/assessment-report/covid-19-vaccine-moderna-epar-public-assessment-report_en.pdf

118. European Medicines Agency (EMA). Moderna Product Information. [Internet]. [cited 2021 Feb 9]. Available from: https://www.ema.europa.eu/en/documents/product-information/covid-19-vaccine-moderna-epar-product-information_en.pdf
119. World Health Organization (WHO) WHO-2019-nCoV-vaccines-SAGE-recommendation-mRNA-1273-background-2021. [Internet]. [cited 2021 Jan 19]. Available from: <https://apps.who.int/iris/handle/10665/339218>
120. World Health Organization (WHO). Interim recommendations for use of the Moderna mRNA-1273 vaccine against COVID-19. Interim guidance 15 January 2021 [Internet]. [cited 2021 Jan 25]. Available from: <https://www.who.int/publications/i/item/interim-recommendations-for-use-of-the-moderna-mrna-1273-vaccine-against-covid-19>
121. UK Medicine and Healthcare Products Regulatory Agency (MHRA). Public Assessment Report Authorisation for Temporary Supply COVID-19 Vaccine AstraZeneca, solution for injection in multidose container COVID-19 Vaccine (ChAdOx1-S [recombinant]) [Internet]. [cited 2021 Jan 6]. Available from: <https://www.gov.uk/government/publications/regulatory-approval-of-covid-19-vaccine-astrazeneca/summary-of-the-public-assessment-report-for-astrazeneca-covid-19-vaccine>
122. European Medicines Agency Committee for Medicinal Products for Human Use (EMA-CHMP). Assessment Report. COVID-19 Vaccine AstraZeneca. January 29 2021 [Internet]. 2021 [cited 2021 Apr 23]. Available from: https://www.ema.europa.eu/en/documents/assessment-report/vaxzevria-previously-covid-19-vaccine-astrazeneca-epar-public-assessment-report_en.pdf
123. European Medicines Agency (EMA). Summary of Product Characteristics. COVID-19 Vaccine (ChAdOx1-S [recombinant]). 2021.
124. World Health Organization. Background document on the AZD1222 vaccine against COVID-19 developed by Oxford University and AstraZeneca. 2021;1222(March). Available from: <https://apps.who.int/iris/bitstream/handle/10665/339882/WHO-2019-nCoV-vaccines-SAGE-recommendation-AZD1222-background-2021.2-eng.pdf?sequence=1&isAllowed=y>
125. World Health Organization (WHO). Interim recommendations for use of the ChAdOx1-S [recombinant] vaccine against COVID-19 (AstraZeneca COVID-19 vaccine AZD1222, SII Covishield, SK Bioscience). Updated 21 Apr 2021.
126. Janssen Biotech, Inc. COVID-19 Vaccine Ad26.COVS.2.S. Sponsor Briefing Document Addendum [Internet]. [cited 2021 Mar 5]. Available from: <https://www.fda.gov/media/146218/download>
127. European Medicines Agency (EMA). COVID-19 vaccine Janssen EPAR product information. [Internet]. [cited 2021 Mar 11]. Available from: https://www.ema.europa.eu/en/documents/product-information/covid-19-vaccine-janssen-epar-product-information_en.pdf
128. European Medicines Agency (EMA). Summary of Product Characteristics. COVID-19 vaccine (Ad26.COVS.2.S [recombinant]). [Internet]. [cited 2021 Mar 24]. Available from: https://www.ema.europa.eu/en/documents/product-information/covid-19-vaccine-janssen-epar-product-information_en.pdf
129. World Health Organization. Background document on the Janssen Ad26.COVS.2.S (COVID-19) vaccine [Internet]. [cited 2021 Apr 6]. Available from: <https://www.who.int/publications/i?healthtopics=b6bd35a3-cf4f-4851-8e80-85cb0068335b>
130. World Health Organization (WHO). Interim recommendations for the use of the Janssen Ad26.COVS.2.S (COVID-19) vaccine. 17 March 2021 [Internet]. [cited 2021 Apr 23]. Available from: <https://www.who.int/publications/i?healthtopics=b6bd35a3-cf4f-4851-8e80-85cb0068335b>
131. Janssen Vaccines & Prevention B.V. A Randomized, Double-blind, Placebo-controlled Phase 3 Study to Assess the Efficacy and Safety of Ad26.COVS.2.S for the Prevention of SARS-CoV-2-mediated COVID-19 in Adults Aged 18 Years and Older. ENSEMBLE. Protocol VAC31 [Internet]. 2020 [cited 2021 Feb 28]. Available from: <https://www.jnj.com/coronavirus/ensemble-1-study-protocol>
132. Food and Health Bureau. The Government of Hong Kong Special Autonomous. Report on Evaluation of Safety, Efficacy and Quality of CoronaVac COVID-19 Vaccine (Vero Cell) Inactivated [Internet]. 2021. [cited 2021 Apr 2]. Available from: https://www.fhb.gov.hk/download/our_work/health/201200/e_evaluation_report_CoronaVac.pdf
133. Food and Health Bureau. The Government of Hong Kong Special Autonomous. COVID-19 Vaccine (Vero Cell) Inactivated Product Information (Detailed). 2021.
134. Badan Pengawas Obat dan Makanan (Indonesia Food and Drug Administration). Fact sheet for health care providers emergency use authorization (EUA) of Coronavac [Internet]. [cited 2021

- Apr 10]. Available from: <http://pionas.pom.go.id/obat-baru/coronavac-suspensi-injeksi-3-mcg05-ml>
135. Palacios R, Patino EG, Pirelli R de O, Conde TRP, Bautista AP. Double-blind, randomized, placebo-controlled phase III clinical trial to evaluate the efficacy and safety of treating healthcare professionals with the adsorbed COVID-19 (inactivated) vaccine manufactures by Sinovac - PROFISCOV: A structured summary of a . *Trials*. 2020;853–5.
 136. Abarca K, Kaergis A, Bueno S, Gonzales P. Efficacy, safety, and immunogenicity of two vaccination schedules of an inactivated vaccine against COVID-19 in Adults (CoronaVac3CL) [Internet]. *Clinicaltrials.Gov*. [cited 2021 Apr 5]. Available from: <https://clinicaltrials.gov/ct2/show/record/NCT04651790>
 137. World Health Organization. Evaluation of COVID-19 vaccine effectiveness. Interim Guidance 17 March 2021. 2021.
 138. Emary K, Golubchik T, Aley P, Ariani C, Angus B. Efficacy of ChAdOx1 nCoV-19 (AZD1222) vaccine against SARS-CoV-2 VOC 20212/01 (B.1.1.7) [Internet]. *SSRN Electronic Journal*. [cited 2021 Mar 8]. Available from: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3779160
 139. Amit S, Regev-Rochay G, Afek A, Kreiss Y, Leshem E. Early rate reductions of SARS-CoV-2 infection and COVID-19 in BNT162b2 vaccine recipients [Internet]. *Lancet*. 2021 [cited 2021 Mar 9]. Available from: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(21\)00448-7/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(21)00448-7/fulltext)
 140. Britton A, KM JS, Edens C, et al. Effectiveness of the Pfizer-BioNTech COVID-19 Vaccine Among Residents of Two Skilled Nursing Facilities Experiencing COVID-19 Outbreaks - Connecticut, December 2020-February 2021. *MMWR Morb Mortal Wkly Rep* [Internet]. 2021;70(11):396–401. Available from: <http://www.epistemonikos.org/documents/4ea6bdb3868dac76a2b6ac17f346c4f0545b9e81>
 141. Cabezas C, Coma E, Mora-Fernandez N, et al. Effects of BNT162b2 mRNA Vaccination on COVID-19 Disease, Hospitalisation and Mortality in Nursing Homes and Healthcare Workers: A Prospective Cohort Study Including 28,594 Nursing Home Residents, 26,238 Nursing Home Staff, and 61,951 Healthcare Workers i. *SSRN* [Internet]. 2021. Available from: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3815682
 142. Chodick G, Tene L, Patalon T, et al. The effectiveness of the first dose of BNT162 b 2 vaccine in reducing SARS-CoV-2 infection 13-24 days after immunization: real-world evidence. *medRxiv* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/1ff25b4cb7b9b006523f8e022be2ea21eccc0c79>
 143. Dagan N, Barda N, Kepten E, et al. BNT162b2 mRNA Covid-19 vaccine in a nationwide mass vaccination setting. *N Engl J Med* [Internet]. 2021. Available from: <https://doi.org/10.1056/NEJMoa2101765>
 144. Guijarro C, Galan IM, Perez-Fernandez E, et al. Dramatic drop of new SARS-CoV-2 infections among health care workers after the first dose of the BNT162b2 mRNA Covid-19 Vaccine. *medRxiv* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/bd8ac8fc1f5bc489ab082fdbfe6150e2e7d6f7bc>
 145. Haas EJ, Angulo FJ, McLaughlin JM, et al. Nationwide Vaccination Campaign with BNT162b2 in Israel Demonstrates High Vaccine Effectiveness and Marked Declines in Incidence of SARS-CoV-2 Infections and COVID-19 Cases, Hospitalizations, and Deaths. *SSRN* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/0ccf39331a250aaef0bc7b095d16cf3c976ec55a>
 146. Hall V, Foulkes S, Saei A, et al. Effectiveness of BNT162b2 mRNA vaccine against infection and COVID-19 vaccine coverage in healthcare workers in England, multicentre prospective cohort study (the SIREN study). *SSRN*. 2021.
 147. Jones N, Rivett L, Seaman S, et al. Single-dose BNT162b2 vaccine protects against asymptomatic SARS-CoV-2 infection. *Elife* [Internet]. 2021;10. Available from: <http://www.epistemonikos.org/documents/2f5773b3ef002738bd289b3fe8401696591231d5>
 148. Monge S, Olmedo C, Alejos B, Lapena M, Sierra MJ, Limia A. Direct and indirect effectiveness of mRNA vaccination against SARS-CoV-2 infection in long-term care facilities in Spain. *medRxiv* [Internet]. 2021. Available from: <https://www.medrxiv.org/content/10.1101/2021.04.08.21255055v1>
 149. Moustsen-Helms IR, Emborg H-D, Nielsen J, et al. Vaccine effectiveness after 1st and 2nd dose of the BNT162b2 mRNA Covid-19 Vaccine in long-term care facility residents and healthcare workers - a Danish cohort study. *medRxiv* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/53c4ee49af74968e43cc779456d74f51c3436dff>

150. Rinott E, Youngster I, Lewis Y. Reduction in COVID-19 Patients Requiring Mechanical Ventilation Following Implementation of a National COVID-19 Vaccination Program - Israel, December 2020-February 2021. *MMWR Morb Mortal Wkly Rep*. 2021. p. 326–8.
151. Zacay G, Shasha D, Bareket R, et al. BNT162b2 Vaccine Effectiveness in Preventing Asymptomatic Infection with SARS-CoV-2 Virus: A Nationwide Historical Cohort Study. SSRN [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/47ca04d87dfe9db54fee471b364c6a07f0d129a0>
152. Bernal JL, Andrews N, Gower C, et al. Early effectiveness of COVID-19 vaccination with BNT162b2 mRNA vaccine and ChAdOx1 adenovirus vector vaccine on symptomatic disease, hospitalisations and mortality in older adults in England [Internet]. *MedRxiv*. 2021 [cited 2021 Mar 9]. Available from: <https://www.medrxiv.org/content/10.1101/2021.03.01.21252652v1>
153. Glampson B, Brittain J, Kaura A, et al. North West London Covid-19 Vaccination Programme: Real-world evidence for Vaccine uptake and effectiveness. *medRxiv* [Internet]. 2021. Available from: <https://www.medrxiv.org/content/10.1101/2021.04.08.21254580v1.full.pdf>
154. Hyams C, Marlow R, Maseko Z, King J, Ward L, Fox K. Assessing the effectiveness of BNT162b2 and ChAdOx1nCoV-19 COVID-19 vaccination in prevention of hospitalisations in elderly and frail adults: a single centre test negative case-control study. SSRN. 2021.
155. Shrotri M, Krutikov M, Palmer T, et al. Vaccine effectiveness of the first dose of ChAdOx1 nCoV-19 and BNT162b2 against SARS-CoV-2 infection in residents of Long-Term Care Facilities (VIVALDI study). *medRxiv* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/5132158a4d58353fd7b74cb64be8f2e6b67efaf7>
156. Vasileiou E, Simpson CR, Robertson C, et al. Effectiveness of first dose of COVID-19 vaccines against hospital admissions in Scotland: national prospective cohort study of 5.4 million people [Internet]. SSRN Electronic Journal. 2021 [cited 2021 Mar 9]. Available from: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3789264
157. Andrejko K, Pry JM, Myers JF, et al. Early evidence of COVID-19 vaccine effectiveness within the general population of California. *medRxiv* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/8040757f20fb5b63a6cd46ac066cb780ae6f8541>
158. Mor V, Gutman R, Yang X, et al. Short-term impact of nursing home SARS-CoV-2 vaccinations on new infections, hospitalizations, and deaths. *J Am Geriatr Soc* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/0d6978157e70cc56490b8161f0baa4d070da14fa>
159. Pawlowski C, Lenehan P, Puranik A, et al. FDA-authorized COVID-19 vaccines are effective per real-world evidence synthesized across a multi-state health system [Internet]. *MedRxiv*. 2021 [cited 2021 Mar 5]. Available from: <https://doi.org/10.1101/2021.02.15.21251623>
160. Roghani A. The Influence of Covid-19 Vaccine on Daily Cases, Hospitalization, and Death Rate in Tennessee: A Case Study in the United States. *medRxiv* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/1eed60fe61d77b4fd5f24cc34a5f3b91d2352773>
161. Thompson M, Burgess J, Naleway A, et al. Interim Estimates of Vaccine Effectiveness of BNT162b2 and mRNA-1273 COVID-19 Vaccines in Preventing SARS-CoV-2 Infection Among Health Care Personnel, First Responders, and Other Essential and Frontline Workers - Eight U.S. Locations, December 2020-March. *MMWR Morb Mortal Wkly Rep* [Internet]. 2021;70(13):495–500. Available from: <http://www.epistemonikos.org/documents/303b1dd4437be9fe0660353ae503f615be077c77>
162. de Faria E, Guedes AR, Oliveira MS, et al. Performance of vaccination with CoronaVac in a cohort of healthcare workers (HCW) - preliminary report. *medRxiv* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/62d22737346dfe78bba89d35c0789a470838c139>
163. Hitchings M, Ranzani OT, Torres MSS, et al. Effectiveness of CoronaVac in the setting of high SARS-CoV-2 P.1 variant transmission in Brazil: A test-negative case-control study. *medRxiv* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/554770e434a57651e335c3b16e4d26e62795ed0a>
164. Luzuriaga JP, Marsico F, Garcia E, et al. Impact of vaccines against COVID-19 on the incidence of new SARS-COV2 infections in health care workers of the Province of Buenos Aires. 2021. Available from: <https://preprints.scielo.org/index.php/scielo/preprint/view/2068/3406>
165. Rivkees S, Roberson S, Blackmore C. Outcomes of COVID-19 Vaccination Efforts in Florida from December 14, 2020 to March 15, 2021 on Older Individuals. *medRxiv* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/24ced7ae24bdd6585ffa245e35617596eb55618a>
166. Dutta S, Kaur RJ, Charan J, et al. Serious adverse events reported from the COVID-19 vaccines: A descriptive study based on WHO database. *medRxiv* [Internet]. 2021. Available from: <https://www.medrxiv.org/content/10.1101/2021.03.23.21253433v1.full.pdf>

167. Mathioudakis A, Ghrew M, Ustianowski A, et al. Self-Reported Real-World Safety and Reactogenicity of COVID-19 Vaccines: A Vaccine Recipient Survey. *Life* (Basel, Switzerland) [Internet]. 2021;11(3). Available from: <https://www.mdpi.com/2075-1729/11/3/249>
168. McMahon D, Amerson E, Rosenbach M, et al. Cutaneous Reactions Reported after Moderna and Pfizer COVID-19 Vaccination: A Registry-Based Study of 414 Cases. *J Am Acad Dermatol* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/640a98a9e5c75d5faf21bba63eb92d115d353023>
169. Mortazavi S. Coronavirus Disease (COVID-19) Vaccination Associated Axillary Adenopathy: Imaging Findings and Follow-Up Recommendations in 23 Women. *AJR Am J Roentgenol* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/47429f8c5d15be0f99b3845912d38d662c15035a>
170. Shimabukuro T, Cole M, JR S. Reports of Anaphylaxis After Receipt of mRNA COVID-19 Vaccines in the US-December 14, 2020-January 18, 2021. *JAMA* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/9e146cd9b0b0e17214584b3f68682e41454a6034>
171. Davido B, Mascitti H, Fortier-Beaulieu M, Jaffal K, P de T. "Blue toes" following vaccination with the BNT162b2 mRNA COVID-19 vaccine. *J Travel Med* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/1066cc91c713cb890ee0b0a2e403b915a1141fbf>
172. García-Grimshaw M, Ceballos-Liceaga SE, Hernández-Vanegas LE, et al. Systemic and Neurologic Adverse Events Among 704,003 First-Dose Recipients of the Pfizer-Biontech (BNT162b2) mRNA COVID-19 Vaccine in Mexico. *SSRN* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/9623d8f06c619da3b9b56f54d7a4adecebbe09e0>
173. Gee J, Marquez P, Su J, Calvert G, Liu R, Myers T, et al. First Month of COVID-19 Vaccine Safety Monitoring — United States, December 14, 2020–January 13, 2021 [Internet]. Vol. 70, Morbidity Mortality Weekly Report. 2021; 70: 283-8 [cited 2021 Mar 22]. Available from: <https://www.cdc.gov/mmwr/volumes/70/wr/pdfs/mm7008e3-H.pdf>
174. Kim S, Wi Y, Yun S, et al. Adverse Events in Healthcare Workers after the First Dose of ChAdOx1 nCoV-19 or BNT162b2 mRNA COVID-19 Vaccination: a Single Center Experience. *J Korean Med Sci* [Internet]. 2021;36(14):e107. Available from: <http://www.epistemonikos.org/documents/48c6eb652f4eefc5627c4e815737f997ea13c49b>
175. Monin-Aldama L, Laing A, Munoz-Ruiz M, et al. Interim results of the safety and immune-efficacy of 1 versus 2 doses of COVID-19 vaccine BNT162b2 for cancer patients in the context of the UK vaccine priority guidelines. *medRxiv* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/310daffd483993141c2e7d5307c8d8daff474069>
176. Riad A, Klugarova J, Koscik M, et al. Prevalence of covid-19 vaccine side effects among healthcare workers in the Czech Republic. *J Clin Med* [Internet]. 2021;10(7). Available from: <http://www.epistemonikos.org/documents/2e6ace3dc0630f1bd3244d375cb81ac390de91f5>
177. Waheed S, Bayas A, Hindi F, Rizvi Z, PS E. Neurological Complications of COVID-19: Guillain-Barre Syndrome Following Pfizer COVID-19 Vaccine. *Cureus* [Internet]. 2021;13(2):e13426. Available from: <http://www.epistemonikos.org/documents/8de11dd4bd7bf0a9cb2bb8ed619bf773c1774f1>
178. CDC. Local Reactions, Systemic Reactions, Adverse Events, and Serious Adverse Events: Moderna COVID-19 Vaccine. *CDC Website* [Internet]. 2021. Available from: <http://www.epistemonikos.org/documents/7c8223e8474331b949d0763a091fde114a9ca943>
179. COVID-19 Response Team FDA. Allergic Reactions Including Anaphylaxis After Receipt of the First Dose of Moderna COVID-19 Vaccine - United States, December 21, 2020-January 10, 2021. *MMWR Morb Mortal Wkly Rep* [Internet]. 2021;70(4):125–9. Available from: <http://www.epistemonikos.org/documents/a06b60b5e02770c92d14b0f86fe85e708af3e13f>
180. Julian J, Mathern D, Fernando D. Idiopathic Thrombocytopenic Purpura and the Moderna Covid-19 Vaccine – A Case Report. *Ann Emerg Med* [Internet]. 2021. Available from: [https://www.annemergmed.com/article/S0196-0644\(21\)00122-0/pdf](https://www.annemergmed.com/article/S0196-0644(21)00122-0/pdf)
181. Kadali R, Janagama R, Peruru S, et al. Adverse effects of COVID-19 mRNA-1273 vaccine: A randomized, cross-sectional study on healthcare workers with detailed self-reported symptoms. *J Med Virol* [Internet]. 2021. Available from: <https://onlinelibrary.wiley.com/doi/epdf/10.1002/jmv.26996?src=getftr>
182. Greinacher A, Thiele T, Warkentin T, Weisser K, Kryle P, Eichinger S. Thrombotic Thrombocytopenia after ChAdOx1 nCov-19 Vaccination. *N Engl J Med*. 2021.
183. Kaur U, Ojha B, Pathak BK, Singh A, Giri KR, Singh A, et al. A prospective observational safety study on ChAdOx1 nCoV- 19 corona virus vaccine (recombinant) use in healthcare workers- first results from India. *medRxiv* [Internet]. 2021. Available from: <https://www.medrxiv.org/content/10.1101/2021.04.03.21254823v1.full.pdf>

184. Mehta P, Mangion S, Benger M, Stanton B, Czuprynska J, Arya R, et al. Cerebral venous sinus thrombosis and thrombocytopenia after COVID-19 vaccination - a report of two UK cases. *Brain Behav Immun* [Internet]. 2021. Available from: <https://www.sciencedirect.com/science/article/pii/S088915912100163X?pes=vor>
185. Schultz NH, Sorvoll IH, Michelsen AE, Munthe LA, Lund-Johansen F, Aanodt A-H. Thrombosis and Thrombocytopenia after ChAdOx1 nCoV-19 Vaccination [Internet]. *N Engl J Med*. 2021 [cited 2021 Apr 9]. Available from: <https://www.nejm.org/doi/full/10.1056/NEJMoa2104882>
186. Wolf M, Bazner H, Luz B, Niehaus L, Bhogal P, Henkes H. Thrombocytopenia and intracranial venous sinus thrombosis after "covid-19 vaccine astrazeneca" exposure. *J Clin Med* [Internet]. 2021;10(8). Available from: <https://www.mdpi.com/2077-0383/10/8/1599>
187. Riad A, Sağıroğlu D, Üstün B, Attia S, Klugar M. Prevalence and Risk Factors of CoronaVac Side Effects: An Independent Cross-Sectional Study Among Healthcare Workers in Turkey [Internet]. SSRN. 2021 [cited 2021 Apr 24]. Available from: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3820571
188. CDC COVID-19 Response Team. Allergic Reactions Including Anaphylaxis After Receipt of the First Dose of Pfizer-BioNTech COVID-19 Vaccine - United States, December 14-23, 2021 [Internet]. *MMWR Morb Mortal Wkly Rep*. 2021 [cited 2021 Apr]. -- incomplete date? Available from: ?
189. CDC COVID-19 Response Team. Allergic Reactions Including Anaphylaxis After Receiving First Dose of Moderna COVID-19 Vaccine - United States December 21, 2020 - January 10, 2021. *MMWR Morb Mortal Wkly Rep*. 2021.
190. CDC. Cases of cerebral venous sinus thrombosis with thrombocytopenia after receipt of the Johnson & Johnson COVID-19 Vaccine. Date?
191. European Medicines Agency (EMA). COVID-19 vaccine safety update. Comirnaty BioNTech Manufacturing GmbH. 28 Jan 2021. [Internet]. [cited 2021 Mar 24]. Available from: https://www.ema.europa.eu/en/documents/covid-19-vaccine-safety-update/covid-19-vaccine-safety-update-comirnaty-january-2021_en.pdf
192. European Medicines Agency. COVID-19 Vaccine AstraZeneca : benefits still outweigh the risks despite possible link to rare blood clots with low blood platelets. [Internet]. [cited 2021 Mar 24]. Available from: <https://www.ema.europa.eu/en/news/covid-19-vaccine-astrazeneca-benefits-still-outweigh-risks-despite-possible-link-rare-blood-clots>
193. European Medicines Association. AstraZeneca's COVID-19 vaccine: EMA finds possible link to very rare cases of unusual blood clots with low blood platelets. Date?
194. EMA Pharmacovigilance Risk Assessment Committee. Signal assessment report on embolic and thrombotic events (SMQ) with COVID-19 Vaccine (ChAdOx1-S [recombinant]) – COVID-19 Vaccine AstraZeneca (Other viral vaccines) [Internet]. [cited 2021 Apr 11]. Available from: https://www.ema.europa.eu/en/documents/prac-recommendation/signal-assessment-report-embolic-thrombotic-events-smq-covid-19-vaccine-chadox1-s-recombinant-covid_en.pdf
195. UK MHRA. Coronavirus vaccine - weekly summary of Yellow Card reporting. Updated 18 Mar 2021 [Internet]. [cited 2021 Mar 24]. Available from: <https://www.gov.uk/government/publications/coronavirus-covid-19-vaccine-adverse-reactions/coronavirus-vaccine-summary-of-yellow-card-reporting>
196. UK MHRA. Coronavirus vaccine - weekly summary of YellowCard reporting. Updated 8 April 2021.
197. Food and Drug Administration Philippines. Reports of Adverse Reaction to COVID-19 Vaccines (01 to 11 April 2021) [Internet]. [cited 2021 Apr 15]. Available from: <https://www.fda.gov.ph/wp-content/uploads/2021/04/Reports-of-Suspected-Adverse-Reaction-to-COVID-19-Vaccines-as-of-11-April-2021-ver-3.pdf>
198. Instituto de Salud Publica de Chile. Reacciones anafilacticas asociadas a la administracion de las vaculas SARS-CoV-2 - Evaluacion de reportes de esavi y recomendaciones para las vacunas Pfizer-BioNTech y Coronavac [Internet]. Nota Informativa De Farmacovigilancia. 2021 [cited 2021 Apr 12]. Available from: <https://www.ispch.cl/wp-content/uploads/2021/03/INF-FARMACOVIGILANCIA.pdf>
199. CDC. FDA and CDC Lift Recommended Pause on Johnson & Johnson (Janssen) COVID-19 Vaccine Use Following Thorough Safety Review [Internet]. [cited 2021 Apr 23]. Available from: <https://www.fda.gov/news-events/press-announcements/fda-and-cdc-lift-recommended-pause-johnson-johnson-janssen-covid-19-vaccine-use-following-thorough>
200. WHO issues its first emergency use validation for a covid-19 vaccine. WHO news release Dec 31 [Internet]. [cited 2021 Jan 4]. Available from: <https://www.who.int/news/item/31-12-2020-who->

- issues-its-first-emergency-use-validation-for-a-covid-19-vaccine-and-emphasizes-need-for-equitable-global-access
201. Food and Drug Administration Philippines. FDA Philippines Grants Emergency Use Authorization to Pfizer-BioNTech COVID19 Vaccine [Internet]. [cited 2021 Feb 25]. Available from: <https://www.fda.gov.ph/wp-content/uploads/2021/02/EUA-Pfizer-Website.pdf>,
 202. Food and Drug Administration Philippines. Emergency Use Authorization (EUA) for COVID-19 Vaccine (ChAdOx1- S[recombinant]) (COVID-19 Vaccine AstraZeneca). Date?
 203. Sputnik V General Information [Internet]. [cited 2021 Feb 24]. Available from: <https://sputnikvaccine.com/about-vaccine/>
 204. Food and Drug Administration Philippines. Emergency Use Authorization (EUA) for Gamaleya National Center of Epidemiology and Microbiology Sputnik V Gam-COVID-Vac COVID-19 Vaccine [Internet]. [cited 2021 Mar 24]. Available from: <https://www.fda.gov.ph/wp-content/uploads/2021/03/EUA-Gamaleya-Website.pdf>
 205. European Medicines Agency (EMA). EMA recommends COVID-19 Vaccine Janssen for authorisation in the EU. [Internet]. [cited 2021 Mar 11]. Available from: <https://www.ema.europa.eu/en/news/ema-recommends-covid-19-vaccine-janssen-authorisation-eu>
 206. WHO adds Janssen vaccine to list of safe and effective emergency tools against COVID-19. 2021.
 207. Food and Drug Administration Philippines. Emergency Use Authorization (EUA) for COVID-19 Vaccine (Ad26.COVS-2 S[recombinant]), Grand River Aseptic Manufacturing, Michigan, USA Site [Internet]. 2021. [cited 2021 Apr 23]. Available from: <https://www.fda.gov.ph/wp-content/uploads/2021/04/EUA-Janssen-Website.pdf>
 208. Food and Drug Administration Philippines. Emergency Use Authorization (EUA) for SARS-CoV-2 Vaccine (Vero Cell) Inactivated [Coronavac] Sinovac Life Sciences Co., Ltd [Internet]. [cited 2021 Mar 24]. Available from: <https://www.fda.gov.ph/wp-content/uploads/2021/03/EUA-SINOVAC-WEBSITE-3-1.pdf>
 209. Tukhvatulin A, Dolzhikova I, Shcheblakov D, Zubkhova O, Dzharullaeva A. An open, non-randomised, 1/2 phase study on the safety, tolerability, and immunogenicity of single dose "Sputnik Light" Vaccine for Prevention of Coronavirus Infection in healthy adults [Internet]. SSRN. 2021 [cited 2021 Sep 3]. Available from: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3886430
 210. Gonzales S, Olszevicki S, Salazar M, Calabria A, Regairaz L, Marin L. Effectiveness of the first component of Gam-COVID-Vac (Sputnik V) on reduction of SARS-CoV-2 confirmed infections, hospitalisations and mortality in patients aged 60-79: a retrospective cohort study in Argentina [Internet]. Vol. 40, *EClinicalMedicine*. 2021 [cited 2021 Oct 30]. Available from: <https://doi.org/10.1016/j.eclinm.2021.101126>
 211. Komissarov A, Dolzhikova I, Efimov G, Logunov D, Mityaeva O, Molodtsov I. Boosting of the SARS-CoV-2-specific immune response after vaccination with single-dose Sputnik Light vaccine [Internet]. medRxiv. 2021 [cited 2021 Oct 30]. Available from: <https://doi.org/10.1101/2021.10.26.21265531>
 212. Saeed U, Uppal S, Pichara Z, Khan A, Rasheed A, Waheed A. Evaluation of SARS-CoV-2 spike antibody levels among Sputnik V first dose vaccinated people in Pakistan: formulation of national anti-COVID-19 mass vaccination strategy [Internet]. Research Square. 2021 [cited 2021 Oct 30]. Available from: <https://doi.org/10.21203/rs.3.rs-480406/v1>
 213. Dolzhikova I, Guschin V, Shcheblyakov D, Tsybin A, Shchetinin A, Pchtoyi A. One-shot immunization with Sputnik Light (the first component of Sputnik V vaccine) is effective against SARS-CoV-2 Delta variant: efficacy data on the use of the vaccine in civil circulation in Moscow [Internet]. medRxiv. 2021 [cited 2021 Oct 30]. Available from: <https://doi.org/10.1101/2021.10.08.21264715>
 214. Food and Drug Administration Philippines. Emergency Use Authorization (EUA) for Gamaleya National Center of Epidemiology and Microbiology Sputnik Light COVID-19 Vaccine [Internet]. [cited 2021 Sep 5]. Available from: <https://www.fda.gov.ph/wp-content/uploads/2021/08/Sputnik-Light-COVID-19-Vaccine.pdf>
 215. COVID19 Vaccine Tracker. Gamaleya: Sputnik Light [Internet]. [cited 2021 Nov 1]. Available from:
 216. Bar-on Y, Goldberg Y, Mandel M, Bodenheimer O, Freedman L, Kalkstein N, et al. BNT162b2 vaccine booster dose protection: A nationwide study from Israel [Internet]. medRxiv. 2021 [cited 2021 Sep 7]. Available from: <https://doi.org/10.1101/2021.08.27.21262679>
 217. Patalon T, Gazit S, Pitzer V, Prunas O, Warren J, Weinberger D. Short Term Reduction in the Odds of Testing Positive for SARS-CoV-2; a Comparison Between Two Doses and Three doses

- of the BNT162b2 Vaccine. medRxiv. 2021.
218. Barda N, Dagan N, Cohen C, MA H, Lipsitch M, IS K, et al. Effectiveness of a third dose of the BNT162b2 mRNA COVID-19 vaccine for preventing severe outcomes in Israel: an observational study. *Lancet* (London, England) [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/f25fb54ff2b477f94228bbe78487fa5fb81044c7>
 219. VRBPAC. September 17, 2021 Meeting Briefing Document - FDA [Internet]. 2021 [cited 2021 Sep 18]. Available from: <https://www.fda.gov/media/152176/download>
 220. Levine-Tiefenbrun M, Yelin I, Alapi H, Katz R, Herzel E, Kuint J. Viral loads of Delta-variant SARS-CoV-2 breakthrough infections after vaccination and booster with BNT162b2 [Internet]. *Nature medicine*. 2021 [cited 2021 Nov 5]. Available from: <https://doi.org/10.1038/s41591-021-01575-4>
 221. Mofaz M, Yechezkel M, Guan G, Brandeau ML, Patalon T, Gazit S, et al. Self-reported and physiological reactions to the third BNT162b2 mRNA COVID-19 (booster) vaccine dose. medRxiv [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/355d4df501c99de8986b72206b3fd258d7faf343>
 222. Niesen MJM, Pawlowski C, O'Horo JC, Challenger DW, Silvert E, Donadio G, et al. Three doses of COVID-19 mRNA vaccination are safe based on adverse events reported in electronic health records. medRxiv [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/bccdc1587a9c07cc64ef3d633f16691c01c65f0c>
 223. Chu L, Montefiori D, Huang W, Nestorova B, Chang Y, Carfi A, et al. Immune Memory Response After a Booster Injection of mRNA-1273 for Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2). medRxiv [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/2ff5ae9b7641ed768b86f07f6b8beb9536aa0bb0>
 224. Wu K, Choi A, Koch M, Ma L, Hill A, Nunna N, et al. Preliminary Analysis of Safety and Immunogenicity of a SARS-CoV-2 Variant Vaccine Booster. medRxiv [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/9a44dd424b51cff2df60d966e00323fd3231408d>
 225. Choi A, Koch M, Wu K, Chu L, Ma L, Hill A, et al. Safety and immunogenicity of SARS-CoV-2 variant mRNA vaccine boosters in healthy adults: an interim analysis. *Nat Med* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/b116c6d1f58f76e7d18c11aed426f5071527ee54>
 226. Flaxman A, Marchevsky NG, Jenkin D, Aboagye J, Angus B, Belij-Rammerstorfer S. Tolerability and immunogenicity after a late second dose or a third dose of ChAdOx1 nCoV-19 (AZD1222) [Internet]. SSRN. 2021 [cited 2021 Aug 8]. Available from: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3873839
 227. Sadoff J, Le Gars M, Cardenas V, Shukarev G, Vaissiere N. Durability of antibody responses elicited by a single dose of Ad26.COV2.S and substantial increase following late boosting [Internet]. medRxiv. 2021 [cited 2021 Sep 7]. Available from: <https://doi.org/10.1101/2021.08.25.21262569>
 228. Pan H, Wu Q, Zeng Q, Zeng G, Yang J, Jiang D. Immunogenicity and safety of a third dose, and immune persistence of CoronaVac vaccine in healthy adults aged 18-59 years: interim results from a double-blind, randomized, placebo-controlled phase 2 clinical trial. medRxiv. 2021.
 229. Li J, Hou L, Guo X, Jin P, Wu S, Zhu J. Heterologous prime-boost immunization with CoronaVac and Convidecia [Internet]. medRxiv. 2021 [cited 2021 Sep 7]. Available from: <https://doi.org/10.1101/2021.09.03.21263062>
 230. Li M, Yang J, Wang L, Wu Q, Wu Z, Zheng W, et al. A booster dose is immunogenic and will be needed for older adults who have completed two doses vaccination with CoronaVac: a randomised, double-blind, placebo-controlled, phase 1/2 clinical trial [Internet]. medRxiv. 2021 [cited 2021 Sep 15]. Available from: <https://doi.org/10.1101/2021.08.03.21261544>
 231. Wang K, Cao Y, Zhou Y, Wu J, Jia Z, Hu Y. A third dose of inactivated vaccine augments the potency, breadth, and duration of anamnestic responses against SARS-CoV-2. medRxiv. 2021.
 232. Mok CKP, Cohen CA, Cheng SMS, Chen C, Kwok K-O, Yiu K, et al. Comparison of the Immunogenicity of BNT162b2 and CoronaVac COVID-19 Vaccines in Hong Kong: An Observational Cohort Study. SSRN [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/c37accfceac5098a5e1308164fffd2fdbc0e94fd>
 233. Barin B, Kasap U, Selçuk F, Volkan E, Uluckan O. Longitudinal Comparison of SARS-CoV-2 Anti-Spike RBD IgG antibody Responses After CoronaVac, BNT162b2, ChAdOx1 nCoV-19 Vaccines and Evaluation of a Single Booster Dose of BNT162b2 or CoronaVac After a Primary CoronaVac Regimen. SSRN [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/3e23dc8b436b4bb7973c2089d8d20f931d6faffd>
 234. Yue L, Xie T, Yang T, Zhou J, Chen H, Zhu H, et al. A third booster dose may be necessary to mitigate neutralizing antibody fading after inoculation with two doses of an inactivated SARS-

- CoV-2 vaccine. J Med Virol [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/6eeb8b98e94a24ccda3173f7e7e5d1c5106d2fda>
235. Liao Y, Zhang Y, Zhao H, Pu J, Zhao Z, Li D, et al. Intensified antibody response elicited by boost suggests immune memory in individuals administered two doses of SARS-CoV-2 inactivated vaccine. *Emerg Microbes Infect* [Internet]. 2021;10(1):1112–5. Available from: <https://dx.doi.org/10.1080/22221751.2021.1937328>
 236. Yue L, Zhou J, Zhou Y, Yang X, Xie T, Yang M, et al. Antibody response elicited by a third boost dose of inactivated SARS-CoV-2 vaccine can neutralize SARS-CoV-2 variants of concern. *Emerg Microbes Infect* [Internet]. 2021;1–9. Available from: <http://www.epistemonikos.org/documents/44649a65700cd4060c0a3ee38e7655c046b9239d>
 237. Li Y, Fang X, Pei R, Fan R, Chen S, Zeng P, et al. Immunogenicity and safety of the homogenous booster shot of a recombinant fusion protein vaccine (V-01) against COVID-19 in healthy adult participants primed with a two-dose regimen. *medRxiv* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/fa9ea94583e966283063d87453097e4d48385c60>
 238. Atmar R, Lyke K, Deming M, Jackson L, Branche A, ElSahly H, et al. Heterologous SARS-CoV-2 Booster Vaccinations – Preliminary Report [Internet]. *MedRxiv*. 2021 [cited 2021 Nov 5]. Available from: <https://doi.org/10.1101/2021.10.10.21264827>
 239. Iketani S, Liu L, Nair M, Mohri H, Wang M, Huang Y, et al. A third COVID-19 vaccine shot markedly boosts neutralizing antibody potency and breadth [Internet]. *MedRxiv*. 2021 [cited 2021 Nov 5]. Available from: <https://doi.org/10.1101/2021.08.11.21261670>
 240. Hoque A, Rahman M, Imam H, Nahar N, Chowdhury FUH. Third dose vaccine With BNT162b2 and its response on Long COVID after Breakthrough infections. *medRxiv* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/fa8cdeb1c29ff1ac00821be30937544c03a5db1d>
 241. Huat N, Lim J, Gill U, de Alwis R, Tan N, Toh J, et al. Differential immunogenicity of homologous versus heterologous boost in Ad26.COVS vaccine recipients [Internet]. *MedRxiv*. 2021 [cited 2021 Nov 12]. Available from: <https://doi.org/10.1101/2021.10.14.21264981>
 242. Sester M, Klemis V, Venhorst A, Halmans L, Schmidt T, Greiß F, et al. Immunogenicity and reactogenicity of heterologous Ad26.COVS and BNT162b2 vaccination. *ResearchSquare* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/375ea1006cb8923d3fe23a63566b9ddc0534350e>
 243. Moghnieh R, Mekdashi R, El-Hassan S, Abdallah D, Jisr T, Bader M, et al. Immunogenicity and reactogenicity of BNT162b2 booster in BBIP-CorV-vaccinated individuals compared with homologous BNT162b2 vaccination: Results of a pilot prospective cohort study from Lebanon. *Vaccine* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/88e5b508c75c86a8869254c1cebe4d4c79b1012f>
 244. Romero-Ibarguengoitia ME, Rivera-Salinas D, Hernandez-Ruiz YG, Armendariz-Vazquez AG, Gonzalez-Cantu A, Barco-Flores IA, et al. Effect of the third dose of BNT162b2 vaccine in quantitative SARS-CoV-2 spike 1-2 IgG antibody titers in healthcare workers. *medRxiv* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/89e8a516684f9ff9ce614d5aa3405d3f37523787>
 245. Liu Y, Zeng Q, Deng C, Li M, Li L, Liu D. Robust induction of B cell and T cell responses by a third dose of inactivated SARS-CoV-2 vaccine. *medRxiv*. 2021.
 246. Sablerolles R, Rietdijk W, Goorhuis B, Postma D, Visser L, Geers D, et al. Immunogenicity and reactogenicity of booster vaccinations after Ad26.COVS priming. *medRxiv* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/8c24d7d3bbe697a739407310aada45cf2920164f>
 247. Patamatamkul S, Thammawat S, Buranrat B. Induction of robust neutralizing antibodies against the COVID-19 Delta variant with ChAdOx1 nCoV-19 or BNT162b2 as a booster following a primary vaccination series with CoronaVac [Internet]. *MedRxiv*. 2021 [cited 2021 Nov 5]. Available from: <https://doi.org/10.1101/2021.09.25.21264099>
 248. Yorsaeng R, Vichaiwattana P, Klinfueng S, Wongsrisang L, Sudhinaraset N, Vongpunsawad S, et al. Immune response elicited from heterologous SARS-CoV-2 vaccination: Sinovac (CoronaVac) followed by AstraZeneca (Vaxzevria) [Internet]. *medRxiv*. 2021 [cited 2021 Sep 30]. Available from: <https://doi.org/10.1101/2021.09.01.21262955>
 249. Singhatiraj E, Rongpirul K, Jongkaewwattana A, Hirankarn N. Intradermal ChAdOx1 Vaccine Following Two CoronaVac Shots: A Case Report. *Vaccines* [Internet]. 2021;9:990. Available from: <https://doi.org/10.3390/v9090990>
 250. Chavarot N, Morel A, Leruez-Ville M, Villain E, Divard G, Burger C. Weak antibody response to 3 doses of mRNA vaccine in kidney transplant recipients treated with belatacept [Internet]. *Am J*

- Transplant. 2021. Available from: <https://doi.org/10.1111/AJT.16814>
251. Kamar N, Abravanel F, Marion O, Couat C, Izopet J, Del Bello A. Three Doses of an mRNA Covid-19 Vaccine in Solid-Organ Transplant Recipients. *N Engl J Med* [Internet]. 2021;385(7):661–2. Available from: <https://dx.doi.org/10.1056/NEJMc2108861>
 252. Bensouna I, Caudwell V, Kubab S, Acquaviva S, Pardon A, Vittoz N, et al. SARS-CoV-2 Antibody Response After a Third Dose of the BNT162b2 Vaccine in Patients Receiving Maintenance Hemodialysis or Peritoneal Dialysis. *Am J Kidney Dis* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/5712a0aecee60f7b858952f03a3d9d14210f5d77>
 253. Ducloux D, Colladant M, Chabannes M, Yannaraki M, Courivaud C. Humoral response after 3 doses of the BNT162b2 mRNA COVID-19 vaccine in patients on hemodialysis. *Kidney Int* [Internet]. 2021;100(3):702–4. Available from: <https://dx.doi.org/10.1016/j.kint.2021.06.025>
 254. Peled Y, Ram E, Lavee J, Segev A, Matezki S, Wieder-Finesod A, et al. Third dose of the BNT162b2 vaccine in heart transplant recipients: Immunogenicity and clinical experience. *J Heart Lung Transplant* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/17effe940234e7478143e97a9c253071efcd4a>
 255. Del Bello A, Abravanel F, Marion O, Couat C, Esposito L, Lavayssiere L. Efficiency of a boost with a third dose of anti-SARS-CoV-2 messenger RNA-based vaccines in solid organ transplant recipients [Internet]. *Am J Transplant*. 2021 [cited 2021 Sep 7]. Available from: <https://doi.org/10.1111/AJT.16775>
 256. Masset C, Kerleau C, Garanddeau C, Ville S, Cantarovich D, Hourmant M. A third injection of BNT162b2 mRNA Covid-19 vaccine in kidney transplant recipients improves the humoral immune response [Internet]. *Kidney Int*. 2021 [cited 2021 Sep 7]. Available from: <https://doi.org/10.1016/j.kint.2021.08.017>
 257. David S, Shamir-Stein N, Gez S, Lerner U, Rahamim-Cohen D, AE Z. Reactogenicity of a third BNT162b2 mRNA COVID-19 vaccine among immunocompromised individuals and seniors - A nationwide survey. *Clin Immunol* [Internet]. 2021;108860. Available from: <http://www.epistemonikos.org/documents/35d4ff071a7e8669998cfe12b3ea46ac7cc20f6b>
 258. Hall V, Ferreira V, Ju T, Jerullo M, Majchzrak-Kita B. Randomized Trial of a Third Dose of mRNA-1273 Vaccine in Transplant Recipients. *N Engl J Med*. 2021.
 259. Benotmane I, Gautier G, Perrin P, Olagne J, Cognard N, Fafi-Kremer S, et al. Antibody Response After a Third Dose of the mRNA-1273 SARS-CoV-2 Vaccine in Kidney Transplant Recipients With Minimal Serologic Response to 2 Doses [Internet]. *JAMA*. 2021 [cited 2021 Aug 28]. Available from: <https://doi.org/10.1001/jama.2021.12339>
 260. Greenberger L, Saltzman L, Senefeld J, Johnson P, DeGennaro L, Nichols G. Anti-spike antibody response to SARS-CoV-2 booster vaccination in patients with B cell-derived hematologic malignancies [Internet]. *Cancer Cell*. 2021 [cited 2021 Nov 5]. Available from: <https://doi.org/10.1016/j.ccell.2021.09.001>
 261. Werbel W, Boyarsky B, Ou M, Massie A, Tobian A, Garonzik-Wang J. Safety and Immunogenicity of a Third Dose of SARS-CoV-2 Vaccine in Solid Organ Transplant Recipients: A Case Series [Internet]. *Ann Intern Med*. 2021 [cited 2021 Sep 7]. Available from: <https://doi.org/10.7326/L21-0282>
 262. Lyski Z, Kim S, Lee D, Sampson D, Raue H, Raghunathan V, et al. Immunogenicity of Pfizer mRNA COVID-19 vaccination followed by J&J adenovirus COVID-19 vaccination in two CLL patients [Internet]. *MedRxiv*. 2021 [cited 2021 Nov 5]. Available from: <https://doi.org/10.1101/2021.09.02.21262146>
 263. Baker M, Mallayosyula V, Davis M, Boyd S, Nadeau K, Robinson W. Effective Viral Vector SARS-CoV-2 Booster Vaccination in a Patient with Rheumatoid Arthritis after Initial Ineffective mRNA Vaccine Response [Internet]. *Arthritis and Rheumatology*. 2021 [cited 2021 Oct 22]. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/art.41978>
 264. Bonelli M, Mrak D, Tobudic S, Sieghart D, Koblishke M, Mandl P, et al. Additional heterologous versus homologous booster vaccination in immunosuppressed patients without SARS-CoV-2 antibody seroconversion after primary mRNA vaccination: a randomized controlled trial. *medRxiv* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/29ed89ed3746bcc6fc81dd6c365069b08e2a2f30>
 265. Caillard S, Thauinat O, Benotmane I, Masset C, Blancho G. Antibody response to a fourth mRNA Covid-19 vaccine boost in weak responder kidney transplant recipients [Internet]. *MedRxiv*. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1101/2021.09.03.21262691>
 266. WHO. Interim statement on COVID-19 vaccine booster doses. Update 4 October 2021 [Internet]. [cited 2021 Oct 7]. Available from: <https://www.who.int/news/item/04-10-2021-interim-statement-on-booster-doses-for-covid-19-vaccination>

267. US CDC. COVID-19 Vaccine Booster Shots. November 9. 2021 [Internet]. [cited 2021 Nov 16]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/booster-shot.html#print>
268. UK JCVI. Update to JCVI advice on booster vaccination in adults, 15 November 2021 [Internet]. [cited 2021 Nov 16]. Available from: <https://www.gov.uk/government/publications/covid-19-booster-vaccine-programme-for-winter-2021-to-2022-jcvi-statement-november-2021/update-to-jcvi-advice-on-booster-vaccination-in-adults-15-november-2021>
269. Wise J. European countries suspend use of Oxford-AstraZeneca vaccine after reports of blood clots. *BMJ*. 2021.
270. Pottegård A, Lund L, Karlstad O. Arterial events, venous thromboembolism, thrombocytopenia, and bleeding after vaccination with Oxford-AstraZeneca ChAdOx1-S in Denmark and Norway: population based cohort study. *BMJ*. 2021;373(n1114).
271. Supasa P, Zhou D, Dejnirattisai W. Reduced neutralization of SARS-CoV-2 B.1.1.7 variant by convalescent and vaccine sera. *Cell*. 2021;184(8):2201-2211.e7.
272. Planas D, Bruel T, Grzelak L, Guivel-Benhassine F, Staropoli I, Porrot F, et al. Sensitivity of infectious SARS-CoV-2 B.1.1.7 and B.1.351 variants to neutralizing antibodies. *Nat Med* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/62eb9c0b52797f914bd1a5bcbe230733f8488088>
273. Hoffmann M, Arora P, Groß R, Seidel A, BF H, AS H, et al. SARS-CoV-2 variants B.1.351 and P.1 escape from neutralizing antibodies. *Cell* [Internet]. 2021; Available from: <https://www.cell.com/action/showPdf?pii=S0092-8674%2821%2900367-6>
274. Emary KR, Golubchick T, Aley PK, Ariani C, Angus B. Efficacy of ChAdOx1 nCoV-19 (AZD1222) Vaccine Against SARS-CoV-2 VOC 202012/01 (B.1.1.7) [Internet]. SSRN. 2021 [cited 2021 Mar 27]. Available from: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3779160
275. Liu X, Shaw R, Stuart A, Greenland M, Aley P, Andrews N. Safety and immunogenicity of heterologous versus homologous prime-boost schedules with an adenoviral vectored and mRNA COVID-19 vaccine (Com-COV): a single-blind, randomised, non-inferiority trial [Internet]. *Lancet*. 2021 [cited 2021 Aug 16]. Available from: [https://doi.org/10.1016/%0DS0140-6736\(21\)01694-9](https://doi.org/10.1016/%0DS0140-6736(21)01694-9)
276. Powell A, Power L, Westrop S, McOwat K, Campbell H, Simmons R. Real-world data shows increased reactogenicity in adults after heterologous compared to homologous prime-boost COVID-19 vaccination, March–June 2021, England. *Euro Surveill*. 2021;26(28):2100634.
277. Normark J, Vikstrom L, Gwon Y, Persson I, Edin A. Heterologous ChAdOx1 nCoV-19 and mRNA-1273 Vaccination [Internet]. *N Engl J Med*. 2021 [cited 2021 Sep 30]. Available from: <https://doi.org/10.1056/NEJMc2110716>
278. Kant R, Dwivedi G, Zaman K, Sahay R, Sapkal G, Kaushal H, et al. tous COVID-19 Vaccine-Mix in Uttar Pradesh, India: Safety and Immunogenicity Assessment of a Heterologous Regime [Internet]. *medRxiv*. 2021 [cited 2021 Sep 3]. Available from: <https://doi.org/10.1101/2021.08.06.21261716>
279. Yorsaeng R, Vichaiwattana P, Klinfueng S, Wongsrisang L, Sudhinaraset N, Vongpunswad S, et al. Immune response elicited from heterologous SARS-CoV-1 2 vaccination: Sinovac (CoronaVac) followed by AstraZeneca (Vaxzevria) [Internet]. *medRxiv*. 2021 [cited 2021 Sep 30]. Available from: <https://doi.org/10.1101/2021.09.01.21262955>
280. Li J, Hou L, Guo X, Jin P, Wu S, Zhu J. Heterologous prime-boost immunization with CoronaVac and Convidecia [Internet]. *medRxiv*. [cited 2021 Sep 7]. Available from: <https://doi.org/10.1101/2021.09.03.21263062>
281. Schmidt T, Klemis V, Schub D, Mihm J, Hielscher F. Immunogenicity and reactogenicity of heterologous ChAdOx1 nCoV-19/mRNA vaccination [Internet]. *Nature medicine*. 2021 [cited 2021 Aug 18]. Available from: <https://doi.org/10.1038/s41591-021-01464-w>
282. Schmidt T, Klemis V, Schub D, Schneitler S, Reichert M, Wilkens H, et al. Cellular immunity predominates over humoral immunity after homologous and heterologous mRNA and vector-based COVID-19 vaccine regimens in solid organ transplant recipients. *Am J Transpl*. 2021;00:1–13.
283. Gram M, Nielsen J, Schelde A, Nielsen K, Mousten-Helms I, Sorensen A. Vaccine effectiveness when combining the ChAdOx1 vaccine as the first dose with an mRNA COVID-19 vaccine as the second dose [Internet]. *medRxiv*. 2021 [cited 2021 Sep 6]. Available from: <https://doi.org/10.1101/2021.07.26.21261130>
284. Borobia A, Carcas A, Olmeda M, Castano L, Bertran M, Perez J. Immunogenicity and reactogenicity of BNT162b2 booster in ChAdOx1-S-primed participants (CombiVacS): a multicentre, open-label, randomised, controlled, phase 2 trial. *Lancet*. 2021;398:121–30.
285. Rose R, Neumann F, Grobe O, Lorentz T, Fickenscher H, Krumbholz A. Heterologous immunisation with vector vaccine as prime followed by mRNA vaccine as boost leads to humoral

- immune response against SARS-CoV-2, which is comparable to that according to a homologous mRNA vaccination scheme [Internet]. medRxiv. 2021 [cited 2021 Aug 12]. Available from: <https://www.medrxiv.org/content/10.1101/2021.07.09.21260251v1.full.pdf>
286. Tenbusch M, Schumacher, Vogel E, Priller P, Held H, Steininger P. Heterologous prime-boost vaccination with ChAdOx1 nCoV-19 and BNT162b2 mRNA [Internet]. medRxiv. 2021 [cited 2021 Aug 16]. Available from: <https://www.medrxiv.org/content/10.1101/2021.07.03.21258887v1.full.pdf>
 287. Benning L, Tollner M, Hidmark A, Schaier M, Nussbag C, Kalble F. Heterologous ChAdOx1 nCoV-19/BNT162b2 Prime-Boost Vaccination Induces Strong Humoral Responses among Health Care Workers. *Vaccines*. 2021;9:857.
 288. Dimeglio C, Herin F, Da-Silva I, Jouglu I, Pradere C, Porcheron M. No Title Heterologous ChAdOx1-S/BNT162b2 vaccination: neutralizing antibody response to SARS-CoV-2 [Internet]. *Clin Inf Dis*. 2021 [cited 2021 Aug 28]. Available from: <https://doi.org/10.1093/cid/ciab705>
 289. Hammerschmidt S, Bosnjak B, Bernhardt G, Friedrichsen M, Ravens I, Dopfer-Jablonka A, et al. Neutralization of the SARS-CoV-2 Delta variant after heterologous and homologous BNT162b2 or ChAdOx1 nCoV-19 vaccination [Internet]. *Cellular & Molecular Immunology*. 2021 [cited 2021 Sep 6]. Available from: <https://doi.org/10.1038/s41423-021-00755-z>
 290. Hillus D, Schwarz T, Tober-Lau P, Hastor H, Thibeault C, Kasper S. Safety, reactogenicity, and immunogenicity of homologous and heterologous prime-boost immunisation with ChAdOx1-nCoV19 and BNT162b2: a prospective cohort study [Internet]. *Lancet*. 2021 [cited 2021 Sep 30]. Available from: [https://doi.org/10.1016/S2213-2600\(21\)00357-X](https://doi.org/10.1016/S2213-2600(21)00357-X)
 291. Vinh D, Gouin J, Cruz-Santiago D, Canac-Marquis M, Bernier S, Bobuef F. Real-world serologic responses to Extended-interval and Heterologous COVID-19 mRNA vaccination in Frail Elderly - Interim report from a prospective observational cohort study [Internet]. medRxiv. 2021 [cited 2021 Sep 30]. Available from: <https://doi.org/10.1101/2021.09.16.21263704>
 292. Behrens G, Cossmann A, Stankov M, Nehlmeier I, Kempf A, Hoffmann M, et al. SARS-CoV-2 delta variant neutralisation after heterologous ChAdOx1-S/BNT162b2 vaccination [Internet]. *Lancet*. 2021 [cited 2021 Sep 3]. Available from: <https://doi.org/10.1038/%0Ds41591-021-01449-9>
 293. Barros-Martins J, SI H, Cossmann A, Odak I, MV S, G MR, et al. Immune responses against SARS-CoV-2 variants after heterologous and homologous ChAdOx1 nCoV-19/BNT162b2 vaccination. *Nat Med* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/eac69a37499fb6a67b7a8c9bdf94ac2286cf7266>
 294. GroB R, Zanon M, Seidel A, Conzelmann C, Gilg A. Heterologous ChAdOx1 nCoV-19 and BNT162b2 prime-boost vaccination elicits potent neutralizing antibody responses and T cell reactivity [Internet]. medRxiv. 2021 [cited 2021 Aug 9]. Available from: <https://www.medrxiv.org/content/10.1101/2021.05.30.21257971v2.full.pdf>
 295. Shaw RH, Stuart A, Greenland M, Liu X, Nguyen Van-Tam JS, Snape MD. Heterologous prime-boost COVID-19 vaccination: initial reactogenicity data. *Lancet* [Internet]. 2021;397(10289):2043–6. Available from: [https://dx.doi.org/10.1016/S0140-6736\(21\)01115-6](https://dx.doi.org/10.1016/S0140-6736(21)01115-6)
 296. WHO. Interim statement on heterologous priming for covid-19 vaccines. Aug 10 2021 [Internet]. [cited 2021 Oct 10]. Available from: <https://www.who.int/news/item/10-08-2021-interim-statement-on-heterologous-priming-for-covid-19-vaccines>
 297. Australian Technical Advisory Group on Immunization. COVID-19 Vaccination ATAGI Clinical Guidance on COVID-19 Vaccine in Australia in 2021. [Internet]. [cited 2021 Sep 30]. Available from: <https://www.health.gov.au/sites/default/files/documents/2021/10/covid-19-vaccination-atagi-clinical-guidance-on-covid-19-vaccine-in-australia-in-2021.pdf>
 298. World Health Organization. Tracking SARS-CoV-2 Variants.
 299. Lopez Bernal J, Andrews N, Gower C, Gallagher E, Simmons R. Effectiveness of Covid-19 Vaccines against the B.1.617.2 (Delta) Variant. *N Engl J Med*. 2021.
 300. Mlcochova P, Kemp S, Dhar M, Papa G, Meng B, Mishra S. SARS-CoV-2 B.1.617.2 Delta variant replication, sensitivity to neutralising antibodies and vaccine breakthrough [Internet]. bioRxiv. 2021 [cited 2021 Aug 19]. Available from: <https://www.biorxiv.org/content/10.1101/2021.05.08.443253v5.full.pdf>
 301. Thiruvengadam R, Awasthi A, Medigeshi G, Bhattacharya S, Mani S. Cellular Immune Responses are Preserved and may Contribute to ChAdOx1 nCoV-19 Vaccine Effectiveness Against Infection due to SARS-CoV-2 B.1.617.2 Delta Variant Despite Reduced Virus Neutralisation. SSRN2. 2021.
 302. Chia P, Ong S, Chiew C, Ang L, Chavatte J. Virological and serological kinetics of SARS-CoV-2 Delta variant vaccine breakthrough infections: a multi-center cohort study [Internet]. medRxiv.

- 2021 [cited 2021 Aug 18]. Available from: <https://www.medrxiv.org/content/10.1101/2021.07.28.21261295v1.full.pdf>
303. Nasreen S, Chung H, He S, Brown K, Gubbay J, Buchan S. Effectiveness of COVID-19 vaccines against variants of concern in Ontario, Canada [Internet]. MedRxiv. 2021 [cited 2021 Aug 18]. Available from: <https://www.medrxiv.org/content/10.1101/2021.06.28.21259420v2.full.pdf>
 304. Sheikh A, McMenamin J, Taylor B, Robertson C. SARS-CoV-2 Delta VOC in Scotland: demographics, risk of hospital admission, and vaccine effectiveness [Internet]. Lancet. 2021 [cited 2021 Aug 9]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8201647/pdf/main.pdf>
 305. Stowe J, Andrews N, Gower C, Gallagher E, Utsi L, Simmons R. Effectiveness of COVID-19 vaccines against hospital admission with the Delta (B.1.617.2) variant [Internet]. [cited 2021 Aug 19]. Available from: https://media.tghn.org/articles/Effectiveness_of_COVID-19_vaccines_against_hospital_admission_with_the_Delta_B._G6gnnqJ.pdf
 306. Buxvoort K, Sy L. Effectiveness of mRNA -1237 against Delta, Mu, and other emerging variants. medRxiv. 2021.
 307. Tang P, Hasan C, Yassine H, Benslimane F, Al Khatib H. BNT162b2 and mRNA-1273 COVID-19 vaccine effectiveness against the Delta (B.1.617.2) variant in Qatar. MedRxiv. 2021.
 308. Tartof S, Slezak J, Fischer H, Hong V, Ackerson B. Effectiveness of mRNA BNT162b2 COVID-19 vaccine up to 6 months in a large integrated health system in the USA: a retrospective cohort study [Internet]. Lancet. 2021 [cited 2021 Oct 5]. Available from: [https://doi.org/10.1016/%0AS0140-6736\(21\)02183-8](https://doi.org/10.1016/%0AS0140-6736(21)02183-8)
 309. Ella R, Reddy S, Blackwelder W, Potdar V, Yadav P, Sarangi V. Efficacy, safety, and lot to lot immunogenicity of an inactivated SARS-CoV-2 vaccine (BBV152): a, double-blind, randomised, controlled phase 3 trial. medRxiv. 2021.
 310. Li X, Huang Y, Jing Q, Zhang C, Qin P, Guan W. Efficacy of inactivated SARS-CoV-2 vaccines against the Delta variant infection in Guangzhou: A test negative case-control real-world study [Internet]. Emerg Microbes Infect. 2021 [cited 2021 Sep 3]. Available from: <https://doi.org/10.1080/22221751.2021.1969291>
 311. Kang M, Yi Y, Li Y, Sun L, Deng A, Hu T, et al. Effectiveness of Inactivated COVID-19 Vaccines Against COVID-19 Pneumonia and Severe Illness Caused by the B.1.617.2 (Delta) Variant: Evidence from an Outbreak in Guangdong, China [Internet]. SSRN. 2021 [cited 2021 Sep 6]. Available from: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3895639
 312. Nanduri S, Pilishvili T, Derado G, Soe MM, Dollard P, Wu H, et al. Effectiveness of Pfizer-BioNTech and Moderna Vaccines in Preventing SARS-CoV-2 Infection Among Nursing Home Residents Before and During Widespread Circulation of the SARS-CoV-2 B.1.617.2 (Delta) Variant - National Healthcare Safety Network, March 1-August. MMWR Morb Mortal Wkly Rep [Internet]. 2021;70(34):1163–6. Available from: <https://dx.doi.org/10.15585/mmwr.mm7034e3>
 313. Davis C, Logan N, Tyson G, Orton R, Harvey W, Hughney J. Reduced neutralisation of the Delta (B.1.617.2) SARS-CoV-2 variant of concern following vaccination. MedRxiv. 2021.
 314. Lustig Y, Zuckerman N, Nemet I, Atari N, Kliker L, Regev-Yochay G. Neutralising capacity against Delta (B.1.617.2) and other variants of concern following Comirnaty (BNT162b2, BioNTech/Pfizer) vaccination in health care workers, Israel [Internet]. Euro surveillance. [cited 2021 Aug 18]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8326656/>
 315. Planas D, Veyer D, Baidaliuk A, Staropoli I, Guivel-Benhassine F. Reduced sensitivity of SARS-CoV-2 variant Delta to antibody neutralization. Nature [Internet]. 2021;596:276–80. Available from: <https://www.nature.com/articles/s41586-021-03777-9>
 316. Edara V, Pinsky B, Suthar M, Lai L, Davis-Gardner M, Floyd K. Infection and Vaccine-Induced Neutralizing-Antibody Responses to the SARS-CoV-2 B.1.617 Variants. N Engl J Med. 2021.
 317. Liu C, Ginn H, Dejnattis W, Ren J, Stuart D, Screaton G. Reduced neutralization of SARS-CoV-2 B.1.617 by vaccine and convalescent serum [Internet]. Cell. 2021 [cited 2021 Aug 19]. p. 4220–36. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8218332/>
 318. Choi A, Koch M, Wu K, Dixon G, Oestreicher J, Legault H. Serum Neutralizing Activity of mRNA-1273 against SARS-CoV-2 Variants [Internet]. bioRxiv. 2021 [cited 2021 Aug 18]. Available from: <https://doi.org/10.1101/2021.06.28.449914>
 319. Pegu A, O'Connell S, Schmidt S, O'Dell S, Talana C. Durability of mRNA-1273 vaccine-induced antibodies against SARS-CoV-2 variants [Internet]. Science. 2021 [cited 2021 Aug 18]. Available from: <https://science.sciencemag.org/content/early/2021/08/11/science.abj4176/tab-pdf>
 320. Sapkal G, Yadav P, Sahay R, Dishpande G, Gupta N. Neutralization of Delta variant with sera of Covishield vaccinees and COVID-19 recovered vaccinated individuals. bioRxiv. 2021.

321. Jongeneelen M, Kaszas K, Veldman D, Huizingh J, van den Vlught R. Ad26.COVS.2 elicited neutralizing activity against Delta and other SARS-CoV-2 variants of concern [Internet]. bioRxiv. 2021 [cited 2021 Aug 19]. Available from: <https://www.biorxiv.org/content/10.1101/2021.07.01.450707v1.full.pdf>
322. Hu J, Wei X, Xiang J, Peng P, Xu F, Wu K. Reduced neutralization of SARS-CoV-2 B.1.617 variant by inactivated and RBD-subunit vaccine. bioRxiv. 2021.
323. Vacharathit V, Aeiwsakun P, Manopwisedjaroen S, Crisaowakarn C, Laopanupong T. SARS-CoV-2 variants of concern exhibit reduced sensitivity to live-virus neutralization in sera from CoronaVac vaccinees and naturally infected COVID-19 patients [Internet]. MedRxiv. 2021 [cited 2021 Aug 18]. Available from: <https://www.medrxiv.org/content/10.1101/2021.07.10.21260232v2.full.pdf>
324. Keeton R, Richardson S, Moyo-Gwete T, Harmanus T, Tincho M, Benede N. Prior infection with SARS-CoV-2 boosts and broadens Ad26.COVS.2 immunogenicity in a variant dependent manner [Internet]. medRxiv. 2021 [cited 2021 Sep 3]. Available from: <https://doi.org/10.1101/2021.07.24.21261037>
325. Gushchin V, Dolzhikova I, Schcheginin A, Odintsova A, Siniavin A, Nikiforova M. against Variants of Concern (VOC: B.1.1.7, B.1.351, P.1, B.1.617.2, B.1.617.3) and Moscow Endemic SARS-CoV-2 Variants. *Vaccines*. 2021;9:779.
326. Noori M, Nejadghaderi S, Arshi S, Carson-Chahhoud K, Ansarin K. Potency of BNT162b2 and mRNA-1273 vaccine-induced neutralizing antibodies against severe acute respiratory syndrome-CoV-2 variants of concern: A systematic review of in vitro studies. *Rev Med Virol*. 2021. p. e2277.
327. US CDC. COVID-19 Information for Pediatric Healthcare Providers [Internet]. [cited 2021 Sep 17]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/pediatric-hcp.html>.
328. American Academy of Pediatrics. Children and COVID-19: State-Level Data Report [Internet]. [cited 2021 Sep 17]. Available from: <https://www.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/children-and-covid-19-state-level-data-report>
329. Wallace M, Woodworth K, Gargano J, Scobie H, Blain A, Bouliia D. The Advisory Committee on Immunization Practices' interim recommendation for use of Pfizer-BioNTech COVID-19 vaccine in adolescents aged 12–15 Years — United States, May 2021. *MMWR Morb Mortal Wkly Rep*. 2021;70(20):749–52.
330. Lv M, Luo X, Shen Q, Lei R, Liu X, Liu E. Safety, immunogenicity, and efficacy of COVID-19 vaccines in children and adolescents: a systematic review. *Vaccines* [Internet]. 2021 [cited 2021 Sep 17];9(10):1102. Available from: <https://doi.org/10.3390/vaccines9101102>.
331. Frenck R, Klein N, Kitchin N, Gurtman A, Absalon J, Lockhart S, et al. Safety, Immunogenicity, and Efficacy of the BNT162b2 Covid-19 Vaccine in Adolescents. *N Engl J Med* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/a5820eb1dc7757b8f2376b5857ab253f3a8c5a62>
332. Ali K, Berman G, Zhou H, Deng W, Faughnan V, Coronado-Voges M, et al. Evaluation of mRNA-1273 SARS-CoV-2 Vaccine in Adolescents. *N Engl J Med* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/71d40df3cbe241937f0ed4b8f0bec9ce89c2ecbd>
333. Han B, Song Y, Li C, Yang W, Ma Q, Jiang Z, et al. Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine (CoronaVac) in healthy children and adolescents: a double-blind, randomised, controlled, phase 1/2 clinical trial. *Lancet Infect Dis* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/b469151f6e2d8574e3f680501b44ed50c82fafdf>
334. Bickel S, Harris C, Huxol H, R M. COVID-19 Vaccination Outcomes at a Pediatric Long-Term Care Facility. *Pediatr Infect Dis J*. 2021;40(7):e281-3.
335. Revon-Riviere G, Ninove L, Min V, Rome A, Coze C, Verschuur A, et al. The BNT162b2 mRNA COVID-19 vaccine in adolescents and young adults with cancer: A monocentric experience. *Eur J Cancer* [Internet]. 2021;154:30–4. Available from: <http://www.epistemonikos.org/documents/8bf6a5f88d3f26a084be7236c5cb9184f670b69a>
336. Delahoy M, Ujamaa D, Whitaker M, A O, Anglin O, Burns E. Hospitalizations associated with

- COVID-19 among children and adolescents — COVID-NET, 14 States, March 1, 2020–August 14, 2021. *MMWR Morb Mortal Wkly Rep* [Internet]. 2021;70(36):1255–60. Available from: <https://doi.org/10.15585/mmwr.mm7036e2>.
337. Siegel D, Reses H, Cool A, Shapiro C, Hsu J, Boehmer T. Trends in COVID-19 cases, emergency department visits, and hospital admissions among children and adolescents aged 0–17 years — United States, August 2020–August 2021. *MMWR Morb Mortal Wkly Rep*. 2021;70:1–6.
 338. Hause AM, Gee J, Baggs J, Abara WE, Marquez P, Thompson D, et al. COVID-19 Vaccine Safety in Adolescents Aged 12–17 Years - United States, December 14, 2020–July 16, 2021. *MMWR Morb Mortal Wkly Rep* [Internet]. 2021;70(31):1053–8. Available from: <https://dx.doi.org/10.15585/mmwr.mm7031e1>
 339. Diaz G, Parsons G, Gering S, Meier A, Hutchinson I, Robicsek A. Myocarditis and pericarditis after vaccination for COVID-19 [Internet]. *JAMA*. 2021 [cited 2021 Sep 5]. Available from: <https://jamanetwork.com/journals/jama/fullarticle/2782900>
 340. Gargano J, Wallace M, Hadler S, Langley G, Su J. Use of mRNA COVID-19 Vaccine After Reports of Myocarditis Among Vaccine Recipients: Update from the Advisory Committee on Immunization Practices — United States, June 2021. *Morbidity Mortality Weekly Report*. 2021.
 341. Pepe S, Gregory A, Denniss A. Myocarditis, pericarditis and cardiomyopathy after COVID-19 vaccination. *Heart Lung Circ*. 2021;30:1425–9.
 342. Das B, Kohli U, Ramachandran P, HH N, Greil G, Hussain T, et al. Myopericarditis following mRNA COVID-19 Vaccination in Adolescents 12 through 18 Years of Age. *J Pediatr* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/80890fa62b48b86edd4a3b974c446c48dbf4e7cb>
 343. Schauer J, Buddhe S, Colyer J, Sagiv E, Law Y. Myopericarditis after the Pfizer mRNA COVID-19 Vaccine in Adolescents. *The Journal of pediatrics*.
 344. Snapiri O, C RD, Shirman N, Weissbach A, Lowenthal A, Ayalon I, et al. Transient Cardiac Injury in Adolescents Receiving the BNT162b2 mRNA COVID-19 Vaccine. *Pediatr Infect Dis J* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/e1edf775714d7c01efaf2c6400720cf7d0a350fa>
 345. Tano E, S SM, Girgis S, Martinez-Fernandez Y, C SV. Perimyocarditis in Adolescents After Pfizer-BioNTech COVID-19 Vaccine. *J Pediatric Infect Dis Soc* [Internet]. 2021; Available from: <http://www.epistemonikos.org/documents/1225cbdc1bb81b538ce18a4d78965ab7e6a2161>
 346. US FDA. Emergency use authorization (EUA) of the Pfizer-Biontech COVID-19 vaccine to prevent coronavirus disease 2019 (COVID019). [Internet]. [cited 2021 May 11]. Available from: <https://www.fda.gov/media/144413/download>
 347. European Medicines Agency. CHMP extension of indication variation assessment [Internet]. [cited 2021 Aug 10]. Available from: https://www.ema.europa.eu/en/documents/variation-report/spikevax-previously-covid-19-vaccine-moderna-epar-chmp-extension-indication-variation-assessment_en.pdf
 348. WHO. COVID-19 advice for the public: Getting vaccinated. July 14, 2021 [Internet]. [cited 2021 Aug 9]. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/covid-19-vaccines/advice>.
 349. Campos-Outcalt D. ACIP recommendations for COVID-19 vaccines—and more. *J Fam Pract*. 2021;70(2):86–92.
 350. National Advisory Committee on Immunization. Recommendation on the use of mRNA COVID-19 vaccines in adolescents 12 to 17 years of age.

351. JCVI statement on COVID-19 vaccination of children and young people aged 12 to 17 years: 4 August 2021.
352. Singapore Ministry of Health. Child vaccination. [Internet]. [cited 2021 Sep 17]. Available from: <https://www.moh.gov.sg/covid-19/vaccination/faqs---children-related-vaccination-matters>
353. PPS-PIDSP Statement on COVID-19 Vaccines for Children. 6 September 2021 [Internet]. [cited 2021 Oct 20]. Available from: <http://www.pidsphil.org/home/wp-content/uploads/2021/09/Updated-PPS-PIDSP-Statement-on-COVID-19-Vaccination-in-Children-Sept-2021.pdf>
354. Marshall M, Ferguson I, Lewis P, Jaggi P, Gagliardo C, JS C. Symptomatic acute myocarditis in 7 adolescents after Pfizer-BioNTech COVID-19 vaccination. Vol. 148, *Pediatrics*. 2021. p. e2021052478.
355. Lane S. Reports of myocarditis and pericarditis following mRNA COVID-19 vaccines: A review of spontaneously reported data from the UK, Europe, and the US. *medRxiv*2. 2021.
356. WHO. The Pfizer BioNTech (BNT162b2) COVID-19 vaccine: What you need to know. September 2, 2021 [Internet]. [cited 2021 Sep 17]. Available from: <https://www.who.int/news-room/feature-stories/detail/who-can-take-the-pfizer-biontech-covid-19--vaccine>
357. Butt A, Sabou-Samra A, Chemaitelly H, Al Khal A, Coyle P, Saleh H. Effectiveness of the SARS-CoV-2 mRNA Vaccines in Pregnant Women [Internet]. *ResearchSquare*. 2021 [cited 2021 Nov 23]. Available from: <https://doi.org/10.21203/rs.3.rs-622782/v1>
358. Dagan N, Barda N, Biron-Shental T, Makov-Assif M, Key C, Kohane I. Effectiveness of the BNT162b2 mRNA COVID-19 vaccine in pregnancy [Internet]. *Nature medicine*. 2021 [cited 2021 Sep 9]. Available from: <https://www.nature.com/articles/s41591-021-01490-8.pdf>
359. Bleicher I, Kadour-Peero E, Sagi-Dain L, Sagi S. Early exploration of COVID-19 vaccination safety and effectiveness during pregnancy: interim descriptive data from a prospective observational study. *Vaccine* [Internet]. 2021;39:6535–8. Available from: <https://doi.org/10.1016/j.vaccine.2021.09.043>
360. Theiler R, Wick M, Mehta R, Weaver A, Virk A, Swift M. Pregnancy and birth outcomes after SARS-CoV-2 vaccination in pregnancy [Internet]. *medRxiv*. 2021 [cited 2021 Dec 8]. Available from: <https://doi.org/10.1101/2021.05.17.21257337>
361. Beharier O, Mayo R, Raz T, Sacks K, Schreiber L, Suissa-Cohen Y, et al. Efficient maternal to neonatal transfer of antibodies against SARS-CoV-2 and BNT162b2 mRNA COVID-19 vaccine. *J Clin Invest* [Internet]. 2021;131(13):e150319. Available from: <https://doi.org/10.1172/JCI150319>
362. Collier A-RY, McMahan K, Yu J, Tostanoski LH, Aguayo R, Ansel J, et al. Immunogenicity of COVID-19 mRNA Vaccines in Pregnant and Lactating Women. *JAMA* [Internet]. 2021;325(23):2370–80. Available from: <https://dx.doi.org/10.1001/jama.2021.7563>
363. Mithal L, Shanes E, Miller E. Cord blood antibodies following maternal coronavirus disease 2019 vaccination during pregnancy [Internet]. *Am J Obstet Gynecol*. 2021 [cited 2021 Nov 23]. Available from: <https://doi.org/10.1016/j.ajog.2021.03.035>
364. Rottenstreich A, Zarbiv G, Oiknine-Dijan E, Zigrion R, Wolf D, Porat S. Efficient Maternofetal Transplacental Transfer of Anti- Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Spike Antibodies After Antenatal SARS-CoV-2 BNT162b2 Messenger RNA Vaccination. *Clin Infect Dis* [Internet]. 2021;73(10):1909–12. Available from: <https://doi.org/10.1093/cid/ciab266>
365. Zdanowski W, Wasniewski T. Evaluation of SARS-CoV-2 Spike Protein Antibody Titers in Cord Blood after COVID-19 Vaccination during Pregnancy in Polish HealthcareWorkers: Preliminary Results. *Vaccines* [Internet]. 2021;9:675–83. Available from: <https://doi.org/10.3390/v9060675>
366. Prabhu M, Murphy E, Sikhu A, Yee J, Singh S, Eng D. Antibody Response to Coronavirus Disease 2019 (COVID-19) Messenger RNA Vaccination in Pregnant Women and Transplacental Passage Into Cord Blood. *Obstet Gynecol*. 2021;138(2):278–80.

367. Shanes E, Otero S, Mithal L, Mupanomunda C, Miller E, Goldstein J. Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Vaccination in Pregnancy. *Obs Gynecol* [Internet]. 2021;138(2):281–3. Available from: <https://doi.org/10.1097/AOG.0000000000004457>
368. Gray K, Brodt E, Atyeo C, Deriso E, Akhmunni B, Young N. Coronavirus disease 2019 vaccine response in pregnant and lactating women: a cohort study. *Am J Obs Gynecol*. 2021;225(303):e1-17.
369. Shook L, Atyeo C, Yonker L, Gasano A, Gray K, Alter G, et al. Durability of anti-Spike antibodies in the infant after maternal COVID-19 vaccination [Internet]. *medRxiv*. 2021 [cited 2021 Nov 23]. Available from: <https://doi.org/10.1101/2021.11.17.21266415>
370. Atyeo C, DeRiso E, Davis C, Bordt E, DeGuzman R, Shook L, et al. COVID-19 mRNA vaccines drive differential Fc-functional profiles in pregnant, lactating, and non-pregnant women [Internet]. *bioRxiv*. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1101/2021.04.04.438404>
371. Peretz S, Regev N, Novic L, Nachshol M, Goffer E, Ben-David A. Short-term outcome of pregnant women vaccinated with BNT162b2 mRNA COVID-19 vaccine. *Ultrasound Obs Gynecol* [Internet]. 2021;58:450–6. Available from: <https://doi.org/10.1002/uog.23729>
372. Atyeo C, Shook L, Brigida S, de Guzman R, Demidkin S, Muir C, et al. Maternal immune response and placental antibody transfer after COVID-19 vaccination across trimester and platforms [Internet]. *medRxiv*. 2021 [cited 2021 Nov 23]. Available from: <https://doi.org/10.1101/2021.11.12.21266273>
373. Nir O, Schwartz A, Touessle-Cohen S, Leibovitch L, Strauss T, Asnaf K, et al. Maternal-neonatal transfer of SARS-CoV-2 immunoglobulin G antibodies among parturient women treated with BNT162b2 messenger RNA vaccine during pregnancy. *Am J Obs Gynecol* [Internet]. 2021;4:100492. Available from: <http://dx.doi.org/10.1016/j.ajogmf.2021.100492>
374. Shimabukuro T, Kim S, Myers T, Moro P, Oduyebo T, Panagiotakopoulos L. Preliminary Findings of mRNA Covid-19 Vaccine Safety in Pregnant Persons. *NEJM* [Internet]. 2021;384(24):2273–82. Available from: <https://doi.org/10.1056/NEJMoa2104983>
375. Kadali R, Janagama R, Peruru S, Racherla S, Tirumala R, Madathala R, et al. Adverse effects of COVID-19 messenger RNA vaccines among pregnant women: a cross-sectional study on healthcare workers with detailed self-reported symptoms [Internet]. *Am J Obstet Gynecol*. 2021 [cited 2021 Nov 23]. Available from: <https://doi.org/10.1016/j.ajog.2021.06.007>
376. Trostle M, Limaye M, Avtushka A, Lighter J, Penfield C. COVID-19 vaccination in pregnancy: early experience from a single institution [Internet]. *American Journal of Obstetrics and Gynecology*. 2021 [cited 2021 Dec 8]. Available from: <https://doi.org/10.1016/j.ajogmf.2021.100464>
377. Magnus M, Gjessing H, Eide H, Wilcox A, Fell D, Haberg S. Covid-19 Vaccination during Pregnancy and First-Trimester Miscarriage [Internet]. *NEJM*. 2021 [cited 2021 Nov 23]. Available from: <https://doi.org/10.1056/NEJMc2114466>
378. Wainstock T, Yoles I, Sergienko R, Sheiner E. Prenatal maternal COVID-19 vaccination and pregnancy outcomes. *Vaccine* [Internet]. 2021;39:6037–40. Available from: <https://doi.org/10.1016/j.vaccine.2021.09.012>
379. Zauche L, Wallace B, Smoots A, Olson C, Oduyebo T, Kim S, et al. Receipt of mRNA COVID-19 vaccines preconception and during pregnancy and risk of self-reported spontaneous abortions, CDC v-safe COVID-19 Vaccine Pregnancy Registry 2020-21 [Internet]. *ResearchSquare*. 2021 [cited 2021 Nov 23]. Available from: <https://doi.org/10.21203/rs.3.rs-798175/v1>
380. Charepe N, Goncalves J, Juliano A, Lopes D, Canhao H, Soares H, et al. COVID-19 mRNA vaccine and antibody response in lactating women: a prospective cohort study. *BMC Pregnancy Childbirth* [Internet]. 2021;21:632. Available from: <https://doi.org/10.1186/s12884-021-04051-6>
381. Valcarce V, Stafford L, Neu J, Cacho N, Parker L, Mueller M, et al. Detection of SARS-CoV-2-Specific IgA in the Human Milk of COVID-19 Vaccinated Lactating Health Care Workers [Internet]. *Breastfeeding Medicine*. 2021 [cited 2021 Nov 23]. Available from: <https://doi.org/10.1089/bfm.2021.0122>
382. Esteve-Palau E, Gonzales-Cuevas A, Guerrero M, Garcia-Terol C, Alvarez M, Garcia-Aranda G.

- Quantification of specific antibodies against SARS-CoV-2 in breast milk of lactating women vaccinated with an mRNA vaccine [Internet]. medRxiv. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1101/2021.04.05.21254819>
383. Juncker H, Mulleners S, van Gils M, de Groot C, Pajkrt D, Korosi A, et al. The Levels of SARS-CoV- 2 Specific Antibodies in Human Milk Following Vaccination. *J Hum Lact* [Internet]. 2021;37(3):477–84. Available from: <https://doi.org/10.1177/08903344211027112>
 384. Friedman M, Kigel A, Bahar Y, Yogeve Y, Dror Y, Lubetsky R. BNT162b2 COVID-19 mRNA vaccine elicits a rapid and synchronized antibody response in blood and milk of breastfeeding women [Internet]. medRxiv. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1101/2021.03.06.21252603>
 385. Schwartz A, Lebovich L, Strauss T, Asraf K, Doolman R. Presence of SARS-CoV-2 antibodies in lactating women and their infants following BNT162b2 messenger RNA vaccine [Internet]. *Am J Obstet Gynecol*. 2021 [cited 2021 Oct 31]. p. 577–9. Available from: <https://doi.org/10.1016/j.ajog.2021.07.016>
 386. Fox A, Norris C, Amanat F, Zolla-Pazner S, Powell R. The vaccine-elicited immunoglobulin profile in milk after COVID-19 mRNA-based vaccination is IgG-dominant and lacks secretory antibodies [Internet]. medRxiv. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1101/2021.03.22.21253831>
 387. Selma-Royo M, Bauer C, Mena-Tudela D, Aguilar-Camprubi L, Perez-Cano F, Parra-Llorca A. Anti-Sars-Cov-2 IgA And IgG In Human Milk After Vaccination Is Dependent On Vaccine Type And Previous Sars-Cov-2 Exposure: A Longitudinal Study [Internet]. medRxiv. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1101/2021.05.20.21257512>
 388. Golan Y, Prahi M, Cassidy A, Wu A, Jigmeddagva U. Immune response during lactation after anti-SARS-CoV2 mRNA vaccine [Internet]. medRxiv. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1101/2021.03.09.21253241>
 389. Low J, Gu Y, Feng M, Amin Z, Lee L, Ng Y, et al. BNT162b2 vaccination induces SARS-CoV-2 specific antibody secretion into human milk with minimal transfer of vaccine mRNA [Internet]. medRxiv. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1101/2021.04.27.21256151>
 390. Perl S, Uzan-Yulzati A, Klainer H, Asiskovich L, Youngster M, Rinott E, et al. SARS-CoV-2–Specific Antibodies in Breast Milk After COVID-19 Vaccination of Breastfeeding Women. *JAMA*. 2021;325(19):2013–4.
 391. Calil V, Palmeira P, Zheng Y, Krebs V, de Carvalho W, Carneiro-Sampaico M. CoronaVac can induce the production of anti-SARSCoV- 2 IgA antibodies in human milk [Internet]. *Clinics*. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.6061/clinics/2021/e3185>
 392. Low J, Lee L, Ng Y, Zhong Y, Amin Z. Breastfeeding mother-child clinical outcomes after COVID-19 vaccination [Internet]. medRxiv. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1101/2021.06.19.21258892>
 393. McLaurin-Jiang S, Garner C, Krusch K, Hale T. Maternal and Child Symptoms Following COVID-19 Vaccination Among Breastfeeding Mothers [Internet]. *Breastfeeding Medicine*. 2021 [cited 2021 Oct 31]. Available from: <https://doi.org/10.1089/bfm.2021.0079>
 394. Bertrand K, Honerkamp-Smith G, Chambers C. Maternal and Child Outcomes Reported by Breastfeeding Women Following Messenger RNA COVID-19 Vaccination. *Breastfeed Med*. 2021;16(9):697–701.
 395. American College of Obstetric-Gynecology. COVID-19 Vaccination Considerations for Obstetric–Gynecologic Care. Practice Advisory. December 3, 2021 [Internet]. [cited 2021 Dec 12]. Available from: https://www.acog.org/clinical/clinical-guidance/practice-advisory/articles/2020/12/covid-19-vaccination-considerations-for-obstetric-gynecologic-care?utm_source=redirect&utm_medium=web&utm_campaign=int
 396. Department of Health. Republic of the Philippines. Can Pregnant Women Get the COVID-19 Vaccine? [Internet]. [cited 2021 Dec 12]. Available from: <https://doh.gov.ph/node/28465>
 397. Philippine Obstetrical and Gynecological Society. Practice Bulletin COVID-19 Vaccination of Pregnant and Breastfeeding Women. August 9, 2021. [Internet]. 2021. [cited 2021 Dec 12].

- Available from: <https://pogsinc.org/practice-bulletins/>
398. Padmanaban U, Mukherjee S, Borse R, Joshi S, Deshmukh R. Phase II Clinical trial for Evaluation of BCG as potential therapy for COVID-19 [Internet]. MedRxiv. 2020 [cited 2021 Mar 6]. Available from: <https://doi.org/10.1101/2020.10.28.20221630>
 399. Giamarellos-Bourboulis E, Tsilika M, Moorlag S, Antonakos N, Kotsaki A, Dominguez-Andres J. Activate : Randomized Clinical Trial of BCG Vaccination against Infection in the Elderly. *Cell*. 2020;183:315–23.
 400. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: An emerging consensus on rating certainty of evidence and strength of recommendations. *Bmj*. 2008;336(7650):924–6.
 401. Deeks Jonathan J, Dinnes J, D'Amico R, Sowden A, Sakarovich C, Song F. Evaluating non-randomised intervention studies. *Health Technol Assess (Rockv)*. 2003;7:27.
 402. Reeves BC, Deeks Jonathan J, Higgins J, Wells G. Including non-randomised studies. In: Higgins J, Green S, editors. *Cochrane handbook for systematic reviews of interventions v502* [Internet]. v. 5.0.2. The Cochrane Collaboration; 2011. Available from: <http://www.cochrane-handbook.org>
 403. de Chaisemartin C, de Chaisemartin I. BCG vaccination in infancy does not protect against COVID-19. Evidence from a natural experiment in Sweden [Internet]. *Clinical Infectious Diseases*. 2020 [cited 2021 Apr 2]. Available from: <https://academic.oup.com/cid/advance-article/doi/10.1093/cid/ciaa1223/5896039>
 404. Hamiel U, Kozer F, Youngster I. SARS-CoV-2 Rates in BCG-Vaccinated and Unvaccinated Young Adults. *JAMA* [Internet]. 2020;323(22):2340–1. Available from: <https://jamanetwork.com/journals/jama/fullarticle/2766182>
 405. Amirlak I, Haddad R, Hardy JD, Khaled NS, Chung MH. Effectiveness of booster BCG vaccination in preventing Covid-19 infection. *MedRxiv*. 2020.
 406. Noval Rivas M, Ebinger J, Wu M, Sun N, Braun J. BCG vaccination history associates with decreased SARS-CoV-2 seroprevalence across a diverse cohort of healthcare workers. *J Clin Invest*. 2020;131(2).
 407. Moorlag S, van Deuren R, van Werkhoven C, Jaeger M, Bedisarun P, Taks E. Safety and COVID-19 Symptoms in Individuals Recently Vaccinated with BCG: a Retrospective Cohort Study. *Cell Rep*. 2020;1(5):100073.
 408. Sharma A, Ahmed A, Baig V, Shakad P, Dalela G, Kacker S. Characteristics and Outcomes of Hospitalized Young Adults with Mild to Moderate Covid-19 at a University Hospital in India. *MedRxiv*. 2020.
 409. WHO. Bacille Calmette-Guérin (BCG) vaccination and COVID-19. Scientific Brief [Internet]. 2020 [cited 2021 Apr 2]. Available from: [https://www.who.int/news-room/commentaries/detail/bacille-calmette-guérin-\(bcg\)-vaccination-and-covid-19](https://www.who.int/news-room/commentaries/detail/bacille-calmette-guérin-(bcg)-vaccination-and-covid-19)
 410. Blanco-Melo D, Nilsson-Payant BE, Liu WC, et al. Imbalanced host response to SARS-CoV-2 drives development of COVID-19. *Cell* 2020;181(5):1036-1045.e9.
 411. Wölfel R, Corman VM, Guggemos W, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature* 2020;581:465-9.
 412. Baum A, Fulton BO, Wloga E, et al. Antibody cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape seen with individual antibodies. *Science* 2020;369:1014-8.
 413. Hansen J, Baum A, Pascal KE, et al. Studies in humanized mice and convalescent humans yield a SARS-CoV-2 antibody cocktail. *Science* 2020;369:1010-4.
 414. O'Brien M, Forleo-Neto E, Musser B, et al. Subcutaneous REGEN-COV antibody combination for Covid-19 Prevention. *NEJM*. 2021 Aug 4. doi:10.1056/NEJMoa2109682.
 415. Gosling H. PMLive.com. US demand for COVID-19 antibody rising fast. [Internet]. 2021. [updated 2021 August 31; cited 2021 September 3]. Available from: https://www.pmlive.com/pharma_news/us_demand_for_covid-19_antibody_treatments_rising_fast_1376021.
 416. Lee B. Forbes.com. Regeneron antibody cocktail for Covid-19 coronavirus gets FDA emergency use authorization. [Internet]. 2020. [updated 2020 November 22, cited 2021 September 3]. Available from: <https://www.forbes.com/sites/brucelee/2020/11/22/regeneron-antibody-cocktail-for-covid-19-coronavirus-gets-fda-emergency-use-authorization/?sh=3310ee75cba8>.
 417. FDA.gov.ph. Emergency Use Authorization (EUA) for Casirivimab + Imdevimab (Ronapreve). [Internet]. 2021. Available from: <https://www.fda.gov.ph/wp-content/uploads/2021/10/EUA-Ronapreve-Website.pdf>. Accessed 8 November 2021.
 418. FDA.gov. FDA authorizes REGEN-COV monoclonal antibody therapy for post-exposure prophylaxis (prevention) for COVID-19. [Internet]. 2021. [updated 2021 August 10; cited 2021 September 3]. Available from: <https://www.fda.gov/drugs/drug-safety-and-availability/fda->

- authorizes-regen-cov-monoclonal-antibody-therapy-post-exposure-prophylaxis-prevention-covid-19.
419. FDA.gov. Reeneron EUA HCP Fact Sheet 09172021. [Internet]. 2021. [updated 2021 September 17; cited 2021 September 27]. Available from: <https://www.fda.gov/media/145611/download>.
 420. Bhimraj A, Morgan RL, Shumaker AH, Laverne V, Baden L, Cheng VC, et al. Infectious Diseases Society of America Guidelines on the Treatment and Management of Patients with COVID-19. Infectious Diseases Society of America 2021; Version 5.1.0. Available at <https://www.idsociety.org/practice-guideline/covid-19-guideline-treatment-and-management/>. Accessed 31 August 2021.
 421. Australian National COVID-19 Clinical Evidence Taskforce. Australian guidelines for the clinical cure of people with COVID-19 v42.1. Available at <https://app.magicapp.org/#/guideline/5571>. Accessed 1 November 2021.
 422. COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. National Institutes of Health. Available at <https://www.covid19treatmentguidelines.nih.gov/>. Accessed 1 November 2021.
 423. World Health Organization. Drugs to prevent COVID-19. 2 March 2021. Available at file:///Users/Carol/Downloads/WHO-2019-nCoV-prophylaxes-2021.1-eng%20(1).pdf. Accessed 24 September 2021 Tordjman S, Chokron S, Delorme R, et al. Melatonin: pharmacology, functions and therapeutic benefits. *Current neuropharmacology*. 2017;15(3):434-443.
 424. Zhang R, Wang X, Ni L, et al. COVID-19: Melatonin as a potential adjuvant treatment. *Life Sciences*. 2020;250:117583.
 425. Zhou Y, Hou Y, Shen J, Huang Y, Martin W, Cheng F. Network-based drug repurposing for novel coronavirus 2019-nCoV/SARS-CoV-2. *Cell Discovery*. 2020;6(1).
 426. Zhou Y, Hou Y, Shen J, et al. A network medicine approach to investigation and population-based validation of disease manifestations and drug repurposing for COVID-19. *PLOS Biology*. 2020;18(11):e3000970.
 427. Foley H, Steel A. Adverse events associated with oral administration of melatonin: A critical systematic review of clinical evidence. *Complementary Therapies in Medicine*. 2019;42:65-81.
 428. Andersen L, Gögenur I, Rosenberg J, Reiter R. The Safety of Melatonin in Humans. *Clinical Drug Investigation*. 2015;36(3):169-175.
 429. Technical guidance publications [Internet]. Who.int. 2020 [cited 29 December 2020]. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance-publications?publicationtypes=d198f134-5eed-400d-922e-1ac06462e676>
 430. Prevention of SARS-CoV-2 | COVID-19 Treatment Guidelines [Internet]. COVID-19 Treatment Guidelines. 2020 [cited 29 December 2020]. Available from: <https://www.covid19treatmentguidelines.nih.gov/overview/prevention-of-sars-cov-2/>
 431. COVID-19 Guideline, Part 2: Infection Prevention [Internet]. Idsociety.org. 2020 [cited 29 December 2020]. Available from: <https://www.idsociety.org/practice-guideline/covid-19-guideline-infection-prevention/>
 432. Efficacy of Melatonin in the Prophylaxis of Coronavirus Disease 2019 (COVID-19) Among Healthcare Workers. - Full Text View - ClinicalTrials.gov [Internet]. Clinicaltrials.gov. 2020 [cited 29 December 2020]. Available from: <https://clinicaltrials.gov/ct2/show/NCT04353128>
 433. Grant WB, Lahore H, McDonnell SL, et al. Evidence that Vitamin D Supplementation Could Reduce Risk of Influenza and COVID-19 Infections and Deaths. *Nutrients*. 2020;12(4).
 434. Xu J, Yang J, Chen J, Luo Q, Zhang Q, Zhang H. Vitamin D alleviates lipopolysaccharide-induced acute lung injury via regulation of the renin-angiotensin system. *Mol Med Rep*. 2017;16(5):7432-8.
 435. Ghasemian R, Shamshirian A, Heydari K, et al. The Role of Vitamin D in The Age of COVID-19: A Systematic Review and Meta-Analysis. *medRxiv*. 2020:2020.06.05.20123554.
 436. Jolliffe DA, Camargo CA, Sluyter JD, et al. Vitamin D supplementation to prevent acute respiratory infections: systematic review and meta-analysis of aggregate data from randomised controlled trials. *medRxiv*. 2020:2020.07.14.20152728.
 437. Annweiler C, Hanotte B, Grandin de l'Eprevier C, Sabatier JM, Lafaie L, Célarier T. Vitamin D and survival in COVID-19 patients: A quasi-experimental study. *J Steroid Biochem Mol Biol*. 2020;204:105771.

438. COVID-19 rapid guideline: vitamin D NICE guideline [NG187] Published date: 17 December 2020: National Institute for Health and Care Excellence; 2020. Available from: <https://www.nice.org.uk/guidance/ng187>.
439. Reducing Asymptomatic Infection With Vitamin D in Coronavirus Disease (RAID-CoV-2) [Internet]. US National Library of Medicine. Available from: <https://clinicaltrials.gov/ct2/show/NCT04476680>.
440. PRevention of COVID-19 With Oral Vitamin D Supplemental Therapy in Essential healthCare Teams (PROTECT). [Internet]. 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04483635>.
441. Vitamin D Supplementation in the Prevention and Mitigation of COVID-19 Infection (VitD-COVID19) [Internet]. 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04482673>.
442. Oral 25-hydroxyvitamin D3 and COVID-19 [Internet]. 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04386850>.
443. Trial of Vitamin D to Reduce Risk and Severity of COVID-19 and Other Acute Respiratory Infections (CORONAVIT) [Internet]. 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04579640>.
444. Vitamin D and COVID-19 Trial (VIVID) [Internet]. 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04536298>.
445. The Effect of D3 on Selected Cytokines Involved in Cytokine Storm in the Covid-19 Uninfected Jordanian People [Internet]. 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04476745>.
446. Vitamin D3 Supplementation to Prevent Respiratory Tract Infections [Internet]. 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04596657>.
447. Reducing Asymptomatic Infection With Vitamin D in Coronavirus Disease (RAID-CoV-2) [Internet]. 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04476680>.
448. te Velthuis AJ, van den Worm SH, Sims AC, Baric RS, Snijder EJ, van Hemert MJ. Zn(2+) inhibits coronavirus and arterivirus RNA polymerase activity in vitro and zinc ionophores block the replication of these viruses in cell culture. *PLoS Pathog*. 2010;6(11):e1001176.
449. Moseley HNB. Current Evidence Supporting the Use of Orally Administered Zinc in the Treatment of COVID-19. *OSF Preprints*. 2020.
450. Lassi ZS, Moin A, Bhutta ZA. Zinc supplementation for the prevention of pneumonia in children aged 2 months to 59 months. *Cochrane Database of Systematic Reviews*. 2016(12).
451. Roth DE, Richard SA, Black RE. Zinc supplementation for the prevention of acute lower respiratory infection in children in developing countries: meta-analysis and meta-regression of randomized trials. *International Journal of Epidemiology*. 2010;39(3):795-808.
452. Mayo-Wilson E, Imdad A, Junior J, Dean S, Bhutta ZA. Preventive zinc supplementation for children, and the effect of additional iron: a systematic review and meta-analysis. *BMJ Open*. 2014;4(6):e004647.
453. Veverka DV, Wilson C, Martinez MA, Wenger R, Tamosuinas A. Use of zinc supplements to reduce upper respiratory infections in United States Air Force Academy cadets. *Complement Ther Clin Pract*. 2009;15(2):91-5.
454. Prasad AS, Beck FW, Bao B, Fitzgerald JT, Snell DC, Steinberg JD, et al. Zinc supplementation decreases incidence of infections in the elderly: effect of zinc on generation of cytokines and oxidative stress. *Am J Clin Nutr*. 2007;85(3):837-44.
455. Zinc Fact Sheet for Health Professionals USA: National Institutes of Health Office of Dietary Supplements; 2020 [updated March 6, 2020. Available from: <https://ods.od.nih.gov/factsheets/Zinc-HealthProfessional/>.
456. Jothimani D, Kailasam E, Danielraj S, Nallathambi B, Ramachandran H, Sekar P, et al. COVID-19: Poor outcomes in patients with zinc deficiency. *Int J Infect Dis*. 2020;100:343-9.
457. Hemilä H. Zinc lozenges and the common cold: a meta-analysis comparing zinc acetate and zinc gluconate, and the role of zinc dosage. *JRSM Open*. 2017;8(5):2054270417694291.
458. National Institutes of Health. 2020. "Zinc Supplementation and COVID-19." COVID-19 Treatment Guidelines. <https://www.covid19treatmentguidelines.nih.gov/adjunctive-therapy/zinc/>.

459. A Randomized Open-label Prophylaxis Trial Among Migrant Workers at High-risk of COVID-19 (DORM Trial) [Internet]. Available from: <https://clinicaltrials.gov/ct2/show/NCT04446104?term=zinc+prophylaxis&type=Intr&cond=COVID-19&draw=2&rank=4>).
460. A Study of Hydroxychloroquine and Zinc in the Prevention of COVID-19 Infection in Military Healthcare Workers (COVID-Milit) [Internet]. Available from: <https://clinicaltrials.gov/ct2/show/NCT04377646>.
461. Zinc Versus Multivitamin Micronutrient Supplementation in the Setting of COVID-19 (ZnCOVID-19) [Internet]. US National Library of Medicine. 2020 [cited January 25, 2021]. Available from: <https://clinicaltrials.gov/ct2/show/NCT04551339>
462. Seet RCS, Quek AML, Ooi DSQ, Sengupta S, Lakshminarasappa SR, Koo CY, et al. Positive impact of oral hydroxychloroquine and povidone-iodine throat spray for COVID-19 prophylaxis: An open-label randomized trial. *Int J Infect Dis*. 2021;106:314-22.
463. Wang M, Cao R, Zhang L, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res*. 2020;30(3):269-271. Available from: doi:10.1038/s41422-020-0282-0 [Accessed 24th February 2021].
464. Zhou D, Dai SM, Tong Q. COVID-19: a recommendation to examine the effect of hydroxychloroquine in preventing infection and progression. *J Antimicrob Chemother*. 2020;75(7):1667-1670. Available from: doi:10.1093/jac/dkaa114 [Accessed 24th February 2021].
465. The RECOVERY Collaborative Group. Effect of Hydroxychloroquine in Hospitalized Patients with Covid-19. *N Engl J Med* 2020;383:2030-40. Available from: <https://www.nejm.org/doi/pdf/10.1056/NEJMoa2022926?articleTools=true> [Accessed 24th February 2021].
466. WHO Solidarity Trial Consortium. Repurposed Antiviral Drugs for Covid-19 — Interim WHO Solidarity Trial Results. *N Engl J Med* 2021;384:497-511. Available at: <https://www.nejm.org/doi/pdf/10.1056/NEJMoa2023184?articleTools=true> [Accessed 24th February 2021].
467. Lewis et al. The efficacy and safety of hydroxychloroquine for COVID-19 prophylaxis: A systematic review and meta- analysis of randomized trials. *PLoS ONE* 16(1): e0244778. Available at: <https://doi.org/10.1371/journal.pone.0244778> [Accessed 24th February 2021]
468. United States National Institutes of Health (US NIH), *COVID-19 Treatment Guidelines: Prevention and Prophylaxis of SARS CoV-2 Infection February 11, 2021*. Available from: <https://www.who.int/publications/i/item/clinical-management-of-covid-19> [Accessed 24th February 2021].
469. World Health Organization (WHO), *Clinical Management of COVID-19: Interim Guidance May 27, 2020*. Available from: <https://www.who.int/publications/i/item/clinical-management-of-covid-19> [Accessed 24th February 2021].
470. Philippine Society for Microbiology and Infectious Diseases Philippine College of Chest Physicians, Philippine College of Physicians Philippine Rheumatology Association, Philippine College of Hematology and Transfusion Medicine, *Interim Guidance on the Clinical Management of Adult Patients with Suspected or Confirmed COVID-19 Infection Version 3.1 July 20, 2020*. Available from: http://philchest.org/wp-content/uploads/2020/07/Final-PCP-PSMID-PCCP-COVID-19-Guidelines-20July2020_1595213265.pdf [Accessed 24th February 2021].
471. Palileo-Villanueva, L., & Cabaluna, I. T. *Should Chloroquine or hydroxychloroquine be used in the prophylaxis of COVID19? Version 3 June 22, 2020*. Available from: <https://www.psmid.org/rapidevidence-review-on-covid-19-management/> [Accessed 24th February 2021]]
472. Cochrane Methods. Risk of Bias. Available at <https://methods.cochrane.org/bias/risk-bias-tool>. [Accessed 24th February 2021]
473. GRADEpro. Available at <https://gradepro.org/> [Accessed 24th February 2021]

474. Boulware et al. A Randomized Trial of Hydroxychloroquine as Postexposure Prophylaxis for Covid-19. *N Engl J Med* 2020; 383:517-525. Available from: DOI: 10.1056/NEJMoa2016638 [Accessed 24th February 2021].
475. Abella et al. Efficacy and Safety of Hydroxychloroquine vs Placebo for Pre-exposure SARS-CoV-2 Prophylaxis Among Health Care Workers. *JAMA Internal Medicine* 2021;181(2):195-202 Available from: doi:doi:10.1001/jamainternmed.2020.6319 [Accessed 24th February 2021].
476. Rajasingham et al. Hydroxychloroquine as Pre-exposure Prophylaxis for Coronavirus Disease 2019 (COVID-19) in Healthcare Workers: A Randomized Trial. *CID* 2020. Available from: DOI: 10.1093/cid/ciaa1571 [Accessed 24th February 2021].
477. Barnabas et al. Hydroxychloroquine as Postexposure Prophylaxis to Prevent Severe Acute Respiratory Syndrome Coronavirus 2 Infection. *Annals of Internal Medicine*. Available from: <https://doi.org/10.7326/M20-6519> [Accessed 24th February 2021].
478. Mitja et al. A Cluster-Randomized Trial of Hydroxychloroquine for Prevention of Covid-19. *N Engl J Med* 2021; 384:417-427. Available from: DOI:10.1056/NEJMoa2021801 [Accessed 24th February 2021].
479. [17] Indian Express, *Preventive use of HCQ in frontline healthcare workers: ICMR study*, June 4, 2020. Available from <https://indianexpress.com/article/india/preventive-use-of-hcq-in-frontline-healthcare-workers-icmr-study-6442948/> [Accessed 24th February 2021]
480. Gentile I, Maraolo AE, Piscitelli P. COVID-19 : Time for Post-Exposure Prophylaxis? 2020;17–9.
481. Park SY, Lee JS, Son JS, et al. Post-exposure prophylaxis for Middle East respiratory syndrome in healthcare workers. *J Hosp Infect* [Internet]. 2019 Jan;101(1):42–6. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0195670118304845>
482. Diaz-Brito V, Leon A, Knobel H, et al. Post-exposure prophylaxis for HIV infection: a clinical trial comparing lopinavir/ritonavir versus atazanavir each with zidovudine/lamivudine. 2012;346:337–46.
483. Jin Y, Zhan Q, Peng Z, et al. Chemoprophylaxis, diagnosis, treatments, and discharge management of COVID-19 : An evidence-based clinical practice guideline (updated version). 2020;1–33.
484. Hellwig MD, Maia A. A COVID-19 prophylaxis? Lower incidence associated with prophylactic administration of ivermectin. *International Journal of Antimicrobial Agents*. 2020 Nov;57(2021).
485. Alam MT, Murshe R, Gomes PF, Masud ZM, Saber S, Chaklader MA et al. Ivermectin as Pre-exposure Prophylaxis for COVID-19 among Healthcare Providers in a Selected Tertiary Hospital in Dhaka – An Observational Study. *European Journal of Medical and Health Sciences*. 2020 Dec;2(6):1-6.
486. Carvallo H, Roberto H, Psaltis A, Contrera V. Study of the Efficacy and Safety of Topical Ivermectin + Iota-Carrageenan in the Prophylaxis against COVID-19 in Health Personnel. *Journal of Biomedical Research and Clinical Investigation*. 2020 Nov;2(1.1007).
487. Caly L, Druce JD, Catton MG, Jans DA, Wagstaff KM. The FDA-approved drug ivermectin inhibits the replication of SARS-CoV-2 *in vitro*. *Antiviral Research*. 2020 Apr;178(2020):1-4.
488. Kory P, Meduri GU, Iglesias J, Varon J, Berkowitz K, Kornfeld H, et al. Review of the Emerging Evidence Demonstrating the Efficacy of Ivermectin in the Prophylaxis and Treatment of COVID-19. Preprint [Internet]. 2020 Nov [cited 2021 Mar 30]. Available from: https://www.researchgate.net/publication/347781954_Review_of_the_Emerging_Evidence_Demonstrating_the_Efficacy_of_Ivermectin_in_the_Prophylaxis_and_Treatment_of_COVID-19.
489. Bryant A, Lawrie TA, Dowswell T, Fordham E, Mitchell S, Hill S, et al. Ivermectin for Prevention and Treatment of COVID-19 Infection: a Systematic Review and Meta-analysis. Preprint [Internet]. 2021 [cited 2021 Apr 5]. Available from: <https://www.researchsquare.com/article/rs-317485/v1>.
490. British Ivermectin Recommendation Development. The BIRD Recommendation on the Use of Ivermectin for COVID-19. Bath, UK: BIRD; 2021 Feb 20 [cited 2021 Mar 30]. Available from <https://www.francesoir.fr/sites/francesoir/files/media-icons/bird-proceedings-02-03-2021-v151.pdf>.

491. Behera P, Patro BK, Singh AK, Chandanshive PD, Ravikumar SR, Prandhan SK, et al. Role of ivermectin in the prevention of SARS-CoV-2 infection among healthcare workers in India: A matched case-control study. *PLoS ONE*. 2021 Feb;16(2):e0247163.
492. Behera P, Patro BK, Padhy BM, Mohapatra PR, Bal SK, Chandanshive PD, et al. Prophylactic role of ivermectin in SARS-CoV-2 infection among healthcare workers. Preprint [Internet]. 2021 [cited 2021 Mar 30]. Available from: <https://www.researchsquare.com/article/rs-208785/v1>.
493. Bernigaud C, Guillemot D, Ahmed-Belkacem A, Grimaldi-Bensouda L, Lespine A, Berry F, et al. Oral ivermectin for a scabies outbreak in a long-term care facility: potential value in preventing COVID-19 and associated mortality. *British Journal of Dermatology*. 2021.
494. Elgazzar A, Eltaweel A, Youssef SA, Hany B, Hafez M, Moussa H. Efficacy and Safety of Ivermectin for Treatment and Prophylaxis of COVID-19 Pandemic. Preprint [Internet]. 2021 [cited 2021 Mar 16]. Available from: <https://www.researchsquare.com/article/rs-100956/v3> doi: <https://doi.org/10.21203/rs.3.rs-100956/v3>.
495. Shoumann WM, Hegazy AA, Nafae RM, Ragab MI, Samra SR, Ibrahim DA, et al. Use of Ivermectin as a potential chemoprophylaxis for COVID-19 in Egypt: A Randomised Clinical Trial. *J Clin Diagn Res*. 2021 Feb;15(2):OC27-32.
496. Chahla RE, Medina Ruiz L, Ortega ES, Morales MF, Barreiro F, George A, et al. A Randomized Trial – Intensive Treatment Based in Ivermectin and Iota-Carrageenan as Pre-exposure Prophylaxis for COVID-19 in Healthcare Agents. Preprint [Internet]. 2021 [cited 2021 Apr 21]. Available from <https://www.medrxiv.org/content/10.1101/2021.03.26.21254398v1>.
497. UpToDate. Ivermectin (systemic): Drug information. UpToDate; 2021 [cited 2021 Mar 30].
498. European Medicines Agency. EMA advises against use of ivermectin for the prevention or treatment of COVID-19 outside of randomized clinical trials. Amsterdam, Netherlands: European Medicines Agency; 2021 Mar 22 [cited 2021 Mar 27]. Available from: <https://www.ema.europa.eu/en/news/ema-advises-against-use-ivermectin-prevention-treatment-covid-19-outside-randomised-clinical-trials>.
499. Alberta Health Services. COVID-19 Scientific Advisory Group Rapid Evidence Report: Ivermectin in the Treatment and Prevention of COVID-19 [Internet]. Alberta, Canada: Alberta Health Services; 2021 Feb 02 [cited 2021 Mar 27]. Available from: <https://www.albertahealthservices.ca/assets/info/ppih/if-ppih-covid-19-sag-ivermectin-in-treatment-and-prevention-rapid-review.pdf>.
500. Infectious Diseases Society of America. Infectious Diseases Society of America Guidelines on the Treatment and Management of Patients with COVID-19 [Internet]. USA: IDSA; 2021 Mar 18 [cited Mar 28]. Available from: <https://www.idsociety.org/globalassets/idsa/practice-guidelines/covid-19/treatment/idsa-covid-19-gl-tx-and-mgmt-v4.1.1.pdf>.
501. Australian Government Department of Health. COVID-19 treatments [Internet]. Australia: Australian Government Department of Health; 2021 Feb 23 [cited 2021 Mar 28]. Available from: <https://www.health.gov.au/news/health-alerts/novel-coronavirus-2019-ncov-health-alert/covid-19-treatments>.
502. World Health Organization. WHO advises that ivermectin only be used to treat COVID-19 within clinical trials [Internet]. Geneva, Switzerland: WHO; 2021 Mar 31 [cited 2021 Apr 5]. Available from: <https://www.who.int/news-room/feature-stories/detail/who-advises-that-ivermectin-only-be-used-to-treat-covid-19-within-clinical-trials>.
503. US Food and Drug Administration. Why You Should Not Use Ivermectin to Treat or Prevent COVID-19 [Internet]. Maryland, USA: US FDA; 2021 Mar 05 [cited 2021 Mar 27] Available from: <https://www.fda.gov/consumers/consumer-updates/why-you-should-not-use-ivermectin-treat-or-prevent-covid-19>.
504. Food and Drug Administration Philippines. FDA Advisory No. 2021-0526 [Internet]. Muntinlupa, Philippines: FDA; 2021 Mar 15 [cited 2021 Mar 27]. Available from: <https://www.fda.gov.ph/wp-content/uploads/2021/03/FDA-Advisory-No.2021-0526.pdf>.
505. Rappler. Hospital gets limited permit to use ivermectin for COVID-19 treatment [Internet]. Pasig City, Philippines: Rappler; 2021 Apr 8 [cited 2021 Apr 11]. Available from:

- <https://www.rappler.com/nation/hospital-gets-limited-permit-use-ivermectin-covid-19-treatment-april-2021>.
506. ABS-CBN. FDA: 2nd hospital given compassionate special permit to use ivermectin [Internet]. Quezon City, Philippines: ABS-CBN; 2021 Apr 16 [cited 2021 Apr 17]/ Available from: <https://news.abs-cbn.com/news/04/16/21/fda-2nd-hospital-given-compassionate-special-permit-to-use-ivermectin>.
 507. Šlapak I, Skoupá J, Strnad P, Horník P. Efficacy of isotonic nasal wash (seawater) in the treatment and prevention of rhinitis in children. *Arch Otolaryngol - Head Neck Surg*. 2008;134(1):67–74.
 508. Tano L, Tano K. A daily nasal spray with saline prevents symptoms of rhinitis. *Acta Otolaryngol*. 2004;124(9):1059–62.
 509. Liu S-C, Tu Y-K, Chien M-N, Chien K-L. Effect of antidiabetic agents added to metformin on glycaemic control, hypoglycaemia and weight change in patients with type 2 diabetes: a network meta-analysis. *Diabetes, Obes Metab* [Internet]. 2012 Sep;14(9):810–20. Available from: <http://doi.wiley.com/10.1111/j.1463-1326.2012.01606.x>
 510. Casale M, Rinaldi V, Sabatino L, Moffa A, Ciccozzi M. Could nasal irrigation and oral rinse reduce the risk for COVID-19 infection? *Int J Immunopathol Pharmacol*. 2020;34:1–3.
 511. Singh S, Sharma N, Singh U, Singh T, Mangal D, Singh V. Nasopharyngeal wash in preventing and treating upper respiratory tract infections: Could it prevent COVID-19? *Lung India*. 2020; 37:246–51.
 512. Baruah B. Could Simultaneous Nasal and Oral Irrigation Be a Nontherapeutic Tool against SARS-CoV-2? *ACS Chem Neurosci*. 2021;0–2.
 513. Panta P, Chatti K, Andhavarapu A. Do saline water gargling and nasal irrigation confer protection against COVID-19? *Explore* [Internet]. 2020 Oct;(January). Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1550830720303128>
 514. Ramalingam S, Graham C, Dove J, Morrice L, Sheikh A. Hypertonic saline nasal irrigation and gargling should be considered as a treatment option for COVID-19. *J Glob Health*. 2020;10(1):19–22
 515. Singh M, Singh M, Jaiswal N, Chauhan A. Heated, humidified air for the common cold [Internet]. Vol. 2017, *Cochrane Database of Systematic Reviews*. John Wiley and Sons Ltd; 2017 [cited 2021 Feb 22]. Available from: <https://pubmed.ncbi.nlm.nih.gov/33232690/>
 516. Pawar S, Bhonsale D, Pawar D, et al. Management of COVID positive medical staff: A medical college and public hospital experience with steam therapy. *Med. Med. Sci*. 2020; 8(6): 056-058.
 517. la Marca G, Barp J, Frenos S, Mugelli A, Galli L, Calistri E, et al. Thermal inactivation of SARS COVID-2 virus: Are steam inhalations a potential treatment? *Life Sci* [Internet]. 2021 Jan 15 [cited 2021 Feb 22];265. Available from: <https://pubmed.ncbi.nlm.nih.gov/33232690/>
 518. Brewster CT, Choong J, Thomas C, Wilson D, Moiemmen N. Steam inhalation and paediatric burns during the COVID-19 pandemic. *Lancet* [Internet]. 2020;395(10238):1690. Available from: [http://dx.doi.org/10.1016/S0140-6736\(20\)31144-2](http://dx.doi.org/10.1016/S0140-6736(20)31144-2)
 519. Lillicrap D, 2020. Disseminated intravascular coagulation in patients with 2019-nCoV pneumonia. *J Thromb Haemost* 18: 786–787.
 520. Al-Samkari H, Karp Leaf RS, Dzik WH, Carlson JCT, Fogerty AE, Waheed A, Goodarzi K, Bendapudi PK, Bornikova L, Gupta S, Leaf DE, Kuter DJ, Rosovsky RP. COVID-19 and coagulation: bleeding and thrombotic manifestations of SARS-CoV-2 infection. *Blood*. 2020 Jul 23;136(4):489-500. doi: 10.1182/blood.202006520. PMID: 32492712; PMCID: PMC7378457.
 521. Kashi M, Jacquin A, Dakhil B, Zaimi R, Mahé E, Tella E, Bagan P. Severe arterial thrombosis associated with Covid-19 infection. *Thromb Res*. 2020 Aug;192:75-77. doi: 10.1016/j.thromres.2020.05.025. Epub 2020 May 16. PMID: 32425264; PMCID: PMC7229939.
 522. Carsana L, Sonzogni A, Nasr A, Rossi RS, Pellegrinelli A, Zerbi P, Rech R, Colombo R, Antinori S, Corbellino M, Galli M, Catena E, Tosoni A, Gianatti A, Nebuloni M. Pulmonary post-mortem findings in a series of COVID-19 cases from northern Italy: a two-centre descriptive study. *Lancet Infect Dis*. 2020 Oct;20(10):1135-1140. doi: 10.1016/S1473-3099(20)30434-5. Epub 2020 Jun 8. PMID: 32526193; PMCID: PMC7279758.

523. Nopp S, Moik F, Jilma B, Pabinger I, Ay C. Risk of venous thromboembolism in patients with COVID-19: A systematic review and meta-analysis. *Res Pract Thromb Haemost*. 2020 Sep 25;4(7):1178–91. doi: 10.1002/rth2.12439. Epub ahead of print. PMID: 33043231; PMCID: PMC7537137.
524. Lodigiani C et al.; Humanitas COVID-19 Task Force, 2020. Venous and arterial thromboembolic complications in COVID-19 patients admitted to an academic hospital in Milan, Italy. *Thromb Res* 191: 9–14.
525. Viecca M, Radovanovic D, Forleo GB, Santus P. Enhanced platelet inhibition treatment improves hypoxemia in patients with severe Covid-19 and hypercoagulability. A case control, proof of concept study. *Pharmacol Res* 2020;158:104950.
526. Meizlish ML, Goshua G, Liu Y, Fine R, Amin K, Chang E, DeFilippo N, Keating C, Liu Y, Mankbadi M, McManus D, Wang SY, Price C, Bona RD, Ochoa Chaar CI, Chun HJ, Pine AB, Rinder HM, Siner JM, Neuberg DS, Owusu KA, Lee AI. Intermediate-dose anticoagulation, aspirin, and in-hospital mortality in COVID-19: A propensity score-matched analysis. *Am J Hematol*. 2021 Apr 1;96(4):471-479. doi: 10.1002/ajh.26102. Epub 2021 Feb 22. PMID: 33476420; PMCID: PMC8013588.
527. Ahmed HY, Papali A, Haile T, Shrestha GS, Schultz MJ, Lundeg G, Akrami KM, For The Covid-Lmic Task Force. Pragmatic Recommendations for the Management of Anticoagulation and Venous Thrombotic Disease for Hospitalized Patients with COVID-19 in Low- and Middle-Income Countries. *Am J Trop Med Hyg*. 2021 Jan 11;104(3 Suppl):99–109. doi: 10.4269/ajtmh.20-1305. Epub ahead of print. PMID: 33432908; PMCID: PMC7957232.

Evidence and Recommendations for the Adjunct Interventions for COVID-19

1. Abd-Elsalam S, Soliman S, Esmail ES, et al. Do zinc supplements enhance the clinical efficacy of hydroxychloroquine? A randomized, multicenter trial. *Biol Trace Elem Res*. 2021; 199: 3642-3646
2. Thomas S, Patel D, Bittel B, et al. Effect of high-dose zinc and ascorbic acid supplementation vs usual care on symptom length and reduction among ambulatory patients with SARS-CoV-2 infection: the COVID A to Z randomized clinical trial. *JAMA Netw Open*. 2021; 4: e210369
3. Kaplan HG, Wang K, Reeves KM, Scanlan JM, Nunn CC, Kieper DA, et al. Resveratrol and zinc in the treatment of outpatients with covid-19 – the reszinate study - a phase 1/2 randomized clinical trial utilizing home patient-obtained nasal and saliva viral sampling. SSRN. 2021 Oct; [cited 2021 Nov 20]. Available from: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3934228.
4. Patel O, Chinni V, El-Khoury J, Perera M, Neto AS, McDonald C, et al. A pilot double-blind safety and feasibility randomized controlled trial of high-dose intravenous zinc in hospitalized COVID-19 patients. (Short Communication) *J Med Virol*. 2021 May;93(5):3261-3267. doi: 10.1002/jmv.26895. Epub 2021 Mar 9. PMID: 33629384; PMCID: PMC8014767.
5. Darban M, Malek F, Memarian M, Gohari A, Kiani A, Emadi A, et al. Efficacy of High Dose Vitamin C, Melatonin and Zinc in Iranian Patients with Acute Respiratory Syndrome due to Coronavirus Infection: A Pilot Randomized Trial. *J Cell Mol Anesth*. 2021;6(2):164-7.DOI: <https://doi.org/10.22037/jcma.v6i2.32182>
6. Abdelmaksoud AA, Ghweil AA, Hassan MH, et al. Olfactory Disturbances as Presenting Manifestation Among Egyptian Patients with COVID-19: Possible Role of Zinc. *Biol Trace Elem Res* 199, 4101–4108 (2021). Available from: <https://doi.org/10.1007/s12011-020-02546-5>
7. Tan CW, Pock L, Kalimuddin S, et al. Cohort study to evaluate the effect of vitamin D, magnesium, and vitamin B12 in combination on progression to severe outcomes in older patients with coronavirus (COVID-19). *Nutrition*. 2020; 79-80. Available from: <https://doi.org/10.1016/j.nut.2020.111017>
8. Dalbeni A, Bevilacqua M, Teani I, Normelli I, Mazzaferri F, Chiarioni G. Excessive vitamin B12 and poor outcome in COVID-19 pneumonia. *Nutr Metab Cardiovasc Dis*. 2021 Mar 10;31(3):774-775. doi: 10.1016/j.numecd.2020.12.005. Epub 2020 Dec 11. PMID: 33549452; PMCID: PMC7834259.

9. Jamalimoghadamsiahkali S, Zaresade B, Koolaji S, SeyedAlinaghi S, Zندهدل A, Tabarestani M, et al. Safety and effectiveness of high-dose vitamin C in patients with COVID-19: a randomized open-label clinical trial. *Eur J of Med Res*. 2021; 26(1):20.
10. Kumari P, Dembra S, Dembra P, Bhawna F, Gul A, Ali B, et al. The role of vitamin C as adjuvant therapy in COVID-19. *Cureus* 2020; 12(11):e11779
11. Zhang J, Rao X, Li Y, Zhu Y, Liu F, Guo G, et al. Pilot trial of high dose vitamin C in critically ill COVID-19 patients. *Ann Intensive Care* 2021; 11:5.
12. Tehrani S, Yadegarynia D, Abrishami A, Moradi H, Gharaei B, Raoufi M, et al. An investigation into the effects of intravenous vitamin C on pulmonary CT findings and clinical outcomes of patients with COVID 19 pneumonia A Randomized Clinical Trial. 2021.
13. Thomas S, Patel D, Bittel B, Wolski K, Wang Q, Kumar A, et al. Effect of high dose zinc and ascorbic acid supplementation vs usual care on symptom length and reduction among ambulatory patients with SARS-CoV-2 infection. The COVID A to Z randomized clinical trial. *JAMA Network Open*. 2021;4(2):e210369
14. Darban M, Malek F, Memarian M, Gohari A, Kiani A, Emadi A, et al. Efficacy of high dose vitamin C, melatonin and zinc in Iranian patients with Acute Respiratory Syndrome due to Coronavirus Infection: A pilot randomized Trial. *Journal of Cellular & Molecular Anesthesia (JCMA)* 2021;6(2).
15. Hakamifard A, Id R, Soltani, Id A, Maghsoudi, Id, et al. The effect of vitamin E and vitamin C in patients with COVID-19 pneumonia; a randomized controlled clinical trial *Correspondence to. *Immunopathol Persa*. 2021;8(1):8.
16. Beigmohammadi MT, Bitarafan S, Hoseindokht A, Abdollahi A, Amoozadeh L, Soltani D. The effect of supplementation with vitamins A, B, C, D, and E on disease severity and inflammatory responses in patients with COVID-19: a randomized clinical trial. *Trials*. 2021 Nov 14;22(1).
17. Murai IH, Fernandes AL, Sales LP, Pinto AJ, Goessler KF, Duran CSC, et al. Effect of a Single High Dose of Vitamin D3 on Hospital Length of Stay in Patients With Moderate to Severe COVID-19: A Randomized Clinical Trial. *JAMA*. 2021; 325: 1053–1060.
18. Castillo ME, Entrenas Costa LM, Vaquero Barrios JM, Alcalá Díaz JF, Miranda JL, Bouillon R, et al. 'Effect of Calcifediol Treatment and best Available Therapy versus best Available Therapy on Intensive Care Unit Admission and Mortality Among Patients Hospitalized for COVID-19: A Pilot Randomized Clinical study'. *J Steroid Biochem Mol Biol*. 2020; 105751.
19. Rastogi A, Bhansali A, Khare N, Suri V, Yaddanapudi N, Sachdeva N, et al. Short term, high-dose vitamin D supplementation for COVID-19 disease: a randomised, placebo-controlled, study (SHADE study). *Postgrad Med J*. Epub ahead of print November 2020. DOI: 10.1136/postgradmedj-2020-139065.
20. Lakkireddy M, Gadiga SG, Malathi RD, Karra ML, Raju ISSVPM, Ragini, et al. Impact of daily high dose oral vitamin D therapy on the inflammatory markers in patients with COVID 19 disease. *Sci Rep*. 2021; 11: 10641.
21. Elamir YM, Amir H, Lim S, Rana YP, Lopez CG, Feliciano NV, et al. A randomized pilot study using calcitriol in hospitalized COVID-19 patients. *Bone*. 2021; 154: 116175.
22. Maghbooli Z, Sahraian MA, Jamalimoghadamsiahkali S, Asadi A, Zarei A, Zندهدل A, et al. Treatment With 25-Hydroxyvitamin D(3) (Calcifediol) Is Associated With a Reduction in the Blood Neutrophil-to-Lymphocyte Ratio Marker of Disease Severity in Hospitalized Patients With COVID-19: A Pilot Multicenter, Randomized, Placebo-Controlled, Double-. *Endocr Pract Off J Am Coll Endocrinol Am Assoc Clin Endocrinol*. Epub ahead of print October 2021. doi: 10.1016/j.eprac.2021.09.016.
23. Soliman AR, Abdelaziz TS, Fathy A. Impact of Vitamin D Therapy on the Progress COVID-19: Six Weeks Follow-Up Study of Vitamin D Deficient Elderly Diabetes Patients. *Proc Singapore Healthc*. 2021; 20101058211041404.
24. Sánchez-Zuno GA, González-Estevez G, Matuz-Flores MG, Macedo-Ojeda G, Hernández-Bello J, Mora-Mora JC, et al. Vitamin D Levels in COVID-19 Outpatients from Western Mexico: Clinical Correlation and Effect of Its Supplementation. *J Clin Med*. 2021; 10(11) 2378. Epub ahead of print May 2021. doi: 10.3390/jcm10112378.
25. Farnoosh G, Akbari qomi M, Badri T, et al. Efficacy of a Low Dose of Melatonin as an Adjunctive Therapy in Hospitalized Patients with COVID-19: A Randomized, Double-blind Clinical Trial. *Authorea Prepr [Internet]*. 2020; Available from: <https://www.authorea.com/doi/full/10.22541/au.160734344.45295921/v1?commit=5be3e7266256468d59e81ff82a1b125710ba7459>
26. Lazarte CC. 2-stage, Randomized, double-blind, placebo-controlled clinical trial on the efficacy and safety of Lagundi (Vitex negundo) tablets/syrup (NIRPROMP formulation) with standard treatment compared to placebo with standard treatment in patients with mild COVID-19 disease

- without comorbidities. [Internet]. Health Research and Development Information Network. 2021. Available from: <https://www.herdin.ph/index.php/registry?view=research&layout=details&cid=2992>
27. Doaei S, Gholami S, Rastgoo S, Gholamalizadeh M, Bourbour F. The effect of omega - 3 fatty acid supplementation on clinical and biochemical parameters of critically ill patients with COVID - 19: a randomized clinical trial. *J Transl Med* [Internet]. 2021;1–9. Available from: <https://doi.org/10.1186/s12967-021-02795-5>
 28. de Alencar JCG, Moreira CL, Müller AD, et al. COVID Register Group. Double-blind, randomized, placebo-controlled trial with N-acetylcysteine for treatment of severe acute respiratory syndrome caused by COVID-19. *Clin Infect Dis*. 2020 Sep 23;ciaa1443. doi: 10.1093/cid/ciaa1443. Epub ahead of print. PMID: 32964918; PMCID: PMC7543361.
 29. Ibrahim H, Perl A, Smith D, et al. Therapeutic blockade of inflammation in severe COVID-19 infection with intravenous N-acetylcysteine. *Clin Immunol*. 2020 Oct;219:108544. doi: 10.1016/j.clim.2020.108544. Epub 2020 Jul 22. PMID: 32707089; PMCID: PMC7374140.
 30. Lu X, Ma Y, He J, Li Y, Zhu H, Yu X. N-acetylcysteine for adults with acute respiratory distress syndrome: A meta-analysis of randomized controlled trials. *Hong Kong Journal of Emergency Medicine*. 2019Jan9;26(5):288–98.
 31. Cohen JB, Hanff TC, William P, et al. Continuation versus discontinuation of renin–angiotensin system inhibitors in patients admitted to hospital with COVID-19: a prospective, randomised, open-label trial. *Lancet Respir Med*. 2021;2(20):1–10.
 32. Lopes RD, Macedo AVS, De Barros E, et al. Effect of Discontinuing vs Continuing Angiotensin-Converting Enzyme Inhibitors and Angiotensin II Receptor Blockers on Days Alive and out of the Hospital in Patients Admitted with COVID-19: A Randomized Clinical Trial. *JAMA - J Am Med Assoc*. 2021;325(3):254–64.
 33. Bikdeli B, Talasaz A, Rashidi F, et al. Intermediate versus standard-dose prophylactic anticoagulation and statin therapy versus placebo in critically-ill patients with COVID-19: Rationale and design of the INSPIRATION/INSPIRATION-S studies. *Thrombosis Research*. 2020 Sept 24. Available from: <https://doi.org/10.1016/j.thromres.2020.09.027>
 34. Bikdeli Behnood, Talasaz A, Rashidi F, et al. Atorvastatin vs Placebo in Patients with COVID-19 Admitted to the ICU: The INSPIRATION-S Trial. In: American College of Cardiology Annual Scientific Session [Internet]; 2021 May 16; Virtual. Available from <https://www.acc.org/education-and-meetings/image-and-slide-gallery/media-detail?id=9DF17809F35C4CA6B149C64C9D700AC6>
 35. Davoodi L, Jafarpour H, Oladi Z et al. Atorvastatin therapy in COVID-19 adult inpatients: A double-blind, randomized controlled trial. *IJC Heart & Vasculature*. 2021 Sept 14. Available from: <https://www.sciencedirect.com/science/article/pii/S2352906721001639?via%3Dihub>
 36. Choi M, Ahn H, Ryu H, et al. Clinical Characteristics and Disease Progression in Early-Stage COVID-19 Patients in South Korea. *J Clin Med*. 2020; 9(6):1959. doi:10.3390/jcm9061959
 37. Esba L, Alqahtani R, Thomas A. Ibuprofen and NSAID Use in COVID-19 Infected Patients Is Not Associated with Worse Outcomes: A Prospective Cohort Study. *Infect Dis Ther*. 2020 Nov 2. Available from: <https://doi.org/10.1007/s40121-020-00363-w>
 38. Kragholm K, Gerds T, Fosbol E et al. Association Between Prescribed Ibuprofen and Severe COVID-19 Infection: A Nationwide Register-Based Cohort Study. *Clin Transl Sci*. 2020; 13:1103–1107. doi:10.1111/cts.12904
 39. Perkins S, Holmes A, Nelson J, et al. Clinical outcomes among COVID-19 patients taking non-steroidal anti-inflammatory drugs. *Ann Emerg Med*. 2020 Oct; 76(4).
 40. Rinnot E, Kozer E, Shapira Y, Bar-Haim A, Youngster I. Ibuprofen use and clinical outcomes in COVID-19 patients. *Clin Microbiol Infect*. 2020; 26 (9):1259.e5-1259.e7.
 41. Samimagham H, Arabi M, Hooshyar D, Kazemijahromi M. The association of non-steroidal anti-inflammatory drugs with COVID-19 severity and mortality. *Arch Clin Infect Dis*. 2020 August; 15(4):e106847.
 42. Viecca M, Radovanovic D, Forleo GB, Santus P. Enhanced platelet inhibition treatment improves hypoxemia in patients with severe Covid-19 and hypercoagulability. A case control, proof of concept study. *Pharmacol Res* 2020;158:104950.
 43. Meizlish ML, Goshua G, Liu Y, Fine R, Amin K, Chang E, DeFilippo N, Keating C, Liu Y, Mankbadi M, McManus D, Wang SY, Price C, Bona RD, Ochoa Chaar CI, Chun HJ, Pine AB, Rinder HM, Siner JM, Neuberg DS, Owusu KA, Lee AI. Intermediate-dose anticoagulation, aspirin, and in-hospital mortality in COVID-19: A propensity score-matched analysis. *Am J Hematol*. 2021 Apr 1;96(4):471-479. doi: 10.1002/ajh.26102. Epub 2021 Feb 22. PMID: 33476420; PMCID: PMC8013588.

44. Ahmed HY, Papali A, Haile T, Shrestha GS, Schultz MJ, Lundeg G, Akrami KM, For The Covid-Lmic Task Force. Pragmatic Recommendations for the Management of Anticoagulation and Venous Thrombotic Disease for Hospitalized Patients with COVID-19 in Low- and Middle-Income Countries. *Am J Trop Med Hyg.* 2021 Jan 11;104(3 Suppl):99–109. doi: 10.4269/ajtmh.20-1305. Epub ahead of print. PMID: 33432908; PMCID: PMC7957232.
45. Carrouel F, Valette M, Gadea E, Esparcieux A, Illes G, Langlois ME, et al. Use of an antiviral mouthwash as a barrier measure in the SARS-CoV-2 transmission in adults with asymptomatic to mild COVID-19: a multicentre, randomized, double-blind controlled trial. *Clinical Microbiology.* 2021;xxx:1-7.
46. Eduardo F de P, Corrêa L, Heller D, Daep CA, Benitez C, Malheiros Z, et al. Salivary SARS-CoV-2 load reduction with mouthwash use: A randomized pilot clinical trial. *Heliyon.* 2021;7(6):1-7.
47. Di Domênico MB, Cesca H, Ponciano THJ, dos Santos RB, Lenz U, Antunes VP, et al. Effectiveness of hydrogen peroxide as auxiliary treatment for hospitalized COVID-19 patients in Brazil: Preliminary results of a randomized double-blind clinical trial. *Epidemiol Health.* 2021;43:1-7.
48. Guenezan J, Garcia M, Strasters D, Jousselin C, Leveque N, Frasca D, et al. Povidone Iodine Mouthwash, Gargle, and Nasal Spray to Reduce Nasopharyngeal Viral Load in Patients with COVID-19: A Randomized Clinical Trial. *JAMA Otolaryngology-Head & Neck Surgery.* 2021;147(4):400-401.
49. Jayaraman BG, Rajan G, Kannian P, Lavanya C, Ravichandran K, Kumarasamy N, et al. Povidone iodine, hydrogen peroxide and chlorhexidine mouthwashes reduce SARS-CoV2 burden in whole mouth fluid and respiratory droplets. *medRxiv [Internet].* 2021;2021.02.25.21252488. Available from: <http://medrxiv.org/content/early/2021/03/01/2021.02.25.21252488.abstract>
50. Seneviratne CJ, Balan P, Ki KKK, et al. Efficacy of commercial mouth-rinses on SARS-CoV-2 viral load in saliva: Randomized Control Trial in Singapore. 2020.
51. Mohamed NA, Baharom N, Wan Sulaiman WS, et al. Early viral Clearance among COVID-19 patients when gargling with povidone-iodine and essential oils – A clinical trial. *medRxiv.* 2020.
52. Figueroa JM, Lombardo ME, Dogliotti A, Flynn LP, Giugliano R, Simonelli G, et al. Efficacy of a Nasal Spray Containing Iota-Carrageenan in the Postexposure Prophylaxis of COVID-19 in Hospital Personnel Dedicated to Patients Care with COVID-19 Disease. *Int J Gen Med.* 2021 Oct 1;14:6277-6286. Doi: 10.2147/IJGM.S328486. PMID: 34629893; PMCID: PMC8493111.
53. Hosseinzadeh A, Tavakolian A, Kia V, Ebrahimi H, Sheibani H, Binesh E, et al. Application of nasal spray containing dimethyl sulfoxide (DMSO) and ethanol during the COVID-19 pandemic may protect healthcare workers: A randomized controlled trial. 2021. Available from: <https://www.medrxiv.org/content/10.1101/2021.07.06.21259749v2>.
54. Shrivastava R, Vijay M, Maneby N, Shrivastava R. Clinical Efficacy of an Osmotic, Antiviral and Anti-inflammatory Polymeric Nasal Film to Treat Covid-19 Early-Phase Respiratory Symptoms. *Open Access J Clin Trials.* 2021 May 18;13:11-20. Doi: 10.2147/OAJCT.S307144
55. Winchester S, John S, Jabbar K, John I. Clinical efficacy of nitric oxide nasal spray (NONS) for the treatment of mild COVID-19 infection. *J Infect.* 2021 Aug;83(2):237-279. Doi: 10.1016/j.jinf.2021.05.009. Epub 2021 May 13. PMID: 33992687; PMCID: PMC8117664.
56. Guenezan J, Garcia M, Strasters D, Jousselin C, Lévêque N, Frasca D, Mimos O. Povidone Iodine Mouthwash, Gargle, and Nasal Spray to Reduce Nasopharyngeal Viral Load in Patients With COVID-19: A Randomized Clinical Trial. *JAMA. Otolaryngol Head Neck Surg.* 2021 Apr 1;147(4):400-401. Doi: 10.1001/jamaoto.2020.5490. PMID: 33538761; PMCID: PMC7863011.
57. Domênico MBD, Collares K, Santos RBD, Lenz U, Antunes VP, Godinho VW, et al. Hydrogen peroxide as an auxiliary treatment for COVID-19 in Brazil: a randomized double-blind clinical trial. *Epidemiol Health.* 2021;43:e2021051. Doi: 10.4178/epih.e2021051. Epub 2021 Aug 3. PMID: 34529913.
58. Arefin MK. Povidone Iodine (PVP-I) Oro-Nasal Spray: An Effective Shield for COVID-19 Protection for Health Care Worker (HCW), for all. *Indian J Otolaryngol Head Neck Surg.* 2021 Apr 8:1-6. Doi: 10.1007/s12070-021-02525-9. Epub ahead of print. PMID: 33846691; PMCID: PMC8026810.
59. Zarabanda D, Vukkadala N, Phillips KM, Qian ZJ, Mfuh KO, Hatter MJ, et al. The Effect of Povidone-Iodine Nasal Spray on Nasopharyngeal SARS-CoV-2 Viral Load: A Randomized Control Trial. *Laryngoscope.* 2021 Nov 1. Doi: 10.1002/lary.29935. Epub ahead of print. PMID: 34724213.

60. Aref ZF, Bazeed SEES, Hassan MH, Hassan AS, Rashad A, Hassan RG, Abdelmaksoud AA. Clinical, Biochemical and Molecular Evaluations of Ivermectin Mucoadhesive Nanosuspension Nasal Spray in Reducing Upper Respiratory Symptoms of Mild COVID-19. *Int J Nanomedicine*. 2021 Jun 15;16:4063-4072. Doi: 10.2147/IJN.S313093. PMID: 34163159; PMCID: PMC8215847.
61. Kasiri H, Rouhani N, Salehifar E, Ghazaeian M, Fallah S. Mometasone furoate nasal spray in the treatment of patients with COVID-19 olfactory dysfunction: A randomized, double blind clinical trial. *Int Immunopharmacol*. 2021 Sep;98:107871. Doi: 10.1016/j.intimp.2021.107871. Epub 2021 Jun 12. PMID: 34147912; PMCID: PMC8196319.
62. Yildiz E, Koca Yildiz S, Kuzu S, Günebakan Ç, Bucak A, Kahveci OK. Comparison of the Healing Effect of Nasal Saline Irrigation with Triamcinolone Acetonide Versus Nasal Saline Irrigation alone in COVID-19 Related Olfactory Dysfunction: A Randomized Controlled Study. *Indian J Otolaryngol Head Neck Surg*. 2021 Jul 10:1-6. Doi: 10.1007/s12070-021-02749-9. Epub ahead of print. PMID: 34277384; PMCID: PMC8272442.

DISCUSSION

Outputs of the Philippine COVID-19 Living CPG Project

CLINICAL PRACTICE QUESTIONS

COVID-19 management issues and questions were collected from the Steering Committee members and Consensus Panelists during the organizational meetings and consensus panel meetings. Management trends and new issues were added to this list as they emerge or as suggested by health policymakers. The topics were reviewed and prioritized regularly. Priority topics were then assigned to the evidence reviewers for new evidence reviews or updating of existing reviews. A total of 139 priority topics were identified.

CONSENSUS MEETINGS, EVIDENCE SUMMARIES, AND RECOMMENDATIONS

For the first phase (December 2020 to May 2021) there were a total of 90 evidence summaries presented with 136 recommendations generated in 26 consensus panel meetings.

For the second phase (October to December 2021), there were a total of 70 evidence summaries presented and 139 recommendations generated during 25 consensus panel meetings. 36 evidence summaries were new reviews while 34 were updates of the Refer to Appendix D for the schedule of panel presentations.

Continued surveillance for new evidence and succeeding updates will be covered by the next phase of the living CPG.

Applicability Issues

The members of the Consensus Panels provided information on the facilitators, barriers, and resource implications for the implementation of the recommendations. They used their expertise and experience to identify these issues, which were discussed in more detail in the *Consensus Issues* section of each evidence summary. These were considered in the final wording of the recommendations. The following subsections summarize the overall discussion of the panelists.

ORGANIZATIONAL CONSIDERATIONS TO IMPLEMENTATION

The availability of testing kits and medical equipment for the screening and diagnostic tests for COVID-19 would likely vary at the regional, provincial, or even municipal/ city level. These issues were especially relevant to RT-PCR testing, rapid antibody, and antigen testing, chest imaging (X-ray, CT-Scan, and ultrasound), and laboratory parameters (LDH, CRP, Ferritin, D-dimer). Clinical risk assessment and using the 14-day symptom test were useful tools for screening for COVID-19, especially if there was a limitation in the availability of screening tests. Specially trained personnel were needed to do the more specialized tests, such as pooled testing using RT-PCR.

Aside from the availability of various testing modalities, there would be some limitations in the availability of treatment and critical care interventions also, most especially those investigational drugs only being accessible through the public via FDA's emergency use authorization. Medical specialists, especially those from infectious diseases, pulmonary medicine, and critical care medicine, were important to effectively lead in the use of these treatments for the management of COVID-19 patients. These limitations would be further compounded by the limitations in available isolation beds, hospital ward beds, and ICU beds.

For non-pharmacologic interventions and proven prophylactic interventions (such as vaccines) for COVID-19, one potentially major barrier was the public's perceptions of these interventions and their actual compliance. This was evident in many instances of violations of the minimum public health standards set by DOH: wearing of face mask and face shields, physical distancing, and hand hygiene. In addition to these, there were rising trends in the use of non-proven prophylactic interventions (such as ivermectin), ineffective medical devices (such as ionizing air filters), and the anti-COVID vaccine movement.

RESOURCE IMPLICATIONS

As a low-middle-income country, our limited resources needed to be allocated and used efficiently. The cost of the tests and interventions being done for COVID-19 management was one important consideration discussed in the panel meetings, especially the investigational drugs (such as remdesivir, tocilizumab) and the highly sophisticated interventions (such as ECMO, hyperbaric oxygen therapy). Health technology assessment should be a key gatekeeping mechanism to ensure that all payments by the government (through PhilHealth) are cost-effective.

Implementation Tools

Selected recommendations from the Philippine COVID-19 Living CPG have been used as a reference to the Unified COVID-19 Algorithms, specifically on testing and management. These algorithms were developed collaboratively by 15 professional organizations and stakeholder institutions. Their complete algorithms were published on the PSMID website (<https://www.psmid.org/unified-covid-19-algorithms-5/>). Healthcare providers, patients, and the public are encouraged to access these algorithms on the main webpage.

Process Evaluation

Using the native website tools and Google analytics, the PSMID website administrator was able to gather various metrics on website visits for the Philippine COVID-19 Living CPG: click trends and download trends. Since the release on March 31, 2021, there have been 246,982 total clicks on the website. Most of these visits from the initial launch were from the National Capital Region, Central Visayas (Region VII), and Davao Region (Region XI), which reflected the areas that had the greatest number of new cases and active cases during the project duration (Figure 7 and Table 6). Furthermore, the list of top evidence summaries clicks revealed the possible topics that CPG users needed the most guidance on (Table 7).

Figure 6. Geographical location of website visitors of the Philippine COVID-19 Living CPG.




Table 6. Top locations of website visitors of the Philippine COVID-19 Living CPG. (as of June 30, 2021)

City	n	Percentage
Quezon City	27,848	18.77
Makati	20,202	13.61
Cebu City	11,112	7.49
Manila	9,706	6.54
Meycauayan	7,994	5.39
Davao City	7,249	4.89
Pasig	3,944	2.66
Paranaque	3,772	2.54
Angeles	3,376	2.28
Antipolo	3,092	2.08
Others	10,191	7.14

Table 7. Top evidence summaries downloaded from the Philippine COVID-19 Living CPG (as of August 8, 2022)

Topic	n
Ivermectin as treatment	4061
Hemoperfusion	4049
Face Shield	3090
Remdesivir	2170
Ivermectin as prophylaxis	2086
Saliva RT-PCR	1980
Lianhua	1815
Rapid Antigen Test	1321
Tocilizumab	1221
14 Day Symptom Based Test	1188

Research Implications

The novel coronavirus, now known as SARS-CoV-2, brought about a disease condition that is new to everyone. Despite the rapidly evolving evidence on COVID-19, many research gaps need to be filled in the management, prevention, and control of this disease. These were identified during the evidence reviews done in this CPG and were documented in the evidence summaries. The following discussion presents a synthesis of these research gaps.

As expected in a novel disease condition, many of the recommendations were answered with low to very low certainty of evidence. These areas are directed to a need for further primary research to be conducted.

While existing studies on the investigational treatment interventions have been able to identify the subset of patients that would benefit best, there is still a need for further studies on dosing, frequency of administration, combinations with other drugs, etc. Aside from the successful experimental drugs, additional research on those drugs showing early promise is urgently required.

Diagnosis and treatment were sometimes overemphasized in the management of COVID-19. Recommendations on vaccination were highly anticipated due to the vaccine roll out, including vaccination for the special populations (i.e. children and pregnant/lactating women) as well as booster vaccination. Equally important were the other prophylactic and non-pharmacologic interventions that are more proximal steps in the national strategy of prevention, detection, isolation, treatment, and reintegration. These studies were also crucial to prove the lack of effectiveness of several interventions that many subscribe to.

Finally, the living CPG methodology used in this project was the first local adoption known to the project team. Research into streamlining the living CPG process is important to make it more efficient. The impact measurement of this living CPG, as described in the *Monitoring and Auditing Criteria* subsection, would be another first study to formally demonstrate the effects of CPG implementation in the country.

Conclusions and Recommendations

Phase 2 of the Philippine COVID-19 Living CPG identified 70 priority questions on COVID-19 management, infection prevention, and control, generated 69 evidence summaries, and came up with 139 recommendations. Thematic areas included in this CPG were screening and diagnosis, treatment, critical care, and respiratory management, non-pharmacologic interventions, vaccines and prophylactic interventions, and adjunct interventions. The CPG recommendations were used in the construction of testing and management algorithms for COVID-19. Process evaluation using website analytics revealed that the CPG recommendations were mostly accessed in regions with the greatest number of new cases and active cases. Furthermore, a list of top evidence summaries accessed indicated topics that CPG users needed the most guidance on, or that remain to be contentious as of the present date.

The main challenges in doing a living CPG for a new disease condition in a pandemic setting were the rapidly evolving evidence and the need to come out with point in time recommendations for clinicians and policymakers. Consensus panels needed to balance the quality and totality of the evidence with the net benefit and the contextual factors related to the implementation of the interventions, i.e., cost, equity, acceptability, and feasibility. Flexibility and adaptability are key in developing a Living CPG, especially in the context of the pandemic. Given this project experience, we recommend the following for the succeeding updating of the Philippine COVID-19 Living CPG:

1. Retain consensus panel members who wish to continue contributing their time and expertise to the COVID-19 Living CPG and to ensure patient representation among all panels.
2. Continue holding capacity building workshops on CPG development, systematic reviews, and evidence-based medicine to increase the pool of skilled evidence reviewers.
3. Allow a longer project cycle for the implementation of the Living CPG development. This will ensure that adequate preparation is done by the task forces and consensus panelists prior to the en banc meeting.
4. Continue the practice of employing one technical coordinator, technical assistant and copy editor per task force to ensure the timely submission of quality evidence summaries and other deliverables.
5. Create an independent oversight committee to oversee that all task forces are complying with the accepted recommendations and a subcommittee to manage the COIs of all task force members. The review and management of conflicts of interest of all project members takes a considerable amount of time hence should be given consideration in the project timeline.
6. Involvement of stakeholders, including policy makers, especially identifying priority topics to be covered by the CPG is highly encouraged.

REFERENCES

1. World Health Organization. WHO Coronavirus (COVID-19) Dashboard. 2021. Retrieved August 20, 2021, from <https://covid19.who.int/>.
2. Philippine Department of Health. DOH COVID-19 Tracker. 2021. Retrieved August 20, 2021, from <https://doh.gov.ph/covid19tracker>.
3. Department of Health Philippines. 2018. Manual for Clinical Practice Guideline Development.
4. Schünemann H, Brożek J, Guyatt G, Oxman A, editors. GRADE handbook for grading quality of evidence and strength of recommendations. Updated October 2013. The GRADE Working Group, 2013. Available from guidelinedevelopment.org/handbook.
5. Brouwers MC, Kerkvliet K, Spithoff K, on behalf of the AGREE Next Steps Consortium. The AGREE Reporting Checklist: a tool to improve reporting of clinical practice guidelines. *BMJ* 2016;352:i1152. doi: 10.1136/bmj.i1152.
6. Akl EA, Meerpohl JJ, Elliott J, Kahale LA, Schünemann HJ; Living Systematic Review Network. Living systematic reviews: 4. Living guideline recommendations. *J Clin Epidemiol*. 2017 Nov;91:47-53. doi: 10.1016/j.jclinepi.2017.08.009. Epub 2017 Sep 11. PMID: 28911999.
7. Dans, AL, Dans, LF, Silvestre, MAA. Eds. Painless Evidence-Based Medicine, Second Edition. John Wiley & Sons Ltd. DOI: 10.1002/9781119196150
8. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality if nonrandomized studies in meta-analyses, 2012. Retrieved August 20, 2021, from http://www.ohri.ca/programs/clinical_epidemiology/nosgen.pdf

APPENDICES

Appendix A. Members of the Philippine COVID-19 Living CPG Task Force

STEERING COMMITTEE

Marissa M. Alejandria, MD, MSc, FPCP, FPSMID
Co-Chair

Professor, University of the Philippines College of Medicine Department of Clinical Epidemiology
Head, Research Implementation and Development Office, College of Medicine, University of the Philippines
Director, Institute of Clinical Epidemiology, National Institutes of Health University of the Philippines
Clinical Professor, Division of Infectious Diseases, Department of Medicine, Philippine General Hospital
Adult Infectious Diseases Specialist, The Medical City
President, Philippine Society for Microbiology and Infectious Diseases

Leonila F. Dans MD, MSc, FPPS, FPRA

Co-Chair
Professor, Department of Clinical Epidemiology, University of the Philippines-Manila
Professor, Department of Pediatrics, University of the Philippines- Philippine General Hospital
Fellow, Philippine Pediatric Society
Fellow, Philippine Rheumatology Association
Faculty, Asia-Pacific Center for Evidence-based Healthcare

Antonio L. Dans, MD, MSc, FPCP

Professor 12, College of Medicine, University of the Philippines- Manila
Faculty, Asia-Pacific Center for Evidence-based Healthcare

Noel L. Espallardo, MD, MSc, FPAFP

Board of Trustees, Philippine Academy of Family Physicians
Technical Specialist, Better Health Program

Aileen R. Espina, RMT, MD, MPH, MHA, FPAFP, CESE

Independent Consultant for Health Systems Strengthening and Disaster Risk Resilience
Member, Medical Advisory Group of Centre Medicales Internationale
Member, Technical and Policy Committee, Philippine Society of Public Health Physicians
Vice President, Foundation for Family Medicine Educators Executive Member at Large, Asia Pacific Regional Council, World Organization of Family Doctors (WONCA)
Chair, WONCA Working Party on Women and Family Medicine

Jemelyn U. Garcia, MD, FPCP, FPSMID

Medical Specialist III, Research Institute for Tropical Medicine
Assistant Secretary, Philippine Society for Microbiology and Infectious Diseases

Evalyn A. Roxas, MD, MPH, FPCP, FPSMID

Clinical Associate Professor, Division of Infectious Diseases, Department of Medicine, University of the Philippines
Associate Professor, Department of Medical Microbiology, College of Public Health, University of the Philippines-Manila
Head, Section of Infectious Diseases, Department of Medicine, Manila Medical Center
President, Philippine Hospital Infection Control Society Inc.

Mario M. Panaligan, MD, FPCP, FACP, FPSMID, FIDSA

Assistant Professor of Medicine, College of Medicine, University of the East, Ramon Magsaysay Memorial Medical Center, Inc.
Medical Specialist II and Head, Section of Infectious Diseases, Department of Medicine, Dr. Jose R. Reyes Memorial Medical Center
Infection Control Coordinator, St. Luke's Medical Center
Member, Data Safety Monitoring Committee, Department of Health, Philippines
Past President, Philippine College of Physicians and Philippine Society for Microbiology and Infectious Diseases

Ivan N. Villespin, MD, MBA, FPCP, FPCCP, FCCP

Associate Professor of Medicine, Faculty of Medicine and Surgery, University of Santo Tomas
Chief Program Officer, Office of Continuing Medical Education, Faculty of Medicine and Surgery, University of Santo Tomas
Active Medical Staff, University of Santo Tomas Hospital
Lead Consultant, Ventilator Development for COVID-19, DOST-EPDC-PCIEERD
Fellow, Philippine College of Physicians; Philippine College of Chest Physicians; American College of Chest Physicians
Board Member, Philippine College of Physicians and Philippine Specialty Board of Internal Medicine

CONSENSUS PANELS

Consensus Panel for Screening and Diagnosis

Florido A. Atibagos Jr., MD, FPSP

Assistant Professor, FEU NRMF Institute of Medicine and UERMMMC College of Medicine
Chairman, Research Management Committee, Jose B. Lingad Memorial Regional Hospital
Residency Training Program Coordinator, Philippine Heart Center

Clemencia D. Bondoc, MD

Municipal Health Officer, Zarraga, Iloilo
Past President, Association of Municipal Health Officers of the Philippines

Jane Eflynn L. Lardizabal-Bunyi, RPh, MD, OHP, DFM, FPAFP, CSPSH

Assistant Professor III, Manila Central University – Filemon D. Tanchoco Medical Foundation
National Treasurer, Philippine Academy of Family Physicians

John Andrew T. Camposano, MD, FPPS, DPIDSP

Medical Specialist, Western Visayas Medical Center, Western Visayas Sanitarium

Virgina de los Reyes, MD, FPCCP, FPCP, FPSSM, MHPED

Section Head, Section of Sleep Medicine, Department of Pulmonary, Critical Care and Sleep Medicine, Lung Center of the Philippines
Training Officer, Pulmonary Fellowship Program, Lung Center of the Philippines
Active consultant, Lung Center of the Philippines; San Juan de Dios Active Consultant; Metro North Hospital
Associate Professor, Ateneo School of Medicine and Public Health

Mary Ann D. Lansang, MD, MSc, FPCP, FPSMID

Clinical Professor, Department of Clinical Epidemiology, College of Medicine, University of the Philippines
Consultant, Infectious Diseases Section, Department of Medicine, The Medical City

Aretha Ann C. Gacutan-Liwag, MD, FPSEDM, FPCP

Quality Management Representative, West Visayas State University Medical Center
Chair, Research Management Committee, Western Visayas Regional Health Research Development Consortium

Marilyn A. Bermudez-Puyot, MD, FPCEM

Consultant, Emergency Department, ACE Medical Center
Medical Specialist I, Department of Emergency Medicine and Acute Care, Pasig City General Hospital

Fatima Johanna T. Santos-Ocampo, MD, FPPS, FPSAAI

Head, Immunodeficiency Council, Philippine Society of Allergy, Asthma and Immunology
Founding Member, Asia Pacific Society of Immunodeficiency
Consultant, Makati Medical Center
Consultant, Asian Hospital and Medical Center

Anelyn L. Reyes, MD, RMT, FPPS, FPIDSP, MHA, MBA

Assistant Professor I, College of Medicine, University of Santo Tomas

Arthur Dessi Roman MD, MTM, FPCP, FPSMID

Board of Council and Treasurer of the Board, Philippine Society for Microbiology and Infectious Diseases
Medical Specialist III and Training Officer, Research Institute for Tropical Medicine
Clinical Associate Professor, Philippine General Hospital

Vernon M. Serafico, MD, FPCP

Asst. Training Officer, Department of Internal Medicine, De Los Santos Medical Center Private
General Internist, Ang Dr. Serafico Medical Clinic
Board of Trustee, Philippine Society of General Internal Medicine
Treasurer, Philippine College of Physician, QC Chapter

Consensus Panel For Treatment

Jenifer R. Otadoy-Agustin, MD, FPCP, FPSAAI

Clinical Associate Professor, Department of Medicine,
Philippine General Hospital
Active Staff, University of Perpetual Help DALTA
Medical Center
Training Officer, Division of Allergy and Immunology,
Department of Medicine, Philippine General Hospital

Mary Ann C. Bunyi, MD, FPPS, FPIDSP

President, Pediatric Infectious Disease Society of the
Philippines
Assistant Professor III, College of Medicine,
Pamantasan ng Lungsod ng Maynila
Consultant, Section of Pediatric Infectious Disease,
Medicine Department, Philippine Children's Medical
Center

Maria Elinore Alba-Concha, MD, FPAFP

Chief Training Officer, Southern Philippines
Medical Center
Member, Research and Publications Committee,
Philippine Academy of Family Physicians

Erwin R. De Mesa, MD, FPOGS, FPIDSOG

Training Officer, Department of OB-GYN, De Los
Santos Medical Center
Head, Section of Infectious Diseases, Department of
OB-GYN, Quirino Memorial Medical Center
Treasurer, Philippine Obstetrical and Gynecological
Society
Immediate Past President, Philippine Infectious
Diseases Society for Obstetrics and Gynecology

Leila R. Ferrer, MD, MAED, CSPSH, FPCGM

Adjunct Faculty, Institute of Aging, National Institutes
of Health, University of the Philippines Manila
Consultant, ACE Medical Center

Sarah May Flores, CSSYB

Supervising Health Program Officer, Disease
Prevention and Control Bureau, Department of Health

Faith Joan C. Mesa-Gaerlan, MD, MS, FPCEM

Clinical Associate Professor, College of Medicine,
University of the Philippines Manila
Training Officer, Department of Emergency Medicine,
Philippine General Hospital

Karl Evans R. Henson, MD, FPCP, FPSMID

Clinical Assistant Professor, Division of Infectious
Diseases, College of Medicine, University of the
Philippines
Training Officer, Infectious Diseases Fellowship
Training Program, The Medical City
Director, Hospital Infection Control and Epidemiology
Center, The Medical City

Maria Encarnita B. Limpin, MD, FPCP, FPSCCM, FPSSM

President, Philippine College of Physicians
Chair, Department of Internal Medicine, Mary
Johnston Hospital
Medical Specialist IV, Philippine Heart Center

Sarah R. Makalinaw, MD, DPPS, DPIDSP

Consultant, Victor R. Potenciano Medical Center
Medical Specialist II, Rizal Medical Center

Roland M. Panaligan, MD, LLM, FPCP, FPCCP

Chair, Department of Medical Ethics, University of
Santo Tomas Faculty of Medicine and Surgery
Training Officer, Adult Pulmonology Fellowship
Training Program, Center for Respiratory Medicine,
University of Santo Tomas Hospital
Chair, Council on Diagnostics and Therapeutics,
Philippine College of Chest Physicians

Rommel B. Punongbayan, RMT, MD, MBA, FPCP, FPSMS, CSPSH, DPCOM

Medical Specialist II, Bulacan Medical Center
Chair, Occupational Health, The Medical City Clark
Vice President, Philippine Society of General
Internal Medicine
Chair Research, Philippine College of Physicians
National
Board of Director, Philippine College of
Occupational Medicine

Rowena Roselle P. Blanco-Santos, MD FPCOM, CMA

Telemedicine Consultant, Pilipinas Shell Petroleum
Corporation
Assistant Professor II, College of Medicine, Our
Lady of Fatima University, Valenzuela City

Consensus Panel for Critical Care and Respiratory Management

Maaliddin B. Biruar, MD, FPCP, FPSN

Senior Medical Director, Global Medical Services,
PAREXEL International
Consultant, National Kidney Transplant Institute
Vice President, Philippine Society of Nephrology
Regent, Philippine College of Physicians

Joseph Adrian L. Buensalido, MD, FPCP, FPSMID

Board Member, Philippine Society for Microbiology
and Infectious Diseases
Infection Prevention and Control Chair, Asian Hospital
and Medical Center Associate Professor of Medicine
and Deputy Training Officer, Division of Infectious
Diseases, Department of Medicine, University of the
Philippines - Philippine General Hospital

Pauline F. Convocar, MD, MCHM, DPBEM, FPCEM, DPCOM

Vice Chair for Patient Services Quality Management
System and Telemedicine, Department of Emergency
Medicine, Southern Philippines Medical Center
Vice-Chair and Residency Training Program Director,
Department of Emergency Medicine, Corazon Locsin
Montelibano Memorial Regional Hospital Section
Immediate Past President & Section on Advocacy
Chair Philippine College of Emergency Medicine
Current Board Member, Asian Society of Emergency
Medicine

Reynaldo C. De Castro Jr., MD, FPPS, FPSHBT, FPSPH

Unit Head, National Confirmatory Center for
Hemoglobinopathies, Institute of Human Genetics,
University of the Philippines Manila
Chairman, Thalassemia Working Group, Philippine
Society of Hematology and Blood Transfusion
Vice President, Philippine Society of Pediatric
Hematology
Consultant, Amang Rodriguez Medical Center

Charito Carbon-De Los Santos, MD, FPPS, FPAPP, FPSCCM

Member, Philippine Foundation for Lung Health,
Research and Development, Inc.
Training Officer, Philippine Heart Center

Phorenice D. Francisco, MD, FPCP, DPSAAI

Visiting Consultant, Ospital ng Maynila Medical Center
Member, Philippine Society of Allergy, Asthma and
Immunology

Juan Javier T. Lichauco, MD, FPCP, FPRA

President, Philippine Rheumatology Association
Chair, Department of Medicine, St. Luke's Medical
Center – Quezon City

Jonathan Go Lim, MD

Consultant, Chong Hua Hospital-Fuente and
Mandaue
Vice chair, Department of Pediatrics, Chong Hua
Hospital-Cebu
Section Head, Infectious Diseases, Department of
Pediatrics, Chong Hua Hospital-Cebu
Member, Research Program Subcommittee,
Pediatric Infectious Disease Society of the
Philippines

Imelda M. Mateo, MD, MBAH, FPCP, FPCCP

Medical Center Chief II, Amang Rodriguez Memorial
Medical Center
Visiting Consultant, Lung Center of the Philippines

Albert L. Rafanan, MD, FPCCP, FPCP, FCCP, FASSM, FPSSM

Chairman, Department of Medicine, Chong Hua
Hospital, Cebu City
Chair, Critical Care Committee, Chong Hua
Hospital, Cebu City
Board Member, Philippine College of Chest
Physicians
Founding Chairman, Philippine Board of Sleep
Medicine

Jeah Alvarez Sabillo, RTRP, RN, MMHoA

Founding Officer and President, Respiratory
Therapy Academy of Critical Care,
Philippines(RTACCP)
Respiratory Therapist III (Supervisor) Division of
Pulmonary and Critical Care Medicine, Philippine
Heart Center

Rowena Marie T. Samares, MD, FPAFP, FPSHPM

Consultant, Silliman University Medical Center
Vice President, Philippine Academy of Family
Physicians Negros Oriental Chapter
Executive Secretary, Philippine Society of Hospice
and Palliative Medicine

Reynaldo G. San Luis III, MD

Consultant, Pope John Paul II Hospital and Medical
Center
Treasurer, Philippine Society of General Internal
Medicine
Member, Philippine Medical Association

Shirley Paras-Whisenhunt, PhD, RN

Visiting Professor, Philippine Christian University
Department Manager, Nursing Research Systems
Management and Accreditation, St. Luke's Medical
Center – Bonifacio Global City

Consensus Panel for Vaccines and Prophylactic Interventions

Ma. Delta B. San Antonio-Aguilar, MD, FPPS, FPIDSP

Active Consultant, Metro Davao Medical and Research Center

Member, Philippine Foundation for Vaccination

Medical Specialist III, Southern Philippines Medical Center

Maria Rhona G. Bergantin, MD, MSc, FPCP, FPSMID

Associate Professor, University of Santo Tomas

Consultant Staff and Training Officer, Section of Infectious Diseases, Department of Medicine, University of Santo Tomas Hospital

Sybil Lizanne R. Bravo, RPh, MD, MSc, FPOGS, FPIDSOG

Chief, Section of OB GYN Infectious Diseases, Philippine General Hospital

President, Philippine Infectious Diseases Society for OB GYN

Fatima Ignacio Gimenez, MD, FPPS, FPIDSP

Training Officer, Pediatric Infectious Disease Section, Philippine Children's Medical Center

Vice President, Pediatric Infectious Disease Society of the Philippines

Chairman, Immunization Committee, Philippine Pediatric Society

PRO, Philippine Foundation for Vaccination

Katrina G. Gomez, MD, MPH

Primary Care Physician, Innovations for Community Health (Kalinga Health)

Edmyr M. Macabulos, MD, MPH, FPCOM

Associate Professor 2, St. Luke's Medical Center College of Medicine

Occupational Health Physician, Pampanga's Best Inc.

Nenacia Ranali Nirena P. Mendoza, MD, FPAFP

Residency Training Officer, Healthway Family Clinic

Member, Committee on Continuing Medical Education, Philippine Academy of Family Physicians

Diana Alcantara-Payawal, MD, DTMH, FPCP, FPSG, FPSFDE

Vice President, Philippine College of Physicians

Professor II, Our Lady of Fatima University

Chair, Department of Internal Medicine, Our Lady of Fatima University Medical Center

Ruth S. Punzalan, MD, MPH, FPAFP

Municipal Health Officer, Tanza, Cavite

Carmela Rosanne A. Remotigue, MD, FPCP

Assistant Professor, Cebu Institute of Medicine

Active Staff, Department of Internal Medicine, Cebu Velez General Hospital

Kim Patrick Salvador Tejano, MD

Medical Officer IV, Disease Prevention and Control Bureau, Department of Health

Felicia Racquel L. Salvador-Tayag, MD

Vice President, Philippine Society of Allergy, Asthma & Immunology

Associate Editor, Philippine Journal of Allergy, Asthma & Immunology

Consultant, Marikina Valley Medical Center

Edmyr M. Macabulos, MD, MPH, FPCOM

Associate Professor II, St. Luke's Medical Center College of Medicine-WHQM

Immediate Past National President, Philippine College of Occupational Medicine

Member, Specialty Board, Philippine College of Occupational Medicine

Gian Carlo Sy Torres, PhD, MAN, RN

Faculty Member, University of Santo Tomas College of Nursing

Chair, Department of Public Affairs, Philippine Nurses Association

Julie Christie Gutierrez Visperas, MD, MHPEd, FPCP, FPCCP

Assistant Professor 3, Faculty of Medicine and Surgery, University of Santo Tomas

Consultant, Center for Respiratory Medicine, University of Santo Tomas Hospital

Consensus Panel for Adjunctive Therapies, Non-Pharmacologic and Infection Control Interventions

Camille Angelica P. Banzon, MD, FPCEM

Secretary General, Philippine College of Emergency Medicine

Medical Specialist I, Ospital ng Makati

Gerard Belimac, MD, MPH

Medical Specialist IV and Program Manager, National HIV, AIDS, and STI Prevention and Control Program, Department of Health

Officer-in-Charge, Policy Division, Systems Integration, Disease Prevention and Control Bureau, Department of Health

Regina P. Berba, MD, MSc

Associate Professor IV, University of the Philippine Manila – Philippine General Hospital

Division Chief, Division of Adult Medicine, Department of Medicine, Philippine General Hospital

Chair, Hospital Infection Control Unit, Philippine General Hospital

Section Head, Infectious Disease Section, The Medical City

Elmer D. Bondoc, RN, MN, PhD

Director, University Research Office, Holy Angel University

Vivien Fe F. Fadrilan-Camacho, MD, MPH, FPAFP

Associate Professor, Department of Environmental and Occupational Health, College of Public Health, University of the Philippines Manila

Faculty, Department of Family and Community Medicine, Philippine General Hospital

Chair, Commission on Specialty Board of Examiners in Family Medicine, Philippine Academy of Family Physicians

Maria Tricia Subido Cariño, MD, FPPS, FPIDSP

Member, Philippine Medical Association

Head, Section of Pediatric Infectious Diseases, Cardinal Santos Medical Center

Medical Specialist III, Research Institute for Tropical Medicine

Member, Antimicrobial Stewardship Committee, University of Perpetual Help Dalta Medical Center

Victoria Isla-Ching, RN, MGM-ESP

Manager, Patient Safety Department, The Medical City

Treasurer and Faculty, Philippine Hospital Infection Control Society

Adviser, Philippine Hospital Infection Control Nurses Association

Anthony F. Cortez, MD

Municipal Health Officer, RHU Bambang, Nueva Vizcaya

President, Association of Municipal Health Officers of the Philippines – Nueva Vizcaya

Radela Yvonne Ramos-Cortes, MD, FPCP, FPSAAI

Associate Professor, College of Medicine, University of St. La Salle, Bacolod City

Consultant, Riverside Medical Center, Bacolod City

Anna Sofia Victoria Tamayo Salazar-Fajardo, MD, MBAH, DPCOM

Medical Director, Perpetual Help Medical Center – Biñan, Laguna

Assistant Section Head, Section of Occupational & Environmental Medicine, Department of Family & Community Medicine, Perpetual Help Medical Center – Biñan, Laguna

Dominga Calalang-Gomez, RN

Founding President and Council of Adviser, Philippine Hospital Infection Control Nurses Association

Founding Member, Past President and Council Adviser, Philippine Hospital Infection Control Society

Joan Mae M. Oliveros, MD, FPAFP

Faculty/Research & COPC Coordinator, Department of Family Medicine, Silliman Medical Center

University Physician, Silliman University

Roberto A. Razo II, MD, FPSP, FPCP

Consultant, Section of Adult Medicine, Department of Medicine, De La Salle University Medical Center

Assistant Professor, De La Salle Medical and Health Sciences Institute

Secretary, Philippine Society of General Internal Medicine

Maria Sonia Salamat, MD, MPH, FPCP, FPSMID

Medical Specialist III, Philippine General Hospital Clinical

Associate Professor, College of Medicine, University of the Philippines

Training Officer, Philippine General Hospital Division of Infectious Diseases

TECHNICAL COORDINATORS

Howell Henrian G Bayona, MSc, CSP-PASP
Speech Pathologist, The Medical City Speech
Pathologist, St. Luke's Medical Center-Global
City

Marie Carmela Lapitan, MD, MS, FPUA, FPCS

Research Professor, Institute of Clinical
Epidemiology, University of the Philippines-
National Institutes of Health

Clinical Professor, Department of Surgery,
College of Medicine, University of the
Philippines

Christopher G. Manalo, MD, FPCEM

Medical Specialist II, Department of Emergency
Medicine, Philippine General Hospital,
University of the Philippines Manila

Carol Stephanie C. Tan-Lim, MD, MScCE, DPPS, DPSAAI

Diplomate, Philippine Pediatric Society and
Philippine Society of Allergy, Asthma and
Immunology

Maria Teresa S. Tolosa, MD, FPDS, D Clin Epi

Assistant Professor, St. Lukes' Medical Center
College of Medicine – WHQM

Epidemiology Consultant, Research and
Biotechnology Group, St. Luke's Medical
Center

Assistant Professor, UERMMMC College of
Medicine

Dan Louie Renz P. Tating, MS(cand), RN

MS Candidate, Department of Clinical
Epidemiology, College of Medicine, University
of the Philippines

TECHNICAL ASSISTANTS

Vaneza Leah A. Espino, MD, DPPS, DPAPP

Attending Physician, Division of Pediatric
Pulmonology, Department of Pediatrics, UP-
PGH

Active Consultant, University of Perpetual Help
Medical Center-Las Pinas & Manilamed
Medical Center Manila

Myzelle Anne J. Infantado, PTRP, MSc (cand.)

Physical Therapy Consultant, ManilaMed Child
Development

Natasha Ann R. Esteban-Ipac, MD, FPPS,
DPSAMS

Medical Specialist III, Division of Adolescent
Medicine, Department of Pediatrics, UP-PGH

Julianne Keane M. Pascual, MD

Project Development Officer III, National
Institutes of Health Clinical Research Center

Michelle Cristine B. Miranda, M.D.

National Projects Coordinator, Division of
Hematology-Oncology, Department of
Pediatrics, UP-PGH

April P. Padua-Zamora, MD, DPPS

Medical Officer IV, Division of Pediatric
Gastroenterology, Hepatology and Nutrition,
Department of Pediatrics, UP-PGH

EVIDENCE REVIEW EXPERTS

Giselle Anne Q. Adajar, MD

University Researcher I, Newborn Screening Reference Center, National Institutes of Health, UP-Manila

Carl Lawrence C. Arenos, MD

Medical Officer III, Department of Medicine, UP-PGH

Carla Marie L. Asis, MD, DPPS

Medical Officer IV, Division of Pediatric Rheumatology, Department of Pediatrics, UP-PGH

Mica Olivine Bastillo-Casillan, MD, DPPS

Medical Specialist I, Pasig City Children's Hospital

Eva I. Bautista, MD, MSc, FPPS

Assistant Professor B, College of Medicine, Far Eastern University-Nicanor Reyes Medical Foundation

Mario Lorenzo L. Bautista, MD

Research Physician, National Institutes of Health

Liza Marie P. Bejemino, MD, DPPS, DPSNM

Neonatal ICU Consultant, Dr. Rafael S. Tumbukon Memorial Hospital, Kalibo, Aklan

John Jefferson V. Besa, MD

Medical Officer III, Philippine General Hospital

Julian M. A. Buban

Medical Intern, Philippine General Hospital

Aldrich Ivan Lois D. Burog, MD, MSc (cand.)

Evidence Reviewer, Living CPG for COVID-19

Ian Theodore Cabaluna, RPh, MD, GDip (Epi)

Medical officer, Wellbridge Health, Inc.,

Carmen Carina G. Cabrera, MD, FPCP, DPSED

Active Consultant, Providence Hospital, Inc.

Timothy Hudson David C. Carandang, MD

Fides Roxanne M. Castor, MD, DPPS

Medical Specialist II and Training Officer, Division of Pediatric Emergency Medicine, Department of Pediatrics, UP-PGH

Mary Anne J. Roldan-Castor, MD, FPPS, FPSAAI

Associate Professor 7, Department of Pediatrics, University of the Philippines College of Medicine

Mary Christine R. Castro, MD, MSc

Executive Director, Nutrition Center of the Philippines

Ina Cathrina R. Chiu, RMT, MD

Medical Officer III, Department of Pediatrics, UP-PGH

Erika Crisostomo, MD

Medical Officer III, Department of Pediatrics, UP-PGH

Marie Gene D. Cruz, MD, DPCP

Internal Medicine Hospitalist, St. Luke's Medical Center - Global City

Patricia Maria Gregoria M. Cuaño, MD, DPCP

Institute of Clinical Epidemiology, National Institutes of Health, University of the Philippines

Dianne Marie Delid-Legaspi, MD, DPPS

Medical Officer IV, Division of Pediatric Rheumatology, Department of Pediatrics, UP-PGH

Lea Roselle O. De Castro, MD, DPCP

Internal Medicine Hospitalist, Chinese General Hospital Internal Medicine Hospitalist, St. Luke's Hospital - BGC

Namnama P. Villarta-De Dios, MD, MSc, DPPS

Medical Specialist II, Department of Pediatrics, Amang Rodriguez Memorial Medical Center

Belen Lardizabal Dofitas, MD, FPDS, MSc Clin Epid

Associate Professor IV, College of Medicine, University of the Philippines
Vice-Chair for Research, Department of Dermatology, Philippine General Hospital

Valentin C. Dones III, PhD

Research Supervisor, Center for Health Research and Movement Science, University of Santo Tomas

Louie Dy, MD

Medical Intern, Philippine General Hospital

Anton Elepano, MD

Medical Officer III, Department of Medicine, UP-PGH

Bryan F. Elvambuena, MD

Medical Officer III, Department of Medicine, UP-PGH

Mar Christopher F. Epetia, MD, DPPS

Medical Officer IV, Division of Pediatric Rheumatology, Department of Pediatrics, UP-PGH

Adrian Ronald A. Espino, MD

Medical Officer IV, DOH Region XII Mlang Distict Hospital

Emmanuel P. Estrella, MD, MSc, FPOA

Deputy Director, Institute of Clinical Epidemiology, National Institutes of Health, UP-Manila

Gina Antonina S. Eubanas, MD, FPDS, D Clin Epi

Assistant Professor, Department of Clinical Research, St. Luke's Medical Center, College of Medicine - WHQM
Head, Section of Dermatology, Department of Internal Medicine, Qualimed - Daniel Mercado Medical Center

Antonio L. Faltado Jr., MD, FPCP, FPSEDM

Chair, Research Ethics Board, Lipa Medix Medical Center
Head, Diabetes Center, Lipa Medix Medical Center
Coordinator, Diabetes Center, Mary Mediatrix Medical Center

Anna Maria Vida P. Garcia, RPh, D Clin Epi

Clinical Scientist, ClinChoice Inc.

Rowena F. Genuino, MD, MSc

Professor, Department of Anatomy, College of Medicine, University of the Philippines
Consultant Dermatologist, Manila Doctors Hospital and Makati Medical Center

Germana Emerita V. Gregorio, MD, PhD, FPPS, FPSPGHN

Consultant, Department of Pediatrics, University of the Philippines Manila College of Medicine Philippine General Hospital

Daniel Y. Guevara, MD, FPCP, DPSN

Active Medical Staff, ManilaMed Medical Center Manila; Premier Medical Center Paranaque

Myzelle Anne J. Infantado, PTRP, MSc (cand.)

PT Consultant, ManilaMed Child Development Enrichment Center and New Beginnings Center for Child Development

Racquel Ibanez, MD, FPCP, FPCCP, D Clin Epi

Medical Specialist, Lung Center of the Philippines and National Center for Mental Health

Marquis Von Angelo Syquiao Go Joson, MD

Medical Officer III, Department of Pediatrics, University of the Philippines - Philippine General Hospital

Anna Angelica Macalalad Josue, MD, FPCP, DPSEDM, MSc (cand)

Medical Specialist, Taguig Pateros District Hospital

Marie Carmela Lapitan, MD, MS, FPUA, FPCS

Research Professor, Institute of Clinical Epidemiology, University of the Philippines-National Institutes of Health
Clinical Professor, Department of Surgery, College of Medicine, University of the Philippines

Furqaan I. Lim, MD, DPPS

Medical Officer IV, Bataan General Hospital and Medical Center

Maria Cristina H. Lozada, MD, DPPS, DPAPP

Medical Specialist III, Division of Pediatric Pulmonology, Department of Pediatrics, UP-PGH

Patricia Marie M. Lusica, MD-MBA

Medical Officer III, Department of Medicine, UP-PGH

Christopher G. Manalo, MD, FPBEM

Medical Specialist II, Department of Emergency Medicine, Philippine General Hospital, University of the Philippines Manila

Mark Jason C. Milan, RN, MD

Medical Officer III, Department of Pediatrics,
UP-PGH

Isabella S. Ocampo, MD, DPPS

Diplomate, Philippine Pediatric Society

Katherine Ruth Oracion-Relato, MD, DPCP

Assistant Professorial Lecturer IV, Pamantasan
ng Lungsod ng Maynila

Marc Andrew O. Perez, MD, DPPS, DPSN, DPNSP

Consultant, Department of Pediatrics, Region 1
Medical Center

Assistant Professor I, College of Medicine,
Lyceum-Northwestern University

Jofermarie O. Pineda RN, MD

Medical Officer IV. Pediatric Infectious Disease
and Tropical Medicine Department, San Lazaro
Hospital

Patricia Pauline M. Remalante-Rayco, MD, FPCP, FPRA

Associate professor, De La Salle Medical and
Health Sciences Institute

Evelyn O. Salido, MD, MSc, FPCP, FPRA

Chair, Division of Rheumatology, Philippine
General Hospital

Christdianzen Grace P. Saroca, MD, DPCP

Adult Cardiology Fellow, St. Luke's Medical
Center- Global City

Aina Fe R. Salem, RN, MD

Medical Officer III, Department of Pediatrics,
UP-PGH

Maria Cristina Z. San Jose, MD, FPNA

Division Chair, Section of Adult Neurology,
Department of Neurosciences, University of the
Philippines - Philippine General Hospital
Associate Professor II, College of Medicine,
University of the Philippines

Maria Vanessa V. Sulit, BSN, RN, MSc (Clinical Epidemiology)

Faculty Member & Workshop Coordinator,
Asia-Pacific Center for Evidence-Based
Healthcare, Inc.

Frangelo Conrad Tampus, MD

Junior Consultant, Lorenzo D Zayco District
Hospital

Carol Stephanie C. Tan-Lim, MD, MScCE, DPPS, DPSAAI

Diplomate, Philippine Pediatric Society and
Philippine Society of Allergy, Asthma and
Immunology

Issa Rufina S. Tang, MD, FPCP, DPSMID

Medical Specialist I, Philippine Orthopedic
Center

Jose Carlo B. Valencia, MD, FPCP, FPSMID

Medical Specialist III, Department of Medicine,
Cagayan Valley Medical Center

Grazielle S. Verzosa, MD, DPPS

Consultant, Department of Pediatrics, East
Avenue Medical Center

Maria Philina P. Villamor, MD, FPCP, FPCCP

Medical Specialist IV, Vicente Sotto Memorial
Medical Center

Cary Amiel G. Villanueva, MD, DPCP

Technical Writer, Institute of Clinical
Epidemiology, National Institutes of Health, UP-
Manila

Paoline Nicole P. Villanueva, RMT, MD**Mithi Kalayaan S. Zamora, MD, FPCP, DPCCP,**

Visiting Consultant, Diliman Doctors Hospital,
ManilaMed Medical Center

FACILITATORS

Screening and Diagnosis

Sandra T. Torres, MD, MScCE, FPCP, FPRA

Active Staff, Section of Rheumatology, Cardinal Santos Medical Center

Lia Aileen M. Palileo-Villanueva, MD, MSc

Associate Professor, College of Medicine, University of the Philippines- Manila

Vice Chair for Undergraduate Training, Department of Medicine, UP-Philippine General Hospital

Treatment

Diana R. Tamondong-Lachica, MD, FPCP

Associate Professor, Division of Adult Medicine, Philippine General Hospital

Vaccines and Prophylactic Interventions

Maria Asuncion A. Silvestre, MD, FPSNB

President, Kalusugan ng Mag-Ina, Inc. (KMI)

Member, Independent Review Group, Early Essential Newborn Care (EENC), WHO, WPRO

Critical Care and Respiratory Management

Bernadette Heizel Manapat-Reyes, MD, MHPed, FPCP, FPRA

Professor, Department of Medicine, College of Medicine, University of the Philippines

Head, Medical Education Unit, College of Medicine, University of the Philippines

Adjunct Interventions and Non-Pharmacologic Interventions

Carlo Irwin Panelo, MD, MA

Professor, Department of Clinical Epidemiology

College of Medicine

University of the Philippines Manila

Carol Stephanie C. Tan-Lim, MD, MScCE, DPPS, DPSAAI

Diplomate, Philippine Pediatric Society and Philippine Society of Allergy, Asthma and Immunology

COPY EDITORS AND TECHNICAL WRITERS

Joyce Anne Ceria-Pereña, RPh, MPM

Christine A. Dator, MD

Kate D. Dunlao, RPh

Ena Lauren F. Farillas, MD

Gian Carlo L. Infante, MD

Mikarla M. Lubat, RND

Maria Regina Naval-Jacinto, MD

Aubrey Melody R. Rocimo, MD

OVERSIGHT COMMITTEE

Dante D. Morales, MD *Chair*

Antonio L. Dans, MD, MSc

Angela A. Du, MD

Joy T. Sanchez, MD

Maria Vanessa Sulit, RN, MSc

Camilo Te Jr, MD

PROJECT STAFF

Project Managers

Melissa A. Dator, MD-MBA, DPPS, DPSN, DPNSP

Attending Physician, Division of Pediatric Nephrology, Department of Pediatrics, Philippine General Hospital

Associate Active Staff, Department of Pediatrics, Makati Medical Center

Reiner Lorenzo J. Tamayo, RN

Research, Policy, and Communications Manager, Innovations for Community Health

Dan Louie Renz P. Tating, MS(cand), RN

MS Candidate, Department of Clinical Epidemiology, College of Medicine, University of the Philippines

TECHNICAL ADVISER

Ian Theodore Cabaluna, RPh, MD, GDip (Epi)

ADMINISTRATIVE STAFF

Rhea M. Anwar

Pauline Bianca N. Aquino

Maria Eleanor L. Candelaria, MPH, RN

Kate D. Dunlao, RPh

Ma. Caroline C. Peralta

Carlos Diego A. Rozul

Rhonaliza P. Tapic

Maria Pamela Tagle

Lailanie Ann C. Tejuco

Appendix B. Decisions of the Oversight Committee – Review of Conflict of Interest

CPG Panel on COVID – 19 Critical Care		
Jeah Alvarez Sabillo	Manageable A	Founder, President & Author/Lead Respiratory Therapy Academy for Critical Care Phils., Respiratory therapist – PHC
Pauline Convozar	Manageable A	Board member – Asian Society of Emergency, Immediate Past Pres. of PCEM
Shirley Whisenhunt	Manageable A	Chairman – CPD of PNA
Juan Javier Lichauco	Manageable A	PI for clinical trials: Infliximab, Tocilizumab, Secukinumab, Cenerimod, Evobrutinib, Speaker for Roche - Cytokine storm webinar, President – PRA, with stocks in Urology Center of the Phils.
Phorenice Francisco	Manageable A	PCP, PSAAI
Rey San Luis	Manageable A	Treasurer- PSGIM, VP – Las Pinas Medical Society, with stocks in Pope John Paul II Hosp.
Reynaldo de Castro	Manageable A	Unit Head Center for Hemoglobinopathies- UP NIH, Chairman- Thalassemia TWG, VP- Phil. Society of Pediatric Hematology, PI for Hemoglobinopathies study
Joseph Adrian Buensalido	Manageable A	Chair TWG – Sepsis CPG, Training officer & Asso. Prof. – UP-PGH, PEDER TF – MMC, Board member – PSMID, Infection Prevention & Control Chair – AHMC, author of haemophilus Influenza Infections in Medscape, stocks in New Sinai MDI Hosp & Makati Med Ctr
Rowena Marie Samares	Manageable A	TWG- Hospice Palliative DOH, Board member – Phil. Society of Hospice, VP – Phil. Acad. Of family Physicians, stocks in COL)
Jonathan Lim	Manageable A	vaccine lectures -Pfizer, Sanofi, MSD, clinical trial PI & Sub I for multinational studies, Treasurer-PPS
Albert Rafanan	Disqualified for (1) inhaled nitric oxide, (2) high flow nasal cannula, (3) hyperbaric oxygen therapy; Manageable A for the other questions	VP- Baga Inc.: med. & lab. Equipment; President- RED Medical Foundation, board member PCCP
Charito delos Santos	Manageable B	>0.1% stocks Binangonan Lakeview Hosp. & Rizal Doctors)
Maaliddin Biruar	Manageable B	Senior Medical Director – Parexel
Imelda Mateo	Manageable B	Amang Rodriguez Medical Center Chief – DOH, Expert Panel CLCPG – Diagnosis & Screening, VP- PCCP

CPG Panel on COVID – 19 Adjunctive Therapies, Non-Pharmaceutical Interventions, Infection Control		
Anthony Cortez	Manageable A	MHO- Bambang, Nueva Vizcaya
Maria Tricia Cariño	Manageable A	PI in collaboration with CDC, ongoing research on COVID serologic assessment, PI for vaccine studies, lecturer for MSD vaccines, Med Sp. III RITM)
Joan Oliveros	Manageable A	member-PAFP
Dominga Gomez	Manageable A	former DOH TWG- infection control, Faculty – DOH training of trainors for infection prevention control, Council of advisers-PHICNA & PHICS
Victoria Isla Ching	Manageable A	Adviser-PHICNA, VP & Faculty - PHIC
Elmer Bondoc	Manageable A	Governor – PNA, Trustee-PREO
Radela Yvonne Ramos-Cortes	Manageable A	Intermed Marketing for Immunomax
Roberto Razo II	Manageable A	Secretary-PSGIM
Maria Sonia Salamat	Manageable A	Co-investigator – Solidarity Trial, TWG Pneumonia CPG, lecturer for COVID-19 diagnostics & therapeutics, Med Sp. III- UP-PGH, member Board of Councils-PSMID)
Vivien Fadrilan-Camacho	Manageable A	Research on host responses in severely COVID, Asst. Prof – CPH-UP, member-HPAAC, technical consultant for 3 private companies – not related to topic
Regina Berba	Manageable A	DOH & WHO studies on COVID, Infection control lectures, Med. Sp. III- UP PGH, Chair- Pharmacovigilance Comm. of PCP, board member-PSMID, stocks in TMC
Anna Sofia Victoria Fajardo	Manageable A	Speakers bureau- PMA, PCOM, PFAP and other companies, Medical Director- PHMC Binan, National Sec.-PCOM
Camille Angelica Banzon	Manageable B	Angelicare Pharmacy owned by spouse
Gerard Belimac	Manageable B	DOH Division Chief- DPCB

CPG Panel on COVID – 19 Treatment		
Ma. Elinore Concha	Manageable A	editorial staff-Filipino Family Physician, Chief training officer SPMC, board member- Foundation for Family Med educators
Ronald Panaligan	Manageable A	PCCP chairman-council on diagnostics & therapeutics, member National Ethics Committee-PCHRD, Medical manager- UAP
Faith Joan Mesa-Gaerlan	Manageable A	Editor in chief- Phil. Journal of E-Med, Training officer- Emergency med UP-PGH, Founding member PCEM, Technical – CPG on Sepsis, Board MAG
Rowena Roselle Blanco-Santos	Manageable A	National board of Directors – PCOM
Mary Ann Bunyi	Manageable A	Deputy Executive Director-PCMC, President- Pediatric Infectious Society of the Phils., Advisory Committee – Pediatrics
Erwin De Mesa	Manageable A	President PIDSOG, active consultant QMMC & QC Gen., Treasurer POGS
Sarah Makalinaw	Manageable A	member PIDSP journal editorial board, consultant IDS-Rizal Medical Center, Technical- CPG on children
Karl Evans Henson	Manageable A	board member -PSMID, Associate Prof. UPCM Division of Infectious Dis., TWG- CPG on sepsis
Rommel Punongbayan	Manageable A	member- TWG COVID 19 Living CPG, BOT PSGIM, PCOM, Advisory board on respiratory division-GSK
Leila Ferrer	Manageable A	stocks in Allied Care Experts Medical Center
Maria Encarnita Limpin	Manageable A	Editorial Board – Respirology, Med Specialist & TRB- PHC, President-PCP, Board member- Phil Society of Critical Care Med, Exec. Director- ASH & FCAP, Chair – PCPCNCD
Jenifer Otadoy-Agustin	Manageable A	Editor in Chief-PJAAI, Webinar on Covid-19 vaccines & adverse reactions, PI for Novartis trials, Associate Clinical Professor & Training Officer-UP-PGH, stocks in Sta. Rosa Hospital
Sarah May Flores	Manageable B	Supervising Health Program Officer-DOH
Marysia Stella Recto	Manageable B	Technical Adviser-Philippine FDA
Anna Melissa Guererro	Manageable B	Unit Head HTA, Division chief- DOH pharma division

CPG Panel on COVID – 19 Vaccines and Prophylaxis		
Ruth Punzalan	Manageable A	(MHO- Tanza Cavite, Executive VP-Assn of MHO in the Phils., President-AMHOP Cavite, VP-AMHOP
Fatima Gimenez	Manageable A	Editorial board- Pediatric Infectious Dis. Journal, stocks in VRPMC, TMC, Visayas Av. Hosp., Chair-Immunization Comm. PPS, PRO-Phil. Foundation for Vaccination
Ranali Mendoza	Manageable A	(Section Editor-The Filipino Family Physician, Training Officer-Healthway
Julie Christie Visperas	Manageable A	Board member- Society of Physiologists
Edmyr Macabulos	Manageable A	Research on Awareness & Attitudes to improve Vaccination in Children, Adviser- Rotary Club, President- Phil. Society of HPN Central Luzon, National Pres. PCOM 2019
Diana Payawal	Manageable A	(APCOLIS Study APASL Task Force 2020, Asst Editor Hepatology Int'l, VP-PCP 2021, Sec.- Board of Regent PCP 2020, Paper on Management of Hepatocellular CA in Covid, Hepa Int'l 2020
Gian Carlos Torres	Manageable A	Editorial- Social Inequities in Covid, Pres- PNA NCR, Board of Director PNA Zone 1
Sybil Lizanne Bravo	Manageable A	Author of handbook of COVID in Pregnancy, member- ACTA Med Journal UPCM, Chief & Training Officer- OB GYN Infectious Dis. & Training Officer PGH & MDH, President- PIDSOG, Member- POGS Committee on Immunization Women
Felicia Racquel Tayag	Manageable A	Associate Editor- Phil. Journal Allergology, Asthma & Immuno, speaker for Menarini, VP-PSAAI, shares <0.1% in Marikina Valley Med Ctr & Metro Antipolo Hosp., TMC
Carmela Remotigue	Manageable A	VP-PCP Central Visayas, PSGIM
Rhona Bergantin	Manageable A	Chair-Publications Comm. PSMID, National Immunization Committee representing academe
Katrina Gomez	Manageable A	CP of LCPG1, Covid Essentials in Family Medicine, Med Officer III-Makati Health Dept.
Ma. Delta San Antonio-Aguilar	Manageable A	PI- Asian Foundation for Tropical Med, Speaker for Pfizer pneumococcal and meningococcal vaccines, speaker for MSD for rotatec and varicella vaccine, Vice Chair for patient COVID services at SPMC, Adviser on Ad Board- Pfizer, >0.1% shares in United Doctors Davao – not yet operational) Manageable B for questions pertaining to Coronavac, mix and match & booster dose (lecture on Sinovac
Kim Patrick Tejano	Manageable B	DOH MO IV Program Manager Nat'l Immunization Program, formulates policies for Immunization program

CPG Panel on COVID – 19 Screening and Diagnosis			
John Andrew Camposano	Manageable A		ADHOC Comm Chair- PIDSP TWG interim CPG for COVID 19
Jane Lardizabal-Bunyi	Manageable A		Med Officer III-Justice Jose Abad Santos Hosp, Treasurer-PAFP
Virginia Delos Reyes	Manageable A		authorship on papers on COVID 19 in LCPG, PI of sponsored clinical trials not directly related to CPG, Med Sp. II-LCP, Exec board-PCCP & ACCP, stocks in Metro North Hosp
Marilyn Puyot	Manageable A		Med Sp. I-Pasig City Gen, stocks in ACE Med Center, Pasig Doctors, Tricity Med Center
Vernon Serafico	Manageable A		Board Member-PSGIM
Mary Ann Lansang	Manageable A		Edit Board Biomed Central, CPG-Covid Diagnostics PCHRD – no honorarium, PCHRD grant for study on SARS COV 2 antibodies, WHO Expert advisory panel on Health science & technology, Board-FACE Inc., PSMID Forum on COVID Diagnostics, stocks in TMC, Chief of Party-USAID
Florido Atibagos	Manageable A		Med Sp. 3-Jose B Lingad
Aretha Gacutan Liwag	Manageable A		grant for mRNA vaccine from Tigermed Inc., Med Sp III- West Visayas State Univ Med Center, Board-UPV
Anelyn Reyes	Manageable A		Board member-PPS, stocks in City of Gen. Trias Doctors Med Center, Dasmariñas City Med Center, South Imus Specialist Hosp.
Fatima Johanna Santos-Ocampo	Manageable A; Manageable B for question genomics		contributor-Immunomodulators as Therap Intervention COVID, Adviser-PHILPOPI, Founder-APSI, Head Immunodeficiency Council-PSAAI, TWG-PCHRD DOSH on identifying local immunobiologicals for emerging infections, stocks in Makati Med; Technical evaluator – COVID 19 immuno-genomic surveillance program
Clemencia Bondoc	Manageable B		Board of Director- Metro Iloilo Hosp, Board-PCOM Iloilo, Past Pres.-AMHOP, MHO- Zarraga, Iloilo
Arthur Dessi Roman	Manageable B		Med Sp. 2-RITM)

Appendix C. General Search Strategy for COVID-19

SEARCH STRATEGY FOR COVID-19:

((("COVID-19" [Supplementary Concept] OR "COVID-19 diagnostic testing" [Supplementary Concept] OR "COVID-19 drug treatment" [Supplementary Concept] OR "COVID-19 serotherapy" [Supplementary Concept] OR "COVID-19 vaccine" [Supplementary Concept] OR "severe acute respiratory syndrome coronavirus 2" [Supplementary Concept] OR "2019-nCoV" OR "2019nCoV" OR "cov 2" OR "Covid-19" OR "sars coronavirus 2" OR "sars cov 2" OR "SARS-CoV-2" OR "severe acute respiratory syndrome coronavirus 2" OR "coronavirus 2" OR "COVID 19" OR "COVID-19" OR "2019 ncov" OR "2019nCoV" OR "corona virus disease 2019" OR "cov2" OR "COVID-19" OR "COVID19" OR "nCov 2019" OR "nCoV" OR "new corona virus" OR "new coronaviruses" OR "novel corona virus" OR "novel coronaviruses" OR "SARS Coronavirus 2" OR "SARS2" OR "SARS-COV-2" OR "Severe Acute Respiratory Syndrome Coronavirus 2") OR ((19[tiab] OR 2019[tiab] OR "2019-nCoV" OR "Beijing" OR "China" OR "Covid-19" OR epidem*[tiab] OR epidemic* OR epidemy OR new[tiab] OR "novel"[tiab] OR "outbreak" OR pandem* OR "SARS-CoV-2" OR "Shanghai" OR "Wuhan") AND ("Coronavirus Infections"[Mesh] OR "coronavirus"[MeSH Terms] OR coronavirus*[all] OR corona-virus*[all] OR cov[tiab] OR pneumonia-virus*[tiab]))) AND 2019/12/1:3000/12/31[PDAT])

SEARCH FILTER FOR RANDOMIZED CONTROLLED TRIALS:

(randomized controlled trial [pt] OR controlled clinical trial [pt] OR randomized [tiab] OR placebo [tiab] OR drug therapy [sh] OR randomly [tiab] OR trial [tiab] OR groups [tiab]) NOT (animals [mh] NOT humans [mh])

SEARCH FILTER FOR SYSTEMATIC REVIEWS AND META-ANALYSES:

((systematic review[ti] OR systematic literature review[ti] OR systematic scoping review[ti] OR systematic narrative review[ti] OR systematic qualitative review[ti] OR systematic evidence review[ti] OR systematic quantitative review[ti] OR systematic meta-review[ti] OR systematic critical review[ti] OR systematic mixed studies review[ti] OR systematic mapping review[ti] OR systematic cochrane review[ti] OR systematic search and review[ti] OR systematic integrative review[ti]) NOT comment[pt] NOT (protocol[ti] OR protocols[ti]) NOT MEDLINE [subset]) OR (Cochrane Database Syst Rev[ta] AND review[pt]) OR systematic review[pt]

Appendix D. Breakdown of Consensus Meetings for Phase 2 of the Philippine COVID-19 Living Clinical Practice Guidelines

CONSENSUS PANEL MEETINGS	
Critical Care (3)	October 26, 2021
	December 1, 2021
	January 3, 2022
Adjunctive Therapies, Non-Pharmaceutical Interventions, Infection Control (4)	October 29, 2021
	November 5, 2021
	December 3, 2021
	December 21, 2021
Treatment (7)	October 12, 2021
	October 21, 2021
	October 28, 2021
	November 8, 2021
	November 18, 2021
	December 6, 2021
	December 20, 2021
Vaccines and Prophylaxis (6)	October 21, 2021
	October 22, 2021
	October 28, 2021
	November 4, 2021
	December 2, 2021
	December 27, 2021
Screening and Diagnosis (5)	November 11, 2021
	November 22, 2021
	November 29, 2021
	December 13, 2021
	December 17, 2021

Appendix E. AGREE Reporting Checklist

This checklist is intended to guide the reporting of clinical practice guidelines.

CHECKLIST ITEM AND DESCRIPTION	REPORTING CRITERIA	Page #
DOMAIN 1: SCOPE AND PURPOSE		
1. OBJECTIVES <i>Report the overall objective(s) of the guideline. The expected health benefits from the guideline are to be specific to the clinical problem or health topic.</i>	<input checked="" type="checkbox"/> Health intent(s) (i.e., prevention, screening, diagnosis, treatment, etc.) <input type="checkbox"/> Expected benefit(s) or outcome(s) <input checked="" type="checkbox"/> Target(s) (e.g., patient population, society)	1
2. QUESTIONS <i>Report the health question(s) covered by the guideline, particularly for the key recommendations.</i>	<input checked="" type="checkbox"/> Target population <input checked="" type="checkbox"/> Intervention(s) or exposure(s) <input checked="" type="checkbox"/> Comparisons (if appropriate) <input checked="" type="checkbox"/> Outcome(s) <input checked="" type="checkbox"/> Health care setting or context	See relevant sections
3. POPULATION <i>Describe the population (i.e., patients, public, etc.) to whom the guideline is meant to apply.</i>	<input checked="" type="checkbox"/> Target population, sex and age <input checked="" type="checkbox"/> Clinical condition (if relevant) <input checked="" type="checkbox"/> Severity/stage of disease (if relevant) <input checked="" type="checkbox"/> Comorbidities (if relevant) <input type="checkbox"/> Excluded populations (if relevant)	1-2
DOMAIN 2: STAKEHOLDER INVOLVEMENT		
4. GROUP MEMBERSHIP <i>Report all individuals who were involved in the development process. This may include members of the steering group, the research team involved in selecting and reviewing/rating the evidence and individuals involved in formulating the final recommendations.</i>	<input checked="" type="checkbox"/> Name of participant <input type="checkbox"/> Discipline/content expertise (e.g., neurosurgeon, methodologist) <input checked="" type="checkbox"/> Institution (e.g., St. Peter's hospital) <input type="checkbox"/> Geographical location (e.g., Seattle, WA) <input checked="" type="checkbox"/> A description of the member's role in the guideline development group	227-238
5. TARGET POPULATION PREFERENCES AND VIEWS <i>Report how the views and preferences of the target population were sought/considered and what the resulting outcomes were.</i>	<input checked="" type="checkbox"/> Statement of type of strategy used to capture patients'/publics' views and preferences (e.g., participation in the guideline development group, literature review of values and preferences) <input checked="" type="checkbox"/> Methods by which preferences and views were sought (e.g., evidence from literature, surveys, focus groups) <input type="checkbox"/> Outcomes/information gathered on patient/public information <input checked="" type="checkbox"/> How the information gathered was used to inform the guideline development process and/or formation of the recommendations	11
6. TARGET USERS <i>Report the target (or intended) users of the guideline.</i>	<input checked="" type="checkbox"/> The intended guideline audience (e.g. specialists, family physicians, patients, clinical or institutional leaders/administrators) <input checked="" type="checkbox"/> How the guideline may be used by its target audience (e.g., to inform clinical decisions, to inform policy, to inform standards of care)	2
DOMAIN 3: RIGOUR OF DEVELOPMENT		

7. SEARCH METHODS <i>Report details of the strategy used to search for evidence.</i>	<input checked="" type="checkbox"/> Named electronic database(s) or evidence source(s) where the search was performed (e.g., MEDLINE, EMBASE, PsychINFO, CINAHL) <input checked="" type="checkbox"/> Time periods searched (e.g., January 1, 2004 to March 31, 2008) <input checked="" type="checkbox"/> Search terms used (e.g., text words, indexing terms, subheadings) <input checked="" type="checkbox"/> Full search strategy included (e.g., possibly located in appendix)	8, 287
8. EVIDENCE SELECTION CRITERIA <i>Report the criteria used to select (i.e., include and exclude) the evidence. Provide rationale, where appropriate.</i>	<input checked="" type="checkbox"/> Target population (patient, public, etc.) characteristics <input checked="" type="checkbox"/> Study design <input checked="" type="checkbox"/> Comparisons (if relevant) <input checked="" type="checkbox"/> Outcomes <input checked="" type="checkbox"/> Language (if relevant) <input type="checkbox"/> Context (if relevant)	8
9. STRENGTHS & LIMITATIONS OF THE EVIDENCE <i>Describe the strengths and limitations of the evidence. Consider from the perspective of the individual studies and the body of evidence aggregated across all the studies. Tools exist that can facilitate the reporting of this concept.</i>	<input checked="" type="checkbox"/> Study design(s) included in body of evidence <input checked="" type="checkbox"/> Study methodology limitations (sampling, blinding, allocation concealment, analytical methods) <input checked="" type="checkbox"/> Appropriateness/relevance of primary and secondary outcomes considered <input checked="" type="checkbox"/> Consistency of results across studies <input checked="" type="checkbox"/> Direction of results across studies <input checked="" type="checkbox"/> Magnitude of benefit versus magnitude of harm <input checked="" type="checkbox"/> Applicability to practice context	8
10. FORMULATION OF RECOMMENDATIONS <i>Describe the methods used to formulate the recommendations and how final decisions were reached. Specify any areas of disagreement and the methods used to resolve them.</i>	<input checked="" type="checkbox"/> Recommendation development process (e.g., steps used in modified Delphi technique, voting procedures that were considered) <input checked="" type="checkbox"/> Outcomes of the recommendation development process (e.g., extent to which consensus was reached using modified Delphi technique, outcome of voting procedures) <input checked="" type="checkbox"/> How the process influenced the recommendations (e.g., results of Delphi technique influence final recommendation, alignment with recommendations and the final vote)	9-15
11. CONSIDERATION OF BENEFITS AND HARMS <i>Report the health benefits, side effects, and risks that were considered when formulating the recommendations.</i>	<input checked="" type="checkbox"/> Supporting data and report of benefits <input checked="" type="checkbox"/> Supporting data and report of harms/side effects/risks <input checked="" type="checkbox"/> Reporting of the balance/trade-off between benefits and harms/side effects/risks <input checked="" type="checkbox"/> Recommendations reflect considerations of both benefits and harms/side effects/risks	9

12. LINK BETWEEN RECOMMENDATIONS AND EVIDENCE <i>Describe the explicit link between the recommendations and the evidence on which they are based.</i>	<input checked="" type="checkbox"/> How the guideline development group linked and used the evidence to inform recommendations <input checked="" type="checkbox"/> Link between each recommendation and key evidence (text description and/or reference list) <input checked="" type="checkbox"/> Link between recommendations and evidence summaries and/or evidence tables in the results section of the guideline	See relevant sections
13. EXTERNAL REVIEW <i>Report the methodology used to conduct the external review.</i>	<input checked="" type="checkbox"/> Purpose and intent of the external review (e.g., to improve quality, gather feedback on draft recommendations, assess applicability and feasibility, disseminate evidence) <input checked="" type="checkbox"/> Methods taken to undertake the external review (e.g., rating scale, open-ended questions) <input checked="" type="checkbox"/> Description of the external reviewers (e.g., number, type of reviewers, affiliations) <input checked="" type="checkbox"/> Outcomes/information gathered from the external review (e.g., summary of key findings) <input checked="" type="checkbox"/> How the information gathered was used to inform the guideline development process and/or formation of the recommendations (e.g., guideline panel considered results of review in forming final recommendations)	16
14. UPDATING PROCEDURE <i>Describe the procedure for updating the guideline.</i>	<input checked="" type="checkbox"/> A statement that the guideline will be updated <input checked="" type="checkbox"/> Explicit time interval or explicit criteria to guide decisions about when an update will occur <input checked="" type="checkbox"/> Methodology for the updating procedure	17
DOMAIN 4: CLARITY OF PRESENTATION		
15. SPECIFIC AND UNAMBIGUOUS RECOMMENDATIONS <i>Describe which options are appropriate in which situations and in which population groups, as informed by the body of evidence.</i>	<input checked="" type="checkbox"/> A statement of the recommended action <input checked="" type="checkbox"/> Intent or purpose of the recommended action (e.g., to improve quality of life, to decrease side effects) <input checked="" type="checkbox"/> Relevant population (e.g., patients, public) <input checked="" type="checkbox"/> Caveats or qualifying statements, if relevant (e.g., patients or conditions for whom the recommendations would not apply) <input checked="" type="checkbox"/> If there is uncertainty about the best care option(s), the uncertainty should be stated in the guideline	See relevant sections
16. MANAGEMENT OPTIONS <i>Describe the different options for managing the condition or health issue.</i>	<input checked="" type="checkbox"/> Description of management options <input checked="" type="checkbox"/> Population or clinical situation most appropriate to each option	See relevant sections

17. IDENTIFIABLE KEY RECOMMENDATIONS <i>Present the key recommendations so that they are easy to identify.</i>	<input checked="" type="checkbox"/> Recommendations in a summarized box, typed in bold, underlined, or presented as flow charts or algorithms <input checked="" type="checkbox"/> Specific recommendations grouped together in one section	See relevant sections and Executive Summary (ix)
DOMAIN 5: APPLICABILITY		
18. FACILITATORS AND BARRIERS TO APPLICATION <i>Describe the facilitators and barriers to the guideline's application.</i>	<input checked="" type="checkbox"/> Types of facilitators and barriers that were considered <input checked="" type="checkbox"/> Methods by which information regarding the facilitators and barriers to implementing recommendations were sought (e.g., feedback from key stakeholders, pilot testing of guidelines before widespread implementation) <input checked="" type="checkbox"/> Information/description of the types of facilitators and barriers that emerged from the inquiry (e.g., practitioners have the skills to deliver the recommended care, sufficient equipment is not available to ensure all eligible members of the population receive mammography) <input checked="" type="checkbox"/> How the information influenced the guideline development process and/or formation of the recommendations	225
19. IMPLEMENTATION ADVICE/TOOLS <i>Provide advice and/or tools on how the recommendations can be applied in practice.</i>	<input checked="" type="checkbox"/> Additional materials to support the implementation of the guideline in practice. For example: <ul style="list-style-type: none"> ○ Guideline summary documents ○ Links to check lists, algorithms ○ Links to how-to manuals ○ Solutions linked to barrier analysis (see Item 18) ○ Tools to capitalize on guideline facilitators (see Item 18) ○ Outcome of pilot test and lessons learned 	222-223
20. RESOURCE IMPLICATIONS <i>Describe any potential resource implications of applying the recommendations.</i>	<input checked="" type="checkbox"/> Types of cost information that were considered (e.g., economic evaluations, drug acquisition costs) <input checked="" type="checkbox"/> Methods by which the cost information was sought (e.g., a health economist was part of the guideline development panel, use of health technology assessments for specific drugs, etc.) <input checked="" type="checkbox"/> Information/description of the cost information that emerged from the inquiry (e.g., specific drug acquisition costs per treatment course)	223 and other relevant sections

	<input checked="" type="checkbox"/> How the information gathered was used to inform the guideline development process and/or formation of the recommendations	
21. MONITORING/ AUDITING CRITERIA <i>Provide monitoring and/or auditing criteria to measure the application of guideline recommendations.</i>	<input checked="" type="checkbox"/> Criteria to assess guideline implementation or adherence to recommendations <input checked="" type="checkbox"/> Criteria for assessing impact of implementing the recommendations <input checked="" type="checkbox"/> Advice on the frequency and interval of measurement <input checked="" type="checkbox"/> Operational definitions of how the criteria should be measured	16, 260
DOMAIN 6: EDITORIAL INDEPENDENCE		
22. FUNDING BODY <i>Report the funding body's influence on the content of the guideline.</i>	<input checked="" type="checkbox"/> The name of the funding body or source of funding (or explicit statement of no funding) <input checked="" type="checkbox"/> A statement that the funding body did not influence the content of the guideline	17
23. COMPETING INTERESTS <i>Provide an explicit statement that all group members have declared whether they have any competing interests.</i>	<input checked="" type="checkbox"/> Types of competing interests considered <input checked="" type="checkbox"/> Methods by which potential competing interests were sought <input checked="" type="checkbox"/> A description of the competing interests <input checked="" type="checkbox"/> How the competing interests influenced the guideline process and development of recommendations	17-18, 239

From: Brouwers MC, Kerkvliet K, Spithoff K, on behalf of the AGREE Next Steps Consortium. The AGREE Reporting Checklist: a tool to improve reporting of clinical practice guidelines. *BMJ* 2016;352:i1152. doi: 10.1136/bmj.i1152.

For more information about the AGREE Reporting Checklist, please visit the AGREE Enterprise website at <http://www.agreetrust.org>.

