

Solutions to Ireland, Rosen “A Classical Introduction to Modern Number Theory”

Richard Ganaye

September 7, 2019

1 Chapter 1

Ex 1.1 *Let a and b be nonzero integers. We can find nonzero integers q and r such that $a = qb + r$ where $0 \leq r < b$. Prove that $(a, b) = (b, r)$.*

Proof. Notation : if a, b are integers in \mathbb{Z} , $a \wedge b$ is the non negative greatest common divisor of a, b , the generator in $\mathbb{N} = \{0, 1, 2, \dots\}$ of the ideal $(a, b) = a\mathbb{Z} + b\mathbb{Z}$.

Let $d \in \mathbb{Z}$.

- If $d \mid a, d \mid b$, then $d \mid a - qb = r$, so $d \mid b, d \mid r$.
- If $d \mid b, d \mid r$, then $d \mid qb + r = a$, so $d \mid a, d \mid b$.

$$\forall d \in \mathbb{Z}, (d \mid b, d \mid r) \iff (d \mid a, d \mid b).$$

If $a = bq + r$, the set of common divisors of a, b is equal to the set of common divisors of b, r .

As $a \wedge b$ is the smallest positive element of this set, so is $b \wedge r$, we conclude that $a \wedge b = b \wedge r$. □

Ex 1.2 *If $r \neq 0$, we can find q_1 and r_1 such that $b = q_1r + r_1$, with $0 \leq r_1 < r$. Show that $(a, b) = (r, r_1)$. This process can be repeated. Show that it must end in finitely many steps. Show that the last nonzero remainder must equal (a, b) . The process looks like*

$$\begin{array}{ll} a = bq + r, & 0 \leq r < b \\ b = q_1r + r_1, & 0 \leq r_1 < r \\ r = q_2r_1 + r_2, & 0 \leq r_2 < r_1 \\ \vdots & \\ r_{k-1} = q_{k+1}r_k + r_{k+1}, & 0 \leq r_{k+1} < r_k \\ r_k = q_{k+2}r_{k+1} & \end{array}$$

Then $r_{k+1} = (a, b)$. This process of finding (a, b) is known as the Euclidian algorithm.

Proof. The Euclidian division of b by r gives $b = q_1r + r_1, 0 \leq r_1 < r$. The result of exercise 1.1 applied to the couple (b, r) shows that

$$b \wedge r = r \wedge r_1.$$

Let $N \in \mathbb{N}$. While the remainders $r_i, i \leq N$, are not equal to 0, we can define the sequences $(q_i), (r_i)$ by

$$r_{-1} = a, r_0 = b, \quad r_{i-1} = q_{i+1}r_i + r_{i+1}, \quad 0 \leq r_{i+1} < r_i \quad 0 \leq i \leq N$$

If no $r_i, i \in \mathbb{N}$, is equal to 0, we can continue this construction indefinitely. So we obtain a strictly decreasing sequence $(r_i)_{i \in \mathbb{N}}$ of positive numbers : it is impossible. Therefore, there exists an index k such as $r_{k+2} = 0$, this is the end of the algorithm.

$$\begin{array}{ll} a = bq + r, & 0 \leq r < b \\ b = q_1r + r_1, & 0 \leq r_1 < r \\ r = q_2r_1 + r_2, & 0 \leq r_2 < r_1 \\ \vdots & \\ r_{k-1} = q_{k+1}r_k + r_{k+1}, & 0 \leq r_{k+1} < r_k \\ r_k = q_{k+2}r_{k+1}, & r_{k+2} = 0 \end{array}$$

From exercise 1, $r_{i-1} \wedge r_i = r_i \wedge r_{i+1}, 0 \leq i \leq k$, so

$$a \wedge b = b \wedge r = \dots = r_k \wedge r_{k+1} = r_{k+1} \wedge r_{k+2} = r_{k+1} \wedge 0 = r_{k+1}.$$

The last non zero remainder is the gcd of a, b . □

Ex 1.3 Calculate (187, 221), (6188, 4709), (314, 159).

Proof. With direct instructions in Python, we obtain :

```
>>> a, b = 187, 221
>>> print("q = ", a//b); a, b = b, a%b; print(a,b)
q = 0
221 187
>>> print("q = ", a // b); a, b = b, a%b; print(a,b)
q = 1
187 34
>>> print("q = ", a // b); a, b = b, a%b; print(a,b)
q = 5
34 17
>>> print("q = ", a // b); a,b = b, a%b; print(a,b)
q = 2
17 0
```

This gives the equalities

$$\begin{aligned} 187 &= 0 \times 221 + 187 \\ 221 &= 1 \times 187 + 34 \\ 187 &= 5 \times 34 + 17 \\ 34 &= 2 \times 17 + 0 \end{aligned}$$

So $187 \wedge 221 = 17$.

With the same instructions, we obtain

$$6188 = 1 \times 4709 + 1479$$

$$4709 = 3 \times 1479 + 272$$

$$1479 = 5 \times 272 + 119$$

$$272 = 2 \times 119 + 34$$

$$119 = 3 \times 34 + 17$$

$$34 = 2 \times 17 + 0$$

$$6188 \wedge 4709 = 17.$$

Finally

$$314 = 1 \times 159 + 155$$

$$159 = 1 \times 155 + 4$$

$$155 = 38 \times 4 + 3$$

$$4 = 1 \times 3 + 1$$

$$3 = 3 \times 1 + 0$$

$$314 \wedge 159 = 1.$$

The Python script which gives the gcd is very concise :

```
def gcd(a,b):  
    a, b = abs(a), abs(b)  
    while b != 0:  
        a, b = b, a % b  
    return a
```

□

Ex 1.4 Let $d = (a, b)$. Show how one can use the Euclidean algorithm to find numbers m and n such that $am + bn = d$. (Hint: In Exercise 2 we have that $d = r_{k+1}$. Express r_{k+1} in terms of r_k and r_{k+1} , then in terms of r_{k-1} and r_{k-2} , etc.).

Proof. With a slight modification of the notations of exercise 2, we note the Euclid's algorithm under the form

$$r_0 = a, r_1 = b, \quad r_i = r_{i+1}q_{i+1} + r_{i+2}, \quad 0 < r_{i+2} < r_{i+1}, \quad 0 \leq i < k, \quad r_k = q_{k+1}r_{k+1}, \quad r_{k+2} = 0$$

We show by induction on i ($i \leq k+1$) the proposition

$$P(i) : \exists (m_i, n_i) \in \mathbb{Z} \times \mathbb{Z}, \quad r_i = am_i + bn_i.$$

• $r_0 = a = 1.a + 0.b$. Define $m_0 = 1, n_0 = 0$. We obtain $r_0 = am_0 + bn_0$, then $P(0)$ is true.

$r_1 = b = 0.a + 1.b$. Define $m_1 = 0, n_1 = 1$. We obtain $r_1 = am_1 + bn_1$, then $P(1)$ is true.

- Suppose for $0 \leq i < k$ the induction hypothesis $P(i)$ et $P(i+1)$:

$$\begin{aligned} r_i &= am_i + bn_i, & m_i, n_i &\in \mathbb{Z}, \\ r_{i+1} &= am_{i+1} + bn_{i+1}, & m_{i+1}, n_{i+1} &\in \mathbb{Z}. \end{aligned}$$

Then $r_{i+2} = r_i - r_{i+1}q_{i+1} = a(m_i - q_{i+1}m_{i+1}) + b(n_i - q_{i+1}n_{i+1})$.

If we define $m_{i+1} = m_i - q_{i+1}m_{i+1}$, $n_{i+1} = n_i - q_{i+1}n_{i+1}$, we obtain $r_{i+2} = am_{i+2} + bn_{i+2}$, $m_{i+2}, n_{i+2} \in \mathbb{Z}$, so $P(i+2)$.

- The conclusion is that $P(i)$ is true for all $i, 0 \leq i \leq k+1$, in particular $r_{k+1} = am_{k+1} + bn_{k+1}$, that is

$$a \wedge b = d = am + bn,$$

where $m = m_{k+1}, n = n_{k+1} \in \mathbb{Z}$. □

Ex 1.5 Find m and n for the pairs a and b given in Ex 1.3

Proof. From exercises 1.3, 1.4, we know that the sequences $(r_i), (m_i), (n_i)$ are given by

$$\begin{aligned} r_0 &= a, r_1 = b \\ m_0 &= 1, m_1 = 0 \\ n_0 &= 0, n_1 = 1 \end{aligned}$$

and for all $i < k$,

$$\begin{aligned} r_{i+2} &= r_i - q_{i+1}r_{i+1} \\ m_{i+2} &= m_i - q_{i+1}m_{i+1} \\ n_{i+2} &= n_i - q_{i+1}n_{i+1} \end{aligned}$$

and for all i

$$r_i = m_i a + n_i b.$$

This gives the direct instructions in Python :

```
>>> a,b = 187, 221
>>> r0,r1,m0,m1,n0,n1 = a,b,1,0,0,1
>>> q = r0//r1;
>>> q = r0//r1; r0,r1,m0,m1,n0,n1 = r1, r0 -q*r1,m1, m0 -q*m1, n1, n0 - q*n1
>>> print(r0,r1,m0,m1,n0,n1)
221 187 0 1 1 0
>>> q = r0//r1; r0,r1,m0,m1,n0,n1 = r1, r0 -q*r1,m1, m0 -q*m1, n1, n0 - q*n1
>>> print(r0,r1,m0,m1,n0,n1)
187 34 1 -1 0 1
>>> q = r0//r1; r0,r1,m0,m1,n0,n1 = r1, r0 -q*r1,m1, m0 -q*m1, n1, n0 - q*n1
>>> print(r0,r1,m0,m1,n0,n1)
34 17 -1 6 1 -5
>>> q = r0//r1; r0,r1,m0,m1,n0,n1 = r1, r0 -q*r1,m1, m0 -q*m1, n1, n0 - q*n1
>>> print(r0,r1,m0,m1,n0,n1)
17 0 6 -13 -5 11
```

So

$$17 = 187 \wedge 221 = 6 \times 187 - 5 \times 221.$$

Similarly

$$17 = 6188 \wedge 4709 = 121 \times 6188 - 159 \times 4709.$$

$$1 = 314 \wedge 159 = -40 \times 314 + 79 \times 159.$$

We obtain the same results with the following Python script :

```
def bezout(a,b):
    """input  : entiers a,b
       output : tuple (x,y,d),
       (x,y) solution de ax+by = d, d = pgcd(a,b)
    """
    (r0,r1)=(a,b)
    (u0,v0) = (1,0)
    (u1,v1) = (0,1)
    while r1 != 0:
        q = r0 // r1
        (r2,u2,v2) = (r0 - q*r1,u0 - q*u1,v0 - q*v1)
        (r0,r1) = (r1,r2)
        (u0,u1) = (u1,u2)
        (v0,v1) = (v1,v2)
    return (u0,v0,r0)
```

□

Ex 1.6 Let $a, b, c \in \mathbb{Z}$. Show that the equation $ax + by = c$ has solutions in integers iff $(a, b) | c$.

Proof. Let $d = a \wedge b$.

- If $ax + by = c, x, y \in \mathbb{Z}$, as $d | a, d | b, d | ax + by = c$.
- Reciprocally, if $d | c$, then $c = dc', c' \in \mathbb{Z}$.

From Prop. 1.3.2., $d\mathbb{Z} = a\mathbb{Z} + b\mathbb{Z}$, so $d = au + bv, u, v \in \mathbb{Z}$, and $c = dc' = a(c'u) + b(c'v) = ax + by$, where $x = c'u, y = c'v$ are integers.

Conclusion :

$$\exists (x, y) \in \mathbb{Z} \times \mathbb{Z}, ax + by = c \iff a \wedge b | c.$$

□

Ex 1.7 Let $d = (a, b)$ and $a = da'$ and $b = db'$. Show that $(a', b') = 1$.

Proof. Suppose $d \neq 0$ (if $d = 0$, then $a = b = 0$, and a', b' are any numbers in \mathbb{Z} and the result may be false, so we must suppose $d \neq 0$).

As $d = am + bn, m, n \in \mathbb{Z}, d = d(a'm + b'n)$, so $1 = a'm + b'n$, which proves $a' \wedge b' = 1$.
conclusion : if $d = a \wedge b \neq 0$, and $a = da', b = db'$, then $a' \wedge b' = 1$.

□

Ex. 1.8 Let x_0 and y_0 be a solution to $ax + by = c$. Show that all solutions have the form $x = x_0 + t(b/d)$, $y = y_0 - t(a/d)$, where $d = (a, b)$ and $t \in \mathbb{Z}$.

Proof. Suppose $a \neq 0, b \neq 0$.

Let x_0 and y_0 be a solution to $ax + by = c$.

If (x, y) is any solution of the same equation,

$$\begin{aligned} ax + by &= c \\ ax_0 + by_0 &= c, \end{aligned}$$

then

$$a(x - x_0) = -b(y - y_0),$$

so

$$\frac{a}{d}(x - x_0) = -\frac{b}{d}(y - y_0).$$

Let $a' = a/d, b' = b/d$: from ex. 1.7, we know that $a' \wedge b' = 1$.

As $a'(x - x_0) = -b'(y - y_0)$, $b' \mid a'(x - x_0)$, and $b' \wedge a' = 1$, so (Gauss' Lemma : prop. 1.1.1) $b' \mid x - x_0$.

There exists $t \in \mathbb{Z}$ such that $x - x_0 = tb'$. Then $a'tb' = -b'(y - y_0)$. As $b \neq 0, b' \neq 0$, so $a't = -(y - y_0)$:

$$\begin{aligned} x &= x_0 + t(b/d) \\ y &= y_0 - t(a/d) \end{aligned}$$

Reciprocally, $a(x_0 + t(b/d)) + b(y_0 - t(a/d)) = ax_0 + by_0 = c$.

Conclusion : if $a \neq 0, b \neq 0$, and $ax_0 + by_0 = c$,

$$ax + by = c \iff \exists t \in \mathbb{Z}, x = x_0 + t(b/d), y = y_0 - t(a/d).$$

□

Ex. 1.9 Suppose that $u, v \in \mathbb{Z}$ and that $(u, v) = 1$. If $u \mid n$ and $v \mid n$, show that $uv \mid n$. Show that this is false if $(u, v) \neq 1$.

Proof. As $u \mid n$, $n = uq, q \in \mathbb{Z}$, so $v \mid n = uq$, and $v \wedge u = 1$, so (Gauss' lemma : prop. 1.1.1), $v \mid q : q = vl, l \in \mathbb{Z}$, and $n = uvl : uv \mid n$.

If the case $u \wedge v \neq 1$, we give the counterexample $6 \mid 18, 9 \mid 18$, but $6 \times 9 \nmid 18$. □

Ex. 1.10 Suppose that $(u, v) = 1$. Show that $(u + v, u - v)$ is either 1 or 2.

Proof. Let $d = (u + v) \wedge (u - v)$. Then $d \mid u + v, d \mid u - v$, so $d \mid 2u = (u + v) + (u - v)$ and $d \mid 2v = (u + v) - (u - v)$. So $d \mid (2u) \wedge (2v) = 2(u \wedge v) = 2$. As $d \geq 0$, $d = 1$ or $d = 2$. □

Ex. 1.11 Show that $(a, a + k) \mid k$.

Proof. Let $d = a \wedge (a + k)$. As $d \mid a, d \mid (a + k)$, $d \mid k = (a + k) - a$.

Conclusion : $a \wedge (a + k) \mid k$. □

Ex. 1.12 Suppose that we take several copies of a regular polygon and try to fit them evenly about a common vertex. Prove that the only possibilities are six equilateral triangles, four squares, and three hexagons.

Proof. Let n be the number of sides of the regular polygon, m the number of sides starting from a summit in the lattice, α the measure of the exterior angle, β the measure of the interior angle (in radians) ($\alpha + \beta = \pi$).

Then $\alpha = 2\pi/n$, $\beta = \pi - 2\pi/n$.

$m\beta = 2\pi$, $m(\pi - 2\pi/n) = 2\pi$, $m(1 - 2/n) = 2$, so

$$\frac{1}{m} + \frac{1}{n} = \frac{1}{2}, \quad m > 0, n > 0. \quad (1)$$

As this equation is symmetric in m, n , we may suppose first $m \leq n$.

In this case $1/m \geq 1/n$, so $2/n \leq 1/2 : n \geq 4$.

If $n > 6$, $1/n < 1/6$, $1/m = 1/2 - 1/n > 1/2 - 1/6 = 1/3$, so $m < 3$, $m \leq 2 : m = 1$ or $m = 2$.

If $m = 1$, $n < 0$: it is impossible. If $m = 2$, $1/n = 0$: also impossible. Therefore $n \leq 6 : 4 \leq n \leq 5$. If $n = 4$, $m = 4$. if $n = 5$, $n = 10/3$: impossible. if $n = 6$, $m = 3$. Using the symetry, the set of solutions of (1) is

$$S = \{(3, 6), (6, 3), (4, 4)\},$$

corresponding with the usual lattices composed of equilateral triangles, squares or hexagons. \square

Ex. 1.13 Let $n_1, n_2, \dots, n_s \in \mathbb{Z}$. Define the greatest common divisor d of n_1, n_2, \dots, n_s and prove that there exist integers m_1, m_2, \dots, m_s such that $n_1m_1 + n_2m_2 + \dots + n_sm_s = d$.

Proof. Let $n_1, n_2, \dots, n_s \in \mathbb{Z}$. The ideal of \mathbb{Z} , $(n_1, \dots, n_s) = n_1\mathbb{Z} + \dots + n_s\mathbb{Z}$ is principal, so there exists an unique $d \in \mathbb{Z}, d \geq 0$ such that

$$n_1\mathbb{Z} + \dots + n_s\mathbb{Z} = d\mathbb{Z} \quad (d \geq 0).$$

We define

$$d = \gcd(n_1, \dots, n_s) \iff n_1\mathbb{Z} + \dots + n_s\mathbb{Z} = d\mathbb{Z} \text{ and } d \geq 0. \quad (2)$$

The characterization of the gcd is

$$d = \gcd(n_1, \dots, n_s) \iff$$

$$(i) \ d \geq 0 \quad (3)$$

$$(ii) \ d \mid n_1, \dots, d \mid n_s \quad (4)$$

$$(iii) \ \forall \delta \in \mathbb{Z}, (\delta \mid n_1, \dots, \delta \mid n_s) \Rightarrow \delta \mid d \quad (5)$$

(\Rightarrow) Indeed, if we suppose (1), then $d \geq 0$, and $n_1 = n_1 \cdot 1 + n_2 \cdot 0 + \dots + n_s \cdot 0 \in n_1\mathbb{Z} + \dots + n_s\mathbb{Z} = d\mathbb{Z}$, so $d \mid n_1$. Similarly $d \mid n_i, 1 \leq i \leq s$ so (i)(ii) are true. if $\delta \mid n_i, 1 \leq i \leq s$, as $d = n_1m_1 + \dots + n_sm_s, m_1, \dots, m_s \in \mathbb{Z}$, then $\delta \mid d$.

(\Leftarrow) Suppose that d verify (i)(ii)(iii). From (ii), we see that $n_i\mathbb{Z} \subset d\mathbb{Z}, i = 1, \dots, s$, so $n_1\mathbb{Z} + \dots + n_s\mathbb{Z} \subset d\mathbb{Z}$.

As \mathbb{Z} is a principal ring, there exists $\delta \geq 0$ such that $n_1\mathbb{Z} + \cdots + n_s\mathbb{Z} = \delta\mathbb{Z}$. $n_i \in n_1\mathbb{Z} + \cdots + n_s\mathbb{Z}$ so $n_i \in \delta\mathbb{Z}$, $i = 1, \dots, s$: $\delta \mid n_1, \dots, \delta \mid n_s$. From (iii), we deduce $\delta \mid d$. As $\delta\mathbb{Z} \subset d\mathbb{Z}$, $d \mid \delta$, with $d \geq 0, \delta \geq 0$. Consequently, $d = \delta$ and $n_1\mathbb{Z} + \cdots + n_s\mathbb{Z} = d\mathbb{Z}, d \geq 0$, so $d = \gcd(n_1, \dots, n_s)$.

At last, as $n_1\mathbb{Z} + \cdots + n_s\mathbb{Z} = d\mathbb{Z}$, there exist integers m_1, m_2, \dots, m_s such that $n_1m_1 + n_2m_2 + \cdots + n_sm_s = d$. \square

Ex. 1.14 Discuss the solvability of $a_1x_1 + a_2x_2 + \cdots + a_rx_r = c$ in integers. (Hint: Use Exercise 13 to extend the reasoning behind Exercise 6.)

Proof. Let $a_1, a_2, \dots, a_r \in \mathbb{Z}$.

Note $\gcd(a_1, a_2, \dots, a_r) = a_1 \wedge a_2 \wedge \cdots \wedge a_r$. The following result generalizes Ex. 6 :

$$\exists(x_1, x_2, \dots, x_r) \in \mathbb{Z}^r, a_1x_1 + a_2x_2 + \cdots + a_rx_r = c \iff a_1 \wedge a_2 \wedge \cdots \wedge a_r \mid c.$$

Let $d = a_1 \wedge a_2 \wedge \cdots \wedge a_r$.

- If $a_1x_1 + a_2x_2 + \cdots + a_rx_r = c$, as $d \mid a_1, \dots, d \mid a_r, d \mid a_1x_1 + a_2x_2 + \cdots + a_rx_r = c$.
- Reciprocally, if $d \mid c$, then $c = dc', c' \in \mathbb{Z}$.

As $d\mathbb{Z} = a_1\mathbb{Z} + a_2\mathbb{Z} + \cdots + a_r\mathbb{Z}$, so $d = a_1m_1 + a_2m_2 + \cdots + a_rm_r, m_1, m_2, \dots, m_r \in \mathbb{Z}$. $c = dc' = a_1(m_1c') + \cdots + a_r(m_rc') = a_1x_1 + \cdots + a_rx_r$, where $x_i = m_ic', i = 1, 2, \dots, r$. \square

Ex. 1.15 Prove that $a \in \mathbb{Z}$ is the square of another integer iff $\text{ord}_p(a)$ is even for all primes p . Give a generalization.

Proof. Suppose $a = b^2, b \in \mathbb{Z}$. Then $\text{ord}_p(a) = 2 \text{ord}_p(b)$ is even for all primes p .

Reciprocally, suppose that $\text{ord}_p(a)$ is even for all primes p . We must also suppose $a > 0$. Let $a = \prod_p p^{a(p)}$ the decomposition of a in primes. As $a(p)$ is even, $a(p) = 2b(p)$ for an integer $b(p)$ function of the prime p . Let $b = \prod_p p^{b(p)}$. Then $a = b^2$.

With a similar demonstration, we obtain the following generalization for each integer $a \in \mathbb{Z}, a > 0$:

$$a = b^n \text{ for an integer } b \in \mathbb{Z} \text{ iff } n \mid \text{ord}_p(a) \text{ for all primes } p. \quad \square$$

Ex. 1.16 If $(u, v) = 1$ and $uv = a^2$, show that both u and v are squares.

Proof. Here $u, v \in \mathbb{N}$, where $\mathbb{N} = \{0, 1, 2, \dots\}$.

For all primes p such that $p \mid u$, $\text{ord}_p(u) + \text{ord}_p(v) = 2 \text{ord}_p(a)$. As $u \wedge v = 1$ and $p \mid u, p \nmid v$, so $\text{ord}_p(v) = 0$. Consequently, $\text{ord}_p(u)$ is even for all prime p such that $p \mid u$. From Exercise 1.15, we can conclude that u is a square. Similarly, v is a square. \square

Ex. 1.17 Prove that the square root of 2 is irrational, i.e., that there is no rational number $r = a/b$ such that $r^2 = 2$.

Proof. Suppose there exists $r \in \mathbb{Q}, r > 0$ such that $r^2 = 2$. Then $r = a/b, a \in \mathbb{N}^*, b \in \mathbb{N}^*$. With $d = a \wedge b, a = da', b = db', a' \wedge b' = 1$, so $r = a'/b', a' \wedge b' = 1$, so we may suppose $r = a/b, a > 0, b > 0, a \wedge b = 1$ and $a^2 = 2b^2$.

a^2 is even, then a is even (indeed, if a is odd, $a = 2k + 1, k \in \mathbb{Z}, a^2 = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1$ is odd).

So $a = 2A, A \in \mathbb{N}$, then $4A^2 = 2b^2, 2A^2 = b^2$.

With the same reasoning, b^2 is even, then b is even : $b = 2B, B \in \mathbb{N}$. $2 \mid a, 2 \mid b$, $2 \mid a \wedge b$, in contradiction with $a \wedge b = 1$.

Conclusion : $\sqrt{2}$ is irrational. \square

Ex. 1.18 Prove that $\sqrt[n]{m}$ is irrational if m is not the n -th power of an integer.

Proof. Here $m \in \mathbb{N}$.

Suppose that $r = \sqrt[n]{m} \in \mathbb{Q}$. As $r \geq 0$, $r = a/b, a \geq 0, b > 0, a \wedge b = 1$, and $r^n = m$, so $a^n = mb^n$.

For all primes p , $n \operatorname{ord}_p(a) = \operatorname{ord}_p(m) + n \operatorname{ord}_p(b)$, so $n \mid \operatorname{ord}_p(m)$.

From Ex. 1.15, we conclude that m is a n -th power.

Conclusion : if $m \geq 0$ is not the n -th power of an integer, $\sqrt[n]{m}$ is irrational. \square

Ex. 1.19 Define the least common multiple of two integers a and b to be an integer m such that $a \mid m, b \mid m$, and m divides every common multiple of a and b . Show that such an m exists. It is determined up to sign. We shall denote it by $[a, b]$.

Proof. As $a\mathbb{Z} \cap b\mathbb{Z}$ is an ideal of \mathbb{Z} , and \mathbb{Z} is a principal ideal domain, there exists a unique $m \geq 0$ such that $a\mathbb{Z} \cap b\mathbb{Z} = m\mathbb{Z}$. So by definition,

$$m = [a, b] \iff a\mathbb{Z} \cap b\mathbb{Z} = m\mathbb{Z} \text{ and } m \geq 0.$$

We may note also $[a, b] = a \vee b$.

characterization of lcm :

$$\begin{aligned} m = a \vee b &\iff \\ (i) \quad &m \geq 0 \\ (ii) \quad &a \mid m, b \mid m \\ (iii) \quad &\forall \mu \in \mathbb{Z}, (a \mid \mu, b \mid \mu) \Rightarrow m \mid \mu \end{aligned}$$

(\Rightarrow) By definition, $m \geq 0$. $m \in m\mathbb{Z} = a\mathbb{Z} \cap b\mathbb{Z}$, so $a \mid m$ and $b \mid m$: (ii) is verified. If $\mu \in \mathbb{Z}$ is such that $a \mid \mu, b \mid \mu$, then $\mu \in a\mathbb{Z} \cap b\mathbb{Z} = m\mathbb{Z}$, so $m \mid \mu$: (iii) is true.

(\Leftarrow) Suppose that m verifies (i),(ii),(iii). Let m' such that $a\mathbb{Z} \cap b\mathbb{Z} = m'\mathbb{Z}, m' \geq 0$. We show that $m = m'$.

As $m' \in a\mathbb{Z} \cap b\mathbb{Z}$, $a \mid m', b \mid m'$, so from (iii) $m \mid m'$. From (ii), we see that $m \in a\mathbb{Z} \cap b\mathbb{Z} = m'\mathbb{Z}$, so $m' \mid m, m \geq 0, m' \geq 0$. The conclusion is $m = m'$ and $a\mathbb{Z} \cap b\mathbb{Z} = m\mathbb{Z}, m \geq 0$, so $m = a \vee b$. \square

Ex. 1.20 Prove the following:

- (a) $\operatorname{ord}_p[a, b] = \max(\operatorname{ord}_p(a), \operatorname{ord}_p(b))$.
- (b) $(a, b)[a, b] = ab$.
- (c) $(a + b, [a, b]) = (a, b)$.

Proof. (a) Let $a = \varepsilon \prod_p p^{a(p)}, b = \varepsilon' \prod_p p^{b(p)}, \varepsilon, \varepsilon' = \pm 1$, and

$$m = \prod_p p^{\max(a(p), b(p))}.$$

Then

- (i) $m \geq 0$.

- (ii) As $a(p) \leq \max(a(p), b(p))$, $p^{a(p)} \mid p^{\max(a(p), b(p))}$, so $a \mid m$. Similarly, $b \mid m$.
 (iii) If $\mu = \varepsilon'' \prod_p p^{c(p)}$ is a common multiple of a and b , then for all primes p , $a(p) \leq c(p)$, $b(p) \leq c(p)$, so $\max(a(p), b(p)) \leq c(p)$, so $m \mid \mu$. m verifies the characterisation of lcm :

$$m = a \vee b = \prod_p p^{\max(a(p), b(p))}.$$

So $\text{ord}_p[a, b] = \max(\text{ord}_p(a), \text{ord}_p(b))$.

(b) Similarly, we prove that

$$a \wedge b = \prod_p p^{\min(a(p), b(p))}.$$

As $\max(a, b) + \min(a, b) = a + b$, we obtain

$$(a \vee b)(a \wedge b) = |ab|.$$

Second proof (without decompositions in primes) :

Let $d = a \wedge b$. If $d = 0$, then $a = b = 0$ and $(a \vee b)(a \wedge b) = ab$.

Suppose now that $d \neq 0$. There exist integers a', b' such that

$$a = da', b = db', a' \wedge b' = 1.$$

Let $m = da'b' : a = da' \mid m$ and $b = db' \mid m$. If μ is a common multiple of a and b , then $d \mid \mu$, and $a' \mid \mu/d, b' \mid \mu/d$. As $a' \wedge b' = 1$, $a'b' \mid \mu/d$ (see Ex.1.9). so $m = da'b' \mid \mu$.

$|m|$ verifies the characterization of lcm (Ex. 1.19), so $a \vee b = |m| = |da'b'| = |ab|/d$.

Conclusion : $(a \vee b)(a \wedge b) = |ab|$.

(c) Let $\delta \in \mathbb{Z}$. If $\delta \mid a, \delta \mid b$, then $\delta \mid a + b$ and $\delta \mid a \vee b$.

Reciprocally, suppose that $\delta \mid a + b, \delta \mid a \vee b$.

Let $a', b' \in \mathbb{Z}$ such that $a = da', b = db', a' \wedge b' = 1$. Then $a \vee b = da'b'$, so

$$\begin{aligned} \delta &\mid d(a' + b'), \\ \delta &\mid da'b'. \end{aligned}$$

Multiplying the first relation by b' and a' , we obtain : $\delta \mid da'b' + db'^2, \delta \mid da'^2 + da'b'$. As $\delta \mid da'b'$, we obtain :

$$\begin{aligned} \delta &\mid db'^2 \\ \delta &\mid da'^2 \end{aligned}$$

As $a'^2 \wedge b'^2 = 1$, $\delta \mid d(a'^2 \wedge b'^2) = d$, so $\delta \mid a, \delta \mid b$.

The set of divisors of a, b is the same that the set of divisors of $a + b, a \vee b$, so

$$(a + b) \wedge (a \vee b) = a \wedge b.$$

□

2 Chapter 2

Ex 2.1 Show that $k[x]$, with k a finite field, has infinitely many irreducible polynomials.

Proof. Suppose that the set S of irreducible polynomials is finite : $S = \{P_1, P_2, \dots, P_n\}$.

Let $Q = P_1 P_2 \cdots P_n + 1$. As S contains the polynomials $x - a, a \in k$, $\deg(Q) \geq q = |k| > 1$. Thus Q is divisible by an irreducible polynomial. As S contains all the irreducible polynomials, there exists $i, 1 \leq i \leq n$, such that $P_i \mid Q = P_1 P_2 \cdots P_n + 1$, so $P_i \mid 1$, and P_i is an unit, in contradiction with the irreducibility of P_i .

Conclusion : $k[x]$ has infinitely many irreducible polynomials. As each polynomial has only a finite number of associates, there exists infinitely many monic irreducible polynomials. \square

Ex. 2.2. Let $p_1, p_2, \dots, p_t \in \mathbb{Z}$ be primes and consider the set of all rational numbers $r = a/b$, $a, b \in \mathbb{Z}$, such that $\text{ord}_{p_i} a \geq \text{ord}_{p_i} b$ for $i = 1, 2, \dots, t$. Show that this set is a ring and that up to taking associates p_1, p_2, \dots, p_t are the only primes.

Proof. Let R the set of such rationals. Simplifying these fractions, we obtain

$$r \in R \iff \exists p \in \mathbb{Z}, \exists q \in \mathbb{Z} \setminus \{0\}, r = \frac{p}{q}, q \wedge p_1 p_2 \cdots p_t = 1.$$

• $1 = 1/1 \in R$.

• if $r, r' \in R$, $r = p/q, r' = p'/q'$, with $q \wedge p_1 p_2 \cdots p_t = 1, q' \wedge p_1 p_2 \cdots p_t = 1$. then $qq' \wedge p_1 p_2 \cdots p_t = 1$, and $r - r' = \frac{pq' - qp'}{qq'}$, $rr' = \frac{pp'}{qq'}$, so $r - r', rr' \in R$.

Thus R is a subring of \mathbb{Q} .

If $r = a/b \in R$ is an unit of R , then $b/a \in R$, so $\text{ord}_{p_i} a = \text{ord}_{p_i}(b)$, $i = 1, \dots, t$. After simplification, $r = p/q$, with $p \wedge p_1 \cdots p_t = 1, q \wedge p_1 \cdots p_t = 1$, and such rationals are all units.

$p_i, 1 \leq i \leq t$, is a prime : if $p_i \mid rs$ in R , where $r = a/b, s = c/d \in R$, then there exists $u = e/f \in R$ such that $rs = p_i u$, with b, d, e relatively prime with p_1, \dots, p_t . Then $acf = p_i bde$. As $p_i \wedge f = 1$, p_i divides a or c in \mathbb{Z} , so p_i divides r or s in R .

If $r = a/b \in R$, with $b \wedge p_1 \cdots p_t = 1$, $a = p_1^{k_1} \cdots p_t^{k_t} v$, $v \in \mathbb{Z}, k_i \geq 0, i = 1, \dots, t$. So $r = up_1^{k_1} \cdots p_t^{k_t}$, where $u = v/b$ is an unit.

Let π be any prime in R . As any element in R , $\pi = up_1^{k_1} \cdots p_t^{k_t}, k_i \geq 0, u = a/b$ an unit. $u^{-1}\pi = p_1^{k_1} \cdots p_t^{k_t}$, so $\pi \mid p_1^{k_1} \cdots p_t^{k_t}$ (in R). As π is a prime in R , $\pi \mid p_i$ for an index $i = 1, \dots, t$. Moreover $p_i \mid p$, so p_i and π are associate.

Conclusion: the primes in R are the associates of p_1, \dots, p_t . \square

Ex. 2.3 Use the formula for $\phi(n)$ to give a proof that there are infinitely many primes.

[Hint: If p_1, p_2, \dots, p_t were all the primes, then $\phi(n) = 1$, where $n = p_1 p_2 \cdots p_t$.]

Proof. Let $\{p_1, \dots, p_t\}$ the finite set of primes, with $p_1 < p_2 < \cdots < p_t$, and $n = p_1 \cdots p_t$. By definition, $\phi(n)$ is the number of integers $k, 1 \leq k \leq n$, such that $k \wedge n = 1$. From the existence of decomposition in primes, if $k \geq 1$, $k = p_1^{k_1} \cdots p_t^{k_t}$, where $k_i \geq 0, i = 1, \dots, t$. So $k \wedge n = 1$ if and only if $k = 1$. Thus $\phi(n) = 1$. The formula for $\phi(n)$ gives $\phi(n) = (p_1 - 1) \cdots (p_t - 1) = 1$. As $p_i \geq 2$, this equation implies that $p_1 = p_2 = \cdots = p_t = 2$, so $t = 1$, and the only prime number is 2. But 3 is also a prime number : this is a contradiction.

Conclusion : there are infinitely many prime numbers. \square

Ex. 2.4 If a is a nonzero integer, then for $n > m$ show that $(a^{2^n} + 1, a^{2^m} + 1) = 1$ or 2 depending on whether a is odd or even.

Proof. Let $d = a^{2^n} + 1 \wedge a^{2^m} + 1$. Then $d \mid a^{2^n} + 1, d \mid a^{2^m} + 1$. So

$$\begin{aligned} a^{2^n} &\equiv -1 \pmod{d} \\ a^{2^m} &\equiv -1 \pmod{d} \end{aligned}$$

As $n > m$, 2^{n-m} is even, so

$$-1 \equiv a^{2^n} = (a^{2^m})^{2^{n-m}} \equiv (-1)^{2^{n-m}} \equiv 1 \pmod{d}.$$

$-1 \equiv 1 \pmod{d}$, then $d \mid 2$ ($d \geq 0$). Thus $d = 1$ or $d = 2$.

If a is even, $a^{2^n} + 1$ is odd, so $d = 1$.

If a is odd, both $a^{2^n} + 1, a^{2^m} + 1$ are even, so $d = 2$. □

Ex. 2.5 Use the result of Ex. 2.4 to show that there are infinitely many primes. (This proof is due to G. Polya.)

Proof. Let $F_n = 2^{2^n} + 1, n \in \mathbb{N}$. We know from Ex. 2.4 that $n \neq m \Rightarrow F_n \wedge F_m = 1$. Define p_n as the least prime divisor of F_n . If $n \neq m, F_n \wedge F_m = 1$, so $p_n \neq p_m$. The application $\varphi : \mathbb{N} \rightarrow \mathbb{N}, n \mapsto p_n$ is injective (one to one), so $\varphi(\mathbb{N})$ is an infinite set of prime numbers. □

Ex. 2.6 For a rational number r let $[r]$ be the largest integer less than or equal to r , e.g., $[\frac{1}{2}] = 0, [2] = 2$, and $[3 + \frac{1}{3}] = 3$. Prove $\text{ord}_p n! = \left[\frac{n}{p}\right] + \left[\frac{n}{p^2}\right] + \left[\frac{n}{p^3}\right] \cdots$.

Proof. The number N_k of multiples m of p^k which are not multiple of p^{k+1} , where $1 \leq m \leq n$, is

$$N_k = \left\lfloor \frac{n}{p^k} \right\rfloor - \left\lfloor \frac{n}{p^{k+1}} \right\rfloor.$$

Each of these numbers brings the contribution k to the sum $\text{ord}_p n! = \sum_{k=1}^n \text{ord}_p k$. Thus

$$\begin{aligned} \text{ord}_p n! &= \sum_{k \geq 1} k \left(\left\lfloor \frac{n}{p^k} \right\rfloor - \left\lfloor \frac{n}{p^{k+1}} \right\rfloor \right) \\ &= \sum_{k \geq 1} k \left\lfloor \frac{n}{p^k} \right\rfloor - \sum_{k \geq 1} k \left\lfloor \frac{n}{p^{k+1}} \right\rfloor \\ &= \sum_{k \geq 1} k \left\lfloor \frac{n}{p^k} \right\rfloor - \sum_{k \geq 2} (k-1) \left\lfloor \frac{n}{p^k} \right\rfloor \\ &= \left\lfloor \frac{n}{p} \right\rfloor + \sum_{k \geq 2} \left\lfloor \frac{n}{p^k} \right\rfloor \\ &= \sum_{k \geq 1} \left\lfloor \frac{n}{p^k} \right\rfloor \end{aligned}$$

Note that $\left\lfloor \frac{n}{p^k} \right\rfloor = 0$ if $p^k > n$, so this sum is finite. □

Ex. 2.7 Deduce from Ex. 2.6 that $\text{ord}_p n! \leq n/(p-1)$ and that $\sqrt[n]{n!} \leq \prod_{p \leq n} p^{1/(p-1)}$.
(The original statement $\prod_{p|n} p^{1/(p-1)}$ was modified.)

Proof.

$$\text{ord}_p n! = \sum_{k \geq 1} \left\lfloor \frac{n}{p^k} \right\rfloor \leq \sum_{k \geq 1} \frac{n}{p^k} = \frac{n}{p} \frac{1}{1 - \frac{1}{p}} = \frac{n}{p-1}$$

The decomposition of $n!$ in prime factors is

$n! = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$ where $\alpha_i = \text{ord}_{p_i} n! \leq \frac{n}{p_i-1}$, and $p_i \leq n$, $i = 1, 2, \dots, k$.

Then

$$\begin{aligned} n! &\leq p_1^{\frac{n}{p_1-1}} p_2^{\frac{n}{p_2-1}} \cdots p_k^{\frac{n}{p_k-1}} \\ \sqrt[n]{n!} &\leq p_1^{\frac{1}{p_1-1}} p_2^{\frac{1}{p_2-1}} \cdots p_k^{\frac{1}{p_k-1}} \\ &\leq \prod_{p \leq n} p^{\frac{1}{p-1}} \end{aligned}$$

(the values of p in this product describe all prime numbers $p \leq n$.) □

Ex. 2.8 Use Exercise 7 to show that there are infinitely many primes.

Proof. If the set \mathbb{P} of prime numbers was finite, we obtain from Ex.2.7, for all $n \geq 2$:

$$\sqrt[n]{n!} \leq C = \prod_{p \in \mathbb{P}} p^{\frac{1}{p-1}},$$

where C is an absolute constant.

Yet $\lim_{n \rightarrow \infty} \sqrt[n]{n!} = +\infty$. Indeed

$$\ln(\sqrt[n]{n!}) = \frac{1}{n}(\ln 1 + \ln 2 + \cdots + \ln n)$$

As \ln is an increasing function,

$$\int_{i-1}^i \ln t \, dt \leq \ln i, \quad i = 2, 3, \dots, n$$

So

$$\int_1^n \ln t \, dt = \sum_{i=2}^n \int_{i-1}^i \ln t \, dt \leq \sum_{i=2}^n \ln i = \sum_{i=1}^n \ln i$$

Thus

$$\ln(\sqrt[n]{n!}) \geq \frac{1}{n} \int_1^n \ln t \, dt = \frac{1}{n}(n \ln n - n + 1) = \ln n - 1 + \frac{1}{n}$$

As $\lim_{n \rightarrow \infty} \ln n - 1 + \frac{1}{n} = +\infty$, $\lim_{n \rightarrow \infty} \ln(\sqrt[n]{n!}) = +\infty$, so $\lim_{n \rightarrow \infty} \sqrt[n]{n!} = +\infty$.

So there exists n such that $\sqrt[n]{n!} \geq C$: this is a contradiction. \mathbb{P} is an infinite set. □

Ex. 2.9 A function on the integers is said to be multiplicative if $f(ab) = f(a)f(b)$ whenever $(a, b) = 1$. Show that a multiplicative function is completely determined by its value on prime powers.

Proof. Let the decomposition of n in prime factors be $n = p_1^{k_1} \cdots p_t^{k_t}$, $p_1 < \cdots < p_t$. As $p_i^{k_i} \wedge p_j^{k_j} = 1$ for $i \neq j$, $i, j = 1, \dots, t$,

$$f(n) = f(p_1^{k_1} \cdots p_t^{k_t}) = f(p_1^{k_1}) \cdots f(p_t^{k_t})$$

(by induction on the number of prime factors.)

So $f(n)$ is completely determined by its value on prime powers. \square

Ex. 2.10 If $f(n)$ is a multiplicative function, show that the function $g(n) = \sum_{d|n} f(d)$ is also multiplicative.

Proof. If $n \wedge m = 1$,

$$\begin{aligned} g(nm) &= \sum_{\delta|nm} f(\delta) \\ &= \sum_{d|n, d'|m} f(dd') \end{aligned}$$

Actually, if $d|n, d'|m$, so $\delta = dd'|nm$, and reciprocally, if $\delta|nm$, as $n \wedge m = 1$, there exist d, d' such that $d|n, d'|m$, and $\delta = dd'$.

If $d|n, d'|m$, with $n \wedge m = 1$, then $d \wedge d' = 1$, so

$$\begin{aligned} g(nm) &= \sum_{d|n} \sum_{d'|m} f(d)f(d') \\ &= \sum_{d|n} f(d) \sum_{d'|m} f(d') \\ &= g(n)g(m) \end{aligned}$$

g is a multiplicative function. \square

Ex. 2.11 Show that $\phi(n) = n \sum_{d|n} \mu(d)/d$ by first proving that $\mu(d)/d$ is multiplicative and then using Ex. 2.9 and 2.10.

Proof. Let's verify that μ is a multiplicative function.

If $n \wedge m = 1$, then $n = p_1^{a_1} \cdots p_l^{a_l}$, $m = q_1^{b_1} \cdots q_r^{b_r}$, where $p_1, \dots, p_l, q_1, \dots, q_r$ are distinct primes. Then the decomposition in prime factors of nm is $nm = p_1^{a_1} \cdots p_l^{a_l} q_1^{b_1} \cdots q_r^{b_r}$. If one of the a_i or one of the b_j is greater than 1, then $\mu(nm) = 0 = \mu(n)\mu(m)$. Otherwise, $n = p_1 \cdots p_l$, $m = q_1 \cdots q_r$, $nm = p_1 \cdots p_l q_1 \cdots q_r$, and $\mu(nm) = (-1)^{l+r} = (-1)^l (-1)^r = \mu(n)\mu(m)$. So

$$\frac{\mu(nm)}{nm} = \frac{\mu(n)}{n} \frac{\mu(m)}{m}.$$

that is, $n \mapsto \frac{\mu(n)}{n}$ is a multiplicative function.

From Ex.2.10, $n \mapsto \sum_{d|n} \frac{\mu(d)}{d}$ is also a multiplicative function, and so is ψ , where ψ is defined by

$$\psi(n) = n \sum_{d|n} \frac{\mu(d)}{d}.$$

To verify the equality $\phi = \psi$, it is sufficient from Ex. 2.9 to verify $\phi(p^k) = \psi(p^k)$ for all prime powers $p^k, k \geq 1$ ($\phi(1) = \psi(1) = 1$).

$$\begin{aligned} \psi(p^k) &= p^k \sum_{d|p^k} \frac{\mu(p^k)}{p^k} \\ &= p^k \left(\frac{\mu(1)}{1} + \frac{\mu(p)}{p} \right) \end{aligned}$$

(The other terms are null.)

So

$$\psi(p^k) = p^k \left(1 - \frac{1}{p} \right) = p^k - p^{k-1} = \phi(p^k).$$

Thus $\phi = \psi$: for all $n \geq 1$,

$$\phi(n) = n \sum_{d|n} \frac{\mu(d)}{d}.$$

□

Ex. 2.12 Find formulas for $\sum_{d|n} \mu(d)\phi(d)$, $\sum_{d|n} \mu(d)^2\phi(d)^2$, and $\sum_{d|n} \mu(d)/\phi(d)$.

Proof. As μ, ϕ are multiplicative, so are $\mu\phi, \mu^2\phi^2, \mu/\phi$. We deduce from Ex. 2.10 that the three following functions F, G, H are multiplicative, defined by

$$F(n) = \sum_{d|n} \mu(d)\phi(d), G(n) = \sum_{d|n} \mu(d)^2\phi(d)^2, H(n) = \sum_{d|n} \mu(d)/\phi(d),$$

so it is sufficient to compute their values on prime powers $p^k, k \geq 1$.

$$\begin{aligned} F(p^k) &= \sum_{i=0}^k \mu(p^i)\phi(p^i) \\ &= \phi(1) - \phi(p) = 1 - (p-1) = 2-p \end{aligned}$$

So $F(n) = \sum_{p|n} (2-p)$.

Similarly,

$$\begin{aligned} G(p^k) &= \sum_{i=0}^k \mu(p^i)^2\phi(p^i)^2 \\ &= \phi(1)^2 + \phi(p)^2 = 1 + (p-1)^2 = p^2 - 2p + 2 \end{aligned}$$

$$\begin{aligned} H(p^k) &= \sum_{i=0}^k \mu(p^i)/\phi(p^i) \\ &= 1/\phi(1) - 1/\phi(p) = 1 - 1/(p-1) = (p-2)/(p-1) \end{aligned}$$

□

Ex. 2.13 Let $\sigma_k(n) = \sum_{d|n} d^k$. Show that $\sigma_k(n)$ is multiplicative and find a formula for it.

Proof. As $n \mapsto n^k$ is multiplicative, then so is σ_k (Ex. 2.10).

• Suppose $k \neq 0$.

If $n = p^\alpha$ is a prime power ($\alpha \geq 1$),

$$\begin{aligned}\sigma_k(p^\alpha) &= \sum_{i=0}^{\alpha} p^{ik} \\ &= \frac{p^{(\alpha+1)k} - 1}{p^k - 1}\end{aligned}$$

• if $k = 0$, $\sigma_0(n)$ is the number of divisors of n .

$$\begin{aligned}\sigma_0(p^\alpha) &= \sum_{i=0}^{\alpha} 1 \\ &= \alpha + 1\end{aligned}$$

Conclusion : if $n = p_1^{\alpha_1} \cdots p_t^{\alpha_t}$ is the decomposition of n in prime factors, then

$$\begin{aligned}\sigma_0(n) &= (\alpha_1 + 1) \cdots (\alpha_t + 1), \\ \sigma_k(n) &= \prod_{i=1}^t \frac{p_i^{(\alpha_i+1)k} - 1}{p_i^k - 1} \quad (k \neq 0).\end{aligned}$$

□

Ex. 2.14 If $f(n)$ is multiplicative, show that $h(n) = \sum_{d|n} \mu(n/d)f(d)$ is also multiplicative.

Proof. We show first that the Dirichlet product $f \circ g$ of two multiplicative functions f, g is multiplicative. Suppose that $n \wedge m = 1$. If $d | n, d' | m$, so $\delta = dd' | nm$, and reciprocally, if $\delta | nm$, as $n \wedge m = 1$, there exist d, d' such that $d | n, d' | m$, and $\delta = dd'$. Thus

$$\begin{aligned}(f \circ g)(nm) &= \sum_{\delta|nm} f(\delta)g\left(\frac{nm}{\delta}\right) \\ &= \sum_{d|n, d'|m} f(dd')g\left(\frac{nm}{dd'}\right) \\ &= \sum_{d|n} \sum_{d'|m} f(d)f(d')g\left(\frac{n}{d}\right)g\left(\frac{m}{d'}\right) \\ &= \sum_{d|n} f(d)g\left(\frac{n}{d}\right) \sum_{d'|m} f(d')g\left(\frac{m}{d'}\right) \\ &= (f \circ g)(n)(f \circ g)(m)\end{aligned}$$

Applying this result with $g = \mu$, we obtain that $n \mapsto h(n) = \sum_{d|n} \mu(n/d)f(d)$ is multiplicative, if f is multiplicative. □

Ex. 2.15 Show that

$$(a) \sum_{d|n} \mu(n/d) \nu(d) = 1 \text{ for all } n.$$

$$(b) \sum_{d|n} \mu(n/d) \sigma(d) = n \text{ for all } n.$$

Proof. Here $\nu = \sigma_0, \sigma = \sigma_1$.

(a) From the Möbius Inversion Theorem, as $\nu(n) = \sum_{d|n} 1 = \sum_{d|n} I(d)$, where $I(n) = 1$ for all $n \geq 1$,

$$1 = I(n) = \sum_{d|n} \mu(n/d) \nu(d).$$

(b) From the same theorem, as $\sigma(n) = \sum_{d|n} d = \sum_{d|n} \text{Id}(d)$, where $\text{Id}(n) = n$ for all $n \geq 1$,

$$n = \text{Id}(n) = \sum_{d|n} \mu(n/d) \sigma(d).$$

□

Ex. 2.16 Show that $\nu(n)$ is odd iff n is a square.

Proof. • If $n = a^2$ is a square, where $a = p_1^{k_1} \cdots p_t^{k_t}$, then $\nu(n) = (2k_1 + 1) \cdots (2k_t + 1)$ is odd.

• Reciprocally, if $n = q_1^{l_1} \cdots q_r^{l_r}$ is odd, then $(l_1 + 1) \cdots (l_r + 1)$ is odd. So each $l_i + 1$ is odd, and then l_i is even, for $i = 1, 2, \dots, r$: n is a square. □

Ex. 2.17 Show that $\sigma(n)$ is odd iff n is a square or twice a square.

Proof. • Note that for all $r \geq 0$, $\sigma(2^r) = 1 + 2 + 2^2 + \cdots + 2^r = 2^{r+1} - 1$ is always odd.

If $p \neq 2$, $\sigma(p^{2k}) = 1 + p + p^2 + \cdots + p^{2k}$ is a sum of $2k + 1$ odd numbers, so is odd.

So if $n = a^2$, or $n = 2a^2, a \in \mathbb{Z}$, $\sigma(n)$ is odd.

• Reciprocally, suppose that $\sigma(n)$ is odd, where $n = p_1^{k_1} p_2^{k_2} \cdots p_t^{k_t}$, with $p_1 = 2 < p_2 < \cdots < p_t$. Then

$$\sigma(n) = (2^{k_1+1} - 1) \frac{p_2^{k_2+1} - 1}{p_2 - 1} \cdots \frac{p_t^{k_t+1} - 1}{p_t - 1}$$

is odd. Then each $\frac{p_i^{k_i+1} - 1}{p_i - 1} = 1 + p_i + \cdots + p_i^{k_i}$ ($i = 2, \dots, t$) is odd. As each $p_i^j, j = 0, \dots, k_i$ is odd, the number of terms $k_i + 1$ is odd, so k_i is even ($i = 2, \dots, t$). Thus n is a square, or twice a square. □

Ex. 2.18 Prove that $\phi(n)\phi(m) = \phi((n, m))\phi([n, m])$.

Proof. Let p_1, \dots, p_r the common prime factors of n and m .

$$\begin{aligned} n &= p_1^{\alpha_1} \cdots p_r^{\alpha_r} q_1^{\lambda_1} \cdots q_s^{\lambda_s} \\ m &= p_1^{\beta_1} \cdots p_r^{\beta_r} s_1^{\mu_1} \cdots s_t^{\mu_t} \end{aligned}$$

where $\alpha_i, \beta_i, \lambda_j, \mu_k \in \mathbb{N}^*$, $1 \leq i \leq r, 1 \leq j \leq s, 1 \leq k \leq t$ (the formula $\phi(p^\alpha) = p^\alpha - p^{\alpha-1}$ is not valid if $\alpha = 0$). Then

$$\begin{aligned} n \wedge m &= p_1^{\gamma_1} \cdots p_r^{\gamma_r} \\ n \vee m &= p_1^{\delta_1} \cdots p_r^{\delta_r} q_1^{\lambda_1} \cdots q_s^{\lambda_s} s_1^{\mu_1} \cdots s_t^{\mu_t}, \end{aligned}$$

where $\gamma_i = \min(\alpha_i, \beta_i), \delta_i = \max(\alpha_i, \beta_i)$ ($\gamma_i \geq 1, \delta_i \geq 1$), $1 \leq i \leq r$. Then

$$\begin{aligned} \phi(n \wedge m) &= \prod_{i=1}^r (p_i^{\gamma_i} - p_i^{\gamma_i-1}) \\ \phi(n \vee m) &= \prod_{i=1}^r (p_i^{\delta_i} - p_i^{\delta_i-1}) \prod_{i=1}^s (q_i^{\lambda_i} - q_i^{\lambda_i-1}) \prod_{i=1}^t (s_i^{\mu_i} - s_i^{\mu_i-1}) \end{aligned}$$

As $\alpha_i + \beta_i = \min(\alpha_i, \beta_i) + \max(\alpha_i, \beta_i) = \gamma_i + \delta_i, 1 \leq i \leq r$, then

$$\begin{aligned} \phi(n)\phi(m) &= \prod_{i=1}^r (p_i^{\alpha_i} - p_i^{\alpha_i-1}) \prod_{i=1}^s (q_i^{\lambda_i} - q_i^{\lambda_i-1}) \prod_{i=1}^r (p_i^{\beta_i} - p_i^{\beta_i-1}) \prod_{i=1}^t (s_i^{\mu_i} - s_i^{\mu_i-1}) \\ &= \prod_{i=1}^r \left[p_i^{\alpha_i+\beta_i} \left(1 - \frac{1}{p_i}\right)^2 \right] \prod_{i=1}^s (q_i^{\lambda_i} - q_i^{\lambda_i-1}) \prod_{i=1}^t (s_i^{\mu_i} - s_i^{\mu_i-1}) \\ &= \prod_{i=1}^r \left[p_i^{\gamma_i+\delta_i} \left(1 - \frac{1}{p_i}\right)^2 \right] \prod_{i=1}^s (q_i^{\lambda_i} - q_i^{\lambda_i-1}) \prod_{i=1}^t (s_i^{\mu_i} - s_i^{\mu_i-1}) \\ &= \prod_{i=1}^r (p_i^{\gamma_i} - p_i^{\gamma_i-1}) \prod_{i=1}^r (p_i^{\delta_i} - p_i^{\delta_i-1}) \prod_{i=1}^s (q_i^{\lambda_i} - q_i^{\lambda_i-1}) \prod_{i=1}^t (s_i^{\mu_i} - s_i^{\mu_i-1}) \\ &= \phi(n \wedge m) \phi(n \vee m) \end{aligned}$$

□

Ex. 2.19 Prove that $\phi(nm)\phi((n, m)) = (n, m)\phi(n)\phi(m)$.

Proof. With the notations of Ex. 2.18,

$$\begin{aligned} \phi(nm) &= \prod_{i=1}^r p_i^{\alpha_i+\beta_i} \left(1 - \frac{1}{p_i}\right) \prod_{i=1}^s q_i^{\lambda_i} \left(1 - \frac{1}{q_i}\right) \prod_{i=1}^t s_i^{\mu_i} \left(1 - \frac{1}{s_i}\right) \\ \phi(n \wedge m) &= \prod_{i=1}^r p_i^{\gamma_i} \left(1 - \frac{1}{p_i}\right) \end{aligned}$$

so

$$\begin{aligned} (n \wedge m)\phi(n)\phi(m) &= \prod_{i=1}^r p_i^{\gamma_i} \prod_{i=1}^r \left[p_i^{\alpha_i+\beta_i} \left(1 - \frac{1}{p_i}\right)^2 \right] \prod_{i=1}^s q_i^{\lambda_i} \left(1 - \frac{1}{q_i}\right) \prod_{i=1}^t s_i^{\mu_i} \left(1 - \frac{1}{s_i}\right) \\ &= \prod_{i=1}^r \left[p_i^{\alpha_i+\beta_i+\gamma_i} \left(1 - \frac{1}{p_i}\right)^2 \right] \prod_{i=1}^s q_i^{\lambda_i} \left(1 - \frac{1}{q_i}\right) \prod_{i=1}^t s_i^{\mu_i} \left(1 - \frac{1}{s_i}\right) \\ &= \phi(nm)\phi(n \wedge m) \end{aligned}$$

Conclusion :

$$(n \wedge m)\phi(n)\phi(m) = \phi(nm)\phi(n \wedge m).$$

□

Ex. 2.20 Prove that $\prod_{d|n} d = n^{\nu(n)/2}$.

Proof. Let

$$n = p_1^{\alpha_1} \cdots p_k^{\alpha_k}$$

the decomposition of n in prime factors.

$$\begin{aligned} \left(\prod_{d|n} d \right)^2 &= \prod_{d|n} d \prod_{d|n} d \\ &= \prod_{d|n} d \prod_{\delta|n} \frac{n}{\delta} \quad (\delta = n/d) \\ &= n^{\nu(n)} \prod_{d|n} d \prod_{d|n} \frac{1}{d} \\ &= n^{\nu(n)} \end{aligned}$$

Conclusion :

$$\prod_{d|n} d = n^{\frac{\nu(n)}{2}}.$$

□

Ex. 2.21 Define $\wedge(n) = \log p$ if n is a power of p and zero otherwise. Prove that $\sum_{d|n} \mu(n/d) \log d = \wedge(n)$. [Hint: First calculate $\sum_{d|n} \wedge(d)$ and then apply the Möbius inversion formula.]

Proof.

$$\begin{cases} \wedge(n) &= \log p & \text{if } n = p^\alpha, \alpha \in \mathbb{N}^* \\ &= 0 & \text{otherwise.} \end{cases}$$

Let $n = p_1^{\alpha_1} \cdots p_t^{\alpha_t}$ the decomposition of n in prime factors. As $\wedge(d) = 0$ for all divisors of n , except for $d = p_j^i, i > 0, j = 1, \dots, t$,

$$\begin{aligned} \sum_{d|n} \wedge(d) &= \sum_{i=1}^{\alpha_1} \wedge(p_1^i) + \cdots + \sum_{i=1}^{\alpha_t} \wedge(p_t^i) \\ &= \alpha_1 \log p_1 + \cdots + \alpha_t \log p_t \\ &= \log n \end{aligned}$$

By Möbius Inversion Theorem,

$$\wedge(n) = \sum_{d|n} \mu\left(\frac{n}{d}\right) \log d.$$

□

Ex. 2.22 Show that the sum of all the integers t such that $1 \leq t \leq n$ and $(t, n) = 1$ is $\frac{1}{2}n\phi(n)$.

Proof. Suppose $n > 1$ (the formula is false if $n = 1$).

$$\text{Let } S = \sum_{1 \leq t \leq n, t \wedge n = 1} t = \sum_{1 \leq t \leq n-1, t \wedge n = 1} t.$$

Using the symmetry $t \mapsto n - t$, as $t \wedge n = 1 \iff (n - t) \wedge n = 1$, we obtain

$$\begin{aligned} 2S &= \sum_{1 \leq t \leq n-1, t \wedge n = 1} t + \sum_{1 \leq t \leq n-1, t \wedge n = 1} t \\ &= \sum_{1 \leq t \leq n-1, t \wedge n = 1} t + \sum_{1 \leq s \leq n-1, (n-s) \wedge n = 1} n - s \quad (s = n - t) \\ &= \sum_{1 \leq t \leq n-1, t \wedge n = 1} t + \sum_{1 \leq t \leq n-1, (n-t) \wedge n = 1} n - t \\ &= \sum_{1 \leq t \leq n-1, t \wedge n = 1} t + \sum_{1 \leq t \leq n-1, t \wedge n = 1} n - t \\ &= \sum_{1 \leq t \leq n-1, t \wedge n = 1} n \\ &= n \text{ Card}\{t \in \mathbb{N} \mid 1 \leq t \leq n - 1, t \wedge n = 1\} \\ &= n\phi(n) \end{aligned}$$

Conclusion :

$$\forall n \in \mathbb{N}^*, \quad \sum_{1 \leq t \leq n, t \wedge n = 1} t = \frac{1}{2} n\phi(n).$$

□

Ex. 2.23 Let $f(x) \in \mathbb{Z}[x]$ and let $\psi(n)$ be the number of $f(j), j = 1, 2, \dots, n$, such that $(f(j), n) = 1$. Show that $\psi(n)$ is multiplicative and that $\psi(p^t) = p^{t-1}\psi(p)$. Conclude that $\psi(n) = n \prod_{p|n} \psi(p)/p$.

Proof. My interpretation of this statement is that $\psi(n)$ is the number of $j, j = 1, 2, \dots, n$, such that $(f(j), n) = 1$ (if f is not one to one, we may obtain a different value).

Let $A_n = \{j \in \mathbb{Z}, 1 \leq j \leq n \mid f(j) \wedge n = 1\}$: then $\psi(n) = |A_n|$. If $f(x) = \sum_{k=0}^d a_k x^k$, note $f_n(x) \in (\mathbb{Z}/n\mathbb{Z})[x]$ the polynomial $f_n(x) = \sum_{k=0}^n [a_k]_n x^k$ (here, we represent the class of $j \in \mathbb{Z}$ in $\mathbb{Z}/n\mathbb{Z}$ by $[j]_n$). We can write without inconvenient $f = f_n$.

Let $B_n = \{a \in \mathbb{Z}/n\mathbb{Z} \mid f(a) \in (\mathbb{Z}/n\mathbb{Z})^*\}$, where $(\mathbb{Z}/n\mathbb{Z})^*$ is the group of invertible elements of $\mathbb{Z}/n\mathbb{Z}$.

Then $u : A_n \rightarrow B_n, j \mapsto [j]_n$ is a bijection.

Indeed u is well defined : if $j \in A_n, f(j) \wedge n = 1$, so $f([j]_n) = [f(j)]_n \in (\mathbb{Z}/n\mathbb{Z})^*$.

u is injective : $[j]_n = [k]_n$ with $1 \leq j \leq n, 1 \leq k \leq n$ implies $j = k$.

u is surjective : if $a \in \mathbb{Z}/n\mathbb{Z}$ verifies $f(a) \in (\mathbb{Z}/n\mathbb{Z})^*$, let j the unique representative of a such that $1 \leq j \leq n$. Then $f(j) \wedge n = 1$, so $u(j) = a$.

Thus

$$\psi(n) = |B_n|, \text{ where } B_n = \{a \in \mathbb{Z}/n\mathbb{Z} \mid f(a) \in (\mathbb{Z}/n\mathbb{Z})^*\}.$$

Suppose $n \wedge m = 1$. Let

$$\varphi : \begin{cases} B_{nm} & \rightarrow B_n \times B_m \\ [j]_{nm} & \mapsto ([j]_n, [j]_m) \end{cases}$$

• φ is well defined : $[j]_{nm} = [k]_{nm} \Rightarrow j \equiv k \pmod{nm} \Rightarrow (j \equiv k \pmod{n}, j \equiv k \pmod{m}) \Rightarrow ([j]_n, [j]_m) = ([k]_n, [k]_m)$.

• φ is injective : if $\varphi([j]_{nm}) = \varphi([k]_{nm})$, then $[j]_n = [k]_n, [j]_m = [k]_m$, so $n \mid j - k, m \mid j - k$. As $n \wedge m = 1, nm \mid j - k$ so $[j]_{nm} = [k]_{nm}$.

• φ is surjective : if $(a, b) \in B_n \times B_m$, there exist $j, k \in \mathbb{Z}, 1 \leq j \leq n, 1 \leq k \leq m$, such that $a = [j]_n, b = [k]_m$. From the Chinese Remainder Theorem, there exists $i \in \mathbb{Z}, 1 \leq i \leq nm$, such that $i \equiv j \pmod{n}, i \equiv k \pmod{m}$. Then $\varphi([i]_{nm}) = ([i]_n, [i]_m) = ([j]_n, [k]_m) = (a, b)$.

Finally, $\psi(nm) = |B_{nm}| = |B_n| |B_m| = \psi(n)\psi(m)$, if $n \wedge m = 1$: ψ is a multiplicative function.

The interval $I = [1, p^t]$ is the disjoint reunion of the p^{t-1} intervals $I_k = [kp+1, (k+1)p]$ for $k = 0, 1, \dots, p^{t-1} - 1$, so $\psi(p^t) = \sum_{k=0}^{p^{t-1}-1} \text{Card } C_k$, where $C_k = \{j \in I_k \mid f(j) \wedge p^t = 1\} = \{j \in I_k \mid f(j) \wedge p = 1\}$.

As $f(j) \wedge p = 1 \iff f(j - kp) \wedge p = 1$, the application $v : C_k \rightarrow C_0, j \mapsto j - kp$ is well defined and is bijective, so $|C_k| = |C_0| = \psi(p)$. Thus $\psi(p^t) = p^{t-1} \text{Card } I_0 = p^{t-1} \psi(p)$:

$$\psi(p^t) = p^{t-1} \psi(p).$$

If $n = \prod_{p|n} p^{t(p)}$, then

$$\begin{aligned} \psi(n) &= \prod_{p|n} \psi(p^{t(p)}) \\ &= \prod_{p|n} p^{t(p)-1} \psi(p) \\ &= n \prod_{p|n} \frac{\psi(p)}{p} \end{aligned}$$

□

Ex. 2.24 Supply the details to the proof of Theorem 3.

As Adam Michalik, I suppose that there is a misprint, we must prove Theorem 4 :

Let k a finite field with q elements.

$\sum q^{-\deg p(x)}$ diverges, where the sum is over all monic irreducible $p(x)$ in $k[x]$.

Proof. Notations :

\mathcal{P} : set of all monic polynomials p in $k[x]$.

\mathcal{P}_n : set of all monic polynomials p in $k[x]$ with $\deg(p) \leq n$.

\mathcal{M} : set of all monic irreducible polynomials p in $k[x]$.

We must prove that $\sum_{p \in \mathcal{M}} q^{-\deg p(x)}$ diverges.

• $\sum_{f \in \mathcal{P}} q^{-\deg f}$ diverges :

$$\begin{aligned} \sum_{f \in \mathcal{P}_n} \frac{1}{q^{\deg f}} &= \sum_{d=0}^n \sum_{\deg(f)=d} \frac{1}{q^d} \\ &= \sum_{d=0}^n \frac{1}{q^d} \text{Card } \{f \in \mathcal{P} \mid \deg(f) = d\} \\ &= \sum_{d=0}^n \frac{1}{q^d} q^d = n + 1. \end{aligned}$$

So $\sum_{f \in \mathcal{P}} q^{-\deg f}$ diverges.

- $\sum_{f \in \mathcal{P}} q^{-2 \deg f}$ converges :

$$\begin{aligned} \sum_{f \in \mathcal{P}_n} q^{-2 \deg(f)} &= \sum_{d=0}^n \sum_{\deg(f)=d} \frac{1}{q^{2d}} \\ &= \sum_{d=0}^n \frac{1}{q^{2d}} \text{Card}\{f \in \mathcal{P} \mid \deg(f) = d\} \\ &= \sum_{d=0}^n \frac{1}{q^d} \\ &\leq \frac{1}{1 - \frac{1}{q}} \end{aligned}$$

As any finite subset of \mathcal{P} is included in some \mathcal{P}_n , $\sum_{f \in \mathcal{P}} q^{-2 \deg f}$ converges.

- $\sum q^{-\deg p(x)}$ diverges :

Let $\mathcal{P}_n = \{p_1, p_2, \dots, p_{l(n)}\}$ the set of all monic irreducible polynomials such that $\deg p_i \leq n$. Let

$$\lambda(n) = \prod_{i=1}^{l(n)} \frac{1}{1 - \frac{1}{q^{\deg(p_i)}}}.$$

For simplicity, we write $l = l(n)$ for a fixed $n \in \mathbb{N}$. Then

$$\begin{aligned} \lambda(n) &= \prod_{i=1}^l \sum_{a_i=0}^{\infty} \frac{1}{q^{a_i \deg p_i}} \\ &= \left(1 + \frac{1}{q^{\deg p_1}} + \frac{1}{q^{2 \deg p_1}} + \dots\right) \times \dots \times \left(1 + \frac{1}{q^{\deg p_l}} + \frac{1}{q^{2 \deg p_l}} + \dots\right) \\ &= \sum_{(a_1, \dots, a_l) \in \mathbb{N}^l} \frac{1}{q^{\deg(p_1^{a_1} \dots p_l^{a_l})}} \end{aligned}$$

Since the monic prime factors of any polynomial $p \in \mathcal{P}_n$ are in \mathcal{P}_n , the decomposition of p is $p = p_1^{a_1} \dots p_l^{a_l}$, so

$$\lambda(n) \geq \sum_{p \in \mathcal{P}_n} \frac{1}{q^{\deg p}} = n + 1.$$

So $\lim_{n \rightarrow \infty} \lambda(n) = \infty$: this is another proof that there exist infinitely many monic irreducible

polynomials in $k[x]$ (cf Ex. 2.1).

$$\begin{aligned}
\log \lambda(n) &= - \sum_{i=1}^{l(n)} \log \left(1 - \frac{1}{q^{\deg p_i}} \right) \\
&= \sum_{i=1}^{l(n)} \sum_{m=1}^{\infty} \frac{1}{mq^{m \deg p_i}} \\
&= \frac{1}{q^{\deg p_1}} + \cdots + \frac{1}{q^{\deg p_{l(n)}}} + \sum_{i=1}^{l(n)} \sum_{m=2}^{\infty} \frac{1}{mq^{m \deg p_i}}
\end{aligned}$$

Yet

$$\begin{aligned}
\sum_{m=2}^{\infty} \frac{1}{mq^{m \deg p_i}} &\leq \sum_{m=2}^{\infty} \frac{1}{q^{m \deg p_i}} \\
&= \frac{1}{q^{2 \deg p_i}} \frac{1}{1 - \frac{1}{q^{\deg p_i}}} \\
&= \frac{1}{q^{2 \deg p_i} - q^{\deg p_i}} \leq \frac{2}{q^{2 \deg p_i}}
\end{aligned}$$

(the last inequality is equivalent to $2 \leq q^{\deg p_i}$). So

$$\log \lambda(n) \leq \frac{1}{q^{\deg p_1}} + \cdots + \frac{1}{q^{\deg p_{l(n)}}} + 2 \left(\frac{1}{q^{2 \deg p_1}} + \cdots + \frac{1}{q^{2 \deg p_{l(n)}}} \right).$$

As $\frac{1}{q^{2 \deg p_1}} + \cdots + \frac{1}{q^{2 \deg p_{l(n)}}}$ is less than the constant $\sum_{f \in \mathcal{P}} q^{-2 \deg f}$, if $\sum_{p \in \mathcal{M}} q^{-\deg p(x)}$ converges, then $\log \lambda(n) \leq C$, where C is a constant, so $\lambda(n) \leq e^C$ for all $n \in \mathbb{N}$, in contradiction with $\lim_{n \rightarrow \infty} \lambda(n) = \infty$.

Conclusion : $\sum_{p \in \mathcal{M}} q^{-\deg p(x)}$ diverges. \square

Ex. 2.25 Consider the function $\zeta(s) = \sum_{n=1}^{\infty} 1/n^s$. ζ is called the Riemann zeta function. It converges for $s > 1$. Prove the formal identity (Euler's identity)

$$\zeta(s) = \prod_p (1 - 1/p^s)^{-1}.$$

Proof. We prove this equality, not only formally, but for all complex value s such that $\operatorname{Re}(s) > 1$.

Let $s \in \mathbb{C}$ and $f(n) = \frac{1}{n^s}$, $n \in \mathbb{N}^*$.

f is completely multiplicative : $f(mn) = f(m)f(n)$ for $m, n \in \mathbb{N}^*$.

Moreover $\sum_{n=1}^{\infty} f(n)$ is absolutely convergent for $\operatorname{Re}(s) > 1$. Indeed, If $s = u + iv$, $u, v \in \mathbb{R}$, $|f(n)| = |n^{-s}| = |e^{-s \log(n)}| = |e^{-u \log(n)} e^{-iv \log(n)}| = e^{-u \log(n)} = \frac{1}{n^u}$, so $\sum_{n=1}^{\infty} |f(n)| = \sum_{n=1}^{\infty} 1/n^u$ converges if $u = \operatorname{Re}(s) > 1$.

With these properties of f (f multiplicative and $\sum_{n=1}^{\infty} f(n)$ absolutely convergent), we will show that

$$\sum_{n=1}^{\infty} f(n) = \prod_p (1 + f(p) + f(p^2) + \cdots).$$

Let $S^* = \sum_{n=1}^{\infty} |f(n)| < \infty$, and $S = \sum_{n=1}^{\infty} f(n) \in \mathbb{C}$. For each prime number p , $\sum_{k=1}^{\infty} |f(p^k)|$ converges (this sum is less than S^*), so $\sum_{k=0}^{\infty} f(p^k)$ converges absolutely. Thus, for $x \in \mathbb{R}$, the two finite products

$$P(x) = \prod_{p \leq x} \sum_{k=0}^{\infty} f(p^k), \quad P^*(x) = \prod_{p \leq x} \sum_{k=0}^{\infty} |f(p^k)|$$

are well defined.

If p, q are two prime numbers, as $\sum_{i=0}^{\infty} f(p^i), \sum_{j=0}^{\infty} f(q^j)$ are absolutely convergent, $(f(p^i)f(q^j))_{(i,j) \in \mathbb{N}^2}$ is sommable, so the sum of these elements can be arranged in any order :

$$\sum_{i=0}^{\infty} f(p^i) \sum_{k=0}^{\infty} f(q^k) = \sum_{(i,j) \in \mathbb{N}^2} f(p^i)f(q^j) = \sum_{(i,j) \in \mathbb{N}^2} f(p^i q^j).$$

If p_1, \dots, p_t are all the prime $p \leq x$, repeating t times these products, we obtain

$$\begin{aligned} P(x) &= \prod_{p \leq x} \sum_{k=0}^{\infty} f(p^k) \\ &= \sum_{i_1=0}^{\infty} f(p_1^{i_1}) \cdots \sum_{i_t=0}^{\infty} f(p_t^{i_t}) \\ &= \sum_{(i_1, \dots, i_t) \in \mathbb{N}^t} f(p_1^{i_1} \cdots p_t^{i_t}) \\ &= \sum_{n \in \Delta} f(n), \end{aligned}$$

where Δ is the set of integers $n \in \mathbb{N}^*$ whose prime factors are not greater than x . Let $\overline{\Delta} = \mathbb{N}^* \setminus \Delta$: this is the set of numbers $n \in \mathbb{N}^*$ such that at least a prime factor is greater than x . So

$$P(x) = \sum_{n \in \Delta} f(n) = S - \sum_{n \in \overline{\Delta}} f(n).$$

Then

$$|P(x) - S| \leq \sum_{n \in \overline{\Delta}} |f(n)| \leq \sum_{n \geq x} |f(n)|.$$

So $\lim_{x \rightarrow +\infty} P(x) = S$, that is

$$\prod_p \sum_{k=0}^{\infty} f(p^k) = \sum_{n=1}^{\infty} f(n).$$

Finally,

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{1}{n^s} &= \prod_p \left(1 + \frac{1}{p^s} + \cdots + \frac{1}{p^{ks}} + \cdots \right) \\ &= \prod_p (1 - 1/p^s)^{-1} \end{aligned}$$

□

Ex. 2.26 Verify the formal identities:

$$(a) \quad \zeta(s)^{-1} = \sum \mu(n)/n^s$$

$$(b) \quad \zeta(s)^2 = \sum \nu(n)/n^s$$

$$(c) \quad \zeta(s)\zeta(s-1) = \sum \sigma(n)/n^s$$

Proof. Without any consideration of convergence :

(a)

$$\begin{aligned} \zeta(s) \sum_{m=1}^{\infty} \frac{\mu(m)}{m^s} &= \sum_{n=1}^{\infty} \frac{1}{n^s} \sum_{m=1}^{\infty} \frac{\mu(m)}{m^s} \\ &= \sum_{n,m \geq 1} \frac{\mu(m)}{n^s m^s} \\ &= \sum_{u=1}^{\infty} \sum_{m|u} \mu(m) \frac{1}{u^s} \quad (u = nm) \\ &= \sum_{u=1}^{\infty} \frac{1}{u^s} \sum_{m|u} \mu(m) \\ &= 1 \end{aligned}$$

Indeed, $\sum_{m|u} \mu(m) = 1$ if $u = 1$, 0 otherwise. So

$$\zeta(s)^{-1} = \sum_{n \in \mathbb{N}^*} \mu(n)/n^s.$$

(b)

$$\begin{aligned} \zeta(s)^2 &= \sum_{n=1}^{\infty} \frac{1}{n^s} \sum_{m=1}^{\infty} \frac{1}{m^s} \\ &= \sum_{n,m \geq 1} \frac{1}{(nm)^s} \\ &= \sum_{u \geq 1} \sum_{n|u} \frac{1}{u^s} \\ &= \sum_{u \geq 1} \frac{1}{u^s} \sum_{n|u} 1 \\ &= \sum_{u \geq 1} \frac{1}{u^s} \nu(u) \end{aligned}$$

So

$$\zeta(s)^2 = \sum_{n=1}^{\infty} \frac{\nu(n)}{n^s}.$$

(c) For $\text{Re}(s) > 2$,

$$\begin{aligned}\zeta(s)\zeta(s-1) &= \sum_{n \geq 1} \frac{1}{n^s} \sum_{m \geq 1} \frac{1}{m^{s-1}} \\ &= \sum_{m, n \geq 1} \frac{m}{(nm)^s} \\ &= \sum_{u \geq 1} \left(\sum_{m|u} m \right) \frac{1}{u^s} \\ &= \sum_{u \geq 1} \frac{\sigma(u)}{u^s}\end{aligned}$$

So

$$\zeta(s)\zeta(s-1) = \sum_{n \geq 1} \frac{\sigma(n)}{n^s}.$$

□

Ex. 2.27 Show that $\sum 1/n$, the sum being over square free integers, diverges. Conclude that $\prod_{p < N} (1 + 1/p) \rightarrow \infty$ as $N \rightarrow \infty$. Since $e^x > 1 + x$, conclude that $\sum_{p < N} 1/p \rightarrow \infty$. (This proof is due to I. Niven.)

Proof. Let $S \subset \mathbb{N}^*$ the set of square free integers.

Let $N \in \mathbb{N}^*$. Every integer n , $1 \leq n \leq N$ can be written as $n = ab^2$, where a, b are integers and a is square free. Then $1 \leq a \leq N$, and $1 \leq b \leq \sqrt{N}$, so

$$\sum_{n \leq N} \frac{1}{n} \leq \sum_{a \in S, a \leq N} \sum_{1 \leq b \leq \sqrt{N}} \frac{1}{ab^2} \leq \sum_{a \in S, a \leq N} \frac{1}{a} \sum_{b=1}^{\infty} \frac{1}{b^2} = \frac{\pi^2}{6} \sum_{a \in S, a \leq N} \frac{1}{a}.$$

So

$$\sum_{a \in S, a \leq N} \frac{1}{a} \geq \frac{6}{\pi^2} \sum_{n \leq N} \frac{1}{n}.$$

As $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges, $\lim_{N \rightarrow \infty} \sum_{a \in S, a \leq N} \frac{1}{a} = +\infty$, so the family $(\frac{1}{a})_{a \in S}$ of the inverse of square free integers is not summable.

Let $S_N = \prod_{p < N} (1 + 1/p)$, and p_1, p_2, \dots, p_l ($l = l(N)$) all prime integers less than N . Then

$$\begin{aligned}S_N &= \left(1 + \frac{1}{p_1}\right) \cdots \left(1 + \frac{1}{p_l}\right) \\ &= \sum_{(\varepsilon_1, \dots, \varepsilon_l) \in \{0,1\}^l} \frac{1}{p_1^{\varepsilon_1} \cdots p_l^{\varepsilon_l}}\end{aligned}$$

We prove this last formula by induction. This is true for $l = 1$: $\sum_{\varepsilon \in \{0,1\}} 1/p_1^{\varepsilon} = 1 + 1/p_1$.

If it is true for the integer l , then

$$\begin{aligned}
\left(1 + \frac{1}{p_1}\right) \cdots \left(1 + \frac{1}{p_l}\right) \left(1 + \frac{1}{p_{l+1}}\right) &= \sum_{(\varepsilon_1, \dots, \varepsilon_l) \in \{0,1\}^l} \frac{1}{p_1^{\varepsilon_1} \cdots p_l^{\varepsilon_l}} \left(1 + \frac{1}{p_{l+1}}\right) \\
&= \sum_{(\varepsilon_1, \dots, \varepsilon_l) \in \{0,1\}^l} \frac{1}{p_1^{\varepsilon_1} \cdots p_l^{\varepsilon_l}} + \sum_{(\varepsilon_1, \dots, \varepsilon_l) \in \{0,1\}^l} \frac{1}{p_1^{\varepsilon_1} \cdots p_l^{\varepsilon_l} p_{l+1}} \\
&= \sum_{(\varepsilon_1, \dots, \varepsilon_l, \varepsilon_{l+1}) \in \{0,1\}^{l+1}} \frac{1}{p_1^{\varepsilon_1} \cdots p_l^{\varepsilon_l} p_{l+1}^{\varepsilon_{l+1}}}
\end{aligned}$$

So it is true for all l .

Thus $S_N = \sum_{n \in \Delta} \frac{1}{n}$, where Δ is the set of square free integers whose prime factors are less than N .

As $\sum 1/n$, the sum being over square free integers, diverges, $\lim_{N \rightarrow \infty} S_N = +\infty$:

$$\lim_{N \rightarrow \infty} \prod_{p < N} \left(1 + \frac{1}{p}\right) = +\infty.$$

$e^x \geq 1 + x$, $x \geq \log(1 + x)$ for $x > 0$, so

$$\log S_N = \sum_{k=1}^{l(N)} \log \left(1 + \frac{1}{p_k}\right) \leq \sum_{k=1}^{l(N)} \frac{1}{p_k}.$$

$\lim_{N \rightarrow \infty} \log S_N = +\infty$ and $\lim_{N \rightarrow \infty} l(N) = +\infty$, so

$$\lim_{N \rightarrow \infty} \sum_{p < N} \frac{1}{p} = +\infty.$$

□

3 Chapter 3

Ex. 3.1 Show that there are infinitely many primes congruent to -1 modulo 6.

Proof. Let n any integer such that $n \geq 3$, and $N = n! - 1 = 2 \times 3 \times \cdots \times n - 1 > 1$.

Then $N \equiv -1 \pmod{6}$. As $N + 2, N + 3, N + 4$ are composite, every prime factor of N is congruent to 1 or -1 modulo 6. If every prime factor of N was congruent to 1, then $N \equiv 1 \pmod{6}$: this is a contradiction because $-1 \not\equiv 1 \pmod{6}$. So there exists a prime factor p of N such that $p \equiv -1 \pmod{6}$.

If $p \leq n$, then $p \mid n!$, and $p \mid N = n! - 1$, so $p \mid 1$. As p is prime, this is a contradiction, so $p > n$.

Conclusion :

for any integer n , there exists a prime $p > n$ such that $p \equiv -1 \pmod{6}$: there are infinitely many primes congruent to -1 modulo 6. □

Ex. 3.2 Construct addition and multiplication tables for $\mathbb{Z}/5\mathbb{Z}$, $\mathbb{Z}/8\mathbb{Z}$, and $\mathbb{Z}/10\mathbb{Z}$.

Proof. More a latex exercise than a mathematical one.

$\mathbb{Z}/5\mathbb{Z}$:

+	0	1	2	3	4	×	0	1	2	3	4
0	0	1	2	3	4	0	0	0	0	0	0
1	1	2	3	4	0	1	0	1	2	3	4
2	2	3	4	0	1	2	0	2	4	1	3
3	3	4	0	1	2	3	0	3	1	4	2
4	4	0	1	2	3	4	0	4	3	2	1

$\mathbb{Z}/8\mathbb{Z}$:

+	0	1	2	3	4	5	6	7	×	0	1	2	3	4	5	6	7
0	0	1	2	3	4	5	6	7	0	0	0	0	0	0	0	0	0
1	1	2	3	4	5	6	7	0	1	0	1	2	3	4	5	6	7
2	2	3	4	5	6	7	0	1	2	0	2	4	6	0	2	4	6
3	3	4	5	6	7	0	1	2	3	0	3	6	1	4	7	2	5
4	4	5	6	7	0	1	2	3	4	0	4	0	4	0	4	0	4
5	5	6	7	0	1	2	3	4	5	0	5	2	7	4	1	6	3
6	6	7	0	1	2	3	4	5	6	0	6	4	2	0	6	4	2
7	7	0	1	2	3	4	5	6	7	0	7	6	5	4	3	2	1

$\mathbb{Z}/10\mathbb{Z}$:

+	0	1	2	3	4	5	6	7	8	9	×	0	1	2	3	4	5	6	7	8	9
0	0	1	2	3	4	5	6	7	8	9	0	0	0	0	0	0	0	0	0	0	0
1	1	2	3	4	5	6	7	8	9	0	1	0	1	2	3	4	5	6	7	8	9
2	2	3	4	5	6	7	8	9	0	1	2	0	2	4	6	8	0	2	4	6	8
3	3	4	5	6	7	8	9	0	1	2	3	0	3	6	9	2	5	8	1	4	7
4	4	5	6	7	8	9	0	1	2	3	4	0	4	8	2	6	0	4	8	2	6
5	5	6	7	8	9	0	1	2	3	4	5	0	5	0	5	0	5	0	5	0	5
6	6	7	8	9	0	1	2	3	4	5	6	0	6	2	8	4	0	6	2	8	4
7	7	8	9	0	1	2	3	4	5	6	7	0	7	4	1	8	5	2	9	6	3
8	8	9	0	1	2	3	4	5	6	7	8	0	8	6	4	2	0	8	6	4	2
9	9	0	1	2	3	4	5	6	7	8	9	0	9	8	7	6	5	4	3	2	1

Python code to generate the latex code to create such an array :

```
n= 10

print('$')
ligne = '\\begin{array}{c|'+ n*'c|'+'}'
print(ligne)
ligne='\\times'
for j in range(n):
    ligne += ' & ' + str(j)
ligne += '\\\\'
ligne += '\\\\'
ligne += ' \\\hline'
print(ligne)
```

```

for i in range(n):
    ligne = str(i)
    for j in range(n):
        ligne += ' & ' + str((i * j) % n)
    ligne += '\\\n'
    ligne += '\\\n'
    print(ligne)
print('\\end{array}')
print('$')

```

□

Ex. 3.3 Let abc be the decimal representation for an integer between 1 and 1000. Show that abc is divisible by 3 iff $a + b + c$ is divisible by 3. Show that the same result is true if we replace 3 by 9. Show that abc is divisible by 11 iff $a - b + c$ is divisible by 11. Generalize to any number written in decimal notation.

Proof. Let $n = \overline{abc}$ the decimal representation of n .

As $10 \equiv 1 \pmod{3}$, $10^3 \equiv 10^2 \equiv 10 \equiv 1 \pmod{3}$, so

$$\begin{aligned}
 3 \mid n &\iff 10^3a + 10^2b + c \equiv 0 \pmod{3} \\
 &\iff a + b + c \equiv 0 \pmod{3} \qquad 3 \mid a + b + c
 \end{aligned}$$

As $10 \equiv 1 \pmod{9}$ the same demonstration is true for the result

$$9 \mid n \iff 9 \mid a + b + c.$$

Similarly, $10 \equiv -1 \pmod{11}$, and $10^2 \equiv 1, 10^3 \equiv -1$, so

$$\begin{aligned}
 11 \mid n &\iff 10^3a + 10^2b + c \equiv 0 \pmod{11} \\
 &\iff a - b + c \equiv 0 \pmod{11}
 \end{aligned}$$

More generally, let $n = \overline{a_l a_{l-1} \cdots a_0}$ is the decimal representation of n .

$10^n \equiv 1 \pmod{9}$, so

$$\begin{aligned}
 9 \mid n &\iff \sum_{k=0}^l a_k 10^k \equiv 0 \pmod{9} \\
 &\iff \sum_{k=0}^l a_k \equiv 0 \pmod{9} \\
 &\iff 9 \mid a_0 + a_1 + \cdots + a_n
 \end{aligned}$$

□

Ex. 3.4 Show that the equation $3x^2 + 2 = y^2$ has no solution in integers.

Proof. If $3x^2 + 2 = y^2$, then $\bar{y}^2 = \bar{2}$ in $\mathbb{Z}/3\mathbb{Z}$.

As $\{-1, 0, 1\}$ is a complete set of residues modulo 3, the squares in $\mathbb{Z}/3\mathbb{Z}$ are $\bar{0} = \bar{0}^2$ and $\bar{1} = \bar{1}^2 = \bar{(-1)}^2$, so $\bar{2}$ is not a square in $\mathbb{Z}/3\mathbb{Z}$: $\bar{y}^2 = \bar{2}$ is impossible in $\mathbb{Z}/3\mathbb{Z}$.

Thus $3x^2 + 2 = y^2$ has no solution in integers.

□

Ex. 3.5 Show that the equation $7x^2 + 2 = y^3$ has no solution in integers.

Proof. If $7x^2 + 2 = y^3$, $x, y \in \mathbb{Z}$, then $y^3 \equiv 2 \pmod{7}$ (so $y \not\equiv 0 \pmod{7}$)

From Fermat's Little Theorem, $y^6 \equiv 1 \pmod{7}$, so $2^2 \equiv y^6 \equiv 1 \pmod{7}$, which implies $7 \mid 2^2 - 1 = 3$: this is a contradiction. Thus the equation $7x^2 + 2 = y^3$ has no solution in integers. \square

Ex. 3.6 Let an integer $n > 0$ be given. A set of integers $a_1, \dots, a_{\phi(n)}$ is called a reduced residue system modulo n if they are pairwise incongruent modulo n and $(a_i, n) = 1$ for all i . If $(a, n) = 1$, prove that $aa_1, aa_2, \dots, aa_{\phi(n)}$ is again a reduced residue system modulo n .

Proof. Let $a_1, \dots, a_{\phi(n)}$ a reduced residue system modulo n .

- As $a \wedge n = 1$ and $a_i \wedge n = 1$, $i = 1, 2, \dots, \phi(n)$, then $aa_i \wedge n = 1$.
- As $a \wedge n = 1$, there exists $a' \in \mathbb{Z}$ such that $aa' \equiv 1 \pmod{n}$. then

$$aa_i \equiv aa_j \Rightarrow a'aa_i \equiv a'aa_j \pmod{n} \Rightarrow a_i \equiv a_j \pmod{n}.$$

So $i \neq j \Rightarrow a_i \not\equiv a_j \Rightarrow aa_i \not\equiv aa_j$:

$aa_1, \dots, aa_{\phi(n)}$ a reduced residue system modulo n .

Note that $\{a_1, a_2, \dots, a_{\phi(n)}\}$ is a reduced residue system modulo n if and only if $\{\overline{a_1}, \overline{a_2}, \dots, \overline{a_{\phi(n)}}\} = U(\mathbb{Z}/n\mathbb{Z})$. \square

Ex. 3.7 Use Ex. 2.6 to give another proof of Euler's theorem, $a^{\phi(n)} \equiv 1 \pmod{n}$ for $(a, n) = 1$.

Proof. The proof is more clear if we stay in $\mathbb{Z}/n\mathbb{Z}$.

$$\text{Let } P = \prod_{x \in U(\mathbb{Z}/n\mathbb{Z})} x$$

(if $\{a_1, \dots, a_{\phi(n)}\}$ is a reduced residue system modulo n , then $\overline{P} = \prod_{i=1}^{\phi(n)} a_i$.)

Let $a \in \mathbb{Z}$ such that $a \wedge n = 1$, then $b = \overline{a} \in U(\mathbb{Z}/n\mathbb{Z})$, and

$$\psi \left\{ \begin{array}{ccc} U(\mathbb{Z}/n\mathbb{Z}) & \rightarrow & U(\mathbb{Z}/n\mathbb{Z}) \\ x & \mapsto & bx \end{array} \right.$$

- $\psi(x) = \psi(x') \Rightarrow bx = bx' \Rightarrow b^{-1}bx = b^{-1}bx' \Rightarrow x = x'$ so ψ is injective.
 - Let $y \in U(\mathbb{Z}/n\mathbb{Z})$. If $x = b^{-1}y$, then $\psi(x) = bb^{-1}y = y$, so ψ is surjective.
- ψ is a bijection, so

$$\prod_{x \in U(\mathbb{Z}/n\mathbb{Z})} bx = \prod_{x \in U(\mathbb{Z}/n\mathbb{Z})} x,$$

that is

$$b^{\phi(n)} \prod_{x \in U(\mathbb{Z}/n\mathbb{Z})} x = \prod_{x \in U(\mathbb{Z}/n\mathbb{Z})} x.$$

As $\prod_{x \in U(\mathbb{Z}/n\mathbb{Z})} x$ is invertible,

$$b^{\phi(n)} = 1.$$

That is $\overline{a}^{\phi(n)} = 1$: for all $a \in \mathbb{Z}$, if $a \wedge n = 1$, then $a^{\phi(n)} \equiv 1 \pmod{n}$. \square

Ex. 3.8 Let p be an odd prime. If $k \in \{1, 2, \dots, p-1\}$, show that there is a unique b_k in this set such that $kb_k \equiv 1 \pmod{p}$. Show that $k \neq b_k$ unless $k = 1$ or $k = p-1$.

Proof. • existence.

As p is prime and $1 \leq k \leq p-1$, $k \wedge p = 1$, so there exist $\lambda_k, \mu_k \in \mathbb{Z}$ such that $\lambda_k p + \mu_k k = 1$. Let $b_k \in \{0, 1, \dots, p-1\}$ such that $b_k \equiv \mu_k \pmod{p}$. Then $kb_k \equiv 1$, and $b_k \not\equiv 0 \pmod{p}$, so $1 \leq b_k \leq p-1$.

• unicity. If $kb_k \equiv kb'_k \pmod{p}$, where $b_k, b'_k \in \{1, 2, \dots, p-1\}$, then $p \mid k(b'_k - b_k)$, and $p \wedge k = 1$, so $p \mid b'_k - b_k$. $b'_k \equiv b_k$, and $b_k, b'_k \in \{1, 2, \dots, p-1\}$, so $b_k = b'_k$.

If p is a prime number, and $k \in \{1, 2, \dots, p-1\}$, there is a unique b_k in $\{1, 2, \dots, p-1\}$ such that $kb_k \equiv 1 \pmod{p}$.

If $k = b_k$, then $k^2 \equiv 1 \pmod{p}$, so $p \mid (k-1)(k+1)$, and p is a prime, thus $p \mid k-1$ or $p \mid k+1$, that is $k \equiv \pm 1 \pmod{p}$. As $1 \leq k \leq p-1$, $k = 1$ or $k = p-1$ (and $1^2 \equiv (p-1)^2 \equiv 1 \pmod{p}$). \square

Ex. 3.9 Use Ex. 3.8 to prove that $(p-1)! \equiv -1 \pmod{p}$. (misprint corrected)

Proof. Each element k in the product $p!$ can be associated with its inverse b_k modulo k , except 1 and $p-1$, which are their own inverse, so

$$p! \equiv 1 \times (p-1) \equiv -1 \pmod{p}.$$

\square

Ex. 3.10 If n is not a prime, show that $(n-1)! \equiv 0 \pmod{n}$, except when $n = 4$.

Proof. Suppose that $n > 1$ is not a prime. Then $n = uv$, where $2 \leq u \leq v \leq n-1$.

• If $u \neq v$, then $n = uv \mid (n-1)! = 1 \times 2 \times \dots \times u \times \dots \times v \times \dots \times (n-1)$ (even if $u \wedge v \neq 1$!).

• If $u = v$, $n = u^2$ is a square.

If u is not prime, $u = st$, $2 \leq s \leq t \leq u-1 \leq n-1$, and $n = u'v'$, where $u' = s, v' = st^2$ verify $2 \leq u' < v' \leq n-1$. As in the first case, $n = u'v' \mid (n-1)!$.

If $u = p$ is a prime, then $n = p^2$.

In the case $p = 2$, $n = 4$ and $n = 4 \nmid (n-1)! = 6$. In the other case $p > 2$, and $(n-1)! = (p^2-1)!$ contains the factors $p < 2p < p^2$, so $p^2 \mid (p^2-1)!, n \mid (n-1)!$.

Conclusion : if n is not a prime, $(n-1)! \equiv 0 \pmod{n}$, except when $n = 4$. \square

Ex. 3.11 Let $a_1, \dots, a_{\phi(n)}$ be a reduced residue system modulo n and let N be the number of solutions to $x^2 \equiv 1 \pmod{n}$. Prove that $a_1 \cdots a_{\phi(n)} \equiv (-1)^{N/2} \pmod{n}$.

Proof. If $n = 2$, then $N = 1$ and the result is false. So we suppose $n > 2$.

Let H the subset of $\mathbb{Z}/n\mathbb{Z}$ of all $x \in \mathbb{Z}/n\mathbb{Z}$ such that $x^2 = 1$:

$$H = \{x \in \mathbb{Z}/n\mathbb{Z} \mid x^2 = 1\}$$

(here $1 = \bar{1}$).

$H \subset U(\mathbb{Z}/n\mathbb{Z})$, and $x \in H, y \in H \Rightarrow x^2 = y^2 = 1 \Rightarrow (xy^{-1})^2 = 1 \Rightarrow xy^{-1} \in H$, so H is a subgroup of $(U(\mathbb{Z}/n\mathbb{Z}), \times)$, and $N = \text{Card } H$.

Each $x \in U(\mathbb{Z}/n\mathbb{Z})$ such that $x \notin H$ can be paired with its inverse x^{-1} , and $xx^{-1} = 1$, so

$$P := \prod_{x \in U(\mathbb{Z}/n\mathbb{Z})} x = \prod_{x \in H} x.$$

If $x \in H$, $-x \in H$.

• If n is odd, each $x = \bar{a} \in H$ ($a \in \mathbb{Z}, 1 \leq a \leq n-1$) satisfy $-x \neq x$: otherwise $2a \equiv 0 \pmod{n}$, $2a = kn, k \in \mathbb{Z}$. As $0 < 2a = kn < 2n$, then $k = 1$, and $n = 2a$ is even, which is in contradiction with the hypothesis.

So we each $x \in H$ can be paired with $-x$ in the product P , and $x(-x) = -1$, so

$$P = \prod_{x \in H} x = (-1)^{N/2}.$$

• If n is even, if $x = \bar{a} \in H$ ($a \in \mathbb{Z}, 1 \leq a \leq n-1$) satisfy $x = -x$, then $0 < a = k\frac{n}{2} < n$, so $a = \frac{n}{2}$, and $x = \overline{(\frac{n}{2})}$ is the only element in $\mathbb{Z}/n\mathbb{Z}$ such that $x = -x$. $2x = \bar{0}$, and $x \in H$, so $2x^2 = \bar{0}, 2 = \bar{0}$: since $n > 2$, this is impossible, so $x \neq -x$ for all $x \in H$, and $\prod_{x \in H} x = (-1)^{N/2}$.

Conclusion: if $n > 2$,

$$\prod_{x \in U(\mathbb{Z}/n\mathbb{Z})} x = (-1)^{N/2}.$$

If $a_1, \dots, a_{\phi(n)}$ is a reduced residue system modulo n , then $\overline{a_1 \cdots a_{\phi(n)}} = P = \prod_{x \in U(\mathbb{Z}/n\mathbb{Z})} x = (-1)^{N/2}$, so

$$a_1 \cdots a_{\phi(n)} \equiv (-1)^{N/2}.$$

□

Ex. 3.12 Let $\binom{p}{k} = \frac{p!}{k!(p-k)!}$ be a binomial coefficient, and suppose p is prime. If $1 \leq k \leq p-1$, show that p divides $\binom{p}{k}$. Deduce $(a+b)^p \equiv a^p + b^p \pmod{p}$.

Proof. $p \mid p! = k!(p-k)!\binom{p}{k}$.

If $1 \leq k \leq p-1$, then for each $i, 1 \leq i \leq k, 1 \leq i < p$, so $i \wedge p = 1$. Thus $\left(\prod_{i=1}^k i\right) \wedge p = 1, k! \wedge p = 1$. Similarly, $p-k < p$, so $\left(\prod_{i=1}^{p-k} i\right) \wedge p = 1, (p-k)! \wedge p = 1$. Thus $p \wedge k!(p-k)! = 1$, and $p \mid p! = k!(p-k)!\binom{p}{k}$, so $p \mid \binom{p}{k}$.

Finally, from binomial formula

$$\begin{aligned} (a+b)^p &= a^p + \sum_{k=1}^{p-1} \binom{p}{k} a^k b^{p-k} + b^p \\ &\equiv a^p + b^p \pmod{p} \end{aligned}$$

□

Ex. 3.13 Use Ex. 3.12 to give another proof of Fermat's theorem, $a^{p-1} \equiv 1 \pmod{p}$ if p does not divide a .

Proof. If we make the induction hypothesis

$$\mathcal{P}(k) \iff \forall(a_1, a_2, \dots, a_k) \in \mathbb{Z}^k, (a_1 + a_2 + \cdots + a_k)^p \equiv a_1^p + a_2^p + \cdots + a_k^p$$

(which is true for $k = 1, k = 2$) then, from induction hypothesis and the case $k = 2$ already proved in Ex 3.12,

$$\begin{aligned}(a_1 + a_2 + \cdots + a_k + a_{k+1})^p &= ((a_1 + a_2 + \cdots + a_k) + a_{k+1})^p \\ &\equiv (a_1 + a_2 + \cdots + a_k)^p + a_{k+1}^p \pmod{p} \\ &\equiv a_1^p + a_2^p + \cdots + a_k^p + a_{k+1}^p \pmod{p}\end{aligned}$$

so $\mathcal{P}(k) \Rightarrow \mathcal{P}(k+1)$:

$$\forall k \in \mathbb{N}^*, \forall (a_1, a_2, \dots, a_k) \in \mathbb{Z}^k, (a_1 + a_2 + \cdots + a_k)^p \equiv a_1^p + a_2^p + \cdots + a_k^p.$$

If we apply this result to the particular case $a_1 = a_2 = \cdots = a_k = 1$, we obtain

$$\forall k \in \mathbb{N}^*, k^p \equiv k \pmod{p}.$$

and $(-k)^p \equiv -k^p \equiv -k \pmod{p}$ (even if $p = 2$), and $0^p = 0$, so

$$\forall k \in \mathbb{Z}, k^p \equiv k \pmod{p}.$$

If $p \nmid a, a \in \mathbb{Z}$, then $p \wedge a = 1$, and $p \mid a^p - a = a(a^{p-1} - 1)$, so $p \mid a^{p-1} - 1, a^{p-1} \equiv 1 \pmod{p}$: this is another proof of Fermat's theorem. \square

Ex. 3.14 Let p and q be distinct odd primes such that $p-1$ divides $q-1$. If $(n, pq) = 1$, show that $n^{q-1} \equiv 1 \pmod{pq}$.

Proof. As $n \wedge pq = 1, n \wedge p = 1, n \wedge q = 1$, so from Fermat's Little Theorem

$$n^{q-1} \equiv 1 \pmod{q}, \quad n^{p-1} \equiv 1 \pmod{p}.$$

$p-1 \mid q-1$, so there exists $k \in \mathbb{Z}$ such that $q-1 = k(p-1)$. Thus

$$n^{q-1} = (n^{p-1})^k \equiv 1 \pmod{p}.$$

$p \mid n^{q-1} - 1, q \mid n^{q-1} - 1$, and $p \wedge q = 1$, so $pq \mid n^{q-1} - 1$:

$$n^{q-1} \equiv 1 \pmod{pq}.$$

\square

Ex. 3.15 For any prime p show that the numerator of $1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{p-1}$ is divisible by p .

Proof. As the result is false for $p = 2$, we must suppose $p > 2$, so p is odd.

$$1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{p-1} = \frac{N}{D}, \text{ where}$$

$$N = (p-1)! + \frac{(p-1)!}{2} + \cdots + \frac{(p-1)!}{p-1}, \quad D = (p-1)!.$$

From Wilson's theorem, $(p-1)! \equiv -1 \pmod{p}$, so in the field $\mathbb{Z}/p\mathbb{Z}$,

$$\overline{N} = (-\overline{1})(\overline{1}^{-1} + \overline{2}^{-1} + \cdots + \overline{p-1}^{-1}).$$

As the application $\varphi : (\mathbb{Z}/p\mathbb{Z})^* \rightarrow (\mathbb{Z}/p\mathbb{Z})^*, x \mapsto x^{-1}$ is bijective (it's an involution),

$$\overline{1} + \overline{2}^{-1} + \cdots + \overline{p-1}^{-1} = \overline{1} + \overline{2} + \cdots + \overline{p-1} = \overline{p} \times \overline{\left(\frac{p-1}{2}\right)} = \overline{0}.$$

So $p \mid N$, and $p \wedge (p-1)! = 1$, that is $p \wedge D = 1$. Thus p divides the numerator of the reduced fraction of N/D . \square

Ex. 3.16 Use the proof of the Chinese Remainder Theorem to solve the system $x \equiv 1 \pmod{7}$, $x \equiv 4 \pmod{9}$, $x \equiv 3 \pmod{5}$.

Proof. Let $m_1 = 7, m_2 = 9, m_3 = 5, m = m_1 m_2 m_3 = 315, n_1 = m/m_1 = m_2 m_3 = 45, n_2 = m_1 m_3 = 35, n_3 = m_1 m_2 = 63$.

If $r_1 = 13, s_1 = -2$, then $r_1 m_1 + s_1 n_1 = 13m_1 - 2m_2 m_3 = 13 \times 7 - 2 \times 45 = 1$,
so $e_1 = s_1 n_1 = -2 \times 45 = -90$ verifies

$$e_1 = -90, \quad e_1 \equiv 1 \pmod{7}, e_1 \equiv 0 \pmod{9}, e_1 \equiv 0 \pmod{5}.$$

If $r_2 = 4, s_2 = -1$, then $r_2 m_2 + s_2 n_2 = 4 \times 9 - 1 \times 35 = 1$,
so $e_2 = s_2 n_2 = -35$ verifies

$$e_2 = -35, \quad e_2 \equiv 0 \pmod{7}, e_2 \equiv 1 \pmod{9}, e_2 \equiv 0 \pmod{5}.$$

If $r_3 = -25, s_3 = 2$, then $r_3 m_3 + s_3 n_3 = -25 \times 5 + 2 \times 63 = 1$,
so $e_3 = s_3 n_3 = 2 \times 63 = 126$ verifies

$$e_3 = 126, \quad e_3 \equiv 0 \pmod{7}, e_3 \equiv 0 \pmod{9}, e_3 \equiv 1 \pmod{5}.$$

Let $x_0 = e_1 + 4e_2 + 3e_3 = 148$: then

$$x_0 = 148, \quad x_0 \equiv 1 \pmod{7}, x_0 \equiv 4 \pmod{9}, x_0 \equiv 3 \pmod{5}.$$

If $x \in \mathbb{Z}$ is any solution of the system, then $7 \mid x - x_0, 9 \mid x - x_0, 5 \mid x - x_0$, with $7 \wedge 9 = 7 \wedge 5 = 9 \wedge 5 = 1$, so $m = 315 \mid x - x_0$:

$$x = 148 + k 315, k \in \mathbb{Z},$$

and all these integers are solutions of the system. □

Ex. 3.17 Let $f(x) \in \mathbb{Z}[x]$ and $n = p_1^{a_1} \cdots p_t^{a_t}$. Show that $f(x) \equiv 0 \pmod{n}$ has a solution iff $f(x) \equiv 0 \pmod{p_i^{a_i}}$ has a solution for $i = 1, \dots, t$.

Proof. If x is such that $f(x) \equiv 0 \pmod{n}$, as $p_i^{a_i} \mid n$, $f(x) \equiv 0 \pmod{p_i^{a_i}}$.

Reciprocally, let x_1, x_2, \dots, x_t such that

$$f(x_1) \equiv 0 \pmod{p_1^{a_1}}$$

...

$$f(x_t) \equiv 0 \pmod{p_t^{a_t}}$$

As $p_i^{a_i} \wedge p_j^{a_j} = 1$ if $i \neq j$, the Chinese Remainder Theorem gives an integer x such that $x \equiv x_i \pmod{p_i^{a_i}}$, $i = 1, 2, \dots, t$. As $f(x) \in \mathbb{Z}[x]$, $f(x) \equiv f(x_i) \equiv 0 \pmod{p_i^{a_i}}$. So $p_i^{a_i} \mid f(x)$, $i = 1, 2, \dots, t$, where $p_i^{a_i} \wedge p_j^{a_j} = 1$ if $i \neq j$, then $n = p_1^{a_1} \cdots p_t^{a_t} \mid f(x)$: x is a solution of $f(x) \equiv 0 \pmod{n}$.

Conclusion : $f(x) \equiv 0 \pmod{n}$ has a solution iff $f(x) \equiv 0 \pmod{p_i^{a_i}}$ has a solution for $i = 1, \dots, t$. □

Ex. 3.18 For $f \in \mathbb{Z}[x]$, let N be the number of solutions to $f(x) \equiv 0 \pmod{n}$ and N_i be the number of solutions to $f(x) \equiv 0 \pmod{p_i^{a_i}}$. Prove that $N = N_1 N_2 \cdots N_t$.

Proof. Note $[x]_n$ the class of x modulo n . Let S the set of solutions in $\mathbb{Z}/n\mathbb{Z}$ of $f(\bar{x}) = 0$, and S_i the set of solutions in $\mathbb{Z}/p_i^{a_i}\mathbb{Z}$ of $f(\bar{x}) = 0$.

(We designate with the same letter the polynomial f in $\mathbb{Z}[x]$ or its reduction in $\mathbb{Z}/n\mathbb{Z}[x]$.)

Let

$$\varphi : \begin{cases} S & \rightarrow S_1 \times S_2 \times \cdots \times S_t \\ [x]_n & \mapsto ([x]_{p_1^{a_1}}, [x]_{p_2^{a_2}}, \dots, [x]_{p_t^{a_t}}) \end{cases}$$

• φ is well defined : if $x \equiv x' \pmod{n}$, then $x \equiv x' \pmod{p_i^{a_i}}$, $i = 1, 2, \dots, t$, so $([x]_{p_1^{a_1}}, [x]_{p_2^{a_2}}, \dots, [x]_{p_t^{a_t}}) = ([x']_{p_1^{a_1}}, [x']_{p_2^{a_2}}, \dots, [x']_{p_t^{a_t}})$. Moreover, we proved in Ex 3.17 that $[x]_n \in S \Rightarrow [x]_{p_i^{a_i}} \in S_i$.

• φ is injective : if $([x]_{p_1^{a_1}}, [x]_{p_2^{a_2}}, \dots, [x]_{p_t^{a_t}}) = ([x']_{p_1^{a_1}}, [x']_{p_2^{a_2}}, \dots, [x']_{p_t^{a_t}})$, then $p_i^{a_i} \mid x' - x$, $i = 1, 2, \dots, t$, so $n \mid x' - x$ and $[x]_n = [x']_n$.

• φ is surjective : if $y = ([x_1]_{p_1^{a_1}}, [x_2]_{p_2^{a_2}}, \dots, [x_t]_{p_t^{a_t}})$ is any element of $S_1 \times S_2 \times \cdots \times S_t$, there exists from Chinese remainder theorem $x \in \mathbb{Z}$ such that $x \equiv x_i \pmod{p_i^{a_i}}$. Then $\varphi([x]_n) = y$ (see Ex. 3.17).

In conclusion, φ is bijective, $N = |S| = |S_1 \times S_2 \times \cdots \times S_t| = N_1 N_2 \cdots N_t$. \square

Ex. 3.19 If p is an odd prime, show that 1 and -1 are the only solutions of $x^2 \equiv 1 \pmod{p^a}$.

Proof.

$$x^2 - 1 \pmod{p^a} \iff p^a \mid (x-1)(x+1).$$

Let $d = (x-1) \wedge (x+1)$: $d = 1$ or $d = 2$.

• If $d = 1$, then x is even (if not, $x-1$ and $x+1$ are even, and $2 \mid d$). As $p^a \mid (x-1)(x+1)$ and $(x-1) \wedge (x+1) = 1$, then $p^a \mid x-1$, or $p^a \mid x+1$, that is

$$x \equiv \pm 1 \pmod{p^a}.$$

• If $d = 2$, then x is odd, and

$$p^a \mid 4 \frac{x-1}{2} \frac{x+1}{2}.$$

As p is an odd prime, $p \wedge 4 = 1$, so $p \mid \frac{x-1}{2} \frac{x+1}{2}$, where $\frac{x-1}{2} \wedge \frac{x+1}{2} = 1$, hence $p^a \mid \frac{x-1}{2} \mid x-1$ or $p^a \mid \frac{x+1}{2} \mid x+1$:

$$x \equiv \pm 1 \pmod{p^a}.$$

$\{-\bar{1}, \bar{1}\}$ is the set of roots of $x^2 - \bar{1}$ in $\mathbb{Z}/p^a\mathbb{Z}$. \square

Ex. 3.20 Show that $x^2 \equiv 1 \pmod{2^b}$ has one solution if $b = 1$, two solutions if $b = 2$, and four solutions if $b \geq 3$.

Proof. Consider the equation $x^2 \equiv 1 \pmod{2^b}$.

• If $b = 1$, $x^2 \equiv 1 \pmod{2} \iff 2 \mid (x-1)(x+1) \iff x \equiv 1 \pmod{2}$: one solution.

• If $b = 2$, as $0^2 \equiv 2^2 \equiv 0 \pmod{4}$, $x^2 \equiv 1 \pmod{4} \iff x \equiv \pm 1 \pmod{4}$: two solutions.

- Suppose $b \geq 3$. The equation has 4 solutions $1, -1, 1 + 2^{b-1}, -1 + 2^{b-1}$.
Indeed, $(\pm 1)^2 \equiv 1 \pmod{2^b}$, and

$$(1 + 2^{b-1})^2 = 1 + 2 \cdot 2^{b-1} + 2^{2b-2} = 1 + 2^b(1 + 2^{b-2}) \equiv 1 \pmod{2^b},$$

and similarly $(-1 + 2^{b-1})^2 \equiv 1 \pmod{2^b}$.

These solutions are incongruent modulo 2^b :

$1 \not\equiv -1 \pmod{2^b}$ and $1 + 2^{b-1} \not\equiv -1 + 2^{b-1}$ (if not, $2^b \mid 2$, so $b \leq 1$).

$1 + 2^{b-1} \equiv -1 \pmod{2^b} \iff 2^b \mid 2 + 2^{b-1} = 2(1 + 2^{b-2})$: so $2 \mid 2^{b-1} \mid (1 + 2^{b-2})$, this is impossible because $1 + 2^{b-2}$ is odd ($b \geq 3$). With the same argument, $-1 + 2^{b-1} \not\equiv 1 \pmod{2^b}$. $1 + 2^{b-1} \equiv 1 \pmod{2^b}$ implies $2^b \mid 2^{b-1}$, so $2 \mid 1$: this is a contradiction, so $1 + 2^{b-1} \not\equiv 1 \pmod{2^b}$, and also $-1 + 2^{b-1} \not\equiv -1 \pmod{2^b}$. There exist at least 4 solutions.

We show that these are the only solutions :

$$\forall x \in \mathbb{Z}, x^2 \equiv 1 \pmod{2^b} \Rightarrow x \equiv \pm 1 \pmod{2^{b-1}}.$$

Indeed, if $x^2 \equiv 1 \pmod{2^b}$, $2^b \mid (x-1)(x+1)$, where $d = (x-1) \wedge (x+1) = 2$.

As in Ex.3.19, if $d = 1$, then $2^b \mid x-1$ or $2^b \mid x+1$, a fortiori $x \equiv \pm 1 \pmod{2^{b-1}}$.

If $d = 2$, then x is odd, and $2^b \mid 4 \frac{x-1}{2} \frac{x+1}{2}$, so $2^{b-2} \mid \frac{x-1}{2} \frac{x+1}{2}$, with $\frac{x-1}{2} \wedge \frac{x+1}{2} = 1$, so $2^{b-2} \mid \frac{x-1}{2}$ or $2^{b-2} \mid \frac{x+1}{2}$, that is $2^{b-1} \mid x-1$ or $2^{b-1} \mid x+1$: $x \equiv \pm 1 \pmod{2^{b-1}}$.

(Alternatively, we can prove this implication by induction.)

Hence every solution of $x^2 \equiv 1 \pmod{2^b}$, $b \geq 3$ is such that $x = \pm 1 + k2^{b-1}$, $k \in \mathbb{Z}$: there exit only four such value in the interval $[0, 2^b[$, namely $1, -1 + 2^{b-1}, 1 + 2^{b-1}, -1 + 2^b$.

Conclusion : if $b \geq 3$, the roots of $x^2 - 1$ in $\mathbb{Z}/2^b\mathbb{Z}$ are $\bar{1}, -\bar{1}, \bar{1} + \bar{2}^{b-1}, -\bar{1} + \bar{2}^{b-1}$. \square

Ex. 3.21 Use Ex. 18-20 to find the number of solutions to $x^2 \equiv 1 \pmod{n}$.

Proof. Let $n = 2^{a_0} p_1^{a_1} \cdots p_k^{a_k}$ the decomposition in prime factors of $n > 1$ ($p_0 = 2 < p_1 < \cdots < p_k$, $a_0 \geq 0, a_i > 0, 1 \leq i \leq k$). Let N the number of solutions of $x^2 \equiv 1 \pmod{n}$, and N_i the number of solutions of $x^2 \equiv 1 \pmod{p_i^{a_i}}, i = 0, 1, \dots, k$. From Ex.3.18, we know that $N = N_0 N_1 \cdots N_k$, where (Ex. 3.19), $N_i = 2, i = 1, 2, \dots, k$, and (Ex.3.20), $N_0 = 1$ if $a_0 = 1$ (or $a_0 = 0$), $N_0 = 2$ if $a_0 = 2$, $N_0 = 4$ if $a_0 \geq 3$.

Conclusion : the number of solutions of $x^2 \equiv 1 \pmod{n}$, where $n = 2^{a_0} p_1^{a_1} \cdots p_k^{a_k}$, is

$$N = 2^k \quad \text{if } a_0 = 0 \text{ or } a_0 = 1$$

$$N = 2^{k+1} \quad \text{if } a_0 = 2$$

$$N = 2^{k+2} \quad \text{if } a_0 \geq 3$$

\square

Ex. 3.22 Formulate and prove the Chinese Remainder Theorem in a principal ideal domain.

Proposition. Let R a principal ideal domain, and $m_1, \dots, m_t \in R$. Suppose that $(m_i, m_j) = 1$ for $i \neq j$ (that is $(m_i) + (m_j) = (1)$, $m_i R + m_j R = R$). Let $b_1, \dots, b_t \in R$ and consider the system of congruences:

$$x \equiv b_1 \pmod{m_1}, x \equiv b_2 \pmod{m_2}, \dots, x \equiv b_t \pmod{m_t}.$$

This system has solutions and any two solutions differ by a multiple of $m_1 m_2 \cdots m_t$.

Proof. Let $m = m_1 m_2 \cdots m_t$, and $n_i = m/m_i$, $i = 1, 2, \dots, t$.

As $(m_1, m_i) = (1)$, we can find $u_i, v_i \in R$ such that $m_1 u_i + m_i v_i = 1$, $i = 2, \dots, t$.

So $1 = \prod_{i=2}^t (m_1 u_i + m_i v_i) = m_1 u + (m_2 \cdots m_t) v$ for some elements $u, v \in R$, thus $(m_1, n_1) = (m_1, m_2 m_3 \cdots m_t) = (1)$, and similarly $(m_i, n_i) = (1)$. So there are $r_i, s_i \in R$ such that $r_i m_i + s_i n_i = 1$. Let $e_i = s_i n_i$. Then $e_i \equiv 1 \pmod{m_i}$ and $e_i \equiv 0 \pmod{m_j}$ for $j \neq i$.

Set $x_0 = \sum_{i=1}^t b_i e_i$. Then we have $x_0 \equiv b_i e_i \equiv b_i \pmod{m_i}$ and so x_0 is a solution.

Suppose that x_1 is another solution. Then $x_1 - x_0 \equiv 0 \pmod{m_i}$ for $i = 1, 2, \dots, t$, in other words m_1, m_2, \dots, m_t divide $x_1 - x_0$, with $(m_i, m_j) = 1$: from lemma 2 generalized to principal rings, m divides $x_1 - x_0$. \square

This result can be generalized to any commutative ring, not necessarily a PID (see S.LANG, Algebra):

Proposition. *Let A a commutative ring. Let $\mathfrak{a}_1, \dots, \mathfrak{a}_n$ be ideals of A such that $\mathfrak{a}_i + \mathfrak{a}_j = A$ for all $i \neq j$. Given elements $x_1, \dots, x_n \in A$, there exists $x \in A$ such that $x \equiv x_i \pmod{\mathfrak{a}_i}$ for all i .*

Ex. 3.23 *Extend the notion of congruence to the ring $\mathbb{Z}[i]$ and prove that $a + bi$ is always congruent to 0 or 1 modulo $1 + i$.*

Proof. If a, b, c are in $\mathbb{Z}[i]$ we say that $a \equiv b \pmod{c}$ if there exists $q \in \mathbb{Z}[i]$ such that $a - b = qc$.

As $i \equiv -1 \pmod{1+i}$, $a + bi \equiv a - b \pmod{1+i}$.

$(1-i)(1+i) = 2$, so $2 \equiv 0 \pmod{1+i}$.

If $a - b$ is even, $a - b = 2k$, $k \in \mathbb{Z} \subset \mathbb{Z}[i]$, so $a - b \equiv 0 \pmod{1+i}$.

If $a - b$ is odd, $a - b = 2k + 1$, $k \in \mathbb{Z}$, so $a - b \equiv 1 \pmod{1+i}$.

Conclusion : for all $z \in \mathbb{Z}[i]$, $z \equiv 0, 1 \pmod{1+i}$. \square

Ex. 3.24 *Extend the notion of congruence to the ring $\mathbb{Z}[\omega]$ and prove that $a + b\omega$ is always congruent to $-1, 0$ or 1 modulo $1 - \omega$.*

Proof. Same definition of congruence in $\mathbb{Z}[\omega]$ as in Ex. 3.23.

$\omega \equiv 1 \pmod{1-\omega}$, so $a + b\omega \equiv a + b \pmod{1-\omega}$.

$0 = 1 - \omega^3 = (1 - \omega)(1 + \omega + \omega^2)$, with $1 - \omega \neq 0$, so $1 + \omega + \omega^2 = 0$. Hence $3 \equiv 0 \pmod{1 - \omega}$.

$a + b \equiv 0, 1, -1 \pmod{3}$, so $a + b \equiv 0, 1, -1 \pmod{1 - \omega}$

For all $z \in \mathbb{Z}[\omega]$, $z \equiv 0, 1, -1 \pmod{1 - \omega}$. \square

Ex. 3.25 *Let $\lambda = 1 - \omega \in \mathbb{Z}[\omega]$. If $\alpha \in \mathbb{Z}[\omega]$ and $\alpha \equiv 1 \pmod{\lambda}$, prove that $\alpha^3 \equiv 1 \pmod{9}$.*

Proof. $\alpha \equiv 1 \pmod{\lambda}$, so $\alpha = 1 + \beta\lambda$, $\beta \in \mathbb{Z}[\omega]$.

$\bar{\lambda} = 1 - \omega^2 = (1 - \omega)(1 + \omega) = -\omega^2(1 - \omega) = -\omega^2\lambda$ (so $\bar{\lambda}$ and λ are associate).

$$\begin{aligned}\alpha^3 - 1 &= (\alpha - 1)(\alpha - \omega)(\alpha - \omega^2) \\ &= (\alpha - 1)(\alpha - 1 + \lambda)(\alpha - 1 + \bar{\lambda}) \\ &= (\alpha - 1)(\alpha - 1 + \lambda)(\alpha - 1 - \omega^2\lambda) \\ &= \beta\lambda(\beta\lambda + \lambda)(\beta\lambda - \omega^2\lambda) \\ &= \lambda^3\beta(\beta + 1)(\beta - \omega^2)\end{aligned}$$

Moreover,

$$\begin{aligned}\beta(\beta + 1)(\beta - \omega^2) &\equiv \beta(\beta + 1)(\beta - 1) \pmod{\lambda} \\ &\equiv 0 \pmod{\lambda}\end{aligned}$$

since $\beta \equiv 0, 1, -1 \pmod{\lambda}$ (see Ex. 3.24).

So $\lambda^4 \mid \alpha^3 - 1$.

As $\lambda\bar{\lambda} = (1 - \omega)(1 - \omega^2) = 1 - \omega - \omega^2 + \omega^3 = 3$, then $\lambda\bar{\lambda} = -\omega^2\lambda^2 = 3$, so λ^2 and 3 are associate : $\lambda^2 = -\omega\lambda^2$. So $9 = (-\omega^2\lambda^2)^2 = \omega\lambda^4$, so $9 \mid \omega^2 9 = \lambda^4 \mid \alpha^3 - 1$.

For all $\alpha \in \mathbb{Z}[\omega]$,

$$\alpha \equiv 1 \pmod{\lambda} \Rightarrow \alpha^3 \equiv 1 \pmod{9}.$$

□

Ex. 3.26 Use Ex. 25 to show that ξ, η, ζ are not zero and $\xi^3 + \eta^3 + \zeta^3 = 0$, then λ divides at least one of the elements ξ, η, ζ .

Proof. Let $\xi, \eta, \zeta \in \mathbb{Z}[\omega] \setminus \{0\}$ such that $\xi^3 + \eta^3 + \zeta^3 = 0$.

With a reductio ad absurdum, suppose that $\lambda \nmid \xi, \lambda \nmid \eta, \lambda \nmid \zeta$.

From Ex. 3.24,

$$\xi \equiv \pm 1 \pmod{\lambda}, \eta \equiv \pm 1 \pmod{\lambda}, \zeta \equiv \pm 1 \pmod{\lambda},$$

and from Ex.3.25,

$$\xi^3 \equiv \pm 1 \pmod{9}, \eta^3 \equiv \pm 1 \pmod{9}, \zeta^3 \equiv \pm 1 \pmod{9},$$

As $\pm 1 \pm 1 \pm 1 \not\equiv 0 \pmod{9}$, this is a contradiction.

Conclusion : if ξ, η, ζ are not zero and $\xi^3 + \eta^3 + \zeta^3 = 0$, then λ divides at least one of the elements ξ, η, ζ .

(consequence : if $x^3 + y^3 + z^3 = 0$, $x, y, z \in \mathbb{Z}$, then $3 \mid xyz$: this is the first case of Fermat's theorem for the exposant 3.) □