

Intelligent Learning and Analysis Systems SS19, Exercise Sheet 2

by Andreas Hene, Niklas Mertens, Richard Palme.
Tutor: Maximilian Thiessen, Group 3

May 3, 2019

Exercise 3

Lemma 1

Each bucket has exactly size $w = \lceil \frac{1}{\varepsilon} \rceil$. \Rightarrow total number of buckets is

$$\frac{m}{w} = \frac{m}{\lceil 1/\varepsilon \rceil} \leq \varepsilon m$$

So the current bucket id is at most εm .

Lemma 2

Proof by induction.

Base Case: Let $b_{\text{current}} = 1$, let $(e, f, \Delta) \in \mathcal{D}$. Since there have been no deletions and no decrements of f until now, we know that f is the true frequency up until this point, i.e. $f = f_e$.

Also, (e, f, Δ) is only deleted when $f \leq 1$, in other words (e, f, Δ) is only deleted when $f_e = f \leq 1 = b_{\text{current}}$ which proves the base case.

Step Case: Let $k > 1$. Suppose that for $b_{\text{current}} < k$ we know that $f_e \leq b_{\text{current}}$ when (e, f, Δ) gets deleted.

Now let $b_{\text{current}} = k$ and $(e, f, \Delta) \in \mathcal{D}$. The true frequency of e in the buckets with ids $\Delta + 1, \dots, b_{\text{current}}$ is $\leq f + (b_{\text{current}} - 1 - \Delta)$, because f gets decremented by 1 at most $b_{\text{current}} - 1 - \Delta$ times.

Let b' be the bucket id where e was deleted the last time, if it exists. Else set $b' = 0$. By the induction hypothesis, the true frequency of e in buckets $1, \dots, b'$ is $\leq b'$, if b' exists. The true frequency in buckets $b' + 1, \dots, \Delta$ is 0. So the true frequency f_e at the current time (bucket id b_{current}) is at most $f_e \leq f + b_{\text{current}} - 1 - \Delta + b'$.

Since $b' \leq \Delta$, we get: $f_e \leq f + b_{\text{current}} - 1$.

If (e, f, Δ) gets deleted, we have $f \leq 1$, so $f_e \leq f + b_{\text{current}} - 1 \leq b_{\text{current}}$.

Lemma 3

If there is no entry for e in \mathcal{D} , then there are 2 cases:

Case 1: There has never been an entry for e in \mathcal{D} .

Then $f_e = 0 \leq \varepsilon m$.

Case 2: There has been an entry for e in \mathcal{D} before.

Let b be the bucket id when e was deleted the last time. Then $f_e \leq b$ by Lemma 2, and $f_e \leq b \leq \varepsilon m$ by Lemma 1.

Lemma 4

$f \leq f_e$, because f gets incremented by 1 at most f_e times.

If the last time e was deleted was after processing bucket b , then by Lemma 3 the true frequency of e in buckets $1, \dots, b$ is at most εm . The true frequency of e in buckets $b + 1, \dots, \Delta$ is 0, and the true frequency in buckets $\Delta + 1, \dots, b_{\text{current}}$ is at most f . So:

$$f_e \leq \varepsilon m + 0 + f = f + \varepsilon m$$

Exercise 4

d_i is the number of elements of \mathcal{D} that were last added to \mathcal{D} during processing of bucket $B - i + 1$. Call the i -th summand the contribution of bucket $B - i + 1$ to the sum. We want that the contribution of all summands does not exceed the total size of buckets $B - j + 1, \dots, B$ (which is jw).

So if element e was created during processing of $B - i + 1$, then e survives the deletion/decrementing process $i - 1$ times. This can only happen if e occurs at least i times in buckets $B - i + 1, \dots, B$. So e is allowed a contribution of i . So the contribution of all the elements last added to \mathcal{D} during processing of bucket $B - i + 1$ is allowed to be id_i , because this way the contribution of all summands can't exceed the total size of $B - j + 1, \dots, B$, i.e.

$$\sum_{i=1}^B id_i \leq jw$$