

National Survey of Sexual Attitudes and Lifestyles 2010-2012 (‘Natsal-3’) Codebook

Last updated: August 2015

For queries regarding the Natsal-3 codebook, please contact:

Dr Catherine Mercer
Centre for Sexual Health and HIV Research
Department of Primary Care & Population Sciences
University College London
Mortimer Market Centre
Capper Street
London
WC1E 6AU

Email: c.mercer@ucl.ac.uk
Tel: 020 7380 2072 (direct line)

CONTENTS

Topics	Face-to-face or CASI	Page number
Survey design	Administrative variables	p4
Geographic characteristics	Administrative variables	p7
General Health	Face-to-face	p9
Alcohol consumption	Face-to-face	p14
Smoking	Face-to-face	p17
Body mass index	Face-to-face	p17
Family history	Face-to-face	p19
Learning about sex	Face-to-face	p23
First sexual experiences	Face-to-face & CASI	p32
Contraception	Face-to-face	p44
Periods and menopause	Face-to-face	p53
CASI eligibility	Administrative variables	p56
Heterosexual behaviour	CASI	p60
Same-sex behaviour	CASI	p68
Heterosexual partnerships	CASI	p75
Same-sex partnerships	CASI	p80
Overlapping partnerships	CASI	p82
Online partnerships	CASI	p82
Total partnerships	CASI	p83
Detailed partnerships	CASI	p85
Travel and sex abroad	CASI	p105
Non-consensual sex	CASI	p108
Paying for sex	CASI	p109
Pregnancy History	CASI	p111
Unplanned pregnancy	CASI	p119
Family formation	CASI	P122
Fertility intentions and infertility	CASI	P127
Sexually Transmitted Infections	CASI	p128
HIV testing	CASI	p143
Sexual function	CASI	p149
Drug Use	CASI	p158
Mood and Wellbeing	CASI	p161
Circumcision	CASI & face-to-face	p162
Attitudes	Face-to-face	p163
Household classification	Face-to-face	p168
Previous live-in partnerships	Face-to-face	p174
Household composition	Face-to-face	p184
Employment	Face-to-face	p186
Qualifications	Face-to-face	p196
Parents occupation	Face-to-face	p201
Ethnicity	Face-to-face	p203
Sexual identity	Face-to-face	p207
Religion	Face-to-face	p208
Internet access	Face-to-face	P209
Urine survey	Face-to-face	p210
Saliva survey	Face-to-face	p214
Interview characteristics	Administrative variables	p215

Codebook notes

- This codebook comprises information on the Natsal-3 dataset, including variables not included in data sets held by the UK Data Archive. It also refers to the Natsal-1 and Natsal-2 data sets.
- Variables are presented in the codebook in order of the corresponding questions in the Natsal-3 questionnaire along with the details of any variables that are derived from these variables.
- If this is an electronic version of the codebook, then use 'Ctrl+F' and type in the variable of interest to go to the relevant section of the codebook
- The Natsal-3 variable name is preceded by the corresponding variable name in Natsal-2 (1999-2001) and Natsal-1 (1990-1991).
- The following characters identify how the Natsal-3 questions and/or responses relate to those used in Natsal-2 and how the Natsal-2 questions relate to those used in Natsal-1. Where applicable, these precede the variable names:
 - U Updated codes for different levels of responses (e.g. variable 'agrp')
 - * Question reworded in Natsal-3 / Natsal-2 since Natsal-2 / Natsal-1 (e.g. variable 'health')
 - N New question introduced in Natsal-3 / Natsal-2 that was not asked in Natsal-2 / Natsal-1
 - R Variable removed from Natsal 2010 that was asked in Natsal-2
 - D Variable derived from responses to 1+ interview question(s)
 - M Question moved in Natsal-3
- It is recommended that users consult the Natsal-3 and -2 technical reports for advice regarding the need for complex survey analyses.
- Missing values are generally coded as -1 if the question is not applicable to the respondent or 9/99/999/9999 if the question was not answered. The 'not answered' codes include respondents who weren't asked a question because they didn't answer a preceding question that was essential for the CAPI filtering. For example, respondents who refuse the CASI section of the interview will be coded as 'not answered' for all CASI questions, unless the question is known to be not applicable. Similarly a respondent will generally be coded as 'not answered' for derived variables if they didn't answer a variable used in the derivation.
- For ease of use, changes corresponding to the Natsal-3 questionnaire are written in blue
- The age range for Natsal-3 was 16-74 years. Therefore in some places in the questionnaire, derived variable definitions have been updated to reflect this. These are not routinely indicated in the codebook.

Making comparisons between the surveys

The version of the Natsal-2 data on the UK data archive was up-to-date at the time of archiving. However, the Natsal team have subsequently updated this dataset with additional variables. For example, the variables `new_finwt`, `newpsu`, `newstrata` (see pages 4 and 5) should be used instead of their old equivalents which are on the archived Natsal-2 data. Such variables are available on request from Dr Catherine Mercer (contact details on front page).

SURVEY DESIGN

(sn) **serial** **SIN2** full serial number

(N) **dateyoi** year of interview

(N) **datemoi** month of interview

(N) **dateoi** date of interview (day set to 01)
sysmiss if missing

Note: The quarters do not refer to a specific year

(R) (N) **quarter** fieldwork quarter

1: May to July

2: August to October

3: November to January

4: February to April

(R) (N) **sector** sector number

Note: Variable 'newpsu' in Natsal-2 was derived to be consistent with the variable 'psu' in Natsal-3. This variable was created subsequent to archiving the Natsal-2 data and is available from Dr Catherine Mercer (contact details on Page 1).

Note: Variable 'psu' should be declared as the PSU for analysis of the core dataset in Stata using Stata's survey commands. See the Natsal-3 Technical Report for further details.

(area) **psu** primary sampling unit (updated version)

(N) **psu_scrm** Scrambled psu identifier

(N) **newpsu** New Natsal-2 primary sampling unit, comparable with Natsal-2

Note: Subsequent to archiving, the Natsal-2 data was reweighted using the census 2001 data (which were not available when the Natsal-2 data were first created). This more accurate weight variable should be used for analysis of Natsal-2. It is available on request from Dr Catherine Mercer (contact details on page 1). It is available for the core sample only.

(N) **newfin_wt** updated weight for Natsal-2 (core sample only)

Note: Variable 'final_wt' should be used as the 'pweight' for analysis of the core dataset in Stata using Stata's survey commands. See the Natsal-2 Technical Report for details of its calculation.

(R) (ND) **final_wt** final weight in the core sample (Natsal-2)

(R) (N) **sample** ethnic cases in core

1: non-ethnic cases in the core

2: ethnic cases in the core used in the ethnic dataset

Note: For more information about the different sample types in Natsal-3 see the Technical Report

(U) **samptype** sample type: core or boost cases

1: Core (16-74)

2: Boost 1 (16-34)

3: Boost 2 (16-29)

Note: Variable 'eth_wt' should be used as the 'pweight' for analysis of the ethnic dataset in Stata using the survey commands.

(R) (ND) **eth_wt** weight for cases in the ethnic dataset

9999 if core sample

SURVEY DESIGN

Notes: 1. Natsal-2: Variable 'total_wt' is 'eth_wt' if the respondent was in the ethnic boost sample or a non-white case in the core sample but is 'final_wt' if the respondent was a white case in the core sample.
2. Natsal-2: Variable 'total_wt' has been scaled so that the weighted number of cases in the 'total' dataset (the core sample and the ethnic boost sample combined) is equal to the unweighted number of cases in the 'total' dataset ($n=12,110$).
3. Natsal-3: Variable 'total_wt' is an overall weight for the total sample (including boost of younger people)

(ND) (U) **total_wt** weight for cases in the 'total' dataset (core and boost samples combined)

Notes: 1. Natsal-2: Variable 'newstrata' (updated since archived version – see below) should be declared as the strata for analysis of the core dataset in Stata using Stata's survey commands.
2. Variable 'strata' in Natsal-1 was derived in Natsal-2 to be consistent with the variable 'strata' in Natsal-2
3. Variable 'strata' is derived from 6 age-groups ('agrp2'), 2 genders ('rsex'), and the 11 regions of 'gor' but consider London as 2 regions: inner and outer London making 12 regions in total, and 144 strata.
4. Variable 'newstrata' in Natsal-2 was derived to be consistent with the variable 'strata' in Natsal-3. This takes into account only the sampling stratification, and does not post-stratify by respondents' age and gender. This variable is available from Dr Catherine Mercer (contact details on Page 1). For more information about the sample design of Natsal please see the technical report.
5. Natsal-2 and Natsal-3: Grouped strata variable stratagrp should be declared as the strata for analysis of the core dataset in Stata using Stata's survey commands when there are too few PSUs in a stratum/strata to use 'newstrata' / 'strata' (Natsal-2 / Natsal-3)
6. If there are still too few PSUs per grouped stratum when using stratagrp, stratagrp2 should be used. Stratagrp3 can be used if the problem persists with stratagrp2.

(N) **strata** Natsal-3 strata variable

(N) **newstrata** New Natsal-2 strata variable, comparable with Natsal-3 variable

(N) **stratagrp** Grouped strata variable (every 2 strata combined)

(N) **stratagrp2** Group strata variable (every 3 strata combined)

(N) **stratagrp3** Group strata variable (every 4 strata combined)

(R) (N) **strata** 144 strata

1: 16-19, male & North East

2: 16-19, male & North West (incl. the old Merseyside region)

3: 16-19, male & Yorkshire & Humberside

4: 16-19, male & East Midlands

5: 16-19, male & West Midlands

6: 16-19, male & South West

7: 16-19, male & Eastern

8: 16-19, male & Inner London

9: 16-19, male & Outer London

10: 16-19, male & South East

11: 16-19, male & Wales

12: 16-19, male & Scotland

13: 20-24, male & North East

14: 20-24, male & North West (incl. the old Merseyside region)

15: 20-24, male & Yorkshire & Humberside

etc....

SURVEY DESIGN

Notes: 1. Variable 'strata2' should be declared as the strata for analysis of the core dataset in Stata using Stata's survey commands when there are too few PSUs in a stratum/strata to use 'strata'. See the Natsal-2 Technical Report for further details.
2. The variable 'strata2' has also been derived for Natsal-1
3. Variable 'strata2' is derived from 3 age-groups ('agrp'), 2 genders ('rsex'), and the 11 regions of 'gor' considering London as 2 regions: inner and outer London, and the 11 regions of 'GOR' but considering London as 2 regions: inner and outer London, so 12 regions in total, and 72 strata in total.

(R) (ND) **strata2** 72 strata

- 1: 16-24, male & North East
- 2: 16-24, male & North West (incl. the old Merseyside region)
- 3: 16-24, male & Yorkshire & Humberside
- 4: 16-24, male & East Midlands
- 5: 16-24, male & West Midlands
- 6: 16-24, male & South West
- 7: 16-24, male & Eastern
- 8: 16-24, male & Inner London
- 9: 16-24, male & Outer London
- 10: 16-24, male & South East
- 11: 16-24, male & Wales
- 12: 16-24, male & Scotland
- etc...

Notes: 1. Variable 'strata3' should be declared as the strata for analysis of the core dataset in Stata using Stata's survey commands when there are too few PSUs in a stratum/strata to use 'strata2'. See the Natsal-2 Technical Report for further details.
2. The variable 'strata3' has also been derived for Natsal-1
3. Variable 'strata3' is derived from 3 age-groups ('agrp'), 2 genders ('rsex'), and the 2 regions of 'london' (Greater London vs. the rest of Britain), so 12 strata in total.

(R) (ND) **strata3** 12 strata

- 1: 16-24, male & rest of Britain
- 2: 16-24, male & London
- 3: 25-34, male & rest of Britain
- 4: 25-34, male & London
- 5: 35-44, male & rest of Britain
- 6: 35-44, male & London
- 7: 16-24, female & rest of Britain
- 8: 16-24, female & London
- 9: 25-34, female & rest of Britain
- 10: 25-34, female & London
- 11: 35-44, female & rest of Britain
- 12: 35-44, female & London

DEMOGRAPHIC CHARACTERISTICS

DEMOGRAPHIC CHARACTERISTICS

Note: The age range for Natsal-3 was 16-74 years. New age categories have been added.

(q1a) **dage** respondent's age at interview, years

(ND) **dagegr** respondent's age at interview, years, grouped

- 1: 16-17 years
- 2: 18-19 years
- 3: 20-24 years
- 4: 25-34 years
- 5: 35-44 years
- 6: 45-54 years
- 7: 55-64 years
- 8: 65-74 years

Note: Variable 'agrp' is derived from dage

(D) **agrp** respondent's age at interview, years, grouped

- 1: 16-24 years
- 2: 25-34 years
- 3: 35-44 years
- 4: 45-54 years
- 5: 55-64 years
- 6: 65-74 years

Note: Variable 'agrp2' is derived from dage

(D) **agrp2** respondent's age at interview, years, grouped

- 1: 16-19 years
- 2: 20-24 years
- 3: 25-29 years
- 4: 30-34 years
- 5: 35-39 years
- 6: 40-44 years
- 7: 45-49 years
- 8: 50-54 years
- 9: 55-59 years
- 10: 60-64 years
- 11: 65-69 years
- 12: 70-74 years

Note: rdobm, rdoby and rdob imputed from dage if not answered

Rdobm respondent's month of birth

- 1 'January'
- 2 'February'
- 3 'March'
- 4 'April'
- 5 'May'
- 6 'June'
- 7 'July'
- 8 'August'
- 9 'September'
- 10 'October'
- 11 'November'
- 12 'December'

Rdoby respondent's year of birth

Rdob respondent's date of birth – day set to 01

SURVEY DESIGN

(ND) **birthcohort** 5-year birth cohort

- 1: 1935-1939 *if* rdoby>=1935 & rdoby<=1939
- 2: 1940-1944 *if* rdoby>=1940 & rdoby<=1944
- 3: 1945-1949 *if* rdoby>=1945 & rdoby<=1949
- 4: 1950-1954 *if* rdoby>=1950 & rdoby<=1954
- 5: 1955-1959 *if* rdoby>=1955 & rdoby<=1959
- 6: 1960-1964 *if* rdoby>=1960 & rdoby<=1964
- 7: 1965-1969 *if* rdoby>=1965 & rdoby<=1969
- 8: 1970-1974 *if* rdoby>=1970 & rdoby<=1974
- 9: 1975-1979 *if* rdoby>=1975 & rdoby<=1979
- 10: 1980-1984 *if* rdoby>=1980 & rdoby<=1984
- 11: 1985-1989 *if* rdoby>=1985 & rdoby<=1989
- 12: 1990-1996 *if* rdoby>=1990 & rdoby<=1996

(q1b) **rsex** respondent's sex

- 1: male
- 2: female

GENERAL HEALTH

GENERAL HEALTH

(q2a*) **health** respondent's opinion of own health

- 1: very good
- 2: good
- 3: fair
- 4: bad
- 5: very bad
- 9: not answered

(q2b*) **disabil** respondent has a longstanding illness, disability or infirmity

- 1: yes
- 2: no
- 9: not answered

Note: Coding for limit changed in Natsal-3 so that those who answered 'no' at disabil are coded as 'no' at limit, not 'not applicable'.

(N) (U) **limit** respondent's activities are limited by their disability

(U – code removed) -1: not applicable if disabil=2

- 1: yes
- 2: no
- 9: not answered

(ND) **disabil2** respondent has a longstanding illness, disability or infirmity

- 1: none if disabil=2
- 2: non-limiting if disabil=1 & limit=2
- 3: limiting if limit=1
- 9: not answered if limit=9

(R) (q2d) **illoracc** respondent has had an illness for 3 months+ in last 5 years

- 1: yes
- 2: no
- 9: not answered

HOSPITAL ATTENDANCE

(R) (N) **outpatyr** hospital out-patient visit in last year (excl. ante-/post-natal)

- 1: yes
- 2: no
- 9: not answered

(R) (N) **daypat** day-ward patient in last year (excl. pregnancy visit)

- 1: yes
- 2: no
- 9: not answered

(R) (N) **inpat** in-patient visit in last year (excl. pregnancy visit)

- 1: yes
- 2: no
- 9: not answered

Note: Illoracc, outpatyr, daypat and inpat which were in Natsal-2 have been removed. These have been replaced by more specific health questions which aim to gain a better understanding of the illnesses potentially important to an individual's sexual life and function. The medication question is important to capture medications which can impact sexual function and may for some illnesses be taken as a marker for severity.

(N) **medcona** respondent has had serious physical health problem listed on Card A

- 1: yes

GENERAL HEALTH

2: no
9: not answered *else if missing*

Note: For multi-code questions this codebook gives the binary variables to be used for analysis. medwcha1..medwcha9 were created directly from the question medwcha. All are coded similarly to medwcha1.

(N) **medwcha1** medical condition reported by respondent: arthritis
1: yes
0: no
9: not answered

And similarly...

(N) **medwcha2** medical condition reported by respondent: heart attack
(N) **medwcha3** medical condition reported by respondent: coronary heart disease / angina / other form of heart disease
(N) **medwcha4** medical condition reported by respondent: hypertension
(N) **medwcha5** medical condition reported by respondent: stroke
(N) **medwcha6** medical condition reported by respondent: diabetes
(N) **medwcha7** medical condition reported by respondent: chronic lung disease
(N) **medwcha8** medical condition reported by respondent: parkinson's disease
(N) **medwcha9** medical condition reported by respondent: epilepsy

(N) **drghheart** currently taking medication for heart disease
1: yes
2: no
9: not answered *else if missing*

(N) **drghhyp** currently taking medication for hypertension
1: yes
2: no
9: not answered *else if missing*

(N) **drbins** currently injecting insulin
1: yes
2: no
9: not answered *else if missing*

(N) **drgrdi** currently taking other medicines, tablets or pills for diabetes
1: yes
2: no
9: not answered *else if missing*

(N) **drgpark** currently taking medicine for parkinsons
1: yes
2: no
9: not answered *else if missing*

(N) **drgepil** currently taking medicine for epilepsy
1: yes
2: no
9: not answered *else if missing*

(N) **medconb** respondent reports male genital health condition listed on Card B
1: yes
2: no
9: not answered *else if missing*
-1: not applicable *if rsex=2*

GENERAL HEALTH

Note: For multi-code questions this codebook gives the binary variables to be used for analysis. medwchb1..medwchb8 were created directly from the question medwchb. All are coded similarly to medwchb1.

(N) **medwchb1** Male genital health: medical condition reported by respondent: prostate enlargement
1: yes
0: no
9: not answered
-1: not applicable if rsex=2

And similarly...

(N) **medwchb2** Male genital health: medical condition reported by respondent: prostate cancer
(N) **medwchb3** Male genital health: medical condition reported by respondent: prostate surgery
(N) **medwchb4** Male genital health: medical condition reported by respondent: bladder surgery
(N) **medwchb5** Male genital health: medical condition reported by respondent: genital surgery
(N) **medwchb6** Male genital health: medical condition reported by respondent: abdominal surgery
(N) **medwchb7** Male genital health: medical condition reported by respondent: broken hip or pelvis bone
(N) **medwchb8** Male genital health: medical condition reported by respondent: hip replacement

(N) **drgpros** currently taking medicine from doctor for prostate disease
1: yes
2: no
9: not answered *else if missing*
-1: not applicable if rsex=2

(N) **medconc** respondent reports female genital health condition listed in Card C
1: yes
2: no
9: not answered *else if missing*
-1: not applicable if rsex=1

Note: For multi-code questions this codebook gives the binary variables to be used for analysis. medwchc1..medwchc7 were created directly from the question medwchc. All are coded similarly to medwchc1.

(N) **medwchc1** Femal genital health: medical condition reported by respondent: Hysterectomy
1: yes
0: no
9: not answered
-1: not applicable if rsex=1

And similarly...

(N) **medwchc2** Femal genital health: medical condition reported by respondent: Bladder surgery
(N) **medwchc3** Femal genital health: medical condition reported by respondent: Genital or gynaecological surgery
(N) **medwchc4** Femal genital health: medical condition reported by respondent: Caesarean section / C section
(N) **medwchc5** Femal genital health: medical condition reported by respondent: Abdominal surgery
(N) **medwchc6** Femal genital health: medical condition reported by respondent: Broken hip or pelvis bone
(N) **medwchc7** Femal genital health: medical condition reported by respondent: Hip replacement

(N) **whnhyst** age when had hysterectomy, years
-1: not applicable if $\neq$ rsex=1
996: not had a hysterectomy if medconc=no or medwchc1 $\neq$ yes
999: not answered *else if missing*

GENERAL HEALTH

(N) **medcond** respondent reported treatment for a medical condition listed in Card D in the last year
1: yes
2: no
9: not answered *else if missing*

Note: For multi-code questions this codebook gives the binary variables to be used for analysis. medwchd1..medwchd7 were created directly from the question medwchd. All are coded similarly to medwchd1.

(N) **medwchd1** Received treatment for medical condition in last year: Back ache lasting for 3 months or longer
1: yes
0: no
9: not answered

And similarly...

(N) **medwchd2** Received treatment for medical condition in last year: Any other muscle or bone disease lasting for 3 months or longer

(N) **medwchd3** Received treatment for medical condition in last year: Depression

(N) **medwchd4** Received treatment for medical condition in last year: Any other mental health condition

(N) **medwchd5** Received treatment for medical condition in last year: Any other neurological condition, apart from Parkinson's disease and epilepsy

(N) **medwchd6** Received treatment for medical condition in last year: Cancer

(N) **medwchd7** Received treatment for medical condition in last year: Any thyroid condition

(N) **medwchd8** Received treatment for medical condition in last year: Any (ovarian/testicular) or pituitary condition

Note: Response options 9 and 10 were only available to women

(N) **medwchd9** Received treatment for medical condition in last year: Polycystic ovaries (Women only)
1: yes
0: no
9: not answered
-1: not applicable *if rsex=1*

(N) **medwchd10** Received treatment for medical condition in last year: IVF or other fertility treatment (Women only)
1: yes
0: no
9: not answered
-1: not applicable *if rsex=1*

Note: the variable drgdep relates to medications taken for depression. Data users may also wish to note the variable depscr which appears later on in the dataset and codebook (page 161) which is based on the PHQ-2 depression screening questionnaire.

(N) **drgdep** currently taking medicine prescribed by a doctor for depression
1: yes
2: no
9: not answered *else if missing*

(N) **drgmh** currently taking medication for mental health condition other than depression
1: yes
2: no
9: not answered *else if missing*

GENERAL HEALTH

Note: Hefunc & heatt are designed to capture limitations to the respondent's life due to health problems. This reflects the fact that the Natsal-3 survey has a higher age limit.

(N) **hefunc** how much difficulty walking up a flight of stairs because of a health problem
-1: not applicable if (health=1 or 2) & disabil=2 medcona=2 & ((rsex=1 & medconb=2) | (rsex=2 & medconc=2) & medcond=no

1: no difficulty

2: some difficulty

3: much difficulty

4: unable to do this

9: not answered *else if missing*

Note: For multi-code questions this codebook gives the binary variables to be used for analysis. heatt1..heatt12 were created directly from the question heatt. All are coded similarly to heatt1.

(N) **heatt1** symptoms that cause difficulty: Chest pain

1: yes

0: no

9: not answered

-1: not applicable *if* hefunc=-1 | hefunc=1

And similarly...

(N) **heatt2** symptoms that cause difficulty: Shortness of breath

(N) **heatt3** symptoms that cause difficulty: Pain

(N) **heatt4** symptoms that cause difficulty: Weakness or fatigue

(N) **heatt5** symptoms that cause difficulty: I am a wheelchair user

(N) **heatt6** symptoms that cause difficulty: Amputation

(N) **heatt7** symptoms that cause difficulty: Visual problems

(N) **heatt8** symptoms that cause difficulty: Tremor or balance problems

(N) **heatt9** symptoms that cause difficulty: Unsteady on the feet

(N) **heatt10** symptoms that cause difficulty: Fear of falling

(N) **heatt11** symptoms that cause difficulty: Incontinence or fear of incontinence

(N) **heatt12** symptoms that cause difficulty: Some other problem or symptom

ANTE-NATAL CARE

Note: The antenatal question (antenat) has been moved into the new fertility section in the self completion.

(R) (N) **antenno** no. of pregnancies antenatal visits were for in last 5 years

-1: not applicable *if* rsex=1

0: *if* antenat=2

99: not answered

(R) (N) **antenyr1** year first attended ante-natal clinic for 1st pregnancy

-1: not applicable *if* rsex=1 | antenat=2

9999: not answered

Similarly for up to 7 pregnancies with additional code of -1 if (antenno<antenyrn)

(R) (N) **antenyr2** year first attended ante-natal clinic for 2nd pregnancy

(R) (N) **antenyr3** year first attended ante-natal clinic for 3rd pregnancy

(R) (N) **antenyr4** year first attended ante-natal clinic for 4th pregnancy

(R) (N) **antenyr5** year first attended ante-natal clinic for 5th pregnancy

(R) (N) **antenyr6** year first attended ante-natal clinic for 6th pregnancy

(R) (N) **antenyr7** year first attended ante-natal clinic for 7th pregnancy

GENERAL HEALTH

Note: Anbld and anwhn which were included in Natsal-2 were removed from the questionnaire in Natsal-3 since these questions had not been used in analysis. These data are no longer needed to triangulate with the HIV data from pregnant women.

(R) (N) **anbld** ever had blood test as part of ante-natal care

-1: not applicable if rsex=1 | antenat=1

1: yes

2: no

9: not answered

Note: The variable 'anwhn' does not apply if female respondents reported ante-natal visit(s) in the last 5 years

(R) (N) **anwhn** blood test as part of ante-natal care since 1 Jan 1988

-1: not applicable if rsex=1 | antenat=1 | anbld=2

1: yes, more than 5 years ago but since 1 Jan 1988

2: no, before 1988 only

9: not answered

ALCOHOL CONSUMPTION

Note: In Natsal-3 there was an instruction to interviewers that respondents not drinking alcohol at the moment due to pregnancy should be coded as 'no' in drink

(N) **drink** ever drink alcohol nowadays

1: yes

2: no

9: not answered

Note: drinkany was removed from the questionnaire in Natsal-3 since very few people responded yes to this question in Natsal-2 and of these very few reported drinking with a frequency of 1-2 times per month or more.

(R) (N) **drinkany** check never drink alcohol nowadays

-1: not applicable if drink=1

1: very occasionally

2: never

9: not answered

Note: eligibility for this Q is slightly different in Natsal-3 since the Q drinkany is no longer asked

(q4a*) **drinkoft** average frequency of alcohol consumption in last 12 months

(U) -1: not applicable if drink=2 -1: not applicable if drinkany=2

1: 5+ days a week

2: 3/4 days a week

3: once/twice a week

4: once/twice a month

5: once/twice in the last 12 months

6: not at all in the last 12 months

9: not answered

GENERAL HEALTH

Note: In Natsal-3 the instruction to exclude parties was taken out of ManyAlc since this may be the only time people drink. ManyAlc is now captured in terms of units instead of drinks and the variable was renamed as ManyAlc2 in Natsal-3.

Note: variable 'manyalc' does not use code 5

(R) (q4b*) manyalc average alcohol intake when do drink (excl. parties/special occasions)

0: drink less frequently than once/twice a month *if* drinkany=2 | drinkoft=5 | 6

1: 1 or 2

2: 3 or 4

3: 5 or 6

4: >6

5: *code not used*

6: less than 1 unit

7: only drink on special occasions/at parties

8: varies too much to say/other answers

9: not answered

(N) (q4b*) manyalc2 average alcohol intake when do drink (excl. special occasions)

0: drink less frequently than once/twice a month *if* drink=2 | drinkoft=5 | 6

1: 1 or 2

2: 3 or 4

3: 5 or 6

4: >6

5: I only drink on special occasions

6: other

9: not answered *else if missing*

Note: Variable 'unitspw' is derived from drinkoft and manyalc:

(U – see below) (ND) unitspw average number of units of alcohol per week

compute drinkoft x manyalc

if drinkoft = 1, 6 days per week on average

if drinkoft = 2, 3.5 days per week on average

if drinkoft = 3, 1.5 days per week on average

if drinkoft = 4, 0.5 days per week on average

if drinkoft = 5, 0 days per week on average

if drinkoft = 6, 0 days per week on average

if manyalc = 0, 0 units on average

if manyalc = 1, 1.5 units on average

if manyalc = 2, 3.5 units on average

if manyalc = 3, 5.5 units on average

if manyalc = 4, 10 units on average

if manyalc = 6, 0.5 units on average

if manyalc = 7, 0 units on average

if manyalc = 8, assume missing

99: *if* drinkoft=9 | manyalc =9

GENERAL HEALTH

Note: Variable 'unitspw' is derived from manyAlc2 instead of manyalc in Natsal-3

unitspw average number of units of alcohol per week
compute drinkoft x manyalc2 compute drinkoft x manyalc
if drinkoft = 1, 6 days per week on average
if drinkoft = 2, 3.5 days per week on average
if drinkoft = 3, 1.5 days per week on average
if drinkoft = 4, 0.5 days per week on average
if drinkoft = 5, 0 days per week on average
if drinkoft = 6, 0 days per week on average
if manyalc2 = 0, 0 units on average
if manyalc2 = 1, 1.5 units on average
if manyalc2 = 2, 3.5 units on average
if manyalc2 = 3, 5.5 units on average
if manyalc2 = 4, 10 units on average
if manyalc2 = 5, 0 units on average
if manyalc = 6, assume missing
99: if drinkoft=9 | manyalc =9

Note: Variable 'alcohol' is derived from unitspw:

(R) (D) **alcohol** average alcoholic consumption per week, grouped
0: none *if unitspw=0*
1: low *if (unitspw>0 & unitspw <=20 & rsex=1) | (unitspw >0 & unitspw<= 15 & rsex=2)*
2: moderate *if (unitspw>20 & unitspw <50 & rsex=1) | (unitspw>15 & unitspw<35 & rsex=2)*
3: high *if (unitspw>=50 & unitspw<99 & rsex=1) | (unitspw>=35 & unitspw<99 & rsex=2)*
9: not answered *if unitspw=99*

Note: Variable 'alcohol2' is derived from unitspw:

(D) **alcohol2** average alcoholic consumption per week, grouped
0: none *if unitspw=0*
1: not more than recommended *if (unitspw>0 & unitspw <=21 & rsex=1) | (unitspw>0 & unitspw<=14 & rsex=2)*
2: more than recommended *if (unitspw>21 & unitspw<99 & rsex=1) | (unitspw>14 & unitspw<99 & rsex=2)*
9: not answered *if unitspw=99*

Notes: 1. Variable 'alcohol3' is derived from unitspw
2. Unlike the derived variables 'alcohol' & 'alcohol2', 'alcohol3' is not sex-specific

(D) **alcohol3** average alcoholic consumption per week, grouped
0: 0 units per week *if unitspw=0*
1: 1-7 units per week *if unitspw>0 & unitspw <=7*
2: 8-14 units per week *if unitspw>= 8 & unitspw <=14*
3: 15-21 units per week *if unitspw>=15 & unitspw <=21*
4: 22+ units per week *if unitspw>=22 & unitspw <99*
9: not answered *if unitspw=99*

(ND) **alcoholn** average alcoholic consumption per week, grouped
0: none *if unitspw=0*
1: low *if (unitspw>0 & unitspw<=21 & rsex==1) | (unitspw>0 & unitspw<=14 & rsex==2)*
2: moderate *if (unitspw>21 & unitspw<=50 & rsex==1) | (unitspw>14 & unitspw<=35 & rsex==2)*
3: high *if (unitspw>50 & unitspw<99 & rsex==1) | (unitspw>35 & unitspw<99 & rsex==2)*
9: not answered *if unitspw=99*

GENERAL HEALTH

Note: Drinknum comes from the FAST measure of hazardous drinking. This question forms the first stage of the FAST measure, after which additional questions are asked of some respondents. These additional questions were not included in Natsal-3 due to restrictions on space.

(N) **drinknum** how often have >6/8 units on one occasion

-1: not applicable if drinkoft=5 | drinkoft=6 | drinkoft=9

1: never if drinknum=1 | drink=2

2: less than monthly

3: monthly

4: weekly

5: daily or almost daily

9: not answered

SMOKING

(q5a*) **smokenow** smoke cigarettes nowadays

1: yes

2: no

9: not answered

(q5b*) **nosmoke** number of cigarettes usually smoked a day

0 if smokenow=2

99: not answered

(q5c*) **exsmoke** ever smoked cigarettes regularly (at least 1 a day)

-1: not applicable if smokenow=1

1: yes

2: no

9: not answered

(D) **smoking** current smoking status

1: non-smoker if (smokenow=2 & exsmoke=2)

2: ex-smoker if exsmoke=1

3: light smoker if (smokenow=1) & (nosmoke>=0 & nosmoke<15)

4: heavy smoker if (smokenow=1) & (nosmoke>=15 & nosmoke <99)

9: not answered if smokenow=9 | nosmoke=99

(ND) **smoker** smoking status

0: never if exsmoke=2

1: ex-smoker if exsmoke=1

2: current smoker if smokenow=1

9: not answered if smokenow=9

WEIGHT, HEIGHT & BODY MASS INDEX

Note: in Natsal-3 interviewers were asked to record pre-pregnancy weights if respondents were pregnant

(q6a3) **kgweight** weight in kilograms

999: not answered

Note: Surewgt has been removed since this has never been used in analysis

(R) (q6b) **surewgt** certainty of weight

-1: not applicable if kgweight=999

1: fairly sure

2: estimate

9: not answered

GENERAL HEALTH

Pregnow has been removed since pregnancy status is captured in the fertility section in the CASI

(R) (N) **pregnow** pregnant at interview
-1: not applicable if rsex =1 | (rsex=2 & kgweight=999)
1: yes
2: no
9: not answered

(q6c2) **cmheight** height in centimetres
999: not answered

Note: Variable 'bodymass' is derived from kgweight and cmheight:

(D) **bodymass** body mass index
*compute kgweight / ((cmheight /100) * (cmheight /100)) if (kgweight<999) & (cmheight<999)*
99: not answered if kgweight=999 | cmheight=999

Notes: 1. Variable 'bmihgroup' is derived from 'bodymass'
2: A Body Mass Index of less than 12 or more than 58 is not considered possible

(R) (ND) **bmigroup** body mass index, grouped
1: bmi <20 kg/m squared if bodymass>=12 & bodymass<20
2: bmi 20-25 kg/m squared if bodymass>=20 & bodymass<25
3: bmi 25 -30 kg/m squared if bodymass>=25 & bodymass<30
4: bmi 30 -40 kg/m squared if bodymass>=30 & bodymass<40
5: bmi 40 kg/m squared + if bodymass>=40 & bodymass<99
9: not answered if bodymass=99

Note: a new BMI grouping variable was created for Natsal-3 in line with the current WHO definitions.

(ND) **bmiwho** body mass index, grouped
1: underweight (bmi <18.5 kg/m squared) if bodymass>=12 & bodymass<18.5
2: normal (bmi 18.5-25 kg/m squared) if bodymass>=18.5 & bodymass<=25
3: overweight (bmi >25-30 kg/m squared) if bodymass>25 & bodymass<=30
4: obese I (bmi >30-35 kg/m squared) if bodymass>30 & bodymass<=35
5: obese II (bmi >35-40 kg/m squared) if bodymass>35 & bodymass<=40
6: obese III (bmi >40 kg/m squared) if bodymass>40 & bodymass<99
9: not answered if bodymass=99

FAMILY HISTORY

FAMILY HISTORY

Note: This section has been redrafted to better capture the diversity in family circumstances. The section no longer specifies siblings when growing up.

Note: For multi-code questions this codebook gives the binary variables to be used for analysis. sibany1..sibany9 were created directly from the question sibany. All are coded similarly to sibany1.

(N) **sibany1** any brothers or sisters: full brother
1: yes
0: no
9: not answered

And similarly...

(N) **sibany2** any brothers or sisters: full sister
(N) **sibany3** any brothers or sisters: adopted brother
(N) **sibany4** any brothers or sisters: adopted sister
(N) **sibany5** any brothers or sisters: half brother
(N) **sibany6** any brothers or sisters: half sister
(N) **sibany7** any brothers or sisters: step brother
(N) **sibany8** any brothers or sisters: step sister
(N) **sibany9** any brothers or sisters: none

(N) **sibbro** no. natural brothers
0: if sibany1=0
99: not answered or if sibany1=9

(N) **sibsis** no. natural sisters
0: if sibany2=0
99: not answered or if sibany2=9

(N) **sibadbr** no. adopted brothers
0: if sibany3=0
99: not answered or if sibany3=9

(N) **sibadsis** no. adopted sisters
0: if sibany4=0
99: not answered or if sibany4=9

(N) **sibhabr** no. half brothers
0: if sibany5=0
99: not answered or if sibany5=9

(N) **sibhasis** no. half sisters
0: if sibany6=0
99: not answered or if sibany6=9

(N) **sibstbr** no. step brothers
0: if sibany7=0
99: not answered or if sibany7=9

(N) **sibstsis** no. step sisters
0: if sibany8=0
99: not answered or if sibany8=9

FAMILY HISTORY

(R) (q7a) **anysibs** any brothers or sisters

1: yes, brother(s) only

2: yes, sister(s) only

3: yes, both

4: no, none

9: not answered

(R) (q7b) **nosibs** number of siblings (incl. half and adopted siblings)

0 if anysibs=4

99: not answered

(R) (q7c) **sibspos** position in family

-1: not applicable if anysibs=4 | nosibs=0

1: oldest

2: youngest

3: in between

9: not answered

(R) (q7e*) **stepsibs** have/had step-brothers or step-sisters

1: yes

2: no

9: not answered

(N) **siblive** live with siblings (natural, half, adopted or step) at any time when growing up

-1: not applicable if sibany9=1 1: yes

2: no

9: not answered or if sibany9=9

(N) **sibpos2** position in family when growing up

-1: not applicable if sibany9=1 | siblive=2

1: oldest

2: youngest

3: in between

9: not answered or if sibany9=9

Note: reference age of bothmapa and ynotboth was changed to 14 to correspond to the TalkMaPa variable. The variable names have been updated accordingly.

(R) (q8a) **bothmapa** lived more or less continuously to age 16 with both natural parents

1: yes

2: no

9: not answered

(N) **bothmapa2** lived more or less continuously to age 14 with both natural parents

1: yes

2: no

9: not answered

FAMILY HISTORY

(R) (q8b U) **ynotboth** why didn't live with both natural parents continuously to age 16

-1: not applicable *if bothmapa=1*

1: divorce/separation

2: death

3: adopted

4: parents never lived together

5: parents were abroad

6: in care

7: other reason

8: lived with grand-parent(s)/other relative

9: left home before age 16

99: not answered

(N) **ynotboth2** why didn't live with both natural parents continuously to age 14

-1: not applicable *if bothmapa2=1*

1: divorce/separation

2: death

3: adopted

4: parents never lived together

5: other

9: not answered *or if bothmapa2=99*

Note: Maorpa removed and a more detailed question (wholive) has been added to capture who the respondent lived with as a child

(R) (q8d U) **maorpa** which parent lived with to age 16

-1: not applicable *if bothmapa=1 | ynotboth=3*

1: mother

2: father

3: other relative

4: other (including in care or fostered)

9: not answered

(N) **wholive** who living with when aged 14

-1: not applicable *if bothmapa==1*

1: Adoptive mother and father

2: Mother and stepfather

3: Father and stepmother

4: Mother only

5: Father only

6: In Local Authority care / foster home

7: Natural mother and father

8: Other (specified)

9: Grandparents/other relative

10: Left home before age of 14

99: not answered

Note: In Natsal-3 one variable, parents, has been derived from wholive not maorpa. Since this affects the definition of the variable, this has been renamed parents3.

Note: Variable 'parents' treats those who were adopted as having lived with both parents till 16

(R) (DN) **parents** family structure to age 16, grouped

1: both natural parents/adopted *if bothmapa=1 | (bothmapa=2 & ynotboth=3)*

2: one natural parent *if maorpa=1 | maorpa=2*

3: neither natural parent *if maorpa=3 | maorpa=4*

9: not answered *if bothmapa=9 | maorpa=9*

FAMILY HISTORY

Note: Variable 'parents2' treats those who were adopted as living with neither natural parent till 16

(R) (DN) **parents2** family structure to age 16, grouped

- 1: both natural parents *if* bothmapa=1
- 2: one natural parent *if* maorpa=1 | maorpa=2
- 3: neither natural parent (incl. adopted) *if* (maorpa=3 | maorpa=4) | ynotboth=3
- 9: not answered *if* bothmapa=9 | maorpa=9

Note: variable 'parents3' is slightly different from parents and uses wholive which says who the respondent was living with at the age of 14. Therefore there is no need for two alternative variables for adopted children.

(ND) **parents3** family structure at age 14, grouped

- 1: both natural parents/adopted *if* bothmapa=1 | (bothmapa=2 & wholive=1) | (bothmapa=2 & wholive=7)
- 2: one natural parent *if* wholive=2 | wholive=3 | wholive=4 | wholive=5
- 3: neither natural parent *if* wholive=6 | wholive=8 | wholive=9 | wholive=10
- 9: not answered *if* bothmapa==9 | wholive==99

(ND) **parents4** family structure at age 14, grouped

- 1: natural/adoptive parents *if* bothmapa2=1 | wholive=1 | wholive=7
- 2: one natural/one step *if* wholive=2 | wholive=3
- 3: single parent *if* wholive=4 | wholive=5
- 4: in care *if* wholive=6
- 5: other *if* wholive>=8 & wholive<=10

CURRENT FAMILY

Note: this information is now collected in the history of pregnancies section (women) and the family formation section (men and some questions men and women) in the CASI.

LEARNING ABOUT SEX

LEARNING ABOUT SEX

Note: eligibility for talkmapa different in Natsal-3

(U) (q10b) **talkmapa** ease with which discussed sex with parents around age 14

(U) -1: not applicable *if bothmapa=2 & (wholive#1 & wholive#2 & wholive#3 & wholive#4 & wholive#5 & wholive#7)*

-1: not applicable *if bothmapa=2 & (maorpa#1 & maorpa#2 & maorpa#9)*

1: easy with one/both

2: difficult

3: didn't discuss with either

4: varied/depended on topic

9: not answered

(ND) **notalk** discussed sex with parents around age 14

-1: not applicable *if talkmapa=-1*

0: discussed with 1+ parent(s) although in some cases difficult *if talkmapa=1 | talkmapa=2 | talkmapa=4*

1: did not discuss with either *if talkmapa=3*

9: not answered *if talkmapa=9*

Note: response options for learns01 have been updated to be relevant in current social context. In particular the increase in the use of the internet. The names of the binary variables have been changed to reflect this change in the list of answer options (Natsal-3 binary variables shown below the Natsal-1 and Natsal-2 ones).

*Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code questions. The following binary variables were created directly from **learns01-learns12** in Natsal-2, and **q17a** in Natsal-1.*

Natsal-2 binary variables derived from Learns01-Learns12 and xlernsex. Note all variables coded in a similar way to sexedmum:

(R) (q17ap D) **sexedmum** learned about sex from mother

-1: not applicable *if learns01=15*

1: yes *if (learns01=1 | learns02=1 | learns03=1 | learns04=1 | learns05=1 | learns06=1 | learns07=1 | learns08=1 | learns09=1 | learns10=1 | learns11=1 | learns12=1) | ((learns01=14 | learns02=14 | learns03=14 | learns04=14 | learns05=14 | learns06=14 | learns07=14 | learns08=14 | learns09=14 | learns10=14 | learns11=14 | learns12=14) & (lernsex1=1 | lernsex2=1 | lernsex3=1))*

0: no *else if (learns01>=1 & learns01<=14)*

9: not answered *if (learns01=98 | learns01=99)*

(R) (q17az D) **sexeddad** learned about sex from father

(R) (q17ax D) **sexedbro** learned about sex from brother(s)

(R) (q17al D) **sexedsis** learned about sex from sister(s)

(R) (q17an D) **sexedrel** learned about sex from other relative(s)

(R) (q17ad D) **sexedsch** learned about sex from lessons at school

(R) (q17aj D) **sexedfrn** learned about sex from friend(s) of a similar age

(R) (q17as D) **sexedfsp** learned about sex from first boy/girlfriend/sexual partner

(R) (q17aa D) **sexeddoc** learned about sex from doctor/nurse/clinic

(R) (q17ak & q17ae D) **sexedtv** learned about sex from television/radio

(R) (ND) **sexedvid** learned about sex from video(s)

(R) (q17ag D) **sexedbks** learned about sex from books

(R) (q17av D) **sexedmag** learned about sex from magazines/newspapers

(R) (q17aq D) **sexedoth** learned about sex from another source

LEARNING ABOUT SEX

Natsal-3 binary variables:

(N) **sexedudoc** learned about sex from doctor/nurse/clinic

0: no

1: yes

9: not answered *if* learnsex = can't remember / not answered

And similarly...

(N) **sexeduintad** learned about sex from internet-sexual advice websites

(N) **sexedusch** learned about sex from lessons at school

(N) **sexeduinto** learned about sex from internet-other

(N) **sexedubmn** learned about sex from books/magazines/newspapers

(N) **sexedufnr** learned about sex from friend(s) of about own age

(N) **sexedutvid** learned about sex from tv/radio/DVDs/videos

(N) **sexedumum** learned about sex from mother (incl. step/adoptive)

(N) **sexedudad** learned about sex from father (incl. step/adoptive)

(N) **sexedufsp** learned about sex from 1st boy/girlfriend/sexual partner

(N) **sexeduintpn** learned about sex from internet-pornographic websites

(N) **sexedupor** learned about sex from pornographic magazines/films

(N) **sexedubsis** learned about sex from brother/sister (incl. half/step/adopted)

(N) **sexeduoth** learned about sex from another source

(N) **sexeduani** learned about sex from animals/growing up on farm

(N) **sexedurel** learned about sex from other relatives

(N) **sexeduofri** learned about sex from older friends

(N) **sexeduofrn** learned about sex from friends (age not specified)

(N) **sexeduexp** learned about sex through own experience

(N) **sexedufile** learned about sex from cinemas/film (other than TV/video)

(N) **sexeduedc** learned about sex from lessons/courses (i.e. formal teaching)

(U – see below) (q17b U) **lernmost** main source of sex education (if more than one source reported)

-1: not applicable *if* learns02=-1

1: mother

2: father

3: brother(s)

4: sister(s)

5: other relative(s)

6: lessons at school

7: friends of about my own age

8: first boy/girl friend/sexual partner

9: doctor/nurse/clinic

10: television/radio

11: videos

12: books

13: magazines/newspapers

14: other

15: can't choose just one source

99: not answered

LEARNING ABOUT SEX

Note: response options for lernmost are as for learnsex but only able to choose one source

lernmost main source of sex education 1 – NEW CODES but question identical

-1: not applicable *if* only one source reported at learnsex

- 1: doctor, nurse or clinic
- 2: Internet – sexual advice websites
- 3: Lessons at school
- 4: Internet – other
- 5: Books / magazines / newspapers
- 6: Friends of about my own age
- 7: Television / radio / DVDs / videos
- 8: Mother (including step or adoptive)
- 9: Father (including step or adoptive)
- 10: First (girlfriend/boyfriend) or sexual partner
- 11: Internet – pornographic websites
- 12: Pornographic magazines / films
- 13: Brother(s) / sister(s) (including half, step, or adoptive)
- 14: Other
- 15: animals/growing up on farm
- 16: Other relative
- 17: Older friends
- 18: Friends (age not specified)
- 19: Through (own) experience
- 20: cinemas/film (other than TV/video)
- 21: Lessons/course (i.e. formal teaching)
- 99: Not answered

Note: in Natsal-3 respondent now has to specify option at lernmost – no option to say >1 so mlernm was not asked.

(R) (N) mlernm01 main sources of sex education 1 (if more than one source reported and can't choose just one)

-1: not applicable *if* (lernmost>=1 & lernmost<=14) | learns02=-1 | (mlernmn=99 & mlernmn-1≠99)

- 1: mother
- 2: father
- 3: brother(s)
- 4: sister(s)
- 5: other relative(s)
- 6: lessons at school
- 7: friends of about my own age
- 8: first boy/girl friend/sexual partner
- 9: doctor/nurse/clinic
- 10: television/radio
- 11: videos
- 12: books
- 13: magazines/newspapers
- 14: other
- 99: not answered

Similarly for up to 2 other sources:

(R) (N) mlernm02 main sources of sex education 2 (if more than one source reported and can't choose just one)

(R) (N) mlernm03 main sources of sex education 3 (if more than one source reported and can't choose just one)

(R) (N) mlernm04 main sources of sex education 4 (if more than one source reported and can't choose just one)

(R) (N) mlernm05 main sources of sex education 5 (if more than one source reported and can't choose just one)

(R) (N) mlernm06 main sources of sex education 6 (if more than one source reported and can't choose just one)

LEARNING ABOUT SEX

(U – see below) (ND) **msedmum** main source of sex education was mother
1: yes if (lerns01=1 & lerns02=-1) | lernmost=1 | (lerns01=1 | lerns02=1 | lerns03=1 | lerns04=1 | lerns05=1 | lerns06=1 | lerns07=1 | lerns08=1 | lerns09=1 | lerns10=1 | lerns11=1)
0: no else if (lerns01>=1 & lerns01<=14 & (lerns02=-1)) | (lernmost>=1 & lernmost<=15) | (mlernm01>=1 & mlernm01<=14)
9: not answered if missing

Note: Variables mseddad ... msedoth are all derived from lernmost and mlernm01 ... mlernm15, and are coded in a similar way to msedmum:

(U – see below) (ND) **mseddad** main source of sex education was father
(R) (ND) **msedbro** main source of sex education was brother(s)
(R) (ND) **msedsis** main source of sex education was sister(s)
(U – see below) (ND) **msedrel** main source of sex education was other relative(s)
(U – see below) (ND) **msedsch** main source of sex education was lessons at school
(U – see below) (ND) **msedfrn** main source of sex education was friend(s) of a similar age
(U – see below) (ND) **msedfsp** main source of sex education was first boy/girlfriend/sexual partner
(U – see below) (ND) **mseddoc** main source of sex education was doctor/nurse/clinic
(R) (ND) **msedtv** main source of sex education was television/radio
(R) (ND) **msedvid** main source of sex education was video(s)
(R) (ND) **msedbks** main source of sex education was book(s)
(R) (ND) **msedmag** main source of sex education was magazines/newspapers
(U – see below) (ND) **msedoth** main source of sex education was another source(s)

Note: in Natsal-3 derivation of these binary variables was slightly different since respondent was only allowed to give one answer at lernmost

(D) **mseddoc** main source of sex education was doctor/nurse/clinic
1: yes if (sexedumum=1 & lernmost=-1) | lernmost=8
0: no else if (sexedumum=0 | (sexedumum=1 & lernmost≠8 & lernmost≠-1))
9: not answered if missing

Note: variables medintad...msededc derived in a similar way to mseddoc using different codes from lernmost

(ND) **msedintad** main source of sex education was internet-sexual advice websites
(D) **msedsch** main source of sex education was lessons at school
(ND) **msedinto** main source of sex education was internet-other
(ND) **msedbmn** main source of sex education was books/magazines/newspapers
(D) **msedfrn** main source of sex education was friend(s) of a similar age
(ND) **msedtvid** main source of sex education was tv/radio/DVDs/videos
(*D) **msedmum** main source of sex education was mother (including step/adoptive)
(*D) **mseddad** main source of sex education was father (including step/adoptive)
(D) **msedfsp** main source of sex education was first boy/girlfriend/sexual partner
(ND) **msedintpn** main source of sex education was internet-pornographic websites
(ND) **msedpor** main source of sex education was pornographic magazines/films
(ND) **msedbsis** main source of sex education was brother/sister (incl. half/step/adopted)
(D) **msedoth** main source of sex education was another source(s)
(ND) **msedani** main source of sex education was animals/growing up on farm
(D) **msedrel** main source of sex education was other relative(s)
(ND) **msedofri** main source of sex education was older friends
(ND) **msedofrn** main source of sex education was friends (age not specified)
(ND) **msedexp** main source of sex education was own experience
(ND) **msedfil** main source of sex education was cinemas/films (other than tv/radio)
(ND) **msededc** main source of sex education was lessons/courses (i.e. formal teaching)

LEARNING ABOUT SEX

Note: Variable sexedsr2 codes respondents who could not choose one main source of sex education as 'other'

(U – see below) (ND) **sexedsr2** main source of sex education, grouped

- 1: mother &/or father *if* lernmost=1 | 2 | (lernmost=-1 & (learns01=1 | lernmost=2))
- 2: lessons at school *if* lernmost=6 | (lernmost=-1 & learns01=6)
- 3: friends of about the same age *if* lernmost=7 | (lernmost=-1 & learns01=7)
- 4: other source *if* (lernmost=3 | lernmost=4 | lernmost=5 | lernmost=8 | lernmost=9 | lernmost=10 | lernmost=11 | lernmost=12 | lernmost=13 | lernmost=14 | lernmost=15) | (lernmost=-1 & (learns01=3 | learns01=4 | learns01=5 | learns01=8 | learns01=9 | learns01=10 | learns01=11 | learns01=12 | learns01=13 | learns01=14 | learns01=15))
- 9: not answered/can't remember *else if missing*

sexedsr2 main source of sex education, grouped

- (*) 1: mother &/or father (incl. adopted) *if* ((mseddmum=1 | mseddad=1) | (mseddmum=-1 & (sexedumum=1 | sexedudad=1)))
- 2: lessons at school *if* (msedsch=1 | (msedsch=-1 & sexedusch=1))
- 3: friends of about the same age *if* (msedfrn=1 | (msedfrn=-1 & sexedufn=1))
- 4: other source *if* ((mseddac=1 | msedintad=1 | msedinto=1 | msedbm=1 | msedvid=1 | msedfsp=1 | msedintpn=1 | msedpor=1 | msedbsis=1 | msedoth=1 | msedani=1 | msedrel=1 | msedofri=1 | msedofrn=1 | msedexp=1 | msedfil=1 | msededc=1) | (mseddac=-1 & (sexedudoc=1 | sexeduintad=1 | sexeduinto=1 | sexedudmn=1 | sexedutvid=1 | sexedufsp=1 | sexeduintpn=1 | sexedupor=1 | sexedubsis=1 | sexeduoth=1 | sexeduani=1 | sexedurel=1 | sexeduofri=1 | sexedofrn=1 | sexeduexp=1 | sexedufil=1 | sexeduedc=1)))
- 9: not answered/can't remember *else if missing*

Note: Variable sexedsr2 codes respondents who could not choose one main source of sex education as 'other'

(U – see below) (ND) **sexedsr2** main source of sex education, grouped

- 1: mother &/or father *if* lernmost=1 | lernmost=2 | (lernmost=-1 & (learns01=1 | learns01=2))
- 2: brother(s) &/or sister(s) *if* lernmost=3 | lernmost=4 | (lernmost=-1 & (learns01=3 | learns01=4))
- 3: lessons at school *if* lernmost=6 | (lernmost=-1 & learns01=6)
- 4: friend(s) of about the same age *if* lernmost=7 | (lernmost=-1 & learns01=7)
- 5: first boy/girl friend/sexual partner *if* lernmost=8 | (lernmost=-1 & learns01=8)
- 6: doctor/nurse/clinic *if* lernmost=9 | (lernmost=-1 & learns01=9)
- 7: media *if* lernmost=10 | lernmost=11 | lernmost=12 | lernmost=13 | (lernmost=-1 & (learns01=10 | learns01=11 | learns01=12 | learns01=13))
- 8: other source *if* (lernmost=5 | lernmost=14 | lernmost=15) | (lernmost=-1 & (learns01=5 | learns01=14 | learns01=15))
- 9: not answered/can't remember *else if missing*

sexedsr2 main source of sex education, grouped

- (*) 1: mother &/or father (incl. adopted) *if* ((mseddmum=1 | mseddad=1) | (mseddmum=-1 & (sexedumum=1 | sexedudad=1)))
- 2: brother &/or sister (incl. half/step/adopted) *if* (msedbsis=1 | (msedbsis=-1 & sexedubsis=1))
- 3: lessons at school *if* (msedsch=1 | (msedsch=-1 & sexedusch=1))
- 4: friends of about the same age *if* (msedfrn=1 | (msedfrn=-1 & sexedufn=1))
- 5: first boy/girlfriend/sexual partner *if* (msedfsp=1 | (msedfsp=-1 & sexedufsp=1))
- 6: doctor/nurse/clinic *if* (mseddac=1 | (mseddac=-1 & sexedudoc=1))
- 7: media *if* ((msedbm=1 | msedvid=1) | (msedbm=-1 & (sexedubmn=1 | sexedutvid=1)))
- 8: sexual advice websites *if* (msedintad=1 | (msedintad=-1 & sexeduintad=1))
- 9: pornography (internet or magazines/films) *if* ((msedintpn=1 | msedpor=1) | (msedintpn=-1 & (sexeduintpn=1 | sexedupor=1)))
- 10: other source *if* ((msedinto=1 | msedoth=1 | msedani=1 | msedrel=1 | msedofri=1 | msedofrn=1 | msedexp=1 | msedfil=1 | msededc=1) | (msedinto=-1 & (sexeduinto=1 | sexeduoth=1 | sexeduani=1 | sexedurel=1 | sexeduofri=1 | sexeduofrn=1 | sexeduexp=1 | sexeduexp=1 | sexedufil=1 | sexeduedc=1)))
- 9: not answered/can't remember *else if missing*

LEARNING ABOUT SEX

Note: in Natsal-3, response options for learnmor have been expanded following responses given in Natsal-2

*Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code questions. The following binary variables were created directly from the question **learnmor** in Natsal-3 and derived from **learnm01-learnm11** in Natsal-2, and **q18b** in Natsal-1.*

Natsal-1 and Natsal-2 binary variables. Note variables knowbbd – knowstd are all coded in a similar way to knowgbd:

(U – see below) (q18ba D) **knowgbd** known more about how girls' bodies develop
0: no else if (learnm01>=1 & learnm01<=12) | learnm01=98
1: yes if learnm01=12 | (learnm01=1 | learnm02=1 | learnm03=1 | learnm04=1 | learnm05=1 | learnm06=1 | learnm07=1 | learnm08=1 | learnm09=1 | learnm10=1 | learnm11=1)
97: would liked to have known more but can't specify if learnm01=97
98: knew enough already if learnm01=98
99: none, not yet ready for sexual experience if learnm01=99
998: can't remember if learnm01=998
999: not answered if missing

Note: Variables knowbbd ... knowstd are all derived from learnm01 learnm11 and are coded in a similar way to knowgbd:

(U – see below) (q18bq D) **knowbbd** known more about how boys' bodies develop
(U – see below) (q18bb D) **knowborn** known more about how babies are born
(U – see below) (q18bh D) **knowsex** known more about sexual intercourse
(U – see below) (q18br D) **knowfp** known more about contraception
(U – see below) (q18bl D) **knowhom** known more about homosexuality/lesbianism
(U – see below) (q18bt D) **knowmast** known more about masturbation
(U – see below) (q18bk D) **knowsati** known more about how to make sex more satisfying
(U – see below) (q18bs D) **knowno** known more about saying 'no' to sex
(U – see below) (q18bm D) **knowfeel** known more about sexual feelings, emotions & relationships
(U – see below) (q18bd D) **knowstd** known more about STDs

Natsal-3 binary variables. Variables knowall-knowborn are all coded in a similar way to knowosp:

Note: if respondent would have liked to know more about all of the list then each variable is coded as 1 (yes)

(N) **knowosp** known more about other sexual practices

0: no

1: yes

97: would liked to have known more but can't specify

98: knew enough already

99: none, not yet ready for sexual experience

999: not answered if missing

(N) **knowall** known more about all these things

(U) **knowgbd** known more about how girls' bodies develop

(U) **knowstd** known more about STDs

(N) **knowsafe** known more about safer sex

(U) **knowsex** known more about sexual intercourse

(U) **knowcond** known more about how to use condom correctly

(U) **knowsati** known more about how to make sex more satisfying

(U) **knowhom** known more about homosexuality/lesbianism

(U) **knowfeel** known more about sexual feelings, emotions & relationships

(U) **knowbbd** known more about how boys' bodies develop

(U) **knowfp** known more about contraception

(U) **knowno** known more about saying 'no' to sex

(U) **knowmast** known more about masturbation

(U) **knowborn** known more about how babies are born

LEARNING ABOUT SEX

(N) **knownsure** known more: not sure what
0: no
1: yes
98: knew enough already
99: none, not yet ready for sexual experience
999: not answered *if* missing

(N) **knownone** known more about nothing, knew enough already
0: no
1: yes
97: would liked to have known more but can't specify
99: none, not yet ready for sexual experience
999: not answered *if* missing

(N) **knownsex** known more: not ready for sex
0: no
1: yes
97: would liked to have known more but can't specify
98: knew enough already
999: not answered *if* missing

(U – see below) (ND) **lernmorg** type of information needed more of before 1st sex
-1: not applicable *if* knowgbd=99 | knowgbd=999
0: knew enough already *if* learnm01=98
1: just biological information *if* (knowgbd=1 | knowbbd=1 | knowborn=1 | knowsex=1) & knowfp=0 & knowhom=0 & knowmast=0 & knowsati=0 & knowno=0 & knowfeel=0 & knowstd=0
2: just risk-reduction information *if* (knowfp=1 | knowstd=1) & knowgbd=0 & knowbbd=0 & knowborn=0 & knowsex=0 & knowhom=0 & knowmast=0 & knowsati=0 & knowno=0 & knowfeel=0
3: just psycho-sexual information *if* (knowhom=1 | knowmast=1 | knowsati=1 | knowno=1 | knowfeel=1) & knowfp=0 & knowstd=0 & knowgbd=0 & knowbbd=0 & knowborn=0 & knowsex=0
4: more than one type of information *else if* missing
9: not answered/can't remember *if* knowgbd=998 | 999

lernmorg type of information needed more of before 1st sex
-1: not applicable *if* knowgbd=99
0: knew enough already *if* knowgbd=98
1: just biological information *if* (knowgbd=1 | knowbbd=1 | knowborn=1 | knowsex=1) & knowosp=0 & knowstd=0 & knowsafe=0 & knowcond=0 & knowsati=0 & knowhom=0 & knowfeel=0 & knowfp=0 & knowno=0 & knowmast=0
2: just risk-reduction information *if* (knowstd=1 | knowsafe=1 | knowcond=1 | knowfp=1) & knowgbd=0 & knowsex=0 & knowbbd=0 & knowborn=0 & knowosp=0 & knowsati=0 & knowhom=0 & knowfeel=0 & knowno=0 & knowmast=0
3: just psycho-sexual information *if* (knowosp=1 | knowsati=1 | knowhom=1 | knowfeel=1 | knowno=1 | knowmast=1) & knowstd=0 & knowsafe=0 & knowcond=0 & knowfp=0 & knowgbd=0 & knowsex=0 & knowbbd=0 & knowborn=0
4: biological and risk-reduction information *if* (knowgbd=1 | knowsex=1 | knowbbd=1 | knowborn=1) & (knowstd=1 | knowsafe=1 | knowcond=1 | knowfp=1) & (knowosp=0 & knowsati=0 & knowhom=0 & knowfeel=0 & knowno=0 & knowmast=0)
5: biological and psycho-sexual information *if* (knowgbd=1 | knowsex=1 | knowbbd=1 | knowborn=1) & (knowosp=1 | knowsati=1 | knowhom=1 | knowfeel=1 | knowno=1 | knowmast=1) & (knowstd=0 & knowsafe=0 & knowcond=0 & knowfp=0)
6: risk-reduction & psycho-sexual *if* (knowstd=1 | knowsafe=1 | knowcond=1 | knowfp=1) & (knowosp=1 | knowsati=1 | knowhom=1 | knowfeel=1 | knowno=1 | knowmast=1) & (knowgbd=0 & knowsex=0 & knowbbd=0 & knowborn=0)
7: all 3 types of information *if* (knowgbd=1 | knowsex=1 | knowbbd=1 | knowborn=1) & (knowstd=1 | knowsafe=1 | knowcond=1 | knowfp=1) & (knowosp=1 | knowsati=1 | knowhom=1 | knowfeel=1 | knowno=1 | knowmast=1)
8: more information needed but not sure what
9: not answered *if* knowgbd=999

LEARNING ABOUT SEX

(U – see below) (ND) **adeqinfo** whether needed more information at 1st sex, grouped
1: needed more information *if* (learnm01>=1 & learnm01<=12) | learnm01=97
2: knew enough *if* learnm01=98
3: not ready for sexual experience *if* learnm01=99
9: not answered/can't remember *if* learnm01=998 | learnm01=999

(U) **adeqinfo** whether needed more information at 1st sex, grouped
1: needed more information *if* (any of knowosp...knowborn =1) | knowosp=97
2: knew enough *if* knowosp=98
3: not ready for sexual experience *if* knowosp=99
9: not answered/can't remember *if* knowosp=998 | knowosp=999

*Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code questions. The following binary variables were created directly from the question **wholern** (and any commonly-occurring other answers at **xwholern**) in Natsal-3, and derived from **wholern1-wholern2** in Natsal-2, and **q18c** in Natsal-1.*

*Natsal-1 and Natsal-2 binary variables. Note variables **dadadfp** – **othadfp** are all coded in a similar way to **mumadfp**:*

(U – see below) (q18cp D) **mumadfp** learned additional info. about sex from mother
1: yes *if* wholern1=1 | wholern1=2 | ((wholern1=14 | wholern2=14) & (wholernx=1))
0: no *else if* wholern1>=1 & wholern1<=14
9: not answered *else if* missing
-1: not applicable *if* wholern1=-1

*Note: Variables **dadadfp** ... **othadfp** are all derived from **wholern1**, **wholern2**, and **wholernx**, and have the same codes as for **mumadfp**:*

(U – see below) (q18cz D) **dadadfp** learned additional info. about sex from father
(R) (q18cx D) **brotadfp** learned additional info. about sex from brother(s)
(R) (q18cl D) **sistadfp** learned additional info. about sex from sister(s)
(U – see below) (q18cn D) **reladfp** learned additional info. about sex from other relative(s)
(U – see below) (q18cd D) **schadfp** learned additional info. about sex from lessons at school
(U – see below) (q18cj D) **frieadfp** learned additional info. about sex from friends of a similar age
(U – see below) (q18cs D) **firpadfp** learned additional info. about sex from first boy/girlfriend/sexual partner
(U – see below) (q18ca D) **docadfp** learned additional info. about sex from doctor/nurse/clinic
(R) (q18ck/q18ce D) **tvadfp** learned additional info. about sex from television/radio
(R) (ND) **vidadfp** learned additional info. about sex from video(s)
(R) (q18cg D) **bookadfp** learned additional info. about sex from books
(R) (q18cv D) **magsadfp** learned additional info. about sex from magazines/newspapers
(U – see below) (q18cq D) **othadfp** learned additional info. about sex from another source(s)

*Natsal-3 binary variables. Note variables **intadadfp** - **reladfp** are all coded in a similar way to **docadfp**:*

(U) **docadfp** wanted additional info. about sex from doctor/nurse/clinic
0: no
1: yes
9: not answered *if* missing | knowosp=999
-1: not applicable *if* knowosp=98 | knowosp=99

*Note: Variables **intadadfp** ... **reladfp** all have the same codes as for **docadfp**:*

(N) **intadadfp** wanted additional info. about sex from internet-sexual advice websites
(U) **schadfp** wanted additional info. about sex from lessons at school
(N) **intoadfp** wanted additional info. about sex from internet-other
(U) **bm nadfp** wanted additional info. about sex from books/ magazines/newspapers
(U) **frieadfp** wanted additional info. about sex from friends of a similar age
(U) **tvidadfp** wanted additional info. about sex from tv/radio/DVDs/videos
(U) **mumadfp** wanted additional info. about sex from mother (incl. step or adoptive)
(U) **dadadfp** wanted additional info. about sex from father (incl. step or adoptive)

LEARNING ABOUT SEX

- (U) **fspadfp** wanted additional info. about sex from first boy/girlfriend/sexual partner
- (N) **intpnadfp** wanted additional info. about sex from internet-pornographic websites
- (N) **poradfp** wanted additional info. about sex from pornography magazines/films
- (N) **bsisadfp** wanted additional info. about sex from brother/sister (incl. half, step or adoptive)
- (U) **othadfp** wanted additional info. about sex from another source(s)
- (U) **reladfp** wanted additional info. about sex from other relative(s)

Note: in Natsal-3 wholerng was removed since it was no longer possible to determine which answer option was given first.

Note: Variable 'wholerng' codes respondents according to their response to their first response to the wholern variables, i.e. wholern1. Thus, this can be considered as their main preferred source of additional information

- (R) (ND) **wholerng** main preferred source of additional sex education, grouped
- 1: not ready for sexual intercourse
 - 0: knew enough already
 - 1: mother &/or father *if* wholern1=1 | wholern1=2 | (wholern1=14 & (wholernx=1 | wholernx=2))
 - 2: brother(s) &/or sister(s) *if* wholern1=3 | wholern1=4 | (wholern1=14 & (wholernx=3 | wholernx=4))
 - 3: lessons at school *if* wholern1=6 | (wholern1=14 & wholernx=6)
 - 4: friends of about the same age *if* wholern1=7 | (wholern1=14 & wholernx=7)
 - 5: first boy/girl friend/sexual partner *if* wholern1=8 | (wholern1=14 & wholernx=8)
 - 6: doctor/nurse/clinic *if* wholern1=9 | (wholern1=14 & wholernx=9)
 - 7: media *if* wholern1=10 | wholern1=11 | wholern1=12 | wholern1=13 | (wholern1=14 & (wholernx=10 | wholernx=11 | wholernx=12 | wholernx=13))
 - 8: other source *if* (wholern1=5) | (wholern1=14 & (wholernx=5 | wholernx=21 | wholernx>=14 & wholernx<=19))
 - 9: not answered/can't remember *else if missing*

FIRST SEXUAL EXPERIENCES

FIRST SEXUAL EXPERIENCES

Note: firstint, xfirstint, nextint, nextage, xnxtage, firstexp, and xfirstexp have been removed from the final dataset in Natsal-3. This was to avoid confusion as the derived variables afsex to afexp are more appropriate to use in analysis.

(R) (q19a) **firstint** age at 1st het. sex. intercourse

95: not sure of age

96: has not happened yet

97: refusal

99: not answered

(R) (q19a1) **xfirstint** estimated at age 1st het. sex. intercourse

-1: not applicable if firstint≠ 95

99: not answered

Note: in Natsal-3 nextint, nextage and xnxtage were moved here in the questionnaire. Previously positioned after xfirstexp

(R) (M) (N) **nextint** had het. sex. intercourse aged 13+ if 1st sex age<13

-1: not applicable if firstint≥13

1: yes

2: not with anyone else since age<13

97: refusal

99: not answered

(R) (M) (q19d) **nextage** age at next het. sex. intercourse if 1st sex age<13

-1: not applicable if firstint≥13

95: not sure of age

96: has not happened since age<13 if nextint =2

97: refusal

99: not answered

(R) (M) (q19d1) **xnxtage** estimated next age at het. sex. intercourse if 1st sex age<13

-1: not applicable if nextage ≠95

99: not answered

(R) (q19b) **firstexp** age at 1st het. sex. experience

95: not sure of age

96: has not happened yet

97: refusal

99: not answered

(R) (q19b1) **xfirstexp** estimated age at 1st het. sex. experience

-1: not applicable if firstexp≠ 95

99: not answered

Note: In Natsal-2 nextint nextage and xnextage were here in the questionnaire. Moved up after xfirstint following respondent comments that previous order was difficult to follow.

FIRST SEXUAL EXPERIENCES

(afsex D) **afsex** age at 1st het. sex. intercourse if aged 13+ (incl. estimates)

(U) if firstint>12 & firstint<75 copy firstint if firstint>12 & firstint<45 copy firstint

(U) if firstint<13 & nextage>12 & nextage<75 copy nextage if firstint<13 & nextage>12 & nextage<45
copy nextage

if firstint=95 & firstint>12 & xfirstint<99 copy xfirstint

if nextage=95 & xnxtage>12 & xnxtage<99 copy xnxtage

if firstint=95 & xfirstint<13 & nextage>12 & nexting<95 copy nextage

if firstint=95 & xfirstint<13 & xnxtage>12 & xnxtage<99 copy xnxtage

90: age<13 and not since if (firstint<13 & nextint=2)

96: has not happened yet if firstint=96

97: refusal if firstint=97

99: not answered if firstint=99 | nextage=99 | (firstint=96 & hetcheck=1)

Note: Variable 'gafsex' in Natsal-1 does not recode those for whom afsex=90 | afsex=97 | afsex=99

(gafsex D) **gafsex** age at 1st het. sex. intercourse if aged 13+ (incl. estimates), grouped

1: 13-15 years if afsex>=13 & afsex <=15

2: 16-17 years afsex>=16 & afsex <=17

3: 18-19 years afsex>=18 & afsex <=19

4: 20-24 years afsex>=20 & afsex <=24

5: 25+ years afsex>=25 & afsex <=44

6: hasn't happened yet afsex=96

(U) -1: if afsex=90 | afsex=97 | afsex=99 sysmis if afsex=90 | afsex=97 | afsex=99

(ND) **afsexgp** age at 1st het. sex. intercourse if aged 13+ (incl. estimates), grouped

1: 13-15 if afsex=13 | afsex=14 | afsex=15

2: 16-17 if afsex=16 | afsex=17

3: 18-19 if afsex=18 | afsex=19

4: 20+ if afsex>=20 & afsex<=44

96: hasn't happened yet/since age12 if afsex=96 | afsex=90

99: refused/not answered if afsex=97 | afsex=99

(afsexall D) **afsexall** age at 1st het. sex. intercourse not limited to age 13+ (incl. estimates)

if firstint<45 copy firstint

if firstint=95 copy xfirstint

96: has not happened yet if firstint=96

97: refusal if firstint=97

99: not answered if firstint=99

Notes: 1. missing cases are not assigned to a missing code in 'hetb416'

2. hetb416 is derived using 'afsexall' rather than 'afsex'

(U) (D) **hetb416** had het. sex. intercourse before age 16

0: no if (afsexall>=16 & afsexall<=44) | afsexall=96

1: yes if afsexall<16

(U) 9: refused / not answered if afsexall=97 | afsexall=99

Notes: 1. missing cases are not assigned to a missing code in 'hetb416a'

2. hetb416a is derived using 'afsex' rather than 'afsexall'

(U) (D) **hetb416a** had het. sex. intercourse before age 16

0: no if (afsex>=16 & afsex<=44) | afsex=90 | afsex=96

1: yes if afsex<16

(U) 9: refused / not answered if afsexall=97 | afsexall=99

(D) **afexp** age at 1st het. sex. experience (incl. estimates)

if firstexp<45 copy firstexp

if firstexp=95 copy xfirstexp

96: has not happened yet if firstexp=96

97: refusal if firstexp=97

99: not answered if firstexp=99

FIRST SEXUAL EXPERIENCES

(R) (q20a) **privacy** private enough to ask questions about 1st het. sex. intercourse aged 13+
1: yes
2: definitely not
9: not answered
-1: not applicable *if firstint=96 | nextage=96*

(R) (q20b) **feaccept** offered & accepted self-completion sheet about 1st het. sex. intercourse
1: sheet accepted
2: prefers to be asked questions
9: not recorded
-1: not applicable *if privacy=-1 | privacy=1*

(R) (q20c) **whyfecd** why self-completion sheet not completed
1: another person (regardless of age) in the room
2: prefers to be asked questions
3: other reason
9: not recorded
-1: not applicable *if privacy=-1 | privacy=1*

(R) (q20d) **fecomp** check completion of self-completion sheet
1: sheet completed, no queries
2: sheet completed after query/ies
3: sheet not completed/refusal
4: sheet completed and entered by interviewer

Notes: 1. if not private enough to ask about first sexual intercourse, these questions were asked in the computer self-completion (CASI) in N Natsal-3. In Natsal-2 & Natsal-1 respondents were asked to complete a paper self-completion (PAPI). For Natsal-3 the variable zmodefexp indicates whether the respondents completed the questions face-to-face or using CASI (see below).

2. in Natsal-3 the remaining questions in the first sexual experiences section were asked either completely in the CAPI, partially CAPI and partially CASI or completely CASI depending on the privacy of the situation. Data collected in these two modes has been combined into single variables which are prefixed with a 'z' to indicate that they originate from both these modes. Variables derived from these do not have a z prefix. The variable zmodefexp gives the mode of completion for each respondent. In Natsal-2 the same variables do not have a z prefix.

3. in Natsal-3 the wording of age1part was changed from 'first new partner' to 'first person' to be consistent with the rest of the questionnaire where use of the word partner is avoided.

(ND) **zmodefexp** interview mode for first sexual experiences, CAPI or CASI interview
-1: Not applicable *if afsex=90 | afsex=96 | afsex=97 | afsex=99*
1: CAPI – private enough *if privacy=1 & privat2=1*
2: CAPI/CASI – initially private but later not private enough *if privacy=1 & privat2=2*
3: CAPI/CASI – initially private enough but forced so NA *if privacy=1 & privat2=-1*
4: CASI – not private enough *if privacy=2 & scaccept#3*
5: No CAPI/CASI – not private enough and refused CASI *if privacy=2 & scaccept=3*

(*) (U) (q21a) **zage1part** age of 1st person (previously partner) if 1st het. sex. intercourse aged 13+
95: estimated partner's age
97: never knew partner's age
98: can't remember partner's age
99: not answered
(U) -1: not applicable *if firstint=96 | nextage=96 | afsex=90 | afsex=96 | afsex=97 | afsex=99 | zmodefexp=5*

Note: in Natsal-3 there is no CASI version of xage1prt since respondents were instructed to approximate an age in zage1part

FIRST SEXUAL EXPERIENCES

(q21a1) **zage1prt** estimated age of 1st partner if 1st het. sex. intercourse aged 13+
99: not answered
-1: not applicable if age1part#95

(ND) **agefsp** age of 1st partner (incl. estimates) if 1st het. sex. intercourse aged 13+
copy age1part if age1part>0 & age1part <95
copy xage1prt if age1part=95
97: never knew partner's age if age1part=97
98: can't remember partner's age if age1part=98
99: not answered if age1part=99 | (age1part=95 & xage1prt=99)
-1: not applicable if age1part=-1

(q21b) **zparts1st** 1st het. sex. intercourse ever for partner too if 1st het. sex. intercourse aged 13+
1: yes, first time
2: think it was first time
3: think it was not first time
4: no, not first time
8: can't remember
9: not answered
(U) -1: not applicable if firstint=96 | nextage=96 | afsex=90 | afsex=96 | afsex=97 | afsex=99 |
zmodexp=5

(q22a) **zbothwill** respondent and partner both willing to have sex if 1st het. sex. intercourse aged 13+
1: both willing
2: respondent more willing
3: partner more willing
9: not answered
(U) -1: not applicable if firstint=96 | nextage=96 | afsex=90 | afsex=96 | afsex=97 | afsex=99 |
zmodexp=5

(q22b) **zprrwill** partner was more willing to have sex if 1st het. sex. intercourse aged 13+
1: respondent also willing
2: respondent had to be persuaded
3: respondent was forced
9: not answered
(U) -1: not applicable if firstint=96 | nextage=96 | afsex=90 | afsex=96 | afsex=97 | afsex=99 |
zmodexp=5 | bothwill=1 | bothwill=2 | bothwill=9

Note: in Natsal-3 prtwill2 renamed partmwill due to changes to variable names in the program. Variables are equivalent.

partmwill (ND) **prtwill2** partner was more willing to have sex if 1st het. sex. intercourse aged 13+ ,
grouped
0: no if bothwill=1 | bothwill=2
1: yes if bothwill=3
9: not answered/can't remember if bothwill=9
-1: not applicable if bothwill=-1

Note: In Natsal-3, respondents who reported the first sex was forced (prtwill=3) were asked a slightly different version of the contraception and relationship questions from other respondents and were no longer asked fstmet, rtime and mainreas
Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code variables. forcond..fornone were created directly from forprec. All variables are coded similarly to forcond.

(N) **forcond** used condom with partner at 1st het sex intercourse aged 13+ if forced
-1: not applicable if prtwill#3

FIRST SEXUAL EXPERIENCES

0: no
1: yes
9: not answered

And similarly...

(N) **forpill** used pill with partner at 1st het sex intercourse aged 13+ if forced

(N) **forecont** used emergency contraception with partner at 1st het sex intercourse aged 13+ if forced

(N) **foroth** used other contraception with partner at 1st het sex intercourse aged 13+ if forced

(N) **fornone** No contraception used with partner at 1st het sex intercourse aged 13+ if forced

(N) **fsprecfgp** Contraception used at 1st het. sex. intercourse aged 13+ if forced, grouped

-1: not applicable if forcond=-1

1: only condom if forcond=1 & forpill=0 & forecont=0 & foroth=0 & fornone=0

2: only pill or only other if (forcond=0 & forpill=1 & forecont=0 & foroth=0 & fornone=0) | (forcond=0 & forpill=0 & forecont=0 & foroth=1 & fornone=0)

3: emergency contraception only if forcond=0 & forpill=0 & forecont=1 & foroth=0 & fornone=0

4: combination of reliable methods (condom/pill/other/EC) if ((forcond=1 & forpill=1) | (forcond=1 & foroth=1) | (forcond=1 & forecont=1) | (forpill=1 & foroth=1) | (forpill=1 & forecont=1))

5: none if forcond=0 & forpill=0 & forecont=0 & foroth=0 & fornone=1

9: not answered if forcond=9

(ND) **contrameth** reliable contraception used at first sex, if first sex not forced

-1: not applicable if fsprecnfgp=-1

1: yes if fsprecnfgp=1 | fsprecnfgp=2 | fsprecnfgp=3 | fsprecnfgp=8

2: no if fsprecnfgp=4 | fsprecnfgp=5 | fsprecnfgp=6 | fsprecnfgp=7

9: not answered if fsprecnfgp=9

(ND) **contramethall** reliable contraception used at first sex

-1: not applicable if het=0

1: yes if contrameth=1 | (zprtwill=3 & fsprecfgp>=1 & fsprecfgp<=4)

2: no if contrameth=2 | (zprtwill=3 & fsprecfgp=5)

9: not answered if het=9 | zmodexp=5 | zbothwill=9 | zprtwill=9 | fsprecfgp=9 | contrameth=9

(N) **zforjmet** type of relationship with partner at 1st het. sex. intercourse aged 13+ if forced

-1: not applicable if prtwill ≠3

1: someone I didn't know

2: recently met

3: known each other for a while, but were not in a steady relationship

4: used to be in a steady relationship, but were not at that time

5: steady relationship

6: living together as a couple / married at the time

7: Something else

9: Not answered

Note: Binary variables fscondom..fsneither originate from the questions anyprec1 (Natsal-2) and anypre2 (Natsal-3). Only the binary variables are shown here.

Note: Variables fscondom ... fscontare are coded in a similar way to fscondom:

(U – see below) (ND) **fscondom** used condom with partner at 1st het. sex. intercourse aged 13+

1: yes if anyprec1=1 | anyprec2=1 | anyprec3=1 | anyprec4=1 | anyprec5=1

0: no else if anyprec1>=1 & <=7

8: can't remember if anyprec1=8

9: not answered if anyprec1=9 or if missing

-1: not applicable if anyprec1=-1 & anyprec2=-1 & anyprec3=-1 & anyprec4=-1 & anyprec5=-1

And similarly...

(U – see below) (ND) **fspill** used contraceptive pill with partner at 1st het. sex. intercourse aged 13+

(U – see below) (ND) **fsother** used other contraception with partner at 1st het. sex. intercourse aged 13+

(U – see below) (ND) **fswithdw** used withdrawal with partner at 1st het. sex. intercourse aged 13+

(U – see below) (ND) **fsrhythm** used rhythm method with partner at 1st het. sex. intercourse aged 13+

FIRST SEXUAL EXPERIENCES

(U – see below) (ND) **fspartdk** didn't use contraception with partner at 1st het. sex. intercourse aged 13+ but partner may have used some (U – see below) (ND) **fsneithr** neither used any contraception with partner at 1st het. sex. intercourse aged 13+
(R) (ND) **fscantre** can't remember if used any contraception with partner at 1st het. sex. intercourse aged 13+

Note: these binary variables were updated in Natsal-3 as there were some new answer options, and those who reported being forced at first sex were asked a different version of this question (see forcond..fornone on page 35).

(U) **fscondom** used condom with partner at 1st het. sex. intercourse aged 13+ & not forced
-1: not applicable if (bothwill#1 & bothwill#2) | (bothwill==3 & (prtwill#1 | prtwill#2))
0: no
1: yes
9: not answered

(U) **fspill** used contraceptive pill with partner at 1st het. sex. intercourse aged 13+ & not forced
(N) **fsecont** used emergency contraception with partner at 1st het. sex. intercourse aged 13+ & not forced
(U) **fsother** used other contraception with partner at 1st het. sex. intercourse aged 13+ & not forced
(U) **fswithdw** used withdrawal with partner at 1st het. sex. intercourse aged 13+ & not forced
(U) **fsrhythm** used rhythm method with partner at 1st het. sex. intercourse aged 13+ & not forced

(U) **fspartdk** didn't use contraception with partner at 1st het. sex. intercourse aged 13+ but partner may have used some

(U) **fsneithr** neither used any contraception with partner at 1st het. sex. intercourse aged 13+

Note: fsprecgp & fsprecg2 renamed in Natsal-3 as these now only correspond to those whose first sexual experience was not forced and in Natsal-3 emergency contraception was included as a separate answer option.

Note: Variable fsprecgp is derived to be consistent with Natsal-1's variable 'q23' but fsprecgp combines 'other' with 'pill'

(R) (q23) **fsprecgp** contraception used at 1st het. sex. intercourse aged 13+, grouped
1: only condom if (fscondom=1 & fspill=0 & fsother=0 & fswithdw=0 & fsrhythm=0 & fspartdk=0 & fsneithr=0 & fscantre=0) else if fscondom=1 & (fswithdw=1 | fsrhythm=1)
2: only 'other' (incl. pill) if fscondom=0 & (fspill=1 | fsother=1) & fswithdw=0 & fsrhythm=0 & fspartdk=0 & fsneithr=0 & fscantre=0 else if fspill=1 & (fswithdw=1 | fsrhythm=1)
3: only used withdrawal if fscondom=0 & fspill=0 & fsother=0 & fswithdw=1 & fsrhythm=0 & fspartdk=0 & fsneithr=0 & fscantre=0 else if fsother=1 & (fswithdw=1 | fsrhythm=1) or anyprec1=4
4: only rhythm method if fscondom=0 & fspill=0 & fsother=0 & fswithdw=0 & fsrhythm=1 & fspartdk=0 & fsneithr=0 & fscantre=0 or anyprec1=5
5: respondent didn't use but partner may have done if fscondom=0 & fspill=0 & fsother=0 & fswithdw=0 & fsrhythm=0 & fspartdk=1 & fsneithr=0 & fscantre=0 or anyprec1=6
6: neither respondent nor partner used contraception if fscondom=0 & fspill=0 & fsother=0 & fswithdw=0 & fsrhythm=0 & fspartdk=0 & fsneithr=1 & fscantre=0 or anyprec1=7
7: used condom & pill/other if (fscondom=1 & fspill=1) | (fscondom=1 & fsother=1) | (fspill=1 & fsother=1) | (fscondom=1 & fspill=1 & fsother=1)
8: can't remember if fscondom=8 & fspill=8 & fsother=8 & fswithdw=8 & fsrhythm=8 & fspartdk=8 & fsneithr=8 & fscantre=1
9: not answered if fscondom=9 & fspill=9 & fsother=9 & fswithdw=9 & fsrhythm=9 & fspartdk=9 & fsneithr=9 & fscantre=9
-1: not applicable if anyprec1=-1

(R) (ND) **fsprecg2** contraception used at 1st het. sex. intercourse aged 13+, grouped
1: used condoms (poss. with other method(s)) if fsprecgp=1 | fsprecgp=7

FIRST SEXUAL EXPERIENCES

- 2: used other reliable (incl. pill) but not condom *if* `fsprecgp=2` | (`fscondom=0` & `fspill=0` & `fsother=1` & `fswithdw=0` & `fsrhythm=0` & `fspartdk=0` & `fsneithr=0` & `fscantre=0`)
3: did not use reliable (withdrawal/rhythm method) *if* `fsprecgp=3` | `fsprecgp=4`
4: respondent didn't use any, don't know about partner *if* `fsprecgp=5` | `fsprecgp=6`
9: can't remember/not answered *if* `fsprecgp=8` | `fsprecgp=9`
-1: not applicable *if* `fsprecgp=-1`

Note: categories for `fsprecnfgp` coded in a hierarchy of reliable methods only, then unreliable methods only, then none options, then combinations of reliable methods, then finally any remaining combinations are coded in ascending order (i.e. else if)

(ND) **fsprecnfgp** contraception used at 1st het. sex. intercourse aged 13+, not forced, grouped

- 1: only condom *if* (`fscondom=1` & `fspill=0` & `fsecont=0` & `fsother=0` & `fswithdw=0` & `fsrhythm=0` & `fspartdk=0` & `fsneithr=0`) *else if* `fscondom=1` & (`fswithdw=1` | `fsrhythm=1`)
2: only pill/'other' *if* (`fscondom=0` & `fspill=1` & `fsecont=0` & `fsother=0` & `fswithdw=0` & `fsrhythm=0` & `fspartdk=0` & `fsneithr=0`) | (`fscondom=0` & `fspill=0` & `fsecont=0` & `fsother=1` & `fswithdw=0` & `fsrhythm=0` & `fspartdk=0` & `fsneithr=0`) *else if* (`fspill=1` & (`fswithdw=1` | `fsrhythm=1` | `fspartdk=1` | `fsneithr=1`)) | (`fsother=1` & (`fswithdw=1` | `fsrhythm=1` | `fspartdk=1` | `fsneithr=1`))
3: only emergency contraception *if* (`fscondom=0` & `fspill=0` & `fsecont=1` & `fsother=0` & `fswithdw=0` & `fsrhythm=0` & `fspartdk=0` & `fsneithr=0`) *else if* `fsecont=1` & (`fswithdw=1` | `fsrhythm=1` | `fspartdk=1` | `fsneithr=1`)
4: only withdrawal method *if* (`fscondom=0` & `fspill=0` & `fsecont=0` & `fsother=0` & `fswithdw=1` & `fsrhythm=0` & `fspartdk=0` & `fsneithr=0`) *else if* `fswithdw=1` & (`fsrhythm=1` | `fspartdk=1` | `fsneithr=1`)
5: only rhythm method *if* (`fscondom=0` & `fspill=0` & `fsecont=0` & `fsother=0` & `fswithdw=0` & `fsrhythm=1` & `fspartdk=0` & `fsneithr=0`) *else if* `fsrhythm=1` & (`fspartdk=1` | `fsneithr=1`)
6: respondent didn't use but partner may have done *if* (`fscondom=0` & `fspill=0` & `fsecont=0` & `fsother=0` & `fswithdw=0` & `fsrhythm=0` & `fspartdk=1` & `fsneithr=0`) *else if* `fspartdk=1` & `fsneithr=1`
7: neither respondent nor partner used contraception *if* (`fscondom=0` & `fspill=0` & `fsecont=0` & `fsother=0` & `fswithdw=0` & `fsrhythm=0` & `fspartdk=0` & `fsneithr=1`)
8: used combination of reliable methods *if* (`fscondom=1` & `fspill=1`) | (`fscondom=1` & `fsecont=1`) | (`fscondom=1` & `fsother=1`) | (`fspill=1` & `fsecont=1`) | (`fspill=1` & `fsother=1`) | (`fsecont=1` & `fsother=1`)
9: not answered *if* (`fscondom=9` & `fspill=9` & `fsecont=9` & `fsother=9` & `fswithdw=9` & `fsrhythm=9` & `fspartdk=9` & `fsneithr=9`)
-1: not applicable *if* `fscondom=-1`

(ND) **fsprecnfg2** contraception used at 1st het. sex summary, grouped

- 1: used condoms (poss. with other method(s)) *if* `fscondom==1`
2: used other reliable (incl. pill) but not condom *if* `fsprecnfgp=2` | `fsprecnfgp=3` | (`fscondom=0` & `fsprecnfgp=8`)
3: did not use reliable (withdrawal/rhythm method) *if* `fsprecnfgp=4` | `fsprecnfgp=5`
4: respondent didn't use any, don't know about partner *if* `fsprecnfgp=6` | `fsprecnfgp=7`
9: can't remember/not answered *if* `fsprecnfgp=9`
-1: not applicable *if* `fsprecnfgp=-1`

FIRST SEXUAL EXPERIENCES

*Note: response options for justmet updated and the same as R1rel in the most recent partners module.
Sex worker now captured in new question about how first met partner which is the same as the question in the most recent partners module*

(R) (q24) **justmet** type of relationship with partner at 1st het. sex. intercourse aged 13+

-1: not applicable if firstint=96 | nextage=96

1: prostitute

2: just met for the first time

3: had met recently

4: known each other for a while, but not a steady relationship

5: steady relationship

6: living together (but not engaged or married)

7: engaged

8: married

9: other

10: work colleague/employer

11: knew well/for a long time

12: rape/sexual abuse

98: can't remember

99: not answered

(N) **zjustmet2** type of relationship with partner at 1st het. sex. intercourse aged 13+ and not forced -
NEW CODES but question the same apart from eligibility

-1: not applicable if (bothwill#1 & bothwill#2) | (bothwill==3 & (prtwill#1 | prtwill#2))

(U) 1: just met for the first time

(U) 2: had met recently

(U) 3: known each other for a while, but not a steady relationship

(U) 4: used to be in a steady relationship but not at the time

5: steady relationship

(U*) 6: living together / married

9: Not answered

(R) (ND) **justmet2** respondent had just met/recently met their 1st partner

0: no if justmet=2 | justmet=3

1: yes if justmet=1 | justmet=4... justmet=12

9: can't remember/not answered if justmet=98 | justmet=99

-1: not applicable if justmet=-1

(N) **justmetN2** respondent had just met/recently met their 1st partner (not forced)

0: no if zjustmet2=3 | zjustmet2=4 | zjustmet2=5 | zjustmet2=6

1: yes if zjustmet2=1 | zjustmet2=2

9: can't remember / not answered if zjustmet2=9

-1: not applicable if (bothwill#1 & bothwill#2) | (bothwill==3 & (prtwill#1 | prtwill#2))

FIRST SEXUAL EXPERIENCES

(N) **zfstmet** how respondent met their 1st partner het. sex. Intercourse aged 13+ and not forced
-1: not applicable if (bothwill#1 & bothwill#2) | (bothwill==3 & (prtwill#1 | prtwill#2))

- 1: school
- 2: university or college
- 3: work / through work
- 4: pub, bar, night club, dance or disco
- 5: Introduced by friends or family
- 6: sports club, faith group, or other organisation or society
- 7: holiday / while travelling
- 8: Internet dating website
- 9: Other dating agency / personal ads
- 10: Online (not through dating website)
- 11: always known each other (for example as family friends or neighbours)
- 12: Neighbour/lived locally/house or flatshare
- 13: Arranged marriage
- 14: public place (e.g. park, cafe, shop, public transport)
- 15: sex worker / prostitute
- 16: Other
- 99: Not answered

Note: in Natsal-3 those reporting that first sex was forced were not asked rtime

(*) (q28) **zrtime** opinion now of timing of 1st het. sex. intercourse aged 13+ and not forced - question the same apart from eligibility

- 1: should have waited longer
- 2: should not have waited so long
- 3: about the right time
- 8: can't remember
- 9: not answered
- 1: not applicable if (bothwill#1 & bothwill#2) | (bothwill==3 & (prtwill#1 | prtwill#2))

Note: in Natsal-3 information was collected on all reasons for first intercourse (the question 'fstres') and then on the main reason (the question 'mainrea2'). Binary variables fstcurious..fstcrem come directly from the multicode question fstres. This was not asked of people reporting that their first sex was forced or for those who reported they were living as a couple with / married to their partner at 1st sex. Variables fstfeel..fstcrem are all coded in a similar way to fstcurious.

fstcurious curious at 1st het sex. Intercourse aged 13+ & not forced

- 1: not applicable if zjustmet=6 | (bothwill#1 & bothwill#2) | (bothwill==3 & (prtwill#1 | prtwill#2))0: no
- 1: yes
- 9: not answered

And similarly...

fstfeel was carried away by feeling at 1st het sex. Intercourse aged 13+ & not forced

fstpeers peers doing it at 1st het sex. Intercourse aged 13+ & not forced

fstfollow natural 'follow on' in the relationship at 1st het sex. Intercourse aged 13+ & not forced

fstdrunk bit drunk at 1st het sex. Intercourse aged 13+ & not forced

fstcann smoked some cannabis at 1st het sex. Intercourse aged 13+ & not forced

fstodrugs taken some other drugs at 1st het sex. Intercourse aged 13+ & not forced

fstlvirg wanted to lose virginity at 1st het sex. Intercourse aged 13+ & not forced

fstlove was in love at 1st het sex. Intercourse aged 13+ & not forced

fstothf other particular factor at 1st het sex. Intercourse aged 13+ & not forced

fstcrem can't remember reason at 1st het sex. Intercourse aged 13+ & not forced

FIRST SEXUAL EXPERIENCES

Note: response options to mainreas were updated in Natsal-3 therefore the variable has been renamed.

Note: Respondents who reported that they were married to their 1st partner are not asked the reason for 1st het. sex. intercourse

(R) (q26b) **mainreas** main reason for intercourse at the time of 1st het. sex. intercourse aged 13+

- 1: curious
- 2: carried away by feelings
- 3: peers doing it
- 4: natural follow-on in the relationship
- 5: a bit drunk at the time
- 6: wanted to lose my virginity
- 7: in love
- 8: other particular reason
- 9: can't chose/more than one reason
- 10: can't remember
- 11: to please partner
- 12: felt ready
- 13: rape/forced
- 99: not answered
- 1: not applicable if firstint=96 | nextage=96 | justmet=8

Note: In Natsal-3, respondents who reported that they were living together as a couple with / married to their 1st partner are not asked the main reason for 1st het. sex intercourse

Note: Those who chose more than one reason for first sex (question 'fstres') were then asked to choose a main read.

(N) **zmainrea2** main reason for intercourse at the time of 1st het. sex. intercourse aged 13+ and not forced and reported more than one reason

-1: not applicable if (zjustmet2=6 | (bothwill#1 & bothwill#2) | (bothwill==3 & (prtwill#1 | prtwill#2))) | reported only one reason at fstcurious..fstcrem)

- 1: curious
- 2: carried away by feelings
- 3: peers doing it
- 4: natural 'follow on' in the relationship
- 5: bit drunk
- (U) 6: smoked some cannabis
- (U) 7: taken some other drugs
- (U) 8: wanted to lose my virginity
- (U) 9: in love
- (U) 10: Other particular factor
- (U) 11: Can't remember
- 99: not answered

(R) (q26) **mmainre1** reason for intercourse at the time of 1st het. sex. intercourse aged 13+ 1

- 1: curious
- 2: carried away by feelings
- 3: peers doing it
- 4: natural follow-on in the relationship
- 5: a bit drunk at the time
- 6: wanted to lose my virginity
- 7: in love
- 8: other particular reason
- 99: not answered
- 1: not applicable if mainreas#9 & mainreas#99

FIRST SEXUAL EXPERIENCES

Similarly for up to 9 reasons, in addition to -1 if mmainren=99 & mmainren-1≠99

- (R) **mmainre2** reason for intercourse at the time of 1st het. sex. intercourse aged 13+ 2
- (R) **mmainre3** reason for intercourse at the time of 1st het. sex. intercourse aged 13+ 3
- (R) **mmainre4** reason for intercourse at the time of 1st het. sex. intercourse aged 13+ 4
- (R) **mmainre5** reason for intercourse at the time of 1st het. sex. intercourse aged 13+ 5
- (R) **mmainre6** reason for intercourse at the time of 1st het. sex. intercourse aged 13+ 6
- (R) **mmainre7** reason for intercourse at the time of 1st het. sex. intercourse aged 13+ 7
- (R) **mmainre8** reason for intercourse at the time of 1st het. sex. intercourse aged 13+ 8
- (R) **mmainre9** reason for intercourse at the time of 1st het. sex. intercourse aged 13+ 9

(U – see below) (q26am D) **curious** main reason had 1st het. sex. intercourse: curious

1: yes if mainreas=1 | (mmainre1=1 | mmainre2=1 | mmainre3=1 | mmainre4=1 | mmainre5=1 | mmainre6=1 | mmainre7=1 | mmainre8=1 | mmainre9=1)

0: no else if (mainreas>=1 & mainreas<=8) | ((mmainre1>=1 & mmainre1<=8) | (mmainre2>=1 & mmainre2<=8) | (mmainre3>=1 & mmainre3<=8) | (mmainre4>=1 & mmainre4<=8) | (mmainre5>=1 & mmainre5<=8) | (mmainre6>=1 & mmainre6<=8) | (mmainre7>=1 & mmainre7<=8) | (mmainre8>=1 & mmainre8<=8) | (mmainre9>=1 & mmainre9<=8))

9: not answered if missing

-1: not applicable if mainreas=-1 & mmainre1=-1

Note: Variables feelings ... othrfact are all derived from mainreas and mmainr1 ... mmainr11 and are coded in a similar way to curious:

(U – see below) (q26ac D) **feelings** main reason had first intercourse: carried away with feelings

(U – see below) (q26af D) **peers** main reason had first intercourse: peers doing it

(U – see below) (q26al D) **followon** main reason had first intercourse: natural follow on

(U – see below) (q26ar D) **drunk** main reason had first intercourse: drunk

(U – see below) (q26ah D) **losevirg** main reason had first intercourse: wanted to lose virginity

(U – see below) (q26ad D) **inlove** main reason had first intercourse: in love

(U – see below) (ND) **othrfact** main reason had first intercourse: other factor(s)

(R) (ND) **pleasprt** main reason had first intercourse: to please partner

(R) (q26aa D) **raped** main reason had first intercourse: raped/forced

Note: Variables curious ... othrfact are all created directly from zmainrea2 and are coded in a similar way to curious for each answer option at zmainrea2

(U) **curious** main reason had 1st het. sex. intercourse: curious

1: yes if zmainrea2=1

0: no if zmainrea2≠1

9: not answered if missing

-1: not applicable if zmainrea2=-1

(U) **feelings** main reason had first intercourse: carried away with feelings

(U) **peers** main reason had first intercourse: peers doing it

(U) **followon** main reason had first intercourse: natural follow on

(U) **drunk** main reason had first intercourse: drunk

(N) **smkcann** main reason had first intercourse: smoked cannabis

(N) **othdrug** main reason had first intercourse: taken other drugs

(U) **losevirg** main reason had first intercourse: wanted to lose virginity

(U) **inlove** main reason had first intercourse: in love

(U) **othrfact** main reason had first intercourse: other factor(s)

(U) **cantrem** main reason had first intercourse: can't remember

(U) **cantchoo** main reason had first intercourse: can't choose a main reason

FIRST SEXUAL EXPERIENCES

Note: There were small changes to the sexual competence variable in Natsal-3 due to i) additional answer options being available at mainrea2 (smoking cannabis and using other drugs); ii) an additional answer option at anypre2 (emergency contraception) and iii) those living together as a couple / married at the time of first sex not being routed into mainrea2.

Note: There were earlier versions of variable 'sexcomp'. This is the latest coding and featured in the Lancet paper by Wellings et al (2001)

(R) (ND) **sexcomp** 'sexual competence' at 1st het. sex. intercourse

1: competent *if* ((bothwill=1) & ((peers=0 & drunk=0 & raped=0 & pleasprt=0) | (peers=-1 & justmet=8)) & (rttime=3 | (rttime=-1 & justmet=8)) & (fscondom=1 | fspill=1))

0: not competent *else if* (bothwill≠8 & bothwill ≠9) & (peers≠9) & (rttime≠8 & rttime≠9) & (fscondom≠8 & fscondom≠9)

9: not answered

-1: not applicable *if* bothwill=-1

(ND) **sexcomp2** 'sexual competence' at 1st het. sex. intercourse

1: competent *if* ((zbothwill=1) & (peers=0 & drunk=0 & smkcann=0 & othdrg=0) & zrttime=3 & (fscondom=1 | fspill=1)) | (zjustmet=6 & peers=-1 & zrttime=3 & zbothwill=1)

0: not competent *else if* (zbothwill ≠9 & peers≠9 & zrttime≠9) & fscondom≠9) | ((zbothwill=2 | zbothwill=3) | (zrttime=1 | zrttime=2) | (fscondom=0 & fspill=0 & zjustmet2>=1 & zjustmet2<=5) | (peers=1 | drunk=1 | smkcann=1 | othdrg=1))

9: not answered

-1: not applicable *if* zbothwill=-1

CONTRACEPTION

CONTRACEPTION

*Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code questions. The following binary variables were created directly from the question **everuse2** in Natsal-3, and derived from **Everus01-Everus13** in Natsal-2, and **q30a** in Natsal-1.*

Natsal-1 and Natsal-2 binary variables for contraception use (ever) are given below:

(R)(ND) **minipill** ever used minipill

1: yes if everus01=1 | everus02=1 | everus03=1 | everus04=1 | everus05=1 | everus06=1 | everus07=1 | everus08=1 | everus09=1 | everus10=1 | everus11=1 | everus12=1 | everus13=1 | ((everus01=20 | everus02=20 | everus03=20 | everus04=20 | everus05=20 | everus06=20 | everus07=20 | everus08=20 | everus09=20 | everus10=20 | everus11=20 | everus12=20 | everus13=20) & (everusd1=1 | everusd2=1 | everusd3=1))

0: no else if everus01>=1 & everus20<=21

9: not answered if missing

-1: not applicable if everus01= -1

*Note: Variables **combpill** ... **otherfp** are all derived from everus01 ... everus20 and everusd1 ... everusd3, and are coded in a similar way to minipill:*

(R) (ND) **combpill** ever used combined pill

(R) (ND) **dkpill** ever used pill (but not sure which)

(ND) **mirena** ever used mirena

(U – see below) (q30ar D) **coil** ever used coil

(U – see below) (q30ad D) **condom** ever used condom

(U – see below) (ND) **femidom** ever used femidom

(U – see below) (q30aq D) **cap** ever used cap

(U – see below) (q30al D) **gels** ever used gels/sprays/pessaries

(R) (q30au D) **sponge** ever used contraceptive sponge

(R) (ND) **persona** ever used persona

(U – see below) (q30ak D) **rhythm** ever used rhythm method/safe period

(U – see below) (q30as D) **withdraw** ever used withdrawal

(U – see below) (ND) **injectns** ever used contraceptive injections

(U – see below) (ND) **implants** ever used contraceptive implants

(U) (ND) **emergency** ever used emergency contraception

(U – see below) (q30ac D) **fsterils** ever used female sterilisation

(U – see below) (q30am D) **msterils** ever used male sterilisation

(R) (q30ab D) **abstain** ever used abstinence

Note: Variable 'otherfp' is not entirely consistent with the corresponding variable in Natsal-1 since more contraceptive methods were available in Natsal-2 for respondents to choose from.

(U – see below) (q30aw D) **otherfp** ever used other contraception

(U – see below) (q30af D) **notused** never used contraception

1: yes if everus01=21

0: no else if everus01>=1 & everus20<=21

9: not answered if missing

-1: not applicable if everus01= -1

*Note: **conpill** was derived in Natsal-2 but is obtained directly from Everuse2 in Natsal-3 since only one pill option was given in Natsal-3.*

(U – see below) (q30aj D) **conpill** ever used any contraceptive pill

1: yes if minipill=1 | combpill=1 | dkpill=1

0: no else if minipill=0 & combpill=0 & dkpill=0

9: not answered if missing

-1: not applicable if everus01= -1

*.Natsal-3 binary variables for contraception use (ever) are given below. Variables **msterils**..**conpatch** are all coded in a similar way to **fsterils**.*

CONTRACEPTION

(U) **notused** never used contraception

-1: not applicable *if* afsex=90 | afsex=96 | afsex=97 | afsex=99

0: Used contraception

1: Never used contraception

9: not answered

(U) **fsterils** ever used female sterilisation

-1: not applicable *if* afsex=90 | afsex=96 | afsex=97 | afsex=99

0: No

1: Yes

9: not answered

And similarly...

(U) **msterils** ever used male sterilisation

(U) **conpill** ever used the pill

(U) **condom** ever used condom

(U) **femidom** ever used femidom

(N) **emcypill** ever used morning after pill

(N) **emcyiud** ever used emergency IUD

(U) **coil** ever used coil

(U) **mirena** ever used mirena

(U) **cap** ever used cap

(U) **injectns** ever used contraceptive injections

(U) **gels** ever used gels/sprays/pessaries

(U) **rhythm** ever used rhythm method/safe period

(U) **withdraw** ever used withdrawal

(U) **implants** ever used contraceptive implants

Note: Variable 'otherfp' is not entirely consistent with the corresponding variable in Natsal-2 since different contraceptive methods were available in Natsal-3 for respondents to choose from.

(U) (q30aw D) **otherfp** ever used other contraception

(N) **conpatch** ever used contraceptive patch

(UD) **emergency** ever used emergency contraception

-1: not applicable *if* afsex=90 | afsex=96 | afsex=97 | afsex=99

0: no *if* emcypill=0 & emcyiud=0

1: yes *if* emcypill=1 | emcyiud=1

9: not answered

*Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code questions. The following binary variables were created directly from the question **usedyr2** in Natsal-3, and derived from **UsedYe01-UsedYe08** in Natsal-2, and **q30b** in Natsal-1. CAPI only allows interviewer to enter method of contraception that has been used ever, or no method.*

Natsal-1 and Natsal-2 binary variables for contraception use (last year) are given below:

(R) (ND) **yminipil** in last year used minipill

1: yes *if* usedye01=1 | ... | usedye08=1

0: no *else if* usedye01>=1 & usedye01<=21

9: not answered *if* missing

-1: not applicable *if* usedye01= -1

Note: Variables ycompill ... yotherfp are coded in a similar way to yminipil:

(R) (ND) **ycompill** in last year used combined pill

(R) (ND) **ydkpill** in last year used pill but not sure which

CONTRACEPTION

(U – see below) (ND) **ymirena** in last year used mirena
(U – see below) (q30br D) **ycoil** in last year used coil
(U – see below) (q30bd D) **ycondom** in last year used condom
(U – see below) (ND) **yfemidom** in last year used femidom
(U – see below) (q30bq D) **ycap** in last year used cap
(U – see below) (q30bl D) **ygels** in last year used gels/sprays/pessaries
(R) (q30bu D) **ysponge** in last year used contraceptive sponge
(R) (ND) **ypersona** in last year used persona
(U – see below) (q30bk D) **yrhythm** in last year used rhythm method/safe period
(U – see below) (q30bs D) **ywithdrw** in last year used withdrawal
(U – see below) (ND) **yinjectn** in last year used contraceptive injections
(U – see below) (ND) **yimplant** in last year used contraceptive implants
(U – see below) (ND) **yemercy** in last year used emergency contraception
(U – see below) (q30bc D) **yfsteril** in last year used female sterilisation
(U – see below) (q30bm D) **ymsteril** in last year used male sterilisation
(R) (q30bb D) **yabstain** in last year used abstinence
(U – see below) (q30bw D) **yothrpf** in last year used other contraception

Note: yconpill was derived in Natsal-2 but is obtained directly from UsedYr2 in Natsal-3 since only one pill option was given in Natsal-3.

(U – see below) (q30bj D) **yconpill** in last year used any contraceptive pill
1: yes if yminipil=1 | ycompill=1 | ydkpill=1
0: no else if yminipil=0 & ycompill=0 & ydkpill=0
9: not answered if missing
-1: not applicable if usedye01= -1

(U – see below) (q30bf D) **ynotused** in last year did not use any contraception
1: yes if usedye01=21
0: no else if usedye01>=1 & usedye08<=21
9: not answered if missing
-1: not applicable if usedye01= -1

Natsal-3 binary variables for contraception use (last year) are given below. Variables ymsteril..yconpatch are all coded in a similar way to yfsteril.

ynotused in last year did not use any contraception
-1: not applicable if afsex=90 | afsex=96 | afsex=97 | afsex=99
0: Used contr. in last year
1: Didn't use contr. in last year
9: not answered

yfsteril in last year used female sterilisation
-1: not applicable if afsex=90 | afsex=96 | afsex=97 | afsex=99
0: no or if fsterils=0
1: yes
9: not answered

ymsteril in last year used male sterilisation
yconpill in last year used contraceptive pill
ycondom in last year used condom
yfemidom in last year used femidom
(N) **yemcypill** in last year used morning after pill
(N) **yemcyiud** in last year used emergency IUD
ycoil in last year used coil
ymirena in last year used mirena
ycap in last year used cap
yinjectn in last year used contraceptive injections
ygels in last year used gels/sprays/pessaries

CONTRACEPTION

yrhythm in last year used rhythm method/safe period
ywithdrw in last year used withdrawal
yimplant in last year used contraceptive implants
yothrfp in last year used other contraception
yconptch in last year used contraceptive patch

(UD) **yemergency** in last year used emergency contraception
-1: not applicable if afsex=90 | afsex=96 | afsex=97 | afsex=99
0: no if yemcypill=0 & yemcyiud=0
1: yes if yemcypill=1 | yemcyiud=1
9: not answered

Note: Variable 'dblcon' was originally called 'double' but this is not allowed as a variable name in the statistical package stata so it was renamed 'dblcon'

(R) (N) **dblcon** used contraceptive methods in combination in last year
1: always used on different occasions
2: sometimes in combination on same occasion
3: always in combination on same occasion
9: not answered
-1: not applicable if everus01=21 | (usedye01=-1 | usedye01=21) | ((usedye01>=1 & usedye01<=21) & (usedye02=-1)) | (ycondom= 1 & at least one other method) | (yabstain=1 & at least one other method) | (yother= 1 & at least one other method)

Note: This question was asked of only some individuals in Natsal-3. For a full list of allowed combinations please see the Natsal-3 questionnaire.

(N) **dblcon2** used contraceptive methods in combination in last year
1: always used on different occasions
2: sometimes in combination on same occasion
3: always in combination on same occasion
9: not answered
-1: not applicable if afsex=90 | afsex=96 | afsex=97 | afsex=99 | (notused=1 | ynotused=1) | ((yfsteril=1 & any(1, yconpill, yemcypill, yemcyiud, ycap, yinjectn, ygels, yrhythm, ywithdrw, yimplant)) | (yconpill=1 & any(1, ycoil, ymirena, yinjectn, yimplant)) | (ycondom=1 & yfemidom=1) OR (yemcypill=1 & any(1, ycoil, yemcyiud)) | (ycoil=1 & any(1, ymirena, yinjectn, yimplant)) | (ymirena=1 & any(1, yinjectn, yimplant))) | (only one method used in last year)

*Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code questions. The following binary variables were obtained directly from the question **usualco2** in Natsal-3, and derived from **usualco1-usualco3** in Natsal-2. CAPI only allows interviewer to enter method of contraception that has been used in the last year, or no method.*

Note: in Natsal-2 respondents were routed into usuaco if they reported using more than one method of contraception in the last year. In Natsal-3, all individuals reporting 1 or more methods in the last year were routed into usualco2. The binary variable names have been changed to reflect this difference.

Natsal-2 binary variables for contraception use (usual method) are given below:

(R) (ND) **uminipil** usually use minipil
1: yes if (usualco1=-1 & usedye01=1) | usualco1=1 | usualco2=1 | usualco3=1
0: no else if (usualco1=-1 & (usedye01>=1 & usedye01<=21)) | (usualco1>=1 & usualco1<=22)
9: not answered if missing
-1: not applicable if yminipil = -1

CONTRACEPTION

Note: Variables ucompill ... uother are all derived from usedye01 (if only one method used in the last year) and usualco1 ...usualco2 (if 2+ methods used in the last year), and are coded in a similar way to uminipil:

- (R) (ND) **ucompill** usually use combined pill
- (R) (ND) **udkpill** usually use pill but not sure which
- (R) (ND) **umirena** usually use mirena
- (R) (ND) **ucoil** usually use coil
- (R) (ND) **ucondom** usually use condom
- (R) (ND) **ufemidom** usually use femidom
- (R) (ND) **ucap** usually use cap
- (R) (ND) **ugels** usually use gels/sprays/pessaries
- (R) (ND) **usponge** usually use contraceptive sponge
- (R) (ND) **upersona** usually use persona
- (R) (ND) **urhythm** usually use rhythm method/safe period
- (R) (ND) **uwithdrw** usually use withdrawal
- (R) (ND) **uinjectn** usually use contraceptive injections
- (R) (ND) **uimplant** usually use contraceptive implants
- (R) (ND) **uemergcy** usually use emergency contraception
- (R) (ND) **ufsteril** usually use female sterilisation
- (R) (ND) **umsteril** usually use male sterilisation
- (R) (ND) **uabstain** usually use abstinence
- (R) (ND) **uother** usually use other contraception

(R) (ND) **unone** not had a usual contraceptive method in the last year (incl. no method)

1: yes *if* usedye01=21 | usualco1=21 | usualco1=22 | usualco2=21 | usualco3=21

0: no *else if* (usedye01>=1 & usedye01<=20) | (usualco1>=1 & usualco1<=20)

9: not answered *else if missing*

-1: not applicable *if* yminipil=-1

(R) (ND) **uconpill** usually use a contraceptive pill

1: yes *if* uminipil=1 | ucompill=1 | udkpill=1

0: no *if* uminipil=0 & ucompill=0 & udkpill=0

9: not answered *if missing*

-1: not applicable *if* uminipil=-1

(R) (UD) **usnone** no contraception method at the moment

-1: not applicable *if* firstint=96 | firstint=97 | firstint=99 | nextage=96 | nextint=9 | nextage=97 | nextage=99

0: using contraception at the moment *if* unotused=0

1: not using contraception method at the moment *if* ynotused=1 | unotused=1

9: not answered

Natsal-3 binary variables are given below.

(N) **usfsteril** usually use female sterilisation

-1: not applicable *if* afsex=90 | afsex=96 | afsex=97 | afsex=99

0: no *or if* yfsteril=0

1: yes

9: not answered

And similarly...

(N) **usmsteril** usually use male sterilisation

(N) **usconpill** usually use contraceptive pill

(N) **uscondom** usually use condom

(N) **usfemidom** usually use femidom

(N) **usemcypill** usually use morning after pill

(N) **usemcyiud** usually use emergency IUD

(N) **uscoil** usually use coil

(N) **usmirena** usually use mirena

CONTRACEPTION

(N) **uscap** usually use cap
(N) **usinjectn** usually use contraceptive injections
(N) **usgels** usually use gels/sprays/pessaries
(N) **usrhythm** usually use rhythm method/safe period
(N) **uswithdrw** usually use withdrawal
(N) **usimplant** usually use contraceptive implants
(N) **usother** usually use other contraception
(N) **usconptch** usually use contraceptive patch
(N) **usnotused** usually uses no contr. method

(N) **uemergency** usually use emergency contraception
-1: not applicable *if* afsex=90 | afsex=96 | afsex=97 | afsex=99
0: no *if* usemcypill=0 & usemcyiud=0
1: yes *if* usemcypill=1 | usemcyiud=1
9: not answered

(R) (N) **whycond** why used condoms in last year
1: to prevent pregnancy
2: to protect against HIV/STDs
3: to prevent pregnancy and to protect against HIV/STDs
9: not answered
-1: not applicable *if* firstint=96 | nextage=96 | ycondom=0

Note: A similar question was asked in Natsal-3 question but the response options are different. This was a multi-code question so it has been converted into binary variables (all coded similarly to whycon21).

(N) **whycon21** why used condoms in last year: to prevent pregnancy
1: yes
0: no
9: not answered
-1: not applicable *if* afsex=90 | afsex=96 | afsex=97 | afsex=99 | ycondom=0

And similarly...

(N) **whycon22** why used condoms in last year: to protect against HIV
(N) **whycon23** why used condoms in last year: to protect against other STIs

Note: ifboth was removed in Natsal-3 since it was not used in any analysis in Natsal-2 and it was decided that it was sufficient to know the reasons for use

(R) (N) **ifboth** main reason for using condoms in last year
1: to prevent pregnancy
2: to protect against HIV/STDs
3: both equally important
4: depends on who the partner is
5: don't know
9: not answered
-1: not applicable *if* firstint=96 | nextage=96 | ycondom=0 | (whycond#3 & whycond#9)

Note: condreas will not be able to be derived in Natsal-3

(R) (ND) **condreas** main reason for using condoms in last year
compute condreas= whycond *if* whycond=1 | whycond=2
compute condreas= ifboth *if* whycond=3
1: to prevent pregnancy
2: to protect against hiv/stds
3: both equally important
4: depends on who the partner is
5: don't know
9: not answered *if* missing
-1: not applicable *if* firstint=96 | nextage=96 | ycondom=0

CONTRACEPTION

(R) (N) **fpsour01** source of contraception advice/supplies in last year 1

- 1: own GP
- 2: another GP at own surgery
- 3: FP clinic at own surgery
- 4: FP clinic, local health authority
- 5: sexual health/contraceptive service for young people
- 6: chemist/pharmacy
- 7: other type of shop
- 8: vending machine
- 9: through the post
- 10: emergency department at hospital
- 11: other type of service
- 12: none of these
- 99: not answered

Similarly for up to 8 sources with the additional code of -1 if fpsour=99 & fpsour-1≠99:

(R) (N) **fpsour02** source of contraception advice/supplies in last year 2

(R) (N) **fpsour03** source of contraception advice/supplies in last year 3

(R) (N) **fpsour04** source of contraception advice/supplies in last year 4

(R) (N) **fpsour05** source of contraception advice/supplies in last year 5

(R) (N) **fpsour06** source of contraception advice/supplies in last year 6

(R) (N) **fpsour07** source of contraception advice/supplies in last year 7

(R) (N) **fpsour08** source of contraception advice/supplies in last year 8

(R) (N) **fpsrcex1** other source of contraceptive advice/supplies in last year 1

- 1: own GP
- 2: another GP at own surgery
- 3: FP clinic at own surgery
- 4: FP clinic, local health authority
- 5: sexual health/contraceptive service for young people
- 6: chemist/pharmacy
- 7: other type of shop
- 8: vending machine
- 9: through the post
- 10: emergency department at hospital
- 11: other type of service
- 12: none of these
- 13: GUM/VD/urinary clinic
- 14: other doctor/hospital/(medical) clinic
- 15: partner provides (if he has them)
- 16: given by parent/relative/friend
- 17: retail outlet/pub (not already stated)
- 99: not answered
- 1: not applicable if fpsour≠11

Similarly for up to 2 other contraceptive sources:

(R) (N) **fpsrcex2** other source of contraceptive advice/supplies in last year 2

(R) (ND) **owngp** source of contraceptive advice/supplies in last year is own GP

- 1: yes if fpsour01=1 | fpsour02=1 | fpsour03=1 | fpsour04=1 | fpsour05=1 | fpsour06=1 | fpsour07=1 | fpsour08=1 | ((fpsour01=11 | fpsour02=11 | fpsour03=11 | fpsour04=11 | fpsour05=11 | fpsour06=11 | fpsour07=11 | fpsour08=11) & (fsrcex1=1 | fsrcex2=1))
- 0: no else if fpsour01>=1 & fpsour01<=12
- 9: not answered if missing

Note: variables anothrgp ... othrserv are all derived from fpsour01 ... fpsour11 and fsrcex1 ... fsrcex3, and are coded in a similar way to owngp:

(R) (ND) **anothrgp** source of contraceptive advice/supplies in last year is other GP at own surgery

(R) (ND) **surfgpcl** source of contraceptive advice/supplies in last year is FP clinic at own surgery

(R) (ND) **lhafpcl** source of contraceptive advice/supplies in last year is local health authority FP clinic

CONTRACEPTION

(R) (ND) **youngcl** source of contraceptive advice/supplies in last year is FP clinic for young people
(R) (ND) **chemist** source of contraceptive advice/supplies in last year is chemist/pharmacy
(R) (ND) **othrshop** source of contraceptive advice/supplies in last year is other type of shop
(R) (ND) **vendmach** source of contraceptive advice/supplies in last year is vending machine
(R) (ND) **thrupost** source of contraceptive advice/supplies in last year is through post
(R) (ND) **aehosptl** source of contraceptive advice/supplies in last year is hospital accident and emergency department
(R) (ND) **othrserv** source of contraceptive advice/supplies in last year is other type of service

(R) (ND) **nonthese** source of contraceptive advice/supplies in last year is none of these
1: yes *if* fpsour01=12
0: no *else if* fpsour01>=1 & fpsour01<=12
9: not answered *if* missing

(R) (ND) **anygp** source of contraceptive advice/supplies in last year is any GP
1: yes *if* (fpsour01=1 | fpsour02=1 | fpsour03=1 | fpsour04=1 | fpsour05=1 | fpsour06=1 | fpsour07=1 | fpsour08=1 | ((fpsour01=11 | fpsour02=11 | fpsour03=11 | fpsour04=11 | fpsour05=11 | fpsour06=11 | fpsour07=11 | fpsour08=11) & (fprcx1=1 | fprcx2=1))) | (fpsour01=2 | fpsour02=2 | fpsour03=2 | fpsour04=2 | fpsour05=2 | fpsour06=2 | fpsour07=2 | fpsour08=2 | ((fpsour01=11 | fpsour02=11 | fpsour03=11 | fpsour04=11 | fpsour05=11 | fpsour06=11 | fpsour07=11 | fpsour08=11) & (fprcx1=2 | fprcx2=2)))
0: no *else if* fpsour01>=1 & fpsour01<=12
9: not answered *if* missing

(R) (ND) **fpsourcg** source of contraceptive advice/supplies in last year, grouped
1: clinical *if* owngp=1 | anothrgp=1 | surgfpcl=1 | lhafpcl=1 | youngcl=1
2: retail only *if* (chemist=1 | othrshop=1 | vendmach=1 | thrupost=1) & (owngp=0 & anothrgp=0 & surgfpcl=0 & lhafpcl=0 & youngcl=0 & aehosptl=0 & othrserv=0 & nonthese=0)
3: other only *if* (othrserv=1 | aehosptl=1) & owngp=0 & anothrgp=0 & surgfpcl=0 & lhafpcl=0 & youngcl=0 & chemist=0 & othrshop=0 & vendmach=0 & thrupost=0 & nonthese=0
4: none of these *if* nonthese=1 & owngp=0 & anothrgp=0 & surgfpcl=0 & lhafpcl=0 & youngcl=0 & chemist=0 & othrshop=0 & vendmach=0 & thrupost=0 & aehosptl=0 & othrserv=0
5: 2+ source (excl. clinical) *else*
9: not answered *if* owngp=9

Note: question slightly revised from FPsource used in Natsal-2. This now specifies that we are interested in where contraception was obtained, ie no longer interested in advice.

Note: This codebook gives only the binary variables to be used for analysis, not the original multicode questions. Variables sownsurg.. sprtnrfp were obtained from the question fpsourc2 in the Natsal-3 questionnaire. Variables are all coded in a similar way to sownsurg.

(N) **sownsurg** source of contraceptive supplies in last year is dr/nurse at GPs surgery
-1: not applicable *if* ynotused#0
0: no
1: yes
9: not answered

(N) **sgumcl** source of contraceptive supplies in last year is GUM clinic
(N) **sfprhcl** source of contraceptive supplies in last year is FP/contraceptive/RH clinic
(N) **snhsancmw** source of contraceptive supplies in last year is NHS ANC/midwife
(N) **sprivdoc** source of contraceptive supplies in last year is private dr/clinic
(N) **syouncl** source of contraceptive supplies in last year is youth advisory clinic
(N) **schemist** source of contraceptive supplies in last year is pharmacy/chemist
(N) **swebsite** source of contraceptive supplies in last year is internet website
(N) **seduserv** source of contraceptive supplies in last year is school/college/university services
(N) **sothrshop** source of contraceptive supplies in last year is petrol station/supermarket/other shop
(N) **svendmach** source of contraceptive supplies in last year is vending machine
(N) **sthrupost** source of contraceptive supplies in last year is mail order
(N) **saehosptl** source of contraceptive supplies in last year is emergency dept at hospital

CONTRACEPTION

- (N) **sothrserv** source of contraceptive supplies in last year is other
(N) **snotgotfp** not got contraception in the last year
(N) **sfriendfp** source of contraceptive supplies in last year is friend
(N) **sparrelfp** source of contraceptive supplies in last year is parent/relative
(N) **sprtnerfp** source of contraceptive supplies in last year is partner

(ND) **fpsorcg** source of contraceptive supplies, grouped

-1: not applicable if ynotused#0

1: clinical if sownsurv=1 | sgumcl=1 | sfprhcl=1 | snhsancmw=1 | sprivdoc=1 | syoungcl=1

2: retail only if (schemist=1 | swebsite=1 | sothrshop=1 | svendmach=1 | sthrupost=1) & (sownsurv=0 & sgumcl=0 & sfprhcl=0 & snhsancmw=0 & sprivdoc=0 & syoungcl=0) & (seduser=0 & saehosptl=0 & sothrserv=0 & sfriendfp=0 & sparrelfp=0 & sprtnerfp=0 & SnotgotFP=0)

3: other only if (saehosptl=1 | sothrserv=1 | sfriendfp=1 | sparrelfp=1 | sprtnerfp=1) & (sownsurv=0 & sgumcl=0 & sfprhcl=0 & snhsancmw=0 & sprivdoc=0 & syoungcl=0 & schemist=0 & swebsite=0 & sothrshop=0 & svendmach=0 & sthrupost=0 & seduser=0 & SnotgotFP=0)

4: education only if seduser=1 & (sownsurv=0 & sgumcl=0 & sfprhcl=0 & snhsancmw=0 & sprivdoc=0 & syoungcl=0 & schemist=0 & swebsite=0 & sothrshop=0 & svendmach=0 & sthrupost=0 & (saehosptl=0 & sothrserv=0 & sfriendfp=0 & sparrelfp=0 & sprtnerfp=0 & SnotgotFP=0)

5: 2+ sources (excl. clinical) else

6: not got contraception supplies in the last year if snotgotfp=1

9: not answered

(R) (N) **fppref** preferred source of professional advice for contraception

1: own GP

2: another GP at own surgery

3: FP clinic at own surgery

4: FP clinic, local health authority

5: sexual health/contraceptive service for young people

6: chemist/pharmacy

7: none of these

9: not answered

Note: question revised from fppref used in Natsal-2 and now asks about getting contraception rather than professional advice for those who have used contraception in the last year. Response options have been updated.

(N) **fppref2** preferred source of contraception

-1: not applicable if ynotused#0

1: doctor / nurse at GPs surgery

2: GUM clinic

3: FP / contraceptive /reproductive health clinic

4: youth advisory clinic

5: pharmacy/chemist

6: NHS or DH website

7: None of these

8: Not needed

9: not answered

PERIODS AND MENOPAUSE

PERIODS AND MENOPAUSE

Note: there is an expanded section on periods and the menopause in Natsal-3 to collect information on HRT which reflects the higher upper age limit

Note: In Natsal-2 periods2 was asked in CASI and the corresponding question periods1 was asked of people not eligible for CASI (before the section on Attitudes). In Natsal-3 Periods2 was asked of all in the face-to-face interview.

(M) **periods2** age when started menstruating

(U) -1: not applicable if rsex=1 -1: not applicable if rsex=1 | scelig=2 | resultsc=1

(U) 97: not started menstruating

99: not answered

Note: Variable periods1 only applies to women not eligible for the self-completion questionnaire or women who only partially completed the self-completion questionnaire

(R) (N) **periods1** age started menstruating

99: not answered

-1: not applicable if rsex=1 | scelig=2 | resultsc=2 | resultsc=3

Note: periods no longer needed in Natsal-3 since all women answer periods2

(R) (s21w D) **periods** age started menstruating

99: not answered

-1: not applicable if rsex=1

Note: in Natsal-3 periodgp was derived from periods2, in Natsal-2 it was derived from periods

(ND) **periodgp** age started menstruating, grouped

1: <=10 years if periods2>0 & periods2<=10

2: 11-12 years if periods2>=11 & periods2<=12

3: 13-14 years if periods2>=13 & periods2<=14

4: >=15 years if periods2>=15 & periods2<=74

5: not started menstruating if periods2=97

9: not answered if periods2=99

-1: not applicable if periods2=-1

Note: in Natsal-3 period13 was derived from periods2

(ND) **period13** started menstruating before age 13

-1: not applicable if periods2=-1

1: <13 years if periods2>0 & periods2<13

2: 13 years if periods2=13

3: >13 years if periods2>13 & periods2<99

4: not started menstruating if periods2=97

9: not answered if periods2=99

(N) **periods3** first day of most recent period

-1: not applicable if rsex=1 or periods2=97

1: last 7 days

2: 7 days – 4 weeks

3: 4 weeks – 6 months

4: 6 months – 1 year

5: 1 year – 5 years

6: longer than 5 years

9: not answered

(N) **perlst** age at most recent period

-1: not applicable if periods3<=4 | periods3=9

99: not answered

PERIODS AND MENOPAUSE

(D) **menopause** menopausal status

-1: not applicable *if* periods3=-1

0: not menopausal *if* (rsex=2 & agrp<=3) | (rsex=2 & agrp>=4 & (periods3=1 | periods3=2 | periods3=3 | periods3=4))

1: menopausal *if* rsex=2 & agrp>=4 & (periods3=5 | periods3=6)

9: not answered *if* periods3=9

(N) **hrtever** ever taken HRT

-1: not applicable *if* periods3<=4 | periods3=9

1: yes

2: no

9: not answered

(N) **hrtstill** still on HRT

-1: not applicable *if* hrtever≠1

1: yes

2: no

9: not answered

SEXUAL ATTRACTION

SEXUAL ATTRACTION

(q31a) **attscale** sexual attraction

- 1: opposite sex only
- 2: more often opposite sex, and at least once with same sex
- 3: about equally often with opposite sex and same sex
- 4: more often same sex, and at least once with opposite sex
- 5: same sex only
- 6: never felt sexually attracted to anyone
- 7: refused
- 9: not answered

(q31b) **expscale** heterosexual/homosexual sexual experience

- 1: opposite sex only
- 2: more often opposite sex, and at least once with same sex
- 3: about equally often with opposite sex and same sex
- 4: more often same sex, and at least once with opposite sex
- 5: same sex only
- 6: never had sexual experience
- 7: refused
- 9: not answered

CASI ELIGIBILITY

SELF-COMPLETION ELIGIBILITY

Note: the variable het indicates whether respondent has had sexual intercourse aged 13+ based on the face-to-face (CAPI) question. See variable 'everhet' for a variable which combines answered from the CAPI with the CASI (self-completion)

(U) (D) **het** summary variable: had het. sex. intercourse aged 13+ (from CAPI questions)

(U) 1: (het. sex. intercourse aged 13+ if (afsex>=13 & afsex <=75) & ((expscale>=1 & expscale <=4) | expscale=7 | expscale=9)

1: het. sex. intercourse aged 13+ if (afsex>=13 & afsex <=44) & (expscale>=1 & expscale <=4 | expscale=7)

0: no het. sex. intercourse ever/since age 13 if (afsex=90 | afsex=96) | ((afsex=97 | afsex=99) & (expscale=5 | expscale=6))

(U) 9: unclassifiable if ((afsex>=13 & afsex<=75) & (expscale=5 | expscale=6)) | ((afsex=97|afsex=99) & (expscale>=1 & expscale <=4 | expscale=7 | expscale=9)

(U) **Sam** (D) **hom** summary variable: had hom. sex. experience ever (based on expscale)

(U) 1: same sex experience ever if (expscale>=2 & expscale <=5

1: hom. sex. experience ever if (expscale>=2 & expscale <=4) | ((afsex=90 | afsex=96 | afsex=97 | afsex=99) & (expscale=5))

(U) 0: no same sex. experience ever if (expscale=1 expscale=6)

0: no hom. sex. experience ever if expscale=1 | ((afsex=90 | afsex=96 | afsex=97 | afsex=99) & (expscale=6))

9: unclassifiable else if missing

Note: in Natsal-3 two variables were made to define eligibility according to i) the Natsal-3 rules (scelign3); and ii) the Natsal-2 rules (scelign2). The latter allows comparisons between the two surveys.

Notes: Variable 'scelig' is derived as in Natsal-1 using a temporary variable 'afexpsc' and dage

(R) (D) **scelig** eligible for self-completion questionnaire

if valid age at 1st het. sex. intercourse, code temporary variable afexpsc as:

(afsex>=13 & afsex<=44) & (expscale=1) afexpsc = 1

(afsex>=13 & afsex<=44) & (expscale=2) afexpsc = 5

(afsex>=13 & afsex<=44) & (expscale=3) afexpsc = 9

(afsex>=13 & afsex<=44) & (expscale=4) afexpsc = 13

(afsex>=13 & afsex<=44) & (expscale=5) afexpsc = 17

(afsex>=13 & afsex<=44) & (expscale=6) afexpsc = 21

(afsex>=13 & afsex<=44) & (expscale=7) afexpsc = 25

if age at 1st het. sex. intercourse was before age 13 and not since, code temporary variable afexpsc as:

(afsex=90) & (expscale=1) afexpsc = 2

(afsex=90) & (expscale=2) afexpsc = 6

(afsex=90) & (expscale=3) afexpsc = 10

(afsex=90) & (expscale=4) afexpsc = 14

(afsex=90) & (expscale=5) afexpsc = 18

(afsex=90) & (expscale=6) afexpsc = 22

(afsex=90) & (expscale=7) afexpsc = 26

if first sex has not happened yet, code temporary variable afexpsc as:

(afsex=96) & (expscale=1) afexpsc = 3

(afsex=96) & (expscale=2) afexpsc = 7

(afsex=96) & (expscale=3) afexpsc = 11

(afsex=96) & (expscale=4) afexpsc = 15

(afsex=96) & (expscale=5) afexpsc = 19

(afsex=96) & (expscale=6) afexpsc = 23

(afsex=96) & (expscale=7) afexpsc = 27

if age at 1st het. sex. intercourse refused, code temporary variable afexpsc as:

(afsex=97) & (expscale=1) afexpsc = 4

CASI ELIGIBILITY

(afsex=97) & (expscale=2) afexpsc = 8
(afsex=97) & (expscale=3) afexpsc = 12
(afsex=97) & (expscale=4) afexpsc = 16
(afsex=97) & (expscale=5) afexpsc = 20
(afsex=97) & (expscale=6) afexpsc = 24
(afsex=97) & (expscale=7) afexpsc = 28

1: eligible if (afexpsc=1 | afexpsc>=5 & afexpsc<=20 | afexpsc=21 | afexpsc=25) | (dage>=18 & afexpsc=2 | afexpsc=3 | afexpsc=4 | afexpsc=26 | afexpsc=27 | afexpsc=28)
2: not eligible if (afexpsc=22 | afexpsc=23 | afexpsc=24) | ((dage=16 | dage=17) & (afexpsc=2 | afexpsc=3 | afexpsc=4 | afexpsc=26 | afexpsc=27 | afexpsc=28))

Note: eligibility for the CASI changed in Natsal-3 and only those who had no sex aged 13+ and no sexual experience were not routed in. Some of these respondents (those who reported no het or same-sex experience within the CASI) were given a shortened set of questions. See the variable askcasi (page 59) for more information.

if valid age at 1st het. sex. intercourse, code temporary variable afexpsc as:

(afsex>=13 & afsex<=75) & (expscale=1) afexpsc = 1
(afsex>=13 & afsex<=75) & (expscale=2) afexpsc = 5
(afsex>=13 & afsex<=75) & (expscale=3) afexpsc = 9
(afsex>=13 & afsex<=75) & (expscale=4) afexpsc = 13
(afsex>=13 & afsex<=75) & (expscale=5) afexpsc = 17
(afsex>=13 & afsex<=75) & (expscale=6) afexpsc = 21
(afsex>=13 & afsex<=75) & (expscale=7 | expscale=9) afexpsc = 25

if age at 1st het. sex. intercourse was before age 13 and not since, code temporary variable afexpsc as:

(afsex=90) & (expscale=1) afexpsc = 2
(afsex=90) & (expscale=2) afexpsc = 6
(afsex=90) & (expscale=3) afexpsc = 10
(afsex=90) & (expscale=4) afexpsc = 14
(afsex=90) & (expscale=5) afexpsc = 18
(afsex=90) & (expscale=6) afexpsc = 22
(afsex=90) & (expscale=7) afexpsc = 26
(afsex=90) & (expscale=9) afexpsc = 29

if first sex has not happened yet, code temporary variable afexpsc as:

(afsex=96) & (expscale=1) afexpsc = 3
(afsex=96) & (expscale=2) afexpsc = 7
(afsex=96) & (expscale=3) afexpsc = 11
(afsex=96) & (expscale=4) afexpsc = 15
(afsex=96) & (expscale=5) afexpsc = 19
(afsex=96) & (expscale=6) afexpsc = 23
(afsex=96) & (expscale=7) afexpsc = 27
(afsex=96) & (expscale=9) afexpsc = 30

if age at 1st het. sex. intercourse refused/ not answered, code temporary variable afexpsc as:

(afsex=97|afsex=99) & (expscale=1) afexpsc = 4
(afsex=97|afsex=99) & (expscale=2) afexpsc = 8
(afsex=97|afsex=99) & (expscale=3) afexpsc = 12
(afsex=97|afsex=99) & (expscale=4) afexpsc = 16
(afsex=97|afsex=99) & (expscale=5) afexpsc = 20
(afsex=97|afsex=99) & (expscale=6) afexpsc = 24
(afsex=97|afsex=99) & (expscale=7) afexpsc = 28
(afsex=97|afsex=99) & (expscale=9) afexpsc = 31

scelign3 eligible for self-completion questionnaire in Natsal-3

afexpsc derived as above. Then:

1: eligible if (afexpsc>=1 & afexpsc<=20 | afexpsc=21 | afexpsc>=25 & afexpsc<=28)
2: not eligible if (afexpsc=22 | afexpsc=23 | afexpsc=24) | (afexpsc>=28 & afexpsc<=31)

CASI ELIGIBILITY

scelig2 eligible for self-completion questionnaire in Natsal-3 using Natsal-2 criteria

1: eligible *if* (afexpsc=1 | afexpsc>=5 & afexpsc<=20 | afexpsc=21 | afexpsc=25) | (dage>=18 & afexpsc=2 | afexpsc=3 | afexpsc=4 | afexpsc=26 | afexpsc=27 | afexpsc=28)
2: not eligible *if* (afexpsc=22 | afexpsc=23 | afexpsc=24) | ((dage=16 | dage=17) & (afexpsc=2 | afexpsc=3 | afexpsc=4 | (afexpsc>=26 & afexpsc<=31)))

(N) **scaccept** eligible respondents accepted the CASI

1: eligible respondent accepted the CASI
2: CASI asked face-to-face by interviewer
3: respondent refused CASI
9: not answered
-1: not applicable *if* not eligible for CASI (scelig=2 | sceligN3=2)

*Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code questions. The following binary variables were created directly from the question **xscacpt** (and any commonly-occurring other answers at **xxscaccept**) in Natsal-3, and derived from **xscace01-04** (and any commonly-occurring other answers at **xscacpt1-3**) in Natsal-2.*

Variables are all coded in a similar way to xsccomp.

xsccomp Reason for CASI refusal: didn't like computers

-1: not applicable *if* scaccept#3
0: no
1: yes
9: not answered

xsceyes Reason for CASI refusal: eyesight problems

xscread Reason for CASI refusal: could not read/write

xscdisab Reason for CASI refusal: other disability

xscobject Reason for CASI refusal: objected to subject

xscconfid Reason for CASI refusal: worried about confidentiality

xsclang Reason for CASI refusal: language problems

xscrttime Reason for CASI refusal: ran out of time

xscboth Reason for CASI refusal: couldn't be bothered

xscoth Reason for CASI refusal: other reason

Note: in Natsal-3 there was no option to complete a paper version of the self-completion section

(R) (N) **offpapi** whether eligible respondents who refused CASI accepted paper version

1: yes, willing to accept paper version of CASI
2: no, not willing to accept paper version of CASI
9: not answered
-1: not eligible for self-completion/accepted CASI *if* scelig=2 | scaccept=1 | scaccept=2 | (scaccept=3 & ((xscace01=1 | xscace01=10) & (xscace02=-1))

(U) (N) **resultsc** eligible respondents accepted CASI

1: CASI only partially completed
2: CASI completed with no help/advice
3: CASI completed with some help/advice
(U) 4: CASI asked face to face by interviewer
-1: not applicable *if* scelig=2

Note: in Natsal-3 reasons for asking the CASI questions face-to-face were collected. The most common of these were coded into binary variables scf2f1..scf2f7

(N) **scf2f1** CASI asked face-to-face because didn't like computers

(N) **scf2f2** CASI asked face-to-face because of eyesight problems

(N) **scf2f3** CASI asked face-to-face because could not read/write

CASI ELIGIBILITY

- (N) **scf2f4** CASI asked face-to-face because of other disability
- (N) **scf2f5** CASI asked face-to-face because of language problems
- (N) **scf2f6** CASI asked face-to-face because asked interviewer to/not fussed/happy to talk about sex openly
- (N) **scf2f7** CASI asked face-to-face because attempted computer but preferred face-to-face with interviewer

Note: In Natsal-3 some respondents who were eligible for the CASI were given a shortened set of questions. These were people who reported that they had not had heterosexual intercourse or same sex experience since age 13. The shortened CASI questionnaire was designed to ensure that respondents were not asked irrelevant questions. The variable askcasi indicates whether the respondent was asked the full or shortened CASI questionnaire. This variable is used in the filtering of questions throughout the CASI as indicated in this codebook. For a list of questions / question modules included in the shortened CASI please see the Natsal-3 questionnaire.

- (N) **askcasi** ask all CASI questions?
 - 1: all CASI questions
 - 2: shortened CASI
 - 1: not applicable if scelign3=2 | scaccept=3

HETEROSEXUAL BEHAVIOUR

The following topics are all within the self-completion section of the Natsal-3 and Natsal-2 questionnaires, which, unlike Natsal-1 was completed by the respondent using a laptop.

HETEROSEXUAL BEHAVIOUR

Note: In Natsal-3 a check question was added to the start of the CASI to confirm whether those who did not report having had sex with an opposite sex partner in the CAPI had actually had sex according to the Natsal-3 definitions.

(N) **hetcheck** ever had het sex.

-1: not applicable if (dmdebut>=13 & dmdebut<=74) | scelig=2 | scaccept=3

1: yes

2: no

9: not answered

(ND) **everhet** summary variable: had het sex. intercourse aged 13+ (using information from CAPI and CASI)

1: het. sex. intercourse aged 13+ if het=1 | hetcheck=1

0: no het. sex. intercourse ever/since age 13 if (het=0 & (hetcheck=-1 | hetcheck=2)) | (het=9 & hetcheck=2)

9: unclassifiable if het=9 & (hetcheck=-1 | hetcheck=9)

Note: in Natsal-3 the sexual practices questions were moved earlier in the questionnaire to improve questionnaire flow. Previously positioned after sex7day.

(M) (s3a) **lastvag** last occasion of vaginal sex

1: in last 7 days

2: between 7 days and 4 weeks

3: between 4 weeks and 6 months

4: between 6 months and 1 year

5: between 1 year and 5 years

6: longer than 5 years

(U) 7: never or if missing & everhet=0 7: never or if missing & (het=0 | hetsc=0)

9: not answered

(U) -1 not applicable if scelign3=-2 -1: not applicable else if scelig=2

Note: variables oralyou ... analsex have the same codes as lastvag

(M) (s3b) **oralyou** last occasion of het. oral sex to partner

(M) (s3c) **oralprt** last occasion of het. oral sex from partner

(M) (s3d) **analsex** last occasion of het. anal sex

Note: Natsal-2 uses slightly different recoding rules to Natsal-1

(M) (R) (s3e) **gencont** last occasion of het. genital contact not involving intercourse

1: in last 7 days

2: between 7 days and 4 weeks

3: between 4 weeks and 6 months ago

4: between 6 months and 1 year ago

5: between 1 year and 5 years ago

6: longer than 5 years ago

7: never or if missing & (expscale=5 & afsex=96 | afsex=97 | afsex=99) | (expscale=6 & afsex=90 | afsex=96 | afsex=97 | afsex=99)

9: not answered

-1: not applicable else if scelig=2

HETEROSEXUAL BEHAVIOUR

Note: gencont2 is the same question as previously but was asked of everyone, even those reporting not having had sex

(N) (s3e) **gencont2** last occasion of het. genital contact not involving intercourse

1: in last 7 days

2: between 7 days and 4 weeks

3: between 4 weeks and 6 months ago

4: between 6 months and 1 year ago

5: between 1 year and 5 years ago

6: longer than 5 years ago

7: never or if missing & scalign3=2 | (expscale=5 & afsex=96 | afsex=97 | afsex=99) | (expscale=6 & afsex=90 | afsex=96 | afsex=97 | afsex=99)

9: not answered

(ND) **vagsxevr** ever had vaginal sex

1: yes if lastvag>=1 & lastvag<=6

0: no if lastvag=7

9: not answered if lastvag=9

-1: not applicable if lastvag=-1

(ND) **vagsxl5y** had vaginal sex in last 5 years

1: yes if lastvag>=1 & lastvag<=5

0: no if lastvag=6 | lastvag=7

9: not answered if lastvag=9

-1: not applicable if lastvag=-1

(ND) **vagsexyr** had vaginal sex in last year

1: yes if lastvag>=1 & lastvag<=4

0: no if lastvag=5 | lastvag=6 | lastvag=7

9: not answered if lastvag=9

-1: not applicable if lastvag=-1

(ND) **vagsexlm** had vaginal sex in last month

1: yes if lastvag=1 | lastvag=2

0: no if lastvag>=3 & lastvag<=7

9: not answered if lastvag=9

-1: not applicable if lastvag=-1

(ND) **ori2hevr** ever given het. oral sex

1: yes if oralyou>=1 & oralyou<=6

0: no if oralyou=7

9: not answered if oralyou=9

-1: not applicable if oralyou=-1

(ND) **ori2hl5y** gave het. oral sex in last 5 years

1: yes if oralyou>=1 & oralyou<=5

0: no if oralyou=6 | oralyou=7

9: not answered if oralyou=9

-1: not applicable if oralyou=-1

(ND) **ori2hyr** gave het. oral sex in last year

1: yes if oralyou>=1 & oralyou<=4

0: no if oralyou=5 | oralyou=6 | oralyou=7

9: not answered if oralyou=9

-1: not applicable if oralyou=-1

HETEROSEXUAL BEHAVIOUR

(ND) **orl2hlm** gave het. oral sex in last month

1: yes *if* oralyou=1 | oralyou=2
0: no *if* oralyou>=3 & oralyou<=7
9: not answered *if* oralyou =9
-1: not applicable *if* oralyou=-1

(ND) **orlfhevr** ever received het. oral sex

1: yes *if* oralprt>=1 & oralprt<=6
0: no *if* oralprt=7
9: not answered *if* oralprt=9
-1: not applicable *if* oralprt=-1

(ND) **orlfhl5y** received het. oral sex in last 5 years

1: yes *if* oralprt>=1 & oralprt<=5
0: no *if* oralprt=6 | oralprt=7
9: not answered *if* oralprt=9
-1: not applicable *if* oralprt=-1

(ND) **orlfhyr** received het. oral sex in last year

1: yes *if* oralprt>=1 & oralprt<=4
0: no *if* oralprt=5 | oralprt=6 | oralprt=7
9: not answered *if* oralprt=9
-1: not applicable *if* oralprt =-1

(ND) **orlfhlm** received het. oral sex in last month

1: yes *if* oralprt=1 | oralprt=2
0: no *if* oralprt>=3 & oralprt<=7
9: not answered *if* oralprt=9
-1: not applicable *if* oralprt =-1

(ND) **anyorevr** ever given/received het. oral sex

1: yes *if* (oralyou>=1 & oralyou<=6) | (oralprt>=1 & oralprt<=6)
0: no *else if* oralyou=7 | oralprt=7
9: not answered
(U) -1 not applicable *if* scelign3=-2 -1: not applicable *else if* scelig=2

(ND) **anyorl5y** gave/received het. oral sex in the last 5 years

1: yes *if* (oralyou>=1 & oralyou<=5) | (oralprt>=1 & oralprt<=5)
0: no *else if* (oralyou=6 | oralyou=7) | (oralprt=6 | oralprt=7)
9: not answered
(U) -1 not applicable *if* scelign3=-2 -1: not applicable *else if* scelig=2

(ND) **anyoral** gave/received het. oral sex in the last year

1: yes *if* (oralyou>=1 & oralyou<=4) | (oralprt>=1 & oralprt<=4)
0: no *else if* (oralyou>=5 & oralyou<=7) | (oralprt>=5 & oralprt<=7)
9: not answered
(U) -1 not applicable *if* scelign3=-2 -1: not applicable *else if* scelig=2

(ND) **anyoralm** gave/received het. oral sex in the last month

1: yes *if* (oralyou=1 | oralyou=2) | (oralprt=1 | oralprt=2)
0: no *else if* (oralyou>=3 & oralyou<=7) | (oralprt>=3 & oralprt<=7)
9: not answered
(U) -1 not applicable *if* scelign3=-2 -1: not applicable *else if* scelig=2

(ND) **analhevr** ever had het. anal sex

1: yes *if* analsex>=1 & analsex<=6
0: no *if* analsex=7
9: not answered *if* analsex=9
-1: not applicable *if* analsex=-1

(ND) **analh5yr** had het. anal sex in last 5 years

- 1: yes *if* analsex>=1 & analsex<=5
- 0: no *if* analsex>=6 & analsex<=7
- 9: not answered *if* analsex=9
- 1: not applicable *if* analsex=-1

(ND) **analhyr** had het. anal sex in last year

- 1: yes *if* analsex>=1 & analsex<=4
- 0: no *if* analsex>=5 & analsex<=7
- 9: not answered *if* analsex=9
- 1: not applicable *if* analsex=-1

(ND) **analhlm** had het. anal sex in last month

- 1: yes *if* analsex=1 | analsex=2
- 0: no *if* analsex>=3 & analsex<=7
- 9: not answered *if* analsex=9
- 1: not applicable *if* analsex=-1

Note in Natsal-3 the variables genhevr..genhlm are based on gencont2, not gencont.

(ND) **genhevr** ever had het. genital contact not involving intercourse

- 1: yes *if* gencont2>=1 & gencont2<=6
- 0: no *if* gencont2=7
- 9: not answered *if* gencont2=9
- 1: not applicable *if* gencont2=-1

(ND) **genhl5yr** had het. genital contact not involving intercourse in last 5 years

- 1: yes *if* gencont2>=1 & gencont2<=5
- 0: no *if* gencont2=6 | gencont2=7
- 9: not answered *if* gencont2=9
- 1: not applicable *if* gencont2=-1

(ND) **genhlyr** had het. genital contact not involving intercourse in last year

- 1: yes *if* gencont2>=1 & gencont2<=4
- 0: no *if* gencont2=5 | gencont2=6 | gencont2=7
- 9: not answered *if* gencont2=9
- 1: not applicable *if* gencont2=-1

(ND) **genhlm** had het. genital contact not involving intercourse in last month

- 1: yes *if* gencont2=1 | gencont2=2
- 0: no *if* gencont2>=3 & gencont2<=7
- 9: not answered *if* gencont2=9
- 1: not applicable *if* gencont2=-1

HETEROSEXUAL BEHAVIOUR

Note:

1. In Natsal-3 routing into sex4wks was dependent on answers to lastvag, oralyou, oralprt and analsex. If any in the last 4 weeks it was asked. This affects no4wks, n1in4wk and new4wks.
2. In Natsal-3 there are a small number (n=40) of cases with sex4wks=0 but lastvag=1 or 2 (i.e. in the last 4 weeks). These cases have been left inconsistent in the data, but data users may want to consider recoding sex4wks to 99 'not answered' for these cases, as has been done in papers by the Natsal team using this variable (e.g. Mercer et al 2013. Changes in Sexual attitudes and lifestyles in Britain through the life course and over time: findings from the National Surveys of Sexual Attitudes and Lifestyles (Natsal). The Lancet: 382; 1781-1794).

(ND) **sex4wks** no. of occasions of het. sex in last 4 weeks

0 if (everhet=1 | everhet=9) & (lastvag>=3 & lastvag<=7) & (oralyou>=3 & oralyou<=7) & (oralprt>=3 & oralprt<=7) & (analsex>=3 & analsex<=7)

995: at least once

996: not had vaginal sex in last 4 wks but other het. practices if (lastvag>=3 & lastvag<=9) & (oralyou<=2 | oralprt<=2 | analsex<=2))

997: not had het. sex yet if everhet=0

999: not answered

-1: not applicable else if scelign3=2

Note: Variable 'sex4wks1' is coded in the same way as the Natsal-1 equivalent 'rs2a' and refers to all respondents eligible for the self-completion section regardless of whether ever had heterosexual sex

(U)(rs2a) **sex4wks1** no. of occasions of het. sex in last 4 weeks (regardless of whether ever had het. sex)

(U) 0 if sex4wks=997 0 if het=0 | (sex4wks2=996 | sex4wks2=997)

(U) 995: at least one if sex4wks=995 | sex4wks=996

(U) 999: not answered or if sex4wks=999 999: not answered or if sex4wks=998

(U) -1: not applicable if scelign3=2 -1: not applicable else if scelig=2

Note: Variable 'sex4wks2' makes use of the new code 997 introduced in Natsal-2 that is assigned to respondents who have not had het. sex

(U) (N) **sex4wks2** no. of occasions of het. sex in last 4 weeks

996: not had vaginal sex in last 4 wks but other het. practices if sex4wks=0 & (lastvag>=3 & lastvag<=9) & (oralyou<=2 | oralprt<=2 | analsex<=2)

997: not had het. sex yet or if missing & het=0

999: not answered or if sex4wks=998

(U) -1 not applicable if scelign3=2 -1: not applicable else if scelig=2

(ND) **sexfreq** no. of occasions of het. sex in the last 4 weeks

1: 0-2 times if sex4wks1>=0 & sex4wks1<=2

2: 3-4 times if sex4wks1>=3 & sex4wks1<=4

3: 5+ times if sex4wks1>=5 & sex4wks1<=994

7: at least once if sex4wks1=995

9: not answered if sex4wks1=999

Note: due to changes to routing hetpsc was removed in Natsal-3. The variable everhet now indicates whether the respondent has had heterosexual sex.

Note: Variable 'hetpsc' considers het. sex as vaginal, oral, and/or anal sex, which is why if 'sex4wks2'=996 then the respondent is still considered as having had het. sex. This issue did not exist for Natsal-1 as 'rs2a', the equivalent of 'sex4wks1', did not have code '996'.

(R) (ND) **hetpsc** summary: had het. sex. intercourse (according to self-completion)

1: had het. sex at least once if sex4wks2>=0 & sex4wks2<997

0: no het. sex if sex4wks2=997

9: not answered unless het=0 | 1 in which case copy het

HETEROSEXUAL BEHAVIOUR

Note: est4wks was not used in analysis and was removed from Natsal-3

(R) (N) **est4wks** certainty of no. of occasions of het. sex in last 4 weeks
1: certain
2: estimate
9: not answered
-1: not applicable *if sex4wks2=0 | sex4wks2=-1 | sex4wks2=997*

Note: routing into no4wks was slightly different in Natsal-3 due to routing change for sex4wks.

Note: Variable no4wks recodes those coded 996 for sex4wks2 to 0 to be consistent with Natsal-1's use of variable 'rs2a' in recoding 'rs2b'

(U) (rs2b) **no4wks** no. of het. sex partners in last 4 weeks
(U) 0 *if (everhet=0 | sex4wks1=0)* 0 *if het=0 | hetsc=0 | sex4wks1=0*
995: at least 1 partner
(R) 996: at least 2 partners
(R) 997: at least 3 partners
999: not answered
(U) -1: not applicable *else if scelgn3=2* -1: not applicable *else if scelig=2*

Note: n1in4wk is used in the derivation of new4wks then dropped from final dataset. New4wks should be used for analysis.

(R) (N) **n1in4wk** last het. sex partner was a new partner (if had 1 het. partner in last 4 weeks)
1: yes
2: no
9: not answered
-1: not applicable *if no4wks≠1 & no4wks≠999*

(rs2c) **new4wks** no. of new het. sex partners in last 4 weeks
0 *if missing & no4wks=1 & n1in4wk=2*
1 *if missing & no4wks=1 & n1in4wk=1*
999: not answered
(U) -1: not applicable *if (everhet=0 | sex4wks=1)* -1: not applicable *if het=0 | hetsc=0 | sex4wks=0*

Note: Routing into cond4wk was different in Natsal-3. Respondents were only asked this question if they reported either vaginal or anal sex in the last 4 weeks at lastvag and analsex

(U) (s2d *U) **cond4wk** condom use for het. sex in last 4 weeks
1: yes, used on every occasion
2: yes, used on some occasions
3: not used in last 4 weeks
(U - answer option removed) 4: not had vaginal/anal sex in last 4 weeks
9: not answered
(U) -1: not applicable *if (everhet=1 & NOT (lastvag=1 | lastvag=2 | analsex=1 | analsex=2)) or else if (everhet=0 | scelign3=2)*
-1: not applicable *if het=0 | hetsc=0 | scelig=2*

Note: Routing into sex7days was different was Natsal-3. Respondents were only asked this question if they reported that one of lastvag, analsex, oralprt or oralyou was in the last 7 days

(U) **sex7day** no. of occasions of het. sex in last 7 days
(U) 0 *if (everhet=0 | sex4wks1=1) or if NOT (lastvag=1 | oralyou=1 | oralprt=1 | analsex=1)*
0 *if het=0 | hetsc=0 | sex4wks2=0*
95: not had vaginal sex in last 7 days but other het. practices *if (lastvag≠1 & (oralyou=1 | oralprt=1 | analsex=1))*
(U – new code) 995: at least once
(U) 999: not answered 99: not answered
(U) -1: not applicable *if (scelign3=2)* -1: not applicable *else if scelig=2*

HETEROSEXUAL BEHAVIOUR

Note: lastvag, oralyou, oralprt and analsex were here in Natsal-2 but have been moved earlier in the questionnaire

(N) **yrcond** had vaginal/ het. anal sex without a condom in last year

1: yes, without a condom

2: no, always with a condom

3: was code 2 but other data suggests code 1

(U) 7: no vaginal/het. anal sex in last year *or if missing & everhet=0* | ((lastvag=5 | lastvag=6 | lastvag=7) & (analsex=5 | analsex=6 | analsex=7))

7: no vaginal/het. anal sex in last year *or if missing & ((het=0 | hetsc=0) | ((lastvag=5 | lastvag=6 | lastvag=7) & (analsex=5 | analsex=6 | analsex=7))*

9: not answered

(U) -1: not applicable *if (scelign3=2 | hetcheck=2)* -1: not applicable *else if scelig=2*

(ND) **yrcond2** had vaginal/ het. anal sex with a condom in last year, grouped

0: no vaginal/het. anal sex in last year *if yrcond=7*

1: yes, vaginal/het. anal sex without a condom *if yrcond=1 | yrcond=3*

2: no, vaginal/het. anal sex always with a condom *if yrcond=2*

9: not answered *if yrcond=9*

-1: not applicable *if yrcond=-1*

(N) **nonocon** no. of vaginal/het. anal partners in last year without condom

0: no unprotected vaginal/het. anal sex in last year *if (yrcond=2 | yrcond=7)*

996: was code 2 for 'yrcond' but other data suggests code 1 *if yrcond=3*

999: not answered

(U) -1: not applicable *if (scelign3=2 | hetcheck=2)* -1: not applicable *else if scelig=2*

(ND) **nonocong** no. of vaginal/ het. anal partners in last year without condom, grouped

0: no unprotected vaginal/het. anal sex in last year *if nonocon=0*

1: 1 *if nonocon=1 | nonocon=996*

2: 2+ *if nonocon>=2 & nonocon<996*

9: not answered *if nonocon=999*

-1: not applicable *else if nonocon=-1*

(N) **masturb** last occasion of masturbation

1: in last 7 days

2: between 7 days and 4 weeks

3: between 4 weeks and 6 months

4: between 6 months and 1 year

5: between 1 year and 5 years

6: longer than 5 years

7: never

9: not answered

(U) -1: not applicable *if (scelign3=2 | everhet=0)* -1: not applicable *else if scelig=2*

(ND) **mastbevr** ever masturbated

1: yes *if masturb>=1 & masturb<=6*

0: no *if masturb=7*

9: not answered *if masturb=9*

-1: not applicable *if masturb=-1*

(ND) **mastbl5y** masturbated in last 5 years

1: yes *if masturb>=1 & masturb<=5*

0: no *if masturb>=6 & masturb<=7*

9: not answered *if masturb=9*

-1: not applicable *if masturb=-1*

HETEROSEXUAL BEHAVIOUR

(ND) **mastbyr** masturbated in the last year

1: yes *if* masturb $\geq$ 1 & masturb $\leq$ 4

0: no *if* masturb $\geq$ 5 & masturb $\leq$ 7

9: not answered *if* masturb=9

-1: not applicable *if* masturb=-1

(ND) **mastblm** masturbated in the last month

1: yes *if* masturb=1 | masturb=2

0: no *if* masturb $\geq$ 3 & masturb $\leq$ 7

9: not answered *if* masturb=9

-1: not applicable *if* masturb=-1

Note: question mastnum added in Natsal-3 to look at frequency of masturbation

(N) **mastnum** no. times masturbated in last week

0 *if* masturb $\geq$ 2 & masturb $\leq$ 7

999: not answered

-1: not applicable *if* masturb=-1

SAME-SEX BEHAVIOUR

SAME SEX BEHAVIOUR

Note: in Natsal-3, all variables in this section use the letters 'sam' instead of 'hom' so as not to cause offense to respondents as the variable names are visible on the CASI screen. All coding for deriving variables in this section has been updated to 'sam' for consistency

eversam (s4a) **everhom** ever had any hom. sex. experience/contact

1: yes *or if missing & sam=1*

2: no *or if missing & sam=0*

9: not answered

(U - code removed) -1: not applicable *else if scelig=2*

frstsam (rs4b) **frsthom** age at first hom. sex. contact/experience

99: not answered

(U) -1: not applicable *if eversam=2* -1: not applicable *if scelig=2 | eversam=2*

genisam (s4c) **genihom** ever had hom. sex. experience with genital contact

1: yes

2: no *or if missing & (sam=0 | eversam=2)*

9: not answered

(U - code removed) -1: not applicable *else if scelig=2*

agegsam (N) **ageghom** age at 1st hom. sex. experience with genital contact

99: not answered

(U) -1: not applicable *if sam=0 | eversam=2 | genisam=2*

-1: not applicable *if scelig=2 | sam=0 | eversam=2 | genisam=2*

(N) **nextageg** had hom. sex. aged 13+ if 1st hom. sex age<13

1: yes

2: no with anybody else since age 13

9: not answered

(U) -1: not applicable *if sam=0 | genisam=2 | (genisam=1 & agegsam>12 & agegsam<99)*

-1: not applicable *if scelig=2 | sam=0 | genisam=2 | (genisam=1 & agegsam>12 & agegsam<99)*

Note: nextasam was added in Natsal-3 to collect age at next same sex if 13+. This is consistent with the opposite sex questions.

(N) **nextasam** age at next hom. sex. intercourse if 1st sex age<13

95: unable to estimate

96: not with anyone since age 13 *if nextageg=2*

99: not answered

(U) -1: not applicable *if sam=0 | eversam=2 | genisam=2 | (agegsam>12 & agegsam<99) |*

Note: variable 'afsam13' is coded in the same way as the Natsal-2 equivalent 'afhom13' and does not make use of the data from the new variable 'nextasam'.

(U) **afsam13** (ND) **afhom13** age 1st same sex. experience if aged 13+

copy agegsam *if agegsam>12 & agegsam<45*

98: had hom. sex. with genital contact <13 & 13+ but unknown age *if agegsam<13 & nextageg=1*

99: not answered *if genisam=1 & agegsam=99 or missing*

-1: not applicable *if scelig=2 | (ageghom<13 & nextageg=2) | genihom=2*

(U) -1: not applicable *if genisam=2 | (agegsam<13 & nextageg=2)*

SAME-SEX BEHAVIOUR

Note: Variable 'afsam13a' makes use of the data from the new variable 'nextasam' introduced in Natsal-3 which asks the age of first homosexual sex after age 13. Note for this reason this variable is not comparable with the Natsal-2 variable afhom13.

(ND) **afsam13a** age 1st hom. sex if aged 13+ including all
copy agegsam if agegsam>12 & agegsam<75
copy nextasam if agegsam>0 & agegsam<13 & nextageg=1
90: age <13 and not since if agegsam>0 & agegsam<13 & nextageg=2
95: unable to estimate
99: not answered if genisam=1 & agegsam=99 or missing
-1: not applicable if genisam=2

(ND) **totafsex** age at first sex (het. intercourse or sam. experience with genital contact)
copy afsex if (afsex<90 & (afsam13a=-1 | afsam13a=90 | (afsam13a>-1 & afsam13a<90 & afsam13a>afsex))) | (afsex<90 & afsam13a<90 & afsam13a>=13 & afsex=afsam13a)
copyafsam13a if afsam13a>=13 & afsam13a<90 & (afsex=90 | afsex=96 | (afsex<90 & afsex>afsam13a))
90: age <13 and not since else if (afsex=90 | afsam13a=90) & afsex#99 & afsam13a#99
96: has not happened yet else if afsex=96 & afsam13a=-1
99: not answered else if afsex=99 | afsam13a=99

(ND) **totafsexgp** grouped age at first het./sam. sex
1: 13-15 if totafsex>=13 & totafsex<=15
2: 16-17 if totafsex=16 | totafsex=17
3: 18-19 if totafsex=18 | totafsex=19
4: 20+ if totafsex>=20 & totafsex<90
96: hasn't happened yet/since age 13 if totafsex=90 | totafsex=96
99: not answered if totafsex=99

(ND) **samsex** ever had same sex sex aged13+
0: no if (afsam13a=-1 & (genisam=2 | agegsam>0 & agegsam<13 & nextageg=2)) | afsam13a=90
1: yes if (afsam13a >=13 & afsam13a <=90) | afsam13a=95 | afsam13a=98
9: missing if genisam=9
-1: not applicable if scelig=2 & eversam=2

Note: samoyou, samopar, samayou, samahim and samgen were after hom7day in Natsal-2 and have been moved for Natsal-3. This is consistent with the changes made to the heterosexual sex section.

(U) (M) **samoyou** (s6a) **homoyou** last occasion of hom. oral sex to partner
1: in last 7 days
2: between 7 days and 4 weeks
3: between 4 weeks and 6 months ago
4: between 6 months and 1 year ago
5: between 1 year and 5 years ago
6: longer than 5 years ago
(U) 7: never or if sam=0 | eversam=2 | genisam=2 | (agegsam>0 & agegsam<13 & nextageg=2)
7: never or if sam=0 | eversam=2 | genisam=2
9: not answered
(U – code removed) -1: not applicable else if scelig=2

Note: Variable homopar has the same codes as homoyou:

(M) **samopar** (s6b) **homopar** last occasion of hom. oral sex from partner

SAME-SEX BEHAVIOUR

(M) **samayou** (s6c) **homayou** last occasion of hom. anal sex to partner

1: in last 7 days

2: between 7 days and 4 weeks

3: between 4 weeks and 6 months ago

4: between 6 months and 1 year ago

5: between 1 year and 5 years ago

6: longer than 5 years ago

(U) 7: never or if rsex=1 & (sam=0 | eversam=2 | genisam=2 | (agegsam>0 & agegsam<13 & nextageg=2))

7: never or if sam=0 | eversam=2 | genisam=2

9: not answered

(U) -1: not applicable if rsex=2 -1: not applicable else if scelig=2 | rsex=2

Note: Variable homahim has the same codes as homayou:

(M) **samahim** (s6d) **homahim** last occasion of hom. anal sex from partner

(ND) **anlhmevr** ever had hom. anal sex

0: no if samayou=7 & samahim=7

1: yes if (samayou>=1 & samayou<=6) | (samahim>=1 & samahim<=6)

9: not answered if samayou=9 | samahim=9

-1: not applicable if samayou=-1 | samahim=-1

(ND) **anlhml5y** had hom. anal sex in last 5 years

0: no if (samayou=6 | samayou=7) & (samahim=6 | samahim=7)

1: yes if (samayou>=1 & samayou<=5) | (samahim>=1 & samahim<=5)

9: not answered if samayou=9 | samahim=9

-1: not applicable if samayou=-1 | samahim=-1

(ND) **anlhmlyr** had hom. anal sex in last year

0: no if (samayou=5 | samayou=6 | samayou=7) & (samahim=5 | samahim=6 | samahim=7)

1: yes if (samayou>=1 & samayou<=4) | (samahim>=1 & samahim<=4)

9: not answered if samayou=9 | samahim=9

-1: not applicable if samayou=-1 | samahim=-1

(ND) **anlhmlm** had hom. anal sex in last month

0: no if (samayou>=3 & samayou<=7) & (samahim>=3 & samahim<=7)

1: yes if (samayou=1 | samayou=2) | (samahim=1 | samahim=2)

9: not answered if samayou=9 | samahim=9

-1: not applicable if samayou=-1 | samahim=-1

Note: in Natsal-3 routing into samgen was different and people routed in even if nextageg=2.

Note: Variable homgen has the same codes as homoyou:

(M) **samgen** s6e **homgen** last occasion of hom. genital contact not involving oral/anal sex

1: in last 7 days

2: between 7 days and 4 weeks

3: between 4 weeks and 6 months ago

4: between 6 months and 1 year ago

5: between 1 year and 5 years ago

6: longer than 5 years ago

7: never or if sam=0 | eversam=2 | genisam=2

9: not answered

(U – code removed) -1: not applicable else if scelig=2

SAME-SEX BEHAVIOUR

Note: in Natsal-3 eligibility for sam4wks changed and was based on one of the practices questions (samoyou, samopar, samayou, samahim) having been in the last 4 weeks. This also affects the routing into samno4w, h1new4w, samnw4w

(U) **sam4wks** (rs5a) **hom4wks** no. of occasions of hom. sex in last 4 weeks

(U) 0 if missing & (sam=0 | eversam=2 | genisam=2 | (agegsam>0 & agegsam<13 & nextageg=2) | (any (samoyou, -1, 2, 4, 5, 6, 7) | any (samopar, -1, 2, 4, 5, 6, 7) | any (samayou, -1, 2, 4, 5, 6, 7) | any (samahim, -1, 2, 4, 5, 6, 7))

(U – new code) 995: at least one

(U) 999: not answered 99: not answered

(U) -1: not applicable *else if scalign3=2* -1: not applicable *else if scelig=2*

Note: Variables 'occs4wks1' and 'occs4wks2' differ in similar ways to 'sex4wks1' and 'sex4wks2' respectively

(U) (DN) **occs4wk1** no. of occasions of het. &/or hom. sex in last 4 weeks

(U) *compute sex4wks1 + sam4wks if ((sex4wks1≠-1 & sex4wks1<995) & (sam4wks≠-1 & sam4wks<995))*

compute sex4wks1 + hom4wks if ((sex4wks1≠-1 & sex4wks1<999) & (sam4wks≠-1 & sam4wks<99))

(U – new code) 995: at least one *if (sex4wks1=995 | sam4wks=995)*

999: not answered *else if missing*

-1: not applicable *if sex4wks1=-1 | sam4wks=-1*

(U) (DN) **occs4wk2** no. of occasions of het. &/or hom. sex in last 4 weeks

(U) *compute sex4wks2 + sam4wks if ((sex4wks2≠-1 & sex4wks2<995) & (genisam=1 & sam4wks≠-1 & sam4wks<995))*

(R) *compute sex4wks2 + sam4wks if ((sex4wks2≠-1 & sex4wks2<996) & (genisam=1 & sam4wks≠-1 & sam4wks<99))*

else copy sex4wks2 if (sex4wks2≠-1 & sex4wks2<996) & genisam=2

(U) *else copy sam4wks2 if sex4wks=997 & (genisam=1 & sam4wks≠-1 & sam4wks<999)*

(R) *else copy sam4wks2 if sex4wks2=997 & (genisam=1 & sam4wks≠-1 & sam4wks<99)*

999: not answered *else if missing*

-1: not applicable *if (sex4wks2=-1 | sam4wks=-1) | (sex4wks2=997 & genisam=2)*

(U) (DN) **occs4wkg** no. of occasions of het. &/or hom. sex in last 4 weeks, grouped

0: none *if occs4wk1=0*

1: 1 *if occs4wk1=1*

2: 2 *if occs4wk1=2*

3: 3 *if occs4wk1=3*

4: 4 *if occs4wk1=4*

5: 5-8 *if occs4wk1>=5 & occs4wk1<=8*

6: 9-12 *if occs4wk1>=9 & occs4wk1<=12*

7: 13-16 *if occs4wk1>=13 & occs4wk1<=16*

8: 17-20 *if occs4wk1>=17 & occs4wk1<=20*

(U) 9: 21+ *if occs4wk1>=21 & occs4wk1<995* 9: 21+ *if occs4wk1>=21 & occs4wk1<999*

(U) 99: not answered *if occs4wk1=999 | occs4wk1=995* 99: not answered *if occs4wk1=999*

-1: not applicable *if occs4wk1 = -1*

(ND) **sex4xlm** four or more sex acts, past 4 weeks

1: yes *if occs4wkg=0 | occs4wkg=1 | occs4wkg=2 | occs4wkg=3*

0: no *if occs4wkg=4 | occs4wkg=5 | occs4wkg=6 | occs4wkg=7 | occs4wkg=8 | occs4wkg=9*

9: not answered *if occs4wkg=99*

(ND) **anysex4wk** any sex in the past 4 weeks

1: yes *if (sex4wks1>=1 & sex4wks1<=996) | (sam4wks>=1 & sam4wks<=995)*

0: no *if sex4wks1=0 & sam4wks1=0*

9: not answered *else if sex4wks1=999 | sam4wks=999*

SAME-SEX BEHAVIOUR

Note: homes4w was not used in analysis and was removed from Natsal-3

(R) (N) **homes4w** certainty of no. of occasions of hom. sex in last 4 weeks

1: certain

2: estimate

9: not answered

-1: not applicable *if* scelig=2 | eversam=2 | sam=0 | genisam=2 | sam4wks=0

Note: Variable 'homno4w' has codes 995... 997 in Natsal-2 but not in Natsal-1

(U) **samno4w** (rs5b) **homno4w** no. of hom. sex. partners in last 4 weeks

0 *if* missing & (eversam=2 | sam=0 | genisam=2 | sam4wks=0)

995: at least 1

(U – code removed) 996: at least 2

(U – code removed) 997: at least 3

999: not answered

(U) -1: not applicable *else if* scelign3=2 -1: not applicable *else if* scelig=2

(DN) **prts4wks** no. of het. &/or hom. partners, last 4 weeks

compute no4wks + samno4w *if* ((no4wks ≠ -1 & no4wks < 995) & (samno4w ≠ -1 & samno4w < 995))

999: not answered *if* no4wks ≥ 995 | samno4w ≥ 995

-1: not applicable *if* no4wks = -1 | samno4w = -1

(N) **h1new4w** last hom. sex. partner was a new partner (if had 1 hom. partner in last 4 weeks)

1: yes

2: no

9: not answered

(U) -1: not applicable *if* scelign3=2 | eversam=2 | sam=0 | genisam=2 | sam4wks=0 | (samno4w > 1 & samno4w < 99)

-1: not applicable *if* scelig=2 | eversam=2 | sam=0 | genisam=2 | sam4wks=0 | (samno4w > 1 & samno4w < 99)

samnw4w (N) **homnw4w** no. of new hom. sex partners in last 4 weeks

999: not answered

(U) -1: not applicable *if* scelign3=2 | eversam=2 | sam=0 | genisam=2 | sam4wks=0 | samno4w = 1

-1: not applicable *if* scelig=2 | eversam=2 | sam=0 | genisam=2 | sam4wks=0 | samno4w = 1

Note: Variable homnew4w is derived from homno4w, h1new4w and homnew4w, and although not derived for Natsal-1 is very similar to rs5c

samnew4w (rs5c* D) **homnew4w** no. of new hom. sex. partners in last 4 weeks

0 *if* samno4w=1 & h1new4w=2

1 *if* samno4w=1 & h1new4w=1

(U) *else copy* samnw4w *if* (samno4w > 1 & samno4w < 999) | samno4w=9995

(R) *else copy* samnw4w *if* sam1yr > 1 & sam1yr < 999

999: not answered *if* missing

(U) -1: not applicable *if* scelign3=2 | sam4wks=0

-1: not applicable *if* scelig=2 | sam4wks=0

Note: in Natsal-3 eligibility for samco4w was based on reporting anal sex in the last 4 weeks (at samayou or samahim). Those who had not had anal sex in the last four weeks were coded as -1 'not applicable'.

(U) **samco4w** (s5d U) **homco4w** condom use for hom. anal sex in last 4 weeks

1: yes, used on every occasion

2: yes, used on some occasions

3: not used in last 4 weeks

(U) *answer option removed* 4: not had anal sex in the last 4 weeks

9: not answered

(U) -1: not applicable *if* scelign3=2 | rsex=2 | eversam=2 | sam=0 | genisam=2 | sam4wks=0 | eversam=9 | (rsex=1 & (samayou≠1 | samayou≠2 | samahim≠1 | samahim≠2))

-1: not applicable *if* scelig=2 | rsex=2 | eversam=2 | sam=0 | genisam=2 | sam4wks=0

SAME-SEX BEHAVIOUR

In Natsal-3 eligibility for usecondm was based on only those who had had het. And/or hom. sex in the last 4 weeks. Natsal-3 syntax was updated to code up all possible combinations of cond4wk and samco4w, and these combinations are set out below (updated variable) even though not all occur in the data. Therefore, although the coding below has been updated for Natsal-3, the actual variable has not changed except that those who have not had vaginal / anal sex in the last 4 weeks are coded as 'not applicable'.

(U – see below) (ND) **usecondm** condom use for het. and/or hom. sex in last 4 weeks

- 1: yes, used on every occasion *if* (cond4wk=1 & samco4w=-1) | (cond4wk=-1 & samco4w=1) | (cond4wk=1 & samco4w=1)
- 2: yes, used on some occasions *if* (cond4wk=2 & samco4w=-1) | (cond4wk=-1 & samco4w=2) | (cond4wk=3 & samco4w=1)
- 3: not used in last 4 weeks *if* (cond4wk=3 & samco4w=-1) | (cond4wk=-1 & samco4w=3)
- 4: not had vaginal/anal sex in last 4 weeks *if* (cond4wk=4 & samco4w=-1) | (cond4wk=-1 & samco4w=4) | (cond4wk=4 & samco4w=1)
- 9: not answered *else*
- 1: not applicable *if* cond4wk=-1 & samco4w=-1

usecondm condom use for het. and/or hom. sex in last 4 weeks

- 1: yes, used on every occasion *if* (cond4wk=1 & samco4w=-1) | (cond4wk=-1 & samco4w=1) | (cond4wk=1 & samco4w=1)
- 2: yes, used on some occasions *if* cond4wk=2 | samco4w=2 | (cond4wk=1 & samco4w=3) | (cond4wk=3 & samco4w=1)
- 3: not used in last 4 weeks *if* (cond4wk=3 & samco4w=-1) | (cond4wk=-1 & samco4w=3) | (cond4wk=3 & samco4w=3)
- 9: not answered *else*
- 1: not applicable *if* cond4wk=-1 & samco4w=-1

(ND) **useco_gp** condom use for het. and/or hom. sex in last 4 weeks, grouped

- 1: used on every occasion of sex in last 4 weeks *if* usecondm=1
- 0: not used on every occasion of sex in last 4 weeks *if* usecondm=2 | usecondm=3
- 9: not answered *if* usecondm=9
- (U) -1 not applicable *if* usecondm=-1 -1: not applicable *if* usecondm=-1 | usecondm=4

Note: in Natsal-3 eligibility for sam7day was based on reporting sex in the last 7 days at samoyou, samohim, samayou or samahim

sam7day (rs5e) **hom7day** no. of occasions of hom. sex in last 7 days

0 *if* missing & (eversam=2 | sam=0 | genisam=2 | sam4wks=0)

(U – new code) 995: at least once

(U) 999: not answered

99: not answered

(U) -1: not applicable *if* scalign3=2

-1: not applicable *if* scelig=2

Note: samoyou, samopar, samayou, samahim, samgen were here in Natsal-2. Moved earlier in the questionnaire in Natsal-3.

(N) **ancom** frequency of condom use for hom. anal sex in last year

1: every time

2: most of times

3: occasionally

4: not at all in the last year

9: not answered

(U) -1: not applicable *if* scalign3=2 | rsex=2 | sam=0 | eversam=2 | genisam=2 | ((samayou=5 | samayou=6 | samayou=7) & (samahim=5 | samahim=6 | samahim=7))

1: not applicable *if* scelig=2 | rsex=2 | sam=0 | eversam=2 | genisam=2 | ((samayou=5 | samayou=6 | samayou=7) & (samahim=5 | samahim=6 | samahim=7))

SAME-SEX BEHAVIOUR

(N) **noancom** no. hom. anal sex partners without condom in last year

99: not answered

(U) -1: not applicable if scalign3=2 | rsex=2 | (ancom=1 | ancom=-1)

-1: not applicable if scelig=2 | rsex=2 | (ancom=1 | ancom=-1)

Note: response options for gaypub were reduced in Natsal-3 so the variable has been renamed gaypub2

(R) (N) **gaypub** frequency go to gay pubs/gay clubs

1: 4 times a week or more

2: 2 or 3 times a week

3: about once a week

4: at least once every 2 weeks

5: at least once a month

6: at least once every 3 months

7: at least once every 6 months

8: at least once a year

9: less often than once a year

10: never

99: not answered

-1: not applicable if scelig=2 | sam=0 | eversam=2 | genisam=2

(N) **gaypub2** frequency go to gay pubs / bars/ clubs

1: 1+ time / week

2: less but at least once / month

3: less but at least twice / year

4: less but at least once / year

5: less than once / year

6: never

-1: not applicable if scalign3=2 | sam=0 | eversam=2 | genisam=2 | rsex=2

HETEROSEXUAL PARTNERSHIPS

HETEROSEXUAL PARTNERSHIPS

Note: in Natsal-3 eligibility for this section was different. For heterosexual sex, this was based on reporting ever having had vaginal, oral or anal sex in the sexual practices questions (lastvag, oralyou, oralprt and analsex), or refusing to answer the practices questions. For same sex sex this was based on reporting ever having had same sex sex (genisam=yes & agegsam>12 or nextasam>12). The introductory variable 'partint' can be used to indicate whether a respondent was asked about their heterosexual and same-sex partner numbers (-1: not applicable, 1: questions asked).

(rs7a) **hetlife** no. of het. sex partners, life

(U) 0 or if missing & everhet=0 0 if missing & (het=0 | hetsc=0)

9995: at least one

(R) 9996: at least two

(R) 9997: at least three

(R) 9999: not answered

(U) -1: not applicable else if scelign3=2 -1: not applicable else if scelig=2

(D) **hettot** no. of het. sexual partners, life, grouped

0: 0 if hetlife=0

1: 1 if hetlife=1

2: 2 if hetlife=2

3: 3-4 if hetlife=3 | hetlife=4

4: 5-9 if hetlife>=5 & hetlife<=9

5: 10+ if hetlife>=10 & hetlife<9995

9: not answered if hetlife>=9995 & hetlife<=9999

(U) -1: not applicable else if scelign3=2 -1: not applicable else if scelig=2

(R) (N) **hetsure** certainty of number of het. sex partners, life

1: certain

2: estimate

9: not answered

-1: not applicable if scelig=2 | hetlife=9999 | (hetlife>=0 & hetlife<5)

Note: hetsure was removed in Natsal-3 and has been replaced by a more detailed question asking those reporting 5+ partners how they arrived at this number

(N) **hetcalc** how arrived at estimate for hetlife

-1: not applicable if (scelign3=2 | hetlife<5 | hetlife>=9995)

1: just knew number

2: counted partners

3: estimated / guessed

4: remembered some and added estimated number for rest

5: other

9: not answered

(rs7b) **het5yrs** no. of het. sex partners, last 5 years

(U) 0 if missing & (everhet=0 | hetlife=0) 0 if missing & (het=0 | hetsc=0 | hetlife=0)

9995: at least one

(R) 9996: at least two

(R) 9997: at least three

9999: not answered

(U) -1: not applicable else if scelign3=2 -1: not applicable if scelig=2

HETEROSEXUAL PARTNERSHIPS

(het5yr D) **het5yrg** no. het. sexual partners, last 5 years, grouped

0: 0 if het5yrs=0

1: 1 if het5yrs=1

2: 2 if het5yrs=2

3: 3-4 if het5yrs=3 | het5yrs=4

4: 5-9 if het5yrs>=5 & het5yrs<=9

5: 10+ if het5yrs>=10 & het5yrs<=9995

9: not answered if het5yrs>=9995

(U) -1: not applicable else if scelig=2

-1: not applicable if scelig=2

(rs7d) **het1yr** no. of het. sex partners, last year

(U) 0 if missing & (everhet=0 | hetlife=0 | het5yrs=0) 0 if missing & (het=0 | het5yrs=0 | hetlife=0 | het5yrs=0)

995: at least one

(U – code removed) 996: at least two

(U – code removed) 997: at least three

999: not answered

(U) -1: not applicable else if scelig=2

-1: not applicable if scelig=2

(D) **het1yrg** no. of het. sex partners, last year, grouped

0: 0 if het1yr=0

1: 1 if het1yr=1

2: 2 if het1yr=2

3: 3-4 if het1yr=3 | het1yr=4

4: 5+ if het1yr>=5 & het1yr<995

9: not answered if het1yr>=995

(U) -1: not applicable else if scelig=2

-1: not applicable if scelig=2

Note: in Natsal-2, the codebook didn't describe the derivation of usex correctly. Specifically, the 995 code was included under the code 0. The codebook has been updated below and differences between the codebook and dataset in Natsal-2 are highlighted below.

(usex D) **usex** had unsafe het. sex in last year (2+ het. partners in last year & no condom use in last year)

1: yes if ((het1yr>=2 & het1yr<995) | het1yr=996 | het1yr=997) & ycondom=0

0: no else if missing

9: not answered if het1yr=999 | ycondom=9

-1: not applicable else if het1yr=-1 | ycondom=-1

Notes: 1. The variable usex2 used data on partners in the last year and condom use in the last 4 weeks as a marker of unsafe sex. It has been removed from the Natsal-3 data and replaced with totusex which uses the last year as the reference period for both partners and condom use. This was felt to be a more useful variable.

2. The variable totusex differs from both usex and usex2, which do not take into account whether or not the respondent reported vaginal or anal sex in the past year. For totusex, respondents who did not report any vaginal or anal sex in the past year are coded as 'no'.

(R) (D) **usex2** had unsafe sex (2+ het. &/or hom. partners in last year & inconsistent condom use in last 4 weeks)

1: yes if (tot1yr>=2 & tot1yr <999) & ((rsex=2 & cond4wk=2 | cond4wk=3) | (rsex=1 & usecondm=2 | usecondm=3) | (rsex=1 & (het1yr=996 | het1yr=997) & (sam1yr=-1 | sam1yr=0) & (cond4wk=2 | cond4wk=3))) | (((het1yr=996 | het1yr=997) & (sam1yr=-1 | sam1yr=0)) & (cond4wk=2 | cond4wk=3)))

0: no if (tot1yr=0 | tot1yr=1) | (rsex=2 & (tot1yr>=2 & tot1yr<999) & (cond4wk=1 | cond4wk=4)) | (rsex=1 & (tot1yr>=2 & tot1yr <999) & (usecondm=1 | usecondm=4)) | (((het1yr=996 | het1yr=997) & (sam1yr=-1 | sam1yr=0)) & (cond4wk=1)) | (rsex=1 & (sam1yr>=2 & sam1yr<995) & sex4wks2=997 & samco4w=4)

9: not answered else if (tot1yr=999 | usecondm=9 | no4wks=999 | samno4w=999 | (sex4wks2=996 & sam1yr=0)) | missing

-1: not applicable if occs4wk2=0 | occs4wk2=-1

HETEROSEXUAL PARTNERSHIPS

(N) **totusex** had unsafe sex in the last year (2+ het &/or hom. partners & no condom use in last year)
1: yes if (het1yr>=2 & het1yr<995 & (vagsexyr=1 | analhmr=1) & ycondom=0) | (sam1yr>=2 & sam1yr<995 & rsex=1 & anlhmr=1 & ancom=4) | (sam1yr=1 & het1yr=1 & (vagsexyr=1 | analhmr=1) & rsex=1 & anlhmr=1 & ycondom=0 & ancom=4)
0: no if ((het1yr<2 | ycondom=1) & (sam1yr<2 | anlhmr=0 | anlhmr=-1 | ancom=1 | ancom=2 | ancom=3)) | (het1yr=1 & sam1yr=1 & (anhlmr=0 | anlhmr=-1 | ycondom=1 | ancom=1 | ancom=2 | ancom=3) | (vagsexyr=0 & analhmr=0) | (het1yr=0 & sam1yr=0)
9 not answered else if (het1yr=999 | sam1yr=999 | ycondom=9 | anlhmr=9 | ancom=9 | het1yr=995 | sam1yr=995 | ((het=0 & het1yr>0) | het=9)

(N) **nonewpt** no. of new het. sex partners, last year, (if had >1 het. partner in last year)

999: not answered

(U) -1: not applicable if scelign3=2 | (het1yr=0 | het1yr=1)

-1: not applicable if scelig=2 | (het1yr=0

| het1yr=1)

(N) **hetnewp** new het. sex partner, last year, (if had 1 het. partner in last year)

1: yes

2: no

9: not answered

(U) -1: not applicable if scelign3=2 | (het1yr>1 & het1yr<9999) | het1yr=0

-1: not applicable if scelig=2 | (het1yr>1 & het1yr<9999) | het1yr=0

Note: in Natsal-2 if het1yr=995 hetnonew was coded as 99 even if there was a valid response at nonewpt/hetnewp. In Natsal-3 all responses at nonewpt & hetnewp were included in hetnonew.

Note: Variable hetnonew is derived from hetnewp and nonewpt:

(ND) **hetnonew** no. of new het. sex partners, last year

0 if het1yr=0 | (het1yr=1 & hetnewp=2)

1 if het1yr=1 & hetnewp=1

(U) copy nonewpt if (het1yr>1 & het1yr<995) | (nonewpt>=0 & mpmewpt<999) copy nonewpt if het1yr>1 & het1yr<995

99: not answered if missing

-1: not applicable if het1yr=-1

(ND) **hetnew2** 1+ new het. sex. partners, last year

0: 0 new het. sex partners if hetnonew=0

1: 1+ new het. sex partner(s) if hetnonew>=1 & hetnonew<999

99: not answered if hetnonew=999

-1: not applicable if hetnonew=-1

(ND) **hetnew3** no. of new het. sex partners in last year, grouped

0: 0 new het. sex partners if hetnonew=0

1: 1 new het. sex partner if hetnonew=1

2: 2+ new het. sex partners if hetnonew>=2 & hetnonew<999

99: not answered if hetnonew=999

-1: not applicable if hetnonew=-1

Note: questions were added to Natsal-3 about oral sex only partners in the last year. These questions were included in Natsal-2 but weren't included in the final dataset as there were some problems with the programme.

The answers to introos, oralone, oralpre have been imputed for respondents who only reported oral sex (not vaginal or anal sex) in the last year.

(N) **introos** any het. sex partners only had oral sex with, last year

1: yes or if (lastvag>=5 & lastvag<=7 & analsex>=5 & analsex<=7 & (oralprt<=4 | oralyou<=4)

2: no or if missing & (het1yr=0 | expscale=6 | ((oralprt=5 | oralprt=6 | oralprt=7) & (oralyou=5 | oralyou=6 | oralyou=7))

9: not answered -1: not applicable if scelig=2

HETEROSEXUAL PARTNERSHIPS

(N) **oralon** no. of oral sex only partners in last year

0 if missing & introos=2

calculate oralon=hetyr if (lastvag>=5 & lastvag<=7 & analsex>=5 & analsex<=7 & (oralprt<=4 | oralyou<=4)

999: not answered

-1: not applicable if introos=-1

(N) **oralpre** previous estimates of no. partners include these oral sex only partners

1: yes, all included or if (lastvag>=5 & lastvag<=7 & analsex>=5 & analsex<=7 & (oralprt<=4 | oralyou<=4)

2: no, not all included or if missing and (introos=1 & ((oralon>0 & oralon<999) & (hetyr>0 & hetyr<995) & oralon>hetyr) | ((oralon>0 & oralon<999) & (hetyr=995 | hetyr=999)))

9: not answered

-1: not applicable if introos=2

Note: h1yrtot was a direct question of respondents about their total number of opposite-sex partners in the last year, including those with whom they had only had oral sex. However, we found a large proportion of respondents gave answers that were inconsistent with hetyr and oralno. We have therefore derived a second variable h1yrtot2 which estimates the number of partners in the last year using hetyr and oralon. We advise data users to treat both variables with caution, as we do not know which is more accurate.

(N) **h1yrtot** no. of het sex partners in last year including oral sex only partners

0 if missing & hetyr=0

copy hetyr if missing & (introos=2 | oralpre=1)

999: not answered

-1: not applicable if scelign3=2

Note: h1yrtot2 assumes that if h1yrtot and oralon are greater than 0 it is valid to have h1yrtot<(hetyr+oralon) because a respondent may have previously reported some, but not all, oral sex partners at hetyr. For other inconsistent answers, h1yrtot has been recalculated.

(ND) **h1yrtot2** no. of het sex partners in last year including oral sex only partners (recalculated for those with inconsistent data)

0 if missing & hetyr=0 & h1yrtot=0 & introos#1

copy hetyr if (introos=2 | oralpre=1)

copy h1yrtot if h1yrtot=hetyr+oralon

copy h1yrtot if hetyr>=0 & hetyr<995 & oralon>=0 & oralon<999 & h1yrtot<(hetyr+oralon) AND h1yrtot>hetyr.

calculate hetyr+oralon else if hetyr>=0 & hetyr<995 & oralon>=0 & oralon<999

999: not answered

-1: not applicable if scelig=2

Note: Natsal-3 included a new question to find out whether respondents had engaged in swinging in the last year.

(N) **swing** engaged in sex with other couples with a het. partner

1: yes

2: no or if missing & (hetyr=0 | hetyr=1 | hetyr=-1) & (h1yrtot=0 & h1yrtot=1 | h1yrtot=-1)

9: not answered

-1: not applicable if scelign3=2

(N) **het3mnt** no. of het. sex partners, last 3 months

(U) 0 if missing & (everhet=0 | hetlife=0 | het5yrs=0 | hetyr=0)

0 if missing & (het=0 | hetsc=0 | hetlife=0 | het5yrs=0 | hetyr=0)

995: at least one

(U – code removed) 996: at least two

(U – code removed) 997: at least three

999: not answered

(U) -1: not applicable else if scelign3=2

-1: not applicable else if scelig=2

HETEROSEXUAL PARTNERSHIPS

(N) **het3mntg** no. of het. sex partners, last 3 months, grouped

0: none *if* het3mnt=0

1: 1 *if* het3mnt=1

2: 2+ *if* het3mnt $\geq$ 2 & het3mnt<995

9: not answered *if* het3mnt $\geq$ 995

-1: not applicable *if* het3mnt=-1

SAME-SEX PARTNERSHIPS

SAME SEX PARTNERSHIPS

*Notes: 1. as above in the samesex sex section, all instances of 'hom' in variable names have been changed to 'sam'.
2. In Natsal-3, people were not routed into this section if they did not report same sex genital contact (genisam=no) or reported that their only same sex experience was before the age of 13 (agegsam=no). Because of this change in the denominator, additional variables have been derived for the Natsal-2 dataset, which have the same denominator as Natsal-3, to enable comparisons. These variables are not available on the archived Natsal-2 data so please contact the Natsal team (via Dr Catherine Mercer) if you require a Natsal-2 dataset which contains these additional variables.*

samlife (rs8a) **homlife** no. of hom. sex partners, life

(U) 0 if sam=0 | eversam=0 | genisam=2 | (nextageg=2 & agegsam>0 & agegsam<13)

0 if hom=0 | eversam=0 | genisam=0

9995: at least one

(U – code removed) 9996: at least two

(U – code removed) 9997: at least three

9999: not answered

(U) -1: not applicable else if scelign3=2 & eversam=2 -1: not applicable else if scelig=2

(R) (N) **homsure** certainty of number of hom. sex partners, life

-1: not applicable if scelig=2 | samlife=9999 | (samlife>=0 & samlife<5)

1: certain

2: estimate

9: not answered

Note: homsure was removed in Natsal-3 and has been replaced by a more detailed question asking those reporting 5+ partners how they arrived at this number

(N) **samcalc** how arrived at estimate for samlife

1: just knew number

2: counted partners

3: estimated / guessed

4: remembered some and added estimated number for rest

5: other

-1: not applicable if (scelign3=2 & eversam=2) | samlife<5 | samlife>9995

samtot (homtot D) **hometot** no. of hom. sex partners, life, grouped

0: 0 if samlife=0

1: 1 if samlife=1

2: 2 if samlife=2

3: 3-4 if samlife=3 | samlife=4

4: 5-9 if samlife>=5 & samlife<=9

5: 10+ if samlife>=10 & samlife<9995

9: not answered if samlife>=9995 & samlife<=9999

(U) -1: not applicable else if scelign3=2 & eversam=2 -1: not applicable else if scelig=2

(R) (D) **homever** ever had hom. sex partner(s)

0: no if genisam=2 & (samlife=-1 | samlife=0)

1: yes if genisam=1 | (samlife>=1 & samlife<=9997)

9: not answered if samlife>=9997

sam5yrs (rs8b) **hom5yrs** no. of hom. sex partners, last 5 years

0 or if hom=0 | eversam=0 | genisam=0 | samlife=0

9995: at least one

9996: at least two

9997: at least three

9999: not answered

(U) -1: not applicable else if scelign3=2 & eversam=2 -1: not applicable if scelig=2

TOTAL PARTNERSHIPS

sam5yrg (D) **hom5yrg** no. of hom. sex partners, last 5 years, grouped

0: 0 if sam5yrs=0

1: 1 if sam5yrs=1

2: 2 if sam5yrs=2

3: 3-4 if sam5yrs=3 | sam5yrs=4

4: 5-9 if sam5yrs>=5 & sam5yrs<=9

5: 10+ if sam5yrs>=10 & sam5yrs<9995

9: not answered if sam5yrs>=9995 & sam5yrs<=9999

(U) -1: not applicable else if scalign3=2 & eversam=2 -1: not applicable if scelig=2

samlst5y (ND) **homlst5y** 1+ hom. sex partner(s), last 5 years

0: no if sam5yrs=0

1: yes if sam5yrs>=1 & sam5yrs<9999

9: not answered if sam5yrs=9999

(U) -1: not applicable else if scalign3=2 & eversam=2 -1: not applicable if scelig=2

sam1yr (rs8d) **hom1yr** no. of hom. sex partners, last year

0 if sam=0 | eversam=0 | genisam=0 | samlife=0 | sam5yrs=0

995: at least one

(U – code removed) 996: at least two

(U – code removed) 997: at least three

999: not answered

(U) -1: not applicable else if scalign3=2 & eversam=2 -1: not applicable if scelig=2

sam1yrg (D) **hom1yrg** no. of hom. sex partners, last year, grouped

0: 0 if sam1yr=0

1: 1 if sam1yr=1

2: 2 if sam1yr=2

3: 3-4 if sam1yr=3 | sam1yr=4

4: 5+ if sam1yr>=5 & sam1yr<995

9: not answered if sam1yr>=995 & sam1yr<=999

(U) -1: not applicable else if scalign3=2 & eversam=2 -1: not applicable if scelig=2

samlst1y (ND) **homlst1y** 1+ hom. sex partner(s), last year

0: no if sam1yr=0

1: yes if sam1yr>=1 & sam1yr<999

9: not answered if sam1yr=9

(U) -1: not applicable else if scalign3=2 & eversam=2 -1: not applicable if scelig=2

snnewwp (N) **hnonewp** no. new hom. sex partners, last year (if had >1 hom. partners in last year)

999: not answered

(U) -1: not applicable if scalign3=2 | (sam1yr=0 | sam1yr=1)

-1: not applicable if scelig=2 | (sam1yr=0 | sam1yr=1)

samnewp (N) **homnewp** new hom. sex partner, last year (if 1 hom. partner in last year)

1: yes

2: no

(U) 9: not answered if sam1yr=999 999: not answered

(U) -1: not applicable else if (scalign3=2 & eversam=2) | (sam1yr>1 & sam1yr<999) | sam1yr=0

-1: not applicable if scelig=2 | (sam1yr=-1 | sam1yr=0 | sam1yr>1)

Note: Variable samnonew is derived from samnewp and hnonewp:

samnonew (ND) **homnonew** no. of new hom. sex partners, last year

0 if sam1yr=0 | (sam1yr=1 & samnewp=2)

1 if sam1yr=1 & samnewp=1

copy hnonewp if sam1yr>1 & sam1yr<995

999: not answered if missing

-1: not applicable if sam1yr=-1

TOTAL PARTNERSHIPS

samnew2 (ND) **homnew2** 1+ new hom. sex partner(s), last year
0: 0 new hom. sex partners *if* samnonew=0
1: 1+ new hom. sex partner(s) *if* samnonew>=1 & samnonew<99
99: not answered *if* samnonew=999
-1: not applicable *if* samnonew=-1

samnew3 (ND) **homnew3** no. of new hom. sex partners, last year, grouped
0: 0 new hom. sex partners *if* samnonew=0
1: 1 new hom. sex partner *if* samnonew=1
2: 2+ new hom. sex partner(s) in last year *if* samnonew>=2 & samnonew<99
99: not answered *if* samnonew=99
-1: not applicable *if* samnonew=-1

sam3mnt (N) **hom3mnt** no. hom. sex partners, last 3 months
0 *if* sam=0 | eversam=2 | genisam=2 | samlife=0 | sam5yrs=0 | sam1yr=0
995: at least one
(U – code removed) 996: at least two
(U – code removed) 997: at least three
999: not answered
(U) -1: not applicable *if* scalign3=2 & eversam=2 --1: not applicable *if* scelig=2

(U – see **sam3mntg** below) (ND) **hom3mntg** no. hom. sex partners, last 3 months, grouped
0 *if* sam3mnt=0
1: 1+ *if* sam3mnt>=1 & sam3mnt<999
else sysmis

(U) **sam3mntg** no. hom. sex partners, last 3 months, grouped
0 incl. not applicable *if* sam3mnt=0 | sam3mnt=-1
1: 1+ *if* sam3mnt>=1 & sam3mnt<995
9: not answered

OVERLAPPING PARTNERSHIPS

Note: additional question on concurrency in the last 5 years (previously concurrency in the last year was estimated from most recent partners data).

Note: further variables on concurrency can be found in the detailed partnerships section (page 102).

(N) **overlp5y** any overlap between partners (ie. Sex with one then another then the first again) in last 5 years
1: yes
2: no *if* (het5yrs+sam5yrs)=1 | (het5yrs+sam5yrs)=0
9: not answered
-1: not applicable *if* scalign3=2 & eversam=2

ONLINE PARTNERS

(N) **netsex** used internet to find sexual partner in last 12 months (if at least one partner ever)
1: yes
2: no
9: not answered
-1: not applicable *if* (scalign3=2 & eversam=2) | (everhet=0 & samsex=0)

TOTAL PARTNERSHIPS

TOTAL PARTNERSHIPS

(ND) **totlife** no. of het. & hom. sex partners, life

compute hetlife + samlife if hetlife>=0 & hetlife<9995 & samlife>=0 & samlife<9995

(U – code added) 9995 if ((hetlife=9999 | hetlife=9995) & (samlife>0 & samlife<9999)) | ((samlife=9999 | samlife=9995) & (hetlife>0 & hetlife<9999))

(U) 9999: not answered else if hetlife=9999 | samlife=9999

9999: not answered if hetlife>=9995 | samlife>=9995

-1: not applicable if hetlife=-1 | samlife=-1

(ND) **tottot** no. of het. & hom. sex partners, life, grouped

0: 0 if totlife=0

1: 1 if totlife=1

2: 2 if totlife=2

3: 3-4 if totlife=3 | totlife=4

4: 5-9 if totlife>=5 & totlife<=9

5: 10+ if totlife>=10 & totlife<9995

9: not answered if totlife>=9995

-1: not applicable if totlife=-1

(ND) **tot5yrs** no. of het. & hom. sex partners, last 5 years

compute het5yrs + sam5yrs if het5yrs>=0 & het5yrs<9995 & sam5yrs>=0 & sam5yrs<9995

(U – code added) 9995 if if ((het5yrs=9999 | het5yrs=9995) & (sam5yrs>0 & sam5yrs<9999)) | ((sam5yrs=9999 | sam5yrs=9995) & (het5yrs>0 & het5yrs<9999))

(U) 9999: not answered else if het5yrs=9999 | sam5yrs=9999

9999: not answered else if het5yrs>9995 | sam5yrs>9995

-1: not applicable if het5yrs=-1 | sam5yrs=-1

(ND) **tot5yrg** no. of het. & hom. sex partners, last 5 years, grouped

0: 0 if tot5yrs=0

1: 1 if tot5yrs=1

2: 2 if tot5yrs=2

3: 3-4 if tot5yrs=3 | tot5yrs=4

4: 5+ if tot5yrs>=5 & tot5yrs<9995

9: not answered if tot5yrs>=9995

-1: not applicable if tot5yrs=-1

(ND) **tot5yrg2** no. of het. & hom. sex partners, last 5 years, grouped

0: 0 if tot5yrs=0

1: 1 if tot5yrs=1

2: 2-4 if tot5yrs>=2 & tot5yrs<=4

3: 5-9 if tot5yrs>=5 & tot5yrs<=9

4: 10+ if tot5yrs>=10 & tot5yrs<9995

9: not answered if tot5yrs>=9995

-1: not applicable if tot5yrs=-1

(ND) **tot1yr** no. of het. & hom. sex partners, last year

compute het1yr + sam1yr if het1yr>=0 & het1yr<995 & sam1yr>=0 & sam1yr<995

(U – code added) 995 if ((het1yr=999 | het1yr=995) & (sam1yr>0 & sam1yr<999)) | ((sam1yr=995 | sam1yr=999) & (hetlife>0 & hetlife<9999))

(U) 999: not answered else if het1yr=999 | sam1yr=999

999: not answered if het1yr>=995 | sam1yr>=995

-1: not applicable if het1yr=-1 | sam1yr=-1

(ND) **tot1yrg** no. of het. & hom. sex partners, last year, grouped

0: 0 if tot1yr=0

1: 1 if tot1yr=1

2: 2 if tot1yr=2

3: 3-4 if tot1yr=3 | tot1yr=4

TOTAL PARTNERSHIPS

4: 5+ *if* tot1yr>=5 & tot1yr<995
9: not answered *if* tot1yr>=995
-1: not applicable *if* tot1yr=-1

(ND) **totnewyr** no. of new het. & hom. sex partners, last year
compute hetnnew + samnnew *if* hetnnew>=0 & hetnnew<999 & samnnew>=0 & samnnew<999
else hetnnew *if* samnnew=-1 & hetnnew>=0 & hetnnew<999
else samnnew *if* hetnnew=-1 & samnnew>=0 & samnnew<999
999: not answered *if* hetnnew=999 | samnnew=999
-1: not applicable *if* hetnnew =-1 | samnnew =-1

(ND) **totnewy2** 1+ new het./hom. sex partners, last year
0: 0 new het./hom. sex partners *if* totnewyr =0
1: 1+ new het./hom. sex partners *if* totnewyr>=1 & totnewyr <999
99: not answered *if* totnewyr=999
-1: not applicable *if* totnewyr=-1

(ND) **totnewy3** no. of new het. & hom. sex partners, last year, grouped
0: 0 new het./hom. sex partners *if* totnewyr =0
1: 1 new het./hom. sex partner *if* totnewyr=1
2: 2+ new het./hom. sex partners *if* totnewyr>=2 & totnewyr<999
99: not answered *if* totnewyr=999
-1: not applicable *if* totnewyr=-1

(ND) **nocondom** no. of het. &/or hom. sex partners without a condom, last year
0: 0 *if* (yrcond=2 & (ancom=-1 | ancom=1)) | (yrcond=7 & (ancom=-1 | ancom=1))
else compute nonocon + noancom *if* (nonocon≠-1 & nonocon<996) & (noancom≠-1 & noancom<99)
else copy nonocon *if* ((nonocon≠-1 & nonocon<996) & (sam1yr=-1 | sam1yr=0) | ((nonocon≠-1 & nonocon<996) & ancom=-1) | ((nonocon≠-1 & nonocon<996) & ancom=1)
else copy noancom *if* ((het1yr=0 | het1yr=-1) & (noancom≠-1 & noancom<99)) | (yrcond=1 & (noancom≠-1 & noancom<99))
else 1 *if* (sam1yr=-1 | sam1yr=0) & het1yr=1 & nonocon=996
999: not answered *if* (sam1yr=-1 | sam1yr=0 & (het1yr>1 & het1yr<999) & nonocon=999) | nonocon=999 | noancom=999

(ND) **nocondmg** no. of het. &/or hom. sex partners without a condom, last year, grouped
0: 0 *if* nocondom=0
1: 1 *if* nocondom=1
2: 2+ *if* nocondom>=2 & nocondom<999
9: not answered *if* nocondom=999

DETAILED PARTNERSHIP HISTORY

TRAVEL & SEX ABROAD

Note: the travel and sex abroad section appeared here in Natsal-2 but has been moved after the most recent partners section in Natsal-3

DETAILED PARTNERSHIP HISTORY

Note: Variables beginning 'r1...' and not ending in a number refer to the most recent sexual partner. These variables are not limited to partners in the last 5 years unlike Natsal-1, which only asks the detailed partnership questions if the respondent reported 2+ partners in the last 5 years.

Note: where het & hetsc are used to derive codes for variables in Natsal-2, in Natsal-3 the new variable everhet (derived from het & hetcheck) was used instead. Similarly, where hom was used in Natsal-2, eversam was used in Natsal-3, and where scelig was used in Natsal-2, scalign3 was used in Natsal-3. The coding below has been updated to reflect this.

Note: Variable r1datey is expressed as 4 digits but the Natsal-1 equivalent is only expressed as 2 digits.

(U – uses everhet instead of hetsc and het) (s10a2) **r1datey** year of most recent sex with most recent partner

9999: not answered

-1: not applicable if sceligN3=2 | ((everhet=0) & sam=0)

Note: in Natsal-3 only asked for month if r1datey was in the last five years

(U – uses everhet instead of hetsc and het) (s10a1) **r1datem** month of most recent sex with most recent partner

1: January

2: February

3: March

4: April

5: May

6: June

7: July

8: August

9: September

10: October

11: November

12: December

(U) 13: unable to estimate the month

(U – code removed) 97: can't remember month or if r1datem=98

99: not answered

-1: not applicable if sceligN3=2 | (everhet=0 & sam=0) | (dateoi - r1datey<5)

(U – uses everhet instead of hetsc) (s10e) **r1sex** sex of most recent partner

1: male or if (rsex=1 & (everhet=0 & samlife>0 & samlife<9999 & totlife>0 & totlife<9999)) | (rsex=2 & (everhet=1 & samlife=0 totlife>0 & totlife<9999))

2: female or if (rsex=2 & (everhet=0 & samlife>0 & samlife<9999 & totlife>0 & totlife<9999)) | (rsex=1 & (everhet=1 & samlife=0 totlife>0 & totlife<9999))

9: not answered

-1: not applicable else if sceligN3=2 | (everhet=0 & sam=0) | (hetlife=0 & samlife=0)

Notes: 1. In Natsal-3, this question was only asked of those who reported having ever had vaginal or anal sex. There is an additional answer option for r1cond1 in Natsal-3. Respondents can now specify that last sex was oral sex.

2. Wave 1 of fieldwork for Natsal-3 there was a CAPI routing error which meant that some eligible respondents were not asked r1cond1. These were respondents who had had vaginal or analsex longer than 5 years ago, but who had not had vaginal or anal sex in the last year. Any such

DETAILED PARTNERSHIP HISTORY

respondents are coded as 9 'not answered' for r1condl. This problem was fixed from wave 2 onwards.

(U) (N) **r1condl** condom used for most recent sex with most recent partner

1: yes

2: no

(U) 3: only had oral sex on the most recent occasion *or if* (lastvag=7 & analsex=7 & (samahim=-1 | samahim=7) & (samayou=-1 | samayou=7) & (oralyou=1 to 6 | oralprt=1 to 6 | (oralyou=7 & oralprt=7 & (samoyou=1 to 6 | samopar=1 to 6)))

9: not answered

(U) -1: not applicable *if* scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0) | (lastvag=7 & analsex=7 & (samahim=-1 | samahim=7) & (samayou=-1 | samayou=7)) | (rsex=2 & r1sex=2)

-1: not applicable *if* scelig=2 | ((het=0 | hetsc=0) & sam=0) | (rsex=2 & r1sex=2)

Note: questions added to establish: i) relationship status at first (r1rel1s) and most recent (r1rel) sex with partner and, ii) for partnerships where most recent sex occurred in the last year, whether they are likely to have sex again. In Natsal-2 this was assumed based on answers to certain questions.

(N) **r1rel** relationship status at most recent sex

1: living together / married / civil partnership

2: steady relationship

3: used to be in steady relationship but not at time

4: known each other for a while but not in steady relationship

5: recently met

6: just met for first time

9: not answered

-1: not applicable *if* sceligN3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0)

(N) **r1sexagn** likely to have sex again if most recent sex <1y ago

1: yes

2: probably

3: probably not

4: no

5: don't know

9: not answered

-1: not applicable *if* sceligN3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0) | dlover1p1=1

Note: in Natsal-3, answer options for r1first were made clearer to reduce question mis-interpretation

(U) (*) (s10b) **r1first** most recent occasion also first occasion with most recent partner

(*) 1: yes, only had sex with him/her once 1: yes, was the first occasion

(*) 2: no, had sex on more than one occasion 2: no, was not the first occasion

9: not answered

(U) -1: not applicable *if* scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0)

-1: not applicable *if* scelig=2 | ((het=0 | hetsc=0) & sam=0)

Note: Variable r1fy is expressed as 4 digits but the Natsal-1 equivalent is only expressed as 2 digits.

(U) (s10c2 *) **r1fy** year of 1st occasion with most recent partner

9999: not answered

(U) -1: not applicable *if* scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0) | r1first=1

-1: not applicable *if* scelig=2 | ((het=0 | hetsc=0) & sam=0) | r1first=1

(U) (s10c1 *) **r1fm** month of 1st occasion with most recent partner

1: January

2: February

3: March

4: April

5: May

DETAILED PARTNERSHIP HISTORY

6: June

7: July

8: August

9: September

10: October

11: November

12: December

(U) 13: unable to estimate the month

(U – code removed) 97: can't remember month or if r1datem=98

99: not answered

(U) -1: not applicable if scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0) | r1first=1 | dfover5p1=1

-1: not applicable if scelig=2 | ((het=0 | hetsc=0) & sam=0) | r1first=1

Note: firstdtp1 combines r1fy with r1fm to give a date format variable. If only had sex with partner on one occasion (r1first=1) then uses r1datey and r1datem to derive date. All missing values are set to sysmiss. If greater understanding of the missing values is needed, the indicator variable firstdtp1ind can be used to see whether the date is missing because it was not given, or because the question was not applicable.

Note: if r1fm/r1datem unknown, month set to June for firstdtp1

(ND) **firstdtp1** date of first sex with P1, most recent partner
sysmis if missing

Note: lastdtp1 uses r1datey and r1datem to derive date of most recent sex with partner. All missing values are set to sysmiss. If greater understanding of the missing values is needed, the indicator variables lastdtp1ind can be used to see whether the date is missing because it was not given, or because the question was not applicable. If r1datem unknown, month set to June.

(ND) **lastdtp1** date of most recent (last) sex with P1, most recent partner
sysmis if missing

Note: dfover1p1 is derived from a combination of date of interview (dateoi), date of first sex with partner (r1fy&r1fm or r1datey&r1datem) i.e. yes if r1fy/r1fm were more than a year before date of interview, or if only had sex once and r1datey/r1datem were more than a year before date of interview. Calculated to be yes if month of sex with partner unknown but possibly in last year (i.e. year is year before interview) and month of interview in July to December (i.e. there is greater likelihood that it was over 1 year ago than less than 1 year ago),

(ND) **dfover1p1** first sex with partner 1 occurred over 1 year ago

-1: not applicable if firstdtp1ind=-1

1: yes, over 1 year ago if (firstdtp1ind=1 & ((dateoi-1)>firstdtp1)) | (firstdtp1ind=2 & ((r1first=1 & r1datey>0 & (dateyoi-r1datey=0)) | (r1first=2 & r1fy>0 & (dateyoi-r1fy=0)) | (r1first=1 & r1datey>0 & (dateyoi-r1datey=1) & range(datemoi 7..12)) | (r1first=2 & r1fy>0 & (dateyoi-r1fy=1) & range(datemoi 7..12))))

2: no, in the last year if (firstdtp1ind=1 & ((dateoi-1)<=firstdtp1)) | (firstdtp1ind=2 & (r1first=1 & r1datey>0 & (dateyoi-r1datey=0)) | (r1first=2 & r1fy>0 & (dateyoi-r1fy=0)) | (r1first=1 & r1datey>0 & (dateyoi-r1datey=1) & range(datemoi 1..6)) | (r1first=2 & r1fy>0 & (dateyoi-r1fy=1) & range(datemoi 1..6))))

9: not answered

Similarly dfover5p1 is derived from a combination of date of interview (dateoi), date of first sex with partner (r1fy&r1fm or r1datey&r1datem)

(ND) **dfover5p1** first sex with partner 1 occurred over 5 years ago

-1: not applicable if firstdtp1ind=-1

1: yes, over 5 years ago

DETAILED PARTNERSHIP HISTORY

2: no, in the last 5 years
9: not answered

Similarly dlover1p1 is derived from a combination of date of interview (dateoi) and date of most recent sex with partner (r1datey & r1datem).

(ND) **dlover1p1** most recent (last) sex with partner 1 occurred over 1 year ago.
-1 not applicable if lastdtp1ind=-1
1: yes, over 1 year ago
2: no, in the last year
9: not answered

Similarly dlover5p1 is derived from a combination of date of interview (dateoi), date of most recent sex with partner (r1datey & r1datem)

(ND) **dlover5p1** most recent (last) sex with partner 1 occurred over 5 years ago
-1: not applicable if firstdtp1ind=-1
1: yes, over 5 years ago
2: no, in the last 5 years
9: not answered

Note: In Natsal-3 respondents reporting that first sex was in same month as interview or the previous calendar month were asked an additional question to determine the length of the sexual relationship. This will allow better interpretation of the number of sex acts variable. Which question is asked depends on whether respondent is likely to have sex with partner again, if r1sexagn=yes/ probably then asked r1mtha; if r1sexagn=no/ probably not/don't know then asked r1mthb

(N) **r1mtha** time since first sex
1: less than 7 days
2: 7 days – 2 weeks
3: 2-4 weeks
4: over 4 weeks
9: not answered
-1: not applicable if scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0) | r1first=1 | (r1sexagn=3 | r1sexagn=4) | dlover1p1=1

(N) **r1mthb** time between first and last sex
1: less than 7 days
2: 7 days – 2 weeks
3: 2-4 weeks
4: over 4 weeks
9: not answered
-1: not applicable if scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0) | r1first=1 | (r1sexagn=1 | r1sexagn=2) | dlover1p1=1

DETAILED PARTNERSHIP HISTORY

Note: 1. In Natsal-3, this question was only asked of those who reported having ever had vaginal or anal sex. There is an additional answer option for r1condf in Natsal-3. Respondents can now specify that first sex was oral sex
 2. Wave 1 of fieldwork for Natsal-3 there was a CAPI routing error which meant that some eligible respondents were not asked r1condf. These were respondents who had had vaginal or analsex longer than 5 years ago, but who had not had vaginal or anal sex in the last year. Any such respondents are coded as 9 'not answered' for r1condf. This problem was fixed from wave 2 onwards.

(U) (N) **r1condf** condom used on 1st occasion with most recent partner

1: yes

2: no

(U) 3: only had oral sex on the first occasion *else if* lastvag=7 & analsex=7 & (samahim=-1 | samahim=7) & (samayou=-1 | samayou=7) & ((oralyou=1 to 6 | oralprt=1 to 6) | (oralyou=7 & oralprt=7 & (samoyou=1 to 6 | samopar=1 to 6)))

9: not answered

(U) -1: not applicable *if* scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0) *else if* (lastvag=7 & analsex=7 & (samahim=-1 | samahim=7) & (samayou=-1 | samayou=7)) | (rsex=2 & r1sex=2)

-1: not applicable *if* scelig=2 | ((het=0 | hetsc=0) & sam=0) | r1first=1 | (rsex=2 & r1sex=2)

Note: r1rel1s collects relationship status at first sex.

(N) **r1rel1s** relationship status at first sex

1: living together / married / civil partnership

2: steady relationship

3: used to be in steady relationship but not at time

4: known each other for a while but not in steady relationship

5: recently met

6: just met for first time

9: not answered

-1: not applicable *if* scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0) | r1first=1

(U) (s10d) **r1ptage** age on 1st occasion with most recent partner

(U) 996: NA – partners age at 1st sex < 13 years

(U) 997: don't know 97: don't know

(U) 999: not answered 99: not answered

(U) -1: not applicable *if* scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0)

-1: not applicable *if* scelig=2 | ((het=0 | hetsc=0) & sam=0)

(ND) **rafsmr** respondent's age in completed years at 1st sex with most recent partner

compute (((r1fy-1900)*12) + r1fm) - (((rdoby-1900)*12) + rdbm) / 12 *if* (r1fy≠-1 & r1fy≠9999) & (r1fm≠-1 & r1fm≠97 & r1fm≠99) & rdbm≠99

else compute (((r1datey-1900)*12) + r1datem) - (((rdoby-1900)*12) + rdbm) / 12 *if* r1first=1 & (r1datey≠-1 & r1datey≠9999) & (r1datem≠-1 & r1datem≠97 & r1datem≠99) & rdbm≠99

else compute r1fy - rdoby *if* (r1fy≠-1 & r1fy≠9999)

else compute r1datey-rdoby *if* r1first=1 & (r1datey≠-1 & r1datey≠9999)

(U) 999: not answered *else if missing* | rafsrmr<0 99: not answered *else if missing* | rafsrmr<0

(U) -1: not applicable *if* scelign3=2 -1: not applicable *if* scelig=2

Notes:

1. *when calculating ad_mrp, if respondent and partner are different sexes then woman's age was subtracted from man's age.*
2. *For most recent partner only, age difference can be computed using respondent's current age and information from the household grid, where MRP date/age data are missing but there is evidence from r1rel that most recent partner is same as current cohabiting partner.*
3. *All age differences of greater than 20 (ad_mrp<-20 | ad_mrp>20) were investigated manually to check consistency of respondent's answers. Where there was evidence from elsewhere in the questionnaire (e.g. cohabitation history) that the respondent had made a data entry error or misunderstood a question, these variables were edited (changed to missing, or changed to most plausible value if there was strong evidence as to what the correct value was).*

DETAILED PARTNERSHIP HISTORY

(ND) **ad_mrp** age difference between respondent & most recent partner
compute (rafsmr-r1ptage) if rsex=1 & rafsmr≠-1 & rafsmr≠996 & rafsmr≠999 & r1ptage≠996 & r1ptage≠997 & r1ptage≠999
compute (r1ptage-rafsmr) if rsex=2 & rafsmr≠-1 & rafsmr≠996 & rafsmr≠999 & r1ptage≠996 & r1ptage≠997 & r1ptage≠999
compute (dage-hhpartage) if rsex=1 & hhpartage≠-1 & hhpartage≠999 & r1rel=1
compute (hhpartage-dage) if rsex=2 & hhpartage≠-1 & hhpartage≠999 & r1rel=1
998: no partners ever if hetlife=0 & samlife=0
999: not answered if rafsmr=999 | rafsmr=996 | r1ptage=999 | r1ptage=997 | r1ptage=996

Note: Natsal-3 collected information on the ethnicity of partners to better describe sexual mixing

(N) **r1ethni** ethnic gp of partner

- 1: white / white British
- 2: Mixed ethnicity
- 3: asian / asian British
- 4: black / black British
- 5: Chinese / other
- 7: not sure
- 9: not answered
- 1: not applicable if scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0)

(N) **r1ethbl** partner's background

- 1: Caribbean
- 2: African
- 3: Other black
- 4: Not sure
- 9: not answered
- 1: not applicable if r1ethni=-1 | r1ethnic=1 | r1ethnic=2 | r1ethni=3 | r1ethni=5 | r1ethni=7

(R) (s10f) **r1relat** relationship with most recent partner

- 1: currently/ever married to each other
- 2: living together but never married
- 3: regular partners but not living together
- 4: not regular partners
- 9: not answered
- 1: not applicable if scelig=2 | ((het=0 | hetsc=0) & sam=0)

Note: In Natsal-3 the answer codes for r1meet were updated.

(U – see below) (N) **r1meet** where first met most recent partner

- 1: at school
- 2: at university/college
- 3: at/through work
- 4: in a pub/café/restaurant/bar/club
- 5: at a social event organised by friend(s)
- 6: through a society/sports club
- 7: on holiday/while travelling
- 8: in a public place (e.g. museum, park)
- 9: through a dating agency/chat line
- 10: have always known each other
- 11: other
- 12: at church
- 13: neighbour/lived locally/shared a flat
- 14: at another public place
- 15: arranged marriage
- 16: prostitute/red light district
- 17: through friends/relatives

DETAILED PARTNERSHIP HISTORY

99: not answered

-1: not applicable if scelig=2 | (het=0 | hetsc=0) & sam=0)

r1meet where met most recent partner – NEW CODES but question identical

1: at school

2: at university/college

3: at/through work

4: in a pub/bar/club/dance/disco

5: introduced by friends / family

6: through a society/sports club

7: on holiday/while travelling

8: internet dating website

9: other dating agency/personal ads

10: online but not through a dating website

11: have always known each other

12: neighbour/lived locally/shared a flat

13: arranged marriage

14: in a public place (e.g. museum, park)

15: sex worker / prostitute

16: other

99: not answered

-1: not applicable if scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0)

(U) (N) **r1live** where most recent partner lived when 1st met

1: same city/town

2: same region, different town

3: same country, different region

4: different country

5: unknown

9: not answered

(U) -1: not applicable if scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0)

-1: not applicable if scelig=2 | ((het=0 | hetsc=0) & sam=0)

(U) (N) **r1known** time between 1st meeting most recent partner and 1st sex

1: 24 hours or less

2: between 1 day and 1 week

3: between 1 week and 4 weeks

4: between 4 weeks and 6 months

5: between 6 months and 1 year

6: between 1 year and 5 years

7: between 5 years and 10 years

8: 10 years or more

99: not answered

(U) -1: not applicable if scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0) | R1Meet=11 | R1Meet=15

-1: not applicable if scelig=2 | ((het=0 | hetsc=0) & sam=0)

Note: Natsal-3 asked respondents whether they thought their partner had had sex with other people in the last 5 years and during the period of their relationship. There were 4 different questions depending on the date of the first and last sexual intercourse (r1pcona..r1pcond). This was only asked if they had had sex with that partner in the last 5 years. The 4 different versions of the question have been combined into one derived variable for each partner (r1pcon1, r1pcon2, r1pcon3, r1pcon4). The variables r1pcona-d are shown here for information, but are not kept in the final dataset. Instead the variables r1pcon1 etc should be used for analysis.

Note: r1pcona – 1st sex >5 years ago (R1FY/R1FM > 5 years ago) & relationship is ongoing (R1sexagn=yes / probably)

(N – shown for info only, see note above) **r1pcona** partner had sex with someone else in last 5 years

1: yes

DETAILED PARTNERSHIP HISTORY

2: probably
3: probably not
4: no
5: prefer not to say
9: not answered
-1: not applicable

Note: r1pconb 1st sex >5 years ago (R1FY/R1FM > 5 years ago) & last sex < 5 years ago (R1datey/R1dateM < 5 years ago) & relationship is not ongoing (R1sexagn=no / probably not / don't know) OR most recent sex >1 year ago (R1datey/R1dateM > 1 year ago)

(N – shown for info only, see note above) **r1pconb** partner had sex with someone else between 5 years ago and last sex

Note: r1pconc 1st sex < 5 years ago (R1FY/R1FM < 5 years ago) & relationship is ongoing (R1sexagn=yes / probably)

(N – shown for info only, see note above) **r1pconc** partner had sex with someone else since they first had sex

Note: r1pcond 1st sex <5 years ago (R1FY/R1FM < 5 years ago) & relationship is not ongoing (R1sexagn=no / probably not / don't know) OR most recent sex >1 year ago (R1datey/R1dateM > 1 year ago)

(N – shown for info only, see note above) **r1pcond** partner had sex with someone else between first and last sex

All same response options as for r1pcona

(ND) **r1pcon1** most recent partner had sex with someone else in last 5 years (between first and most recent sex with respondent)

1: yes if r1pcona=1 | r1pconb=1 | r1pconc=1 | r1pcond=1
2: probably if r1pcona=2 | r1pconb=2 | r1pconc=2 | r1pcond=2
3: probably not if r1pcona=3 | r1pconb=3 | r1pconc=3 | r1pcond=3
4: no if r1pcona=4 | r1pconb=4 | r1pconc=4 | r1pcond=4
5: prefer not to say if r1pcona=5 | r1pconb=5 | r1pconc=5 | r1pcond=5
9: not answered if r1pcona=9 | r1pconb=9 | r1pconc=9 | r1pcond=9
-1: not applicable if r1pcona=-1 & r1pconb=-1 & r1pconc=-1 & r1pcond=-1

Note: mrpos was added to Natsal-3 to look in more detail at oral sex only partnerships where both first and last sex where oral sex and the respondent reported having had anal or vaginal sex ever in the practices questions.

(N) **mrpos** was this an oral sex only partner

1: yes
2: no
9: not answered
-1: not applicable if scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0) | (rsex=2 & r1sex=2) | (lastvag≠1, 2, 3, 4, 5 & analsex≠1, 2, 3, 4, 5 & samayou≠1, 2, 3, 4, 5 & samahim≠1, 2, 3, 4, 5) | (r1condl=1 | r1condl=2) | (r1first=2 & (r1condf=1 | r1condf=2))

Note: Variables r1datem ... r1known and their codes also apply to 2 next most recent partners if the respondent reported 2+ partners in the last 5 years. Consequently there are the additional codes of -1 if tot5yrs < 2 for 2nd most recent partner and -1 if tot5yrs < 3 for 3rd most recent partner.

2nd most recent partner

r1datey2
r1datem2

r1sex2

(U) **r1condl2**

(N) **r1sxagn2**

DETAILED PARTNERSHIP HISTORY

(N) r1rel2
(*) r1first2
r1fy2
r1fm2
(N) firstdtp2
(N) firstdtp2ind
(N) lastdtp2
(N) lastdtp2ind
(N) dfover1p2
(N) dlover1p2
(N) dfover5p2
(N) dlover5p2
(N) r1mtha2
(N) r1mthb2
(U) r1condf2
(N) r1rel1st2
r1ptage2
rafs2mr
(N) ad_2mrp
(R) r1relat2
(N) r1ethni2
(N) r1ethbl2
r1meet2
r1live2
r1known2
(N) r1pcon2
(N) mrpos2

3rd most recent partner:

r1datey3
r1datem3
r1sex3
(U) r1condl3
(N) r1rel3
(N) r1sxagn3
(*) r1first3
r1fy3
r1fm3
(N) firstdtp3
(N) firstdtp3ind
(N) lastdtp3
(N) lastdtp3ind
(N) dfover1p3
(N) dlover1p3
(N) dfover5p3
(N) dlover5p3
(N) r1mtha3
(N) r1mthb3
(U) r1condf3
(N) r1rel1st3
r1ptage3
rafs3mr
(N) ad_3mrp
(R) r1relat3
(N) r1ethni3
(N) r1ethbl3
r1meet3
r1live3
r1known3
(N) r1pcon3

DETAILED PARTNERSHIP HISTORY

(N) mrpos3

Note: Variables beginning 'r1' and ending 4 are only asked if the respondent has had both heterosexual and homosexual partnerships, and details of both heterosexual and homosexual partnerships have not been reported as the most recent, 2nd most recent or 3rd most recent partnerships. This is determined by the responses to r1sex, r1sex2, and r1sex3. These variables are not limited to the last 5 years.

(N) **r1datey4** year of most recent sex with most recent other sexuality partner

9999: not answered

-1: not applicable *else if* r1sex4=-1

(N) **r1datem4** month of most recent sex with most recent other sexuality partner

1: January

2: February

3: March

4: April

5: May

6: June

7: July

8: August

9: September

10: October

11: November

12: December

(U) 13: I am unable to estimate the month

(U – code removed) 97: can't remember month *or if* r1datem4=98

99: not answered

-1: not applicable *else if* r1sex4=-1

(U) (N) **r1sex4** sex of most recent other sexuality partner

1: male

2: female

9: not answered

(U) -1: not applicable *if* scelign3=2 | (everhet=0 & eversam=2) | (hetlife=0 & samlife=0) | ((everhet=1 | everhet=9) & samlife=0) | (everhet=0 & (samlife>0 & samlife<=9999)) | ((everhet=1 & (samlife>0 & samlife<9999)) & (r1sex#9 & r1sex2#9 & r1sex3#9) & (((r1sex2#-1) & (r1sex3=-1) & (r1sex#r1sex2)) | ((r1sex3#-1) & ((r1sex#r1sex2) | (r1sex#r1sex3) | (r1sex2#r1sex3))))

-1: not applicable *if* scelig=2 | ((het=0 | hetsc=0) & sam=0) | (hetsc=1 & samlife=0) | (hetsc=0 & samlife>0 & samlife<9999) | ((hetsc=1 & (samlife>0 & samlife<9999)) & (r1sex#9 & r1sex2#9 & r1sex3#9) & (((r1sex2#-1) & (r1sex3=-1) & (r1sex#r1sex2)) | ((r1sex3#-1) & ((r1sex#r1sex2) | (r1sex#r1sex3) | (r1sex2#r1sex3))))

(U) (N) **r1condl4** condom used for most recent sex with most recent other sexuality partner

1: yes

2: no

(U) 3: only had oral sex on the most recent occasion *or if* (lastvag=7 & analsex=7 & (samahim=-1 | samahim=7) & (samayou=-1 | samayou=7) & (oralyou=1 to 6 | oralprt=1 to 6 | (oralyou=7 & oralprt=7 & (samoyou=1 to 6 | samopar=1 to 6)))

9: not answered

(U) -1: not applicable *if* (lastvag=7 & analsex=7 & (samahim=-1 | samahim=7) & (samayou=-1 | samayou=7)) | r1sex4=-1 | (rsex=2 & r1sex4=2)

-1: not applicable *else if* (r1sex4=-1) | (rsex=2 | r1sex4=2)

Note: questions added to establish i) relationship status at last sex and, ii) where sex occurred in the last year, whether they are likely to have sex again. In Natsal-2 this was assumed based on answers to certain questions.

(N) **r1rel4** relationship status at most recent sex

1: living together / married / civil partnership

2: steady relationship

DETAILED PARTNERSHIP HISTORY

3: used to be in steady relationship but not at time
4: known each other for a while but not in steady relationship
5: recently met
6: just met for first time
9: not answered
-1: not applicable *if* r1sex4=-1

(N) **r1sexagn4** likely to have sex again if most recent sex <1y ago

1: yes
2: probably
3: probably not
4: no
5: don't know
9: not answered
-1: not applicable *if* dlover1p4=1 | r1sex4=-1

Note: in Natsal-3, answer options for r1first4 were made clearer to reduce question mis-interpretation

(*) (N) **r1first4** most recent sex with most recent other sexuality partner was first occasion

(*) 1: yes, only had sex with him/her once 1: yes, was the first occasion

(*) 2: no, had sex on more than one occasion 2: no, was not the first occasion 9: not answered

-1: not applicable *else if* r1sex4=-1

(N) **r1fy4** year of 1st occasion with most recent other sexuality partner

9999: not answered

-1: not applicable *if* r1sex4=-1 | r1first4=1

(N) **r1fm4** month of 1st occasion with most recent other sexuality partner

1: January
2: February
3: March
4: April
5: May
6: June
7: July
8: August
9: September
10: October
11: November
12: December

(U) 13: unable to estimate the month

(U – code removed) 97: can't remember month *or if* r1fm4=98

99: not answered

(U) -1: not applicable *if* r1sex4=-1 | r1first4=1 | dfover5p4=1

-1: not applicable *if* r1sex4=-1 | r1first4=1

Note: firstdtp4 combines r1fy4 with r1fm4 to give a date format variable. If only had sex with partner on one occasion (r1first4=1) then uses r1datey4 and r1datem4 to derive date. All missing values are set to sysmiss. If greater understanding of the missing values is needed, the indicator variable firstdtp4ind can be used to see whether the date is missing because it was not given, or because the question was not applicable.

Note: if r1fm4/r1datem4 unknown, month set to June for firstdtp4

(ND) **firstdtp4** date of first sex with P4, most recent other sexuality partner
sysmis *if* missing

Note: lastdtp4 uses r1datey4 and r1datem4 to give a date format variable. All missing values are set to sysmiss. If greater understanding of the missing values is needed, the indicator variable lastdtp4ind can

DETAILED PARTNERSHIP HISTORY

be used to see whether the date is missing because it was not given, or because the question was not applicable. If r1datem4 unknown, month set to June.

(ND) **lastdtp4** date of most recent (last) sex with P4, most recent other sexuality partner
sysmis if missing

Note: dfover1p4 is derived from a combination of date of interview (dateoi), date of first sex with partner (r1fy4&r1fm4 or r1datey4&r1datem4) i.e. yes if r1fy4/r1fm4 were more than a year before date of interview, or if only had sex once and r1datey4/r1datem4 were more than a year before date of interview. Calculated to be yes if month of sex with partner unknown but possibly in last year (i.e. year is year before interview) and month of interview in July to December (i.e. there is greater likelihood that it was over 1 year ago than less than 1 year ago),

(ND) **dfover1p4** first sex with partner 4 occurred over 1 year ago

-1: not applicable if firstdtp4ind=-1

1: yes, over 1 year ago if (firstdtp4ind=1 & ((dateoi-1)>firstdtp4)) | (firstdtp4ind=2 & ((r1first4=1 & r1datey4>0 & (dateyoi-r1datey4=0)) | (r1first4=2 & r1fy4>0 & (dateyoi-r1fy4=0)) | (r1first4=1 & r1datey4>0 & (dateyoi-r1datey4=1) & range(datemoi 7.. 12)) | (r1first4=2 & r1fy4>0 & (dateyoi-r1fy4=1) & range(datemoi 7..12)))

2: no, in the last year if (firstdtp4ind=1 & ((dateoi-1)<=firstdtp4)) | (firstdtp4ind=2 & (r1first4=1 & r1datey4>0 & (dateyoi-r1datey4=0)) | (r1first4=2 & r1fy4>0 & (dateyoi-r1fy4=0)) | (r1first4=1 & r1datey4>0 & (dateyoi-r1datey4=1) & range(datemoi 1..6)) | (r1first4=2 & r1fy4>0 & (dateyoi-r1fy4=1) & range(datemoi 1..6)))

9: not answered

Similarly dfover5p4 is derived from a combination of date of interview (dateoi), date of first sex with partner (r1fy4&r1fm4 or r1datey4&r1datem4)

(ND) **dfover5p4** first sex with partner 4 occurred over 5 years ago

-1: not applicable if firstdtp4ind=-1

1: yes, over 5 years ago

2: no, in the last 5 years

9: not answered

Similarly dlover1p4 is derived from a combination of date of interview (dateoi) and date of most recent sex with partner (r1datey4 & r1datem4).

(ND) **dlover1p4** most recent (last) sex with P4 occurred over 1 year ago.

-1 not applicable if lastdtp4ind=-1

1: yes, over 1 year ago

2: no, in the last year

9: not answered

Similarly dlover5p4 is derived from a combination of date of interview (dateoi), date of most recent sex with partner (r1datey4 & r1datem4)

(ND) **dlover5p4** most recent (last) sex with partner 4 occurred over 5 years ago

-1: not applicable if firstdtp4ind=-1

1: yes, over 5 years ago

2: no, in the last 5 years

9: not answered

Note: In Natsal-3 if first sex was in same month as interview or previous calendar month, a question has been added to determine the length of the sexual relationship. This will allow better interpretation of the number of sex acts variable. The question asked depends on whether respondent is likely to have sex with partner again

(N) **r1mtha4** time since first sex

1: less than 7 days

2: 7 days – 2 weeks

DETAILED PARTNERSHIP HISTORY

3: 2-4 weeks
4: over 4 weeks
9: not answered
-1: not applicable if r1sex4=-1 | r1first4=1 | r1sexagn4=3 | r1sexagn4=4 | dfover1p4=1

(N) **r1mthb4** time between first and last sex

1: less than 7 days
2: 7 days – 2 weeks
3: 2-4 weeks
4: over 4 weeks
9: not answered
-1: not applicable if r1sex4=-1 | r1first4=1 | r1sexagn4=1 | r1sexagn4=2 | dfover1p4=1

Note: There is an additional answer option for r1cond1 in Natsal-2. Respondents can now specify that last sex was oral sex

(U) (N) **r1condf4** condom used on 1st occasion with most recent other sexuality partner

1: yes
2: no
(U) 3: only had oral sex on the first occasion or if (lastvag=7 & analsex=7 & (samahim=-1 | samahim=7) & (samayou=-1 | samayou=7) & (oralyou=1 to 6 | oralprt=1 to 6 | (oralyou=7 & oralprt=7 & (samoyou=1 to 6 | samopar=1 to 6)))
9: not answered
(U) -1: not applicable if (lastvag=7 & analsex=7 & (samahim=-1 | samahim=7) & (samayou=-1 | samayou=7)) | r1sex4=-1 | (rsex=2 & r1sex4=2) | r1first4=1
-1: not applicable if r1sex4=-1 | r1first4=1 | (rsex=2 & r1sex4=2)

Note: r1rel1st4 collects relationship status at first sex. This replaces the old variable r1relat

(N) **r1rel1s4** relationship status at most recent sex

1: living together / married / civil partnership
2: steady relationship
3: used to be in steady relationship but not at time
4: known each other for a while but not in steady relationship
5: recently met
6: just met for first time
9: not answered
-1: not applicable if r1sex=-1 | r1first4=1

(U) (N) **r1ptage4** age on 1st occasion with most recent other sexuality partner

(U) 996: NA – partners age at 1st sex <13 years

(U) 997: don't know 97: don't know

(U) 999: not answered 99: not answered

-1: not applicable if r1sex4=-1

(ND) **rafs4mr** respondent's age in completed years at 1st sex with most recent other sexuality partner
compute (((r1fy4-1900)*12) + r1fm4) - (((rdoby-1900)*12) + rdobm4) / 12 if (r1fy4≠-1 & r1fy4≠9999) & (r1fm4≠-1 & r1fm4≠97 & r1fm4≠99) & rdobm4≠99

else compute (((r1datey4-1900)*12) + r1datem4) - (((rdoby4-1900)*12) + rdobm4) / 12 if r1first4=1 & (r1datey4≠-1 & r1datey4≠9999) & (r1datem4≠-1 & r1datem4≠97 & r1datem4≠99) & rdobm4≠99

else compute r1fy - rdoby if (r1fy≠-1 & r1fy≠9999)

else compute r1datey-rdoby if r1first=1 & (r1datey≠-1 & r1datey≠9999)

(U) 999: not answered else if missing | rafs4mr<0

99: not answered else if missing | rafs4mr<0

-1: not applicable if scelig=2 | r1sex4=-1

(ND) **ad_4mrp** age difference between respondent & 4th most recent partner

compute (rafs4mr-r1ptage4) if rsex=1 & rafs4mr≠-1 & rafs4mr≠996 & rafs4mr≠999 & r1ptage4≠996 & r1ptage4≠997 & r1ptage4≠999

DETAILED PARTNERSHIP HISTORY

compute (r1ptage4-rafs4mr) if rsex=2 & rafs4mr#-1 & rafs4mr#996 & rafs4mr#999 & r1ptage4#996 & r1ptage4#997 & r1ptage4#999
999: not answered *if rafs4mr=999 | rafs4mr=996 | r1ptage4=999 | r1ptage4=997 | r1ptage4=996*
998: not applicable (only had partners of one sex, or already reported partners of both sexes in first 3 loops) else

Note: Natsal-3 collected more information on sexual mixing and asked about ethnicity

(N) **r1ethni4** ethnic gp of partner

- 1: white / white British
- 2: Mixed ethnicity
- 3: asian / asian British
- 4: black / black British
- 5: Chinese / other
- 7: not sure
- 9: not answered
- 1: not applicable *if r1sex4=-1*

(N) **r1ethblk4** partners background

- 1: Caribbean
- 2: African
- 3: Other black
- 4: Not sure
- 9: not answered
- 1: not applicable *if r1ethni=-1 | r1ethni=1 | r1ethni=2 | r1ethni=3 | r1ethni=5 | r1ethni=7*

(R) (N) **r1relat4** relationship with most recent other sexuality partner

- 1: currently/ever married
- 2: living together
- 3: regular partners but not living together
- 4: not regular partners
- 9: not answered
- 1: not applicable *if r1sex4=-1*

Note: In Natsal-3 the answer codes for r1meet4 were updated to be more relevant currently.

(U – see below) (N) **r1meet4** where first met most recent other sex partner

- 1: at school
- 2: at university/college
- 3: at/through work
- 4: in a pub/café/restaurant/bar/club
- 5: at a social event organised by friend(s)
- 6: through a society/sports club
- 7: on holiday/while travelling
- 8: in a public place (e.g. museum, park)
- 9: through a dating agency/chat line
- 10: have always known each other
- 11: other
- 12: at church
- 13: neighbour/lived locally/shared a flat
- 14: at another public place
- 15: arranged marriage
- 16: prostitute/red light district
- 17: through friends/relatives
- 99: not answered
- 1: not applicable *if r1sex4=-1 | r1relat=1 | r1relat=2 | r1relat=3*

r1meet4 where met most recent partner – NEW CODES but question identical

- 1: at school
- 2: at university/college

DETAILED PARTNERSHIP HISTORY

- 3: at/through work
- 4: in a pub/bar/club/dance/disco
- 5: introduced by friends / family
- 6: through a society/sports club
- 7: on holiday/while travelling
- 8: internet dating website
- 9: other dating agency/personal ads
- 10: online but not through a dating website
- 11: have always known each other
- 12: neighbour/lived locally/shared a flat
- 13: arranged marriage
- 14: in a public place (e.g. museum, park)
- 15: sex worker / prostitute
- 16: other
- 99: not answered
- 1: not applicable *if* r1sex4=-1

(U) (N) **r1live4** where most recent other sexuality partner lived when 1st met

- 1: same city/town
- 2: same region, different town
- 3: same country, different region
- 4: different country
- 9: not answered

(U) -1: not applicable *if* r1sex4=-1

-1: not applicable *if* r1sex4=-1 | r1relat=1 | r1relat=2 | r1relat=3

DETAILED PARTNERSHIP HISTORY

(U) (N) **r1known4** time between 1st meeting most recent other sexuality partner and 1st sex

1: 24 hours or less

2: between 1 day and 1 week

3: between 1 week and 4 weeks

4: between 4 weeks and 6 months

5: between 6 months and 1 year

6: between 1 year and 5 years

7: between 5 years and 10 years

8: 10 years or more

99: not answered

(U) -1: not applicable if r1sex4=-1 | r1meet4=11 | r1meet4=15

-1: not applicable if r1sex4=-1 | r1relat=1 | r1relat=2 | r1relat=3

Note: Natsal-3 asked respondents whether they thought their partners had had sex with other people during their relationship period in the last 5 years. The question asked depends on the date of the first and last sexual intercourse. The variable r1pcon4 should be used for analysis (see full note on page 91).

(N – shown for information only) r1pcon4 partner had sex with someone else in last 5 years

1: yes

2: probably

3: probably not

4: no

5: prefer not to say

9: not answered

-1: not applicable

(N – shown for information only) r1pconb4 partner had sex with someone else between 5 years ago and last sex

(N – shown for information only) r1pconc4 partner had sex with someone else since they first had sex

(N – shown for information only) r1pcond4 partner had sex with someone else between first and last sex

All same response options as for r1pcon4

(ND) **pr1con4** partner had sex with someone else in last 5 years (between first and most recent sex with respondent).

1: yes if r1pcon4=1 | r1pconb4=1 | r1pconc4=1 | r1pcond4=1

2: probably if r1pcon4=2 | r1pconb4=2 | r1pconc4=2 | r1pcond4=2

3: probably not if r1pcon4=3 | r1pconb4=3 | r1pconc4=3 | r1pcond4=3

4: no if r1pcon4=4 | r1pconb4=4 | r1pconc4=4 | r1pcond4=4

5: prefer not to say if r1pcon4=5 | r1pconb4=5 | r1pconc4=5 | r1pcond4=5

9: not answered if r1pcon4=9 | r1pconb4=9 | r1pconc4=9 | r1pcond4=9

-1: not applicable if r1pcon4=-1 & r1pconb4=-1 & r1pconc4=-1 & r1pcond4=-1

Note: mrpos was added to Natsal-3 to look in more detail at oral sex only partnerships.

(N) **mrpos4** was this an oral sex only partner

1: yes

2: no

9: not answered

-1: not applicable if r1sex4=-1

Note: in Natsal-2 assumptions were made re concurrency for analysis. In Natsal-3 respondents were asked a direct question about concurrency if they reported that one partnership ended the same month another began.

(N) **mrpolap** any concurrency between partnerships

1: yes – there was overlap

2: no

3: not sure

4: prefer not to say

9: not answered

-1: not applicable if tot5yr<2 | ((if lastdtp2≠-1 then lastdtp2≠firstdtp1) & (if lastdtp3≠-1 then lastdtp3≠firstdtp2) & (if lastdtp4≠-1 then lastdtp4≠firstdtp3))

DETAILED PARTNERSHIP HISTORY

Note: Variables `cmc_3mrf` etc. are used to create concurrency indicators. They express the date of first and last sex in terms of the number of months since the beginning of 1900

(R) (ND) **cmc_3mrf** date of first sex with 3rd most recent partner in last 5 years in CMCs
compute $((r1fy3-1900)*12) + r1fm3$ *if* $r1first3=2$ & $(r1fy3\neq -1$ & $r1fy3\neq 9999)$ & $(r1fm3\neq -1$ & $r1fm3\neq 97$ & $r1fm3\neq 99)$
else compute $((r1datey3-1900)*12) + r1datem3$ *if* $r1first3=1$ & $(r1datey3\neq -1$ & $r1datey3\neq 9999)$ & $(r1datem3\neq -1$ & $r1datem3\neq 97$ & $r1datem3\neq 99)$
 9999: not answered *else if* missing
 -1: not applicable *if* $tot5yrs < 3$

(R) (ND) **cmc_2mrf** date of first sex with 2nd most recent partner in last 5 years in CMCs
compute $((r1fy2-1900)*12) + r1fm2$ *if* $r1first2=2$ & $(r1fy2\neq -1$ & $r1fy2\neq 9999)$ & $(r1fm2\neq -1$ & $r1fm2\neq 97$ & $r1fm2\neq 99)$
else compute $((r1datey2-1900)*12) + r1datem2$ *if* $r1first2=1$ & $(r1datey2\neq -1$ & $r1datey2\neq 9999)$ & $(r1datem2\neq -1$ & $r1datem2\neq 97$ & $r1datem2\neq 99)$
 9999: not answered *else if* missing
 -1: not applicable *if* $tot5yrs < 2$

(R) (ND) **cmc_1mrf** date of first sex with most recent partner in CMCs
compute $((r1fy-1900)*12) + r1fm$ *if* $r1first=2$ & $(r1fy\neq -1$ & $r1fy\neq 9999)$ & $(r1fm\neq -1$ & $r1fm\neq 97$ & $r1fm\neq 99)$
else compute $((r1datey-1900)*12) + r1datem$ *if* $r1first=1$ & $(r1datey\neq -1$ & $r1datey\neq 9999)$ & $(r1datem\neq -1$ & $r1datem\neq 97$ & $r1datem\neq 99)$
 9999: not answered *else if* missing

(R) (ND) **cmc_3mrl** date of last sex with 3rd most recent partner in last 5 years in CMCs
compute $((r1datey3-1900)*12) + r1datem3$ *if* $r1datey3\neq -1$ & $(r1datey3\neq 9999$ & $r1datey3\neq -1)$ & $(r1datem3\neq -1$ & $r1datem3\neq 97$ & $r1datem3\neq 99)$
 9999: not answered *else if* missing
 -1: not applicable *if* $tot5yrs < 3$

(R) (ND) **cmc_2mrl** date of last sex with 2nd most recent partner in last 5 years in CMCs
compute $((r1datey2-1900)*12) + r1datem2$ *if* $(r1datey2\neq -1$ & $r1datey2\neq 9999)$ & $(r1datem2\neq -1$ & $r1datem2\neq 97$ & $r1datem2\neq 99)$
 9999: not answered *else if* missing
 -1: not applicable *if* $tot5yrs < 2$

(R) (ND) **cmc_1mrl** date of last sex with most recent partner in CMCs
compute $((r1datey-1900)*12) + r1datem$ *if* $(r1datey\neq -1$ & $\neq 9999)$ & $(r1datem\neq -1$ & $\neq 97$ & $\neq 99)$
 9999: not answered *else if* missing

DETAILED PARTNERSHIP HISTORY

The following notes apply to all concurrency variables.

- Notes:
1. Partnerships are considered concurrent if the month and year of 1st sex with the more recent partner is prior to the month and year of last sex with the former partner.
 2. Concurrency is assumed not to have taken place if the last sex with partner x occurred in the same month that the first sex with partner x+1 occurred.
 3. See notes in the Lancet paper by Johnson et al (2001) for details of the assumptions made where it could not be ascertained whether someone reporting 2 partners was serially monogamous or had concurrent partnerships

*Note: variables similar to **concurt1**..**concurt** and **concurtly** have been derived for Natsal-3. These variables are comparable but have been derived using the variables **lastdtp1** and **firstdtp1**, rather than **cmc_3mrf**..**cmc_1mrl** therefore they have been renamed and are shown below as new variables.*

(R) (ND) **concurt1** most recent and 2nd most recent partners in last 5 years were concurrent
 1: yes if (cmc_1mrf < cmc_2mrl) & (cmc_1mrf#-1 & cmc_1mrf#9999) & (cmc_2mrl#-1 & cmc_2mrl#9999) | (r1first=2 & ((cmc_1mrf=9999 & cmc_2mrl=9999) | (cmc_1mrf=9999 & (cmc_2mrl#-1 & cmc_2mrl#9999)) | (cmc_2mrl=9999 & (cmc_1mrf#-1 & cmc_1mrf#9999))) & (r1fy#9999 & r1datey2#9999) & (r1fy#r1datey2) & (r1fy<r1datey2))
 0: no if ((cmc_1mrf >= cmc_2mrl) & (cmc_1mrf#-1 & cmc_1mrf#9999) & (cmc_2mrl#-1 & cmc_2mrl#9999)) | (r1first=1 & (cmc_1mrf=9999 | cmc_2mrl=9999)) | (r1first=2 & ((cmc_1mrf=9999 & cmc_2mrl=9999) | (cmc_1mrf=9999 & (cmc_2mrl#-1 & cmc_2mrl#9999)) | (cmc_2mrl=9999 & (cmc_1mrf#-1 & cmc_1mrf#9999))) & (r1fy#9999 & r1datey2#9999) & (r1fy#r1datey2) & (r1fy>r1datey2))
 9: not answered else if tot5yrs>=2 & tot5yrs<9999
 -1: not applicable if tot5yrs<2

(N) **mrpconcl5y1** most recent and 2nd most recent partners in last 5 years were concurrent (*comparable with concurt1*)

1: yes if (((Firstdtp1ind=1 & Lastdtp2ind=1) | (Firstdtp1ind=2 & Lastdtp2ind=2)) & (Firstdtp1<Lastdtp2)) | (((Firstdtp1ind=1 & Lastdtp2ind=2) | (Firstdtp1ind=2 & Lastdtp2ind=1)) & R1first=2 & R1FY<R1dateY2)
 0: no if r1first=1 | (((Firstdtp1ind=1 & Lastdtp2ind=1) | (Firstdtp1ind=2 & Lastdtp2ind=2)) & (Firstdtp1>=Lastdtp2)) | (((Firstdtp1ind=1 & Lastdtp2ind=2) | (Firstdtp1ind=2 & Lastdtp2ind=1)) & R1first=2 & R1FY>=R1dateY2)
 9: not answered
 -1: not applicable if tot5yrs<2

(R) (ND) **concurt2** 2nd most recent and 3rd most recent partners in last 5 years were concurrent
 1: yes if (cmc_2mrf < cmc_3mrl) & (cmc_2mrf#-1 & #9999) & (cmc_3mrl#-1 & #9999) | (r1first2=2 & ((cmc_2mrf=9999 & cmc_3mrl=9999) | (cmc_2mrf=9999 & (cmc_3mrl#-1 & #9999)) | (cmc_3mrl=9999 & (cmc_2mrf#-1 & #9999))) & (r1fy2#9999 & r1datey3#9999) & (r1fy2#r1datey3) & (r1fy2<r1datey3))
 0: no if ((cmc_2mrf >= cmc_3mrl) & (cmc_2mrf#-1 & #9999) & (cmc_3mrl#-1 & #9999)) | (r1first2=1 & (cmc_2mrf=9999 | cmc_3mrl=9999)) | (r1first2=2 & ((cmc_2mrf=9999 & cmc_3mrl=9999) | (cmc_2mrf=9999 & (cmc_3mrl#-1 & #9999)) | (cmc_3mrl=9999 & (cmc_2mrf#-1 & #9999))) & (r1fy2#9999 & r1datey3#9999) & (r1fy2#r1datey3) & (r1fy2>r1datey3))
 9: not answered else if tot5yrs>=3 & <9999
 -1: not applicable if tot5yrs<3

(N) **mrpconcl5y2** 2nd most recent and 3rd most recent partners in last 5 years were concurrent (*comparable with concurt2*)

1: yes if (((Firstdtp2ind=1 & Lastdtp3ind=1) | (Firstdtp2ind=2 & Lastdtp3ind=2)) & (Firstdtp2<Lastdtp3)) | (((Firstdtp2ind=1 & Lastdtp3ind=2) | (Firstdtp2ind=2 & Lastdtp3ind=1)) & R1first2=2 & R1FY2<R1dateY3)
 0: no if r1first2=1 | (((Firstdtp2ind=1 & Lastdtp3ind=1) | (Firstdtp2ind=2 & Lastdtp3ind=2)) & Firstdtp2>=Lastdtp3) | (& ((Firstdtp2ind=1 & Lastdtp3ind=2) | (Firstdtp2ind=2 & Lastdtp3ind=1)) & R1first2=2 & R1FY2>=R1dateY3)
 9: not answered
 -1: not applicable if tot5yrs<3

(R) (D) **concurt** concurrent partnerships in the last 5 years

1: yes if (concurt1=1 | concurt2=1) | (concurt1=1 & concurt2=-1)
 0: no if concurt1=0 & concurt2=0
 9: not answered else if missing
 -1: not applicable if concurt1=-1 & concurt2=-1

DETAILED PARTNERSHIP HISTORY

(N) **mrpconcl5y** concurrent partnerships in the last 5 years (*comparable with concur*)1: yes if
MRPConcL5Y1=1 | MRPConcL5Y2=1
0: no if (MRPConcL5Y1=0 & MRPConcL5Y2=0) | (MRPConcL5Y1=0 & MRPConcL5Y2=-1) |
(MRPConcL5Y1=-1 & MRPConcL5Y2=0)
9: not answered
-1: not applicable if (MRPConcL5Y1=-1 & MRPConcL5Y2=-1)

Note: For those respondents unable to provide both date components but the partnerships did not occur in the same year, then concurrency is assumed not to be possible. For the remaining respondents who had at least 2 partners in the year prior to interview, then these partners are assumed to be distributed between the serial monogamous and concurrent categories with equal probabilities to those whose dates are known. See notes in the Lancet paper by Johnson et al (2001) for further details.

(U – uses **mrpconcl5y** instead of **concur**, see below) (D) **I5ypartn** partnership history in the last 5 years
0: 0 partners if tot5yrs=0
1: 1 partner (monogamous) if tot5yrs=1
2: 2+ partners (serial monogamous) if (tot5yrs>=2 & tot5yrs<9999) & concur=1
3: 2+ partners (concurrent) if (tot5yrs>=2 & tot5yrs<9999) & concur=0
4: 2+ partners (unknown concurrency) if (tot5yrs>=2 & tot5yrs<9999) & concur=9
9: not answered for variables used to derive 'concur' else if concur=9 | tot5yrs=9999
-1: not applicable if tot5yrs=-1

I5ypartn partnership history in the last 5 years
0: 0 partners if tot5yrs=0
1: 1 partner (monogamous) if tot5yrs=1
2: 2+ partners (serial monogamous) if mrpconcl5y=0
3: 2+ partners (concurrent) if mrpconcl5y=1
4: 2+ partners (unknown concurrency) if (tot5yrs>=2 & tot5yrs<9999) & mrpconcl5y=9
9: not answered else if mrpconcl5y=9 | tot5yrs=9999
-1: not applicable if tot5yrs=-1

(R) (ND) **concrtrly** concurrent partnerships in the year before interview
0: 2+ partners (serial monogamy) if (cmc_1mrf>=cmc_2mrl) if ((cmc_1mrf=-1 & cmc_1mrf#9999) & (cmc_2mrl=-1 & cmc_2mrl#9999) & (cmc_2mrl>=(cmc_doi-12))) | ((cmc_2mrl=9999 & (cmc_1mrf=-1 & cmc_1mrf#9999)) & (r1fy=-1 & r1fy#9999 & r1datey2=-1 & r1datey2#9999) & (r1first=2) & (r1datey2>=(dateyoi-1)) & (r1fy>r1datey2)) | ((cmc_1mrf=9999 & cmc_2mrl=9999) | (cmc_1mrf=9999 & (cmc_2mrl=-1 & cmc_2mrl#9999))) & (r1fy=-1 & r1fy#9999 & r1datey2=-1 & r1datey2#9999) & (r1first=2) & (r1datey2>=(dateyoi-1)) & (r1fy>r1datey2))
1: 2+ partners (concurrent) if (cmc_1mrf<cmc_2mrl) if ((cmc_1mrf=-1 & cmc_1mrf#9999) & (cmc_2mrl=-1 & cmc_2mrl#9999) & (cmc_2mrl>=(cmc_doi-12))) | ((cmc_2mrl=9999 & (cmc_1mrf=-1 & cmc_1mrf#9999)) & (r1fy=-1 & r1fy#9999 & r1datey2=-1 & r1datey2#9999) & (r1first=2) & (r1datey2>=(dateyoi-1)) & (r1fy<r1datey2)) | ((cmc_1mrf=9999 & cmc_2mrl=9999) | (cmc_1mrf=9999 & (cmc_2mrl=-1 & cmc_2mrl#9999))) & (r1fy=-1 & r1fy#9999 & r1datey2=-1 & r1datey2#9999) & (r1first=2) & (r1datey2>=(dateyoi-1)) & (r1fy<r1datey2)) | ((cmc_3mrl=9999 & (cmc_2mrl=-1 & cmc_2mrl#9999)) & (r1fy2=-1 & r1fy2#9999 & r1datey3=-1 & r1datey3#9999) & (r1first2=2) & (r1datey3>=(dateyoi-1)) & (r1fy2<r1datey3)) | (((cmc_2mrf=9999 & cmc_3mrl=9999) | (cmc_2mrf=9999 & (cmc_3mrl=-1 & cmc_3mrl#9999))) & (r1fy2=-1 & r1fy2#9999 & r1datey3=-1 & r1datey3#9999) & (r1first2=2) & (r1datey3>=(dateyoi-1)) & (r1fy2<r1datey3))
2: 2+ partners (unknown concurrency) else if (tot1yr>=2 & tot1yr<999)
9: not answered else if tot1yr=999
-1: not applicable if tot1yr<2

DETAILED PARTNERSHIP HISTORY

(N) **mrpconclly1** concurrency between 1st & 2nd most recent partners in the year before interview
1: yes if (Dlover1P2=2 & ((FirstdtP1ind=1 & LastdtP2ind=1) | (FirstdtP1ind=2 & LastdtP2ind=2)) & (FirstdtP1<LastdtP2)) | (Dlover1P2=2 & ((FirstdtP1ind=1 & LastdtP2ind=2) | (FirstdtP1ind=2 & LastdtP2ind=1)) & R1first=2 & R1FY<R1dateY2)

0: no if r1first=1 | dlover1p2=1 | dlover1p1=1 | (Dlover1P2=2 & ((FirstdtP1ind=1 & LastdtP2ind=1) | (FirstdtP1ind=2 & LastdtP2ind=2)) & FirstdtP1>=LastdtP2) | (Dlover1P2=2 & ((FirstdtP1ind=1 & LastdtP2ind=2) | (FirstdtP1ind=2 & LastdtP2ind=1)) & R1first=2 & R1FY>=R1dateY2)

9: not answered

-1: not applicable if tot1yr<2

(N) **mrpconclly2** concurrency between 2nd & 3rd most recent partners in the year before interview
1: yes if (Dlover1P3=2 & ((FirstdtP2ind=1 & LastdtP3ind=1) | (FirstdtP2ind=2 & LastdtP3ind=2)) & (FirstdtP2<LastdtP3)) | (Dlover1P3=2 & ((FirstdtP2ind=1 & LastdtP3ind=2) | (FirstdtP2ind=2 & LastdtP3ind=1)) & R1first2=2 & R1FY2<R1dateY3)

0: no if r1first2=1 | Dlover1P2=1 | Dlover1P3=1 | (Dlover1P3=2 & ((FirstdtP2ind=1 & LastdtP3ind=1) | (FirstdtP2ind=2 & LastdtP3ind=2)) & FirstdtP2>=LastdtP3) | (Dlover1P3=2 & ((FirstdtP2ind=1 & LastdtP3ind=2) | (FirstdtP2ind=2 & LastdtP3ind=1)) & R1first2=2 & R1FY2>=R1dateY3)

9: not answered

-1: not applicable if tot1yr<3

(N) **mrpconclly** concurrent partnerships in the year before interview (*comparable with conctrly*)

0: 2+ partners (serial monogamy) if & (MRPConcLY1=0 & MRPConcLY2=0) | (MRPConcLY1=0 & MRPConcLY2=-1) | (MRPConcLY1=-1 & MRPConcLY2=0)

1: 2+ partners (concurrent) if MRPConcLY1=1 | MRPConcLY2=1

2: 2+ partners (unknown concurrency) else if (tot1yr>=2 & tot1yr<999)

9: unknown number of partners in last year

-1: not applicable if (MRPConcLY1=-1 & MRPConcLY2=-1)

Note: For those respondents unable to provide both date components but the partnerships did not occur in the same year, then concurrency is assumed not to be possible. For the remaining respondents who had at least 2 partners in the year prior to interview, then these partners are assumed to be distributed between the serial monogamous and concurrent categories with equal probabilities to those whose dates are known.

(U – uses **mrpconclly** instead of **conctrly**, see below) (ND) **lypartn** partnership history in last year

0: 0 partners if tot1yr=0

1: 1 partner (monogamous) if tot1yr=1

2: 2+ partners (serial monogamous) if (tot1yr>=2 & tot1yr<999) & conctrly=0

3: 2+ partners (concurrent) if (tot1yr>=2 & tot1yr<999) & conctrly=1

4: 2+ partners (unknown concurrency) if (tot1yr>=2 & tot1yr<999) & conctrly=2

9: unknown number of partners in the last year else if tot1yr=999 | conctrly=9

-1: not applicable for: tot1yr if tot1yr=-1

lypartn partnership history in last year

0: 0 partners if tot1yr=0

1: 1 partner (monogamous) if tot1yr=1

2: 2+ partners (serial monogamous) if mrpconclly=0

3: 2+ partners (concurrent) if mrpconclly=1

4: 2+ partners (unknown concurrency) if mrpconclly=2

9: unknown number of partners in the last year else if tot1yr=999 | mrpconclly=9

-1: not applicable if tot1yr=-1

(ND) **lypartn2** concurrent partnerships in last year

0: no if lypartn=1 | lypartn=2

1: yes if lypartn=3

2: 2+ partners but unknown concurrency if lypartn=4

9: unknown number of partners in the last year if lypartn=9

-1: not applicable if lypartn=-1 | lypartn=0

TRAVEL AND SEX ABROAD

TRAVEL & SEX ABROAD

Note: In Natsal-3 these questions placed here to improve flow. Previously they appeared before the Most Recent Partners section

(M) (N) **travel** travelled outside of UK, last 5 years

1: yes

2: no

9: not answered

(U) -1: not applicable if *scelign3=2* | (*het5yrs=0* & *sam5yrs=0*)

-1: not applicable if *scelig=2* | (*het5yrs=0* & *sam5yrs=0*)

(M) (N) **trvlsex** any new sex partners while outside UK, last 5 years

1: yes

2: no or if (*sam5yrs=0* & *het5yrs=0*) | *travel=2*

9: not answered

(U) -1: not applicable if *scelign3=2*

-1: not applicable else if *scelig=2*

(M) (N) **nosextl** no. of new sex partners while outside UK, last 5 years

9999: not answered

-1: not applicable if *trvlsex=2* | *trvlsex=-1*

(ND) **nosextl2** no. of new sex partners while outside UK, last 5 years, grouped

0 if *trvlsex=2*

1 if *nosextl=1*

2: 2+ if *nosextl>=2* & *nosextl<9999*

9: not answered if *trvlsex=1* & *nosextl=9999*

-1: not applicable if *trvlsex=-1*

*Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code questions. The following binary variables were created directly from the question **cntry1** in Natsal-3, and derived (as shown) from **cntry101-cntry110** in Natsal-2.*

(ND) **uk1** origin of new partner(s) while outside UK, last 5 years: UK

-1: not applicable if *cntry101=-1* & *cntry102=-1* & *cntry103=-1* & *cntry104=-1* & *cntry105=-1* & *cntry106=-1* & *cntry107=-1* & *cntry108=-1* & *cntry109=-1* & *cntry110=-1*

1: yes if (*cntry101=0* | *cntry102=0* | *cntry103=0* | *cntry104=0* | *cntry105=0* | *cntry106=0* | *cntry107=0* | *cntry108=0* | *cntry109=0* | *cntry110=0*) | ((*cntry101=9* | *cntry102=9* | *cntry103=9* | *cntry104=9* | *cntry105=9* | *cntry106=9* | *cntry107=9* | *cntry108=9* | *cntry109=9* | *cntry110=9*) & (*cntry1x1=0* | *cntry1x2=0* | *cntry1x3=0*))

0: no else if *cntry101>=0* & *cntry101<=9*

9: not answered if missing

*Variables **othereu1** ...other cs1, are derived from **cntry101** ... **cntry110** and **cntry1x1** ...**cntry1x3**, and are coded in a similar way to **uk1**:*

(ND) **othereu1** origin of new partner(s) while outside UK, last 5 years: another EU country

(ND) **oznz1** origin of new partner(s) while outside UK, last 5 years: Australia/New Zealand

(ND) **northam1** origin of new partner(s) while outside UK, last 5 years: North America

(ND) **southam1** origin of new partner(s) while outside UK, last 5 years: South America

(ND) **caribbn1** origin of new partner(s) while outside UK, last 5 years: Caribbean

(ND) **asia1** origin of new partner(s) while outside UK, last 5 years: Asia

(ND) **mideast1** origin of new partner(s) while outside UK, last 5 years: Middle East, North Africa

(R) (ND) **ssafrca1** origin of new partner(s) while outside UK, last 5 years: Sub Saharan Africa

(N) **othafrica1** origin of new partner(s) while outside UK, last 5 years: African countries (other than North Africa)

(ND) **othercs1** origin of new partner(s) while outside UK, last 5 years: other country/region

(N) **dkcntry1** origin of new partner(s) while outside UK, last 5 years: don't know which country/region

TRAVEL AND SEX ABROAD

(U) (ND) **sexabsum** new partner(s) while outside UK, in last 5 years, grouped

0: no *if* trvlsex=2

(U) 1: yes, (all) from UK *if* trvlsex=1 & (othereu1#1 | oznz1#1 | northam1#1 | southam1#1 | carribn1#1 | asia1#1 | mideast1#1 | othafrika1#1 | othercs1#1)

1: yes, (all) from UK *if* trvlsex=1 & (othereu1#1 | oznz1#1 | northam1#1 | southam1#1 | carribn1#1 | asia1#1 | mideast1#1 | ssafrc1#1 | othercs1#1)

(U) 2: yes, 1+ new partner from a country other than UK *if* trvlsex=1 & (othereu=1 | oznz=1 | northam=1 | southam=1 | carribn=1 | asia=1 | mideast=1 | othafrika=1 | othercs=1)

2: yes, 1+ new partner from a country other than UK *if* trvlsex=1 & (othereu=1 | oznz=1 | northam=1 | southam=1 | carribn=1 | asia=1 | mideast=1 | ssafrc1=1 | othercs=1)

9: not answered *if* trvlsex=9

-1: not applicable *if* trvlsex=-1

(M) (N) **sexabrd** new partner while in UK but partner from outside UK, last 5 years

1: yes

2: no *or if* (sam5yrs=0 & het5yrs=0)

9: not answered

(U) -1: not applicable *else if* scelign3=2 -1: not applicable *else if* scelig=2

(M) (N) **nosexab** no. of new sex partners while in UK but partner(s) from outside UK, last 5 years

9999: not answered

(U) -1: not applicable *if* sexabrd=2 | scelig=2

-1: not applicable *if* sexabrd=2 | scelig=2

(M) (N) **nosexab2** no. of new partners while in UK but partner(s) from outside UK, last 5 years, grouped

0: 0 *if* sexabrd=2

1: 1 *if* nosexab=1

2: 2+ *if* nosexab>1 & nosexab<9999

9: not answered *if* nosexab=9999

-1: not applicable *if* sexabrd=-1

*Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code questions. The following binary variables were created directly from the question **cntry2** in Natsal-3, and derived (as shown) from **cntry201..cntry210** in Natsal-2.*

(ND) **othereu2** origin of new partner(s) in UK, last 5 years, another eu country

1: yes *if* (cntry201=1 | cntry202=1 | cntry203=1 | cntry204=1 | cntry205=1 | cntry206=1 | cntry207=1 | cntry208=1) | ((cntry201=9 | cntry202=9 | cntry203=9 | cntry204=9 | cntry205=9 | cntry206=9 | cntry207=9 | cntry208=9) & cntry2x1=1) |

0: no *else if* cntry201>=0 & cntry201<=9

9: not answered *if* missing

-1: not applicable *if* cntry201=-1 & cntry202=-1 & cntry203=-1 & cntry204=-1 & cntry205=-1 & cntry206=-1 & cntry207=-1 & cntry208=-1

Variables oznz2... dkcrs are derived from cntry201 ... cntry208, and cntry2x1, and are coded in a similar way to othereu2:

(ND) **oznz2** origin of new partner(s) in UK, last 5 years: Australia/New Zealand

(ND) **northam2** origin of new partner(s) in UK, last 5 years: North America

(ND) **southam2** origin of new partner(s) in UK, last 5 years: South America

(ND) **caribbn2** origin of new partner(s) in UK, last 5 years: the Caribbean

(ND) **asia2** origin of new partner(s) in UK, last 5 years: Asia

(ND) **mideast2** origin of new partner(s) in UK, last 5 years: the Middle East, North Africa

(R) (ND) **ssafrc2** origin of new partner(s) in UK, last 5 years: Sub-Saharan Africa

(N) **othafrika2** origin of new partner(s) in UK, last 5 years: African countries (other than North Africa)

(ND) **othercs2** origin of new partner(s) in UK, last 5 years: another country

(R) (ND) **dkcrs2** origin of new partner(s) in UK in last 5 years: can't remember/don't know

(N) **dkctry2** origin of new partner(s) in UK in last 5 years: don't know which region or country

TRAVEL AND SEX ABROAD

Note: Variable forsexsp combines responses to questions about sex while the respondent was abroad (i.e. otheru1....) with responses to questions about sex in Britain with people from other countries (i.e. otheru2....) to derive a variable for whether/not reported new partner(s) from outside UK in last 5 years

(U) (ND) **forsexsp** new sexual partner(s) from outside UK, last 5 years

(U) 1: yes if sexabsum=2 | (otheru2=1 | oznz2=1 | northam2=1 | southam2=1 | carribn2=1 | asia2=1 | mideast2=1 | othafrika2=1 | othercs2=1)

1: yes if sexabsum=2 | (otheru2=1 | oznz2=1 | northam2=1 | southam2=1 | carribn2=1 | asia2=1 | mideast2=1 | ssafrica2=1 | othercs2=1)

0: no if tot5yrs>=1 & tot5yrs<9999

9: not answered if tot5yrs=9999

-1: not applicable if tot5yrs=0 | tot5yrs=-1

PAYING FOR SEX

NONCONSENSUAL SEX

Note: this section was added in Natsal-3

(N) **viotry** anyone tried to have sex with them against their will, aged 13+

- 1: yes
- 2: no
- 3: don't know
- 9: not answered
- 1: not applicable *if* scelign3=2 | askcasi=2

(N) **viosuc** anyone had sex with them against their will, aged 13+

- 1: yes
- 2: no *or if* viotry=2
- 3: don't know *or if* viotry=3
- 9: not answered
- 1: not applicable *if* scelign3=2 | askcasi=2

Note: the variables viotry2 and viosuc2 were used in analysis for Macdowall et al (2013) Lifetime prevalence, associated factors, and circumstances of non-volitional sex in women and men in Britain: findings from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3). These variables recode those with no lifetime partners (who weren't asked the question) as 'no' rather than excluding them.

(N) **viotry2** anyone tried to have sex with them against their will, aged 13+

- 1: yes *if* viotry=1
- 0: no *if* viotry=2 | viotry=-1
- 9: not answered *if* viotry=3 | viotry=9

(N) **viosuc2** anyone had sex with them against their will, aged 13+

- 1: yes *if* viosuc=1
- 0: no *if* viosuc=2 | viosuc=-1
- 9: not answered *if* viosuc=3 | viosuc=9

(N) **viowhen** age when last had sex against will, aged 13+

- 9999: not answered
- 1: not applicable *if* scelign3=2 | askcasi=2 | viosuc=2

(N) **viowho** relationship with person who had sex against will, aged 13+

- 1: someone were/had been in relationship with
- 2: family member or friend
- 3: known but not family member or friend
- 4: someone didn't know
- 5: other
- 9: not answered
- 1: not applicable *if* scelign3=2 | askcasi=2 | viosuc=2

(N) **viotalk** talk to anyone about sex against will, aged 13+

- 1: yes *or if* viopol=1
- 2: no
- 9: not answered
- 1: not applicable *if* scelign3=2 | askcasi=2 | viosuc=2

(N) **viopol** talk to police about sex against will, aged 13+

- 1: yes
- 2: no
- 9: not answered
- 1: not applicable *if* scelign3=2 | askcasi=2 | viosuc=2

PAYING FOR SEX

Note: in Natsal-3 these questions were asked of both men and women. In Natsal-3 these were not asked of people reporting never having had sex.

Note: in Natsal-2 & Natsal-1 These questions were only asked to men

Note: In Natsal-1, respondents with missing data are recoded to 'no' if Het=0 | LastVag=7

(U) (s13am) **everpd** ever paid money for het. sex

1: yes

(U) 2: no or if everhet=0

2: no or if everpd=9 & (rsex=1 & (het=0 | hetsc=0))

9: not answered

(R) -1: not applicable else if scelig=2 | rsex=2

(U) (s13bm) **lastpay** when last paid money for het. sex

1: in last 7 days

2: at least 7 days ago but less than 4 weeks

3: at least 4 weeks ago but less than 1 year

4: at least 1 year ago but less than 5 years

5: at least 5 years

9: not answered

(U) -1: not applicable if everpd=2 -1: not applicable else if scelig=2 | rsex=2 | everpd=2

(U) (rs13c1m) **nopaid** total no. of different women / men paid money to have sex with

0 if rsex=1 & (het=0 | hetsc=0 | everpd=2)

999: not answered

(U) -1: not applicable if everpd=2

-1: not applicable rsex=2 else if scelig=2

(U) (N) **payprev** het. sex partner total includes women / men paid for sex

1: yes, included

2: no, not included

9: not answered

(U) -1: not applicable if everpd=2

-1: not applicable if rsex=2 | everpd=2 or else if scelig=2

Note: Questions about paying for same-sex sex were only asked of men

Note: as above, all instances of 'hom' in variable names were changed to sam in Natsal-3

sampaid (s14am) **hompaid** ever paid money for hom. sex

1: yes

2: no or if rsex=1 & (sam=0 | eversam=2)

9: not answered

(U) -1: not applicable if rsex=2

-1: not applicable if rsex=2 | else if scelig=2

samplst (s14bm) **homplst** when last paid money for hom. sex

1: in last 7 days

2: at least 7 days ago but less than 4 weeks

3: at least 4 weeks ago but less than 1 year

4: at least 1 year ago but less than 5 years

5: at least 5 years

9: not answered

-1: not applicable if sampaid=-1 | sampaid=2

sampdno (rs14c1m) **hompdno** total no. of different men paid money to have sex with

0 if rsex=1 & sampaid=2

999: not answered

PAYING FOR SEX

-1: not applicable *if* sampaid=-1 | sampaid=2

sampdp (N) **hompdp** hom. sex partner total includes men paid for sex

1: yes, included

2: no, not included

9: not answered

-1: not applicable *if* sampaid=-1 | sampaid=2

Note: in Natsal-3 additional questions were asked about paying for sex abroad if reported paying for sex at least once (everpd=yes | sampaid=yes).

(N) **payabrd** ever paid money for sex outside UK

1: yes

2: no *or if* everpd=2 & ((rsex=1 & sampaid=2) | rsex=2)

9: not answered

*Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code questions. The following binary variables were created directly from the question **payabwhr**. All variables have the same codes as **pothereu**.*

(N) **pothereu** Region paid for sex outside UK: another EU country

1: yes

0: no

9: not answered

-1: not applicable *if* payabrd=2

And similarly...

(N) **poznsz** Region paid for sex outside UK: Australia/New Zealand

(N) **pnortham** Region paid for sex outside UK: North America

(N) **psoutham** Region paid for sex outside UK: South America

(N) **pcaribbn** Region paid for sex outside UK: the Caribbean

(N) **pasia** Region paid for sex outside UK: Asia

(N) **pmideast** Region paid for sex outside UK: the Middle East, North Africa

(N) **pothafrica** Region paid for sex outside UK: African countries (other than North Africa)

(N) **pothercs** Region paid for sex outside UK: another country

(N) **pdcktry** Region paid for sex outside UK: don't know which region or country

(U) (ND) **evrpdany** paid money for het./hom. sex, ever

(U) 1: yes *if* everpd=1 | sampaid=1

1: yes *if* rsex=1 & (everpd=1 | sampaid=1)

(U) 2: no *else if* everpd=2 | sampaid=2

2: no *else if* rsex=1 & everpd=2 | sampaid=2 | ((het=0 | hetsc=0) & sam=0)

9: not answered *if* everpd=9 | sampaid=9

(R) -1: not applicable *if* scelig=2 | rsex=2

(U) (ND) **paid5yrs** paid money for het./hom. sex, last 5 years

(U) 1: yes *if* (lastpay>=1 & <=4) | (samplst>=1 & samplst<=4)

1: yes *if* rsex=1 & (lastpay>=1 & <=4) | (samplst>=1 & samplst<=4)

(U) 0: no *else if* evrpdany=2 | ((lastpay=5 & (samplst#9)) | (samplst=5 & (lastpay#9)))

0: no *else if* rsex=1 & (evrpdany=2 | ((lastpay=5 & (samplst#9)) | (samplst=5 & (lastpay#9))))

9: not answered *if* lastpay=9 | samplst=9

(R) -1: not applicable *if* lastpay=-1 & samplst=-1

(U) (ND) **paid1yr** paid for het./hom. sex, last year

(U) 1: yes *if* (lastpay>=1 & <=3) | (samplst>=1 & samplst<=3)

1: yes *if* rsex=1 & (lastpay>=1 & <=3) | (samplst>=1 & samplst<=3)

(U) 0: no *else if* (evrpdany=2 | ((lastpay=4 | 5 & (samplst#9)) | (samplst=4 | 5 & (lastpay#9))))

0: no *else if* rsex=1 & (evrpdany=2 | ((lastpay=4 | 5 & (samplst#9)) | (samplst=4 | 5 & (lastpay#9))))

9: not answered *if* (lastpay=9 | samplst=9)

(R) -1: not applicable *if* lastpay=-1 & samplst=-1

PREGNANCY HISTORY

HISTORY OF PREGNANCIES

Note: This was a newly developed section in Natsal-3. Asked only of women. The information collected in this section was previously collected in the gynaecological history section in Natsal-3.

(N) **everpreg** Ever been pregnant

- 1: yes
- 2: no or if (rsex=2 & (scelign3=2 | askcasi=2 | lastvag=7))
- 9: not answered
- 1: not applicable if rsex=1

(N) **nopreg** no. times pregnant

- 0 if everpreg=2
- 99 'Not answered'
- 1: not applicable if rsex=1 | (rsex=2 & (scelign3=2 | askcasi=2 | lastvag=7))

Note: the loop zpregoc! (where !=loop number) to abweek is repeated for each pregnancy, up to a maximum of 20 loops. Variables are shown for pregnancy 1 below. Variable names are suffixed with the loop (pregnancy) number for pregnancies 2-20.

Note: xpregoc! is a derived variable which combines data from questions pregout and pregolst in the questionnaire.

(N D) **zpregoc1** outcome of 1st preg

- 1: miscarriage
- 2: abortion
- 3: stillbirth
- 4: single live birth
- 5: multiple live birth
- 6: currently pregnant
- 7: never been pregnant if everpreg=2
- 9: not answered
- 1: not applicable if (rsex=1) or for if !=nopreg (loops2-20).

Note: at later pregnancies, also coded as not applicable if have been pregnant, but number of pregnancies is less than loop number for zpregoc.

(N) **pregml** no. children gave birth to in multi-birth

- 999: not answered
- 1: not applicable if rsex=1 | (rsex=2 & (zpregoc1≠5 & zpregoc1≠9))

c(N) **lbsex** gender of single live birth

- 1: male
- 2: female
- 9: not answered
- 1: not applicable if rsex=1 | (rsex=2 & (zpregoc1≠4 & zpregoc1≠9))

PREGNANCY HISTORY

(N) **lbmonth** month single live birth born

-1: not applicable *if* rsex=1 | (rsex=2 & zpregoc1#4 & zpregoc1#9))

1: january

2: february

3: march

4: april

5: may

6: june

7: july

8: august

9: september

10: october

11: november

12: december

99: not answered

(N) **lbyear** year single live birth born

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#4 & zpregoc1#9))

9999: not answered

(N) **lb1sex1** gender 1st multiple birth child (pregnancy 1)

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#5 & zpregoc1#9))

1: male

2: female

(N) **lb1sex2** gender 2nd multiple birth child (pregnancy 1)

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#5 & zpregoc1#9))

1: male

2: female

(N) **lb1sex3** gender 3rd multiple birth child (pregnancy 1)

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#5 & zpregoc1#9))

1: male

2: female

(N) **lb1sex4** gender 4th multiple birth child (pregnancy 1)

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#5 & zpregoc1#9))

1: male

2: female

(N) **lb1sex5** gender 5th multiple birth child (pregnancy 1)

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#5 & zpregoc1#9))

1: male

2: female

(N) **lb1sex6** gender 6th multiple birth child (pregnancy 1)

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#5 & zpregoc1#9))

1: male

2: female

(N) **lb1sex7** gender 7th multiple birth child (pregnancy 1)

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#5 & zpregoc1#9))

1: male

2: female

(N) **lb1sex8** gender 8th multiple birth child (pregnancy 1)

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#5 & zpregoc1#9))

1: male

2: female

PREGNANCY HISTORY

(N) **lbmmth** month multiple live births born

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#5 & zpregoc1#9))

1: january

2: february

3: march

4: april

5: may

6: june

7: july

8: august

9: september

10: october

11: november

12: December

99: not answered

(N) **lbmyea** year multiple live births born

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#5 & zpregoc1#9))

9999: not answered

The variables alldob1 to alldob20 combine lbyear with lbmyea for each of the 20 loops. All missing values are set to sysmiss. If greater understanding of the missing values is needed, the indicator variables allyrs1 to allyrs20 can be used to see whether the date is missing because it was not given, or it is not applicable.

Note: only month and year collected, day set to 01 in alldob1-alldob20. If month unknown, month set to 06.

(ND) **Alldob1** DOB of 1st single/multiple live birth

Similarly for up to 20 pregnancies:

(ND) **Alldob2** DOB of 2nd single/multiple live birth

(ND) **Alldob3** DOB of 3rd single/multiple live birth

(ND) **Alldob4** DOB of 4th single/multiple live birth

(ND) **Alldob5** DOB of 5th single/multiple live birth

(ND) **Alldob6** DOB of 6th single/multiple live birth

(ND) **Alldob7** DOB of 7th single/multiple live birth

(ND) **Alldob8** DOB of 8th single/multiple live birth

(ND) **Alldob9** DOB of 9th single/multiple live birth

(ND) **Alldob10** DOB of 10th single/multiple live birth

(ND) **Alldob11** DOB of 11th single/multiple live birth

(ND) **Alldob12** DOB of 12th single/multiple live birth

(ND) **Alldob13** DOB of 13th single/multiple live birth

(ND) **Alldob14** DOB of 14th single/multiple live birth

(ND) **Alldob15** DOB of 15th single/multiple live birth

(ND) **Alldob16** DOB of 16th single/multiple live birth

(ND) **Alldob17** DOB of 17th single/multiple live birth

(ND) **Alldob18** DOB of 18th single/multiple live birth

(ND) **Alldob19** DOB of 19th single/multiple live birth

(ND) **Alldob20** DOB of 20th single/multiple live birth

(N) **prgage** age at miscarriage, abortion or still birth

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#1 | zpregoc1#2 | zpregoc1#3))

999: not answered

(N) **abweek** no. weeks pregnant at abortion

-1: not applicable *if* rsex=1 | (rsex=2 & (zpregoc1#2))

99: not answered

PREGNANCY HISTORY

(ND) **pregomisc** Ever had a pregnancy that ended in miscarriage
1 yes *if any of zpregoc1 to zpregoc20=1*
2 no *if zpregoc1 to zpregoc20 ≠ 1 & ≠ 9 & ≠ -1*
9 not answered
-1 not applicable *if zpregoc1=-1*

(ND) **prmiscagf** Age at first miscarriage
99 not answered
-1 not applicable *if pregomisc=2 | pregomisc=-1*

(ND) **pregosb** Ever had a pregnancy that ended in a stillbirth
1 yes *if any of zpregoc1 to zpregoc20=3*
2 no *if zpregoc1 to zpregoc20 ≠ 3 & ≠ 9 & ≠ -1*
9 not answered
-1 not applicable *if zpregoc1=-1*

(ND) **prsbagf** Age at first stillbirth
99 not answered
-1 not applicable *if pregosb=2 | pregosb=-1*

(ND) **pregoab** Ever had a pregnancy that ended in an abortion
1: yes *if any of zpregoc1 to zpregoc20=2*
2: no *if zpregoc1 to zpregoc20 ≠ 2 & ≠ 9 & ≠ -1*
9: not answered
-1: not applicable *if zpregoc1=-1*

Note: prabagf is similar to the Natsal-2 variable abageall (see p118) but the variable name has been changed to reflect changes to the questions about pregnancies in Natsal-3.

(ND) **prabagf** Age at first abortion
99: not answered
-1: not applicable *if pregoab=2 | pregoab=-1*

Note: manyab2 is similar to the Natsal-2 variable manyab (see p117) but the variable name has been changed to reflect changes to the questions about pregnancies in Natsal-3.

(ND) **manyab2** Number of abortions, ever
99: not answered
-1: not applicable *if pregoab=2 | pregoab=-1*

(ND) **manyab2gp** Number of abortions, ever, grouped
0 *if manyab2=0*
1 *if manyab2=1*
2: 2+ *if manyab2>=2 & manyab2<=99*
9: not answered *if manyab2=99*
-1: not applicable *if manyab2=-1*

Note: prabagl is similar to the Natsal-2 variable abageal2 (see p118) but the variable name has been changed to reflect changes to the questions about pregnancies in Natsal-3.

(ND) **prabagl** Age at last abortion
99: not answered
-1: not applicable *if pregoab=2 | pregoab=-1*

Note: abort18_2 is similar to the Natsal-2 variable abort18 (see p118118) but the variable name has been changed to reflect changes to the questions about pregnancies in Natsal-3. As in Natsal-2, denominator is women aged 18+

PREGNANCY HISTORY

(ND) **abort18_2** Had an abortion before age 18yrs

0: No *if* dage>=18 & (pregoab=2 | (prabagf>=18 & prabagf<99))

1: Yes *if* dage>=18 & (prabagf>0 & prabagf<18)

9: Not answered

-1: not applicable *if* pregoab=-1 | dage<18

Note: abort20_2 is similar to the Natsal-2 variable abort20 (see p118118) but the variable name has been changed to reflect changes to the questions about pregnancies in Natsal-3. As in Natsal-2, denominator is women aged 20+

(ND) **abort20_2** Had an abortion before age 20yrs

0: No *if* dage>=20 & (pregoab=2 | (prabagf>=20 & prabagf<99))

1: Yes *if* dage>=20 & (prabagf>0 & prabagf<20)

9: Not answered

-1: not applicable *if* pregoab=-1 | dage<20

Note: abort15 is an approximate variable for abortion in the last 5 years. We only have participants' age at abortion, therefore for some participants we are unable to tell whether the abortion was in the last 5 years, or just over 5 years ago. This variable only codes people as 'yes' if we know that it was definitely in the past 5 years, therefore it is an underestimate.

(ND) **abort15** had an abortion in last 5 years (approximate)

-1: not applicable *if* rsex=1

1: yes *if* (prabagl>0 & prabagl<99) & (dage-prabagl<=4)

0: no *if* rsex=2 & (pregoab=2 | (pregoab=1 & prabagl≠99 & (dage-prabagl>4)))

9: not answered *if* prabagl=99

(ND) **agfstpreg** Age at first pregnancy outcome

copy prmiscagf *if* zpregoc1=1

copy prabagf *if* zpregoc1=2

copy prsbagf *if* zpregoc1=3

copy age1ch (see p124) *if* zpregoc1=4 | zpregoc1=5

copy dage *if* zpregoc1=6

99: not answered

-1: not applicable *if* zpregoc1=-1 | zpregoc1=7

(ND) **pregob418** pregnancy outcome before 18

0: no *if* rsex=2 & dage>=18 & ((agfstpreg>=18 & agfstpreg<56) | everpreg=2)

1: yes *if* rsex=2 & dage>=18 & ((agfstpreg>=13 & agfstpreg<18)

-1: not applicable *if* rsex=1 | dage<18

9: not answered *else if* agfstpreg=99

(D) **lastpreg** outcome of most recent pregnancy

-1: not applicable *if* everpreg=2 | rsex=1 |

1: live birth *if* (nopreg!= & zpregoc!=4 | zpregoc!=5)

2: stillbirth *if* (nopreg!= & zpregoc!=3)

3: miscarriage *if* (nopreg!= & zpregoc!=1)

4: abortion *if* (nopreg!= & zpregoc!=2)

5: currently pregnant *if* (nopreg!= & zpregoc!=6)

9: not answered *if* (nopreg!= & zpregoc!=9) | nopreg=9

Note: dtlastpry and dtlastprm come directly from a program variable used in the routing. However, this is an approximate variable and may be incorrect for some respondents e.g. if their pregnancy history data was edited. Therefore it should be treated with caution, and not assumed to be 100% accurate.

(N) **dtlastpry** year of last pregnancy

sysmis *if* missing

PREGNANCY HISTORY

(N) **dtlastprm** month of last pregnancy

- 1: january
- 2: february
- 3: march
- 4: april
- 5: may
- 6: june
- 7: july
- 8: august
- 9: september
- 10: october
- 11: november
- 12: December
- 99: not answered
- 1: not applicable *if* rsex=1 | (rsex=2 & everpreg=2)

(N) **dtlastpr** date of last pregnancy (day set to 01 for all cases)

Note: Antenat was previously in the face to face section of the questionnaire after the general health questions. Eligibility changed for this question in Natsal-3 and it was only asked of women who may have been pregnant in the last 5 years.

(U) (M) (N) **antenat2** ante-natal visit to hospital/GP surgery in last 5 years

- 1: not applicable *if* rsex=1 | dage>50 | dtlastpr>dateoi – 5 years
- 1: yes
- 2: no
- 9: not answered

PREGNANCY HISTORY

GYNAECOLOGICAL HISTORY

Note: in Natsal-3 this section was completely remodelled and information was collected on each pregnancy. These questions were only asked of women. This new section is called History of Pregnancies. Some previously directly-asked variables from this section were derived using the data collected in the history of pregnancies section.

Note: These questions were only asked to women

(R) (s13aw) **miscarr** ever had a pregnancy that ended in miscarriage or still birth

- 1: yes
- 2: no or if rsex=2 & (het=0 | hetsc=0 | lastvag=7)
- 9: not answered
- 1: not applicable if rsex=1 | scelig=2

(R) (N) **whnmisc** (last) miscarriage or still birth

- 1: in last year
- 2: between 1 and 5 years
- 3: between 5 and 10 years
- 4: longer than at least 10 years
- 9: not answered
- 1: not applicable if scelig=2 | rsex=1 | miscarr=2

Note: Variable 'whnmisc2' was derived for consistency between Natsal-1 and Natsal-2.

(R) (s13bw U) **whnmisc2** last miscarriage or still birth

- 1: in last year
- 2: between 1 and 5 years
- 3: longer than 5 years if whnmisc=3 | whnmisc=4
- 9: not answered
- 1: not applicable if scelig=2 | rsex=1 | miscarr=2

(R) (sb5 D) **mcsbl5y** miscarriage/still birth, last 5 years

- 0: no if miscarr=2 | (whnmisc=3 | whnmisc=4)
- 1: yes if whnmisc=1 | whnmisc=2
- 9: not answered if whnmisc=9
- 1: not applicable if miscarr=-1

(R) (s14aw) **abort** had an abortion, ever

- 1: yes
- 2: no or if rsex=2 & (het=0 | hetsc=0 | lastvag=7)
- 9: not answered
- 1: not applicable if rsex=1 | scelig=2

(R) (N) **manyab** no. of abortions, ever

- 0 if rsex=2 & (het=0 | hetsc=0 | abort=2)
- 99: not answered
- 1: not applicable if scelig=2 | rsex=1

(R) (N) **ageab** age at abortion (if had one ever)

- 99: not answered
- 1: not applicable if scelig=2 | rsex=1 | (rsex=2 & (manyab=0 | manyab>1))

Notes: 1. There was a problem with the wording of the question 'ageabfir' because the question asked 'What age were you when you had the termination?', i.e. it doesn't specify the first termination. However, the next question, 'ageablas' is clearer so hopefully this would have clarified the meaning of 'ageabfir'.

2. In the raw dataset the age for 'ageabfir' was later than 'ageablas' for 3 cases, so these ages were swapped for these 3 cases (serial numbers: 1087281, 1369231, 2316351)

PREGNANCY HISTORY

(R) (N) **ageabfir** age at 1st abortion (if had 2+ ever)

99: not answered

-1: not applicable *if* scelig=2 | rsex=1 | (rsex=2 & (manyab=0 | manyab=1))

(R) (N) **ageablas** age at last abortion (if had 2+ ever)

99: not answered

-1: not applicable *if* scelig=2 | rsex=1 | (rsex=2 & (manyab=0 | manyab=1))

(R) (ND) **ageabal1** age at 1st abortion

copy ageab *if* manyab=1 & (ageab≠-1 & ageab<99)

else ageabfir *if* manyab>1 & manyab<99 & (ageabfir≠-1 & ageabfir<99)

99: not answered *else if* rsex=2 & (abort=1 | abort=9)

-1: not applicable *if* rsex=1 | (rsex=2 & abort=2)

(R) (ND) **ageabal2** age at last abortion

copy ageabal2 *if* manyab=1 & (ageab≠-1 & ageab<99)

else ageablas *if* manyab>1 & <99 & (ageablas≠-1 & ageablas<99)

99: not answered *else if* rsex=2 & (abort=1 | abort=9)

-1: not applicable *if* rsex=1 | (rsex=2 & abort=2)

(R) (ND) **abort18** had an abortion before age 18

0: no *if* dage>=18 & abort=2 | (manyab=1 & (ageab≠-1 & ageab>=18 & ageab<99)) | (manyab>1 & manyab<99 & (ageabfir≠-1 & ageabfir>=18 & ageabfir<99))

1: yes *if* (manyab=1 & (ageab≠-1 & ageab<18)) | (manyab>1 & manyab<99 & (ageabfir≠-1 & ageabfir<18))

9: not answered *else if* missing

-1: not applicable *if* rsex=1 | (rsex=2 & (dage=16 | dage=17))

(R) (ND) **abort20** had an abortion before age 20

0: no *if* dage>=20 & abort=2 | (manyab=1 & (ageab≠-1 & ageab>=20 & ageab<99)) | (manyab>1 & manyab<99 & (ageabfir≠-1 & ageabfir>=20 & ageabfir<99))

1: yes *if* (manyab=1 & (ageab≠-1 & ageab<20)) | (manyab>1 & manyab<99 & (ageabfir≠-1 & ageabfir<20))

9: not answered *else if* missing

-1: not applicable *if* rsex=1 | (rsex=2 & (dage=16 | dage=17 | dage=18 | dage=19))

(R) (ND) **abortl5y** had an abortion, last 5 years

0: no *if* abort=2 | (dage-ageabal2>5)

1: yes *if* (dage-ageabal2<=5)

9: not answered *else if* missing

-1: not applicable *if* rsex=1

(R) (ND) **abortlyr** had an abortion, last year

0: no *if* abort=2 | (dage-ageabal2>1)

1: yes *if* (dage-ageabal2<=1)

9: not answered *else if* missing

-1: not applicable *if* rsex=1

Note: the questions about fertility problems (infertl and medhelp) appeared here in Natsal-1 and Natsal-2 but in Natsal-3 were moved to the end of the Fertility intentions and infertility section.

UNPLANNED PREGNANCY

UNPLANNED PREGNANCY

Note: Natsal-3 asked the London Measure of Unplanned Pregnancy (<http://measure.ascody.co.uk>) for pregnancies within the last year. This was only asked of women who had a pregnancy within the last year (i.e. those who were pregnant at the time of interview, gave birth in the last year, or had another pregnancy outcome in the last year.

*Note: the variable **lmupscore** should be used for analysis.*

*Note: **pregyr** is an approximate variable derived automatically in the CASI which uses date of birth of live births, and age at miscarriage, abortion, and still birth to calculate whether the respondent was pregnant in the last year*

(N) **pregyr** Pregnant in the last year

0: no if FOR MOST RECENT PREGNANCY: ((zpregoc=1 | zpregoc=2 | zpregoc=3) & (dage – prgage>1) & lmupscr==2) & ((zpregoc=4 | zpregoc=5) & (dateoi – alldob>1) & zpregoc≠6

1: yes if FOR MOST RECENT PREGNANCY: ((zpregoc=1 | zpregoc=2 | zpregoc=3) & (dage – prgage≤1) & lmupscr==1) | ((zpregoc=4 | zpregoc=5) & (dateoi – alldob≤1) | zpregoc=6

9: not answered

-1: not applicable if rsex=1 | everpreg=2

(ND) **pa** experienced pregnancy with known outcome in the past year, if aged <45.

-1: not applicable if rsex=1 | (rsex=2 & dage>44)

0: no if rsex=2 & dage≤44 & (pregyr=0 | pregyr=-1 | lastpreg=5)

1: yes if rsex=2 & dage≤44 & pregyr=1 & lastpreg>0 & lastpreg<5

9: not answered if pregyr=9

*Note: **lmupscr** is a direct screening question to ascertain whether the respondent was pregnant in the last 12 months, if it is not clear from other questions. **Maxprage** is a program variable which is the age given at prgage for the most recent pregnancy (where outcome was miscarriage, abortion, or stillbirth).*

(N) **lmupscr** been pregnant in last 12 months (if not clear from pregnancy questions)

-1: not applicable if rsex=1 | everpreg=2 | zpregoc=6 (any loop) | (maxprage<dage-1) | (for all loops)(zpregoc=-1 | zpregoc=4 | zpregoc=5)

1: yes

2: no

9: not answered

(N) **lmup1** contraception use in month became pregnant

-1: not applicable if rsex=1 | everpreg=2 | pregyr=0 | lmupscr=2

1: not using contraception

2: using contraception but not on every occasion

3: always used contraception but know method had failed at least once

4: always used contraception

9: not answered

(N) **lmup2** in terms of being a mother, pregnancy has happened

-1: not applicable if rsex=1 | everpreg=2 | pregyr=0 | lmupscr=2

1: right time

2: ok but not quite right time

3: wrong time

9: not answered

(N) **lmup3** intentions to get pregnant just before became pregnant

-1: not applicable if rsex=1 | everpreg=2 | pregyr=0 | lmupscr=2

1: intended to get pregnant

2: intentions kept changing

3: did not intend to get pregnant

9: not answered

UNPLANNED PREGNANCY

(N) **Imup4** feelings towards baby just before became pregnant

-1: not applicable *if* rsex=1 | everpreg=2 | pregyr=0 | Imupscr=2

1: wanted to have baby

2: mixed feelings about having baby

3: did not want to have a baby

9: not answered

(N) **Imup5** discussion about pregnancy with male partner before became pregnant

-1: not applicable *if* rsex=1 | everpreg=2 | pregyr=0 | Imupscr=2

1: agreed would like to get pregnant

2: discussed children but hadn't agreed to get pregnant

3: never discussed having children

9: not answered

Note: binary variables Imupfold..Imupnone were created directly from the multi-code question Imup6.

(N) **Imupfolic** did anything to improve health before became pregnant: took folic acid

-1: not applicable *if* rsex=1 | everpreg=2 | pregyr=0 | Imupscr=2

0: no

1: yes

9: not answered

Note: Imupsmok to Imupnone have same response options as Imupfolic

(N) **Imupsmok** did anything to improve health before became pregnant: stopped/cut down smoking

(N) **Imupdrk** did anything to improve health before became pregnant: stopped/cut down drinking

(N) **Imuphlth** did anything to improve health before became pregnant: at more healthily

(N) **Imupmad** did anything to improve health before became pregnant: sought medical / health advice

(N) **Imupoact** did anything to improve health before became pregnant: some other action

(N) **Imupnone** did nothing to improve health before becoming pregnant

(N) **Imupscore** London Measure of Unplanned Pregnancy score.

sysmis if missing

Note: in Wellings et al (2013) The prevalence of unplanned pregnancy and associated factors in Britain: findings from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3), analysis of planstat and unplanned was restricted to those with a pregnancy with a known outcome in the past year (using the variable pregyrkw).

(ND) **planstat** pregnancy planning status (pregnancies in the last year)

-1: not applicable *if* rsex=1 | everpreg=2 | pregyr=0 | Imupscr=2

1: unplanned *if* Imupscore>=1 & Imupscore<=3

2: ambivalent *if* Imupscore>=4 & Imupscore<=9

3: planned *if* Imupscore>=10 & Imupscore<=12

9: not answered *else if missing*

(ND) **unplanned** unplanned pregnancy (pregnancies in the last year)

-1: not applicable *if* rsex=1 | everpreg=2 | pregyr=0 | Imupscr=2

0: no *if* planstat=2 | planstat=3

1: yes *if* planstat=1

9: not answered *if* planstat=9

UNPLANNED PREGNANCY

*Note: pregrisk derivation is simplified for purposes of the codebook. r1ptagenow1-3 represents the respondent's most recent 3 partners (in the past year) ages now. This is calculated by combining the following: $\text{int}((r1ptage*12+(((dateyoi*12)+datemoi)-((r1datey*12)+r1datem)))/12)$*

(ND) **pregrisk** pregnancy risk

1: no vaginal sex in the past year *if* lastvag>=5 & lastvag≠9

2: intention to conceive or pregnant current/past year *if* fertint=1 | (rsex==2 & pregyr=1) | (lmupscore≠-1 & (lastvag==9 | lastvag<=5)

3: permanent contraception/no longer fertile *if* (rsex=2 & (yfsteril=1 | fsteril=1)) | (rsex=1 & (ymsteril=1 | msteril=1)) | medwchc1=1 | (rsex=2 & ymsteril=1 & het1yr=1) | (rsex=1 & yfsteril=1 & het1yr=1) | (agrp>=4 & (periods3=5 | periods3=6) & rsex=2) | (agrp>=5 & rsex=2) | (het1yr=3 & dlover1p1=2 & r1ptagenow1>=50 & dlover1p2=2 & r1ptagenow2>=50 & dlover1p3=2 & r1ptagenow3>=50) | (het1yr=2 & dlover1p1=2 & r1ptagenow1>=50 & dlover1p2=2 & r1ptagenow2>=50) | (het1yr=1 & dlover1p1=2 & r1ptagenow1>=50)

4: effective contraception used *if* (ycondom=1 | yfemidom=1 | yconpill=1 | ycoil=1 | ymirena=1 | yinjectn=1 | yimplant=1) & pregrisk≠5

5: less effective contraception used *if* yemcyiud=1 | yemcypill=1 | ygels=1 | yrhythm=1 | ycap=1 | ywithdrw=1

6: no contraception used *if* pregrisk≠1 | pregrisk≠2 | pregrisk≠3 | pregrisk≠4 | pregrisk≠5 | pregrisk≠9

9: not answered *if* lastvag=9

FAMILY FORMATION

FAMILY FORMATION

Note: anychild, nochild, dobchm, dobchy were asked of all respondents in the face to face section in Natsal-1 and Natsal-2. In Natsal-3 these questions were moved to the CASI and were asked of men only (women were asked about their children in the pregnancy history module). only asked of men who reported ever having vaginal sex. The variable names have been changed to reflect this difference.

Note: stillbirths are no longer included in these questions which will affect comparability with Natsal-2 & Natsal-1

(R) (q14a) **anychild** any natural children

- 1: yes
- 2: no
- 9: not answered

(R) (q14c) **nochild** number of natural children (incl. stillborn and died)

- 0 if anychild=2
- 99: not answered

(R) (N) **dobchm01** month of birth of 1st born natural child

-1: not applicable if anychild=2 | (nochild < dobchmn)

- 1: January
- 2: February
- 3: March
- 4: April
- 5: May
- 6: June
- 7: July
- 8: August
- 9: September
- 10: October
- 11: November
- 12: December
- 99: not answered

Similarly for up to 10 natural children:

(RM) (N) **dobchm02** month of birth of 2nd born natural child

(RM) (N) **dobchm03** month of birth of 3rd born natural child

(RM) (N) **dobchm04** month of birth of 4th born natural child

(RM) (N) **dobchm05** month of birth of 5th born natural child

(RM) (N) **dobchm06** month of birth of 6th born natural child

(RM) (N) **dobchm07** month of birth of 7th born natural child

(RM) (N) **dobchm08** month of birth of 8th born natural child

(RM) (N) **dobchm09** month of birth of 9th born natural child

(RM) (N) **dobchm10** month of birth of 10th born natural child

(RM) (N) **dobchy01** year of birth of 1st born natural child

- 9999: not answered
- 1: not applicable if anychild=2 | (nochild < dobchyn)

Similarly for up to 10 natural children:

(RM) (N) **dobchy02** year of birth of 2nd born natural child

(RM) (N) **dobchy03** year of birth of 3rd born natural child

(RM) (N) **dobchy04** year of birth of 4th born natural child

(RM) (N) **dobchy05** year of birth of 5th born natural child

(RM) (N) **dobchy06** year of birth of 6th born natural child

(RM) (N) **dobchy07** year of birth of 7th born natural child

(RM) (N) **dobchy08** year of birth of 8th born natural child

(RM) (N) **dobchy09** year of birth of 9th born natural child

(RM) (N) **dobchy10** year of birth of 10th born natural child

FAMILY FORMATION

Note: Variables 'cmc_ch1' .. 'cmc_ch10' report the date of birth of all natural children in terms of the number of months since the beginning of 1900.

(RND) **cmc_ch1** century-month-code of month and year of birth of 1st child

-1: not applicable if dobchy01=-1

Compute $\text{cmc_ch1} = ((\text{dobchy01}-1900) * 12) + \text{dobchm01}$ if $(\text{dobchy01} \neq -1 \ \& \ \text{dobchy01} \neq 9999) \ \& \ (\text{dobchm01} \neq -1 \ \& \ \text{dobchm01} \neq 99)$

9999: not answered if dobchy01=9999 | dobchm01=99

Similarly for up to 10 natural children:

(RND) **cmc_ch2** century-month-code of month and year of birth of 2nd child

(RND) **cmc_ch3** century-month-code of month and year of birth of 3rd child

(RND) **cmc_ch4** century-month-code of month and year of birth of 4th child

(RND) **cmc_ch5** century-month-code of month and year of birth of 5th child

(RND) **cmc_ch6** century-month-code of month and year of birth of 6th child

(RND) **cmc_ch7** century-month-code of month and year of birth of 7th child

(RND) **cmc_ch8** century-month-code of month and year of birth of 8th child

(RND) **cmc_ch9** century-month-code of month and year of birth of 9th child

(RND) **cmc_ch10** century-month-code of month and year of birth of 10th child

(N) **anychm** any natural children (excl. stillbirths) – men only

-1: if rsex=2

1: yes

2: no or if lastvag=7

9: not answered

(ND) **anychall** any natural children (excl. stillbirths)

1: yes if $(\text{rsex}=1 \ \& \ \text{anychm}=1) \ | \ (\text{rsex}=2 \ \& \ (\text{zpregoc}=4 \ | \ \text{zpregoc}=5 \ \text{for any loop}))$

2: no if $(\text{rsex}=1 \ \& \ \text{anychm}=2) \ | \ (\text{rsex}=2 \ \& \ (\text{zpregoc} \neq 4 \ \& \ \text{zpregoc} \neq 5 \ \& \ \text{zpregoc} \neq 9 \ \text{for any loop}))$

9: not answered if nochall=999

Note: nochall was a direct question for men who answered anychm=1, and is derived for women using information about live births from the pregnancy module.

(N) **nochall** number of natural children (excl. stillbirths)

0 if anychall=2

999 "not answered"

(ND) **prevparity** number of children before most recent pregnancy (women only)

-1: not applicable if rsex=1

0: 0 if $(\text{rsex}=2 \ \& \ \text{nochall}=0 \ \& \ \text{everpreg}=2) \ | \ (\text{rsex}=2 \ \& \ \text{nochall}=0 \ \& \ \text{lastpreg} \neq 1 \ \& \ \text{lastpreg} \neq 9) \ | \ (\text{rsex}=2 \ \& \ \text{lastpreg}=1 \ \& \ \text{nochall}=1)$

1: 1 if $(\text{rsex}=2 \ \& \ \text{nochall}=1 \ \& \ \text{lastpreg} \neq 1 \ \& \ \text{lastpreg} \neq 9) \ | \ (\text{rsex}=2 \ \& \ \text{lastpreg}=1 \ \& \ \text{nochall}=2)$

2: 2+ if $(\text{rsex}=2 \ \& \ \text{nochall}=2 \ \& \ \text{lastpreg} \neq 1 \ \& \ \text{lastpreg} \neq 9) \ | \ (\text{rsex}=2 \ \& \ \text{nochall} \geq 3)$

9: not answered if

Note: All missing values are set to sysmiss for dobcm01 to dobcm12. Day is set to 01 for all cases.

(ND) **dobcm01** Date of birth of 1st born natural child (male respondents only, excl stillborns)

Similarly for up to 12 natural children:

(ND) **dobcm02** Date of birth of 2nd born natural child (male respondents only, excl stillborns)

(ND) **dobcm03** Date of birth of 3rd born natural child (male respondents only, excl stillborns)

(ND) **dobcm04** Date of birth of 4th born natural child (male respondents only, excl stillborns)

(ND) **dobcm05** Date of birth of 5th born natural child (male respondents only, excl stillborns)

(ND) **dobcm06** Date of birth of 6th born natural child (male respondents only, excl stillborns)

(ND) **dobcm07** Date of birth of 7th born natural child (male respondents only, excl stillborns)

(ND) **dobcm08** Date of birth of 8th born natural child (male respondents only, excl stillborns)

(ND) **dobcm09** Date of birth of 9th born natural child (male respondents only, excl stillborns)

(ND) **dobcm10** Date of birth of 10th born natural child (male respondents only, excl stillborns)

FAMILY FORMATION

(ND) **dobcm11** Date of birth of 11th born natural child (male respondents only, excl stillborns)
(ND) **dobcm12** Date of birth of 12th born natural child (male respondents only, excl stillborns)

(N) **dobcm01** Year of birth of 1st born natural child (male respondents only, excl stillborns)
-1: not applicable if rsex=2 | nochall=0
9999: not answered

Similarly for up to 12 natural children:

(N) **dobcm02** Year of birth of 2nd born natural child (male respondents only, excl stillborns)
(N) **dobcm03** Year of birth of 3rd born natural child (male respondents only, excl stillborns)
(N) **dobcm04** Year of birth of 4th born natural child (male respondents only, excl stillborns)
(N) **dobcm05** Year of birth of 5th born natural child (male respondents only, excl stillborns)
(N) **dobcm06** Year of birth of 6th born natural child (male respondents only, excl stillborns)
(N) **dobcm07** Year of birth of 7th born natural child (male respondents only, excl stillborns)
(N) **dobcm08** Year of birth of 8th born natural child (male respondents only, excl stillborns)
(N) **dobcm09** Year of birth of 9th born natural child (male respondents only, excl stillborns)
(N) **dobcm10** Year of birth of 10th born natural child (male respondents only, excl stillborns)
(N) **dobcm11** Year of birth of 11th born natural child (male respondents only, excl stillborns)
(N) **dobcm12** Year of birth of 12th born natural child (male respondents only, excl stillborns)

(N) **dobcmm01** Month of birth of 1st born natural child (male respondents only, excl stillborns)
-1: not applicable if rsex=2 | nochall=0
1: January
2: February
3: March
4: April
5: May
6: June
7: July
8: August
9: September
10: October
11: November
12: December
99: not answered

Similarly for up to 12 natural children:

(N) **dobcmm02** Month of birth of 2nd born natural child (male respondents only, excl stillborns)
(N) **dobcmm03** Month of birth of 3rd born natural child (male respondents only, excl stillborns)
(N) **dobcmm04** Month of birth of 4th born natural child (male respondents only, excl stillborns)
(N) **dobcmm05** Month of birth of 5th born natural child (male respondents only, excl stillborns)
(N) **dobcmm06** Month of birth of 6th born natural child (male respondents only, excl stillborns)
(N) **dobcmm07** Month of birth of 7th born natural child (male respondents only, excl stillborns)
(N) **dobcmm08** Month of birth of 8th born natural child (male respondents only, excl stillborns)
(N) **dobcmm09** Month of birth of 9th born natural child (male respondents only, excl stillborns)
(N) **dobcmm10** Month of birth of 10th born natural child (male respondents only, excl stillborns)
(N) **dobcmm11** Month of birth of 11th born natural child (male respondents only, excl stillborns)
(N) **dobcmm12** Month of birth of 12th born natural child (male respondents only, excl stillborns)

Note: Variable 'age1chil' is derived from the respondent's month and year of birth, expressed here as 'cmc_dob' the number of months since January 1900. These variables are not available in the archived dataset for confidentiality reasons.

(R) (q14b) **age1chil** age at birth of 1st child
-1: not applicable if anychild=2
Compute (cmc_ch1- cmc_dob)/12) if (cmc_ch1≠-1 & cmc_ch1≠9999)
99: not answered if cmc_ch1=9999

FAMILY FORMATION

(N) **age1ch** age at birth of 1st child

-1: not applicable if anychall=2

Compute date difference in years between dobcm01 and rdob if rsex=1

Compute date difference in years between alldob01 and rdob if rsex=2

99: not answered

(U – see below) (ND) **child18** had a child before age 18

-1: not applicable if dage=16 | dage=17

0: no if dage>=18 & ((age1chil>=18 & age1chil <99) | anychild=2) 1: yes if dage>=18 & (age1chil >0 & age1chil <18)

9: not answered if cmc_ch1=9999

(U) **child18** had a child before age 18

-1: not applicable if dage=16 | dage=17

0: no if dage>=18 & ((age1ch>=18 & age1ch<99) | anychall=2)

1: yes if dage>=18 & (age1ch >0 & age1ch<18)

9: not answered if age1ch=99

(U – see below) (ND) **child20** had a child before age 20

-1: not applicable if dage=16 | dage=17 | dage=18 | dage=19

0: no if dage>=20 & ((age1chil>=20 & age1chil <99) | anychild=2)

1: yes if dage>=20 & (age1chld >0 & age1chld<20)

9: not answered if cmc_ch1=9999

(U) **child20** had a child before age 20

-1: not applicable if dage>=16 & dage<20

0: no if dage>=20 & ((age1ch>=20 & age1ch<99) | anychall=2)

1: yes if dage>=20 & (age1ch >0 & age1ch<20)

9: not answered if age1ch=99

(ND) **nochb435** did not have any natural children by the age of 35 (those aged 35+ only)

-1: not applicable if dage<=34

0: had children before age 35 else if age1ch>=0 & age1ch<=34

1: did not have children before age 35 else if (age1ch>=35 & age1ch<=99) | (age1ch=-1)

9: not answered if age1ch=99

Note: samemf is now split into two questions, one for men and one for women since information on children is collected in different ways

(R) (q14e) **samemf** natural children have same other parent

-1: not applicable if anychild=2 | nochild=1

1: yes both/all the same other parent

2: no, different other parent(s)

9: not answered

(N) **samemfm** children have same mother

1: yes

2: no

9: not answered

-1: not applicable if rsex=2 | anychall=2 | nochall=1

Note: an equivalent question 'samemff' was asked of women, however there was a filter error in the CASI which meant that this question did not work, therefore the variable has not been included in the final data.

Note: Variable q15a in Natsal-1 asked respondents if they had any adopted or step children, whereas Natsal-2 asked about adopted children in a separate question to the question about step children.

(U) (M) **adopchld** (q15a*) **adopchil** any adopted children1: yes

2: no

9: not answered

FAMILY FORMATION

(U) -1: not applicable if `scelign3=2` | `askcasi=2`

(U) (M) **stepchld** (q15a*) **stepchil** any step-children

1: yes

2: no

9: not answered

(U) -1: not applicable if `scelign3=2` | `askcasi=2`

FERTILITY INTENTIONS

FERTILITY INTENTIONS AND INFERTILITY

Note: Additional questions were asked about fertility intentions in Natsal-3

(N) **fertint** how respondent feels about having (more) children

-1: not applicable *if* scelig3=2 | (rsex=2 & dage>=50) | (zpregoc=6 for any loop)

1: definitely like (more) and currently trying

2: definitely like (more) but not currently trying

3: might like (more) but not sure yet

4: definitely not like (more)

5: don't know

9: not answered

Note: codes 1-7 are the original answer options for fertwhy, codes 8-12 are additional codes that have been added to the data from back-coding the 'other' answers (open text question in the questionnaire).

(N) **fertwhy** main reason not currently trying to have (more) children

1: don't feel ready

2: don't have suitable partner

3: partner unwilling

4: financial

5: work / study

6: partner's work / study

7: other

9: not answered

8: health problems/medical reasons

9: age – too old

10: age – too young

11: recently had child/waiting until children are older

12: wanted to be married first

-1: not applicable *if* fertint=-1 | fertint=1 | fertint=3 | fertint=4 | fertint=5

Note: In Natsal-2 the variables infertl and medhelp appeared in the gynaecological history section. In Natsal-3 these questions weren't asked of respondents who had never had vaginal sex (lastvag=never). The question about infertility was updated to refer to a 1 year period in Natsal-3, rather than 6 months, therefore the variable name has been changed.

Note: Variables 'infertl' and 'medhelp' refer to both men and women:

(R) (M) (s15a) **infertl** pregnancy attempt, ever, for 6+ months

1: yes

2: no *or if* het=0 | hetsc=0 | lastvag=7

9: not answered

-1: not applicable *else if* scelig=2

(N) **infert1y** pregnancy attempt, ever, for 12+ months

1: yes

2: no

9: not answered

-1: not applicable *if* scelig3=2 | askcasi=2 | lastvag=7

(U) (M) (s15b) **medhelp** sought medical/professional help about infertility, ever

1: yes

2: no *if* het=0 | hetsc=0

9: not answered

(U) -1: not applicable *if* scelig3=2 | askcasi=2 | lastvag=7

-1: not applicable *else if* scelig=2

SEXUALLY TRANSMITTED INFECTIONS

SEXUALLY TRANSMITTED INFECTIONS

Note: Eligibility for this section changed in Natsal-3. Only asked of people reporting 1+ lifetime partner at hetlife and samlife

(N) **stdtrt** if thought had STI, where would first go for diagnosis and/or treatment (if had any lifetime partners)

- 1: GP
- 2: GUM clinic
- 3: NHS Family planning clinic / contraceptive clinic / reproductive health clinic
- 4: NHS Antenatal clinic / midwife
- 5: Private non-NHS clinic or doctor
- 6: Pharmacy / chemist
- 7: Internet site offering treatment
- 8: Youth advisory clinic (e.g. Brook clinic)
- 9: Hospital accident and emergency (A&E) department
- 10: Somewhere else
- 1: not applicable *if* (hetlife=0 & samlife=0) | dage>44

Note: during the Natsal-3 data editing process, a small number of cases were identified that were not eligible for this question, but data from elsewhere in the questionnaire suggests that they had attended a STD clinic, ever. These cases were recoded as 'yes' at stdclin.

(*U) (s16a) **stdclin** STD clinic attendance, ever *Ever attended a sexual health (GUM) clinic (if had any lifetime partners)*

- 1: yes
- 2: no
- 9: not answered
- (U) -1: not applicable *if* (hetlife=0 & samlife=0) -1: not applicable *if* scelig=2

Note: codes for whnclin categorised into broader periods which is how data were categorised for analysis in Natsal-2. The question itself is identical.

(U – see below) (N) **whnclin** when (last) attended a STD clinic

- 1: in last 4 weeks
- 2: between 4 weeks and 3 months
- 3: between 3 months and 1 year
- 4: between 1 and 2 years
- 5: between 2 and 3 years
- 6: between 3 and 4 years
- 7: between 4 and 5 years
- 8: longer than 5 years
- 99: not answered
- 1: not applicable *if* scelig=2 | stdclin=2

whnclin when (last) attended a STD clinic (if had any lifetime partners)

- 1: less than one year ago
- 2: between 1 and 5 years ago
- 3: between 5 and 10 years ago
- 4: more than 10 years ago
- 9: not answered
- 1: not applicable *if* stdclin=-1 | stdclin=2

(U – see below) (rs16 D) **attendcl** when (last) attended a STD clinic

- 1: never *if* stdclin=2
- 2: in last year *if* whnclin=1 | whnclin=2 | whnclin=3
- 3: 1-5 years *if* whnclin=4 | whnclin=5 | whnclin=6 | whnclin=7
- 4: over 5 years *if* whnclin=8
- 5: attended but not answered when *if* stdclin=1 & whnclin=99
- 9: not answered *if* whnclin=99
- 1: not applicable *if* whnclin=-1

SEXUALLY TRANSMITTED INFECTIONS

attendcl when (last) attended a sexual health (GUM) clinic

- 1: never *if* stdclin=2
- 2: less than 1 year ago *if* whnclin=1
- 3: between 1 and 5 years ago *if* whnclin=2
- 4: over 5 years ago *if* whnclin=3 | whnclin=4
- 5: attended but not answered when *if* stdclin=1 & whnclin=9
- 9: not answered *if* whnclin=9
- 1: not applicable *if* whnclin=-1

Note: Variable 'whnclin2' uses the same codes as in Natsal-1. There are no codes 0, 1 or 2:

(U – see below) (s16b UD) **whnclin2** when (last) attended a STD clinic

- 3: in last year *if* whnclin=1 | whnclin=2 | whnclin=3
- 4: between 1 and 5 years *if* whnclin=4 | whnclin=5 | whnclin=6 | whnclin=7
- 5: longer than 5 years *if* whnclin=8
- 9: not answered *if* whnclin=99
- 1: not applicable *if* whnclin=-1

whnclin2 when (last) attended a STD clinic

- 3: in last year *if* whnclin=1
- 4: between 1 and 5 years *if* whnclin=2
- 5: longer than 5 years *if* whnclin=3 | whnclin=4
- 9: not answered *if* whnclin=9
- 1: not applicable *if* whnclin=-1

Note: Variable is consistent with the definition of 'whnclin' & uses the same codes as in Natsal-1:

(U) (ND) **whncl5yr** attended a STD clinic, last 5 years

- 0: no *if* whnclin=8 | (whnclin=-1 & scelg=2)
- 1: yes *if* whnclin>=1 & whnclin<=7)
- 9: not answered *if* whnclin=99
- 1: not applicable *else if* whnclin=-1

(U) **whncl5yr** attended a STD clinic, last 5 years

- 0: no *if* whnclin2=5 | stdclin=2
- 1: yes *if* whnclin2=3 | whnclin2=4
- 9: not answered *if* whnclin2=9
- 1: not applicable *else if* hetlife=0 & samlife=0

- Notes:*
1. This codebook gives only the binary variables to be used for analysis, not the original multi-code questions (for original questions, see questionnaire documentation). The following binary variables were created directly from the question **diagnos** in Natsal-3, and derived from **diagnos1-diagnos7** in Natsal-2. All are coded similarly to chlamydia unless indicated.
 2. The question 'diagnos' was updated in Natsal-3 so more common STIs appear higher up in the list of response options. Additionally, in Natsal-3 the question specifies including infections that were not transmitted sexually which may increase reports of thrush.
 3. Those who were not eligible for the self-completion section are coded as 'not applicable' for variables chlamydia..eversti3. However, as most of these participants have not had any sexual partners (hence not being eligible for the self-completion) data users may wish to recode these cases as 'no' for these variables before analysis so that they are included in the denominator.

chlamydia (ND) **chlamyda** ever diagnosed with chlamydia (if had any lifetime partners)

- 1: yes
- 0: no *or if* (hetlife=0 & samlife=0)
- 9: not answered
- 98: had a STD but can't remember which one *if* dkstd=1 & chlamydia=0
- 1: not applicable *if* scelg3=2

SEXUALLY TRANSMITTED INFECTIONS

And similarly...

(ND) **gonorrh** ever diagnosed with gonorrhoea (if had any lifetime partners)
(ND) **warts** ever diagnosed with genital warts (if had any lifetime partners)
(ND) **syphilis** ever diagnosed with syphilis (if had any lifetime partners)
(ND) **trich** ever diagnosed with trichomonas (if had any lifetime partners)
(ND) **herpes** ever diagnosed with genital herpes (if had any lifetime partners)
(N) **plicecrab** ever diagnosed with public lice / crabs (if had any lifetime partners)
(N) **hepatitis** ever diagnosed with hepatitis B (if had any lifetime partners)

Note: nsungu and epididy were answer options only available to men

(ND) **nsungu** ever diagnosed with NSU/NGU (if had any lifetime partners)

1: yes

0: no or if (hetlife=0 & samlife=0)

9: not answered

98: had a STD but can't remember which one if dkstd=1 & nsungu=0

-1: not applicable if scelign3=2 | rsex=2

And similarly..

(N) **epididy** ever diagnosed with epididymitis (if had any lifetime partners)

Note: pid, thrush, and bacvagin were answer options only available to women

(ND) **pid** ever diagnosed with pelvic inflammatory disease (if had any lifetime partners)

1: yes

0: no or if (hetlife=0 & samlife=0)

9: not answered

98: had a STD but can't remember which one if dkstd=1 & pid=0

-1: not applicable if scelign3=2 | rsex=1

And similarly..

(ND) **thrush** ever diagnosed with thrush (if had any lifetime partners)

(N) **bacvagin** ever diagnosed with bacterial vaginosis (if had any lifetime partners)

(ND) **dkstd** ever diagnosed with a STD but can't remember which one (if had any lifetime partners)

(N) **nostd** no sexually transmitted infections ever diagnosed (if had any lifetime partners)

1: yes or if hetlife=0 & samlife=0

0: no

9: not answered

-1: not applicable if scelign3=2

Note: In Natsal-3 the question diagnos (and resulting binary variables dkstd to nostd) cover a greater number of infections than in Natsal-2. As eversti, eversti2, eversti3 are derived from these binary variables, this change may affect the comparability of these variables between the two survey years.

(U – see below) (ND) **eversti** ever diagnosed with a STI

1: yes if herpes=1 | trich=1 | gonorrh=1 | syphilis=1 | chlamydia=1 | nsungu=1 | warts=1 | pid=1 | dkstd=1

0: no if herpes=-1 | herpes=0 | herpes=98

9: not answered if herpes=9

-1: not applicable if herpes=-1

eversti ever diagnosed with a STI (Natsal-3 definition, including thrush as an STI)

1: yes if herpes=1 | trich=1 | gonorrh=1 | syphilis=1 | chlamydia=1 | plicecrab=1 | hepatitis=1 | epididy=1 | nsungu=1 | warts=1 | pid=1 | thrush=1 | bacvagin=1 | dkstd=1

0: no if nostd =1

9: not answered if nostd=9

-1: not applicable if nostd=-1

SEXUALLY TRANSMITTED INFECTIONS

Note: Variables eversti2 and eversti3 do not count thrush as a STI

(U – see below) (ND) **eversti2** ever been diagnosed with a STI (excl. thrush)

1: yes if herpes=1 | trich=1 | gonorrh=1 | syphilis=1 | chlamyda=1 | nsungu=1 | warts=1 | pid=1 | dkstd=1

0: no if herpes=-1 | herpes=0 | herpes=98

97: only STI had is thrush if thrush=1

9: not answered if herpes=9

-1: not applicable if herpes=-1

eversti2 ever been diagnosed with a STI (Natsal-3 definition, excl. thrush)

1: yes if herpes=1 | trich=1 | gonorrh=1 | syphilis=1 | chlamydia=1 | plicecrab=1 | hepatitis=1 | epididy=1 |

nsungu=1 | warts=1 | pid=1 | bacvagin=1 | dkstd=1

0: no if nostd =1

97: only had is thrush if thrush=1 & (all other STIs ≠ 1)

9: not answered if nostd=9

-1: not applicable if nostd=-1

(U – see below) (ND) **eversti3** ever been diagnosed with a STI (excl. thrush)

1: yes if herpes=1 | trich=1 | gonorrh=1 | syphilis=1 | chlamyda=1 | nsungu=1 | warts=1 | pid=1 | dkstd=1

0: no if herpes=-1 | herpes=0 | herpes=98 | eversti2=97

9: not answered if herpes=9

-1: not applicable if herpes=-1

eversti3 ever been diagnosed with a STI (Natsal-3 definition, coding thrush as non-STI)

1: yes if herpes=1 | trich=1 | gonorrh=1 | syphilis=1 | chlamydia=1 | plicecrab=1 | hepatitis=1 | epididy=1 |

nsungu=1 | warts=1 | pid=1 | bacvagin=1 | dkstd=1

0: no if nostd =1 | (thrush=1 & all other STIs ≠ 1)

9: not answered if nostd=9

-1: not applicable if nostd=-1

Note: evervirl and everbact have not been derived for Natsal-3

(R) (ND) **evervirl** ever been diagnosed with a viral STI

1: yes if herpes=1 | warts=1

0: no else if eversti=0 | eversti=1

98: had a STI but can't remember which one

9: not answered if herpes=9

-1: not applicable if herpes=-1

(R) (ND) **everbact** ever been diagnosed with a bacterial STI

1: yes if trich=1 | gonorrh=1 | syphilis=1 | chlamyda=1 | nsungu=1 | pid=1

0: no else if eversti=0 | eversti=1

98: had a STI but can't remember which one

9: not answered if herpes=9

-1: not applicable if herpes=-1

Note: STI diagnosis questions were expanded in Natsal-3. Questions were also included on Chlamydia testing and HPV vaccination.

(N) **whnchlam** when last diagnosed with Chlamydia

1: Less than 1 year ago

2: Between 1 and 5 years ago

3: Between 5 and 10 years ago

4: More than 10 years ago

9: not answered

-1: not applicable if chlamydia=-1 | chlamydia=0

SEXUALLY TRANSMITTED INFECTIONS

Note: chtstwh1& chtstwy1 only asked of those aged 16-44 years

(N) **chtstwh1** where last tested for Chlamydia (if last diagnosis in last year)

- 1: General practice (GP) surgery
- 2: Sexual health clinic (GUM clinic)
- 3: NHS Family planning clinic / contraceptive clinic / reproductive health clinic
- 4: Ante-natal Clinic / midwife
- 5: Private non-NHS clinics or doctor
- 6: Youth advisory clinic (e.g. Brook clinic)
- 7: School / college / university
- 8: Termination of pregnancy (abortion) clinic
- 9: Hospital accident and emergency (A&E) department
- 10: Pharmacy / chemist
- 11: Internet
- 12: Other non-health care place e.g. youth club, festival, bar
- 13: Somewhere else
- 99: not answered
- 1: not applicable if chlamydia=-1 | chlamydia=0 | whnchlam=2 | whnchlam=3 | whnchlam=4 | dage>44

(N) **chtstwy1** why last tested for Chlamydia

- 1: I had symptoms
- 2: My partner had symptoms
- 3: I was notified because a partner diagnosed with Chlamydia
- 4: I wanted a general sexual health check-up
- 5: Check up after previous positive test
- 6: I had no symptoms but I was worried about Chlamydia risk
- 7: I was offered a routine test
- 8: Other
- 99: not answered
- 1: not applicable if chlamydia=-1 | chlamydia=0 | whnchlam=2 | whnchlam=3 | whnchlam=4 | dage>44

Note: chtrtlgr asked of those aged 16-44 years and those aged 45+ years if diagnosis in last 5 years

(N) **chTrtLgr** where last treated for chlamydia

- 1: General practice (GP) surgery
- 2: Sexual health clinic (GUM clinic)
- 3: NHS Family planning clinic / contraceptive clinic / reproductive health clinic
- 4: Ante-natal Clinic / midwife
- 5: Private non-NHS clinics or doctor
- 6: Pharmacy / chemist
- 7: internet
- 8: Youth advisory clinic (e.g. Brook clinic)
- 9: Termination of pregnancy (abortion) clinic
- 10: Hospital accident and emergency (A&E) department
- 11: Somewhere else
- 99: not answered
- 1: not applicable if chlamydia=0 | chlamydia=-1 | (dage>44 & (whnchlam=3 | whnchlam=4))

Note: chlmtst to chlofref were asked of those aged 16-44 years who haven't had a Chlamydia diagnosis in the last year. They were also asked of those aged 16-25 years in the CASI who had not had sex (and therefore not asked the question about chlamydia diagnosis)

(N) **chlmtst** tested for chlamydia in the last year (if not diagnosed with chlamydia in last year, age 16-44)

- 1: yes
- 2: no
- 9: not answered
- 1: not applicable if (dage<25 & scelign3=2) | (dage>=25 & dage<=44 & (chlamydia=-1 | whnchlam=1)) | dage>44

SEXUALLY TRANSMITTED INFECTIONS

(N) (D) **cttestly** tested for chlamydia in the last year

-1: not applicable *if* dage>44 | scelign3=2 | totlife=0

1: yes *else if* chlmttest=1 | whnchlam=1

0: no *else if* chlmttest=2 | whnchlam>1 & whnchlam<4

9: not answered *else if* scaccept=3 | whnchlam=9 | chlmttest=9

(N) **chltstwh** where last tested for chlamydia (if not diagnosed with chlamydia in last year, age 16-44)

1: General practice (GP) surgery

2: Sexual health clinic (GUM clinic)

3: NHS Family planning clinic / contraceptive clinic / reproductive health clinic

4: Ante-natal Clinic / midwife

5: Private non-NHS clinics or doctor

6: Youth advisory clinic (e.g. Brook clinic)

7: School / college / university

8: Termination of pregnancy (abortion) clinic

9: Hospital accident and emergency (A&E) department

10: Pharmacy / chemist

11: Internet

12: Other non-health care place e.g. youth club, festival, bar

13: Somewhere else

99: not answered

-1: not applicable *if* chlmttest=2 | chlmttest=-1

(N) **chltstw** why last tested for chlamydia (if not diagnosed with chlamydia in last year, age 16-44)

1: had symptoms

2: partner had symptoms

3: notified because a partner diagnosed with Chlamydia

4: wanted a general sexual health check-up

5: Check up after previous positive test

6: no symptoms but I was worried about Chlamydia risk

7: offered a routine test

8: Other

9: not answered

-1: not applicable *if* chlmttest=2 | chlmttest=-1

Note: chlofref asked of those aged 16-44 who have not been tested for chlamydia in the last year

(N) **chlofref** offered and refused Chlamydia test (if not diagnosed with chlamydia in last year, age 16-44)

1: yes

2: no

9: not answered

-1: not applicable *if* chlamydia=-1 | chlmttest=1 | whnchlam=1 | dage>44

Note: questions whn-X (where X is STI reported at gonorrh to nsungu) were asked of all who reported X. Questions wher-X were only asked of those reporting X who were aged 16-44, or those aged 45+ reporting being diagnosed with X in the last 5 years (at whn-X)

(N) **whngono** when last diagnosed with Gonorrhoea

1: Less than 1 year ago

2: Between 1 and 5 years ago

3: Between 5 and 10 years ago

4: More than 10 years ago

9: not answered

-1: not applicable *if* gonorrh=-1 | gonorrh=0

SEXUALLY TRANSMITTED INFECTIONS

(N) **whergono** where last treated for Gonorrhoea (if aged <45 or diagnosed in last 5 years)

1: GP

2: GUM clinic

3: NHS Family planning clinic / contraceptive clinic / reproductive health clinic

4: Ante-natal Clinic / midwife

5: Private non-NHS clinics or doctor

6: Pharmacy / chemist

7: internet

8: Youth advisory clinic (e.g. Brook clinic)

9: Termination of pregnancy (abortion) clinic

10: Hospital accident and emergency (A&E) department

11: Somewhere else

99: not answered

-1: not applicable if gonorrh=-1 | gonorrh=0 | (dage>44 & (whngono=3 | whngono=4))

Note: whngono and whergono repeated for each of the following STI diagnoses

(N) **whnwarts**

(N) **wherwart** (if aged <45 or diagnosed in last 5 years)

(N) **whnsyphl**

(N) **whersyph** (if aged <45 or diagnosed in last 5 years)

(N) **whntric**

(N) **whertric** (if aged <45 or diagnosed in last 5 years)

(N) **whnherp**

(N) **wherherp** (if aged <45 or diagnosed in last 5 years)

(N) **whnnsu**

(N) **whernsu** (if aged <45 or diagnosed in last 5 years)

Note: Variables whndiag1 ... whndiag8, diagno1... diagno8, and diagwhe1... diagwhe8, are STD-specific such that if respondent reported only having had chlamydia (diagnos1=5) then whndiag1, diagno1, and diagwhe1 will not have been answered but whndiag5, diagno5, and diagwhe5 will (hopefully) have been reported

(R) (N) **whndiag1** year last diagnosed with genital herpes

-1: not applicable if herpes=0 | herpes=-1

9: not answered

Note: Variables whndiag2 ... whndiag8 are coded in a similar way to whndiag1:

(R) (N) **whndiag2** year last diagnosed with trichomonas

(R) (N) **whndiag3** year last diagnosed with gonorrhoea

(R) (N) **whndiag4** year last diagnosed with syphilis

(R) (N) **whndiag5** year last diagnosed with chlamydia

(R) (N) **whndiag6** year last diagnosed with NS/NG urethritis

(R) (N) **whndiag7** year last diagnosed with genital warts

(R) (N) **whndiag8** year last diagnosed with PID

Notes:

1. variables ending ‘_l5y’ and ‘_l9y’ are coded as 1 if the difference between the year of interview and the year of (most recent) diagnosis is <=5 or 0 | 1, respectively. Thus, someone diagnosed (e.g.) January 1998 but who was interviewed in December 1999 would be coded as being diagnosed in the last year, which isn’t strictly the case. This method may result in an over-estimation but the alternative will result in an under-estimation, and since we are likely to be under-estimating the prevalence anyway, it was decided that a slight over-estimation would be preferable.
2. These variables have been replaced by ct5y etc – see below.

(R) (ND) **tv_l5y** diagnosed with trichomonas in the 5 calendar years before interview

0: no if ((whndiag2≠-1 & whndiag2≠9) & ((dateyoi-whndiag2)>5)) | trich=0

1: yes (whndiag2≠-1 & whndiag2≠9) & ((dateyoi-whndiag2)<=5)

9: not answered if whndiag2=9

-1: not applicable if scelig=2

SEXUALLY TRANSMITTED INFECTIONS

Note: Variables 'gc_15y' ... 'pid_15y' are coded in a similar way to 'tv_15y'

(R) (ND) **gc_15y** diagnosed with gonorrhoea in the 5 calendar years before interview

(R) (ND) **ct_15y** diagnosed with chlamydia in the 5 calendar years before interview

(R) (ND) **syph_15y** diagnosed with syphilis in the 5 calendar years before interview

(R) (ND) **nsu_15y** diagnosed with NSU/NGU in the 5 calendar years before interview

(R) (ND) **pid_15y** diagnosed with PID in the 5 calendar years before interview

(R) (ND) **wart_15y** diagnosed with genital warts in the 5 calendar years before interview

(R) (ND) **hsv_15y** diagnosed with genital warts in the 5 calendar years before interview

(R) (ND) **bact_15y** diagnosed with a bacterial STI in the 5 calendar years before interview

1: yes if tv_15y=1 | gc_15y=1 | ct_15y=1 | syph_15y=1 | nsu_15y=1 | pid_15y=1

0: no else if (everbact=1 & (tv_15y=0 & gc_15y=0 & ct_15y=0 & syph_15y=0 & nsu_15y=0 & pid_15y=0)) | everbact=0

98: not answered when last diagnosed with a bacterial STI else if everbact=1 & (tv_15y=9 | gc_15y=9 | ct_15y=9 | syph_15y=9 | nsu_15y=9 | pid_15y=9)

99: not answered whether ever diagnosed with a bacterial STI else if everbact=9 | everbact=98

-1: not applicable if everbact=-1

(R) (ND) **virl_15y** diagnosed with a viral STI in the 5 calendar years before interview

1: yes if hsv_15y=1 | wart_15y=1

0: no else if (evervirl=1 & (hsv_15y=0 & wart_15y=0)) | evervirl=0

98: not answered when last diagnosed with a viral STI else if evervirl=1 & (hsv_15y=9 | wart_15y=9)

99: not answered whether ever diagnosed with a viral STI else if evervirl=9 | 98

-1: not eligible for self-completion if evervirl=-1

(R) (ND) **any_15y** diagnosed with any STI in the 5 calendar years before interview

1: yes (excl. thrush) if hsv_15y=1 | wart_15y=1 | tv_15y=1 | gc_15y=1 | ct_15y=1 | syph_15y=1 | nsu_15y=1 | pid_15y=1

0: no else if (eversti2=1 & (hsv_15y=0 | wart_15y=0 | tv_15y=0 | gc_15y=0 | ct_15y=0 | syph_15y=0 | nsu_15y=0 | pid_15y=0)) | eversti2=0

98: not answered when last diagnosed with a STI (excl. thrush) else if eversti2=1 & (hsv_15y=9 | wart_15y=9 | tv_15y=9 | gc_15y=9 | ct_15y=9 | syph_15y=9 | nsu_15y=9 | pid_15y=9)

99: not answered whether ever diagnosed with a STI (excl. thrush) else if eversti2=9 | eversti2=98

-1: not eligible for self-completion if eversti2=-1

Note: In Natsal-2 respondents were asked for year of most recent diagnosis for each STI. In Natsal-3 respondents were asked 'how long ago' with banded response options. Therefore the Natsal-3 variables are more accurate than the Natsal-2 variables, which are an overestimate (see note on page 134).

(ND) **ct5y** diagnosed with chlamydia, last 5 years

1: yes if whnchlam=1 | whnchlam=2

0: no if chlamydia=0 | whnchlam=3 | whnchlam=4

-1: not applicable if hetlife=0 & samlife=0

9: not answered else if chlamydia=9 | whnchlam=9

Note: variables gc5y..nsu5y are coded in a similar to ct5y

(ND) **gc5y** diagnosed with gonorrhoea, last 5 years

(ND) **warts5y** diagnosed with genital warts, last 5 years

(ND) **syph5y** diagnosed with syphilis, last 5 years

(ND) **tric5y** diagnosed with trichomonas, last 5 years

(ND) **herp5y** diagnosed with herpes, last 5 years

(ND) **nsu5y** diagnosed with NSU/NGU, last 5 years

SEXUALLY TRANSMITTED INFECTIONS

Note: Natsal-3 variable any_15y2 differs from any_15y (hence new variable name) as follows:

- 1. In Natsal-2 respondents were asked when they were diagnosed with the following STIs: trichomonas, gonorrhoea, chlamydia, syphilis, NSU/NGU, pelvic inflammatory disease, genital warts, herpes. In Natsal-3 timing was asked for all of these infections except for pelvic inflammatory disease*
- 2. In Natsal-2 respondents were asked for year of most recent diagnosis. In Natsal-3 respondents were asked 'how long ago' with banded response options. Therefore the Natsal-3 variables are more accurate than the Natsal-2 variables, which are an overestimate (see note on page 134).*

(ND) **any_15y2** diagnosed with any STI (Natsal-3 definition) excl thrush in 5 years before interview

1: yes if (whnchlam=1 | whnchlam=2 | whngono=1 | whngono=2 | whnsyphl=1 | whnsyphl=2 | whnherp=1 | whnherp=2 | whnwarts=1 | whnwarts=2 | whntric=1 | whntric=2 | whnnsu=1 | whnnsu=2)

0: no if ((whnchlam=-1 | whnchlam=3 | whnchlam=4) & (whngono=-1 | whngono=3 | whngono=4) & (whnsyphl=-1 | whnsyphl=3 | whnsyphl=4) & (whnherp=-1 | whnherp=3 | whnherp=4) & (whnwarts=-1 | whnwarts=3 | whnwarts=4) & (whntric=-1 | whntric=3 | whntric=4) & (whnnsu=-1 | whnnsu=3 | whnnsu=4))

9: not answered else if (whnchlam=9 | whngono=9 | whnsyphl=9 | whnherp=9 | whnwarts=9 | whntric=9 | whnnsu=9)

Note: in Natsal-3 it was not possible to derive variables of the number of years since diagnosis of the various STIs since year of diagnosis was not collected. Respondents were instead given answer categories.

Note: Variables 'yrsherpe' ... 'yrsbact' are measured as the difference between the year of the interview and the reported year (last) diagnosed with herpes, i.e. a difference of 0 year may underestimate the true time by up to 364 days

(R) (ND) **yrsherpe** number of years since (last) diagnosed with genital herpes

compute dateyoi-whndiag1 if whndiag1≠-1 & whndiag1≠9

99: not answered if whndiag1=9

-1: not eligible for self-completion/never diagnosed with genital herpes if whndiag1=-1

Note: Variables 'yrstrich' and 'yrspid' are coded in a similar way to 'yrsherpe'

(R) (ND) **yrstrich** number of years since (last) diagnosed with trichomonas

(R) (ND) **yrsgonor** number of years since (last) diagnosed with gonorrhoea

(R) (ND) **yrsyphl** number of years since (last) diagnosed with syphilis

(R) (ND) **yrschlam** number of years since (last) diagnosed with chlamydia

(R) (ND) **yrsnsung** number of years since (last) diagnosed with NSU/NGU

(R) (ND) **yrswarts** number of years since (last) diagnosed with genital warts

(R) (ND) **yrspid** number of years since (last) diagnosed with PID

(R) (ND) **tv_lyr** diagnosed with trichomonas in the calendar year before interview

0: no if ((whndiag2≠-1 & whndiag2≠9) & ((dateyoi-whndiag2)>1)) | trich=0

1: yes (whndiag2≠-1 & ≠9) & ((dateyoi-whndiag2)<=1)

9: not answered if whndiag2=9

-1: not applicable if scelig=2

Note: Variables gc_lyr ... pid_lyr are coded in a similar way to tv_lyr

(R) (ND) **gc_lyr** diagnosed with gonorrhoea in the calendar year before interview

(R) (ND) **ct_lyr** diagnosed with chlamydia in the calendar year before interview

(R) (ND) **syph_lyr** diagnosed with syphilis in the calendar year before interview

(R) (ND) **nsu_lyr** diagnosed with NSU/NGU in the calendar year before interview

(R) (ND) **pid_lyr** diagnosed with PID in the calendar year before interview

(R) (ND) **wart_lyr** diagnosed with genital warts in the calendar year before interview

(R) (ND) **hsv_lyr** diagnosed with herpes in the calendar year before interview

(R) (ND) **bact_lyr** diagnosed with a bacterial STI (excl. thrush) in the calendar year before interview

1: yes if tv_lyr=1 | gc_lyr=1 | ct_lyr=1 | syph_lyr=1 | nsu_lyr=1 | pid_lyr=1

0: no else if (everbact=1 & (tv_lyr=0 & gc_lyr=0 & ct_lyr=0 & syph_lyr=0 & nsu_lyr=0 & pid_lyr=0)) | everbact=0

98: not answered when last diagnosed with a bacterial STI else if everbact=1 & (tv_lyr=9 | gc_lyr=9 | ct_lyr=9 | syph_lyr=9 | nsu_lyr=9 | pid_lyr=9)

99: not answered whether ever diagnosed with a bacterial STI else if everbact=9 | everbact=98

-1: not applicable if everbact=-1

SEXUALLY TRANSMITTED INFECTIONS

(R) (ND) **virl_1yr** diagnosed with a viral STI in the calendar year before interview

1: yes if hsv_1yr=1 | wart_1yr=1

0: no else if (evervirl=1 & (hsv_1yr=0 & wart_1yr=0)) | evervirl=0

98: not answered when last diagnosed with a viral STI else if evervirl=1 & (hsv_1yr=9 | wart_1yr=9)

99: not answered whether ever diagnosed with a viral STI else if evervirl=9 | evervirl=98

-1: not applicable if evervirl=-1

(R) (ND) **any_1yr** diagnosed with a STI (excl. thrush) in the calendar year before interview

1: yes if hsv_1yr=1 | wart_1yr=1 | tv_1yr=1 | gc_1yr=1 | ct_1yr=1 | syph_1yr=1 | nsu_1yr=1 | pid_1yr=1

0: no else if (eversti2=1 & (hsv_1yr=0 | wart_1yr=0 | tv_1yr=0 | gc_1yr=0 | ct_1yr=0 | syph_1yr=0 | nsu_1yr=0 | pid_1yr=0)) | eversti2=0

98: not answered when last diagnosed with a STI (excl. thrush) else if eversti2=1 & (hsv_1yr=9 | wart_1yr=9 | tv_1yr=9 | gc_1yr=9 | ct_1yr=9 | syph_1yr=9 | nsu_1yr=9 | pid_1yr=9)

99: not answered whether ever diagnosed with a STI (excl. thrush) else if eversti2=9 | eversti2=98

-1: not applicable if eversti2=-1

Note: In Natsal-2 respondents were asked for year of most recent diagnosis. In Natsal-3 respondents were asked 'how long ago' with banded response options. Therefore the Natsal-3 variables are more accurate than the Natsal-2 variables, which are an overestimate (see note on page 134).

(ND) **ct1y** diagnosed with chlamydia, last year

1: yes if whnchlam=1 | whnchlam=2

0: no if chlamydia=0 | whnchlam=3 | whnchlam=4

-1: not applicable if hetlife=0 & samlife=0

9: not answered else if chlamydia=9 | whnchlam=9

Note: variables gc1y..nsu1y are coded in a similar to ct1y

(ND) **gc1y** diagnosed with gonorrhoea, last year

(ND) **warts1y** diagnosed with genital warts, last year

(ND) **syph1y** diagnosed with syphilis, last year

(ND) **tric1y** diagnosed with trichomonas, last year

(ND) **herp1y** diagnosed with herpes, last year

(ND) **nsu1y** diagnosed with NSU/NGU, last year

Note: Natsal-3 variable any_1yr2 differs from any_1yr (hence new variable name) as follows:

- 1. In Natsal-2 respondents were asked when they were diagnosed with the following STIs: trichomonas, gonorrhoea, chlamydia, syphilis, NSU/NGU, pelvic inflammatory disease, genital warts, herpes. In Natsal-3 timing was asked for all of these infections except for pelvic inflammatory disease*
- 2. In Natsal-2 respondents were asked for year of most recent diagnosis. In Natsal-3 respondents were asked 'how long ago' with banded response options. Therefore the Natsal-3 variables are more accurate than the Natsal-2 variables, which are an overestimate (see note on page 134).*

any_1yr2 diagnosed with an STI (Natsal-3 definition) excl thrush in year before interview

1: yes if (whnchlam=1 | whngono=1 | whnsyphl=1 | whnherp=1 | whnwarts=1 | whntric=1 | whnnsu=1)

0: no if ((whnchlam=-1 | (whnchlam>=2 & whnchlam<=4)) & ((whngono=-1 | (whngono>=2 & whngono<=4)) & ((whnsyphl=-1 | (whnsyphl>=2 & whnsyphl<=4)) & ((whnherp=-1 | (whnherp>=2 & whnherp<=4)) & ((whnwarts=-1 | (whnwarts>=2 & whnwarts<=4)) & ((whntric=-1 | (whntric>=2 & whntric<=4)) & ((whnnsu=-1 | (whnnsu>=2 & whnnsu<=4))

9: not answered else if (whnchlam=9 | whngono=9 | whnsyphl=9 | whnherp=9 | whnwarts=9 | whntric=9 | whnnsu=9)

SEXUALLY TRANSMITTED INFECTIONS

(R) (ND) **nostis** number of STIs ever diagnosed with (excl. thrush), grouped
0: none *if* (diagnos1=11 | diagnos1=9) & diagnos2=-1
1: 1 *if* ((diagnos1≠-1 & diagnos1≠9 & diagnos1≠11 & diagnos1≠99) & diagnos2=-1) | ((diagnos1≠-1 & diagnos1≠9 & diagnos1≠11 & diagnos1≠99) & diagnos2=-1 | diagnos2=9) | (diagnos1=9 & diagnos2≠-1 & diagnos2≠9 & diagnos2≠11 & diagnos2≠99)
2: 2+ *if* (diagnos1≠-1 & diagnos1≠9 & diagnos1≠11 & diagnos1≠99 & diagnos2≠-1 & diagnos2≠9 & diagnos2≠11 & diagnos2≠99)
9: not answered *if* diagnos1=99
-1: not applicable *if* diagnos1=-1

Note: In Natsal-3 variables very similar to whenherp-whnwarts were directly obtained from the interview, so they have not been derived.

(R) (ND) **whnherpe** time since (last) diagnosed with genital herpes, grouped
-1: not applicable *if* herpes=-1
0: never *if* herpes=0
1: in same calendar year as interview *if* yrsherpe=1
2: 1-5 calendar years before interview *if* yrsherpe>=1 & yrsherpe<=5
3: >5 calendar years before interview *if* yrsherpe>5 & yrsherpe<99
9: not answered *if* yrsherpe=99

Note: Variables whntrich...whnpid are coded in a similar way to whnherpe

(R) (ND) **whntrich** time since (last) diagnosed with trichomonas, grouped
(R) (ND) **whngonor** time since (last) diagnosed with gonorrhoea, grouped
(R) (ND) **whnsyphl** time since (last) diagnosed with syphilis, grouped
(R) (ND) **whnchlam** time since (last) diagnosed with chlamydia, grouped
(R) (ND) **whnnsung** time since (last) diagnosed with NSU/NGU, grouped
(R) (ND) **whnwarts** time since (last) diagnosed with genital warts, grouped
(R) (ND) **whnpid** time since (last) diagnosed with PID, grouped

Note: in Natsal-3 respondents were not asked how many times they were diagnosed with STIs. Very few people had had more than one diagnosis in Natsal-2.

Note: respondents are only asked the number of times diagnosed with genital herpes, gonorrhoea, and chlamydia, thus diagno2 ... diagno5 are sti-specific

(R) (N) **diagno2** no. of times diagnosed with trichomonas, last 5 years
0 *if* trich=0 | (whndiag2>1900 & whndiag2<1995)
98: not answered but trichomonas diagnosis in last 5 years *if* whndiag2>=1995
99: not answered
-1: not applicable *if* trich=-1

Note: Variables diagno3...diagno5 are coded in a similar way to diagno2

(R) (N) **diagno3** no. of times diagnosed with gonorrhoea, last 5 years
(R) (N) **diagno4** no. of times diagnosed with syphilis, last 5 years
(R) (N) **diagno5** no. of times diagnosed with chlamydia, last 5 years

(R) (ND) **no_trich** no. of episodes of trichomonas, last 5 years, grouped
0: none *if* diagno2=0
1: once *if* diagno2=1
2: more than once *if* diagno2>=2 & diagno2<98
9: not answered *if* diagno2=98 | diagno2=99
-1: not applicable *if* diagno2=-1

Note: Variables no_gonor ... no_chlam are coded in a similar way to no_trich

(R) (ND) **no_gonor** no. of episodes of gonorrhoea, last 5 years, grouped
(R) (ND) **no_syph** no. of episodes of syphilis, last 5 years, grouped
(R) (ND) **no_chlam** no. of episodes of chlamydia, last 5 years, grouped

SEXUALLY TRANSMITTED INFECTIONS

Note: in Natsal-3 information was collected on where STIs were diagnosed for Chlamydia, Gonorrhoea, genital warts, syphilis, trichomonas vaginalis, genital herpes, NSU/NGU in separate questions detailed above.

Note: Variables diagwhe2... diagwhe5 are STI-specific so respondents were only asked where they were (last) treated for genital herpes, gonorrhoea, syphilis, and chlamydia, in the last 5 years

(R) (N) **diagwhe2** where (last) treated for trichomonas, last 5 years

1: gp surgery

2: nhs vd/std/sexual health clinic

3: nhs family planning clinic

4: private clinic/doctor

5: somewhere else

98: not answered but have had trichomonas in last 5 years if whndiag2>=1995

99: not answered

-1: not applicable if trich=-1 | 0 | (whndiag2>1900 & whndiag2<1995)

Note: Variables diagwhe3... diagwhe5 are coded in a similar way to diagwhe2

(R) (N) **diagwhe3** where (last) treated for gonorrhoea, last 5 years

(R) (N) **diagwhe4** where (last) treated for syphilis, last 5 years

(R) (N) **diagwhe5** where (last) treated for chlamydia, last 5 years

Note: information was collected in Natsal-3 on cervical screening and HPV vaccination (women only) and on current STI symptoms. The STI symptom data will link to the biological testing results.

(N) **everpap** when was last cervical smear (if female and aged >24)

-1: not applicable if rsex=1 | dimage<25

1: never had one

2: less than 3 years ago

3: between 3 and 5 years ago

4: between 5 and 10 years ago

5: more than 10 years ago

9: not answered

(N) **hpvever** been vaccinated against HPV (if female and aged>30)

-1: not applicable if rsex=1 | dimage>=30

1: yes – all 3 doses

2: yes – 1/2 doses but not all 3

3: no

9: not answered

(N) **hvpvcomp** intend to complete the other doses

-1: not applicable if rsex=1 | dimage>=30 | hpvever=1 | hpvever=3

1: yes

2: no

9: not answered

(N) **hpvofref** ever offered vaccination (if not received vaccination)

-1: not applicable if rsex=1 | dimage>=30 | hpvever=1 | hpvever=2

1: yes but refused

2: no never offered

9: not answered

SEXUALLY TRANSMITTED INFECTIONS

(N) **hpvwhere** where offered / received vaccination (women outside vaccination programme)

-1: not applicable *if* rsex=1 | dage>=30 | rdob later than 01/09/1990 | hpvever=3

1: GP

2: GUM clinic

3: private clinic / doctor including chemist / pharmacy

4: somewhere else

9: not answered

*Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code questions. The following binary variables were created directly from the question **stisymph** (new question in Natsal-3).*

(N) **mstipainuri** STI symptoms in the last month (men): pain, burning or stinging passing urine

-1: not applicable *if* rsex=2 | dage>44 | (hetlife=0 & samlife>0)

0: no

1: yes

9: not answered

mstipassmor..mstinone are all coded in a similar way to mstipainuri

(N) **mstipassmor** STI symptoms in the last month (men): more frequent urination

(N) **mstigenwart** STI symptoms in the last month (men): genital wart/lump

(N) **mstigenulc** STI symptoms in the last month (men): genital ulcer/sore

(N) **mstipdisch** STI symptoms in the last month (men): penile discharge

(N) **mstipaintest** STI symptoms in the last month (men): painful testicles

(N) **mstinone** STI symptoms in the last month (men): none

*The following binary variables were created directly from the question **stisymph** (new question in Natsal-3).*

(N) **fstipainuri** STI symptoms in the last month (women): pain, burning or stinging passing urine

-1: not applicable *if* rsex=1 | dage>44 | (hetlife=0 & samlife>0)

0: no

1: yes

9: not answered

fstipassmor..fstinone are all coded in a similar way to fstipainuri

(N) **fstipassmor** STI symptoms in the last month (women): more frequent urination

(N) **fstigenwart** STI symptoms in the last month (women): genital wart/lump

(N) **fstigenulc** STI symptoms in the last month (women): genital ulcer/sore

(N) **fstivdisch** STI symptoms in the last month (women): abnormal vaginal discharge

(N) **fstiodour** STI symptoms in the last month (women): odourous vaginal discharge

(N) **fstivpainsex** STI symptoms in the last month (women): vaginal pain during sex

(N) **fstibleedper** STI symptoms in the last month (women): bleeding between periods

(N) **fstibleedsex** STI symptoms in the last month (women): bleeding after sex

(N) **fstiabdopain** STI symptoms in the last month (women): lower abdominal / pelvic pain

(N) **fstinone** STI symptoms in the last month (women): none

(ND) **anystisymph** any STI symptoms in the last month (men)

-1: not applicable *if* mstipainuri=-1

1: yes *if* mstipainuri =1 | mstipassmor =1 | mstigenwart =1 | mstigenulc =1 | mstipdisch =1 | mstipaintest=1

0: no *if* mstipainuri =0 & mstipassmor =0 & mstigenwart =0 & mstigenulc =0 & mstipdisch =0 & mstipaintest=0

mstipaintest=0

9: not answered *if* mstipainuri=9

(ND) **anystisymph** any STI symptoms in the last month (women)

-1: not applicable *if* fstipainuri=-1

1: yes *if* fstipainuri=1 | fstipassmor=1 | fstigenwart=1 | fstigenulc=1 | fstivdisch=1 | fstiodour=1 |

fstivpainsex=1 | fstibleedper=1 | fstibleedsex=1 | fstiabdopain=1

0: no *if* fstipainuri=0 & fstipassmor=0 & fstigenwart=0 & fstigenulc=0 & fstivdisch=0 & fstiodour=0 &

fstivpainsex=0 & fstibleedper=0 & fstibleedsex=0 & fstiabdopain=0

9: not answered *if* fstipainuri=9

SEXUALLY TRANSMITTED INFECTIONS

(ND) **anystisypm** any STI symptoms in the last month (all)
compute anystisypm=anystisypm if rsex=1
compute anystisypm=anystisypm if rsex=1

INJECTING DRUG USE

Note: in Natsal-3 the questions in this section were moved to a new section on general drug use at the end of the CASI

Note: in Natsal-3 inject was modified to only include non-prescribed drugs or other substances

(R) (s17a) **inject** ever injected drugs or other substances
1: yes
2: no
9: not answered
-1: not applicable *if scelig=2*

(R) (s17b1) **prescrb** injected drugs prescribed by a doctor for a medical condition
1: all prescribed
2: some prescribed
3: none prescribed
9: not answered
-1: not applicable *if scelig=2 | inject=2*

(R) (s17b *U) **medcond1** medical condition 1 for which the drugs were prescribed
1: diabetes
2: allergic disorders
3: chronic pain
4: thrombosis/embolus
5: infections
6: psychiatric illness
7: drug abuse
8: other
99: not answered
-1: not applicable *if scelig=2 | inject=2 | prescrb=3 | (medcondn=99 & medcondn-1≠99)*

Similarly for up to 3 medical conditions:

(R) (s17b *U) **medcond2** medical condition 2 for which the drugs were prescribed

(R) (s17b *U) **medcond3** medical condition 3 for which the drugs were prescribed

(R) (q17b1 D) **diabetes** injected prescribed drugs for diabetes
1: yes *if medcond1=1 | medcond2=1 | medcond3=1*
0: no *else if medcond1>=1 & medcond1<=8*
9: not answered *if missing*
-1: not applicable *if medcond1=-1 & medcond2=-1 & medcond3=-1*

Note: Variables allergy ... othcond are derived from medcond1 ... medcond3, and are coded in a similar way to diabetes:

(R) (q17bb *D) **allergy** injected prescribed drugs for allergies

SEXUALLY TRANSMITTED INFECTIONS

- (R) (ND) **chnrpain** injected prescribed drugs for chronic pain
- (R) (ND) **thrombos** injected prescribed drugs for thrombosis
- (R) (ND) **infectns** injected prescribed drugs for infections
- (R) (ND) **psychiat** injected prescribed drugs for psychiatric illness
- (R) (ND) **addictn** injected prescribed drugs for drug addiction
- (R) (q17be D) **othconds** injected prescribed drugs for other medical condition(s)

Note: in Natsal-3 the variable drugs was directly obtained from the questionnaire and is now the variable inject.

Note: Variable drugs in Natsal-1 is coded 2 for 'yes' and 1 for 'no', which is inconsistent with the other 'yes/no' variables, so the codes must be reversed in Natsal-1 to be consistent with Natsal-2:

(R) (D) **drugs** ever injected non-prescribed drugs

1: yes *if* inject=1 & (prescrb=2 | prescrb=3)

2: no *if* inject=2 | (inject=1 & prescrb=1)

9: not answered *if* missing

-1: not applicable *if* scelig=2

HIV TESTING

HIV TESTING

Note: in Natsal-3 questions slightly revised to reflect change in age of respondents and donation questions not asked of 16 year olds as too young to give blood.

(R) (N) **donblod** donated blood since beginning of 1986

-1: not applicable if scelig=2

1: yes

2: no

9: not answered

(N) **donblod2** donated blood

1: yes

2: no

9: not answered

-1: not applicable if scelign3=2 | dage=16

(R) (N) **whendon** when last donated blood (since the beginning of 1986)

1: in last year

2: between 1 and 2 years ago

3: between 2 and 5 years ago

4: longer than 5 years ago but since the beginning of 1986

9: not answered

-1: not applicable if scelig=2 | donblod=2

(N) **whendon2** when last donated blood

1: in last year

2: between 1 and 2 years ago

3: between 2 and 5 years ago

4: longer than 5 years and 10 years ago

5: longer than 10 years ago but since the beginning of 1986

6: longer ago than the beginning of 1986

9: not answered

-1: not applicable if scelign3=2 | dimage=16 | donblod2=2

Note: in Natsal-3 no longer thought to be necessary to have a prefer not to answer question for HIV testing.

(U) (N) **hivtsty** ever had a blood test for HIV other than when giving blood

1: yes

2: no

3: maybe/not sure

(U – code removed) 4: prefer not to answer

9: not answered

(U) -1: not applicable if scelign3=2 | donblod2=2 | dimage=16

-1: not applicable if scelig=2 | donblod=2

(U) (N) **hivtstn** ever had a blood test for HIV (if never given blood)

1: yes

2: no

3: maybe/not sure

(U – code removed) 4: prefer not to answer

9: not answered

(U) -1: not applicable if scelign3=2 | donblod2=1

-1: not applicable if scelig=2 | donblod=1

HIV TESTING

Note: Variable hivtest refers to whether/not ever had a HIV test excluding when giving blood.

(U) (ND) **hivtest** ever had a blood test for HIV (excluding when given blood)

1: yes if ((donblod=1 | donblod=9) & hivtsty=1) | ((donblod=2 | donblod=9) & hivtstn=1)

2: no if ((donblod=1 | donblod=9) & hivtsty=2) | ((donblod=2 | donblod=9) & hivtstn=2)

3: maybe/not sure if ((donblod=1 | donblod=9) & hivtsty=3) | ((donblod=2 | donblod=9) & hivtstn=3)

(U – code removed) 4: prefer not to answer if ((donblod=1 | donblod=9) & hivtsty=4) | ((donblod=2 | donblod=9) & hivtstn=4)

9: not answered if missing if ((donblod=1 | donblod=9) & hivtsty=9) | ((donblod=2 | donblod=9) & hivtstn=9)

(U) -1: not applicable if scalign3=2 -1: not applicable if scelig=2

*Note: This codebook gives only the binary variables to be used for analysis, not the original multi-code questions. The following binary variables were created directly from the question **whytest** in Natsal-3, and derived (as shown) from **whytest1-whytest5** in Natsal-2. Testinsr to testothr are coded in the same way as testpreg.*

(s20bb D) **testpreg** HIV test reason: pregnancy

1: yes if whytest1=1 | whytest2=1 | whytest3=1 | whytest4=1 | whytest5=1

0: no else if whytest1>=1 & whytest1<=5

9: not answered if missing

(U) -1: not applicable if scalign3=2 | hivtest=2 | hivtest=3

-1: not applicable if whytest1=-1 & whytest2=-1 & whytest3=-1 & whytest4=-1 & whytest5=-1

(s20bc D) **testinsr** HIV test reason: travel/insurance/mortgage

(N) **testschk** HIV test reason: sexual health check

(ND) **testhchk** HIV test reason: general health check

(N) **testcond** HIV test reason: stopped using condoms

(ND) **testrisk** HIV test reason: concern of risk to self/partner

(N) **testdoc** HIV test reason: doctor's advice

Note: Variable testothr is derived differently to Natsal-1's definition of 'other reason' for HIV test

(R) (ND) **testothr** HIV test reason: other reason(s)

1: yes if whytest1=5 | whytest2=5 | whytest3=5 | whytest4=5 | whytest5=5

0: no else if whytest1>=1 & whytest1<=5

9: not answered if missing

-1: not applicable if whytest1=-1 & whytest2=-1 & whytest3=-1 & whytest4=-1 & whytest5=-1

Note: The 'other reason' variable testothr2 is not directly comparable with Natsal-2's testothr because more response options were available for respondents to choose from at whytest in Natsal-3, hence it has been renamed.

(N) **testothr2** HIV test reason: other reason

HIV TESTING

Note: Variable 'testoth2' is consistent with Natsal-1's definition of 'other reason' for HIV test:

(U – see below) (s20bd D) **testoth2** HIV test reason: other (incl. general health check & concern about risk)

1: yes if testhchk=1 | testrisk=1 | testothr=1

0: no else if testhchk=0 | testhchk=1 | testrisk=0 | testrisk=1 | testothr=0 | testothr=1

9: not answered if testhchk=9 & testrisk=9 & testothr=9

-1: not applicable if testhchk=-1 & testrisk=-1 & testothr=-1

testoth2 HIV test reason: other (incl. general health check, concern about risk, sexual health check, stopped using condoms, doctor's advice)

1: yes if testhchk=1 | testrisk=1 | testothr=1 | testschk=1 | testcond=1 | testdoc=1

0: no else if testhchk=0 | testhchk=1 | testrisk=0 | testrisk=1 | testothr=0 | testothr=1 | testschk=0 | testcond=0 | testdoc=0

9: not answered if testhchk=9 & testrisk=9 & testothr=9

-1: not applicable if testhchk=-1 & testrisk=-1 & testothr=-1

(N) **whntest** when was blood test for HIV (last one if more than one)

1: in the last year

2: between 1 year and 2 years ago

3: between 2 years and 5 years ago

4: longer than 5 years ago

9: not answered

(U) -1: not applicable if scelign3=2 | hivtest=2 -1: not applicable if scelign=2 | hivtest=2 | hivtest=3 | hivtest=4

Note: Variable 'whntest2' is consistent with Natsal-1's definition of s20c in terms of time frames. However, in Natsal-1 question s20c was asked to everyone –including those who only had a HIV test as a blood donor. Consequently, comparisons between surveys need to ensure that Natsal-1 data are limited to those who had an HIV test aside from as a blood donor.

(s20c D) **whntest2** when was blood test for HIV (last one if more than one)

1: in last year if whntest=1

2: 1-2 years ago if whntest=2

3: >2 years ago if whntest=3 | whntest=4

9: not answered if whntest=9

-1: not applicable if whntest=-1

Note: Variable 'hivtest5' is equivalent to variable 's20a' in Natsal-1 in that it refers to having a HIV test in the last 5 years not ever. Unlike Natsal-1, this derivation of hivtest5 excludes from the numerator those who only had a HIV test as a blood donor

(s20a D*) **hivtest5** had a blood test for HIV, last 5 years

1: yes if whntest=1 | whntest=2 | whntest=3

0: no if whntest=4 | hivtest=2

9: not answered if hivtest=9 | hivtest=3 | whntest=9

-1: not applicable if hivtest=-1

(ND) **hivtestly** had a blood test for HIV, last year

1: yes if whntest=1

0: no if whntest=2 | whntest=3 | whntest=4 | hivtest=2

9: not answered if hivtest=9 | hivtest=3 | whntest=9

-1: not applicable if hivtest=-1

HIV TESTING

(U – see below) (N) **whertest** where had blood test for HIV (last one if more than one)

- 1: GP surgery
- 2: NHS VD/STD/sexual health clinic
- 3: NHS family planning clinic
- 4: private clinic or doctor
- 5: somewhere else
- 9: not answered
- 1: not applicable *if* scelig=2 | hivtest=2 | hivtest=3 | hivtest=4

whertest where had blood test for HIV (last one if more than one) – NOTE question identical but answer options have changed

- 1: GP surgery
- 2: GUM clinic
- 3: NHS family planning clinic
- 4: antenatal clinic
- 5: private clinic or doctor
- 6: internet site offering postal kit
- 7: youth advisory clinic
- 8: termination of pregnancy clinic
- 9: hospital A&E
- 10: somewhere else
- 99: not answered
- 1: not applicable *if* scelig=2 | hivtest=2

Note: Natsal-2 did not ask respondents what the result was, just whether or not they were given the result.

(N) **retest** given result of blood test for HIV

- 1: yes
- 2: no
- 9: not answered
- 1: not applicable *if* scelig=2 | hivtest=2

SEXUAL FUNCTION

SEXUAL PROBLEMS

Note: this section was extensively revised in Natsal-3 and a psychometric measure of sexual function was developed. This section has been renamed as sexual function and appears below the Natsal-2 Sexual problems section (page 118).

(R) (N) **ifdysf** any sex 'problems' lasting 1+ month(s)+, last year

1: yes

2: no

9: not answered

-1: not applicable if scelig=2

(R) (N) **whdysf1** sex 'problem' 1 lasting 1+ month(s), last year

1: lacked interest

2: anxiety about sexual performance

3: unable to experience orgasm

4: premature orgasm

5: physically painful intercourse

6: problems achieving/maintaining erections (*only if male*)

7: trouble lubricating (*only if female*)

9: not answered

-1: not applicable if scelig=2 | ifdysf=2 | (whdysfn=9 & whdysfn-1 ≠9)

Similarly for up to 6 problems:

(R) (N) **whdysf2** sex 'problem' 2 lasting 1+ month(s), last year

(R) (N) **whdysf3** sex 'problem' 3 lasting 1+ month(s), last year

(R) (N) **whdysf4** sex 'problem' 4 lasting 1+ month(s), last year

(R) (N) **whdysf5** sex 'problem' 5 lasting 1+ month(s), last year

(R) (N) **whdysf6** sex 'problem' 6 lasting 1+ month(s), last year

(R) (ND) **lackint** lacked interest in having sex for 1+ month(s), last year

1: yes if whdysf1=1 | whdysf2=1 | whdysf3=1 | whdysf4=1 | whdysf5=1 | whdysf6=1

0: no else if whdysf1>=1 & whdysf1<=7

9: not answered if missing

-1: not applicable if whdysf1=-1 & whdysf2=-1 & whdysf3=-1 & whdysf4=-1 & whdysf5=-1 & whdysf6=-1

Note: Variables anxiety ... painsic are derived from whdysf1 ... whdysf6 in a similar way to lackint:

(R) (ND) **anxiety** anxiety about sexual performance for 1+ month(s), last year

(R) (ND) **orgasmno** unable to experience orgasm for 1+ month(s), last year

(R) (ND) **preorgsm** premature orgasm for 1+ month(s), last year

(R) (ND) **painsic** physically painful intercourse for 1+ month(s), last year

(R) (ND) **erection** problems achieving/maintaining erections for 1+ month(s), last year

1: yes if whdysf1=6 | whdysf2=6 | whdysf3=6 | whdysf4=6 | whdysf5=6 | whdysf6=6

0: no else if whdysf1>=1 & whdysf1<=7

9: not answered if missing

-1: not applicable if (whdysf1=-1 & whdysf2=-1 & whdysf3=-1 & whdysf4=-1 & whdysf5=-1 & whdysf6=-1) |

rsex=2

(R) (ND) **lubricat** trouble lubricating for 1+ month(s), last year

1: yes if whdysf1=7 | whdysf2=7 | whdysf3=7 | whdysf4=7 | whdysf5=7 | whdysf6=7

0: no else if whdysf1>=1 & whdysf1<=7

9: not answered if missing

-1: not applicable if (whdysf1=-1 & whdysf2=-1 & whdysf3=-1 & whdysf4=-1 & whdysf5=-1 & whdysf6=-1) |

rsex=1

SEXUAL FUNCTION

(R) (ND) **noofprob** no. of sexual 'problems' lasting 1+ months(s), last year
count (lackint anxiety orgasmno preorgsm painsic erection lubricat) if (lackint=0 | 1 & anxiety=0 | 1 & orgasmno=0 | 1 & preorgsm=0 | 1 & painsic=0 | 1 & erection=0 | 1 & lubricat=0 | 1)
9: not answered if lackint=9 | anxiety=9 | orgasmno=9 | preorgsm=9 | painsic=9 | erection=9 | lubricat=9
-1: not applicable if scelig=2

Note: Variables longdys1 ... longdys7 are sex-problem-specific variables

(R) (N) **longdys1** length of time lacked interest in sex

- 1: at least 1 month but less than 3 months
- 2: at least 3 months but less than 6 months
- 3: at least 6 months but less than 1 year
- 4: 1 year or longer
- 9: not answered
- 1: not applicable if lackint=-1 | lackint=0

Note: longdys2 ... longdys7 are coded in a similar way to longdys1

(R) (N) **longdys2** length of time experienced anxiety about sexual performance

(R) (N) **longdys3** length of time been unable to experience orgasm

(R) (N) **longdys4** length of time experienced premature orgasm

(R) (N) **longdys5** length of time experienced physically painful intercourse

(R) (N) **longdys6** length of time had problems achieving/maintaining erections

(R) (N) **longdys7** length of time had trouble lubricating

(R) (ND) **longdysg** length of time experienced any sexual 'problem(s)', last year

0: no sexual 'problem(s)' lasting 1+ month(s), last year if ifdysf=2

1: at least 1 month but less than 3 months if longdys1=1 | longdys2=1 | longdys3=1 | longdys4=1 | longdys5=1 | longdys6=1 | longdys7=1

2: at least 3 months but less than 6 months else if longdys1=2 | longdys2=2 | longdys3=2 | longdys4=2 | longdys5=2 | longdys6=2 | longdys7=2

3: at least 6 months but less than 1 year else if longdys1=3 | longdys2=3 | longdys3=3 | longdys4=3 | longdys5=3 | longdys6=3 | longdys7=3

4: 1 year or longer else if longdys1=4 | longdys2=4 | longdys3=4 | longdys4=4 | longdys5=4 | longdys6=4 | longdys7=4

9: not answered if ifdys1=9

-1: not eligible if ifdys1=-1

(R) (ND) **longdy1g** lacked interest in having sex for 6+ months, last year

1: yes if longdys1=3 | longdys1=4

0: no if longdys1=-1 | longdys1=1 | longdys1=2

9: not answered if longdys1=9

-1: not applicable if ifdysf=-1

Note: Variables longdy2g ... longdy7g are derived in a similar way to longdy1g

(R) (N) **longdy2g** anxiety about sexual performance for 6+ months, last year

(R) (N) **longdy3g** unable to experience orgasm for 6+ months, last year

(R) (N) **longdy4g** experienced premature orgasm for 6+ months, last year

(R) (N) **longdy5g** experienced physically painful intercourse for 6+ months, last year

(R) (N) **longdy6g** problems achieving/maintaining erections for 6+ months, last year

(R) (N) **longdy7g** trouble lubricating for 6+ months, last year

(R) (ND) **noofpro2** no. of sexual 'problems' lasting 6+ months(s), last year

count (longdy1g longdy2g longdy3g longdy4g longdy5g longdy6g longdy7g) if (longdy1g=0 | longdy1g=1 & longdy2g=0 | longdy2g=1 & ... & longdy7g=0 | longdy7g=1)

9: not answered if longdy1=9 | ... | longdy7g=9

-1: not applicable if scelig=2

SEXUAL FUNCTION

(R) (N) **avoisex** avoided sex during last year because of sexual 'problem(s)'

1: yes

2: no

9: not answered

-1: not applicable if scelig=2 | ifdysf=2

Note: Variables contact1...contact4 are not sexual-problem-specific as question asks 'have you contacted any of the following for help with either/any of the/these conditions?'

(R) (N) **contact1** contact 1 for sexual 'problem(s)', last year

-1: not applicable if scelig=2 | ifdysf=2 | (contactn=99 & contactn-1 ≠99)

1: GP

2: VD/STD/sexual health clinic

3: psychiatrist/psychologist

4: marriage counsellor

5: other type of clinic/doctor

6: contacted a help-line

7: someone else

8: none of these

99: not answered

Similarly for up to 4 contacts:

(R) (N) **contact2** contact 2 for sexual 'problem(s)', last year

(R) (N) **contact3** contact 3 for sexual 'problem(s)', last year

(R) (N) **contact4** contact 4 for sexual 'problem(s)', last year

(R) (ND) **spgp** contacted GP about sexual 'problem(s)', last year

1: yes if contact1=1 | contact2=1 | contact3=1 | contact4=1

0: no else if contact1>=1 & contact1<=8

9: not answered if missing

-1: not applicable if contact1=-1 & contact2=-1 & contact3=-1 & contact4=-1

Note: Variables spstdcl ... spnone are derived in a similar way to spgp:

(R) (ND) **spstdcl** contacted VD/STD clinic about sexual 'problem(s)', last year

(R) (ND) **sppsyh** contacted a psychiatrist/psychologist about sexual 'problem(s)', last year

(R) (ND) **spmarc** contacted marriage counsellor about sexual 'problem(s)', last year

(R) (ND) **spothrgp** contacted other type of clinic/doctor about sexual 'problem(s)', last year

(R) (ND) **sphelp** contacted a help-line about sexual 'problem(s)', last year

(R) (ND) **spelse** contacted someone else about sexual 'problem(s)', last year

(R) (ND) **spnone** contacted none of these sources about sexual 'problem(s)', last year

(R) (N) **ifcon1** NHS/private GP contacted about sexual 'problem(s)', last year

1: NHS

2: private

3: other (e.g. voluntary organisation)

4: don't know

9: not answered

-1: not applicable if scelig=2 | ifdysf=2 | spgp=0 | spgp=-1

Note: Variables ifcon2...ifcon5 are sex-problem contact-specific variables and are coded in a similar way to ifcon1:

(R) (N) **ifcon2** NHS/private VD/STD clinic contacted about sexual 'problem(s)', last year

(R) (N) **ifcon3** NHS/private psychiatrist/psychologist contacted about sexual 'problem(s)', last year

(R) (N) **ifcon4** NHS/private marriage counsellor contacted about sexual 'problem(s)', last year

(R) (N) **ifcon5** NHS/private other type of clinic/doctor contacted about sexual 'problem(s)', last year

SEXUAL FUNCTION

SEXUAL FUNCTION

The 17-item Natsal-SF measure provides an indication of an individual's level of sexual function, taking into account reported function problems, the relational context as well as levels of satisfaction and distress. As such it adopts a psycho-social perspective, recognising that sexual function is about much more than the experience of biomedical difficulties with sexual response.

Not all variables listed in this section are part of the measure. The table shows which variables are included in the measure and who is routed to them.

Variables that form the SF measure	Who is included
<i>sfprb</i>	<i>All respondents who reported sex in the last year (het1yr>0 or sam1yr>0), except for those aged 16-17 who reported only oral sex (ever)</i>
<i>sfrlnt, sfrlikes, sfrpdf, sfrem</i>	<i>All respondents who reported sex in the last year (het1yr>0 or sam1yr>0) and who were in a relationship that has lasted for all of the last year (based on information about most recent partners), except for those aged 16-17 who reported only oral sex (ever)</i>
<i>SFFavoid, spinternet to spnohelp</i>	<i>All, except those aged 16-17 who reported only oral sex (ever)</i>
<i>SFFSatis, SFFDst</i>	<i>All in CASI</i>

Please note: *Only people who reported sex in the last year and were in a relationship answer all questions on the measure. Individuals who have had sex in the last year but who have not been in a relationship do not answer the four questions on partnership and individuals who have not been sexually active in the last year only answer SFFSatis and SFFDst, but their missing answers (what they would have answered had they been in a relationship or been sexually active) are estimated using sophisticated modelling techniques.*

Scoring the measure – derived variables:

Respondents are categorised as either low functioning, normal functioning, or not sexually active. The cut off point for low function is determined by the lowest gender-specific quintile.

This approach was used to create the derived variables given on the next page (lowsfscore..sdmean).

For further information about how the conceptual basis of the measure, please see Mitchell et al 2011. For information about the development and validation of the measure, please see Mitchell et al. 2012.

Mitchell K., Ploubidis, G., Datta, J., Wellings, K. (2012) *The Natsal-SF: A validated measure of sexual function for use in community surveys. European Journal of Epidemiology. DOI 10.1007/s10654-012-9697-3*

Mitchell, K. & Wellings, K. (2011). *Measuring sexual function in community surveys: Development of a conceptual framework. Journal of Sex Research. DOI:10.1080/00224499.2011.621038*

SEXUAL FUNCTION

Natsal-SF measure variables

(N) (D) **sfscore** Natsal sexual function measure score (lower scores = lower function)

99999: not available

(N) (D) **lowerfuncofall** Low function according to the Natsal Sexual Function Measure

0: 'Normal' function ((sfscore>-0.718 & rsex==1 & sfscore!=99999 & tot1yr!=999 & tot1yr!=0) | (sfscore>-0.772 & rsex==2 & sfscore!=99999 & tot1yr!=999 & tot1yr!=0))

1: 'Low' function ((sfscore<=-0.718 & rsex==1 & sfscore!=99999 & tot1yr!=999 & tot1yr!=0) | (sfscore<=-0.772 & rsex==2 & sfscore!=99999 & tot1yr!=999 & tot1yr!=0))

8: Not sexually active (sfscore!=99999 & tot1yr!=999 & tot1yr==0)

9: Not available (sfscore==99999 | tot1yr==999)

Other Natsal sexual function variables

Note: Respondents aged 16 & 17 who had only had oral sex were only asked sffsatis, sfdst, frequent and easyoua/easyoub as other questions were felt to be irrelevant to this group.

(N) **hafsex** health condition / disability in last year that has affected sexual activity or enjoyment

1: yes

2: no

9: not answered

-1: not applicable if scalign3=2 | (hetlife=0 & samlife=0) | (hetlife=999 & samlife=9999) | (dage<18 & lastvag=7 & analsex=7 & (samahim=7 | samahim=-1) & (samayou=7 | samayou=-1))

(N) **mafsex** medications in last year that have limited sexual activity or enjoyment

1: yes

2: no

9: not answered

-1: not applicable else if scalign3=2 | (hetlife=0 & samlife=0) | (hetlife=999 & samlife=9999) | (dage<18 & lastvag=7 & analsex=7 & (samahim=7 | samahim=-1) & (samayou=7 | samayou=-1))

Note: respondents were only asked about sexual difficulties if had had sex in the year before interview.

Note: Binary variables lackint3m..premorg3m were created directly from the question sfrpb. All vars are coded similarly to lackint3m.

(N) **lackint3m** lacked interest in having sex for 3+ months, last year

1: yes

0: no

9: not answered

-1: not applicable if scalign3=2 | (het1yr<1 & sam1yr<1) | (het1yr=999 & sam1yr=999) | (dage<18 & lastvag=7 & analsex=7 & (samahim=7 | samahim=-1) & (samayou=7 | samayou=-1))

And similarly...

(N) **lackenj3m** lacked enjoyment in having sex for 3+ months, last year

(N) **anxsex3m** anxiety during sex for 3+ months, last year

(N) **painsex3m** physical pain as a result of sex for 3+ months, last year

(N) **nonarou3m** no excitement / arousal during sex for 3+ months, last year

(N) **nolgorg3m** no orgasm or took a long time to reach orgasm during sex for 3+ months, last year

(N) **premorg3m** reached orgasm more quickly than would have liked during sex for 3+ months, last year

lubric3m was only a response option for women

(N) **lubric3m** uncomfortably dry vagina for 3+ months, last year

1: yes

SEXUAL FUNCTION

0: no

9: not answered

-1: not applicable *if* $\text{scelign3}=2 \mid (\text{het1yr}<1 \ \& \ \text{sam1yr}<1) \mid (\text{het1yr}=999 \ \& \ \text{sam1yr}=999) \mid (\text{dage}<18 \ \& \ \text{lastvag}=7 \ \& \ \text{analsex}=7 \ \& \ (\text{samahim}=7 \mid \text{samahim}=-1) \ \& \ (\text{samayou}=7 \mid \text{samayou}=-1)) \mid \text{rsex}=1$

erect3m was only a response option for men

(N) **erect3m** trouble achieving / maintaining erections for 3+ months, last year

1: yes

0: no

9: not answered

-1: not applicable *if* $\text{scelign3}=2 \mid (\text{het1yr}<1 \ \& \ \text{sam1yr}<1) \mid (\text{het1yr}=999 \ \& \ \text{sam1yr}=999) \mid (\text{dage}<18 \ \& \ \text{lastvag}=7 \ \& \ \text{analsex}=7 \ \& \ (\text{samahim}=7 \mid \text{samahim}=-1) \ \& \ (\text{samayou}=7 \mid \text{samayou}=-1)) \mid \text{rsex}=2$

(N) **nosfprb3m** no experience of any of these for 3+ months, last year

1: yes

0: no

9: not answered

-1: not applicable *if* $\text{scelign3}=2 \mid (\text{het1yr}<1 \ \& \ \text{sam1yr}<1) \mid (\text{het1yr}=999 \ \& \ \text{sam1yr}=999) \mid (\text{dage}<18 \ \& \ \text{lastvag}=7 \ \& \ \text{analsex}=7 \ \& \ (\text{samahim}=7 \mid \text{samahim}=-1) \ \& \ (\text{samayou}=7 \mid \text{samayou}=-1))$

(ND) **oneplus3m** experienced one or more sexual response problems

-1: not applicable *if* $\text{lackint3m}=-1$

0: no *if* $\text{rsex}=1 \ \&$

$(\text{lackint3m}+\text{lackenj3m}+\text{anxsex3m}+\text{painsex3m}+\text{nonarou3m}+\text{nolgorg3m}+\text{premorg3m}+\text{erect3m}=0) \ \& \ \text{lackint3m} \neq -1 \ \& \ \text{lackint3m} \neq 9) \mid (\text{rsex}=2 \ \&$

$(\text{lackint3m}+\text{lackenj3m}+\text{anxsex3m}+\text{painsex3m}+\text{nonarou3m}+\text{nolgorg3m}+\text{premorg3m}+\text{lubric3m}=0) \ \& \ \text{lackint3m} \neq -1 \ \& \ \text{lackint3m} \neq 9)$

1: yes *if* $\text{rsex}=1 \ \&$

$(\text{lackint3m}+\text{lackenj3m}+\text{anxsex3m}+\text{painsex3m}+\text{nonarou3m}+\text{nolgorg3m}+\text{premorg3m}+\text{erect3m} \geq 1) \ \& \ \text{lackint3m} \neq -1 \ \& \ \text{lackint3m} \neq 9) \mid (\text{rsex}=2 \ \&$

$(\text{lackint3m}+\text{lackenj3m}+\text{anxsex3m}+\text{painsex3m}+\text{nonarou3m}+\text{nolgorg3m}+\text{premorg3m}+\text{lubric3m} \geq 1) \ \& \ \text{lackint3m} \neq -1 \ \& \ \text{lackint3m} \neq 9)$

9: not answered *if* $\text{lackint3m}=9$

(D) **twoplus3m** experienced two or more sexual response problems

-1: not applicable *if* $\text{lackint3m}=-1$

0: no *if* $\text{rsex}=1 \ \&$

$(\text{lackint3m}+\text{lackenj3m}+\text{anxsex3m}+\text{painsex3m}+\text{nonarou3m}+\text{nolgorg3m}+\text{premorg3m}+\text{erect3m}=0 \ \text{or} \ 1) \ \& \ \text{lackint3m} \neq -1 \ \& \ \text{lackint3m} \neq 9) \mid (\text{rsex}=2 \ \&$

$(\text{lackint3m}+\text{lackenj3m}+\text{anxsex3m}+\text{painsex3m}+\text{nonarou3m}+\text{nolgorg3m}+\text{premorg3m}+\text{lubric3m}=0 \ \text{or} \ 1) \ \& \ \text{lackint3m} \neq -1 \ \& \ \text{lackint3m} \neq 9)$

1: yes *if* $\text{rsex}=1 \ \&$

$(\text{lackint3m}+\text{lackenj3m}+\text{anxsex3m}+\text{painsex3m}+\text{nonarou3m}+\text{nolgorg3m}+\text{premorg3m}+\text{erect3m} \geq 2) \ \& \ \text{lackint3m} \neq -1 \ \& \ \text{lackint3m} \neq 9) \mid (\text{rsex}=2 \ \&$

$(\text{lackint3m}+\text{lackenj3m}+\text{anxsex3m}+\text{painsex3m}+\text{nonarou3m}+\text{nolgorg3m}+\text{premorg3m}+\text{lubric3m} \geq 2) \ \& \ \text{lackint3m} \neq -1 \ \& \ \text{lackint3m} \neq 9)$

9: not answered *if* $\text{lackint3m}=9$

Note: sfprlng, SFprfrq, sfprbds, asked for each reported problem with suffix 2 – 9. wording for sfprfrq is slightly different to be applicable but coding is the same.

(N) **sfprlng** how long experiencing lacking interest

1: 3-6 months

2: 6 months – 1 year

3: 1 – 5 years

4: 5+ years

9: not answered

-1: not applicable *if* $\text{lackint3m}=-1 \mid \text{lackint3m}=0$

SEXUAL FUNCTION

- (N) **sfprlng2** how long experiencing lacking enjoyment
- (N) **sfprlng3** how long experiencing anxiousness
- (N) **sfprlng4** how long experiencing physical pain
- (N) **sfprlng5** how long experiencing lack of excitement / arousal during sex
- (N) **sfprlng6** how long experiencing taking a long time to orgasm / not orgasming
- (N) **sfprlng7** how long experiencing orgasming too quickly
- (N) **sfprlng8** how long experiencing uncomfortably dry vagina
- (N) **sfprlng9** how long experiencing trouble getting / keeping erection

(N) **sfprfrq** frequency experiencing lacking interest

- 1: always
- 2: very often
- 3: sometimes
- 4: not very often
- 9: not answered
- 1: not applicable *if* lackint3m=-1 | lackint3m=0

(N) **sfprfrq2** frequency experiencing lacking enjoyment

- (N) **sfprfrq3** frequency experiencing anxiousness
- (N) **sfprfrq4** frequency experiencing physical pain
- (N) **sfprfrq5** frequency experiencing lack of excitement / arousal during sex
- (N) **sfprfrq6** frequency experiencing taking a long time to orgasm / not orgasming
- (N) **sfprfrq7** frequency experiencing orgasming too quickly
- (N) **sfprfrq8** frequency experiencing uncomfortably dry vagina
- (N) **sfprfrq9** frequency experiencing trouble getting / keeping erection

(N) **sfprbds** degree of distress about lacking interest

- 1: not at all distressed
- 2: little distressed
- 3: fairly distressed
- 4: very distressed9: not answered
- 1: not applicable *if* lackint3m=-1 | lackint3m=0

(N) **sfprbds2** degree of distress about lacking enjoyment

- (N) **sfprbds3** degree of distress about anxiousness
- (N) **sfprbds4** degree of distress about physical pain
- (N) **sfprbds5** degree of distress about lack of excitement / arousal during sex
- (N) **sfprbds6** degree of distress about taking a long time to orgasm / not orgasming
- (N) **sfprbds7** degree of distress about orgasming too quickly
- (N) **sfprbds8** degree of distress about uncomfortably dry vagina
- (N) **sfprbds9** degree of distress about trouble getting / keeping erection

Note: sfrint – relsat are asked only of respondents who have had sex in the last year AND are in a relationship that has lasted for all of the last year.

(N) **sfrint** share same level of interest in having sex as partner

- 1: agree strongly
- 2: agree
- 3: neither agree or disagree
- 4: disagree
- 5: disagree strongly
- 9: not answered
- 1: not applicable *if* scelign3=2 | ((het1yr=0 | het1yr=999) & (sam1yr=0 | sam1yr=999)) | NOT (((r1rel=1 | r1rel=2) & (r1sexagn=1 | r1sexagn=2) & dfover1p1=1) | ((r1rel2=1 | r1rel2=2) & (r1sexagn2=1 | r1sexagn2=2) & dfover1p2=1) | ((r1rel3=1 | r1rel3=2) & (r1sexagn3=1 | r1sexagn3=2) & dfover1p3=1))

Note: sfrlikes and sfrpdfhave same answer options and coding as sfrint

SEXUAL FUNCTION

(N) **sflikes** share same sexual likes and dislikes as partner
(N) **sfrpdf** partner experienced sexual difficulties, last year

(N) **sfrem** emotionally close to partner when have sex

- 1: always
- 2: most of the time
- 3: sometimes
- 4: not very often
- 5: hardly ever
- 9: not answered
- 1: not applicable *if sfrint=-1*

(ND) **sfrint2** participant does not share same level of interest in sex as partner (binary)

- 1: not applicable *if sfrint=-1*
- 0: no/other *if sfrint=1 | sfrint=2 | sfrint=3*
- 1: yes/other *if sfrint=4 | sfrint=5*
- 9: not answered *if sfrint=-1*

(ND) **sflikes2** participant does not share same likes and dislikes as partner (binary)

- 1: not applicable *if sflikes=-1*
- 0: no/other *if sflikes=1 | sflikes=2 | sflikes=3*
- 1: yes *if sflikes=4 | sflikes=5*
- 9: not answered *if sflikes=-1*

(ND) **sfpdf2** partner experienced sexual difficulties, last year (binary)

- 1: not applicable *if sfpdf=-1*
- 0: no/other *if sfpdf=1 | sfpdf=2 | sfpdf=3*
- 1: yes *if sfpdf=4 | sfpdf=5*
- 9: not answered *if sfpdf=-1*

(ND) **sfrem2** participant does not emotionally close to partner when have sex (binary)

- 1: not applicable *if sfem=-1*
- 0: no/other *if sfem=1 | sfem=2 | sfem=3*
- 1: yes *if sfem=4 | sfem=5*
- 9: not answered *if sfem=-1*

(N) **relsat** how happy in relationship with partner (scale of 1 – 7; happy – unhappy)

- 9: not answered
- 1: not applicable *if sfrint=-1*

(ND) **relsat2** happy with relationship

- 1: not applicable *if relsat=-1*
- 1: yes *if relsat=1 | relsat=2*
- 0: other *if relsat=3 | relsat=4 | relsat=5 | relsat=6 | relsat=7*
- 9: not answered *if relsat=9*

Note: respondents are asked to think about their relationship in the past year when answering sffsatis..sffavoid.

(N) **sffsatis** “Feel satisfied with sex life”

- 1: agree strongly
- 2: agree
- 3: neither agree or disagree
- 4: disagree
- 5: disagree strongly
- 9: not answered
- 1: not applicable *if scelign3=2 | ((hetlife=0 | hetlife=9999) & (samlife=0 | samlife=9999))*

Note: sfdst has same answer options and coding as sffsatis

(N) **sffdst** “Feel distressed / worried by sex life”

(N) **sffavoid** “Avoided sex because of sexual difficulties – theirs or partners”

SEXUAL FUNCTION

1: agree strongly
2: agree
3: neither agree or disagree
4: disagree
5: disagree strongly
9: not answered

-1: not applicable *if* scelign3=2 | ((hetlife=0 | hetlife=9999) & (samlife=0 | samlife=9999)) | (dage<18 & (lastvag=-1 | lastvag=7) & (analsex=-1 | analsex=7) & (samahim=-1 | samahim=7) & (samayou=-1 | samayou=7))

(ND) **sffsatis2** dissatisfied with sex life

-1: not applicable *if* sffsatis=-1
0: no *if* sffsatis=1 | sffsatis=2 | sffsatis=3
1: yes *if* sffsatis=4 | sffsatis=5
9: not answered *if* sffsatis=9

(ND) **sffdst2** distressed or worried about sex life

-1: not applicable *if* sffdst=-1
0: no *if* sffdst=3 | sffdst=4 | sffdst=5
1: yes *if* sffdst=1 | sffdst=2
9: not answered *if* sffdst=9

(ND) **sffavoid2** avoided sex because of own or partner's sexual difficulties

-1: not applicable *if* sffavoid=-1
0: no *if* sffavoid=3 | sffavoid=4 | sffavoid=5
1: yes *if* sffavoid=1 | sffavoid=2
9: not answered *if* sffavoid=9

Notes: binary variables availackint..avoionone were created directly from the multi-code question sfavprb. All vars are coded similarly to availackint.

(N) **availackint** lack of interest in sex causing avoidance of sex.

1: yes
0: no
9: not answered
-1: not applicable *if* sffavoid=-1 | sffavoid=3 | sffavoid=4 | sffavoid=5

(N) **availackenj** lack of enjoyment in sex causing avoidance of sex

(N) **avoianxsex** felt anxious during sex causing avoidance of sex

(N) **avoipainsex** feared/felt physical pain during sex causing avoidance of sex

(N) **avoiononarou** no excitement/arousal causing avoidance of sex

(N) **avoiononlyorg** no orgasm or took a long time to reach orgasm causing avoidance of sex

(N) **avoipremorg** reached orgasm more quickly than would have liked causing avoidance of sex

Note: answer option availubric only available to female respondents

(N) **availubric** uncomfortable dry vagina causing avoidance of sex

1: yes
0: no
9: not answered
-1: not applicable *if* sffavoid=-1 | sffavoid=3 | sffavoid=4 | sffavoid=5 | rsex=1

Note: answer option avoierect only available to male respondents

(N) **avoierect** trouble achieving / maintaining erections causing avoidance of sex

1: yes
0: no
9: not answered
-1: not applicable *if* sffavoid=-1 | sffavoid=3 | sffavoid=4 | sffavoid=5 | rsex=2

SEXUAL FUNCTION

- (N) **avoipartprb** partner had sexual difficulties causing avoidance of sex
(N) **avoione** none of these problems caused avoidance of sex

Notes: binary variables spfamfr..spnohelp were created directly from the multi-code question sfhwch. All vars are coded similarly to spfamfr.

- (N) **spfamfr** contacted family member/friend about sexual advice
1: yes
0: no
9: not answered
-1: not applicable *if* scelign3=2 | | ((hetlife=0 | hetlife=9999) & (samlife=0 | samlife=9999)) | (dage<18 & (lastvag=-1 | lastvag=7) & (analsex=-1 | analsex=7) & (samahim=-1 | samahim=7) & (samayou=-1 | samayou=7))

- (N) **spinternet** got information and support from the internet about sexual advice
(N) **spbookslea** got self-help books / information leaflets about sexual advice
(N) **spselfh** contacted self-help groups about sexual advice
(N) **sphelpi** contacted helpline about sexual advice
(N) **spgp** contacted GP / family doctor about sexual advice
(N) **spgumsti** contacted sexual health/GUM/STI clinic about sexual advice
(N) **sppsych** contacted psychiatrists/psychologist about sexual advice
(N) **sprelc** contacted relationship counsellor about sexual advice
(N) **spothrgp** contacted other type of clinic or doctor about sexual advice
(N) **spnohelp** have not sought any help about sexual advice

- (ND) **sphelp** sought help or advice for sex life in the past year
-1: not applicable *if* spnohelp=-1
0: no *if* spnohelp=1
1: yes *if* spnohelp=0
9: not answered *if* spnohelp=9

Note: frequent was moved here in Natsal-3. It was at the end of the attitudes section in Natsal-2. Now only asked of those reporting at least one partner in the last year (het1yr and sam1yr)

- (MU) (N) **frequent** preference for frequency of sex now
1: to have sex much more often
2: to have sex a bit more often
3: it is about right as it is
4: to have sex less often
5: to have sex much less often
9: not answered/don't know
(U) -1: not applicable *if* scelign3=2 | ((het1yr=0 | het1yr=999) & (sam1yr=0 | sam1yr=999))

Note: easyyou was included in Natsal-2 at the end of the attitudes section. This was split into two questions for Natsal-3 depending on the number of lifetime partners the respondent reported

- (N) **easyyoua** how easy to talk about sex, 1 lifetime partner
1: always easy with my partner
2: always difficult with my partner
3: depends. Sometimes easy, sometimes difficult
9: not answered
-1: not applicable *if* scelign3=2 | ((hetlife=0 & samlife=0) | hetlife>1 | samlife>1 | (hetlife=1 & samlife=1))

- (N) **easyoub** how easy to talk about sex, 2+ lifetime partners
1: easy with spouse/reg. partner but difficult with a new partner
2: easy with a new partner but difficult with a spouse/reg. partner
3: easy with any partner
4: difficult with any partner
5: depends. Sometimes easy, sometimes difficult

SEXUAL FUNCTION

9: not answered

-1: not applicable *if* scelign3=2 | (hetlife=0 | (hetlife=9999) & (samlife=0 | samlife=9999)) | (hetlife=1 & (samlife=0 | samlife=-1 | samlife=9999)) | (samlife=1 & (hetlife=0 | hetlife=-1 | hetlife=9999))

(ND) **easyou** always find it easy to talk about sex with partners

-1: not applicable *if* easyoua=-1 | easyoub=-1

0: no/other *if* easyoua=2 | easyoua=3 | easyoub=1 | easyoub=2 | easyoub=4 | easyoub=5

1: yes *if* other *if* easyoua=1 | easyoub=3

9: not answered *if* easyoua=9 | easyoub=9

VIAGRA

Note: question medper was asked of respondents in the CASI who were aged >17, or respondents in the CASI aged 16-17 who reported vaginal or (het/sam) anal sex ever.

(N) **medper** taken drugs to assist sexual performance

1: yes

2: no

9: not answered

-1: not applicable *if* scelign3=2 | ((hetlife=0 & samlife=0) | (dage<18 & (lastvag=7 | lastvag=9) & (analsex=7 | analsex=9) & (samahim=-1 | samahim=7 | samahim=9) & (samayou=-1 | samayou=7 | samayou=9)))

(N) **medpwhn** last time took drugs to assist sexual performance

0: did not take sexual performance medications *if* medper=2

1: last 7 days

2: 7 days – 4 weeks

3: 4 weeks – 1 year

4: more than 1 year

9: not answered

-1: not applicable *if* merper=-1

DRUG USE

DRUGS

Note: Binary variables drcannabis..drnone were created directly from the question druguse. All vars are coded similarly to drcannabis. These questions refer to non-injected drugs. Drugsever and drugsyr are derived variables for use of any drugs (injected plus non-injected) – see page 160.

(N) **drcannabis** drugs taken ever (non injected): cannabis

1: yes

0: no

9: not answered

-1: not applicable *if* scelign3=2

(N) **drampheta** drugs taken ever (non injected): amphetamines

(N) **drcocaine** drugs taken ever (non injected): cocaine

(N) **drcrack** drugs taken ever (non injected): crack

(N) **drecstasy** drugs taken ever (non injected): ecstasy

(N) **drnoniher** drugs taken ever (non injected): non injected heroine

(N) **dracidlsd** drugs taken ever (non injected): acid / lsd

(N) **drcrysmeth** drugs taken ever (non injected): crystal meth

(N) **dramylnit** drugs taken ever (non injected): amyl nitrates

(N) **drothnonpre** drugs taken ever (non injected): other non prescribed drugs

(N) **drnone** drugs taken ever (non injected): none of these

(N) **drcan12m** cannabis, last 12 months

1: yes

2: no *or if* drcannabis=0

9: not answered

-1: not applicable *if* scelign3=2

(N) **drcan4w** cannabis, last 4 weeks

1: yes

2: no *or if* drcannabis=0

9: not answered

-1: not applicable *if* scelign3=2

(N) **drg12m** other drugs, last 12 months

1: yes

2: no *or if* drampheta=0 & drcocaine=0 & drcrack=0 & drecstasy=0 & drnoniher=0 & dracidlsd=0 & drcrysmeth=0 & dramylnit=0 & drothnonpre=0

9: not answered

-1: not applicable *if* scelign3=2

(N) **drg4w** other drugs, last 4 weeks

1: yes

2: no *or if* drampheta=0 & drcocaine=0 & drcrack=0 & drecstasy=0 & drnoniher=0 & dracidlsd=0 & drcrysmeth=0 & dramylnit=0 & drothnonpre=0

9: not answered

-1: not applicable *if* scelign3=2

(N) **inject2** ever injected non-prescribed drugs / other substances

1: yes

2: no

9: not answered

-1: not applicable *if* scelign3=2

DRUG USE

Notes: 1. The question wording differs slightly to that used in Natsal-1, which asked 'When was the last time you injected yourself with drugs or other substances?' since this question was asked to respondents who reported injecting all or some prescribed drugs rather than just those who reported injecting non-prescribed drugs. It is therefore vital to exclude respondents in Natsal-1 who reported injecting only prescribed drugs when making comparisons with Natsal-2's *wheninj* variable.

2. Unfortunately this inconsistency in question wording was not realised until after the publication of the paper by Johnson et al in the Lancet on 1 Dec 2001. Published estimates of the change in prevalence between Natsal-1 and Natsal-2 were therefore under-estimated.

3. The HPA requested more precise estimates of the injecting period to create a period of risk. Answer options for *wheninj* were therefore reduced in Natsal-3 and additional questions are asked about age at first and last injection.

(MU – see below) (s17c) **wheninj** last time injected non-prescribed drugs/other substances

- 1: in the last 7 days
- 2: between 7 days and 4 weeks
- 3: between 4 weeks and 1 year
- 4: between 1 and 5 years
- 5: longer than 5 years
- 9: not answered
- 1: not applicable if *scelig*=2 | *inject*=2 | *prescrb*=1

(s17c) **wheninj** last time injected non-prescribed drugs/other substances – NOTE question identical but answer options have changed

- 1: in the last 7 days
- 2: between 7 days and 4 weeks
- 3: between 4 weeks and 1 year
- 4: longer than 1 year
- 9: not answered
- 1: not applicable if *scelign3*=2 | *inject2*=2

(R) (ND) **inject5y** injected non-prescribed drugs/other substances, last 5 years

- 0: no if *drugs*=2 | *wheninj*=5
- 1: yes if *wheninj*=1 | *wheninj*=2 | *wheninj*=3 | *wheninj*=4
- 9: not answered if *wheninj*=9
- 1: not applicable if *drugs*=-1

(U) (ND) **inject1y** injected non-prescribed drugs/other substances, last year

- (U) 0: no if *inject2*=2 | *whninj*=4 0: no if *drugs*=2 | *wheninj*=4 | *wheninj*=5
- 1: yes if *wheninj*=1 | *wheninj*=2 | *wheninj*=3
- 9: not answered if *wheninj*=9
- (U) -1: not applicable if *inject2*=-1 -1: not applicable if *drugs*=-1

(N) **injaglas** age when last injected if >1 year ago

- 99: not answered
- 1: not applicable if *scelign3*=2 | *inject2*=2 | *wheninj*=1 | *wheninj*=2 | *wheninj*=3

(N) **injaglasall** age when last injected (all)

- compute *injaglasall*=*dage* if *wheninj*=1 | *wheninj*=2 | *wheninj*=3
- compute *injaglasall*=*injaglas* if *wheninj*=4
- 99: not answered
- 1: not applicable if *scelign3*=2 | *inject2*=2

(N) **injagfst** age when first injected

- 99: not answered
- 1: not applicable if *scelign3*=2 | *inject2*=2

DRUG USE

Note: In Natsal-1, all respondents who reported to have ever injected drugs, regardless of whether or not they were prescribed drugs, were asked whether or not they had ever shared a needle etc. In comparison, in Natsal-2 only those who reported injecting non-prescribed drugs were asked about sharing needles. It is therefore important in Natsal-1 to exclude respondents who reported injecting only prescribed drugs when making comparisons with Natsal-2's 'needle' variable.

(s17d) **needle** ever shared a needle or other equipment used for injecting

1: yes

2: no

9: not answered

(U) -1: not applicable if scalign3=2 | inject2=2 -1: not applicable if scelig=2 | inject=2 | prescib=1

(N) **drugsever** ever taken drugs, including injected drugs

1: yes if drnone=0 | inject2=1

0: no if drnone=1 & inject2=2

9: not answered if drnone=9

-1: not applicable if drnone=-1

(N) **drugsyr** taken drugs in last year, including injected drugs

1: yes if drcan12m=1 | drg12m=1 | inject1y=1

0: no if drcan12m=2 & drg12m=2 & inject1y=0

9: not answered if drcan12m=9

-1: not applicable if drcan12m=-1

(ND) **drugsyr2** drug use in the past year

-1: not applicable if drugsyr=-1

0: no if drugsyr=0

1: yes, cannabis only if drugsyr=1 & drg12m≠1 & inject1y≠1

2: yes, drugs other than cannabis if drugsyr=1 & (drg12m=1 | inject1y=1)

9: not answered if drugsyr=9

MOOD AND WELLBEING

MOOD AND WELL-BEING

Notes: 1. Mood1 and Mood2 form a composite measure, called PHQ2, used to screen for depressive symptoms in the last two weeks, and should not normally be used as independent variables. Many studies have described and validated the use of PHQ2¹.

2. Mood1 and Mood2 are scored 0-3 for answers 'not at all' to 'nearly every day' to give a total score (mscore) for the measure of 0-6. Although PHQ2 is derived from PHQ9, it was not designed to measure severity of depression and the variable mscore should not be used as such. Most studies use a threshold of 3 or more to define 'current depression', which strikes an appropriate balance between sensitivity and specificity. The derived variable depscr uses this threshold and is the variable recommended for analysis.

(N) **mood1** how frequent bothered by "little interest / pleasure in doing things", last 2 weeks

- 1: not at all
- 2: several days
- 3: more than half days
- 4: nearly every day
- 9: not answered
- 1: not applicable if scalign3=2

(N) **mood2** how frequent bothered by "feeling down, depressed or hopeless", last 2 weeks

- 1: not at all
- 2: several days
- 3: more than half days
- 4: nearly every day
- 9: not answered
- 1: not applicable if scalign3=2

Note: mood1 and mood2 are scored 0-3 for answers not at all – nearly every day

(ND) **mscore** Total score on PHQ-2 (i.e. score on mood1 + score on mood2)

- 9: not answered if mood1=9 | mood2=9
- 1: not applicable if mood1=-1

(ND) **depscr** Screen positive for current depression

- 0: no if mscore>=0 & mscore<=2
- 1: yes if mscore>=3 & mscore<=6
- 9: not answered if mood1=9 | mood2=9
- 1: not applicable if mood1=-1

¹ Examples: Arroll B, Khin N, Kerse N. Screening for depression in primary care with two verbally asked questions: cross sectional study. BMJ. 2003 Nov 15;327(7424):1144-6. PubMed PMID: 14615341; PubMed Central PMCID: PMC261815.

Kroenke K, Spitzer RL, Williams JB. The Patient Health Questionnaire-2: validity of a two-item depression screener. Med Care. 2003 Nov;41(11):1284-92. PubMed PMID: 14583691.

CIRCUMCISION

CIRCUMCISION

Note: Questions on circumcision were only asked to men

Note: Variable circum2 only applies to men eligible for the self-completion questionnaire.

Note: circum1 and circum2 are not included in the Natsal-3 dataset, as the derived variable 'circum' should be used for analysis.

(RU) (N) **circum2** circumcised

1: yes

2: no

(U) 3: don't know

9: not answered

(U) -1: not applicable if rsex=2 | scelig=2 -1: not applicable if rsex=2 | scelig=2

Note: Variable circum1 only applies to men not eligible for the self-completion questionnaire or men who only partially completed the self-completion questionnaire

(RU) (N) **circum1** circumcised

1: yes

2: no

3: question not understood

9: not answered

(U) -1: not applicable if rsex=2 | circum2=1 | circum2=2 | circum2=3 | circum2=9 | resultsc=2 | resultsc=3 | resultsc=4 | scelig=2

-1: not applicable if rsex=2 | circum2≠9 | (resultsc=2 | resultsc=3) | scelig=2

(s21m D) **circum** circumcised

1: yes if circum1=1 | circum2=1

2: no if circum1=2 | circum2=2

9: question not understood/not answered if circum1=3 | circum1=9 | circum2=3 | circum2=9

-1: not applicable if rsex=2

ATTITUDES

The following topics were all asked face-to-face by the interviewer.

ATTITUDES

Note: the attitudes section was extensively revised in Natsal-3 since a lot of the previous questions were felt to be outdated. New questions were drafted to attempt to capture more relevant concerns in current society.

(R) (q37a) **impfaith** importance for a successful marriage/long-term relationship: faithfulness
1: very important
2: quite important
3: not very important
4: not at all important
8: don't know *or if code 98 (the usual code for don't know)*
9: not answered

*Note: Variables **impinc** ... **impaste** and have the same codes as **impfaith**:*

(R) (q37b) **impinc** importance for a successful marriage/long-term relationship: an adequate income
(R) (q37c) **impresp** importance for a successful marriage/long-term relationship: mutual respect & appreciation
(R) (q37d) **imprelig** importance for a successful marriage/long-term relationship: shared religious beliefs
(R) (q37e) **impsexre** importance for a successful marriage/long-term relationship: a happy sexual relationship
(R) (q37f) **impshare** importance for a successful marriage/long-term relationship: sharing household chores
(R) (q37g) **impchild** importance for a successful marriage/long-term relationship: having children
(R) (q37h) **impaste** importance for a successful marriage/long-term relationship: tastes & interests in common

*Note: Variables **rwpremar** ... **rwabort** have code 8 'depend/don't know', which is an addition since Natsal-1*

(R) (q39a) **rwpremar** opinion of sex before marriage: right/wrong
1: always wrong
2: mostly wrong
3: sometimes wrong
4: rarely wrong
5: not wrong at all
8: depends/don't know *or if code 98 (the usual code for don't know)*
9: not answered

*Note: Variables **rwpremar** ... **rwabort** have the same codes as **rwpremar**:*

(q39b) **rwadult** opinion of adultery when married: right/wrong
(R) (q39c) **rwcohab** opinion of adultery when cohabiting: right/wrong
(R) (q39d) **rwregp** opinion of unfaithfulness: right/wrong
(q39e) **rwcasual** opinion of one night stands: right/wrong
rwsamm (q39f) **rwhomm** opinion of sexual relations between two adult men: right/wrong
rwsamf (q39g) **rwhomf** opinion of sexual between two adult women: right/wrong
(R) (q39h) **rwabort** opinion of abortion: right/wrong

(R) (q40c) **agaffec** company/affection more important than sex in a marriage/relationship: agree/disagree
1: agree strongly
2: agree
3: neither agree nor disagree
4: disagree
5: disagree strongly
8: don't know *or if code 98 (the usual code for don't know)*
9: not answered

*Note: Variables **agaffec** ... **agseximp** have the same codes as **agaffec**:*

ATTITUDES

- (R) (q40d) **agorgasm** sex, no orgasm not really satisfying for men: agree/disagree
(R) (q40e) **agorgasf** sex, no orgasm not really satisfying for women: agree/disagree
(R) (q40g) **agseximp** sex most important part of marriage: agree/disagree

(N) **snnolov** it's ok to have sex with someone without being in love with them

- 1: agree strongly
- 2: agree
- 3: neither agree nor disagree
- 4: disagree
- 5: disagree strongly
- 8: don't know
- 9: not answered

Note: snpres – snsexed have same response options as snnolov

(N) **snpres** people are under a lot of pressure to have sex nowadays

(N) **snold** it's natural for people to want less sex as they get older

(N) **snsexdrv** men have a naturally higher sex drive than women

(N) **snmedia** there's too much sex in the media these days

(N) **snearly** young people start having sex too early

(N) **sngayadp** gay men should be able to adopt children

(N) **snlesadp** lesbians should be able to adopt children

(N) **snsexed** teaching young people about sexual matters encourages them to have sex

(N) **ch1625** respondent has children now aged 16-25 and had contact with them when growing up

- 1: yes
- 2: no
- 9: not answered
- 1: not applicable if anychm≠1 & stepchil≠1 & adopchil≠1 & nochall≠0

(N) **talkch** ease of talking to children about sexual matters when growing up

- 1: easy
- 2: difficult
- 3: easy with some, difficult with others
- 4: depends on topic
- 5: didn't discuss
- 9: not answered
- 1: not applicable if ch1625=-1 | ch1625=2

(N) **uagesex** respondent's perception of proportion of young people having sex before 16

- 1: less than $\frac{1}{4}$
- 2: about $\frac{1}{4}$
- 3: about $\frac{1}{2}$
- 4: about $\frac{3}{4}$
- 5: more than $\frac{3}{4}$
- 8: don't know
- 9: not answered

ENJOYMENT OF SEX

Note: Frequent was here in Natsal-2 but was moved to the CASI sexual function section in Natsal-3

(R) (N) **satisfy** sexual satisfaction now

- 1: always enjoy it
- 2: enjoy it most of the time
- 3: do not often enjoy it
- 4: never enjoy it
- 5: don't have sex these days
- 9: not answered/don't know

ATTITUDES

Note: Easyyou was split into two questions in Natsal-3 which now appear in the sexual function section of the CASI.

(R) (q41b*) **easyyou** how easy to talk about sex

- 1: easy with spouse/reg. partner but difficult with a new partner
- 2: easy with a new partner but difficult with a spouse/reg. partner
- 3: easy with any partner
- 4: difficult with any partner
- 5: depends/can't say/don't know
- 9: not answered

LIFESTYLE

(R) (q42a U) **idealnow** ideal sexual lifestyle now

- 1: prefer to have no sex activity
- 2: no regular partners but casual partners when I feel like it
- 3: a few regular partners
- 4: one regular partner but not living together
- 5: not married, but living with a partner and with some sex activity outside the partnership
- 6: not married, but living with a partner, and no other sex partners
- 7: married, with some sex activity outside the marriage
- 8: married, with no other sex partners
- 9: have no ideal/none of these/don't know
- 99: not answered

Note: Variable 'ideal5yr' has the same codes as idealnow:

(R) (N) **ideal5yr** ideal sexual lifestyle in 5 years time

SAFER SEX

Note: In Natsal-3 a lot of the safer sex relating to HIV questions were removed since they would not have been relevant to the younger respondents.

(R) (q44a*) **anychang** changed sexual lifestyle because of concerns about HIV/STI risk

- 1: yes
- 2: no
- 3: lifestyle changed but not because of AIDS
- 9: not answered

(R) (q44b) **howchan1** how changed lifestyle because of concern about HIV/STI risk 1

- 1: having fewer partners
- 2: finding out more about the person before having sex
- 3: using a condom
- 4: not having sex
- 5: sticking to one partner
- 6: avoiding some sexual practices
- 7: other change(s)
- 9: not answered
- 1: not applicable if anychang=2 | anychang=3 | howchan1 ≠ 9

Similarly for at most 7 changes:

(R) **howchan2** how changed lifestyle because of concern about HIV/STI risk 2

(R) **howchan3** how changed lifestyle because of concern about HIV/STI risk 3

(R) **howchan4** how changed lifestyle because of concern about HIV/STI risk 4

(R) **howchan5** how changed lifestyle because of concern about HIV/STI risk 5

(R) **howchan6** how changed lifestyle because of concern about HIV/STI risk 6

(R) **howchan7** how changed lifestyle because of concern about HIV/STI risk 7

ATTITUDES

(R) (q44bd D) **rrsfewrp** risk reduction strategy: having fewer partners

1: yes *if* howchan1=1 | howchan2=1 | howchan3=1 | howchan4=1 | howchan5=1 | howchan6=1 | howchan7=1

0: no *if* (anychang=1 & howchan1>=1 & howchan1<=7) | anychang=2 | anychang=3

9: not answered *if* missing

Note: Variables rrsknosh ... rrsother are coded in a similar way to rrsfewrp:

(R) (q44bl D) **rrsknosh** risk reduction strategy: finding out more about partner before sex

(R) (q44bk D) **rrscondm** risk reduction strategy: using condoms

(R) (q44bc D) **rrsabstn** risk reduction strategy: not having sex

(R) (q44bx D) **rrsmonog** risk reduction strategy: sticking to one partner

(R) (q44bq D) **rrsavoid** risk reduction strategy: avoiding some sexual practices

(R) (q44bn D) **rrsother** risk reduction strategy: other

RISK PERCEPTION

(q45a) **rskyou** HIV/AIDS risk: to self

1: greatly at risk

2: quite a lot

3: not very much

4: not at all at risk

8: don't know

9: not answered *if* missing

(N) (q45a) **stirisk** other STI risk: to self

1: greatly at risk

2: quite a lot

3: not very much

4: not at all at risk

8: don't know

9: not answered *if* missing

(R) q45b **rskprom** HIV/AIDS risk: people who have many heterosexual partners

1: greatly at risk

2: quite a lot at risk

3: not very much at risk

4: not at all at risk

5: depends on whether they use condoms/practise safe sex (*this is documented as code 9 in the Technical Report*)

8: don't know

9: not answered *if* missing

Note: Variables rskmarr ... rskhomf have the same codes as rskprom:

(R) (q45c) **rskmarr** HIV/AIDS risk: married couples who only have sex with each other

(R) (q45d) **rskmocc** HIV/AIDS risk: married couples who occasionally have sex with other partner(s)

(R) (q45e) **rskhomm** HIV/AIDS risk: male homosexuals

(R) (q45f) **rskhomf** HIV/AIDS risk: female homosexuals

(R) (N) **rskdrugs** HIV/AIDS risk: those who inject illegal drugs

1: greatly at risk

2: quite a lot at risk

3: not very much at risk

4: not at all at risk

5: depends (e.g. on whether they share needles) (*this is documented as code 9 in the technical report*)

8: don't know

9: not answered *if* missing

ATTITUDES

(R) (N) **cfgbusa** perception of HIVrisk in USA relative to Britain

- 1: much more likely
- 2: somewhat more likely
- 3: about the same as Britain
- 4: somewhat less likely
- 5: much less likely
- 8: can't say
- 99: not answered

*Note: Variables **cfgbswi** ... **cfgbthai** have the same codes as **cfgbusa**:*

(R) (N) **cfgbswi** perception of HIVrisk in Switzerland relative to Britain

(R) (N) **cfgbken** perception of HIVrisk in Kenya relative to Britain

(R) (N) **cfgbneth** perception of HIVrisk in the Netherlands relative to Britain

(R) (N) **cfgbthai** perception of HIVrisk in Thailand relative to Britain

HOUSEHOLD CLASSIFICATION

HOUSEHOLD CLASSIFICATION

Note: hhsz-reIToR were later on in the questionnaire in Natsal-2

(M) (c4a) **hhsz** no. of people who live regularly in household (incl. respondent)
(U – code removed, 9995 code used instead) 99: not answered
9995: at least one if hhsz is not answered

Note: household member 1 refers to the respondent

(M) (N) **gender** gender of household member 1

1: male

2: female

9: not answered

-1: not applicable if hhsz < gendern

Similarly for up to 14 (previously 12) household members:

(M) (N) **gender2** gender of household member 2

(M) (N) **gender3** gender of household member 3

(M) (N) **gender4** gender of household member 4

(M) (N) **gender5** gender of household member 5

(M) (N) **gender6** gender of household member 6

(M) (N) **gender7** gender of household member 7

(M) (N) **gender8** gender of household member 8

(M) (N) **gender9** gender of household member 9

(M) (N) **gender10** gender of household member 10

(M) (N) **gender11** gender of household member 11

(M) (N) **gender12** gender of household member 12

Note: Household member 1 refers to the respondent

(M) (N) **ageof** age of household member 1

999: not answered

-1: not applicable if hhsz < ageofn

Similarly for up to 14 (previously 12) household members:

(M) (N) **ageof2** age of household member 2

(M) (N) **ageof3** age of household member 3

(M) (N) **ageof4** age of household member 4

(M) (N) **ageof5** age of household member 5

(M) (N) **ageof6** age of household member 6

(M) (N) **ageof7** age of household member 7

(M) (N) **ageof8** age of household member 8

(M) (N) **ageof9** age of household member 9

(M) (N) **ageof10** age of household member 10

(M) (N) **ageof11** age of household member 11

(M) (N) **ageof12** age of household member 12

Note: There is no 'reitor' as this would simply be the respondent themselves.

(U – see below) (M) (N) **reitor2** relationship with household member 2

1: spouse/partner

2: son/daughter (incl. step/adopted/in-law)

3: parent (incl. step/in-law)

4: brother/sister (incl. half/in-law)

5: grandparent

6: grandchild

7: other relative

8: non-relative

99: not answered

-1: not applicable if hhsz < gendern

HOUSEHOLD CLASSIFICATION

(U) **reltor2** relationship with household member 2

- 1: spouse / civil partner
- 2: cohabiting partner
- 3: son/daughter
- 4: step-son / -daughter
- 5: foster child
- 6: son-in-law / daughter-in-law
- 7: parent / guardian
- 8: step-parent
- 9: foster parent
- 10: parent-in-law
- 11: brother / sister (incl adopted)
- 12: half brother / sister
- 13: foster brother / sister
- 14: brother- / sister-in-law
- 15: grandchild
- 16: grandparent
- 17: other relative
- 18: other non-relative

Similarly for up to 14 (previously 12) household members:

- (U) (M) (N) **reltor3** relationship with household member 3
- (U) (M) (N) **reltor4** relationship with household member 4
- (U) (M) (N) **reltor5** relationship with household member 5
- (U) (M) (N) **reltor6** relationship with household member 6
- (U) (M) (N) **reltor7** relationship with household member 7
- (U) (M) (N) **reltor8** relationship with household member 8
- (U) (M) (N) **reltor9** relationship with household member 9
- (U) (M) (N) **reltor10** relationship with household member 10
- (U) (M) (N) **reltor11** relationship with household member 11
- (U) (M) (N) **reltor12** relationship with household member 12

Note: Spoushh..cosamhh are variables generated automatically by the CAPI based on answers to reltor and gender questions. Civpthh, coopphh, and cosamhh are coded the same as spoushh.

(N) **spoushh** spouse in the household

- 1: yes
- 2: no

(N) **civpthh** civil partner in the household

- (N) **coopphh** cohabiting opposite-sex partner in the household
- (N) **cosamhh** cohabiting same-sex partner in the household

Note: partnerhh..nonrelhh (who respondent lives with) are not mutually exclusive. These variables are combined in the final variable rhold, which sets out different types of household structure.

(ND) **partnerhh** respondent lives with spouse/civil partner/cohabiting partner

- 1: yes if spoushh=1 | civpthh=1 | coopphh=1 | cosamhh=1
- 0: no if spoushh=0 & civpthh=0 & coopphh=0 & cosamhh=0

Note: hhpartage..nonrelhh are derived using repeat commands for all loops of reltor, thus !=1...12 (command repeated for each loop).

HOUSEHOLD CLASSIFICATION

(ND) **hhpartage** spouse/civil partner/cohabiting partner's age
compute hhpartage=ageof! if reltor!=1 | reltor!=2
-1: not applicable *if* spoushh=2 & civpthh=2 & coopphh=2 & cosamhh=2
999: not answered *if* (reltor!=1 | reltor!=2) & ageof!=999

(ND) **child12** respondent lives with child aged<12y (incl. adopted/step/fostered)
1: yes *if* (reltor!>=3 & reltor!<=5 & ageof!<12)
9: not answered *else if* reltor!=99
0: no *else*

(ND) **child1217** respondent lives with child aged12-17y (incl. adopted/step/fostered)
1: yes *if* (reltor!>=3 & reltor!<=5 & ageof!>=12 & ageof!<=17)
9: not answered *else if* reltor!=99
0: no *else*

(ND) **childm17** respondent lives with child aged18+ (incl. adopted/step/fostered)
1: yes *if* (reltor!>=3 & reltor!<=5 & ageof!>=18 & ageof!<=999)
9: not answered *else if* reltor!=99
0: no *else*

(ND) **parenthh** respondent lives with parent/s (incl step/foster/parents-in-law)
1: yes *if* reltor!>=7 & reltor!<=10
9: not answered *else if* reltor!=99
0: no *else*

(ND) **othrelhh** respondent lives with other relatives (incl. siblings)
1: yes *if* reltor!>=11 & reltor!<=17
9: not answered *else if* reltor!=99
0: no *else*

(ND) **nonrelhh** respondent lives with other relatives
1: yes *if* reltor!=18
9: not answered *else if* reltor!=99
0: no *else*

(ND) **rhhold** respondent's household structure at time of interview.
1: lives alone *if* hhsz=1
2: lives with child/ren aged <12y *if* child12=1 & child1217=0 & childm17=0 & parenthh=0 & othrelhh=0 & nonrelhh=0
3: lives with child/ren aged >=12y *if* child12=0 & (child1217=1 | childm17=1) & parenthh=0 & othrelhh=0 & nonrelhh=0
4: lives with child/ren aged <12y and >=12y *if* child12=1 & (child1217=1 | childm17=1) & parenthh=0 & othrelhh=0 & nonrelhh=0
5: lives with partner *if* partnerhh=1 & child12=0 & child1217=0 & childm17=0 & parenthh=0 & othrelhh=0 & nonrelhh=0
6: lives with partner and child/ren aged<12y *if* partnerhh=1 & child12=1 & child1217=0 & childm17=0 & parenthh=0 & othrelhh=0 & nonrelhh=0
7: lives with partner and child/ren aged >=12y *if* partnerhh=1 & child12=0 & (child1217=1 | childm17=1) & parenthh=0 & othrelhh=0 & nonrelhh=0
8: lives with partner and children aged <12y and >=12y *if* partnerhh=1 & child12=1 & (child1217=1 | childm17=1) & parenthh=0 & othrelhh=0 & nonrelhh=0
9: lives with parents *if* partnerhh=0 & child12=0 & child1217=0 & childm17=0 & parenthh=1 & othrelhh=0 & nonrelhh=0
10: lives with parents and child/ren (any age) *if* partnerhh=0 & (child12=1 | child1217=1 | childm17=1) & parenthh=1 & othrelhh=0 & nonrelhh=0
11: lives with parents and partner *if* partnerhh=1 & child12=0 & child1217=0 & childm17=0 & parenthh=1 & othrelhh=0 & nonrelhh=0

HOUSEHOLD CLASSIFICATION

12: lives with parents and partner and child/ren (any age) *if* partnerhh=1 & (childl12=1 | child1217=1 | childm17=1) & parenthh=1 & othrelhh=0 & nonrelhh=0
 13: lives with parents and other relatives (incl. siblings) *if* partnerhh=0 & childl12=0 & child1217=0 & childm17=0 & parenthh=1 & othrelhh=1 & nonrelhh=0
 14: lives with parents and other relatives (incl. siblings) and child/ren (any age) *if* partnerhh=0 & (childl12=1 | child1217=1 | childm17=1) & parenthh=1 & othrelhh=1 & nonrelhh=0
 15: lives with parents and other relatives and partner *if* partnerhh=1 & childl12=0 & child1217=0 & childm17=0 & parenthh=1 & othrelhh=1 & nonrelhh=0
 16: lives with parents and other relatives and partner and child/ren (any age) *if* partnerhh=1 & (childl12=1 | child1217=1 | childm17=1) & parenthh=1 & othrelhh=1 & nonrelhh=0
 17: lives with other relatives (incl. siblings) *if* partnerhh=0 & childl12=0 & child1217=0 & childm17=0 & parenthh=0 & othrelhh=1 & nonrelhh=0
 18: lives with child/ren (any age) and other relatives (incl. siblings) *if* partnerhh=0 & (childl12=1 | child1217=1 | childm17=1) & parenthh=0 & othrelhh=1 & nonrelhh=0
 19: lives with partner and other relatives (incl. siblings) *if* partnerhh=1 & childl12=0 & child1217=0 & childm17=0 & parenthh=0 & othrelhh=1 & nonrelhh=0
 20: lives with partner and children (any age) and other relatives (incl. siblings) *if* partnerhh=1 & (childl12=1 | child1217=1 | childm17=1) & parenthh=0 & othrelhh=1 & nonrelhh=0
 21: lives with non-relatives *if* partnerhh=0 & childl12=0 & child1217=0 & childm17=0 & parenthh=0 & othrelhh=0 & nonrelhh=1
 22: lives with child/ren (any age) and non-relatives *if* partnerhh=0 & (childl12=1 | child1217=1 | childm17=1) & parenthh=0 & othrelhh=0 & nonrelhh=1
 23: lives with partner and non-relatives *if* partnerhh=1 & childl12=0 & child1217=0 & childm17=0 & parenthh=0 & othrelhh=0 & nonrelhh=1
 24: lives with partner and child/ren (any age) and non-relatives *if* partnerhh=1 & (childl12=1 | child1217=1 | childm17=1) & parenthh=0 & othrelhh=0 & nonrelhh=1
 25: lives with non-relatives and relatives (incl. parents) *if* (partnerhh=1 | childl12=1 | child1217=1 | childm17=1 | parenthh=1 | othrelhh=1) & nonrelhh=1
 26: not answered (data missing for 1+ household member) *if* (hhsz=9995 | partnerhh=9 | childl12=9 | child1217=9 | childm17=9 | parenthh=9 | othrelhh=9 | nonrelhh=9)

(U – see below) (c1a) **marstat** marital status at present

- 1: married & living with spouse
- 2: living with a partner of opposite sex as a couple
- 3: living with a partner of same sex as a couple
- 4: widowed
- 5: divorced
- 6: separated
- 7: single & never been married
- 9: not answered

(U) **marstat** marital status at present – question identical but answer options different. Response options 7, 8 and 9 were spontaneous only.

- 1: single & never married
- 2: married & living with spouse *or if* marstat not answered & spoushh=1
- 3: in registered same sex civil partnership & living with partner *or if* marstat not answered & civpthh=1
- 4: separated but still legally married
- 5: divorced
- 6: widowed
- 7: separated but still legally in same-sex civil partnership
- 8: formerly a same sex civil partnership but now legally dissolved
- 9: surviving civil partner, partner having died
- 99: not answered

(U – see below) (D) **ms** married status, grouped

- 1: married *if* marstat=1
- 2: cohabitation, opposite sex *if* marstat=2

HOUSEHOLD CLASSIFICATION

- 3: cohabitation, same sex *if* marstat=3
- 4: widowed *if* marstat=4
- 5: divorced/separated *if* marstat=5 | marstat=6
- 6: single and never been married *if* marstat=7
- 9: not answered

ms married status, grouped

- 1: married and living with spouse *if* marstat=2
- 2: in a registered same sex civil partnership and living with partner *if* marstat=3
- 3: cohabitation, opposite sex *if* copphh=1
- 4: cohabitation, same sex *if* cosamhh=1
- 5: Widowed (incl from same-sex civil partnership) *if* marstat=6 | marstat=9
- 6: divorced / separated, including from a same-sex civil partnership *if* marstat=4 | marstat=5 | marstat=7 | marstat=8
- 7: single and never married or in a same sex civil partnership *else if* marstat=1
- 9: not answered *if* marstat=99

(U – see below) (D) **ms2** married status, grouped

- 1: married & living with spouse *if* marstat=1
- 2: cohabitation (opposite sex & same sex) *if* marstat=2 | marstat=3
- 3: previously married (widowed, divorced & separated) *if* marstat=4 | marstat=5 | marstat=6
- 4: single and never been married *if* marstat=7
- 9: not answered

ms2 marital status, grouped

- 1: married / civil partnership & living with spouse / partner *if* ms=1 | ms=2
- 2: cohabitation (opposite or same-sex) *if* ms=3 | ms=4
- 3: previously married / in a civil partnership (widowed, divorced, separated) *if* ms=5 | ms=6
- 4: single and never married *if* ms=7
- 9: not answered *if* ms=9

(U – see below) (D) **ms3** married status, grouped

- 1: married & living with spouse *if* marstat=1
- 2: cohabitation, opposite sex *if* marstat=2
- 3: cohabitation, same sex *if* marstat=3
- 4: previously married (widowed, divorced & separated) *if* marstat=4 | marstat=5 | marstat=6
- 5: single and never been married *if* marstat=7
- 9: not answered

HOUSEHOLD CLASSIFICATION

ms3 married status, grouped

- 1: married and living with spouse *if* ms=1
- 2: in a registered same-sex civil partnership and living with partner *if* ms=2
- 3: cohabitation, opposite sex *if* ms=2
- 4: cohabitation, same sex *if* ms=4
- 5: previously married/same-sex civil partnership (widowed, divorced, separated) *if* ms=5 | ms=6
- 6: single and never married *if* ms=7
- 9: not answered *if* ms=9

(ND) **relstat** relationship status at time of interview

- 1: married and living with spouse *if* ms=1 | ms=2
- 2: in a registered same-sex civil partnership and living with partner *if* ms=3 | ms=4
- 3: living with partner *if* (r1rel=1 & (r1sexagn=1 | r1sexagn=2)) | (r1rel2=1 & (r1sexagn2=1 | r1sexagn2=2)) | (r1rel3=1 & (r1sexagn3=1 | r1sexagn3=2)) | (r1rel4=1 & (r1sexagn4=1 | r1sexagn4=2))
- 4: in a 'steady' and ongoing relationship but not living together *if* (r1rel=2 & (r1sexagn=1 | r1sexagn=2)) | (r1rel2=2 & (r1sexagn2=1 | r1sexagn2=2)) | (r1rel3=2 & (r1sexagn3=1 | r1sexagn3=2)) | (r1rel4=2 & (r1sexagn4=1 | r1sexagn4=2))
- 5: widowed and not in a 'steady' and ongoing relationship in the last year *else if* ms=5
- 6: divorced / separated and not in a 'steady' and ongoing relationship in the last year *else if* ms=6
- 7: previously lived with partner as a non-married couple but not now in a 'steady' and ongoing relationship in the last year *else if* ms=7 & pplive=1
- 8: never married or lived with partner, and not in a 'steady' and ongoing relationship in the last year *else if* ms=7 & pplive=2
- 9: not answered *if* (ms>=5 & ms<=9 & r1rel=9 & r1datey not in the last year)

(ND) **relstatgp** relationship status at time of interview, grouped

- 1: Living with partner (includes married / civil partnership) *if* relstat=1 | relstat=2 | relstat=3
- 2: In a steady relationship, not living with partner *if* relstat=4
- 3: Not in a steady relationship, previously cohabited (includes previously married / civil partner) *if* relstat=5 | relstat=6 | relstat=7
- 4: Not in a steady relationship, never cohabited *if* relstat=8
- 9: not answered *if* relstat=9

(ND) **relstatgp2** relationship status at interview

- 1: married or civil partnership *if* relstat=1 | relstat=2
- 2: living with a partner *if* relstat=3
- 3: in a 'steady' relationship but not living together *if* relstat=4
- 4: not in a 'steady relationship' *if* relstat=5 | relstat=6 | relstat=7 | relstat=8
- 9: not answered *if* relstat=9

LIVE-IN PARTNERSHIPS

PREVIOUS LIVE-IN PARTNERSHIPS

(R) (c1b) **page** age of live-in partner, last birthday

99: not answered

-1: not applicable if marstat=4 | marstat=5 | marstat=6 | marstat=7

(c2a*U) **pplive** previously lived with someone as a couple

1: yes

2: no

9: not answered

-1: not applicable if marstat≠7 & marstat≠9

(N) **ppnum** no. of times lived with someone as a couple

99: not answered

-1: not applicable if pplive=2

Note: Questions about previous partnerships were updated in Natsal-3 so that respondents were only asked questions about a maximum of 5 previous cohabitations (rather than 13 as in Natsal-2), as this was felt to be sufficient for analysis. In Natsal-2 the questions started with the first live-in partnership and worked forwards to the most recent. In Natsal-3 the questions start with the most recent live-in partnership and work backwards to the first. In Natsal-3 the fifth loop of questions (ppystrt5..pp5months) refers either to the fifth most recent cohabiting partnership if ppnum=5 or the first ever cohabiting partnership if ppnum>5.

Note: The Natsal-2 variables are shown below the Natsal-3 variables (Natsal-2 variables start page 180).

Natsal-3 PREVIOUS LIVE-IN PARTNERSHIP QUESTIONS

(N) **pp1ystrt** year started living with most recent live-in partner

9999: not answered

-1: not applicable if ppnum=-1

Note: in Natsal-3 pp1mstrt was only asked if this was in the last 5 years

(N) **pp1mstrt** month started living with most recent live-in partner

1: January

2: February

3: March

4: April

5: May

6: June

7: July

8: August

9: September

10: October

11: November

12: December

97: can't remember month

99: not answered

-1: not applicable if ppnum=-1 | (dateyoi – ppystrt > 5)

Note: pp1strt is a date-format variable based on pp1ystrt and pp1mstrt. Day is set to 01 for all cases, and month is set to June if missing. All missing values are set as system missing. The indicator variable pp1strt_ind can be used if more information is needed about the missing values.

(N) **pp1strt** date most recent live-in partnership started (missing estimated: day=1st, month=June)
sysmis if missing

LIVE-IN PARTNERSHIPS

(N) **pp1ms** most recent live-in partnership: type of union at start of cohabitation

- 1: married
- 2: civil partnership
- 3: living as a couple *or if* pp1ms=not answered and marstat=1
- 9: not answered
- 1: not applicable *if* ppnum=-1

(N) **pp1mar** later married / entered into civil partnership with most recent live-in partner

- 1: yes
- 2: no *or if* pp1mar=9 & marstat=7
- 9: not answered
- 1: not applicable *if* ppnum=-1 | pp1ms=1 | pp1ms=2

(N) **pp1ymar** year married / entered civil partnership most recent partner (if lived together before marriage)

- 9999: not answered
- 1: not applicable *if* ppnum=-1 | pp1ms=1 | pp1ms=2 | pp1mar=2

(N) **pp1mmar** month married / entered civil partnership most recent partner (if lived together before marriage)

- 1: January
- 2: February
- 3: March
- 4: April
- 5: May
- 6: June
- 7: July
- 8: August
- 9: September
- 10: October
- 11: November
- 12: December
- 97: can't remember
- 99: not answered
- 1: not applicable *if* ppnum=-1 | pp1ms=1 | pp1ms=2 | pp1mar=2

Note: Due to a CAPI filtering error pp1sex was not asked of those with marstat='single, never married'. In most cases this can be imputed, as shown below. Cases that could not be imputed are coded as 9 'not answered'.

(N) **pp1sex** sex of most recent live-in partner

- 1: heterosexual *or if* spoushh=1 | coopphh=1 | marstat=2 | (ppnum>=1 & ppnum<99 & eversam=2 & (marstat=4 | marstat=5 | marstat=6))
- 2: homosexual *or if* civpthh=1 | cosamhh=1 | marstat=3 | (ppnum>=1 & ppnum<99 & eversam=1 & (marstat=1 & hetlife=0) | marstat=7 | marstat=8 | marstat=9))
- 9: not answered (*includes those not able to impute*)
- 1: not applicable *if* ppnum=-1

Note: code 96 'still living with partner' only available for pp1ystop, not previous partnerships (pp2ystop etc).

(N) **pp1ystop** year stopped living with most recent live-in partner (if ever did)

- 96: still living with partner *or if* ppnum>=1 & ppnum<99 & (marstat=2 | marstat=3 | spoushh=1 | civpthh=1 | coopphh=1 | cosamhh=1)
- 9999: not answered
- 1: never lived with someone as a couple *if* pplive=2

LIVE-IN PARTNERSHIPS

(N) **pp1months** no. months lived with partner for if started and stopped cohab in same year
96: still living with partner
99: not answered
-1: not applicable if $ppnum \geq 1$ & $ppnum < 99$ & $pp1ystrt > 0$ & $pp1ystrt < 9999$ & $pp1ystop > 0$ & $pp1ystop < 9999$ & $pp1ystop \neq 96$ & $pp1ystrt \neq pp1ystop$

Notes: 1. Reasons cohabiting relationship ended was only asked for the most recent live-in partnership (that was not ongoing at the time of interview) i.e. loop 1 if there was no spouse / civil partner / cohabiting partner in the household (variables $pp1why1..pp1why25$) or loop 2 if there was a spouse / civil partner / cohabiting partner in the household and the respondent had at least one other previous cohabiting partnership (variables $pp2why1..pp2why24$).

2. Respondents who said that they were widowed ($marstat=6$) were not asked $ppwhy$ to avoid causing distress. This is indicated in the variable $pp1why25$. These respondents have not been coded as 'death of partner' ($pp1why2$) because we cannot be certain that the spouse who died was the most recent cohabiting partner.

3. A CAPI routing error meant that $ppwhy$ was only asked in loop 1 for waves 1-4 (first half) of fieldwork, therefore there is no information for wave 1-4 respondents who were married / in a civil partnership / cohabiting relationship at the time of interview about why their 2nd most recent cohabitation ended.

4. The binary variables $pp1why1..pp1why24$ were created directly from the question $ppwhy$. All variables coded in the same way as $pp1why1$.

(N) **pp1why1** Reason most recent cohabiting relationship ended: relationship has not ended1: mentioned
0: not mentioned
9: not answered
-1: not applicable if $ppnum=-1$

(N) **pp1why2** Reason most recent cohabiting relationship ended: death of partner
(N) **pp1why3** Reason most recent cohabiting relationship ended: difficulties with sex life
(N) **pp1why4** Reason most recent cohabiting relationship ended: other
(N) **pp1why5** Reason most recent cohabiting relationship ended: domestic violence
(N) **pp1why6** Reason most recent cohabiting relationship ended: different interests / nothing in common
(N) **pp1why7** Reason most recent cohabiting relationship ended: unfaithfulness / adultery
(N) **pp1why8** Reason most recent cohabiting relationship ended: arguments
(N) **pp1why9** Reason most recent cohabiting relationship ended: not having children
(N) **pp1why10** Reason most recent cohabiting relationship ended: grew apart
(N) **pp1why11** Reason most recent cohabiting relationship ended: moved because of change in circumstances
(N) **pp1why12** Reason most recent cohabiting relationship ended: lack of respect / appreciation
(N) **pp1why13** Reason most recent cohabiting relationship ended: not sharing housework enough
(N) **pp1why14** Reason most recent cohabiting relationship ended: money problems
(N) **pp1why15** Reason most recent cohabiting relationship ended: drink / drugs / gambling problems
(N) **pp1why16** Reason most recent cohabiting relationship ended: mental health or related problems
(N) **pp1why17** Reason most recent cohabiting relationship ended: problem with children
(N) **pp1why18** Reason most recent cohabiting relationship ended: never at home
(N) **pp1why19** Reason most recent cohabiting relationship ended: problems with parents / in-laws / family
(N) **pp1why20** Reason most recent cohabiting relationship ended: age problem
(N) **pp1why21** Reason most recent cohabiting relationship ended: another relationship involved
(N) **pp1why22** Reason most recent cohabiting relationship ended: lived in / moved to a different country
(N) **pp1why23** Reason most recent cohabiting relationship ended: changed mind / feelings / personality
(N) **pp1why24** Reason most recent cohabiting relationship ended: partner left without explanation

(N) **pp1why25** Reason most recent cohabiting relationship ended: respondent is a widow
1: mentioned if $marstat=6$
0: not mentioned if $ppnum > 0$ & $ppnum < 99$ & $marstat \neq 6$
9: not answered
-1: not applicable if $ppnum=-1$

LIVE-IN PARTNERSHIPS

Note: Variables pp2ystrt2 ... ppwhyx5 refer to the 2nd most recent live-in partner:

(N) **pp2ystrt** year starting living with 2nd most recent live-in partner

9999: not answered

-1: not applicable if ppnum<2

Note: in Natsal-3 pp2mstrt was only asked if this was in the last 5 years

(N) **pp2strtm** month started living with 2nd most recent live-in partner

1: January

2: February

3: March

4: April

5: May

6: June

7: July

8: August

9: September

10: October

11: November

12: December

97: can't remember

99: not answered

-1: not applicable if ppnum<2 | (dateyoi – pp2ystrt > 5)

Note: pp2strt is a date-format variable based on pp2ystrt and pp2mstrt. Day is set to 01 for all cases, and month is set to June if missing. All missing values are set as system missing. The indicator variable pp2strt_ind can be used if more information is needed about the missing values.

(N) **pp2strt** date second most recent live-in partnership started (missing estimated: day=1st, month=June)
sysmis if missing

pp2ms 2nd most recent live-in partnership: type of union at start of cohabitation

1: married

2: civil partnership

3: living as a couple

9: not answered

-1: not applicable if ppnum<2

(N) **pp2mar** later married / entered into civil partnership with 2nd most recent live-in partner

1: yes

2: no

9: not answered

-1: no applicable if (ppnum=-1 | ppnum<2) | pp2ms=1

(N) **pp2ymar** year married / entered civil partnership 2nd most recent partner (if lived together before marriage)

9999: not answered

-1: not applicable if ppnum<2 | pp2ms=1 | pp2ms=2 | pp2mar=2

LIVE-IN PARTNERSHIPS

(N) **pp2mmar** month married / entered civil partnership 2nd most recent partner (if lived together before marriage)

1: January

2: February

3: March

4: April

5: May

6: June

7: July

8: August

9: September

10: October

11: November

12: December

97: can't remember

99: not answered

-1: not applicable if ppnum<2 | pp2ms=1 | pp2ms=2 | pp2mar=2

Note: As with pp1sex, pp2sex was not asked of those with marstat='single, never married' due to a CAPI filtering error. In most cases this can be imputed, as shown below. Cases that could not be imputed are coded as 9 'not answered'.

(N) **pp2sex** sex of 2nd most recent live-in partner

1: heterosexual or if ppnum>=2 & ppnum<99 & eversam=2 & ((marstat=1 & hetlife>0 & hetlife<9999) | marstat=2 | marstat=4 | marstat=5 | marstat=6)

2: homosexual or if ppnum>=2 & ppnum<99 & marstat=1 & eversam=1 & (hetlife=0 | (hetlife>0 & hetlife<9999 & (marstat=3 | marstat=7 | marstat=8 | marstat=9)))

9: not answered

-1: not applicable if ppnum<2

(N) **pp2ystop** year stopped living with second most recent partner

9999: not answered

-1: <2 live-in partnerships if (ppnum<2)

(N) **pp2months** no. months lived with 2nd partner for if started and stopped cohab in same year

99: not answered

-1: not applicable if ppnum>=1 & ppnum<99 & pp1ystrt>0 & pp1ystrt<9999 & pp1ystop>0 & pp1ystop<9999 & pp1ystop#96 & pp1ystrt#pp1ystop

Notes: 1) Reasons 2nd most recent cohabiting relationship ended was only asked if the most recent partnership was ongoing, and the respondent had at least 2 cohabiting relationships ever.

2) Respondents who said that they were widowed (marstat=6) were not asked ppwhy to avoid causing distress. This is indicated in the variable pp1why25.

3) A CAPI routing error meant that ppwhy was only asked in loop 1 for waves 1-4 (first half) of fieldwork, therefore there is no information for wave 1-4 respondents who were married / in a civil partnership / cohabiting relationship at the time of interview about why their 2nd most recent cohabitation ended.

4) The binary variables pp2why1..pp2why24 were created directly from the question ppwhy. All variables coded in the same way as pp2why1.

(N) **pp2why1** Reason most recent cohabiting relationship ended: relationship has not ended1: mentioned

0: not mentioned

9: not answered

-1: not applicable if ppnum=-1

LIVE-IN PARTNERSHIPS

And similarly...

- (N) **pp2why2** Reason second most recent cohabiting relationship ended: death of partner
- (N) **pp2why3** Reason second most recent cohabiting relationship ended: difficulties with sex life
- (N) **pp2why4** Reason second most recent cohabiting relationship ended: other
- (N) **pp2why5** Reason second most recent cohabiting relationship ended: domestic violence
- (N) **pp2why6** Reason second most recent cohabiting relationship ended: different interests / nothing in common
- (N) **pp2why7** Reason second most recent cohabiting relationship ended: unfaithfulness / adultery
- (N) **pp2why8** Reason second most recent cohabiting relationship ended: arguments
- (N) **pp2why9** Reason second most recent cohabiting relationship ended: not having children
- (N) **pp2why10** Reason second most recent cohabiting relationship ended: grew apart
- (N) **pp2why11** Reason second most recent cohabiting relationship ended: moved because of change in circumstances
- (N) **pp2why12** Reason second most recent cohabiting relationship ended: lack of respect / appreciation
- (N) **pp2why13** Reason second most recent cohabiting relationship ended: not sharing housework enough
- (N) **pp2why14** Reason second most recent cohabiting relationship ended: money problems
- (N) **pp2why15** Reason second most recent cohabiting relationship ended: drink / drugs / gambling problems
- (N) **pp2why16** Reason second most recent cohabiting relationship ended: mental health or related problems
- (N) **pp2why17** Reason second most recent cohabiting relationship ended: problem with children
- (N) **pp2why18** Reason second most recent cohabiting relationship ended: never at home
- (N) **pp2why19** Reason second most recent cohabiting relationship ended: problems with parents / in-laws / family
- (N) **pp2why20** Reason second most recent cohabiting relationship ended: age problem
- (N) **pp2why21** Reason second most recent cohabiting relationship ended: another relationship involved
- (N) **pp2why22** Reason second most recent cohabiting relationship ended: lived in / moved to a different country
- (N) **pp2why23** Reason second most recent cohabiting relationship ended: changed mind / feelings / personality
- (N) **pp2why24** Reason second most recent cohabiting relationship ended: partner left without explanation

Notes: 1. Variables *pp1str* ... *pp1months* and their codes also apply to live-in partnerships 3 ... 5
 2. The fifth loop of live-in partnership questions relates to the first ever cohabiting relationship if *ppnum*>5.

Note: variable *age1cohb* is derived differently in Natsal-3 because of changes to the questions about previous partnerships. However, it is comparable with the Natsal-2 variables and therefore has not been renamed.

Note: Variable *age1cohb* is coded as 99 if respondents can't remember the month they first started cohabiting.

(U – see below) (D) **age1cohb** respondent's age when started living with first live-in partner, completed years

compute (((((pp1str-1900)*12) + ppm1str) - (((rdob1-1900)*12) + rdob1m)) / 12 if ((pp1str≠-1 & pp1str≠9999) & (ppm1str≠-1 & ppm1str≠97 & ppm1str≠99) & (rdob1m≠99))

99: not answered *else if* missing

-1: not applicable *if* *pplive*=2 | *pp1str*=-1

age1cohb respondent's age when started living with first live-in partner, completed years

compute *age1cohb* = (pp1str – rdob) *if* *ppnum*=1

compute *age1cohb* = (pp2str – rdob) *if* *ppnum*=2

compute *age1cohb* = (pp3str – rdob) *if* *ppnum*=3

compute *age1cohb* = (pp4str – rdob) *if* *ppnum*=4

compute *age1cohb* = (pp5str – rdob) *if* *ppnum*>=5

999: not answered *else if* missing

-1: not applicable *if* *pplive*=2 | *ppnum*=-1

(D) **cohab420** lived with someone before age 20
 1: yes *if* age1cohb>0 & age1cohb<20
 0: no *if* (age1cohb>=20 & age1cohb<99) | (age1cohb=-1)
 9: not answered *else if* missing
 -1: not applicable *if* dage<20

Natsal-2 PREVIOUS LIVE-IN PARTNERSHIPS QUESTIONS

(R) (N) **ppystrt** year started living with first live-in partner
 9999: not answered
 -1: not applicable *if* ppnum=-1

(R) (N) **ppmstrt** month started living with first live-in partner
 1: January
 2: February
 3: March
 4: April
 5: May
 6: June
 7: July
 8: August
 9: September
 10: October
 11: November
 12: December
 97: can't remember month
 99: not answered
 -1: not applicable *if* ppnum=-1

(R) (N) **ppms** first live-in partnership: type of union at start of cohabitation
 1: married
 2: living as a couple *or if* ppms=9 & marstat=7
 9: not answered
 -1: not applicable *if* ppnum=-1

(R) (N) **ppmar** later married first live-in partner
 1: yes
 2: no *if* pppmar=9 & marstat=7
 9: not answered
 -1: not applicable *if* ppnum=-1 | ppms=1

Note: Variable age1cohb is coded as 99 if respondents can't remember the month they first started cohabiting.

(ND) **age1cohb** respondent's age when started living with first live-in partner, completed years

$$\text{compute } (((\text{ppystrt}-1900)*12) + \text{ppmstrt}) - (((\text{rdoby}-1900)*12) + \text{rdobm}) / 12 \text{ if } ((\text{ppystrt} \neq -1 \ \& \ \text{ppystrt} \neq 9999) \ \& \ (\text{ppmstrt} \neq -1 \ \& \ \text{ppmstrt} \neq 97 \ \& \ \text{ppmstrt} \neq 99) \ \& \ (\text{rdobm} \neq 99))$$

 99: not answered *else if* missing
 -1: not applicable *if* pplive=2 | ppystrt=-1

(ND) **cohab420** lived with someone before age 20
 1: yes *if* age1cohb>0 & age1cohb<20
 0: no *if* (age1cohb>=20 & age1cohb<99) | (age1cohb=-1)
 9: not answered *else if* missing
 -1: not applicable *if* dage<20

LIVE-IN PARTNERSHIPS

(R) (N) **ppymar** year married first partner (if lived together before marriage)

9999: not answered

-1: not applicable if (ppmar=-1 | ppmar=2) | ppms=1

(R) (N) **ppmmar** month married first partner (if lived together before marriage)

1: January

2: February

3: March

4: April

5: May

6: June

7: July

8: August

9: September

10: October

11: November

12: December

97: can't remember

99: not answered

-1: not applicable if (ppmar=-1 | ppmar=2) | ppms=1

Notes: ppsex is not asked if currently single, even if have previously cohabited. This is because of a filter error in the CAPI. Unfortunately the group most likely to have cohabited with someone of the same sex (those reporting 'single, never married' at 'marstat') were not asked the sex of each partner they cohabited with. However, in most cases this can be imputed. If a respondent reports homosexual experience and never had sex with someone of the opposite sex, we can assume cohabiting partners were same sex (and vice versa.) However, if they reported same sex experience and have had sex with heterosexual partner(s), then we can't impute a value for ppsex.

(R) (N) **ppsex** sex of first live-in partner

1: heterosexual or if ppmar=1 | ppms=1

2: homosexual

9: not answered

-1: not applicable if marstat=7

(R) (N) **ppystop** year stopped living with first live-in partner (if ever did)

9999: not answered

-1: never lived with someone as a couple if pplive=2

-2: still living with partner if (ppnum=1 & marstat=1 | marstat=2 | marstat=3)

(R) (N) **ppmstop** month stopped living with first live-in partner (if ever did)

1: January

2: February

3: March

4: April

5: May

6: June

7: July

8: August

9: September

10: October

11: November

12: December

97: can't remember

99: not answered

-1: never lived with someone as a couple if pplive=2

-2: still living with partner if (ppnum=1 & marstat=1 | marstat=2 | marstat=3)

LIVE-IN PARTNERSHIPS

Note: Variables 'ppwhy...' only refer to the last previous cohabitation, not all previous cohabitations

(R) (N) **ppwhy01** reason 1 why live-in partnership 1 ended

1: unfaithfulness/adultery

2: money problems

3: difficulties with sex life

4: different interests/nothing in common

5: grew apart

6: not having children

7: lack of respect/appreciation

8: domestic violence

9: arguments

10: not sharing housework enough

11: moved because of change in circumstances

12: death of partner

13: other

99: not answered

-1: not applicable *if* ((ppnum>2) | (ppnum=1 & marstat=1 | marstat=2 | marstat=3 | marstat=4) | (ppnum=2 & marstat=4 | marstat=5 | marstat=6 | marstat=7) | (marstat=7 & pplive=2) | (ppwhyn=99 & ppwhyn-1≠99))

Similarly for up to 11 reasons:

(R) (N) **ppwhy02** reason 2 why live-in partnership 1 ended

(R) (N) **ppwhy03** reason 3 why live-in partnership 1 ended

(R) (N) **ppwhy04** reason 4 why live-in partnership 1 ended

(R) (N) **ppwhy05** reason 5 why live-in partnership 1 ended

(R) (N) **ppwhy06** reason 6 why live-in partnership 1 ended

(R) (N) **ppwhy07** reason 7 why live-in partnership 1 ended

(R) (N) **ppwhy08** reason 8 why live-in partnership 1 ended

(R) (N) **ppwhy09** reason 9 why live-in partnership 1 ended

(R) (N) **ppwhy10** reason 10 why live-in partnership 1 ended

(R) (N) **ppwhy11** reason 11 why live-in partnership 1 ended

(R) (N) **ppwhyx1** other reason 1 why live-in partnership 1 ended

1: unfaithfulness/adultery

2: money problems

3: difficulties with sex life

4: different interests/nothing in common

5: grew apart

6: not having children

7: lack of respect/appreciation

8: domestic violence

9: arguments

10: not sharing household chores enough

11: moved because of change in circumstances

12: death of partner

13: other

14: drink, drugs or gambling problem

15: mental health or related problem

16: problem with children/step-children

17: never at home (e.g. always out with friends)

18: problems with parents/in-laws/family

19: age-related problems (e.g. big age difference)

20: another relationship involved

21: lived in/moved to a different country/area

22: still in relationship, but stopped living together

23: change of mind/feelings/personality

24: partner just left without any explanation

99: not answered

-1: not applicable *if* ((ppnum>2) | (ppnum=1 & marstat=1 | marstat=2 | marstat=3 | marstat=4) | (ppnum=2 & marstat=4 | marstat=5 | marstat=6 | marstat=7) | (marstat=7 & pplive=2) | (ppwhyn=99 & ppwhyn-1≠99) | (ppwhy01... ppwhy11≠13))

LIVE-IN PARTNERSHIPS

Similarly for at most 3 other reasons:

(R) (N) **ppwhyx2** other reason 2 why live-in partnership 1 ended

(R) (N) **ppwhyx3** other reason 3 why live-in partnership 1 ended

Notes: 1. Variables *ppystrtn ... ppwhyxn* and their codes repeated for live-in partnerships 2 ... 12
2. Only 'ppwhy...' variables with valid codes are included in the final dataset, so the number of 'ppwhy...' variables varies by partnership number.

HOUSEHOLD COMPOSITION

HOUSEHOLD COMPOSITION

Note: In Natsal-2 hhszize-RelToR were here in the questionnaire. Moved to start of the household section in Natsal-3 and used to input into the relationship questions.

Note: Unlike in Natsal-1, variable 'tenure' in Natsal-2 has the additional code of 8.

(U – see below) (c5a U) **tenure** own/rent accommodation

- 1: own outright/with mortgage or loan
- 2: rent from council
- 3: rent from Housing Association
- 4: rent from private landlord
- 5: tied to job (including rent free)
- 6: squat
- 7: other
- 8: shared ownership (pays part rent and part mortgage)
- 9: lives rent-free
- 99: not answered

tenure own/rent accommodation – question wording identical but updated codes

- 1: own outright
- 2: buying with mortgage or loan
- 3: shared ownership (pays part rent and part mortgage)
- 4: Rent it
- 5: lives rent-free
- 9: not answered

Note: Variable 'tenure2' is consistent with Natsal-1's definition.

(R) (c5a) **tenure2** own/rent accommodation

- 1: own outright/with mortgage or loan
- 2: rent from council
- 3: rent from Housing Association
- 4: rent from private landlord
- 5: tied to job
- 6: squat
- 7: other (incl. shared ownership (pays part rent and part mortgage) and lives rent-free if tenure=7 | tenure=8 | tenure=9
- 9: not answered/don't know if tenure=99

(N) **livehere** always lived in village/town/city

- 1: yes
- 2: no
- 9: not answered

(c6c) **whenmove** age when moved to present village/town/city

- 1: not applicable if livehere=1
- 99: not answered

(R) (c6b) **bornin** country of birth

- 1: England (incl. Channel Islands)
- 2: Wales
- 3: Scotland
- 4: Northern Ireland/Eire
- 7: another country
- 9: not answered

(R) (D) **ebornin** global region of birth

- 1: Europe, EU country
- 2: Europe, non-EU country (incl. Eastern Europe, Russia, ...)
- 3: Australia/New Zealand

HOUSEHOLD COMPOSITION

- 4: North America
- 5: South America (incl. Mexico)
- 6: Caribbean countries
- 7: India, Pakistan, Bangladesh
- 8: China (incl. Hong Kong and Taiwan)
- 9: other Asia (incl. Thailand, Singapore and Indonesia)
- 10: Middle East and North Africa
- 11: West Africa
- 12: Central and Eastern Africa
- 13: South Africa
- 99: not answered

(R) (ND) **bornin_g** country/region of birth

- 1: UK *if* bornin=1 | bornin=2 | bornin=3
- 2: Ireland *if* bornin=4
- 3: other Europe *if* bornin=7 & (ebornin=1 | ebornin=2)
- 4: Caribbean *if* bornin=7 & ebornin=6
- 6: North Africa *if* bornin=7 & ebornin=10
- 7: Central/East Africa *if* bornin=7 & ebornin=12
- 8: West Africa *if* bornin=7 & ebornin=11
- 9: South Africa *if* bornin=7 & ebornin=13
- 10: India *if* bornin=7 & ebornin=7 & whasian=1
- 11: Pakistan *if* bornin=7 & ebornin=7 & whasian=2
- 12: other *if* bornin≠9 & ebornin≠99
- 99: not answered *else*

(R) (c6c) **agebrit** age first came to live in Britain

- 99: not answered/can't remember
- 1: not applicable *if* bornin=1 | bornin=2 | bornin=3

QUALIFICATIONS

EMPLOYMENT

Note: questions about employment were updated for Natsal-3 to collect information needed to code national statistics socioeconomic classification (NS-SEC) For more information on NS-SEC see: <http://www.ons.gov.uk/ons/guide-method/harmonisation/primary-set-of-harmonised-concepts-and-questions/index.html>

(R) (N) **ractiv01** occupation last week 1

- 1: full-time education (incl. if on holiday)
- 2: government training/employment scheme
- 3: paid employment (incl. self-employment)
- 4: waiting to take up paid work already obtained
- 5: unemployed and registered for benefit
- 6: unemployed, not registered, but looking for a job
- 7: unemployed, wanting a job, but not actively looking for a job
- 8: long-term sickness/disability
- 9: looking after the home/family
- 10: doing something else
- 11: retired
- 99: not answered
- 1: no other codes applicable *if ractivn=99 & ractivn-1≠99*

Similarly for up to 4 occupation codes:

(R) (N) **ractiv02** occupation last week 2

(R) (N) **ractiv03** occupation last week 3

(R) (N) **ractiv04** occupation last week 4

(R) (N) **ractivx1** other occupation last week 1

- 1: full-time education (incl. if on holiday)
- 2: government training scheme
- 3: paid employment (incl. self-employment)
- 4: waiting to take up paid work already obtained
- 5: unemployed and registered for benefit
- 6: unemployed not registered but looking for a job
- 7: unemployed wanting a job but not looking for one
- 8: long-term sickness/disability
- 9: looking after the home/family
- 10: doing something else
- 11: retired
- 12: temporarily sick/disabled
- 13: early retirement
- 99: not answered
- 1: no other codes applicable *if ractivn≠10*

Similarly for up to 3 other occupation codes:

(R) (N) **ractivx2** other occupation last week 2

(R) (N) **ractivx3** other occupation last week 3

QUALIFICATIONS

Note: Variable 'ractivhi' is derived by the CASI from 'ractiv01' ... 'ractiv04' and 'ractivx1' ... 'ractivx3'

- (R) (c7 D) **ractivhi** highest occupation code
- 1: full-time education (incl. if on holiday)
 - 2: government training/employment scheme
 - 3: paid employment (incl. self-employment)
 - 4: waiting to take up paid work already obtained
 - 5: unemployed and registered for benefit
 - 6: unemployed not registered but looking for a job
 - 7: unemployed wanting a job but not looking for a job
 - 8: long-term sickness/disability
 - 9: looking after the home/family
 - 10: doing something else
 - 11: retired
 - 12: temporarily sick/disabled
 - 13: early retirement
 - 99: not answered

Note: binary variables rfted..rother were created directly from the question ractiv in Natsal-3. All variables coded in the same way as rfted except where indicated. These binary variables were then used to create the derived variable ractivhi2.

- (N) **rfted** Occupation last week: Full-time education
- 1: yes
 - 0: no
 - 9: not answered

And similarly..

- (N) **rjpdwork** Occupation last week: Paid employment
- (N) **rgovtr** Occupation last week: Government scheme for employment training
- (N) **rueunpaid** Occupation last week: Unpaid work for a business that respondent or relative owns
- (N) **rwaitwork** Occupation last week: Waiting to take up paid work already obtained
- (N) **ruelookn** Occupation last week: Unemployed: looking for paid work / Government training scheme
- (N) **rtempsick** Occupation last week: Unemployed: Intending to look for work but prevented by temporary sickness or injury

Note: the response option 'long term sickness or disability' was only available to those within working age. The retirement age for men during the study period was 65. The retirement age for women changed over the study period: it was 60 in Natsal-3, 61 in 2011, and 62 in 2012. Therefore the binary variable rpermsick has an additional code of -1 for those who were not eligible for this response option.

- (N) **rpermsick** Occupation last week: long term sickness or disability
- 1: yes
 - 0: no
 - 9: not answered
 - 1: not applicable if (rsex=1 & dage>=66) | (rsex=2 & ((dage>=60 & dateyoi=2010) | (dage>=61 & dateyoi=2011) | (dage>=62 & dateyoi=2012)))

- (N) **retired** Occupation last week: Retired
- (N) **rhome** Occupation last week: Looking after home / family
- (N) **rother** Occupation last week: Doing something else

QUALIFICATIONS

(N) (D) **ractivhi2** highest occupational coding

- 1: full-time education
- 2: paid employment
- 3: government scheme for employment or training
- 4: unpaid work for a business that respondent or a relative owns
- 5: waiting to take up paid work already obtained
- 6: unemployed: looking for paid work / government training scheme
- 7: unemployed: intending to look for work but prevented by temporary sickness or injury
- 8: long term sickness or disability
- 9: retired
- 10: looking after home / family
- 11: doing something else
- 99: not answered

(N) **ractivhi3** highest occupational coding – comparable with Natsal-2 variable ractivhi

- 1: full-time education *if* rfted=1
- 2: government training / employment scheme *if* rgovtr=1
- 3: paid employment (incl. self-employment) *if* rjpdwork=1
- 4: waiting to take up paid work already obtained *if* rwaitwork=1
- 5: unemployed, looking for work or training *if* ruelookn=1
- 6: long term sickness / disability *if* rpermsick=1
- 7: looking after the home or family *if* rhome=1
- 8: doing something else *if* rother=1 | rueunpaid=1 | ractivhi2=8
- 9: retired *if* rretired=1
- 10: temporarily sick / disabled *if* rtempsick=1
- 99: not answered *if* ractivhi2=99

(ND) **employed** employment status at interview

- 1: employed *if* ractivhi2=2
- 2: in full-time education *if* ractivhi2=1
- 3: unemployed *if* ractivhi2=3 | ractivhi2=4 | ractivhi2=5 | ractivhi2=6 | ractivhi2=7 | ractivhi2=8 | ractivhi2=10 | ractivhi2=11
- 4: retired *if* ractivhi2=9
- 9: not answered *if* ractivhi2=99

(U – see below) (c9a*) **hours** hours worked per week, main job

- 1: 35 or more
- 2: 10, to less than 35
- 3: less than 10
- 9: not answered
- 1: not applicable *if* ractiv#3 & ractiv#99

hours hours worked per week, main job – question wording identical but updated codes

- 1: 50 or more
- 2: 35 to 49
- 3: 10, to less than 35
- 4: less than 10
- 9: not answered
- 1: not applicable *if* rjpdwork#1 & ractivhi2#99

Note: rlastjob was here in Natsal-2. Moved later in Natsal-3

(c9c U) **revraway** ever away with job for 1+ night(s) at a time

- 1: yes
- 2: no
- 9: not answered

(U) -1: not applicable *if* hours=-1 | hours=4 | hours=9

-1: not applicable *if* ractiv01#99 & ((ractivn=3 & hours=3) | (ractivn#3))

QUALIFICATIONS

(N) **rngtaway** no. of nights away with job in last year

999: not answered

-1: not applicable *if* revraway=2 | revraway=-1

Note: nights was added to determine eligibility for the saliva sample. Night workers were not eligible

(N) **rnights** usually work night shifts

1: yes

2: no

9: not answered

-1: not applicable *if* rjpdwork≠-1 & ractivhi2=99 | hours=4

Note: rlastjob was earlier in Natsal-2

(M) (c8 U) **rlastjob** when last had a paid job of 10+ hours a week

1: never

2: only occasional jobs (e.g. holiday/saturday jobs)

3: within past 6 months

4: at least 6 months but less than 1 year ago

5: at least 1 year but less than 5 years ago

6: at least 5 years but less than 10 year ago

7: at least 10 years but less than 20 year ago

8: at least 20 years ago

99: not answered

(U) -1: not applicable *if* hours=1 | hours=2 | hours=3 | rwaitwork=1

-1: not applicable *if* (ractivn=3 & hours=1 | hours=2) | ractivn=4

(R) (c10e) **rempstat** employee/self employed

1: employee

2: self-employed

9: not answered

-1: not applicable *if* rlastjob=1 | rlastjob=2 | rlastjob=7 | rlastjob=8

(R) (N) **rdirectr** director of a limited company

1: yes

2: no

9: not answered

-1: not applicable *if* (rlastjob=1 | rlastjob=2 | rlastjob=7 | rlastjob=8) | (rempstat=1)

(R) (N) **rmanage** manager/foreman/supervisor

1: manager

2: foreman/supervisor

3: other employee

9: not answered

-1: not applicable *if* (rlastjob=1 | rlastjob=2 | rlastjob=7 | rlastjob=8) | (rempstat=2)

(R) (N) **rempnum** no. of employees at (last) place of work (incl. respondent)

1: 1-2

2: 3-24

3: 25-499

4: 500+

9: not answered

-1: not applicable *if* (rlastjob=1 | rlastjob=2 | rlastjob=7 | rlastjob=8) | (rempstat=2 & rdirectr=2)

(R) (N) **rsenum** no. of employees respondent has/had

1: none

2: 1-24

3: 25-499

4: 500+

9: not answered

-1: not applicable *if* (rlastjob=1 | rlastjob=2 | rlastjob=7 | rlastjob=8) | (rempstat=1) | (rempstat=2 & rdirectr=1)

QUALIFICATIONS

Notes: 1. See the following website for the International Standard Occupation Codes (SOC):
<http://www.ons.gov.uk/ons/guide-method/classifications/archived-standard-classifications/standard-occupational-classification-2000/index.html>

2. Natsal-2 collected information to 3 digits, which corresponds to the minor group level, e.g.

Major group	2	Professional occupations
Sub major group	21	Science and technology professionals
Minor group	211	Science professionals

3. Natsal-3 data has socioeconomic variables for NS-SEC, and SOC2010 and SOC 2000 to allow comparisons with Natsal-2. Only the derived grouped variables used for analysis are included in the data. For NS-SEC, those in full time education have been given a separate code, rather than being left uncoded or classified based on their previous / part time jobs. See the following for more information:

<http://www.ons.gov.uk/ons/guide-method/classifications/current-standard-classifications/soc2010/index.html>

(N) **rnssecgp_6** respondent's NS-SEC (7 groups)

- 1: managerial & professional occupations
- 2: intermediate occupations
- 3: small employers and own account workers
- 4: lower supervisory and technical occupations
- 5: semi-routine and routine occupations
- 6: never worked or long term unemployed *if* rlastjob=1 | rlastjob=2
- 7: no job of 10+ hours/week or not in the last 10 years *if* rlastjob=7 | rlastjob=8
- 8: student in full-time education
- 99: not classifiable

(N) **rnssecgp_4** respondent's NS-SEC (4 groups)

- 1: managerial & professional occupations
- 2: intermediate occupations
- 3: semi-routine / routine occupations
- 4: never worked / no job of 10+ hrs/week / not in the last 10 years *if* rlastjob=1 | rlastjob=2 | rlastjob=7 | rlastjob=8
- 5: student in full-time education
- 99: not classifiable

(R) (N) **rsoc** respondent's International Standard Occupation Code

- 1: not applicable *if* rlastjob<3 | rlastjob>6
- 999: not available/can't code occupation

(R) (N) **rsoc_gp** respondent's standard occupation classification, grouped

- 1: not applicable if not had job in last 10 years
- 1: managers and senior officials
- 2: professional occupations
- 3: associate professional and technical occupations
- 4: administrative and secretarial occupations
- 5: skilled trade occupations
- 6: personal service occupations
- 7: sales and customer service occupations
- 8: process, plant and machine occupations
- 9: elementary occupations
- 999: not available / can't code occupation

QUALIFICATIONS

(N) **rsoc2000_9** Respondent's SOC 2000 classification (9 major groups)

1: managers and senior officials

2: professional occupations

3: associate professional and technical occupations

4: administrative and secretarial occupations

5: skilled trades occupations

6: personal service occupations

7: sales and customer service occupations

8: process, plant and machine operatives

9: elementary occupation

99: not coded *if* unable to classify occupation

-1: not applicable *if* rlastjob=1 | rlastjob=2 | (rlastjob>6 & rlastjob<99) | hours=4 | (rjpdwork=0 & rwaitwork=0) OR did not answer one or more questions needed to code occupation

QUALIFICATIONS

(N) **rsoc2000_7** Respondent's SOC 2000 classification (7 major groups)

1: managers and senior officials

2: professionals

3: associate professional & technical

4: administrative, secretarial, personal, sales & customer services

5: skilled trades

6: process, plant & machine operatives

7: elementary occupations

99: not coded *if* unable to classify occupation

-1: not applicable *if* rlastjob=1 | rlastjob=2 | (rlastjob>6 & rlastjob<99) | hours=4 | (rjpdwork=0 & rwaitwork=0) OR did not answer one or more questions needed to code occupation

(N) **rsoc2000_4** Respondent's SOC 2000 classification (4 major groups)

1: managers, professional, technical

2: administrative, sales, customer services, skilled trades

3: process, plant & machine operatives, elementary occupations

4: other

99: not coded *if* unable to classify occupation

-1: not applicable *if* rlastjob=1 | rlastjob=2 | (rlastjob>6 & rlastjob<99) | hours=4 | (rjpdwork=0 & rwaitwork=0) OR did not answer one or more questions needed to code occupation

(N) **rsoc2010_9** Respondent's SOC2010 group (9 major groups)

1: managers, directors and senior officials

2: professional occupations

3: associate professional and technical occupations

4: administrative and secretarial occupations

5: skilled trades occupations

6: caring, leisure and other service occupations

7: sales and customer service occupations

8: process, plant and machine operatives

9: elementary occupations

99: not coded *if* unable to classify occupation

-1: not applicable *if* rlastjob=1 | rlastjob=2 | (rlastjob>6 & rlastjob<99) | hours=4 | (rjpdwork=0 & rwaitwork=0) OR did not answer one or more questions needed to code occupation

(N) **rsoc2010_7** Respondent's SOC2010 group (7 major groups)

1: managers, directors and senior officials

2: professionals

3: associate professional and technical

4: administrative, secretarial, personal, sales & customer services

5: skilled trades

6: process, plant & machine operatives

7: elementary occupations

99: not coded *if* unable to classify occupation

-1: not applicable *if* rlastjob=1 | rlastjob=2 | (rlastjob>6 & rlastjob<99) | hours=4 | (rjpdwork=0 & rwaitwork=0) OR did not answer one or more questions needed to code occupation

(N) **rsoc2010_4** Respondent's SOC2010 group (4 major groups)

1: managers, professional, technical

2: administrative, sales, customer services, skilled trades

3: process, plant & machine operatives, elementary occupations

4: other

99: not coded *if* unable to classify occupation

-1: not applicable *if* rlastjob=1 | rlastjob=2 | (rlastjob>6 & rlastjob<99) | hours=4 | (rjpdwork=0 & rwaitwork=0) OR did not answer one or more questions needed to code occupation

QUALIFICATIONS

Note: The following categories are taken from the website:

http://www.statistics.gov.uk/methods_quality/ns_sec/continuity.asp

(R) (N) **rseg** respondent's socio-economic group

- 1.1: employers in industry, commerce, etc. - large establishments
- 1.2: managers in government, industry, commerce, etc. - large establishments
- 2.1: employers in industry, commerce, etc. - small establishments
- 2.2: managers in industry, commerce, etc. - small establishments
- 3.0: professional workers - self-employed
- 4.0: professional workers - employees
- 5.1: intermediate non-manual workers: ancillary workers and artists
- 5.2: intermediate non-manual workers: foremen and supervisors
- 6.0: junior non-manual workers
- 7.0: personal service workers
- 8.0: foremen and supervisors - manual
- 9.0: skilled manual workers
- 10.0: semi-skilled manual workers
- 11.0: unskilled manual workers
- 12.0: own account workers (except professional)
- 13.0: farmers - employers and managers
- 14.0: farmers - own account
- 15.0: agricultural workers
- 16.0: members of armed forces
- 999.9: not available
- 1: not applicable *if* rlastjob<3 | rlastjob>6

(R) (c10sc/soclas U) **rsc** respondent's social class

- 1: professional/managerial
- 2: managerial/technical
- 3.1: skilled non-manual
- 3.2: skilled manual
- 4: partly skilled
- 5: unskilled
- 6: unclassifiable/army/other
- 1: not applicable *if* rlastjob<3 | rlastjob>6

(R) (ND) **rsc_gp** respondent's social class is manual/non-manual

- 0: manual *if* rsc=1 | rsc=2 | rsc=3.1
- 1: non-manual *if* rsc=1 | rsc=2 | rsc=3.2
- 98: unclassifiable/army/other *if* rsc=6
- 9: not answered *if* rsc=9
- 1: not applicable *if* rsc=-1

(R) (ND) **rsc_gp2** respondent's social class, grouped

- 1: I/II *if* rsc=1 | rsc=2
- 2: III non-manual/III manual *if* rsc=3.1 | rsc=3.2
- 3: IV/V *if* rsc=4 | rsc=5
- 9: not answered *if* rsc=9
- 1: not applicable/unclassifiable *if* rsc=-1 | rsc=98

(R) (ND) **rsc_kw** respondent's social class, grouped

- 1: I/II/III non-manual/III manual *if* rsc=1 | rsc=2 | rsc=3.1 | rsc=3.2
- 2: IV/V/ unemployed *if* rsc=4 | rsc=5 | ((rsc=6 | rsc=-1) & (ractivhi=5 | ractivhi=6 | ractivhi=7))
- 9: not answered/not applicable/unclassifiable *else*

Notes: 1. Variable, 'rsic' categorises people by the nature of their industry.

2. Details on the category codes are given at the website:

http://www.statistics.gov.uk/methods_quality/sic/contents.asp

(R) (N) **rsic** respondent's Standard Industrial Classification

QUALIFICATIONS

(R) (N) **cie** who has the largest income from employment, state benefits, investments, or other sources

1: respondent

2: partner

3: both equally

9: not answered

-1: not applicable if marstat=4 | marstat=5 | marstat=6 | marstat=7

Note: The questions on employment and occupation are repeated for the respondent's partner, with the same codes, in addition to: -1 if marstat=4 | marstat=5 | marstat=6 | marstat=7 Natsal-1/Natsal-2 or if spoushh#1 & civpthh#1 & coopphh#1 & cosamhh#1 & marstat#2 & marstat#3 in Natsal-3

(R) (c11) **pactiv01** partner's occupation last week 1

Similarly for up to 3 occupations:

(R) (c11) **pactiv02** partner's occupation last week 2

(R) (c11) **pactiv03** partner's occupation last week 3

(R) (c11) **pactivx1** partner's other occupation last week 1

Similarly for up to 2 other occupations:

(R) (c11) **pactivx2** partner's other occupation last week 2

(R) (c11) **pactivhi** partner's highest occupation code

(N) **pfted** Partner's occupation last week: Full-time education

1: yes

0: no

9: not answered

(N) **pjpdwork** Partner's occupation last week: Paid employment

(N) **pgovtr** Partner's occupation last week: Government scheme for employment training

(N) **pueunpaid** Partner's occupation last week: Unpaid work for a business that respondent or relative owns

(N) **pwaitwork** Partner's occupation last week: Waiting to take up paid work already obtained

(N) **puelookn** Partner's occupation last week: Unemployed: looking for paid work / Government training scheme

(N) **ptempsick** Partner's occupation last week: Unemployed: Intending to look for work but prevented by temporary sickness or injury

Note: the response option 'long term sickness or disability' was only available to those within working age. The retirement age for men during the study period was 65. The retirement age for women changed over the study period: it was 60 in Natsal-3, 61 in 2011, and 62 in 2012. Therefore the binary variable ppermsick has an additional code of -1 for those whose partners were not eligible for this response option, based on reltor/gender/ageof variables.

(N) **ppermsick** Partner's occupation last week: long term sickness or disability

1: yes

0: no

9: not answered

-1: not applicable if (spoushh#1 & civpthh#1 & coopphh#1 & cosamhh#1 & marstat#2 & marstat#3) |

(reltor!=1 | reltor!=2) & ((gender!=1 & ageof!>=66) | (gender!=2 & ((ageof!>=60 & dateyoi=2010) |

(ageof!>=61 & dateyoi=2011) | (ageof!>=62 & dateyoi=2012))))

(N) **pretired** Partner's occupation last week: Retired

(N) **phome** Partner's occupation last week: Looking after home / family

(N) **pother** Partner's occupation last week: Doing something else

(N) **pactivhi2** Partner's highest occupational coding

(N) **pactivhi3** Partner's highest occupational coding – comparable with Natsal-2 variable pactivhi

QUALIFICATIONS

(U) (c13a*) **phours** hours worked per week, main job, partner
phours hours worked per week, main job – question wording identical but updated codes
1: 50 or more
2: 35 to 49
3: 10, to less than 35
4: less than 10
9: not answered
-1: not applicable if spoushh#1 & civpthh#1 & coopphh#1 & cosamhh#1 & marstat#2 & marstat#3

Note: plastjob was here in Natsal-2. Moved later in Natsal-3

(c13c U) **pevraway** partner ever away with job for 1+ night at a time
(N) **pngtaway** no. of nights partner away with job in last year

(M) (c12 U) **plastjob** when partner last had a paid job of 10+ hours a week

(c14e) **pempstat** partner is/was an employee or self-employed
(R) (N) **pdirctr** partner is/was the director of a limited company
(R) (N) **pmanage** partner is/was a manager/foreman/supervisor

(R) (c14g) **pempnum** no. of employees at partner's (last) place of work

(R) (c14g) **psenum** no. of employees partner has/had

(N) **pnsecgp_6** partner's NS-SEC code (7 groups)
(N) **pnsecgp_4** partner's NS-SEC code (4 groups)

Note: Variable 'psoc' has the additional code of -1 if cie=-1 | cie=1 | cie=9

(R) (N) **psoc** partner's International Standard Occupation Code

(N) **psoc2010_9** partner's SOC2010 classification (9 major groups)
(N) **psoc2010_7** partner's SOC2010 classification (7 major groups)
(N) **psoc2010_4** partner's SOC2010 classification (4 major groups)

(N) **psoc2000_9** partner's SOC2000 classification (9 major groups)
(N) **psoc2000_7** partner's SOC2000 classification (7 major groups)
(N) **psoc2000_4** partner's SOC2000 classification (4 major groups)

(R) (N) **pseg** partner's socio-economic group
(R) (N) **psc** partner's social class
(R) (ND) **psc_gp** partner is in a manual/non-manual social class
(R) (ND) **psc_gp2** partner's social class, grouped
(R) (N) **psic** partner's Standard Industrial Classification

Note: Variable 'sc' is derived as outlined in the Natsal-1 codebook but also makes use of the variable cie', which was not asked in Natsal-1

(R) (sc D) **sc** household social class
if (ms# 1 & ms#2 & ms#3) then rsc
else (ms=1 | ms=2 | ms=3) & (cie=1 | cie=3) then rsc
else (ms=1 | ms=2 | ms=3) & (cie=2) then psc
else sc = 9
1: professional/managerial
2: managerial/technical
3.1: skilled non-manual
3.2: skilled manual
4: partly skilled
5: unskilled
6: unclassified/army/other
9: not answered
-1: not applicable

QUALIFICATIONS

Note: A proxy for household NS-SEC was derived in Natsal-3, taking the higher value from respondent's NS-SEC or partner's NS-SEC.

(N) **hhnssecgp_6** household NS-SEC group (7 groups)

- 1: managerial & professional occupations *if* rnssecgp_6=1 | pnssecgp_6=1
- 2: intermediate occupations *else if* rnssecgp_6=2 | pnssecgp_6=2
- 3: small employers and own account workers *else if* rnssecgp_6=3 | pnssecgp_6=3
- 4: lower supervisory and technical occupations *else if* rnssecgp_6=4 | pnssecgp_6=4
- 5: semi-routine and routine occupations *else if* rnssecgp_6=5 | pnssecgp_6=5
- 6: never worked or long term unemployed *else if* rnssecgp_6=6 | pnssecgp_6=6
- 7: no job of 10+ hours/week or not in the last 10 years *else if* rnssecgp_6=7 | pnssecgp_6=7
- 8: student in full-time education
- 99: not classifiable *else if* rnssecgp_6=9 & pnssecgp_6=9

(N) **hhnssecgp_4** household NS-SEC (4 groups)

- 1: managerial & professional occupations *if* rnssecgp_4=1 | pnssecgp_4=1
- 2: intermediate occupations *else if* rnssecgp_4=2 | pnssecgp_4=2
- 3: semi-routine / routine occupations *else if* rnssecgp_4=3 | pnssecgp_4=3
- 4: never worked / no job of 10+ hrs/week / not in the last 10 years *else if* rnssecgp_4=4 | pnssecgp_4=4
- 5: student in full-time education
- 99: not classifiable

(N) **income** household income per year pre tax (inc benefits, pensions etc)

- 1: <2,500
- 2: 2,500-4,999
- 3: 5,000-9,999
- 4: 10,000-19,999
- 5: 20,000-29,999
- 6: 30,000-39,999
- 7: 40,000-49,999
- 8: 50,000+
- 99: not answered

QUALIFICATIONS

Note: Variable 'tedage' has the additional code of 95 in Natsal-2's ethnic boost dataset

(U) (N) **tedage** age completed continuous full-time education

(U – code removed) 95: never went to school

96: not yet finished *or if* rfted=1

(U) 97: never went to school

99: not answered/can't remember

(c15a) **anyexam** passed any exams/got any qualifications

- 1: yes
- 2: no, none
- 9: not answered

Notes:

- 1. Variable 'educ' does not apply to respondents who were aged 16 at interview as they may not have (yet) had an opportunity to take any exams/gain any qualifications
- 2. Natsal-1's variable 'educ' is equivalent to Natsal-2's derived variable 'educ3'
- 3. Respondents who left school aged 17+ with no qualifications are coded as 'not answered' in educ2.

QUALIFICATIONS

(U) (ND) **educ** left school at 16 with/without passing any exams/gaining any qualifications

1: left school at 16 without passing any exams/gaining any qualifications *if* dage>16 & tedage>0 & tedage<17 & anyexam=2

2: left school at 16 having passed some exams/gained some qualifications *if* dage>16 & tedage>0 & tedage<17 & anyexam=1

(U) 3: left school aged 17+ *if* dage>16 & tedage>=17 & tedage<=96

3: left school aged 17+ *if* dage>16 & tedage>=17 & tedage=96 | tedage=95 9: not answered/don't know *else if* missing

(U) -1: not applicable *if* dage=16

(ND) **educ2** left school at 16 with/without passing any exams/gaining any qualifications (version 2)

1: left school at or before 16 without passing any exams/gaining any qualifications *if* dage>16 & tedage>0 & tedage<17 & anyexam=2

2: left school having passed some exams/gained some qualifications *if* dage>16 & ((tedage>0 & tedage<17 & anyexam=1) | (tedage>=17 & tedage<=96) & anyexam=1))

9: not answered/don't know *else if* missing

(U) -1: not applicable *if* dage=16 | tedage=97 -1: not applicable *if* dage=16

(c15b*U) **exams** highest academic qualification

1: degree level qualification

2: A-levels

3: AS-levels

4: SLC Higher Grade

5: O-level, 1975 or earlier

6: O-level, after 1975 A-C

7: O-level, after 1975 D-E

8: GCSE grades A-C

9: GCSE grades D-G

10: CSE grade 1

11: CSE grade 2-5

12: CSE Ungraded

13: SLC Lower

14: SUPE Lower or Ordinary

15: School certificate

16: Foreign qualification

(U – response option removed) 17: other educational qualification

99: not answered

-1: not applicable *if* anyexam=2

(ND) **exams2** highest academic qualification, grouped

1: degree level qualification *if* exams=1

2: A-levels/AS-levels/SLC Higher Grade *if* exams=2 | exams=3 | exams=4

(U) 3: O-level/other *if* exams>=5 & exams<=16 3: O-level/other *if* exams>=5 & exams<=17

4: none *if* exams=-1

9: not answered/don't know *if* exams=99

(N) **anyqual** passed any non-academic exams/qualifications

1: yes

2: no, none

9: not answered

Note: In Natsal-3 the question 'quals' was converted into binary variables, each relating to a different response option. All are coded in the same way as quteach.

(N) **quteach** non-academic exams passed/qualifications gained: teaching qualification

1: yes

0: no or *if* anyqual=2

9: not answered

QUALIFICATIONS

(N) **qunurse** non-academic exams passed/qualifications gained: nursing qualification
 (N) **quhnhnd** non-academic exams passed/qualifications gained: HNC/HND
 (N) **quoncond** non-academic exams passed/qualifications gained: ONC/OND
 (N) **qucandgfull** non-academic exams passed/qualifications gained: City and Guilds, full
 (N) **qucandgadv** non-academic exams passed/qualifications gained: City and Guilds, advanced
 (N) **qucandgcra** non-academic exams passed/qualifications gained: City and Guilds, craft
 (N) **qunvqlev5** non-academic exams passed/qualifications gained: NVQ Level 5
 (N) **qunvqlev4** non-academic exams passed/qualifications gained: NVQ Level 4
 (N) **qunvqlev3** non-academic exams passed/qualifications gained: NVQ Level 3
 (N) **qunvqlev2** non-academic exams passed/qualifications gained: NVQ Level 2
 (N) **qunvqlev1** non-academic exams passed/qualifications gained: NVQ Level 1
 (N) **quappren** non-academic exams passed/qualifications gained: recognised trade apprenticeship completed
 (N) **quclercomm** non-academic exams passed/qualifications gained: clerical or commercial qualification
 (N) **quothqual** non-academic exams passed/qualifications gained: other vocational qualification

(R) (c15b*U) **quals01** non-academic exams passed/qualifications gained 1

1: teaching qualification
 2: nursing qualification
 3: HNC/HND
 4: ONC/OND
 5: City and Guilds, Full
 6: City and Guilds, Advanced
 7: City and Guilds, Craft
 8: NVQ Level 5
 9: NVQ Level 4
 10: NVQ Level 3
 11: NVQ Level 2
 12: NVQ Level 1
 13: recognised trade apprenticeship completed
 14: clerical or commercial qualification
 15: other vocational or professional qualification
 99: not answered
 -1: not applicable *if* anyqual=2 | (qualsn=99 & qualsn-1≠99)

Similarly for up to 11 non-academic qualifications:

(R) (c15b*U) **quals02** non-academic exams passed/qualifications gained 2
 (R) (c15b*U) **quals03** non-academic exams passed/qualifications gained 3
 (R) (c15b*U) **quals04** non-academic exams passed/qualifications gained 4
 (R) (c15b*U) **quals05** non-academic exams passed/qualifications gained 5
 (R) (c15b*U) **quals06** non-academic exams passed/qualifications gained 6
 (R) (c15b*U) **quals07** non-academic exams passed/qualifications gained 7
 (R) (c15b*U) **quals08** non-academic exams passed/qualifications gained 8
 (R) (c15b*U) **quals09** non-academic exams passed/qualifications gained 9
 (R) (c15b*U) **quals10** non-academic exams passed/qualifications gained 10
 (R) (c15b*U) **quals11** non-academic exams passed/qualifications gained 11

QUALIFICATIONS

Note: Variables 'qualsx1', 'qualsx2' and 'qualsx3' do not use code 15

(R) (c15b*U) **qualsx1** Other non-academic exams passed/qualifications gained 1

- 1: teaching qualification
- 2: nursing qualification
- 3: HNC/HND
- 4: ONC/OND
- 5: City and Guilds, Full
- 6: City and Guilds, Advanced
- 7: City and Guilds, Craft
- 8: NVQ Level 5
- 9: NVQ Level 4
- 10: NVQ Level 3
- 11: NVQ Level 2
- 12: NVQ Level 1
- 13: recognised trade apprenticeship completed
- 14: clerical or commercial qualification
- 16: other vocational qualification
- 17: NVQ level not specified
- 18: nursery nurse examination board qualification
- 19: qualifications obtained through military
- 20: other professional qualification
- 99: not answered
- 1: not applicable *if qualsn#15*

Similarly for up to 3 other non-academic qualifications:

(R) (c15b*U) **qualsx2** Other non-academic exams passed/qualifications gained 2

(R) (c15b*U) **qualsx3** Other non-academic exams passed/qualifications gained 3

Note: Variable 'educ3' is equivalent to variable 'educ' in Natsal-1

Note: the coding of educ3 has been updated in this codebook to reflect the fact that binary vars quteach..quothqual are used, rather than the Natsal-2 vars qualsx1..qualsx3. However, the variable is essentially the same so the variable name has not been changed.

(U – see below) (educ D) **educ3** highest academic/non-academic qualification, grouped

- 1: degree *if exams=1*
- 2: higher education, <degree level but A-level/equivalent *if exams=2 | exams=3 | exams=4 | (quals01>=1 & quals01<=6) | ... | (quals11>=1 & quals11<=6) | (qualsx1>=1 & qualsx1<=6) | (qualsx2>=1 & qualsx2<=6) | (qualsx3>=1 & qualsx3<=6) | (quals01>=18 & quals01<=20) | similarly up to: | (quals11>=18 & quals11<=20) | (qualsx1>=18 & qualsx1<=20) | similarly up to: | (qualsx3>=18 & qualsx3<=20)*
- 3: GCSE, O-level or equivalent *if exams>=5 & exams<=15 | (quals01>=7 & quals01<=17) | similarly up to | (quals11>=7 & quals11<=17) | (qualsx1>=7 & qualsx1<=17) | similarly up to | (qualsx3>=7 & qualsx3<=17)*
- 4: foreign or other *if exams=16 | exams=17*
- 5: none *if anyexam=2*
- 9: not answered/don't know *else if missing*

QUALIFICATIONS

(U) **educ3** highest academic/non-academic qualification, grouped

- 1: degree *if* exams=1
- 2: higher education, <degree level but A-level/equivalent *if* exams=2 | exams=3 | exams=4 | quteach=1 | qunurse=1 | quhnchnd=1 | quoncond=1 | qucandgfull=1 | qucandgadv=1
- 3: GCSE, O-level or equivalent *if* (exams>=5 & exams<=15) | qucandgcra=1 | qunvqlev5=1 | qunvqlev4=1 | qunvqlev3=1 | qunvqlev2=1 | qunvqlev1=1 | quappren=1 | quclercomm=1 | quothqual=1
- 4: foreign or other *if* exams=16 | exams=17
- 5: none *if* anyexam=2
- 9: not answered/don't know *else if* missing

Note: educ4 classifies people aged 17+ into three groups, according to their highest academic qualification. It is assumed that people who already have a qualification typically gained at age 16 (GCSEs or equivalent), who are still in full-time education, will go on to achieve a further qualification. Those with foreign qualifications are coded as 'not answered' as we do not have information on what level these qualifications are. Those aged 17+ in full-time education who do not have a qualification typically gained at age 16 are coded as 'not answered' as we do not have information on what level of qualification they are studying towards.

(N) **educ4** highest academic qualification

- 1: not applicable *if* dage=16
- 1: no academic qualifications *if* dage>16 & tedage#96 & tedage#99 & anyexam=2
- 2: academic qualifications typically gained at age 16 *if* dage>16 & exams>=5 & exams<=15 & tedage#96
- 3: studying for/attained further academic qualifications *if* dage>16 & ((exams>0 & exams<5) | (tedage=96 & exams>=5 & exams<=15))
- 9: not answered *else if* dage>16 & (exams=99 | exams=16 | tedage=96 | tedage=99)

(R) (N) **qualyr** year obtained most recent academic/non-academic qualification

- 9999: not answered
- 1: not applicable *if* anyexam=2 & anyqual=2

(c16) **typesch** last school was mixed/single-sex

- 1: mixed school
- 2: single sex school
- 3: single sex school to 6th form but mixed 6th form
- 9: not answered/can't remember

(ND) **singlsex** last school was single-sex

- 0: no *if* typesch=1 | typesch=3
- 1: yes *if* typesch=2
- 9: not answered/can't remember *if* typesch=9

(N) **boarder** any of secondary schooling as a boarder

- 1: yes
- 2: no
- 9: not answered

PARENTS OCCUPATION

PARENT'S OCCUPATION

Notes: 1. Variables *par1occ ... par3occ* refer to the respondent's father's occupation if they lived with both parents, else the parent that the respondent lived with

2. In Natsal-3 age of *par1occ* variable was 14 to tie up with *wholive*.

3: Change of mode of administration: In Natsal-2 respondents were asked to code their parent's occupation from the list. After feedback that respondents found it difficult to determine the correct code, this was changed in Natsal-3 so that respondents told the interviewer their parent's occupation and the interviewer coded it to one of the options on the list. Where the interviewer was unsure of what to code, they recorded the open text answer at *xpar1occ* and this was backcoded to the codeframe by head office staff where possible.

(*) (N) **par1occ** parent's occupation when respondent was aged 14 (previously 16)

- 1: farmer/farm manager
- 2: farm worker
- 3: skilled manual worker
- 4: semi-skilled/unskilled manual worker
- 5: professional/technical worker
- 6: manager/administrator
- 7: clerical
- 8: sales
- 9: never had job/housewife/permanently sick/disabled
- 10: other
- 11: can't say
- 12: Armed Forces
- 13: didn't have a job at the time/in full-time education
- 99: not answered/don't know
- 1: not applicable if (bothmapa=2 & ynotboth=3) | (bothmapa=2) & (maorpa=3 | maorpa=4))

(U – see below) (N) **par2occ** was parent self-employed/employee

- 1: self-employed/had own business/farm
- 2: worked for someone else
- 9: not answered/don't know
- 1: not applicable if *par1occ*=-1 | (*par1occ*>=9 & *par1occ*<=13)

par2occ was parent self-employed/employee

- 1: employee
- 2: self-employed
- 9: not answered/don't know

(*U – see below) (N) **par3occ** was parent a supervisor/foreman

- 1: supervisor/foreman of manual workers
- 2: supervisor/foreman of non-manual workers
- 3: neither
- 9: not answered/don't know
- 1: not applicable if *par1occ*=-1 | (*par1occ*>=9 & *par1occ*<=13)

par3occ responsible for supervising others

- 1: yes
- 2: no

PARENTS OCCUPATION

Note: due to changes to the response options for par3occ, it was not possible to derive parsc, parsc_gp or parsc2 for Natsal-3. Instead use vars parsc3 or parsc_kw.

Note: Variable 'parsc' is automatically derived from responses to 'par1occ' and 'par3occ'

(R) (DN) **parsc** parent(s) social class

- 1: professional/managerial
- 2: managerial/technical
- 3: skilled non-manual
- 4: skilled manual
- 5: partly skilled/unskilled
- 6: never had a job
- 7: unclassifiable/army/other
- 8: can't say
- 9: not answered/don't know

(U) -1: not applicable *if bothmapa2=2 & (wholive<2 | wholive>5)*

-1: not applicable *if bothmapa=2 & (ynotboth=3 | (maorpa#1 & maorpa#2 & maorpa#-1 & maorpa#9))*

(R) (DN) **parsc_gp** parents social class, grouped

- 0: manual *if parsc=4 | parsc=5*
- 1: non-manual *if parsc=1 | parsc=2 | parsc=3*
- 9: not answered/don't know *if parsc=9*
- 98: unclassified/army/other *if parsc>=6 & parsc<=8*
- 1: not applicable *if parsc=-1*

(R) (DN) **parsc2** parents social class, grouped

- 1: I/II *if parsc=1 | parsc=2*
- 2: III (skilled manual/non-manual) *if parsc=3 | parsc=4*
- 3: IV/V *if parsc=5*
- 9: not answered/don't know/unclassifiable *if parsc>=6 & parsc<=9*
- 1: not applicable *if parsc=-1*

(ND) **parsc3** parents social class, grouped

- 1: professional / managerial *if par1occ=1*
- 2: technical / skilled *if par1occ= 1 | par1occ=3 | par1occ=6 | par1occ=7 | par1occ=8 | (par3occ=1 & (par2occ=2 | par1occ=4))*
- 3: partly skilled / unskilled *if par3occ=2 & (par1occ=2 | par1occ=4)*
- 4: never had a job *if par1occ=9*
- 5: unclassifiable / army / other *if par1occ=10 | par1occ=12 | par1occ=13*
- 6: can't say *if par2occ=11*
- 9: not answered *else if missing*
- 1: not applicable *if par1occ=-1*

(U –see below) (DN) **parsc_kw** parents social class, grouped

- 1: I/II/III *if parsc>=1 & parsc<=4*
- 2: IV/V or unemployed *if parsc=5 | parsc=6 | ((parsc=7 | parsc=8) & (par1occ=9))*
- else* -1: not applicable *if parsc=-1*
- else* 9: not answered/can't say/unclassifiable

parsc_kw parents social class, grouped

- 1: I/II/III *if parsc3=1 | parsc3=2*
- 2: IV/V or unemployed *if parsc3=3 | parsc3=4 | (par1occ=9 & (parsc3=5 | parsc3=6))*
- else* -1: not applicable *if parsc3=-1*
- else* 9: not answered/can't say/unclassifiable

ETHNICITY

ETHNICITY

Note: ethnic was moved here Natsal-3. After religion in Natsal-2. Only the Office for National Statistics harmonised question on ethnicity was asked in Natsal-3. In Natsal-2 there were several other ethnicity questions because of the ethnic minority boost sample. These have been removed in Natsal-3.

(U – see below) (U) **ethnic** ethnic group that the respondent considers themselves belonging to

- 1: white
- 2: mixed
- 3: asian/asian british
- 4: black/black british
- 5: Chinese
- 6: other

ethnic ethnic group that respondent considers themselves belonging to

- 1: white british
- 2: white irish
- 3: white other
- 4: mixed white and black Caribbean
- 5: mixed white and black African
- 6: mixed white and asian
- 7: mixed other
- 8: Asian/Asian British: indian
- 9: Asian/Asian British: Pakistani
- 10: Asian/Asian British: bangladeshi
- 11: Asian/Asian British: other
- 12: black/black British: Caribbean
- 13: black/black British: African
- 14: black/black British: other
- 15: Chinese
- 16: other ethnic group
- 99: not answered

(ND) **ethnicgrp** ethnic group that respondent considers themselves to belong to, grouped – comparable with ethnic in Natsal-2

- 1: white *if ethnic*>=1 & *ethnic*<=3
- 2: mixed *if ethnic*>=4 & *ethnic*<=7
- 3: asian/asian british *if ethnic*>=8 & *ethnic*<=11
- 4: black/black british *if ethnic*>=12 & *ethnic*<=14
- 5: Chinese *if ethnic*=15
- 6: other *if ethnic*=16
- 9: not answered

(R) (N) **whwhite** cultural background of white respondents

- 1: British
- 2: Irish
- 3: other background
- 9: not answered
- 1: not applicable *if ethnic*≠1 & *ethnic*≠9

(R) (N) **whmixed** cultural background of respondents from a mixed ethnic group

- 1: white British & black Caribbean
- 2: white British & black African
- 3: white British & Asian
- 4: other mixed background
- 9: not answered
- 1: not applicable *if ethnic*≠2 & *ethnic*≠9

ETHNICITY

(R) (N) **whasian** cultural background of Asian respondents

- 1: Indian
- 2: Pakistani
- 3: Bangladeshi
- 4: other Asian background
- 9: not answered
- 1: not applicable *if ethnic#3 & ethnic#9*

Note: Variable 'whmixas' is only in the ethnic boost dataset

(R) (N) **whmixas** cultural background of white/asian respondents

- 1: white British & Indian
- 2: white British & Pakistani
- 3: white British & other Asian background
- 9: not answered
- 9999: not asked in Natsal-2 core questionnaire
- 1: not applicable *if ethnic#3 & ethnic#9*

(R) (N) **whblack** cultural background of black respondents

- 1: Caribbean
- 2: African
- 3: other black background
- 9: not answered
- 1: not applicable *if ethnic#4 & ethnic#9*

Note: Variable 'afrcoun' only had more codes available in the Ethnic Boost questionnaire compared to the Core questionnaire. Thus, the Core has been recoded to have codes consistent with the Ethnic Boost – see variable 'afrcoun2'

(R) (N) **afrcoun** country of origin of black African respondents

- 1: Democratic Republic of Congo (Zaire)
- 2: Ghana
- 3: Kenya
- 4: Nigeria
- 5: Republic of the Congo (Congo/Brazzaville)
- 6: South Africa
- 7: Tanzania
- 8: Uganda
- 9: Zambia
- 10: Zimbabwe
- 11: other African country
- 12: Somalia
- 13: Sierra Leone
- 14: Angola
- 99: can't remember/not answered
- 1: not applicable *if whblack#2 & whblack#9*

Note: Variable 'afrcoun2' has codes used in the core questionnaire for both the core and the ethnic boost datasets.

(R) (ND) **afrcoun2** country of origin of black African respondents, grouped

- 1: Democratic Republic of Congo (Zaire) *if afrcoun=1*
- 2: Ghana *if afrcoun=2*
- 3: Kenya *if afrcoun=3*
- 4: Nigeria *if afrcoun=4*
- 5: South Africa *if afrcoun=6*
- 6: Uganda *if afrcoun=8*
- 7: Zambia *if afrcoun=9*
- 8: Zimbabwe *if afrcoun=10*
- 9: other African country *if afrcoun=5 | afrcoun=7 | afrcoun=11 | afrcoun=12 | afrcoun=13 | afrcoun=14*
- 99: not answered *if afrcoun=99*
- 1: not applicable *if afrcoun=-1*

ETHNICITY

Note: Variable 'whacar' is only available in the ethnic boost questionnaire

(R) (N) **whacar** country/island of origin for Caribbean respondents

- 1: Bahama Islands
- 2: Bermuda
- 3: Cuba
- 4: Dominican Republic
- 5: Jamaica
- 6: Haiti
- 7: Antigua, Monserrat & other Leeward Islands
- 8: Puerto Rico
- 9: Trinidad and Tobago
- 10: Barbados, and other Windwards Islands
- 11: other
- 12: Guyana
- 1: not applicable *if whmixed#1 & whblack#1*
- 99: can't remember/not answered *else if missing*
- 9999: not asked in Natsal-2 core questionnaire

Note: Variable 'ethnic2' is the most detailed variable on self-reported ethnicity

(R) (ND) **ethnic2** ethnic group that the respondent considers themselves belonging to

- 1: white, British *if ethnic=1 & whwhite=1*
- 2: white, Irish *if ethnic=1 & whwhite=2*
- 3: white, other background *if ethnic=1 & whwhite=3*
- 4: mixed ethnic group, white British & black Caribbean *if ethnic=2 & whmixed=1*
- 5: mixed ethnic group, white British & black African *if ethnic=2 & whmixed=2*
- 6: mixed ethnic group, white British & Asian *if ethnic=2 & whmixed=3*
- 7: mixed ethnic group, other mixed background *if ethnic=2 & whmixed=4*
- 8: Asian/Asian British, Indian *if ethnic=3 & whasian=1*
- 9: Asian/Asian British, Pakistani *if ethnic=3 & whasian=2*
- 10: Asian/Asian British, Bangladeshi *if ethnic=3 & whasian=3*
- 11: Asian/Asian British, other Asian background *if ethnic=3 & whasian=4*
- 12: black/black British, Caribbean *if ethnic=3 & whblack=1*
- 13: black/black British, African *if ethnic=3 & whblack=2*
- 14: black/black British, other black background *if ethnic=3 & whblack=3*
- 15: Chinese *if ethnic=5*
- 16: other ethnic group *if ethnic=6*
- 17: Middle Eastern *if ethnic=7*
- 99: not answered *if ethnic=9*

(R) (ND) **ethnic3** ethnic group respondent considers themselves belonging to, grouped

- 1: white *if ethnic2>=1 & ethnic2<=3*
- 2: mixed ethnic group *if ethnic2>=4 & ethnic2<=7*
- 3: Asian *if ethnic2>=8 & ethnic2<=11*
- 4: black African *if ethnic2=13*
- 5: black Caribbean *if ethnic2=12*
- 6: other ethnic group (incl. Chinese and Middle Eastern) *if (ethnic2>=14 & ethnic2<=17) | whasian=4*
- 9: not answered *if ethnic2=99*

(R) (ND) **ethnic4** ethnic group respondent considers themselves belonging to, grouped

- 1: white *if ethnic=1*
- 2: black *if ethnic=4*
- 3: Asian *if ethnic=3*
- 4: other *if ethnic=2 | ethnic=5 | ethnic=6 | ethnic=7*
- 9: not answered *if ethnic=9*

ETHNICITY

(R) (ND) **ethnic5** ethnic group respondent considers themselves belonging to, grouped

- 0: white *if* ethnic=1
- 1: else *if* ethnic≠1 & ethnic≠9
- 9: not answered *if* ethnic=9

(R) (ND) **ethnic6** ethnic group respondent considers themselves belonging to, grouped

- 1: white *if* ethnic3=1
- 2: black Caribbean *if* ethnic3=5 | (ethnic2=14 & ebornin=6) | (whacar≠-1 & whacar≠99)
- 3: black African *if* ethnic2=4 | (ethnic2=14 & ebornin>=10 & ebornin<=13) | (afrcoun≠-1 & afrcoun≠99)
- 4: Indian *if* ethnic2=8
- 5: Pakistani *if* ethnic2=9
- 6: other *else if* ethnic2≠99
- 9: not answered *else*

(R) (ND) **ethnic7** ethnic group respondent considers themselves belonging to, grouped

- 1: white *if* ethnic3=1
- 2: black Caribbean *if* ethnic3=5
- 3: black other *if* ethnic2=14
- 4: black African *if* ethnic3=4
- 5: Indian *if* ethnic2=8
- 6: Pakistani *if* ethnic2=9
- 7: other *else if* ethnic2≠99
- 9: not answered *else*

Notes:

1. Variable 'ethnic8' was derived by National Centre for Social Research to assign those in the ethnic boost sample to one of the four specific ethnic groups.
2. If a respondent stated that they were e.g. half Indian and half Black African, the individual case was studied, and on the basis of place of birth, verbatim description of 'other' ethnicity, religion, languages spoken, name etc. NatCen allocated them to just one group.
3. This variable is also derived for the Core dataset.
4. Serial numbers beginning 5.... refer to cases in the ethnic boost; all others are cases in the core sample
5. See the Natsal-2 codebook for full details of the coding of ethnic8 and ethnic9

(R) (ND) **ethnic8** derived ethnic group

- 1: Black Caribbean
- 2: Black African
- 3: Indian
- 4: Pakistani
- 5: white
- 6: other or ethnic group not clear
- 9: refused

Note: Variable 'ethnic9' was derived by the National Centre for Social Research to make the assumptions used in deriving 'Ethnic8' explicit.

(R) (ND) **ethnic9** derived ethnic group with assumptions made explicit

- 1: Black Caribbean
- 3: Black African
- 4: recoded as Black African
- 5: Indian *if*
- 6: recoded as Indian
- 7: Pakistani
- 8: recoded as Pakistani
- 9: white
- 10: recoded as white
- 11: other, or ethnic group not clear

ETHNICITY

Note: Variable 'ethnic10' is similar to 'ethnic 8' but categorises white respondents, 'other' and 'not answered' into just one category

(R) (ND) **ethnic10** derived ethnic group

- 1: black African *if* ethnic8=1
- 2: black Caribbean *if* ethnic8=2
- 3: Indian *if* ethnic8=3
- 4: Pakistani *if* ethnic8=4
- 5: else (predominantly white but also 'other' and 'not answered') *else*

Note: Variable 'ethnic11' was derived to be consistent with Natsal-1's ethnicity variable

(R) (c17 D) **ethnic11** ethnic group respondent considers themselves belonging to, grouped

- 1: black *if* ethnic2=12 | ethnic2=13 | ethnic2=14
- 2: white *if* ethnic2=1 | ethnic2=2 | ethnic2=3
- 3: Indian *if* ethnic2=8
- 4: Pakistani *if* ethnic2=9
- 5: Bangladeshi *if* ethnic2=10
- 6: Chinese *if* ethnic2=15
- 7: other Asian *if* ethnic2=11
- 8: other *if* ethnic2=4 | ethnic2=5 | ethnic2=6 | ethnic2=7 | ethnic2=16 | ethnic2=17
- 9: not answered *if* ethnic2=99

Note: Variables 'pethnic'... 'pwhblack' are derived for the respondent's partner. The same codes are used as for the respondent in addition to: -1 if marstat>=4 & marstat<=7

(R) (N) **pethnic** ethnic group respondent considers partner belongs to, grouped

(R) (N) **pwhwhite** cultural background of white partners

(R) (N) **pwhmixed** cultural background of mixed ethnic group partners

(R) (N) **pwhasian** cultural background of british asian partners

(R) (N) **pwhblack** cultural background of british african partners

Note: Variables 'pethnic2'... 'pethnic7' are derived for the respondent's partner. The same codes are used as for the respondent in addition to: -1 if marstat>=4 & marstat<=7

(R) (ND) **pethnic2** ethnic group respondent considers partner belongs to

(R) (ND) **pethnic3** ethnic group respondent considers partner belongs to, grouped

(R) (ND) **pethnic4** ethnic group respondent considers partner belongs to, grouped

(R) (ND) **pethnic5** ethnic group respondent considers partner belongs to, grouped

(R) (ND) **pethnic6** ethnic group respondent considers partner belongs to, grouped

(R) (ND) **pethnic7** ethnic group respondent considers partner belongs to, grouped

SEXUAL IDENTITY

(N) **sexid** sexual identity

- 1: heterosexual / straight
- 2: gay / lesbian
- 3: bisexual
- 4: other
- 9: not answered

RELIGION

RELIGION

(q12d) **religimp** importance of religion and religious beliefs now

- 1: very important
- 2: fairly important
- 3: not very important
- 4: not important at all
- 9: not answered

(N) **belrelig** belong to any religion now

- 1: yes
- 2: no, none
- 9: not answered

Note: codes 16-19 were removed from whirel in Natsal-3 because the numbers were insufficient to be useful for analysis. Instead, these respondents remain coded as 'other non-christian' (the response option selected by the respondent) .

(q13ab) **whirel** which religion belongs to

- 1: Christian, no denomination
- 2: Roman Catholic
- 3: Church of England/Anglican
- 4: United Reformed Church
- 5: Congregational
- 6: Baptist
- 7: Methodist
- 8: Presbyterian/Church of Scotland
- 9: other Christian
- 10: Hindu
- 11: Jew
- 12: Islam/Muslim
- 13: Sikh
- 14: Buddhist
- 15: other non-Christian
- (U – code removed) 16: Jehovah's Witness
- (U – code removed) 17: 7th Day Adventist
- (U – code removed) 18: Rastafarian
- (U – code removed) 19: Pentecostal
- 99: not answered
- 1: not applicable *if* belrelig=2

(ND) **religion** religion, grouped

- 1: none *if* belrelig=2
- 2: Christian, Church of England/Anglican *if* whirel=3
- 3: Christian, Roman Catholic *if* whirel=2
- (U) 4: Christian, other *if* whirel=1 | (whirel>=4 & whirel<=9)
- 4: Christian, other *if* whirel=1 | (whirel>=4 & whirel<=9) | whirel=16 | whirel=17 | whirel=19
- (U) 5: non-Christian *if* (whirel>=10 & whirel<=15)
- 5: non-Christian *if* (whirel>=10 & whirel<=15) | whirel=18
- 9: not answered *if* missing

RELIGION

(ND) **religio2** religion, grouped

- 1: none *if* belrelig=2
- 2: Christian, Church of England/Anglican *if* whirel=3
- 3: Roman Catholic *if* whirel=2
- 4: (U) Christian, other *if* whirel=1 | (whirel>=4 & whirel<=9)
- 4: Christian, other *if* whirel=1 | (whirel>=4 & whirel<=9) | whirel=16 | whirel=17 | whirel=19
- 5: Muslim *if* whirel=12
- 6: Hindu *if* whirel=10
- (U) 7: non-Christian *if* whirel=11 | whirel=13 | whirel=14 | whirel=15
- 7: non-Christian *if* whirel=11 | whirel=13 | whirel=14 | whirel=15 | whirel=18
- 9: not answered *else if* missing

(q13c) **oftrelig** frequency of attendance at religious services/meetings

- 1: once a week or more
- 2: less often but at least once in 2 weeks
- 3: less often but at least once a month
- 4: less often but at least twice a year
- 5: less often but at least once a year
- 6: less often
- 7: never or practically never
- 8: varies
- 9: not answered
- 1: not applicable *if* belrelig=2

INTERNET ACCESS

(N) **netacc** access to internet at home

- 1: yes
- 2: no
- 9: not answered

(N) **intagain** respondent agreed to recontact for future studies

- 1: yes
- 2: no
- 1: not applicable *if* partial interview

(N) **email** respondent gave email address for recontact

- 1: yes
- 2: no
- 1: not applicable *if* intagain#1
- 3: not asked – new question (*question introduced shortly after the start of fieldwork wave 3*)

URINE SURVEY

URINE SURVEY

Note: urine samples were collected from a sub-set of respondents in Natsal-2 and Natsal-3. The variables have different names in the two surveys due to differences in eligibility criteria, and tests carried out. Natsal-3 variables are shown first in blue below, then Natsal-2 variables.

Note: selection for the urine sample in Natsal-3 was as follows:

- All 16-17 year olds
- All 18-24 year olds who reported 1+ sexual partners, ever
- A random selection of 25-34 year olds who reported 1+ sexual partners, ever
- All men aged 16-44 with same-sex partners in the 5 years before interview

Note: The urine test result variables (ct_posconfirmed..hiv_positive) and urine weights (urine_wt) are available on a separate anonymised dataset, which can be accessed by application to the UK data archive.

(N) **urselect** respondent selected for urine sample

1: respondent selected

2: respondent not selected

(N) **urinuse** useable urine sample

1: yes – recorded in CAPI, consent form received, sample received

2: yes – sample collected in error, but respondent eligible, consent form received, sample received

3: no – problem with consent form, sample ID, or sample not received, or leaked

4: no – sample collected in error, respondent not eligible

-1: not applicable if urinsamp=2 | urinsamp=-1

Note: Additional weights were created for analysis of the urine test results. These combine the survey weights (total_wt) with additional weights which account for differential selection and non-response to the urine sample. All analysis of the urine data should use these weights. For more information about how these urine weights were created see the Natsal-3 technical report: www.natsal.ac.uk/natsal-3/methodology.

(N) **urine_wt** weight for analysis of the urine data

(N) **ct_posconfirmed** chlamydia test results from urine sample

1: chlamydia detected

0: chlamydia not detected

-1: not applicable

Note: gc_posconfirmed was subsequently replaced by gc_pos2 after an additional (2nd) confirmatory request identified an additional positive case.

(N) **gc_posconfirmed** gonorrhoea test results from urine sample

1: gonorrhoea detected

0: gonorrhoea not detected

-1: not applicable

(N) **gc_pos2** gonorrhoea test results from urine sample

1: gonorrhoea detected

0: gonorrhoea not detected

-1: not applicable

(N) **mg_posconfirmed** *Mycoplasma genitalium* test results from urine sample

1: M.gen detected

0: M. gen not detected

-1: not applicable

URINE SURVEY

Note: Stored Natsal-2 urine samples were tested for HPV. These results are not on the archived Natsal-2 dataset, but are available on request.

(N) **hvp_positive** HPV test results from urine sample

1: HPV (any type) detected

0: No HPV detected

2: Equivocal / indeterminate

-1: not applicable

(N) **HPV_type** HPV type detected in urine sample

(N) **HPV_6** HPV 6 result from urine sample

1: HPV type 6 detected

0: HPV type 6 not detected

-1: not applicable

Variables HPV_11..HPV_82 are all coded similarly to HPV_6

(N) **HPV_11** HPV 11 result from urine sample

(N) **HPV_16** HPV 16 result from urine sample

(N) **HPV_18** HPV 18 result from urine sample

(N) **HPV_26** HPV 26 result from urine sample

(N) **HPV_31** HPV 31 result from urine sample

(N) **HPV_33** HPV 33 result from urine sample

(N) **HPV_35** HPV 35 result from urine sample

(N) **HPV_39** HPV 39 result from urine sample

(N) **HPV_45** HPV 45 result from urine sample

(N) **HPV_51** HPV 51 result from urine sample

(N) **HPV_52** HPV 52 result from urine sample

(N) **HPV_53** HPV 53 result from urine sample

(N) **HPV_56** HPV 56 result from urine sample

(N) **HPV_58** HPV 58 result from urine sample

(N) **HPV_59** HPV 59 result from urine sample

(N) **HPV_66** HPV 66 result from urine sample

(N) **HPV_68** HPV 68 result from urine sample

(N) **HPV_70** HPV 70 result from urine sample

(N) **HPV_73** HPV 73 result from urine sample

(N) **HPV_82** HPV 82 result from urine sample

Note: types coded as 'low risk' (HPV_LR) were: 6, 11, 26, 53, 66, 70, 73, 82

Types coded as 'high risk' (HPV_HR) were: 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 68

(N) **HPV_LR** 1+ low risk HPV strain detected in urine sample

1: LR-HPV detected

0: LR-HPV not detected

-1: not applicable

(N) **HPV_HR** 1+ high risk HPV strain detected in urine sample

1: HR-HPV detected

0: HR-HPV not detected

-1: not applicable

(N) **hiv_positive** HIV result from urine sample

1: HIV detected

0: HIV not detected

-1: not applicable

(N) **tv_posconfirmed** *Trichomonas vaginalis* test result from urine sample

1: TV detected

0: TV not detected

-1: not applicable

URINE SURVEY

9: indeterminate/equivocal

Note: Respondents were only eligible for the urine survey if (1) their address is a urine sample point; (2) they were aged over 18 and have had some sexual experience

(R) (N) **urefcode** reason for refusing to provide a urine sample

- 1: no reason for test (e.g. with present life-style)
- 2: worried about confidentiality
- 3: worried about effect on insurance
- 4: already been tested for chlamydia
- 5: on medication/health problems
- 6: menstruating
- 7: felt unable to pass urine
- 8: did not want to know the test result
- 9: no time
- 10: should not be part of the survey
- 11: would not participate in this kind of test
- 12: too personal/embarrassed
- 13: no reason given (incl. just didn't want to)
- 14: other reason given
- 99: not answered
- 1: not applicable *if* not eligible for urine study or agreed to provide a sample

(R) (N) **urinsamp** urine sample provided

- 1: yesc
- 2: no
- 1: not applicable *if* not eligible for urine study or did not agree to provide a sample

(R) (N) **ct_lab** chlamydia test results from lab

- 8: no test done/no result
- 7: other text
- 6: LCR equivocal
- 1: not applicable *if* urinsamp=2 | urinsamp=-1
- 1: no chlamydia detected
- 2: chlamydia detected

(R) (N) **c_result** chlamydia test result from urine sample

- 0: no chlamydia detected
- 1: chlamydia detected
- 9: chlamydia result not known/not eligible for urine study *else if missing*

Note: The original urine weighting variable for Natsal-2 was 'urine_wt'. This variable was updated (in 2014) and the correct variable to use for any subsequent analysis is 'newurine_wt'. The changes are as follows:

- *newurine_wt3 uses the updated survey weights newfin_wt, rather than final_wt*
- *newurine_wt3 non-response weighting model was created using more up-to-date methodology (in terms of the recoding of small categories, building a separate model for men and women, and the trimming and scaling of the weights).*

This may result in small changes to the prevalence estimates, compared with those published in the original Natsal-2 papers.

Note: Variable 'Urine_Wt' should be used as the pweight for analysis in Stata of the urine survey data using the survey commands

(use newurine_wt3 instead – see note) (N) **urine_wt** weight for analysis of the urine data

- 1: not applicable *if* not eligible for urine study
- 0: eligible for urine study but refused to provide a urine sample/result not known
- >0: eligible for urine study, provided a urine sample and result known

URINE SURVEY

Note: when analysing change over time between Natsal-2 and Natsal-3 urine data, care should be taken to ensure the denominator is the same for both surveys. The Natsal team typically restrict analysis to those reporting at least one sexual partner ever: ((hetlife>0 & hetlife<9999) | (samlife>0 & samlife<9999)) in Natsal-3 and ((hetlife>0 & hetlife<9999) | (homlife>0 & homlife<9999)). This excludes a small number of cases in Natsal-2 with chlamydia / HPV results but no sexual partners ever.

Note: Stored urine samples from Natsal-2 were subsequently tested for Human Papillomavirus (HPV). The HPV data for Natsal-2 are available on request from Cath Mercer (contact details on front page).

SALIVA SURVEY

SALIVA SURVEY

Note: in Natsal-3 a sub-sample of respondents aged 18-74 were also eligible for a saliva sample to be tested for testosterone. Selection was as follows:

- *A random selection of 18-74 year olds who did not report working mainly night shifts (nights#1)*
- *From fieldwork wave 4 onwards all 35-74 year olds who did not report working mainly night shifts.*

The saliva sample data can be accessed by application to the UK data archive.

Note: see the Natsal technical report (<http://www.natsal.ac.uk/natsal-3/methodology.aspx>) for full details of saliva samples collected, and numbers excluded because they were unusable. The variable 'saliva' indicates all samples with a valid, useable, testosterone result.

(N) **saliva** valid testosterone result

1 yes

else missing

Note: Additional weights were created for analysis of the saliva test results. These combine the survey weights (total_wt) with additional weights which account for differential selection and non-response to the saliva sample. All analysis of the saliva data should use these weights. For more information about how these weights were created see the Natsal-3 technical report: www.natsal.ac.uk/natsal-3/methodology.

(N) **saliva_wt** weight for use in saliva analysis

(N) **testo** salivary testosterone level (pmol/litre)

999993: lab error (not tested)

999994: <6.5

999995: insufficient sample

999996: na/unsuitable

999997: no/unknown consent

999998: chromatography problem

999999: no sample

INTERVIEW CHARACTERISTICS

INTERVIEW CHARACTERISTICS

Note: in Natsal-3 we included data for some 'partial' interviews i.e. where respondents had stopped the interview before the last question. Partial interviews were only included if the respondent had answered at least up to expscale. See the Natsal-3 technical report for further details.

(N) **outcome** interview outcome

110: full interview

210: partial interview

Note: Variables langi01...langi06 are only available in the ethnic boost survey.

(R) (N) **langi01** languages spoken by the respondent 1

1: English

2: Gujarati

3: Hindi

4: Punjabi

5: Urdu

6: Bengali

7: Sylheti

8: Patois/Creole

9: French

10: other

99: not answered

-1: not applicable *if* in core sample/no other languages spoken

Similarly for up to 6 languages:

(R) (N) **langi02** languages spoken by the respondent 2

(R) (N) **langi03** languages spoken by the respondent 3

(R) (N) **langi04** languages spoken by the respondent 4

(R) (N) **langi05** languages spoken by the respondent 5

(R) (N) **langi06** languages spoken by the respondent 6

Note: Variables clangi1...clangi3 are only available in the ethnic boost survey.

(R) (N) **clangi1** other languages spoken by the respondent 1

1: English

2: Gujarati

3: Hindi

4: Punjabi

5: Urdu

6: Bengali

7: Sylheti

8: Patois/Creole

9: French

10: (code not used)

11: Yoruba

12: Twi/Ga

13: Swahili

14: Somali

15: Arabic

16: other European language

17: other African language (e.g. Ibo, Shona)

18: other Asian language (e.g. Hinko, Canton)

99: not answered

-1: not applicable *if* in core sample/no other languages spoken

Similarly for up to 3 other languages:

(R) (N) **clangi2** other languages spoken by the respondent 2

(R) (N) **clangi3** other languages spoken by the respondent 3

INTERVIEW CHARACTERISTICS

Note: Variable LangM is only available in the ethnic boost surv.

(R) (N) **langm** main language spoken

- 1: English
- 2: Gujarati
- 3: Hindu
- 4: Punjabi
- 5: Urdu
- 6: Bengali
- 7: Sylheti
- 8: Patois/Creole
- 9: French
- 10: other
- 99: not answered
- 1: not applicable *if* in core sample/respondent only speaks one language

Notes: 1. Variable 'intlang' is only available in the ethnic boost survey

2. Only respondents reporting their ethnicity as Indian or Pakistani were asked 'intlang'.

(R) (N) **intlang** proportion of interview in English

- 1: wholly in English
- 2: partly in English, partly in another language
- 3: wholly in another language
- 1: not applicable *if* in core sample/(whasian#1 & whasian#2)

Note: Variable 'ilangwh' is only available in the ethnic boost survey.

(R) (N) **ilangwh** other language used for interview

- 1: Punjabi
- 2: Urdu
- 9: not answered
- 1: not applicable *if* in core sample/(intlang=2 | intlang=2)

(c20a) **others** anyone else present in home at all during interview

- 1: yes
- 2: no
- 3: interview conducted elsewhere (e.g. garage, car)
- 9: not answered

(c20b) **whothere** anyone else present during any part of interview & (possible) able to overhear

- 1: yes
- 2: no
- 9: not answered
- 1: not applicable *if* others=2 | others=3

(c20c1) (d63) **homespart** spouse/partner present for some/all of the interview

- 1: present throughout
- 2: present some of the time
- 3: may have overheard all/part
- 4: passing through only
- 5: not present (incl. not applicable)
- 9: not answered
- 1: not applicable *if* (others=2 | others=3) | whothere=2

Note: Variable d63 codes apply to the following variables d64 ... d68:

(c20c2) (d64) **homepars** parent(s) present for some/all of the interview

(c20c3) (d65) **home0to5** child(ren) aged 0-5 years present for some/all of the interview

(c20c4) (d66) **home6to15** child(ren) aged 6-15 years present for some/all of the interview

(c20c5) (d67) **home16to21** young adult(s) aged 16-21 years present for some/all of the interview

(c20c6) (d68) **home22** other adult(s) aged 22+ years present for some/all of the interview

INTERVIEW CHARACTERISTICS

(c20d) **scseen** anyone else in household look at/discuss any of CASI

1: yes, looked at/read/completed together

2: yes, discussed only

3: no

4: questionnaire not completed

9: not answered

-1: not applicable *if* (others=2 | others=3) | whothere=2

(c21a1) **langprob** difficulties with interview because of language problems

1: yes, severe

2: yes, some

3: no problem

9: not answered

Note: Variable langprob codes apply to the following variables litprob and cogprob:

(c21a2) **litprob** difficulties with interview because of literacy problems

(N) **cogprob** difficulties with interview because of cognitive problems

(c21b) **readcard** interviewer read out show cards

1: yes, all/most

2: yes, some

3: no

9: not answered

(c21c) **embarras** interviewer's opinion of how embarrassed the respondent was

1: very embarrassed/ill at ease

2: somewhat embarrassed/ill at ease

3: only slightly embarrassed/ill at ease

4: not at all embarrassed/ill at ease

9: not answered

(R) (N) **difficlt** interviewer's opinion of how difficult it was to persuade the respondent to participate

1: very difficult

2: fairly difficult

3: not very difficult

4: not at all difficult

9: not answered

(N) **willing** respondent's initial willingness to participate in CASI

1: very willing

2: fairly willing

3: not at all willing

9: not answered

-1: not applicable *if* scelig=2

(R) (N) **areatype** interviewer's perception of area-type

1: urban/city centre

2: small country town centre

3: suburban residential

4: rural residential/village centre

5: rural agricultural with isolated dwellings/small hamlets

9: not answered

INTERVIEW CHARACTERISTICS

(R) (N) **bldtype** interviewer's perception of residential building type

- 1: terraced houses
- 2: semi-detached houses
- 3: detached houses
- 4: mixed housing
- 5: low rise flats (5 storeys or less)
- 6: high rise flats (blocks over 5 storeys)
- 7: flats with commercial (flats/maisonettes over shops)
- 8: flats –mixed (high and low rise)
- 9: mixed houses and flats
- 99: not answered

(R) (N) **typdwell** interviewer's perception of household dwelling type

- 1: whole house/bungalow: detached
- 2: whole house/bungalow: semi-detached
- 3: whole house/bungalow: terraced
- 4: purpose built flat/maisonette: basement
- 5: purpose built flat/maisonette: 4th floor
- 6: converted flat/maisonette: part-house/rooms in house
- 7: dwelling with business premises
- 8: caravan/houseboat
- 9: other
- 99 not answered

(R) (N) **ethmix** ethnic mix area

- 1: predominantly white
- 2: predominantly black/brown
- 3: mixed
- 9: not answered

(N) **adults** number of people in the household eligible for Natsal-2

(U) 9995: at least one

(R) (N) **nofdu** number of occupied dwelling units

99: missing

(R) (U) **stdreg** standard region

- 1: North
- 2: Yorkshire & Humberside
- 3: East Midlands
- 4: East Anglia
- 5: South East (excl. London)
- 6: South West
- 7: West Midlands
- 8: North West
- 9: Wales
- 10: Scotland
- 11: Greater London

Notes: 1. There are some discrepancies between the coding of stdreg and gor as counties are distributed slightly differently between them. For example, 'North East' under gor includes Cleveland, Durham, Northumberland and Tyne and Wear, while the stdreg 'North' includes Cleveland, Durham, Northumberland, Tyne and Wear, and Cumbria.
2. stdreg is the older classification and it was used in Natsal-1 but gor is the more widely used now. Only gor is available for the Natsal-3 data
3. Data for gor was obtained from the Office for National Statistics and originally included the region 'Merseyside' but this region was merged with 'North West' since 'Merseyside' no longer exists as a region, thus there is no code 3 for gor.

INTERVIEW CHARACTERISTICS

(D) **gor** Government Office Region

- 1: North East
- 2: North West (incl. the old 'Merseyside' region)
- 4: Yorkshire & Humber
- 5: East Midlands
- 6: West Midlands
- 7: South West
- 8: Eastern
- 9: London
- 10: South East
- 11: Wales
- 12: Scotland

(ND) **gor2** Government Office Region (grouped)

- 1: England *if gor≠11 & gor≠12*
- 2: Scotland *if gor=12*
- 3: Wales *if gor=11*

(N) **gor_1** Government Office Region with inner & outer London specified

- 1: North East
- 2: North West (incl. the old 'Merseyside' region)
- 4: Yorkshire & Humber
- 5: East Midlands
- 6: West Midlands
- 7: South West
- 8: Eastern
- 9: Inner London
- 10: Outer London
- 11: South East
- 12: Wales
- 13: Scotland

(ND) **london** respondent lives in Greater London (as defined by 'gor')

- 0: elsewhere *if gor≠9*
- 1: Greater London *if gor=9*

(N) **npopdens** local population density for those living in England – quintiles

1=least dense 5=most dense

Note: snpopdens and wnpopdens are coded similarly to npopdens

(N) **snpopdens** local population density for those living in Scotland – quintiles

(N) **wnpopdens** local population density for those living in Wales – quintiles

Notes:

1. *Natsal-3 data includes a combined Index of Multiple Deprivation (IMD) quintile variable for England, Scotland and Wales. (This replaces the variable imdgbq from Natsal-2.) This is based on an adjusted IMD score which is designed to be broadly comparable across nations, unlike the individual country IMD scores which are not directly comparable.*
2. *The adjusted IMD score was generated using a method by Payne and Abel (2012) which is described in:
Payne and Abel (2012) UK indices of multiple deprivation – a way to make comparisons across constituent countries easier, Health Statistics Quarterly (53): 22-37
It is based on the income and employment of the individual country IMD scores combined with the co-efficients and residual values from a linear regression of income and employment on the overall IMD score for each country.*
3. *The combined scores were generated using the most up-to-date scores for each country at the time. These were IMD 2010 for England, IMD 2011 for Wales, and IMD 2009 for Scotland.*
4. *The quintiles for the individual countries are also available (qimd, qwimd, qsimd) for those conducting analysis on just one country.*

INTERVIEW CHARACTERISTICS

(N) **adj_imd_quintile** adjusted IMD quintiles for England, Scotland and Wales combined
1=least deprived 5=most deprived.

(N) **qimd** IMD quintile for England (2010)
1=least deprived 5=most deprived

(N) **qwind** IMD quintile for Wales (2011)
1=least deprived 5=most deprived

(N) **qsimd** IMD quintile for Scotland (2009)
1=least deprived 5=most deprived

Notes: 1. As at 23 July 2003, a PDF report on how the deprivation index is derived can be downloaded from the website:
http://www.odpm.gov.uk/stellent/groups/odpm_urbanpolicy/documents/sectionhomepage/odpm_urbanpolicy_page.hcsp
2. **imdgbq** is not available for the Scottish respondents in the Ethnic Boost dataset (n=26/949)

(R) (ND) **imdgbq** Index of Multiple Deprivation, quintiles
1: first quintile (least deprived)
2: second quintile
3: third quintile
4: fourth quintile
5: fifth quintile (most deprived)
9: not available

Notes: 1. In comparison to 'imdgbq', the codes for 'jarmenq' are the opposite way round, i.e. the first quintile is the most deprived for 'jarmenq' unlike 'imdgbq'
2. Jarmen quintiles were not available for the Scottish respondents.

(R) (ND) **jarmenq** Jarmen index, quintiles
1: first quintile (most deprived)
2: second quintile
3: third quintile
4: fourth quintile
5: fifth quintile (least deprived)
9: not available if gor=12

Note: the Natsal-3 data include the 2011 national statistics Output Area Classification codes. For information about how to use these see: <http://www.ons.gov.uk/ons/guide-method/geography/products/area-classifications/ns-area-classifications/index/cluster-summaries/output-areas/index.html>

(N) **oac2011** ONS output area classification (2011 version)

A previous version of the data also included the 2001 version of the OAC ('oac_old'), for more information about the 2001 version see: <http://www.ons.gov.uk/ons/guide-method/geography/products/area-classifications/ns-area-classifications/index/index.html>

(N) **oac_old** ONS output area classification (2001 version)

Note: The Natsal-3 data include ONS urban/rural indicator variables. For more information about these, see the ONS website for England & Wales: <http://www.ons.gov.uk/ons/guide-method/geography/products/area-classifications/rural-urban-definition-and-la/rural-urban-definition-england-and-wales-/index.html>;
or the Scottish Executive website for Scotland:
<http://www.scotland.gov.uk/Topics/Statistics/About/Methodology/UrbanRuralClassification>

(N) **urindew** urban / rural indicator (England & Wales)
1: urban >=10k - sparse
2: Town & Fringe - sparse

INTERVIEW CHARACTERISTICS

- 3: Village - sparse
- 4: Hamlet and isolated dwelling- sparse
- 5: urban>=10k – less sparse
- 6: Town & Fringe – less sparse
- 7: Village – less sparse
- 8: Hamlet & isolated dwelling – less sparse
- 9: In Scotland

(N) **urinds** urban/rural indicator (Scotland)

- 1: large urban area (125, 000+)
- 2: Other urban area (10,000-125,000)
- 3: Accessible small town (3,000-10,000)
- 4: Remote small town (3,000-10,000)
- 5: Very remote small town (3,000-10,000)
- 6: accessible rural (<3,000)
- 7: Remote rural (<3,000)
- 8: Very remote rural (<3,000)
- 9: In England / Wales

(R) (N) **acorn6** ACORN code, 6 categories

- 1: thriving
- 2: expanding
- 3: rising
- 4: settling
- 5: aspiring
- 6: striving
- 9: not available

(R) (N) **acorn17** ACORN code, 17 categories

- 1: wealthy achievers
- 2: affluent greys
- 3: prosperous pensioners
- 4: affluent executives
- 5: well-off workers
- 6: affluent urbanites
- 7: prosperous professionals
- 8: better-off executives
- 9: comfortable middle agers
- 10: skilled workers
- 11: new home owners
- 12: white collar workers
- 13: older people
- 14: council residents, better-off
- 15: council residents, unemployed
- 16: council residents, hardship
- 17: multi-ethnic, low income
- 99: not available

Note: Variable 'ons' is only available for respondents in the core dataset resident in England.

(R) (ND) **ons** Office for National Statistics area code

- 0: Wales/Scotland
- 1: rural –coast and country
- 2: rural –mixed urban and rural
- 3: prospering – growth areas
- 4: prospering –most prosperous
- 5: maturer – services and education
- 6: maturer – resort and retirement

INTERVIEW CHARACTERISTICS

- 7: urban centres –mixed areas
- 8: urban centres –manufacturing
- 9: mining and industrial –ports and industry
- 10: mining and industrial –coalfields
- 11: inner London
- 1: not applicable *if* ethnic boost dataset

Note: Variable 'onsshort' is only available for respondents in the core dataset resident in England.

(R) (ND) **onsshort** Office for National Statistics area code, grouped

- 0: Wales/Scotland
- 1: rural
- 2: prospering
- 3: maturer
- 4: urban centres
- 5: mining and industrial
- 6: inner London
- 1: not applicable *if* ethnic boost dataset