
Assignment/Problem Set 1

Heinrich Dinkel

ID: 1140339107

E-mail: heinrich.dinkel@sjtu.edu.com.cn

Notes

- Das geht noch einfacher. 2
- Es ist eine Drehmatrix welche sich einfach um den n fachen Winkel weiter dreht. 2

1 Exercise 1

1 I shorten the notation by using \sin as a replacement for $\sin(x)$ and \cos as a replacement for $\cos(x)$. We use the binet formula to get the power of A^n . We first estimate the eigenvectors, then concatenate them into matrix P and use this matrix to compute the diagonal matrix B , which can be used to directly take the power for the matrix A . Afterwards we need to reverse the operations, which we had done on achieving B to get A^n .

$$\begin{aligned} A^n &= \begin{pmatrix} \cos & \sin \\ -\sin & \cos \end{pmatrix}^n, v = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, (A - \lambda I)v = 0 \\ &\begin{pmatrix} \cos & \sin \\ -\sin & \cos \end{pmatrix} - \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} = \\ &(\cos^2 - \lambda) + \sin^2 = 0 \\ &\lambda_1 = i \sin + \cos \\ &\lambda_2 = -i \sin + \cos \end{aligned}$$

Now we achieved the eigenvalues, so we can estimate the eigenvectors. The eigenvector λ_1

$$\begin{pmatrix} \cos & \sin \\ -\sin & \cos \end{pmatrix} - \begin{pmatrix} i \sin + \cos & 0 \\ 0 & i \sin + \cos \end{pmatrix} = \begin{pmatrix} i & \\ & 1 \end{pmatrix}$$

Symmetrically the vector for λ_2 is:

$$\begin{pmatrix} -i \\ 1 \end{pmatrix}$$

The matrix P is just a concatenation of λ_1 and λ_2 .

$$P = \begin{pmatrix} i & -i \\ 1 & 1 \end{pmatrix} \tag{1}$$

To calculate an diagonal matrix, we use the equation $B = P^{-1}AP$ to transform A into a diagonal matrix.

$$\begin{aligned}
 P^{-1} &= \frac{1}{\det(P)} \text{adj}(P) = \\
 &= \frac{1}{2i} \begin{pmatrix} 1 & i \\ -1 & i \end{pmatrix} = \begin{pmatrix} \frac{1}{2i} & \frac{1}{2} \\ -\frac{1}{2i} & \frac{1}{2} \end{pmatrix} \\
 P^{-1}AP &= \begin{pmatrix} \frac{1}{2i} & \frac{1}{2} \\ -\frac{1}{2i} & \frac{1}{2} \end{pmatrix} \begin{pmatrix} \cos & \sin \\ -\sin & \cos \end{pmatrix} \begin{pmatrix} i & -i \\ 1 & 1 \end{pmatrix} = \\
 &= \begin{pmatrix} \left(\frac{\cos}{2i} - \frac{\sin}{2} \right) i + \frac{\sin}{2i} + \frac{\cos}{2} & \left(\frac{\cos}{2i} - \frac{\sin}{2} \right) (-i) + \frac{\sin}{2i} + \frac{\cos}{2} \\ \left(-\frac{\cos}{2i} - \frac{\sin}{2} \right) i + -\frac{\sin}{2i} + \frac{\cos}{2} & \left(-\frac{\cos}{2i} - \frac{\sin}{2} \right) (-i) - \frac{\sin}{2i} + \frac{\cos}{2} \end{pmatrix} = \\
 &= \begin{pmatrix} \cos - i \sin & 0 \\ 0 & \cos + i \sin \end{pmatrix}
 \end{aligned}$$

Now we can apply the power operation on the matrix.

$$B^n = \begin{pmatrix} (\cos - i \sin)^n & 0 \\ 0 & (\cos + i \sin)^n \end{pmatrix}$$

Now we can apply the matrix exponential onto A, since $A = PB^nP^{-1}$.

$$\begin{aligned}
 &\begin{pmatrix} i & -i \\ 1 & 1 \end{pmatrix} \begin{pmatrix} (\cos - i \sin)^n & 0 \\ 0 & (\cos + i \sin)^n \end{pmatrix} \begin{pmatrix} \frac{1}{2i} & \frac{1}{2} \\ -\frac{1}{2i} & \frac{1}{2} \end{pmatrix} = \\
 &\begin{pmatrix} (\cos - i \sin)^n i & (\cos + i \sin)^n (-i) \\ (\cos - i \sin)^n & (\cos + i \sin)^n \end{pmatrix} \begin{pmatrix} \frac{1}{2i} & \frac{1}{2} \\ -\frac{1}{2i} & \frac{1}{2} \end{pmatrix} = \\
 &\frac{1}{2} \begin{pmatrix} (\cos - i \sin)^n + (\cos + i \sin)^n & (\cos - i \sin)^n i - (\cos + i \sin)^n i \\ i(\cos + i \sin)^n - i(\cos - i \sin)^n & (\cos - i \sin)^n + (\cos + i \sin)^n \end{pmatrix} = A^n
 \end{aligned}$$

Das geht noch einfacher.

$$A^n = \begin{pmatrix} \cos(n \cdot x) & \sin(n \cdot x) \\ -\sin(n \cdot x) & \cos(n \cdot x) \end{pmatrix} \quad (2)$$

Es ist eine Drehmatrix welche sich einfach um den n fachen Winkel weiter dreht.

2 Like in exercise 1 we begin by getting the eigenvalues and eigenvectors and the matrix P.

$$A = \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \text{ Eigenvalues of } A : \begin{pmatrix} 1 - \lambda & 1 \\ -1 & 1 - \lambda \end{pmatrix} = (1 - \lambda)^2 + 1 = 0$$

$$\lambda_1 = 1 + i$$

$$\lambda_2 = 1 - i$$

$$\text{Eigenvector of } \lambda_1, \lambda_2 \left[\begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} - \begin{pmatrix} 1 + i & 0 \\ 0 & 1 + i \end{pmatrix} \right] \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0$$

$$v_1 = \begin{pmatrix} 1 \\ i \end{pmatrix}$$

$$v_2 = \begin{pmatrix} 1 \\ -i \end{pmatrix}$$

Since we computed the eigenvectors v_1 and v_2 , we continue by calculating the diagonal matrix B .

$$\begin{aligned}
P &= \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix} \Rightarrow P^{-1} = \frac{1}{\det(P)} \begin{pmatrix} -i & -1 \\ -i & 1 \end{pmatrix} = \frac{1}{-2i} \begin{pmatrix} -i & -1 \\ -i & 1 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & \frac{1}{2i} \\ \frac{1}{2} & -\frac{1}{2i} \end{pmatrix} \\
B &= \begin{pmatrix} \frac{1}{2} & \frac{1}{2i} \\ \frac{1}{2} & -\frac{1}{2i} \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & \frac{1}{2i} \\ \frac{1}{2} & -\frac{1}{2i} \end{pmatrix} \begin{pmatrix} 1+i & 1-i \\ -1+i & -1-i \end{pmatrix} = \begin{pmatrix} 1+i & 0 \\ 0 & 1-i \end{pmatrix} \\
B^n &= \begin{pmatrix} (1+i)^n & 0 \\ 0 & (1-i)^n \end{pmatrix} \\
A^n &= PB^nP^{-1} = \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix} \begin{pmatrix} (1+i)^n & 0 \\ 0 & (1-i)^n \end{pmatrix} \begin{pmatrix} \frac{1}{2} & \frac{1}{2i} \\ \frac{1}{2} & -\frac{1}{2i} \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix} \begin{pmatrix} \frac{(1+i)^n}{2} & \frac{(1+i)^n}{-2i} \\ \frac{(1-i)^n}{2} & \frac{(1-i)^n}{-2i} \end{pmatrix} = \\
&= \frac{1}{2} \begin{pmatrix} (1+i)^n + (1-i)^n & (1+i)^n - (1-i)^n \\ i(1+i)^n - i(1-i)^n & (1+i)^n + (1-i)^n \end{pmatrix} = A^n
\end{aligned}$$

2 Exercise 2

1 Computing $A^{-1}B$.

$$\begin{aligned}
A^{-1} &= \frac{1}{\det(A)} \begin{pmatrix} 4 & 5 & 0 \\ 2 & 3 & 1 \\ 2 & 7 & -3 \end{pmatrix} = -\frac{1}{24} \begin{pmatrix} -16 & 15 & 5 \\ 8 & -12 & -4 \\ 8 & -18 & 2 \end{pmatrix} \\
AB &= \frac{1}{12} \begin{pmatrix} 12 & 0 & -30 & 95 \\ 0 & 12 & 24 & -52 \\ 0 & 0 & 0 & -46 \end{pmatrix}
\end{aligned}$$

2 Computing CA^{-1}

$$CA^{-1} = \begin{pmatrix} 4 & 5 & 0 \\ 2 & 3 & 1 \\ 2 & 7 & 9 \\ -2 & 3 & 7 \end{pmatrix} \left(-\frac{1}{24}\right) \begin{pmatrix} -16 & 15 & 5 \\ 8 & -12 & -4 \\ 8 & -18 & 2 \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ -12 & 27 & 0 \\ -14 & 24 & 1 \end{pmatrix}$$

3 Exercise 3

We want to proof the following equation:

$$\text{adj}(AB) = \text{adj}(B)\text{adj}(A) \quad (3)$$

It is already known that $A\text{adj}(A) = \det(A)I$, so

$$\text{adj}(A) = A^{-1}\det(A)I \quad (4)$$

Also it is known that

$$\det(AB) = \det(A)\det(B) \quad (5)$$

We substitute 4 into 3, by using 5.

$$\begin{aligned}
\text{adj}(B)\text{adj}(A) &= \det(B)B^{-1}I\det(A)A^{-1}I = \\
&= \det(A)\det(B)B^{-1}IA^{-1}I = \det(AB)(AB)^{-1}
\end{aligned}$$

Using $C = AB$ we get:

$$\det(C)C^{-1} = \text{adj}(C) \Rightarrow \text{adj}(AB) = \text{adj}(A)\text{adj}(B)$$

4 Exercise 4

Assuming having a matrix A with dimensions of $n \times m$.
We assume that there exists two ranks, $\text{rowrank}(A)$ and $\text{colrank}(A)$.

$$A = \begin{pmatrix} a_{11} & \dots & a_{n1} \\ \vdots & \ddots & \vdots \\ a_{1m} & \text{hdots} & a_{nm} \end{pmatrix}$$

If we modify A to A' by doing Gaussian elimination, A' will have exactly r non-zero rows, which will be denoted as $\text{rowrank}(A) = r$. By doing the transpose A'^T , we can still observe that the rowrank is r times non-zero rows. Since the columns in the transpose are the rows of the non-transpose, it leads to:

$$\text{rowrank}(A') = \text{rowrank}(A'^T) = \text{colrank}(A') = \text{colrank}(A'^T) = \text{rowrank}(A) = \text{colrank}(A)$$

I show that this behaviour is working as well for the square AA^T matrix. Since it is already proofed that the column rank is the row rank, we can use the basic formula for ranks: $\text{rank}(AB) \leq \min(\text{rank}(A), \text{rank}(B))$.

$$\text{rank}(AA^T) \leq \min(\text{rank}(A), \text{rank}(A^T)) \Rightarrow \text{rank}(A) = \text{rank}(A^T) = \text{rank}(AA^T) \quad (6)$$

5 Exercise 5

To proof $\text{rank}(A^n) = \text{rank}(A^{n+1})$ we use recursion.

$$\text{rank}(AB) \leq \min(\text{rank}(A), \text{rank}(B)) \quad (7)$$

By the use of 7 we get the following recursion:

$$\begin{aligned} \text{rank}(A^{n+1}) &= \\ \text{rank}(A^n A) &\leq \min(\text{rank}(A^n), \text{rank}(A)) \\ \text{rank}(A^n) &= \text{rank}(A^{n-1} A) \leq \min(\text{rank}(A^{n-1}), \text{rank}(A)) \\ &\vdots \\ \text{rank}(AA) &= \min(\text{rank}(A), \text{rank}(A)) = \text{rank}(A) \end{aligned}$$

So after this recursion, it can be seen that $\text{rank}(A^{n+1}) = \text{rank}(A^n)$