

Your Name: _____ netid: _____
 Name: _____ netid: _____
 Name: _____ netid: _____
 Name: _____ netid: _____
 Name: _____ netid: _____

Group #:

ECE 120 Worksheet 4: Karnaugh Maps

Before you begin today's discussion, be sure that you are familiar with terminology, such as literals, minterms, maxterms, canonical forms, implicants, prime implicants, and K-maps. Also be sure that you know how to find canonical SOP and POS forms for Boolean functions, and that you know how to use K-maps. To check these skills, you can make up a truth table at random, find a Boolean expression for the function, then check your result by writing a truth table for your expression.

K-maps

- Think about *Karnaugh map*, or *K-map*, as an alternative representation of truth table that
 - Lists cells in Gray code order, and
 - Each cell corresponds to a row of the truth table
- Two-variable Boolean function example:
 - four possible values, $m_0 - m_4$, which can be arranged into a Karnaugh map as follows:

Truth table for a 2-variable function

x	y	f(x,y)
0	0	m_0
0	1	m_1
1	0	m_2
1	1	m_3

Corresponding K-map representation

		y	
		0	1
x	0	m_0	m_1
	1	m_2	m_3
		f(x,y)	

- Similarly, we can obtain K-maps for 3- and 4-variable Boolean functions

		yz			
		00	01	11	10
x	0	m_0	m_1	m_3	m_2
	1	m_4	m_5	m_7	m_6
		f(x,y,z)			

		yz			
		00	01	11	10
wx	00	m_0	m_1	m_3	m_2
	01	m_4	m_5	m_7	m_6
	11	m_{12}	m_{13}	m_{15}	m_{14}
	10	m_8	m_9	m_{11}	m_{10}
		f(w,x,y,z)			

Problem 1

a) Fill in 3-variable K-maps for the functions shown below:

		yz			
		00	01	11	10
x	0	1	1	0	0
	1	1	1	0	0

$f(x,y,z)=y'$

		yz			
		00	01	11	10
x	0	0	0	0	0
	1	1	1	1	1

$f(x,y,z)=x$

		yz			
		00	01	11	10
x	0	0	1	0	0
	1	0	0	0	0

$f(x,y,z)=x'y'z$

		yz			
		00	01	11	10
x	0	0	0	0	1
	1	0	0	0	1

$f(x,y,z)=yz'$

		yz			
		00	01	11	10
x	0	0	0	0	0
	1	1	0	0	1

$f(x,y,z)=xz'$

		yz			
		00	01	11	10
x	0	0	1	0	1
	1	1	0	0	1

$f(x,y,z)=x'y'z+yz'+xz'$

b) Fill in 4-variable K-maps for the functions shown below:

		yz			
		00	01	11	10
wx	00	0	1	0	0
	01	1	0	1	1
	11	0	0	0	1
	10	0	1	0	0


$f(w,x,y,z)=w'xz'+x'y'z+w'xy+xyz'$

		yz			
		00	01	11	10
wx	00	1	1	0	1
	01	0	0	1	1
	11	0	0	1	1
	10	1	1	0	1


$f(w,x,y,z)=x'y'+yz'+xy$

Function simplification


- K-maps is a great tool for simplifying Boolean expressions
- A product term is an *implicant* of a function if the function has the value 1 for all minterms of the product term
 - In terms of K-map, implicants correspond to *all* legal loops
- An implicant is a *prime implicant* if it is not contained within a larger implicant
 - In terms of K-map, prime implicants correspond to *all biggest* loops
 - Example:



- An SOP (or POS) expression is *minimal* if
 - It has the minimum number of product (sum) terms, and
 - Among expressions with minimum number of terms, it has fewest literals
- A minimal SOP expression is a sum of prime implicants.
- Minimal SOP example:



Minimal POS example:



Use the area heuristic—number of literals plus the number of operators, not including complemented literals—to find minimal expressions in today's problems. Choosing a minimal number of prime implicants from a K-map will give you minimal solutions in this sense.

Problem 2

- a) Find all minimal SOP expressions from the following K-maps. Circle the corresponding loops on the K-maps and write down the Boolean expressions. Note that min SOP may not be unique!

		yz			
		00	01	11	10
wx	00	1	0	1	1
	01	1	0	0	1
	11	1	1	1	0
	10	0	0	1	1

Min SOP:

$$\underline{x'y + w'z'} + wxy' + wxz$$

min SOP is not unique,
there is another way to
cover m_{12}, m_{13}, m_{15} :

$$\underline{x'y + w'z'} + xy'z' + wxz$$

		yz			
		00	01	11	10
wx	00	0	1	0	0
	01	0	1	1	1
	11	1	1	1	0
	10	0	0	1	0

Min SOP:

$$wxy' + w'y'z + wyz + w'xy$$

(all essential PIs)

- b) Find minimal POS examples from the following K-maps. Circle the corresponding loops on the K-maps and write down the Boolean expressions:

		yz			
		00	01	11	10
wx	00	1	1	0	0
	01	1	1	1	1
	11	1	1	0	1
	10	1	1	0	0

Min POS:

$$\underline{(w' + y' + z')(x + y')}$$

		yz			
		00	01	11	10
wx	00	0	1	0	0
	01	0	1	1	1
	11	1	1	1	0
	10	0	0	1	0

Min POS:

$$\underline{(w + y + z)(w + x + y')} \bullet$$

$$\underline{(w' + y' + z)(w' + x + y)}$$

- c) Use the K-map below to calculate both a minimal SOP expression and a minimal POS expression for F .

		AB			
		00	01	11	10
CD	00	0	0	0	0
	01	1	1	0	0
	11	0	1	0	0
	10	0	0	0	0

Min SOP: $F(A,B,C,D) = A'BD + A'C'D$

Min POS: $F(A,B,C,D) = A'(B+C')D$