

## Foundations of Artificial Intelligence

Prof. Dr. J. Boedecker, Prof. Dr. W. Burgard, Prof. Dr. F. Hutter, Prof. Dr. B. Nebel  
T. Schulte, R. Rajan, S. Adriaensen, K. Sirohi  
Summer Term 2021

University of Freiburg  
Department of Computer Science

### Exercise Sheet 6 — Solutions

#### Exercise 6.1 (Semantics of Predicate Logic)

Consider the Interpretation  $\mathcal{I} = \langle \mathcal{D}, \cdot^{\mathcal{I}} \rangle$  with

- $\mathcal{D} = \{0, 1, 2, 3\}$
- $even^{\mathcal{I}} = \{0, 2\}$
- $odd^{\mathcal{I}} = \{1, 3\}$
- $lessThan^{\mathcal{I}} = \{(0, 1), (0, 2), (0, 3), (1, 2), (1, 3), (2, 3)\}$
- $two^{\mathcal{I}} = 2$
- $plus^{\mathcal{I}} : \mathcal{D} \times \mathcal{D} \rightarrow \mathcal{D}, plus^{\mathcal{I}}(a, b) = (a + b) \bmod 4$

and the variable assignment  $\alpha = \{(x, 0), (y, 1)\}$ .

Decide for the following formulae  $\theta_i$  if  $\mathcal{I}$  is a model for  $\theta_i$  under  $\alpha$ , i.e. if  $\mathcal{I}, \alpha \models \theta_i$ .

Explain your answer by formally applying the semantics.

- (a)  $\theta_1 = odd(y) \wedge even(two)$
- (b)  $\theta_2 = \forall x (even(x) \vee odd(x))$
- (c)  $\theta_3 = \forall x \exists y lessThan(x, y)$
- (d)  $\theta_4 = \forall x (even(x) \Rightarrow \exists y lessThan(x, y))$
- (e)  $\theta_5 = \forall x (odd(x) \Rightarrow even(plus(x, y)))$

#### Solution:

- (a)  $\theta_1^{\mathcal{I}} =$   
 $(odd(y) \wedge even(two))^{\mathcal{I}} =$   
 $odd^{\mathcal{I}}(y^{\mathcal{I}}) \wedge even^{\mathcal{I}}(two^{\mathcal{I}}) =$   
 $odd^{\mathcal{I}}(1) \wedge even^{\mathcal{I}}(2) =$   
 $\top \wedge \top = \top$ , since  $1 \in odd^{\mathcal{I}}$  and  $2 \in even^{\mathcal{I}}$ .  
Thus,  $\mathcal{I}, \alpha \models \theta_1$ .

- (b) Let  $\theta_2 = \forall x \phi_2$  with  $\phi_2 = even(x) \vee odd(x)$ .  
Then  $\mathcal{I}, \alpha \models \theta_2$  iff  $\mathcal{I}, \alpha[x/d](\phi_2) = \top$  for all  $d \in \mathcal{D}$ .  
Notice that here  $x$  is bound by a quantifier and therefore the variable assignment in the definition of  $\alpha$  is overridden by  $\alpha[x/d]$ .

Since our universe  $\mathcal{D}$  is finite (and small), we can evaluate the above for all  $d \in \mathcal{D}$ :

$$\begin{aligned}\mathcal{I}, \alpha[x/0](\phi_2) &= \text{even}^{\mathcal{I}}(0) \vee \text{odd}^{\mathcal{I}}(0) = \top \vee \perp = \top \\ \mathcal{I}, \alpha[x/1](\phi_2) &= \text{even}^{\mathcal{I}}(1) \vee \text{odd}^{\mathcal{I}}(1) = \perp \vee \top = \top \\ \mathcal{I}, \alpha[x/2](\phi_2) &= \text{even}^{\mathcal{I}}(2) \vee \text{odd}^{\mathcal{I}}(2) = \top \vee \perp = \top \\ \mathcal{I}, \alpha[x/3](\phi_2) &= \text{even}^{\mathcal{I}}(3) \vee \text{odd}^{\mathcal{I}}(3) = \perp \vee \top = \top\end{aligned}$$

Thus,  $\mathcal{I}, \alpha \models \theta_2$ .

- (c) Let  $\theta_3 = \forall x \exists y \phi_3$  with  $\phi_3 = \text{lessThan}(x, y)$ .  
Then  $\mathcal{I}, \alpha \models \theta_3$  iff for all  $d_1 \in \mathcal{D}$  there exists a  $d_2 \in \mathcal{D}$  so that  $\mathcal{I}, \alpha[x/d_1, y/d_2](\phi_3) = \top$ .  
In this case it is clear that for  $d_1 = 3$  no such  $d_2$  exists, more formally,  $\text{lessThan}^{\mathcal{I}}(3, d_2) = \perp$  for all  $d_2 \in \mathcal{D}$  and therefore  $\mathcal{I}, \alpha \not\models \theta_3$ .
- (d) Let  $\theta_4 = \forall x \phi_4$  with  $\phi_4 = \text{even}(x) \Rightarrow \exists y \text{ lessThan}(x, y)$ .  
Then  $\mathcal{I}, \alpha \models \theta_4$  iff  $\mathcal{I}, \alpha[x/d_1](\phi_4) = \top$  for all  $d_1 \in \mathcal{D}$ .  
We again consider all cases, starting odd numbers, i.e.,  $d_1 \in \{1, 3\}$ :  
 $(\text{even}(x) \Rightarrow \exists y \text{ lessThan}(x, y))^{\mathcal{I}} = \neg \text{even}^{\mathcal{I}}(d_1) \vee \dots = \neg \perp \vee \dots = \top \vee \dots = \top$ .  
For  $d_1 \in \{0, 2\}$  we have  $\neg \text{even}^{\mathcal{I}}(d_1) = \perp$ , thus we have to find a corresponding  $d_2 \in \mathcal{D}$  with  $\mathcal{I}, \alpha[x/d_1, y/d_2](\text{lessThan}(x, y)) = \top$ . Here, we have  $(0, 1) \in \text{lessThan}^{\mathcal{I}}$  and  $(2, 3) \in \text{lessThan}^{\mathcal{I}}$ .  
Summing up all of the above, we showed that  $\mathcal{I}, \alpha \models \theta_4$ .
- (e) Let  $\theta_5 = \forall x \phi_5$  with  $\phi_5 = \text{odd}(x) \Rightarrow \text{even}(\text{plus}(x, y))$ .  
Then  $\mathcal{I}, \alpha \models \theta_5$  iff  $\mathcal{I}, \alpha[x/d](\phi_5) = \top$  for all  $d \in \mathcal{D}$ .  
 $\mathcal{I}, \alpha[x/d](\phi_5) = \neg \text{odd}^{\mathcal{I}}(d) \vee \text{even}^{\mathcal{I}}(\text{plus}^{\mathcal{I}}(d, 1))$   
Notice that here  $y$  is a free variable and therefore the assignment  $[y/1]$  is applied.  
Now we consider all cases for  $d$ :  
 $\mathcal{I}, \alpha[x/0](\phi_5) = \neg \text{odd}^{\mathcal{I}}(0) \vee \text{even}^{\mathcal{I}}(\text{plus}^{\mathcal{I}}(0, 1)) = \neg \text{odd}^{\mathcal{I}}(0) \vee \text{even}^{\mathcal{I}}(1) = \top \vee \perp = \top$   
 $\mathcal{I}, \alpha[x/1](\phi_5) = \neg \text{odd}^{\mathcal{I}}(1) \vee \text{even}^{\mathcal{I}}(\text{plus}^{\mathcal{I}}(1, 1)) = \neg \text{odd}^{\mathcal{I}}(1) \vee \text{even}^{\mathcal{I}}(2) = \perp \vee \top = \top$   
 $\mathcal{I}, \alpha[x/2](\phi_5) = \neg \text{odd}^{\mathcal{I}}(2) \vee \text{even}^{\mathcal{I}}(\text{plus}^{\mathcal{I}}(2, 1)) = \neg \text{odd}^{\mathcal{I}}(2) \vee \text{even}^{\mathcal{I}}(3) = \top \vee \perp = \top$   
 $\mathcal{I}, \alpha[x/3](\phi_5) = \neg \text{odd}^{\mathcal{I}}(3) \vee \text{even}^{\mathcal{I}}(\text{plus}^{\mathcal{I}}(3, 1)) = \neg \text{odd}^{\mathcal{I}}(3) \vee \text{even}^{\mathcal{I}}(0) = \perp \vee \top = \top$   
Thus,  $\mathcal{I}, \alpha \models \theta_5$ .

### Exercise 6.2 (Semantics of Predicate Logic)

Consider the Interpretation  $\mathcal{I} = \langle \mathcal{D}, \cdot^{\mathcal{I}} \rangle$  and the variable assignment  $\alpha$ :

- $\mathcal{D} = \{a, b, c\}$
- $P^{\mathcal{I}} = \{(a, a), (a, b), (b, a), (b, b), (b, c), (c, a)\}$
- $Q^{\mathcal{I}} = \{a, b\}$
- $R^{\mathcal{I}} = \{(a, a), (a, b), (a, c), (b, c), (c, b)\}$

- $\alpha = \{(v, a), (w, b)\}$

Decide for the following formulae  $\theta_i$  if  $\mathcal{I}$  is a model for  $\theta_i$  under  $\alpha$ , i.e. if  $\mathcal{I}, \alpha \models \theta_i$ . Explain your answer by formally applying the semantics.

- (a)  $\theta_1 = \forall x (P(x, w) \Rightarrow Q(x))$
- (b)  $\theta_2 = \exists x (R(v, x) \Rightarrow P(x, x))$
- (c)  $\theta_3 = \forall x \forall y (R(x, y) \iff Q(y))$
- (d)  $\theta_4 = \left[ \neg \forall x \forall y (Q(y) \vee P(x, y)) \right] \wedge \left[ \exists z (Q(z) \vee P(w, z)) \right]$

**Solution:**

- (a) Let  $\theta_1 = \forall x \phi_1$  with  $\phi_1 = (P(x, w) \Rightarrow Q(x))$ .  
Then  $\mathcal{I}, \alpha \models \theta_1$  iff  $\mathcal{I}, \alpha[x/d](\phi_1) = \top$  for all  $d \in \mathcal{D}$ .  
Notice that here  $x$  is bound by a quantifier and therefore the variable assignment in the definition of  $\alpha$  is overridden by  $\alpha[x/d]$ .  
Also  $w$  is a free variable and therefore the assignment  $[w/b]$  is applied.  
Since our universe  $\mathcal{D}$  is finite (and small), we can evaluate the above for all  $d \in \mathcal{D}$ :  

$$\mathcal{I}, \alpha[x/a](\phi_1) = \neg P^{\mathcal{I}}(a, b) \vee Q^{\mathcal{I}}(a) = \perp \vee \top = \top$$

$$\mathcal{I}, \alpha[x/b](\phi_1) = \neg P^{\mathcal{I}}(b, b) \vee Q^{\mathcal{I}}(b) = \perp \vee \top = \top$$

$$\mathcal{I}, \alpha[x/c](\phi_1) = \neg P^{\mathcal{I}}(c, b) \vee Q^{\mathcal{I}}(c) = \top \vee \perp = \top$$
 Thus,  $\mathcal{I}, \alpha \models \theta_1$ .
- (b) Let  $\theta_2 = \exists x \phi_2$  with  $\phi_2 = (R(v, x) \Rightarrow P(x, x))$ .  
Then  $\mathcal{I}, \alpha \models \theta_2$  if there exists a  $d \in \mathcal{D}$  so that  $\mathcal{I}, \alpha[x/d](\phi_2) = \top$ .  
Also  $v$  is a free variable and therefore the assignment  $[v/a]$  is applied.  
One case which can be applied is  $d = b$ .  

$$\mathcal{I}, \alpha[x/b](\phi_2) = \neg R^{\mathcal{I}}(a, b) \vee P^{\mathcal{I}}(b, b) = \perp \vee \top = \top$$
 Thus  $\mathcal{I}, \alpha \models \theta_2$ .
- (c) Let  $\theta_3 = \forall x \forall y \phi_3$  with  $\phi_3 = (R(x, y) \iff Q(y))$ .  
Then  $\mathcal{I}, \alpha \models \theta_3$  iff  $\mathcal{I}, \alpha[x/d1][y/d2](\phi_3) = \top$  for all  $d1$  and  $d2 \in \mathcal{D}$ .  
However, it can does not satisfy by assigning  $x=a$  and  $y = c$ ,  

$$\mathcal{I}, \alpha[x/a][y/c](\phi_3) = (R^{\mathcal{I}}(a, c) \iff Q^{\mathcal{I}}(c)) = \perp \iff \top = \perp$$
 Hence  $\mathcal{I}, \alpha \not\models \theta_3$ .
- (d)  $\mathcal{I}, \alpha \models \theta_4$   
Let  $\theta_4 = \neg \theta_4^a \wedge \theta_4^b$  with  

$$\theta_4^a = \forall x \forall y (Q(y) \vee P(x, y)) \text{ and } \theta_4^b = \exists z (Q(z) \vee P(w, z))$$
 Thus,  $\mathcal{I}, \alpha$  must model  $\theta_4^b$  but not  $\theta_4^a$ .  

$$\mathcal{I}, \alpha \not\models \theta_4^a, \text{ since for } y = c, c \notin Q^{\mathcal{I}} \text{ and } (c, c) \notin P^{\mathcal{I}}$$

$$\mathcal{I}, \alpha \models \theta_4^b, \text{ since, for instance, } a \in Q^{\mathcal{I}}.$$