

MT-1004

Linear Algebra

Fall 2023

Week # 4

National University of Computer and Emerging Sciences

September 14, 2023

Section 1.8

Introduction to Linear Transformations

Motivation

Let A be an $m \times n$ matrix.

Dr Ali and Dr Sara

Motivation

Let A be an $m \times n$ matrix. For the matrix equation $Ax = b$ we have learned to describe

Motivation

Let A be an $m \times n$ matrix. For the matrix equation $Ax = b$ we have learned to describe

- ▶ the solution set: all x in \mathbb{R}^n making the equation true.

Motivation

Let A be an $m \times n$ matrix. For the matrix equation $Ax = b$ we have learned to describe

- ▶ the solution set: all x in \mathbb{R}^n making the equation true.
- ▶ the column span: the set of all b in \mathbb{R}^m making the equation consistent.

Motivation

Let A be an $m \times n$ matrix. For the matrix equation $Ax = b$ we have learned to describe

- ▶ the solution set: all x in \mathbb{R}^n making the equation true.
- ▶ the column span: the set of all b in \mathbb{R}^m making the equation consistent.

It turns out these two sets are very closely related to each other.

Motivation

Let A be an $m \times n$ matrix. For the matrix equation $Ax = b$ we have learned to describe

- ▶ the solution set: all x in \mathbb{R}^n making the equation true.
- ▶ the column span: the set of all b in \mathbb{R}^m making the equation consistent.

It turns out these two sets are very closely related to each other.

In order to understand this relationship, it helps to think of the matrix A as a *transformation* from \mathbb{R}^n to \mathbb{R}^m .

Motivation

Let A be an $m \times n$ matrix. For the matrix equation $Ax = b$ we have learned to describe

- ▶ the solution set: all x in \mathbb{R}^n making the equation true.
- ▶ the column span: the set of all b in \mathbb{R}^m making the equation consistent.

It turns out these two sets are very closely related to each other.

In order to understand this relationship, it helps to think of the matrix A as a *transformation* from \mathbb{R}^n to \mathbb{R}^m .

It's a special kind of transformation called a *linear transformation*.

Motivation

Let A be an $m \times n$ matrix. For the matrix equation $Ax = b$ we have learned to describe

- ▶ the solution set: all x in \mathbb{R}^n making the equation true.
- ▶ the column span: the set of all b in \mathbb{R}^m making the equation consistent.

It turns out these two sets are very closely related to each other.

In order to understand this relationship, it helps to think of the matrix A as a *transformation* from \mathbb{R}^n to \mathbb{R}^m .

It's a special kind of transformation called a *linear transformation*.

This is also a way to understand the *geometry of matrices*.

Transformations

Definition


A **transformation** (or **function** or **map**) from \mathbb{R}^n to \mathbb{R}^m is a rule T that assigns to each vector x in \mathbb{R}^n a vector $T(x)$ in \mathbb{R}^m .

Transformations

Definition

A **transformation** (or **function** or **map**) from \mathbb{R}^n to \mathbb{R}^m is a rule T that assigns to each vector x in \mathbb{R}^n a vector $T(x)$ in \mathbb{R}^m .

- ▶ \mathbb{R}^n is called the **domain** of T (the inputs).




\mathbb{R}^n
domain

Transformations


Definition

A **transformation** (or **function** or **map**) from \mathbb{R}^n to \mathbb{R}^m is a rule T that assigns to each vector x in \mathbb{R}^n a vector $T(x)$ in \mathbb{R}^m .

- ▶ \mathbb{R}^n is called the **domain** of T (the inputs).
- ▶ \mathbb{R}^m is called the **codomain** of T (the outputs).



\mathbb{R}^n
domain



\mathbb{R}^m
codomain


Transformations

Definition

A **transformation** (or **function** or **map**) from \mathbb{R}^n to \mathbb{R}^m is a rule T that assigns to each vector x in \mathbb{R}^n a vector $T(x)$ in \mathbb{R}^m .

- ▶ \mathbb{R}^n is called the **domain** of T (the inputs).
- ▶ \mathbb{R}^m is called the **codomain** of T (the outputs).
- ▶ For x in \mathbb{R}^n , the vector $T(x)$ in \mathbb{R}^m is the **image** of x under T .

Notation: $x \mapsto T(x)$.




Transformations

Definition

A **transformation** (or **function** or **map**) from \mathbb{R}^n to \mathbb{R}^m is a rule T that assigns to each vector x in \mathbb{R}^n a vector $T(x)$ in \mathbb{R}^m .

- ▶ \mathbb{R}^n is called the **domain** of T (the inputs).
- ▶ \mathbb{R}^m is called the **codomain** of T (the outputs).
- ▶ For x in \mathbb{R}^n , the vector $T(x)$ in \mathbb{R}^m is the **image** of x under T .
Notation: $x \mapsto T(x)$.
- ▶ The set of all images $\{T(x) \mid x \text{ in } \mathbb{R}^n\}$ is the **range** of T .



Transformations


Definition

A **transformation** (or **function** or **map**) from \mathbb{R}^n to \mathbb{R}^m is a rule T that assigns to each vector x in \mathbb{R}^n a vector $T(x)$ in \mathbb{R}^m .

- ▶ \mathbb{R}^n is called the **domain** of T (the inputs).
 - ▶ \mathbb{R}^m is called the **codomain** of T (the outputs).
 - ▶ For x in \mathbb{R}^n , the vector $T(x)$ in \mathbb{R}^m is the **image** of x under T .
- Notation:** $x \mapsto T(x)$.
- ▶ The set of all images $\{T(x) \mid x \text{ in } \mathbb{R}^n\}$ is the **range** of T .

Notation:

$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m$ means T is a transformation from \mathbb{R}^n to \mathbb{R}^m .



Transformations


Definition

A **transformation** (or **function** or **map**) from \mathbb{R}^n to \mathbb{R}^m is a rule T that assigns to each vector x in \mathbb{R}^n a vector $T(x)$ in \mathbb{R}^m .

- ▶ \mathbb{R}^n is called the **domain** of T (the inputs).
 - ▶ \mathbb{R}^m is called the **codomain** of T (the outputs).
 - ▶ For x in \mathbb{R}^n , the vector $T(x)$ in \mathbb{R}^m is the **image** of x under T .
- Notation:** $x \mapsto T(x)$.
- ▶ The set of all images $\{T(x) \mid x \text{ in } \mathbb{R}^n\}$ is the **range** of T .

Notation:

$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m$ means T is a transformation from \mathbb{R}^n to \mathbb{R}^m .



It may help to think of T as a “machine” that takes x as an input, and gives you $T(x)$ as the output.

Functions from Calculus

Many of the functions you know and love have domain and codomain \mathbb{R} .

Dr Ali and Dr Sara

Functions from Calculus

Many of the functions you know and love have domain and codomain \mathbb{R} .

$$\sin: \mathbb{R} \longrightarrow \mathbb{R} \quad \sin(x) = \left(\begin{array}{l} \text{the length of the opposite edge over the} \\ \text{hypotenuse of a right triangle with angle} \\ x \text{ in radians} \end{array} \right)$$

Functions from Calculus

Many of the functions you know and love have domain and codomain \mathbb{R} .

$$\sin: \mathbb{R} \longrightarrow \mathbb{R} \quad \sin(x) = \left(\begin{array}{l} \text{the length of the opposite edge over the} \\ \text{hypotenuse of a right triangle with angle} \\ x \text{ in radians} \end{array} \right)$$

Note how I've written down the *rule* that defines the function \sin .

Functions from Calculus

Many of the functions you know and love have domain and codomain \mathbb{R} .

$$\sin: \mathbb{R} \longrightarrow \mathbb{R} \quad \sin(x) = \left(\begin{array}{l} \text{the length of the opposite edge over the} \\ \text{hypotenuse of a right triangle with angle} \\ x \text{ in radians} \end{array} \right)$$

Note how I've written down the *rule* that defines the function \sin .

$$f: \mathbb{R} \longrightarrow \mathbb{R} \quad f(x) = x^2$$

Functions from Calculus

Many of the functions you know and love have domain and codomain \mathbb{R} .

$$\sin: \mathbb{R} \longrightarrow \mathbb{R} \quad \sin(x) = \left(\begin{array}{l} \text{the length of the opposite edge over the} \\ \text{hypotenuse of a right triangle with angle} \\ x \text{ in radians} \end{array} \right)$$

Note how I've written down the *rule* that defines the function \sin .

$$f: \mathbb{R} \longrightarrow \mathbb{R} \quad f(x) = x^2$$

Note that “ x^2 ” is sloppy (but common) notation for a function: it doesn't have a name!

Functions from Calculus

Many of the functions you know and love have domain and codomain \mathbb{R} .

$$\sin: \mathbb{R} \longrightarrow \mathbb{R} \quad \sin(x) = \left(\begin{array}{l} \text{the length of the opposite edge over the} \\ \text{hypotenuse of a right triangle with angle} \\ x \text{ in radians} \end{array} \right)$$

Note how I've written down the *rule* that defines the function \sin .

$$f: \mathbb{R} \longrightarrow \mathbb{R} \quad f(x) = x^2$$

Note that " x^2 " is sloppy (but common) notation for a function: it doesn't have a name!

You may be used to thinking of a function in terms of its graph.

$(x, \sin x)$

•

•

x

Functions from Calculus

Many of the functions you know and love have domain and codomain \mathbb{R} .

$$\sin: \mathbb{R} \longrightarrow \mathbb{R} \quad \sin(x) = \left(\begin{array}{l} \text{the length of the opposite edge over the} \\ \text{hypotenuse of a right triangle with angle} \\ x \text{ in radians} \end{array} \right)$$

Note how I've written down the *rule* that defines the function \sin .

$$f: \mathbb{R} \longrightarrow \mathbb{R} \quad f(x) = x^2$$

Note that “ x^2 ” is sloppy (but common) notation for a function: it doesn't have a name!

You may be used to thinking of a function in terms of its graph.

$(x, \sin x)$

•

•

x

The horizontal axis is the domain, and the vertical axis is the codomain.

Functions from Calculus

Many of the functions you know and love have domain and codomain \mathbb{R} .

$$\sin: \mathbb{R} \longrightarrow \mathbb{R} \quad \sin(x) = \left(\begin{array}{l} \text{the length of the opposite edge over the} \\ \text{hypotenuse of a right triangle with angle} \\ x \text{ in radians} \end{array} \right)$$

Note how I've written down the *rule* that defines the function \sin .

$$f: \mathbb{R} \longrightarrow \mathbb{R} \quad f(x) = x^2$$

Note that " x^2 " is sloppy (but common) notation for a function: it doesn't have a name!

You may be used to thinking of a function in terms of its graph.

$(x, \sin x)$

•

•

x

The horizontal axis is the domain, and the vertical axis is the codomain.

This is fine when the domain and codomain are \mathbb{R} , but it's hard to do when they're \mathbb{R}^2 and \mathbb{R}^3 !

Functions from Calculus

Many of the functions you know and love have domain and codomain \mathbb{R} .

$$\sin: \mathbb{R} \longrightarrow \mathbb{R} \quad \sin(x) = \left(\begin{array}{l} \text{the length of the opposite edge over the} \\ \text{hypotenuse of a right triangle with angle} \\ x \text{ in radians} \end{array} \right)$$

Note how I've written down the *rule* that defines the function \sin .

$$f: \mathbb{R} \longrightarrow \mathbb{R} \quad f(x) = x^2$$

Note that “ x^2 ” is sloppy (but common) notation for a function: it doesn't have a name!

You may be used to thinking of a function in terms of its graph.

$(x, \sin x)$

•

•

x

The horizontal axis is the domain, and the vertical axis is the codomain.

This is fine when the domain and codomain are \mathbb{R} , but it's hard to do when they're \mathbb{R}^2 and \mathbb{R}^3 ! You need ____ dimensions to draw that graph.

Functions from Calculus

Many of the functions you know and love have domain and codomain \mathbb{R} .

$$\sin: \mathbb{R} \longrightarrow \mathbb{R} \quad \sin(x) = \left(\begin{array}{l} \text{the length of the opposite edge over the} \\ \text{hypotenuse of a right triangle with angle} \\ x \text{ in radians} \end{array} \right)$$

Note how I've written down the *rule* that defines the function \sin .

$$f: \mathbb{R} \longrightarrow \mathbb{R} \quad f(x) = x^2$$

Note that “ x^2 ” is sloppy (but common) notation for a function: it doesn't have a name!

You may be used to thinking of a function in terms of its graph.

$(x, \sin x)$

•

•

x

The horizontal axis is the domain, and the vertical axis is the codomain.

This is fine when the domain and codomain are \mathbb{R} , but it's hard to do when they're \mathbb{R}^2 and \mathbb{R}^3 ! You need five dimensions to draw that graph.

Matrix Transformations

Most of the transformations we encounter in this class will come from (surprise) matrices!

Dr Ali and Dr Sara

Matrix Transformations

Most of the transformations we encounter in this class will come from (surprise) matrices!

Definition

Let A be an $m \times n$ matrix. The **matrix transformation** associated to A is the transformation

$$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m \quad \text{defined by} \quad T(x) = Ax.$$

Matrix Transformations

Most of the transformations we encounter in this class will come from (surprise) matrices!

Definition

Let A be an $m \times n$ matrix. The **matrix transformation** associated to A is the transformation

$$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m \quad \text{defined by} \quad T(x) = Ax.$$

$$A_{m \times n} X_{n \times 1}$$

In other words, T takes the vector x in \mathbb{R}^n to the vector Ax in \mathbb{R}^m .

Matrix Transformations

Most of the transformations we encounter in this class will come from (surprise) matrices!

Definition

Let A be an $m \times n$ matrix. The **matrix transformation** associated to A is the transformation

$$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m \quad \text{defined by} \quad T(x) = Ax.$$

$$A_{m \times n} X_{n \times 1}$$

In other words, T takes the vector x in \mathbb{R}^n to the vector Ax in \mathbb{R}^m .

- The *domain* of T is

Matrix Transformations

Most of the transformations we encounter in this class will come from (surprise) matrices!

Definition

Let A be an $m \times n$ matrix. The **matrix transformation** associated to A is the transformation

$$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m \quad \text{defined by} \quad T(x) = Ax.$$

$$A_{m \times n} X_{n \times 1}$$

In other words, T takes the vector x in \mathbb{R}^n to the vector Ax in \mathbb{R}^m .

- The *domain* of T is \mathbb{R}^n , which is the number of _____ of A .

Matrix Transformations

Most of the transformations we encounter in this class will come from (surprise) matrices!

Definition

Let A be an $m \times n$ matrix. The **matrix transformation** associated to A is the transformation

$$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m \quad \text{defined by} \quad T(x) = Ax.$$

$$A_{m \times n} X_{n \times 1}$$

In other words, T takes the vector x in \mathbb{R}^n to the vector Ax in \mathbb{R}^m .

- The *domain* of T is \mathbb{R}^n , which is the number of *columns* of A .

Matrix Transformations

Most of the transformations we encounter in this class will come from (surprise) matrices!

Definition

Let A be an $m \times n$ matrix. The **matrix transformation** associated to A is the transformation

$$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m \quad \text{defined by} \quad T(x) = Ax.$$

$$A_{m \times n} X_{n \times 1}$$

In other words, T takes the vector x in \mathbb{R}^n to the vector Ax in \mathbb{R}^m .

- ▶ The *domain* of T is \mathbb{R}^n , which is the number of *columns* of A .
- ▶ The *codomain* of T is

Matrix Transformations

Most of the transformations we encounter in this class will come from (surprise) matrices!

Definition

Let A be an $m \times n$ matrix. The **matrix transformation** associated to A is the transformation

$$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m \quad \text{defined by} \quad T(x) = Ax.$$

$$A_{m \times n} X_{n \times 1}$$

In other words, T takes the vector x in \mathbb{R}^n to the vector Ax in \mathbb{R}^m .

- ▶ The *domain* of T is \mathbb{R}^n , which is the number of *columns* of A .
- ▶ The *codomain* of T is \mathbb{R}^m , which is the number of ____ of A .

Matrix Transformations

Most of the transformations we encounter in this class will come from (surprise) matrices!

Definition

Let A be an $m \times n$ matrix. The **matrix transformation** associated to A is the transformation

$$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m \quad \text{defined by} \quad T(x) = Ax.$$

$$A_{m \times n} X_{n \times 1}$$

In other words, T takes the vector x in \mathbb{R}^n to the vector Ax in \mathbb{R}^m .

- ▶ The *domain* of T is \mathbb{R}^n , which is the number of *columns* of A .
- ▶ The *codomain* of T is \mathbb{R}^m , which is the number of *rows* of A .

Matrix Transformations

Most of the transformations we encounter in this class will come from (surprise) matrices!

Definition

Let A be an $m \times n$ matrix. The **matrix transformation** associated to A is the transformation

$$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m \quad \text{defined by} \quad T(x) = Ax.$$

$$A_{m \times n} X_{n \times 1}$$

In other words, T takes the vector x in \mathbb{R}^n to the vector Ax in \mathbb{R}^m .

- ▶ The *domain* of T is \mathbb{R}^n , which is the number of *columns* of A .
- ▶ The *codomain* of T is \mathbb{R}^m , which is the number of *rows* of A .
- ▶ The *range* of T is the set of all images of T :

$$T(x) = Ax = \begin{pmatrix} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = x_1 v_1 + x_2 v_2 + \cdots + x_n v_n.$$

This is the

Matrix Transformations

Most of the transformations we encounter in this class will come from (surprise) matrices!

Definition

Let A be an $m \times n$ matrix. The **matrix transformation** associated to A is the transformation

$$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m \quad \text{defined by} \quad T(x) = Ax.$$

$$A_{m \times n} X_{n \times 1}$$

In other words, T takes the vector x in \mathbb{R}^n to the vector Ax in \mathbb{R}^m .

- ▶ The *domain* of T is \mathbb{R}^n , which is the number of *columns* of A .
- ▶ The *codomain* of T is \mathbb{R}^m , which is the number of *rows* of A .
- ▶ The *range* of T is the set of all images of T :

$$T(x) = Ax = \begin{pmatrix} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = x_1 v_1 + x_2 v_2 + \cdots + x_n v_n.$$

This is the *column span* of A . It is a span of vectors in the codomain.

Matrix Transformations

Most of the transformations we encounter in this class will come from (surprise) matrices!

Definition

Let A be an $m \times n$ matrix. The **matrix transformation** associated to A is the transformation

$$T: \mathbb{R}^n \longrightarrow \mathbb{R}^m \quad \text{defined by} \quad T(x) = Ax.$$

$$A_{m \times n} X_{n \times 1}$$

In other words, T takes the vector x in \mathbb{R}^n to the vector Ax in \mathbb{R}^m .

- ▶ The *domain* of T is \mathbb{R}^n , which is the number of *columns* of A .
- ▶ The *codomain* of T is \mathbb{R}^m , which is the number of *rows* of A .
- ▶ The *range* of T is the set of all images of T :

$$T(x) = Ax = \begin{pmatrix} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = x_1 v_1 + x_2 v_2 + \cdots + x_n v_n.$$

This is the *column span* of A . It is a span of vectors in the codomain.

Your life will be much easier if you just remember these.

Matrix Transformations

Example

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$.

Dr Ali and Dr Sara

Matrix Transformations

Example

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

Matrix Transformations

Example

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

Matrix Transformations

Example

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

► If $u = \begin{pmatrix} 3 \\ 4 \end{pmatrix}$ then $T(u) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 7 \\ 4 \\ 7 \end{pmatrix}.$

Matrix Transformations

Example

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

► If $u = \begin{pmatrix} 3 \\ 4 \end{pmatrix}$ then $T(u) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 7 \\ 4 \\ 7 \end{pmatrix}$.

► Let $b = \begin{pmatrix} 7 \\ 5 \\ 7 \end{pmatrix}$. Find v in \mathbb{R}^2 such that $T(v) = b$.

Matrix Transformations

Example

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

► If $u = \begin{pmatrix} 3 \\ 4 \end{pmatrix}$ then $T(u) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 7 \\ 4 \\ 7 \end{pmatrix}$.

► Let $b = \begin{pmatrix} 7 \\ 5 \\ 7 \end{pmatrix}$. Find v in \mathbb{R}^2 such that $T(v) = b$.

Matrix Transformations

Example

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

► If $u = \begin{pmatrix} 3 \\ 4 \end{pmatrix}$ then $T(u) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 7 \\ 4 \\ 7 \end{pmatrix}$.

► Let $b = \begin{pmatrix} 7 \\ 5 \\ 7 \end{pmatrix}$. Find v in \mathbb{R}^2 such that $T(v) = b$. Is there more than one?

We want to find v such that $T(v) = Av = b$. We know how to do that:

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix} v = \begin{pmatrix} 7 \\ 5 \\ 7 \end{pmatrix} \xrightarrow[\text{augmented matrix}]{\text{~~~~~}} \left(\begin{array}{cc|c} 1 & 1 & 7 \\ 0 & 1 & 5 \\ 1 & 1 & 7 \end{array} \right) \xrightarrow[\text{reduce}]{\text{~~~~~}} \left(\begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & 5 \\ 0 & 0 & 0 \end{array} \right).$$

This gives $x = 2$ and $y = 5$, or $v = \begin{pmatrix} 2 \\ 5 \end{pmatrix}$ (unique). In other words,

$$T(v) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ 5 \end{pmatrix} = \begin{pmatrix} 7 \\ 5 \\ 7 \end{pmatrix}.$$

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

- Is there any c in \mathbb{R}^3 such that there is more than one v in \mathbb{R}^2 with $T(v) = c$?

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

- Is there any c in \mathbb{R}^3 such that there is more than one v in \mathbb{R}^2 with $T(v) = c$?

Translation: is there any c in \mathbb{R}^3 such that the solution set of $Ax = c$ has more than one vector v in it?

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

- Is there any c in \mathbb{R}^3 such that there is more than one v in \mathbb{R}^2 with $T(v) = c$?

Translation: is there any c in \mathbb{R}^3 such that the solution set of $Ax = c$ has more than one vector v in it?

The solution set of $Ax = c$ is a translate of the solution set of $Ax = b$ (from before), which has ___ vector in it.

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

- Is there any c in \mathbb{R}^3 such that there is more than one v in \mathbb{R}^2 with $T(v) = c$?

Translation: is there any c in \mathbb{R}^3 such that the solution set of $Ax = c$ has more than one vector v in it?

The solution set of $Ax = c$ is a translate of the solution set of $Ax = b$ (from before), which has one vector in it.

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

- Is there any c in \mathbb{R}^3 such that there is more than one v in \mathbb{R}^2 with $T(v) = c$?

Translation: is there any c in \mathbb{R}^3 such that the solution set of $Ax = c$ has more than one vector v in it?

The solution set of $Ax = c$ is a translate of the solution set of $Ax = b$ (from before), which has one vector in it. So the solution set to $Ax = c$ has only one vector.

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

- Is there any c in \mathbb{R}^3 such that there is more than one v in \mathbb{R}^2 with $T(v) = c$?

Translation: is there any c in \mathbb{R}^3 such that the solution set of $Ax = c$ has more than one vector v in it?

The solution set of $Ax = c$ is a translate of the solution set of $Ax = b$ (from before), which has one vector in it. So the solution set to $Ax = c$ has only one vector. So no!

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

- Is there any c in \mathbb{R}^3 such that there is more than one v in \mathbb{R}^2 with $T(v) = c$?

Translation: is there any c in \mathbb{R}^3 such that the solution set of $Ax = c$ has more than one vector v in it?

The solution set of $Ax = c$ is a translate of the solution set of $Ax = b$ (from before), which has one vector in it. So the solution set to $Ax = c$ has only one vector. So no!

- Find c such that there is *no* v with $T(v) = c$.

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

- Is there any c in \mathbb{R}^3 such that there is more than one v in \mathbb{R}^2 with $T(v) = c$?

Translation: is there any c in \mathbb{R}^3 such that the solution set of $Ax = c$ has more than one vector v in it?

The solution set of $Ax = c$ is a translate of the solution set of $Ax = b$ (from before), which has one vector in it. So the solution set to $Ax = c$ has only one vector. So no!

- Find c such that there is *no* v with $T(v) = c$.

Translation: Find c such that $Ax = c$ is inconsistent.

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

- Is there any c in \mathbb{R}^3 such that there is more than one v in \mathbb{R}^2 with $T(v) = c$?

Translation: is there any c in \mathbb{R}^3 such that the solution set of $Ax = c$ has more than one vector v in it?

The solution set of $Ax = c$ is a translate of the solution set of $Ax = b$ (from before), which has one vector in it. So the solution set to $Ax = c$ has only one vector. So no!

- Find c such that there is *no* v with $T(v) = c$.

Translation: Find c such that $Ax = c$ is inconsistent.

Translation: Find c not in the column span of A (i.e., the range of T).

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

- Is there any c in \mathbb{R}^3 such that there is more than one v in \mathbb{R}^2 with $T(v) = c$?

Translation: is there any c in \mathbb{R}^3 such that the solution set of $Ax = c$ has more than one vector v in it?

The solution set of $Ax = c$ is a translate of the solution set of $Ax = b$ (from before), which has one vector in it. So the solution set to $Ax = c$ has only one vector. So no!

- Find c such that there is *no* v with $T(v) = c$.

Translation: Find c such that $Ax = c$ is inconsistent.

Translation: Find c not in the column span of A (i.e., the range of T).

We could draw a picture, or notice: $a \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + b \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} a+b \\ b \\ a+b \end{pmatrix}$.

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

- Is there any c in \mathbb{R}^3 such that there is more than one v in \mathbb{R}^2 with $T(v) = c$?

Translation: is there any c in \mathbb{R}^3 such that the solution set of $Ax = c$ has more than one vector v in it?

The solution set of $Ax = c$ is a translate of the solution set of $Ax = b$ (from before), which has one vector in it. So the solution set to $Ax = c$ has only one vector. So no!

- Find c such that there is *no* v with $T(v) = c$.

Translation: Find c such that $Ax = c$ is inconsistent.

Translation: Find c not in the column span of A (i.e., the range of T).

We could draw a picture, or notice: $a \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + b \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} a+b \\ b \\ a+b \end{pmatrix}$. So anything in the column span has the same first and last coordinate.

Matrix Transformations

Example, continued

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$.

- Is there any c in \mathbb{R}^3 such that there is more than one v in \mathbb{R}^2 with $T(v) = c$?

Translation: is there any c in \mathbb{R}^3 such that the solution set of $Ax = c$ has more than one vector v in it?

The solution set of $Ax = c$ is a translate of the solution set of $Ax = b$ (from before), which has one vector in it. So the solution set to $Ax = c$ has only one vector. So no!

- Find c such that there is *no* v with $T(v) = c$.

Translation: Find c such that $Ax = c$ is inconsistent.

Translation: Find c not in the column span of A (i.e., the range of T).

We could draw a picture, or notice: $a \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + b \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} a+b \\ b \\ a+b \end{pmatrix}$. So

anything in the column span has the same first and last coordinate. So $c = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ is not in the column span (for example).

Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$.

Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$.

Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$. Then

$$T \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} x \\ y \\ 0 \end{pmatrix}.$$

Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$. Then

$$T \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} x \\ y \\ 0 \end{pmatrix}.$$

This is


Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$. Then

$$T \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} x \\ y \\ 0 \end{pmatrix}.$$

This is *projection onto the xy -axis*. Picture:



Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

Dr Ali and Dr Sara

Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

Dr Ali and Dr Sara

Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$. Then

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -x \\ y \end{pmatrix}.$$

Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$. Then

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -x \\ y \end{pmatrix}.$$

This is


Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$. Then

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -x \\ y \end{pmatrix}.$$

This is *reflection over the y-axis*. Picture:



Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

Dr Ali and Dr Sara

Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

Dr Ali and Dr Sara

Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$. Then

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ -y \end{pmatrix}.$$

Then

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -x \\ -y \end{pmatrix}.$$

Then

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} y \\ x \end{pmatrix}.$$

Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$. Then

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ -y \end{pmatrix}.$$

This is

Then

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -x \\ -y \end{pmatrix}.$$

This is

Then

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} y \\ x \end{pmatrix}.$$

This is

Matrix Transformations

Geometric example

Let $A = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$. Then

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ -y \end{pmatrix}.$$

This is *reflection over the x-axis*.

Let $A = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$. Then

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -x \\ -y \end{pmatrix}.$$

This is *reflection about origin*.

Let $A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$. Then

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} y \\ x \end{pmatrix}.$$


This is *reflection about the line $y = x$* .

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$. (T is called a **shear**.)

Poll

What does T do to this sheep?

Hint: first draw a picture what it does to the box *around* the sheep.




Poll

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and let $T(x) = Ax$, so $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$. (T is called a **shear**.)

Poll

What does T do to this sheep?

Hint: first draw a picture what it does to the box *around* the sheep.



Linear Transformations

Recall: If A is a matrix, u, v are vectors, and c is a scalar, then

$$A(u + v) = Au + Av \quad A(cv) = cAv.$$

Linear Transformations

Recall: If A is a matrix, u, v are vectors, and c is a scalar, then

$$A(u + v) = Au + Av \quad A(cv) = cAv.$$

So if $T(x) = Ax$ is a matrix transformation then,

$$T(u + v) = T(u) + T(v) \quad T(cv) = cT(v).$$

Linear Transformations

Recall: If A is a matrix, u, v are vectors, and c is a scalar, then

$$A(u + v) = Au + Av \quad A(cv) = cAv.$$

So if $T(x) = Ax$ is a matrix transformation then,

$$T(u + v) = T(u) + T(v) \quad T(cv) = cT(v).$$

This property is so special that it has its own name.

Linear Transformations

Recall: If A is a matrix, u, v are vectors, and c is a scalar, then

$$A(u + v) = Au + Av \quad A(cv) = cAv.$$

So if $T(x) = Ax$ is a matrix transformation then,

$$T(u + v) = T(u) + T(v) \quad T(cv) = cT(v).$$

This property is so special that it has its own name.

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **linear** if it satisfies the above equations for all vectors u, v in \mathbb{R}^n and all scalars c .

Linear Transformations

Recall: If A is a matrix, u, v are vectors, and c is a scalar, then

$$A(u + v) = Au + Av \quad A(cv) = cAv.$$

So if $T(x) = Ax$ is a matrix transformation then,

$$T(u + v) = T(u) + T(v) \quad T(cv) = cT(v).$$

This property is so special that it has its own name.

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **linear** if it satisfies the above equations for all vectors u, v in \mathbb{R}^n and all scalars c .

In other words, T “respects” addition and scalar multiplication.

Linear Transformations

Recall: If A is a matrix, u, v are vectors, and c is a scalar, then

$$A(u + v) = Au + Av \quad A(cv) = cAv.$$

So if $T(x) = Ax$ is a matrix transformation then,

$$T(u + v) = T(u) + T(v) \quad T(cv) = cT(v).$$

This property is so special that it has its own name.

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **linear** if it satisfies the above equations for all vectors u, v in \mathbb{R}^n and all scalars c .

In other words, T “respects” addition and scalar multiplication.

Check: if T is linear, then

$$T(0) = 0 \quad T(cu + dv) = cT(u) + dT(v)$$

for all vectors u, v and scalars c, d .

Linear Transformations

Recall: If A is a matrix, u, v are vectors, and c is a scalar, then

$$A(u + v) = Au + Av \quad A(cv) = cAv.$$

So if $T(x) = Ax$ is a matrix transformation then,

$$T(u + v) = T(u) + T(v) \quad T(cv) = cT(v).$$

This property is so special that it has its own name.

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **linear** if it satisfies the above equations for all vectors u, v in \mathbb{R}^n and all scalars c .

In other words, T “respects” addition and scalar multiplication.

Check: if T is linear, then

$$T(0) = 0 \quad T(cu + dv) = cT(u) + dT(v)$$

for all vectors u, v and scalars c, d . More generally,

$$T(c_1 v_1 + c_2 v_2 + \cdots + c_n v_n) = c_1 T(v_1) + c_2 T(v_2) + \cdots + c_n T(v_n).$$

Linear Transformations

Recall: If A is a matrix, u, v are vectors, and c is a scalar, then

$$A(u + v) = Au + Av \quad A(cv) = cAv.$$

So if $T(x) = Ax$ is a matrix transformation then,

$$T(u + v) = T(u) + T(v) \quad T(cv) = cT(v).$$

This property is so special that it has its own name.

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **linear** if it satisfies the above equations for all vectors u, v in \mathbb{R}^n and all scalars c .

In other words, T “respects” addition and scalar multiplication.

Check: if T is linear, then

$$T(0) = 0 \quad T(cu + dv) = cT(u) + dT(v)$$

for all vectors u, v and scalars c, d . More generally,

$$T(c_1 v_1 + c_2 v_2 + \cdots + c_n v_n) = c_1 T(v_1) + c_2 T(v_2) + \cdots + c_n T(v_n).$$

In engineering this is called **superposition**.

Linear Transformations

Dilation

Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $T(x) = 1.5x$. Is T linear? Check:

Linear Transformations

Dilation

Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $T(x) = 1.5x$. Is T linear? Check:

$$T(u + v) = 1.5(u + v) = 1.5u + 1.5v = T(u) + T(v)$$

$$T(cv) = 1.5(cv) = c(1.5v) = c(Tv).$$

Linear Transformations

Dilation

Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $T(x) = 1.5x$. Is T linear? Check:

$$T(u + v) = 1.5(u + v) = 1.5u + 1.5v = T(u) + T(v)$$

$$T(cv) = 1.5(cv) = c(1.5v) = c(Tv).$$

So T satisfies the two equations, hence T is linear.

Linear Transformations

Dilation

Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $T(x) = 1.5x$. Is T linear? Check:

$$T(u + v) = 1.5(u + v) = 1.5u + 1.5v = T(u) + T(v)$$

$$T(cv) = 1.5(cv) = c(1.5v) = c(Tv).$$

So T satisfies the two equations, hence T is linear.

This is called **dilation** or **scaling** (by a factor of 1.5). Picture:

Linear Transformations

Dilation


Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $T(x) = 1.5x$. Is T linear? Check:

$$T(u + v) = 1.5(u + v) = 1.5u + 1.5v = T(u) + T(v)$$

$$T(cv) = 1.5(cv) = c(1.5v) = c(Tv).$$

So T satisfies the two equations, hence T is linear.

This is called **dilation** or **scaling** (by a factor of 1.5). Picture:



Linear Transformations

Rotation

Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -y \\ x \end{pmatrix}.$$

Is T linear? Check:

Linear Transformations

Rotation

Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -y \\ x \end{pmatrix}.$$

Is T linear? Check:

$$T \left(\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} + \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right) = \begin{pmatrix} -u_2 \\ u_1 \end{pmatrix} + \begin{pmatrix} -v_2 \\ v_1 \end{pmatrix} = \begin{pmatrix} -(u_2 + v_2) \\ (u_1 + v_1) \end{pmatrix} = T \begin{pmatrix} u_1 + u_2 \\ v_1 + v_2 \end{pmatrix}$$

$$T \left(c \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right) = T \begin{pmatrix} cv_1 \\ cv_2 \end{pmatrix} = \begin{pmatrix} -cv_2 \\ cv_1 \end{pmatrix} = c \begin{pmatrix} -v_2 \\ v_1 \end{pmatrix} = cT \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}.$$

Linear Transformations

Rotation

Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -y \\ x \end{pmatrix}.$$

Is T linear? Check:

$$T \left(\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} + \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right) = \begin{pmatrix} -u_2 \\ u_1 \end{pmatrix} + \begin{pmatrix} -v_2 \\ v_1 \end{pmatrix} = \begin{pmatrix} -(u_2 + v_2) \\ (u_1 + v_1) \end{pmatrix} = T \begin{pmatrix} u_1 + u_2 \\ v_1 + v_2 \end{pmatrix}$$

$$T \left(c \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right) = T \begin{pmatrix} cv_1 \\ cv_2 \end{pmatrix} = \begin{pmatrix} -cv_2 \\ cv_1 \end{pmatrix} = c \begin{pmatrix} -v_2 \\ v_1 \end{pmatrix} = cT \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}.$$

So T satisfies the two equations, hence T is linear.

Linear Transformations

Rotation

Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by


$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -y \\ x \end{pmatrix}.$$

Is T linear? Check:

$$T \left(\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} + \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right) = \begin{pmatrix} -u_2 \\ u_1 \end{pmatrix} + \begin{pmatrix} -v_2 \\ v_1 \end{pmatrix} = \begin{pmatrix} -(u_2 + v_2) \\ u_1 + v_1 \end{pmatrix} = T \begin{pmatrix} u_1 + v_1 \\ u_2 + v_2 \end{pmatrix}$$

$$T \left(c \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right) = T \begin{pmatrix} cv_1 \\ cv_2 \end{pmatrix} = \begin{pmatrix} -cv_2 \\ cv_1 \end{pmatrix} = c \begin{pmatrix} -v_2 \\ v_1 \end{pmatrix} = c T \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}.$$

So T satisfies the two equations, hence T is linear. This is called **rotation** (by 90°). Picture:



Linear Transformations

Rotation

Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -y \\ x \end{pmatrix}.$$


Is T linear? Check:

$$T \left(\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} + \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right) = \begin{pmatrix} -u_2 \\ u_1 \end{pmatrix} + \begin{pmatrix} -v_2 \\ v_1 \end{pmatrix} = \begin{pmatrix} -(u_2 + v_2) \\ u_1 + v_1 \end{pmatrix} = T \begin{pmatrix} u_1 + u_2 \\ v_1 + v_2 \end{pmatrix}$$

$$T \left(c \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right) = T \begin{pmatrix} cv_1 \\ cv_2 \end{pmatrix} = \begin{pmatrix} -cv_2 \\ cv_1 \end{pmatrix} = c \begin{pmatrix} -v_2 \\ v_1 \end{pmatrix} = cT \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}.$$

So T satisfies the two equations, hence T is linear. This is called **rotation** (by 90°). Picture:

$$T \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} -2 \\ 1 \end{pmatrix}$$



Linear Transformations

Rotation

Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -y \\ x \end{pmatrix}.$$


Is T linear? Check:

$$T \left(\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} + \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right) = \begin{pmatrix} -u_2 \\ u_1 \end{pmatrix} + \begin{pmatrix} -v_2 \\ v_1 \end{pmatrix} = \begin{pmatrix} -(u_2 + v_2) \\ u_1 + v_1 \end{pmatrix} = T \begin{pmatrix} u_1 + u_2 \\ v_1 + v_2 \end{pmatrix}$$

$$T \left(c \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right) = T \begin{pmatrix} cv_1 \\ cv_2 \end{pmatrix} = \begin{pmatrix} -cv_2 \\ cv_1 \end{pmatrix} = c \begin{pmatrix} -v_2 \\ v_1 \end{pmatrix} = c T \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}.$$

So T satisfies the two equations, hence T is linear. This is called **rotation** (by 90°). Picture:

$$T \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} -2 \\ 1 \end{pmatrix}$$
$$T \begin{pmatrix} -1 \\ 1 \end{pmatrix} = \begin{pmatrix} -1 \\ -1 \end{pmatrix}$$



Linear Transformations

Rotation

Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -y \\ x \end{pmatrix}.$$

Is T linear? Check:

$$T \left(\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} + \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right) = \begin{pmatrix} -u_2 \\ u_1 \end{pmatrix} + \begin{pmatrix} -v_2 \\ v_1 \end{pmatrix} = \begin{pmatrix} -(u_2 + v_2) \\ u_1 + v_1 \end{pmatrix} = T \begin{pmatrix} u_1 + u_2 \\ v_1 + v_2 \end{pmatrix}$$


$$T \left(c \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right) = T \begin{pmatrix} cv_1 \\ cv_2 \end{pmatrix} = \begin{pmatrix} -cv_2 \\ cv_1 \end{pmatrix} = c \begin{pmatrix} -v_2 \\ v_1 \end{pmatrix} = cT \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}.$$

So T satisfies the two equations, hence T is linear. This is called **rotation** (by 90°). Picture:

$$T \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} -2 \\ 1 \end{pmatrix}$$

$$T \begin{pmatrix} -1 \\ 1 \end{pmatrix} = \begin{pmatrix} -1 \\ -1 \end{pmatrix}$$

$$T \begin{pmatrix} 0 \\ -2 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$$



Section 1.9

The Matrix of a Linear Transformation

Unit Coordinate Vectors

Definition

The **unit coordinate vectors** in \mathbb{R}^n are

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}.$$

Unit Coordinate Vectors

Definition

The **unit coordinate vectors** in \mathbb{R}^n are

This is what e_1, e_2, \dots mean,
for the rest of the class.

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}.$$

•


Unit Coordinate Vectors

Definition

The **unit coordinate vectors** in \mathbb{R}^n are

This is what e_1, e_2, \dots mean,
for the rest of the class.

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}.$$




Unit Coordinate Vectors

Definition

The **unit coordinate vectors** in \mathbb{R}^n are

This is what e_1, e_2, \dots mean,
for the rest of the class.

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}.$$



Note: if A is an $m \times n$ matrix with columns v_1, v_2, \dots, v_n , then $Ae_i = v_i$ for $i = 1, 2, \dots, n$:


Unit Coordinate Vectors

Definition

The **unit coordinate vectors** in \mathbb{R}^n are

This is what e_1, e_2, \dots mean,
for the rest of the class.

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}.$$



Note: if A is an $m \times n$ matrix with columns v_1, v_2, \dots, v_n , then $Ae_i = v_i$ for $i = 1, 2, \dots, n$: multiplying a matrix by e_i gives you the i th column.

$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 4 \\ 7 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ 5 \\ 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 6 \\ 9 \end{pmatrix}$$

Linear Transformations are Matrix Transformations

Recall: A matrix A defines a linear transformation T by $T(x) = Ax$.

Dr Ali and Dr Sara

Linear Transformations are Matrix Transformations

Recall: A matrix A defines a linear transformation T by $T(x) = Ax$.

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Let

$$A = \begin{pmatrix} \left| \begin{array}{c} T(e_1) \\ \vdots \end{array} \right| & \left| \begin{array}{c} T(e_2) \\ \vdots \end{array} \right| & \cdots & \left| \begin{array}{c} T(e_n) \\ \vdots \end{array} \right| \end{pmatrix}.$$

This is an _____ matrix,

Linear Transformations are Matrix Transformations

Recall: A matrix A defines a linear transformation T by $T(x) = Ax$.

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Let

$$A = \begin{pmatrix} | & | & & | \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ | & | & & | \end{pmatrix}.$$

This is an $m \times n$ matrix, and T is the matrix transformation for A : $T(x) = Ax$.

Linear Transformations are Matrix Transformations

Recall: A matrix A defines a linear transformation T by $T(x) = Ax$.

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Let

$$A = \begin{pmatrix} | & | & & | \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ | & | & & | \end{pmatrix}.$$

This is an $m \times n$ matrix, and T is the matrix transformation for A : $T(x) = Ax$.

The matrix A is called the **standard matrix** for T .

Linear Transformations are Matrix Transformations

Recall: A matrix A defines a linear transformation T by $T(x) = Ax$.

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Let

$$A = \begin{pmatrix} | & | & & | \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ | & | & & | \end{pmatrix}.$$

This is an $m \times n$ matrix, and T is the matrix transformation for A : $T(x) = Ax$.

The matrix A is called the **standard matrix** for T .

Take-Away

Linear transformations are the same as matrix transformations.

Linear Transformations are Matrix Transformations

Recall: A matrix A defines a linear transformation T by $T(x) = Ax$.

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Let

$$A = \begin{pmatrix} | & | & & | \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ | & | & & | \end{pmatrix}.$$

This is an $m \times n$ matrix, and T is the matrix transformation for A : $T(x) = Ax$.

The matrix A is called the **standard matrix** for T .

Take-Away

Linear transformations are the same as matrix transformations.

Dictionary

Linear Transformations are Matrix Transformations

Recall: A matrix A defines a linear transformation T by $T(x) = Ax$.

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Let

$$A = \begin{pmatrix} | & | & & | \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ | & | & & | \end{pmatrix}.$$

This is an $m \times n$ matrix, and T is the matrix transformation for A : $T(x) = Ax$.

The matrix A is called the **standard matrix** for T .

Take-Away

Linear transformations are the same as matrix transformations.

Dictionary

Linear transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ \rightsquigarrow $m \times n$ matrix $A = \begin{pmatrix} | & | & & | \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ | & | & & | \end{pmatrix}$

Linear Transformations are Matrix Transformations

Recall: A matrix A defines a linear transformation T by $T(x) = Ax$.

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Let

$$A = \begin{pmatrix} | & | & & | \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ | & | & & | \end{pmatrix}.$$

This is an $m \times n$ matrix, and T is the matrix transformation for A : $T(x) = Ax$.

The matrix A is called the **standard matrix** for T .

Take-Away

Linear transformations are the same as matrix transformations.

Dictionary

$$\begin{array}{ll} \text{Linear transformation} & \\ T: \mathbb{R}^n \rightarrow \mathbb{R}^m & \rightsquigarrow m \times n \text{ matrix } A = \begin{pmatrix} | & | & & | \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ | & | & & | \end{pmatrix} \\ T(x) = Ax & \\ T: \mathbb{R}^n \rightarrow \mathbb{R}^m & \longleftarrow m \times n \text{ matrix } A \end{array}$$

Linear Transformations are Matrix Transformations

Continued

Why is a linear transformation a matrix transformation?

Dr Ali and Dr Sara

Linear Transformations are Matrix Transformations

Continued

Why is a linear transformation a matrix transformation?

Suppose for simplicity that $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2$.

$$\begin{aligned} T \begin{pmatrix} x \\ y \\ z \end{pmatrix} &= T \left(x \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + y \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + z \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right) \\ &= T(xe_1 + ye_2 + ze_3) \\ &= xT(e_1) + yT(e_2) + zT(e_3) \\ &= \begin{pmatrix} | & | & | \\ T(e_1) & T(e_2) & T(e_3) \\ | & | & | \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} \\ &= A \begin{pmatrix} x \\ y \\ z \end{pmatrix}. \end{aligned}$$

Linear Transformations are Matrix Transformations

Example

Before, we defined a **dilation** transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $T(x) = 1.5x$.
What is its standard matrix?

$$\left. \begin{aligned} T(e_1) &= 1.5e_1 = \begin{pmatrix} 1.5 \\ 0 \end{pmatrix} \\ T(e_2) &= 1.5e_2 = \begin{pmatrix} 0 \\ 1.5 \end{pmatrix} \end{aligned} \right\} \Rightarrow A = \begin{pmatrix} 1.5 & 0 \\ 0 & 1.5 \end{pmatrix}.$$

Linear Transformations are Matrix Transformations

Example

Before, we defined a **dilation** transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $T(x) = 1.5x$.
What is its standard matrix?

$$\left. \begin{aligned} T(e_1) &= 1.5e_1 = \begin{pmatrix} 1.5 \\ 0 \end{pmatrix} \\ T(e_2) &= 1.5e_2 = \begin{pmatrix} 0 \\ 1.5 \end{pmatrix} \end{aligned} \right\} \Rightarrow A = \begin{pmatrix} 1.5 & 0 \\ 0 & 1.5 \end{pmatrix}.$$

Check:

$$\begin{pmatrix} 1.5 & 0 \\ 0 & 1.5 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} =$$

Linear Transformations are Matrix Transformations

Example

Before, we defined a **dilation** transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $T(x) = 1.5x$. What is its standard matrix?

$$\left. \begin{aligned} T(e_1) &= 1.5e_1 = \begin{pmatrix} 1.5 \\ 0 \end{pmatrix} \\ T(e_2) &= 1.5e_2 = \begin{pmatrix} 0 \\ 1.5 \end{pmatrix} \end{aligned} \right\} \Rightarrow A = \begin{pmatrix} 1.5 & 0 \\ 0 & 1.5 \end{pmatrix}.$$

Check:

$$\begin{pmatrix} 1.5 & 0 \\ 0 & 1.5 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1.5x \\ 1.5y \end{pmatrix} =$$

Linear Transformations are Matrix Transformations

Example

Before, we defined a **dilation** transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $T(x) = 1.5x$.
What is its standard matrix?

$$\left. \begin{aligned} T(e_1) &= 1.5e_1 = \begin{pmatrix} 1.5 \\ 0 \end{pmatrix} \\ T(e_2) &= 1.5e_2 = \begin{pmatrix} 0 \\ 1.5 \end{pmatrix} \end{aligned} \right\} \Rightarrow A = \begin{pmatrix} 1.5 & 0 \\ 0 & 1.5 \end{pmatrix}.$$

Check:

$$\begin{pmatrix} 1.5 & 0 \\ 0 & 1.5 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1.5x \\ 1.5y \end{pmatrix} = 1.5 \begin{pmatrix} x \\ y \end{pmatrix} = T \begin{pmatrix} x \\ y \end{pmatrix}.$$

Linear Transformations are Matrix Transformations

Example

Question


What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?

Linear Transformations are Matrix Transformations

Example

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




Linear Transformations are Matrix Transformations

Example

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




Linear Transformations are Matrix Transformations

Example

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




Linear Transformations are Matrix Transformations

Example

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




$$T(e_1) = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




$$T(e_2) = e_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}.$$

Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?



$$T(e_3) = \begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix}.$$

Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?

$$\left. \begin{aligned} T(e_1) &= \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \\ T(e_2) &= \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \\ T(e_3) &= \begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix} \end{aligned} \right\} \Rightarrow A =$$

Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?

$$\left. \begin{aligned} T(e_1) &= \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \\ T(e_2) &= \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \\ T(e_3) &= \begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix} \end{aligned} \right\} \Rightarrow A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

Onto Transformations

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **onto** (or **surjective**) if the range of T is equal to \mathbb{R}^m (its codomain).

Onto Transformations

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **onto** (or **surjective**) if the range of T is equal to \mathbb{R}^m (its codomain). In other words, each b in \mathbb{R}^m is the image of *at least one* x in \mathbb{R}^n :

Onto Transformations

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **onto** (or **surjective**) if the range of T is equal to \mathbb{R}^m (its codomain). In other words, each b in \mathbb{R}^m is the image of *at least one* x in \mathbb{R}^n : every possible output has an input.

Onto Transformations

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **onto** (or **surjective**) if the range of T is equal to \mathbb{R}^m (its codomain). In other words, each b in \mathbb{R}^m is the image of *at least one* x in \mathbb{R}^n : every possible output has an input. Note that *not* onto means

Onto Transformations


Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **onto** (or **surjective**) if the range of T is equal to \mathbb{R}^m (its codomain). In other words, each b in \mathbb{R}^m is the image of *at least one* x in \mathbb{R}^n : every possible output has an input. Note that *not* onto means there is some b in \mathbb{R}^m which is not the image of any x in \mathbb{R}^n .

Onto Transformations

Definition



A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **onto** (or **surjective**) if the range of T is equal to \mathbb{R}^m (its codomain). In other words, each b in \mathbb{R}^m is the image of *at least one* x in \mathbb{R}^n : every possible output has an input. Note that *not* onto means there is some b in \mathbb{R}^m which is not the image of any x in \mathbb{R}^n .



Onto Transformations

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **onto** (or **surjective**) if the range of T is equal to \mathbb{R}^m (its codomain). In other words, each b in \mathbb{R}^m is the image of *at least one* x in \mathbb{R}^n : every possible output has an input. Note that *not* onto means there is some b in \mathbb{R}^m which is not the image of any x in \mathbb{R}^n .



Characterization of Onto Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto

Characterization of Onto Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto
- ▶ $T(x) = b$ has a solution for every b in \mathbb{R}^m

Characterization of Onto Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto
- ▶ $T(x) = b$ has a solution for every b in \mathbb{R}^m
- ▶ $Ax = b$ is consistent for every b in \mathbb{R}^m

Characterization of Onto Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto
- ▶ $T(x) = b$ has a solution for every b in \mathbb{R}^m
- ▶ $Ax = b$ is consistent for every b in \mathbb{R}^m
- ▶ The columns of A span \mathbb{R}^m

Characterization of Onto Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto
- ▶ $T(x) = b$ has a solution for every b in \mathbb{R}^m
- ▶ $Ax = b$ is consistent for every b in \mathbb{R}^m
- ▶ The columns of A span \mathbb{R}^m
- ▶ A has a pivot in every ____

Characterization of Onto Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto
- ▶ $T(x) = b$ has a solution for every b in \mathbb{R}^m
- ▶ $Ax = b$ is consistent for every b in \mathbb{R}^m
- ▶ The columns of A span \mathbb{R}^m
- ▶ A has a pivot in every row

Characterization of Onto Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto
- ▶ $T(x) = b$ has a solution for every b in \mathbb{R}^m
- ▶ $Ax = b$ is consistent for every b in \mathbb{R}^m
- ▶ The columns of A span \mathbb{R}^m
- ▶ A has a pivot in every row

Question

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is onto, what can we say about the relative sizes of n and m ?

Characterization of Onto Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto
- ▶ $T(x) = b$ has a solution for every b in \mathbb{R}^m
- ▶ $Ax = b$ is consistent for every b in \mathbb{R}^m
- ▶ The columns of A span \mathbb{R}^m
- ▶ A has a pivot in every row

Question

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is onto, what can we say about the relative sizes of n and m ?

Answer: T corresponds to an _____ matrix A .

Characterization of Onto Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto
- ▶ $T(x) = b$ has a solution for every b in \mathbb{R}^m
- ▶ $Ax = b$ is consistent for every b in \mathbb{R}^m
- ▶ The columns of A span \mathbb{R}^m
- ▶ A has a pivot in every row

Question

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is onto, what can we say about the relative sizes of n and m ?

Answer: T corresponds to an $m \times n$ matrix A .

Characterization of Onto Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto
- ▶ $T(x) = b$ has a solution for every b in \mathbb{R}^m
- ▶ $Ax = b$ is consistent for every b in \mathbb{R}^m
- ▶ The columns of A span \mathbb{R}^m
- ▶ A has a pivot in every row

Question

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is onto, what can we say about the relative sizes of n and m ?

Answer: T corresponds to an $m \times n$ matrix A . In order for A to have a pivot in every row, it must have *at least as many* columns as rows: $m \leq n$.

Characterization of Onto Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto
- ▶ $T(x) = b$ has a solution for every b in \mathbb{R}^m
- ▶ $Ax = b$ is consistent for every b in \mathbb{R}^m
- ▶ The columns of A span \mathbb{R}^m
- ▶ A has a pivot in every row

Question

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is onto, what can we say about the relative sizes of n and m ?

Answer: T corresponds to an $m \times n$ matrix A . In order for A to have a pivot in every row, it must have *at least as many* columns as rows: $m \leq n$.

$$\begin{pmatrix} \color{red}{1} & 0 & \star & 0 & \star \\ 0 & \color{red}{1} & \star & 0 & \star \\ 0 & 0 & 0 & \color{red}{1} & \star \end{pmatrix}$$

Characterization of Onto Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto
- ▶ $T(x) = b$ has a solution for every b in \mathbb{R}^m
- ▶ $Ax = b$ is consistent for every b in \mathbb{R}^m
- ▶ The columns of A span \mathbb{R}^m
- ▶ A has a pivot in every row

Question

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is onto, what can we say about the relative sizes of n and m ?

Answer: T corresponds to an $m \times n$ matrix A . In order for A to have a pivot in every row, it must have *at least as many* columns as rows: $m \leq n$.

$$\begin{pmatrix} \color{red}{1} & 0 & \star & 0 & \star \\ 0 & \color{red}{1} & \star & 0 & \star \\ 0 & 0 & 0 & \color{red}{1} & \star \end{pmatrix}$$

For instance, \mathbb{R}^2 is “too small” to map *onto* \mathbb{R}^3 .

One-to-one Transformations

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **one-to-one** (or **into**, or **injective**) if different vectors in \mathbb{R}^n map to different vectors in \mathbb{R}^m .

One-to-one Transformations

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **one-to-one** (or **into**, or **injective**) if different vectors in \mathbb{R}^n map to different vectors in \mathbb{R}^m . In other words, each b in \mathbb{R}^m is the image of *at most one* x in \mathbb{R}^n :

One-to-one Transformations

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **one-to-one** (or **into**, or **injective**) if different vectors in \mathbb{R}^n map to different vectors in \mathbb{R}^m . In other words, each b in \mathbb{R}^m is the image of *at most one* x in \mathbb{R}^n : different inputs have different outputs.

One-to-one Transformations

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **one-to-one** (or **into**, or **injective**) if different vectors in \mathbb{R}^n map to different vectors in \mathbb{R}^m . In other words, each b in \mathbb{R}^m is the image of *at most one* x in \mathbb{R}^n : different inputs have different outputs. Note that *not* one-to-one means

One-to-one Transformations


Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **one-to-one** (or **into**, or **injective**) if different vectors in \mathbb{R}^n map to different vectors in \mathbb{R}^m . In other words, each b in \mathbb{R}^m is the image of *at most one* x in \mathbb{R}^n : different inputs have different outputs. Note that *not* one-to-one means different vectors in \mathbb{R}^n have the same image.

One-to-one Transformations

Definition


A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **one-to-one** (or **into**, or **injective**) if different vectors in \mathbb{R}^n map to different vectors in \mathbb{R}^m . In other words, each b in \mathbb{R}^m is the image of *at most one* x in \mathbb{R}^n : different inputs have different outputs. Note that *not* one-to-one means different vectors in \mathbb{R}^n have the same image.



One-to-one Transformations

Definition

A transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **one-to-one** (or **into**, or **injective**) if different vectors in \mathbb{R}^n map to different vectors in \mathbb{R}^m . In other words, each b in \mathbb{R}^m is the image of *at most one* x in \mathbb{R}^n : different inputs have different outputs. Note that *not* one-to-one means different vectors in \mathbb{R}^n have the same image.



Characterization of One-to-One Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one

Characterization of One-to-One Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one
- ▶ $T(x) = b$ has one or zero solutions for every b in \mathbb{R}^m

Characterization of One-to-One Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one
- ▶ $T(x) = b$ has one or zero solutions for every b in \mathbb{R}^m
- ▶ $Ax = b$ has a unique solution or is inconsistent for every b in \mathbb{R}^m

Characterization of One-to-One Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one
- ▶ $T(x) = b$ has one or zero solutions for every b in \mathbb{R}^m
- ▶ $Ax = b$ has a unique solution or is inconsistent for every b in \mathbb{R}^m
- ▶ $Ax = 0$ has a unique solution

Characterization of One-to-One Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one
- ▶ $T(x) = b$ has one or zero solutions for every b in \mathbb{R}^m
- ▶ $Ax = b$ has a unique solution or is inconsistent for every b in \mathbb{R}^m
- ▶ $Ax = 0$ has a unique solution
- ▶ The columns of A are linearly independent

Characterization of One-to-One Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one
- ▶ $T(x) = b$ has one or zero solutions for every b in \mathbb{R}^m
- ▶ $Ax = b$ has a unique solution or is inconsistent for every b in \mathbb{R}^m
- ▶ $Ax = 0$ has a unique solution
- ▶ The columns of A are linearly independent
- ▶ A has a pivot in every _____.

Characterization of One-to-One Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one
- ▶ $T(x) = b$ has one or zero solutions for every b in \mathbb{R}^m
- ▶ $Ax = b$ has a unique solution or is inconsistent for every b in \mathbb{R}^m
- ▶ $Ax = 0$ has a unique solution
- ▶ The columns of A are linearly independent
- ▶ A has a pivot in every column.

Characterization of One-to-One Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one
- ▶ $T(x) = b$ has one or zero solutions for every b in \mathbb{R}^m
- ▶ $Ax = b$ has a unique solution or is inconsistent for every b in \mathbb{R}^m
- ▶ $Ax = 0$ has a unique solution
- ▶ The columns of A are linearly independent
- ▶ A has a pivot in every column.

Question

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is one-to-one, what can we say about the relative sizes of n and m ?

Characterization of One-to-One Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one
- ▶ $T(x) = b$ has one or zero solutions for every b in \mathbb{R}^m
- ▶ $Ax = b$ has a unique solution or is inconsistent for every b in \mathbb{R}^m
- ▶ $Ax = 0$ has a unique solution
- ▶ The columns of A are linearly independent
- ▶ A has a pivot in every column.

Question

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is one-to-one, what can we say about the relative sizes of n and m ?

Answer: T corresponds to an $m \times n$ matrix A .

Characterization of One-to-One Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one
- ▶ $T(x) = b$ has one or zero solutions for every b in \mathbb{R}^m
- ▶ $Ax = b$ has a unique solution or is inconsistent for every b in \mathbb{R}^m
- ▶ $Ax = 0$ has a unique solution
- ▶ The columns of A are linearly independent
- ▶ A has a pivot in every column.

Question

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is one-to-one, what can we say about the relative sizes of n and m ?

Answer: T corresponds to an $m \times n$ matrix A . In order for A to have a pivot in every column, it must have *at least as many rows as columns*: $n \leq m$.

Characterization of One-to-One Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one
- ▶ $T(x) = b$ has one or zero solutions for every b in \mathbb{R}^m
- ▶ $Ax = b$ has a unique solution or is inconsistent for every b in \mathbb{R}^m
- ▶ $Ax = 0$ has a unique solution
- ▶ The columns of A are linearly independent
- ▶ A has a pivot in every column.

Question

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is one-to-one, what can we say about the relative sizes of n and m ?

Answer: T corresponds to an $m \times n$ matrix A . In order for A to have a pivot in every column, it must have *at least as many rows as columns*: $n \leq m$.

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

Characterization of One-to-One Transformations

Theorem

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one
- ▶ $T(x) = b$ has one or zero solutions for every b in \mathbb{R}^m
- ▶ $Ax = b$ has a unique solution or is inconsistent for every b in \mathbb{R}^m
- ▶ $Ax = 0$ has a unique solution
- ▶ The columns of A are linearly independent
- ▶ A has a pivot in every column.

Question

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is one-to-one, what can we say about the relative sizes of n and m ?

Answer: T corresponds to an $m \times n$ matrix A . In order for A to have a pivot in every column, it must have *at least as many rows as columns*: $n \leq m$.

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

For instance, \mathbb{R}^3 is “too big” to map *into* \mathbb{R}^2 .

Questions

- ▶ What matrix transforms $(1, 0)$ into $(2, 5)$ and transforms $(0, 1)$ to $(1, 3)$?
- ▶ What matrix transforms $(2, 5)$ to $(1, 0)$ and $(1, 3)$ to $(0, 1)$? Why does no matrix transform $(2, 6)$ to $(1, 0)$ and $(1, 3)$ to $(0, 1)$?
- ▶ What transformation take x_1 to Ax_1 and x_2 to Ax_2 .

$$x_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \rightarrow Ax_1 = \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix} \quad \text{and} \quad x_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \rightarrow Ax_2 = \begin{pmatrix} 4 \\ 6 \\ 8 \end{pmatrix}.$$

Means whenever we have transformation of two (independent) vectors in \mathbb{R}^2 we would be able to find transformation matrix.

- ▶ What matrix has the effect of rotating every vector through 90° and then projecting the result onto the x -axis?
- ▶ What matrix represents projection onto the x -axis followed by projection onto the y -axis?

Questions

- ▶ The matrix $A = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$ produces a stretching in the x direction. Draw the circle $x^2 + y^2 = 1$ and sketch around it the points $(2x, y)$ that result from multiplication by A . What shape is that curve?
- ▶ What 3×3 matrices represent the transformations that
 - (a) project every vector onto the x - y plane?
 - (b) reflect every vector through the x - y plane?
 - (c) rotate the x - y plane through 90° , leaving the z -axis alone?
 - (d) rotate the x - y plane, then x - z , then y - z , through 90° ?
 - (e) carry out the same three rotations, but each one through 180° ?
- ▶ Every straight line remains straight after a linear transformation. If z is halfway between x and y , show that Az is halfway between Ax and Ay .