

Chapter 5

Eigenvalues and Eigenvectors

Section 5.1

Eigenvectors and Eigenvalues

Eigenvectors and Eigenvalues

Let $T : V \rightarrow V$ be a linear transformation. Suppose for some scalar λ and $0 \neq v \in V$,

$$T(v) = \lambda v$$

Then v is an **eigenvector** for T and λ is its **eigenvalue**.

Definition

Let A be an $n \times n$ matrix.

Eigenvalues and eigenvectors are only for square matrices.

1. An **eigenvector** of A is a nonzero vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} .

Eigenvectors and Eigenvalues

Let $T : V \rightarrow V$ be a linear transformation. Suppose for some scalar λ and $0 \neq v \in V$,

$$T(v) = \lambda v$$

Then v is an **eigenvector** for T and λ is its **eigenvalue**.

Definition

Let A be an $n \times n$ matrix.

Eigenvalues and eigenvectors are only for square matrices.

1. An **eigenvector** of A is a *nonzero* vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} .

Eigenvectors and Eigenvalues

Let $T : V \rightarrow V$ be a linear transformation. Suppose for some scalar λ and $0 \neq v \in V$,

$$T(v) = \lambda v$$

Then v is an **eigenvector** for T and λ is its **eigenvalue**.

Definition

Let A be an $n \times n$ matrix.

Eigenvalues and eigenvectors are only for square matrices.

1. An **eigenvector** of A is a *nonzero* vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} . In other words, Av is a multiple of v .

Eigenvectors and Eigenvalues

Let $T : V \rightarrow V$ be a linear transformation. Suppose for some scalar λ and $0 \neq v \in V$,

$$T(v) = \lambda v$$

Then v is an **eigenvector** for T and λ is its **eigenvalue**.

Definition

Let A be an $n \times n$ matrix.

Eigenvalues and eigenvectors are only for square matrices.

1. An **eigenvector** of A is a *nonzero* vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} . In other words, Av is a multiple of v .
2. An **eigenvalue** of A is a number λ in \mathbf{R} such that the equation $Av = \lambda v$ has a nontrivial solution.

Eigenvectors and Eigenvalues

Let $T : V \rightarrow V$ be a linear transformation. Suppose for some scalar λ and $0 \neq v \in V$,

$$T(v) = \lambda v$$

Then v is an **eigenvector** for T and λ is its **eigenvalue**.

Definition

Let A be an $n \times n$ matrix.

Eigenvalues and eigenvectors are only for square matrices.

1. An **eigenvector** of A is a *nonzero* vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} . In other words, Av is a multiple of v .
2. An **eigenvalue** of A is a number λ in \mathbf{R} such that the equation $Av = \lambda v$ has a *nontrivial* solution.

Eigenvectors and Eigenvalues

Let $T : V \rightarrow V$ be a linear transformation. Suppose for some scalar λ and $0 \neq v \in V$,

$$T(v) = \lambda v$$

Then v is an **eigenvector** for T and λ is its **eigenvalue**.

Definition

Let A be an $n \times n$ matrix.

Eigenvalues and eigenvectors are only for square matrices.

1. An **eigenvector** of A is a *nonzero* vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} . In other words, Av is a multiple of v .
2. An **eigenvalue** of A is a number λ in \mathbf{R} such that the equation $Av = \lambda v$ has a *nontrivial* solution.

If $Av = \lambda v$ for $v \neq 0$, we say λ is the **eigenvalue for** v , and v is an **eigenvector for** λ .

Eigenvectors and Eigenvalues

Let $T : V \rightarrow V$ be a linear transformation. Suppose for some scalar λ and $0 \neq v \in V$,

$$T(v) = \lambda v$$

Then v is an **eigenvector** for T and λ is its **eigenvalue**.

Definition

Let A be an $n \times n$ matrix.

Eigenvalues and eigenvectors are only for square matrices.

1. An **eigenvector** of A is a *nonzero* vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} . In other words, Av is a multiple of v .
2. An **eigenvalue** of A is a number λ in \mathbf{R} such that the equation $Av = \lambda v$ has a *nontrivial* solution.

If $Av = \lambda v$ for $v \neq 0$, we say λ is the **eigenvalue for** v , and v is an **eigenvector for** λ .

Note: Eigenvectors are by definition nonzero. Eigenvalues may be equal to zero.

Eigenvectors and Eigenvalues

Let $T : V \rightarrow V$ be a linear transformation. Suppose for some scalar λ and $0 \neq v \in V$,

$$T(v) = \lambda v$$

Then v is an **eigenvector** for T and λ is its **eigenvalue**.

Definition

Let A be an $n \times n$ matrix.

Eigenvalues and eigenvectors are only for square matrices.

1. An **eigenvector** of A is a *nonzero* vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} . In other words, Av is a multiple of v .
2. An **eigenvalue** of A is a number λ in \mathbf{R} such that the equation $Av = \lambda v$ has a *nontrivial* solution.

If $Av = \lambda v$ for $v \neq 0$, we say λ is the **eigenvalue for** v , and v is an **eigenvector for** λ .

Note: Eigenvectors are by definition nonzero. Eigenvalues may be equal to zero.

Eigenvectors and Eigenvalues

Let $T : V \rightarrow V$ be a linear transformation. Suppose for some scalar λ and $0 \neq v \in V$,

$$T(v) = \lambda v$$

Then v is an **eigenvector** for T and λ is its **eigenvalue**.

Definition

Let A be an $n \times n$ matrix.

Eigenvalues and eigenvectors are only for square matrices.

1. An **eigenvector** of A is a *nonzero* vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} . In other words, Av is a multiple of v .
2. An **eigenvalue** of A is a number λ in \mathbf{R} such that the equation $Av = \lambda v$ has a *nontrivial* solution.

If $Av = \lambda v$ for $v \neq 0$, we say λ is the **eigenvalue for v** , and v is an **eigenvector for λ** .

Note: Eigenvectors are by definition nonzero. Eigenvalues may be equal to zero.

Eigenvectors and Eigenvalues

Let $T : V \rightarrow V$ be a linear transformation. Suppose for some scalar λ and $0 \neq v \in V$,

$$T(v) = \lambda v$$

This is the definition of an eigenvector and eigenvalue.

Definition

Let A be an $n \times n$ matrix.

Eigenvalues and eigenvectors are only for square matrices.

1. An **eigenvector** of A is a *nonzero* vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} . In other words, Av is a multiple of v .
2. An **eigenvalue** of A is a number λ in \mathbf{R} such that the equation $Av = \lambda v$ has a *nontrivial* solution.

If $Av = \lambda v$ for $v \neq 0$, we say λ is the **eigenvalue for** v , and v is an **eigenvector for** λ .

Note: Eigenvectors are by definition nonzero. Eigenvalues may be equal to zero.

Eigenvectors and Eigenvalues

Let $T : V \rightarrow V$ be a linear transformation. Suppose for some scalar λ and $0 \neq v \in V$,

$$T(v) = \lambda v$$

Definition

Let A be an $n \times n$ matrix.

Eigenvalues and eigenvectors are only for square matrices.

1. An **eigenvector** of A is a *nonzero* vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} . In other words, Av is a multiple of v .
2. An **eigenvalue** of A is a number λ in \mathbf{R} such that the equation $Av = \lambda v$ has a *nontrivial* solution.

If $Av = \lambda v$ for $v \neq 0$, we say λ is the **eigenvalue for** v , and v is an **eigenvector for** λ .

Note: Eigenvectors are by definition nonzero. Eigenvalues may be equal to zero.

This is the most important definition in the course.

Verifying Eigenvectors

Example

$$A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \quad v = \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix}$$

Verifying Eigenvectors

Example

$$A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \quad v = \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix}$$

Multiply:

$$Av = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix} = \begin{pmatrix} 32 \\ 8 \\ 2 \end{pmatrix} = 2v$$

Verifying Eigenvectors

Example

$$A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \quad v = \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix}$$

Multiply:

$$Av = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix} = \begin{pmatrix} 32 \\ 8 \\ 2 \end{pmatrix} = 2v$$

Hence v is an eigenvector of A , with eigenvalue $\lambda = 2$.

Verifying Eigenvectors

Example

$$A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \quad v = \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix}$$

Multiply:

$$Av = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix} = \begin{pmatrix} 32 \\ 8 \\ 2 \end{pmatrix} = 2v$$

Hence v is an eigenvector of A , with eigenvalue $\lambda = 2$.

Verifying Eigenvectors

Example

$$A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \quad v = \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix}$$

Multiply:

$$Av = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix} = \begin{pmatrix} 32 \\ 8 \\ 2 \end{pmatrix} = 2v$$

Hence v is an eigenvector of A , with eigenvalue $\lambda = 2$.

Example

$$A = \begin{pmatrix} 2 & 2 \\ -4 & 8 \end{pmatrix} \quad v = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Verifying Eigenvectors

Example

$$A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \quad v = \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix}$$

Multiply:

$$Av = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix} = \begin{pmatrix} 32 \\ 8 \\ 2 \end{pmatrix} = 2v$$

Hence v is an eigenvector of A , with eigenvalue $\lambda = 2$.

Example

$$A = \begin{pmatrix} 2 & 2 \\ -4 & 8 \end{pmatrix} \quad v = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Multiply:

$$Av = \begin{pmatrix} 2 & 2 \\ -4 & 8 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \end{pmatrix} = 4v$$

Verifying Eigenvectors

Example

$$A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \quad v = \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix}$$

Multiply:

$$Av = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix} = \begin{pmatrix} 32 \\ 8 \\ 2 \end{pmatrix} = 2v$$

Hence v is an eigenvector of A , with eigenvalue $\lambda = 2$.

Example

$$A = \begin{pmatrix} 2 & 2 \\ -4 & 8 \end{pmatrix} \quad v = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Multiply:

$$Av = \begin{pmatrix} 2 & 2 \\ -4 & 8 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \end{pmatrix} = 4v$$

Hence v is an eigenvector of A , with eigenvalue $\lambda = 4$.

Verifying Eigenvectors

Example

$$A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \quad v = \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix}$$

Multiply:

$$Av = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix} = \begin{pmatrix} 32 \\ 8 \\ 2 \end{pmatrix} = 2v$$

Hence v is an eigenvector of A , with eigenvalue $\lambda = 2$.

Example

$$A = \begin{pmatrix} 2 & 2 \\ -4 & 8 \end{pmatrix} \quad v = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Multiply:

$$Av = \begin{pmatrix} 2 & 2 \\ -4 & 8 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \end{pmatrix} = 4v$$

Hence v is an eigenvector of A , with eigenvalue $\lambda = 4$.

Poll

Which of the vectors

A. $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ B. $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ C. $\begin{pmatrix} -1 \\ 1 \end{pmatrix}$ D. $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$ E. $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$

are eigenvectors of the matrix $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$?

What are the eigenvalues?

Poll

Which of the vectors

A. $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ B. $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ C. $\begin{pmatrix} -1 \\ 1 \end{pmatrix}$ D. $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$ E. $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$

are eigenvectors of the matrix $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$?

What are the eigenvalues?

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

eigenvector with eigenvalue 2

Poll

Which of the vectors

A. $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ B. $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ C. $\begin{pmatrix} -1 \\ 1 \end{pmatrix}$ D. $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$ E. $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$

are eigenvectors of the matrix $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$?

What are the eigenvalues?

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

eigenvector with eigenvalue 2

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = 0 \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

eigenvector with eigenvalue 0

Poll

Which of the vectors

A. $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ B. $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ C. $\begin{pmatrix} -1 \\ 1 \end{pmatrix}$ D. $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$ E. $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$

are eigenvectors of the matrix $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$?

What are the eigenvalues?

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

eigenvector with eigenvalue 2

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = 0 \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

eigenvector with eigenvalue 0

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} -1 \\ 1 \end{pmatrix} = 0 \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

eigenvector with eigenvalue 0

Poll

Which of the vectors

A. $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ B. $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ C. $\begin{pmatrix} -1 \\ 1 \end{pmatrix}$ D. $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$ E. $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$

are eigenvectors of the matrix $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$?

What are the eigenvalues?

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

eigenvector with eigenvalue 2

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = 0 \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

eigenvector with eigenvalue 0

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} -1 \\ 1 \end{pmatrix} = 0 \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

eigenvector with eigenvalue 0

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \end{pmatrix}$$

not an eigenvector

Poll

Which of the vectors

A. $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ B. $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ C. $\begin{pmatrix} -1 \\ 1 \end{pmatrix}$ D. $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$ E. $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$

are eigenvectors of the matrix $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$?

What are the eigenvalues?

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

eigenvector with eigenvalue 2

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = 0 \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

eigenvector with eigenvalue 0

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} -1 \\ 1 \end{pmatrix} = 0 \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

eigenvector with eigenvalue 0

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \end{pmatrix}$$

not an eigenvector

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

is never an eigenvector

Verifying Eigenvalues

Question: Is $\lambda = 3$ an eigenvalue of $A = \begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix}$?

Verifying Eigenvalues

Question: Is $\lambda = 3$ an eigenvalue of $A = \begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix}$?

In other words, does $Av = 3v$ have a nontrivial solution?

Verifying Eigenvalues

Question: Is $\lambda = 3$ an eigenvalue of $A = \begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix}$?

In other words, does $Av = 3v$ have a nontrivial solution?

... does $Av - 3v = 0$ have a nontrivial solution?

Verifying Eigenvalues

Question: Is $\lambda = 3$ an eigenvalue of $A = \begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix}$?

In other words, does $Av = 3v$ have a nontrivial solution?

... does $Av - 3v = 0$ have a nontrivial solution?

... does $(A - 3I)v = 0$ have a nontrivial solution?

Verifying Eigenvalues

Question: Is $\lambda = 3$ an eigenvalue of $A = \begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix}$?

In other words, does $Av = 3v$ have a nontrivial solution?

... does $Av - 3v = 0$ have a nontrivial solution?

... does $(A - 3I)v = 0$ have a nontrivial solution?

We know how to answer that!

Verifying Eigenvalues

Question: Is $\lambda = 3$ an eigenvalue of $A = \begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix}$?

In other words, does $Av = 3v$ have a nontrivial solution?

... does $Av - 3v = 0$ have a nontrivial solution?

... does $(A - 3I)v = 0$ have a nontrivial solution?

We know how to answer that! Row reduction!

Verifying Eigenvalues

Question: Is $\lambda = 3$ an eigenvalue of $A = \begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix}$?

In other words, does $Av = 3v$ have a nontrivial solution?

... does $Av - 3v = 0$ have a nontrivial solution?

... does $(A - 3I)v = 0$ have a nontrivial solution?

We know how to answer that! Row reduction!

$$A - 3I = \begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix} - 3 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} -1 & -4 \\ -1 & -4 \end{pmatrix}$$

Row reduce:

$$\begin{pmatrix} -1 & -4 \\ -1 & -4 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 1 & 4 \\ 0 & 0 \end{pmatrix}$$

Parametric form: $x = -4y$; parametric vector form: $\begin{pmatrix} x \\ y \end{pmatrix} = y \begin{pmatrix} -4 \\ 1 \end{pmatrix}$.

Does there exist an eigenvector with eigenvalue $\lambda = 3$? Yes! Any nonzero multiple of $\begin{pmatrix} -4 \\ 1 \end{pmatrix}$. Check:

$$\begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix} \begin{pmatrix} -4 \\ 1 \end{pmatrix} = \begin{pmatrix} -12 \\ 3 \end{pmatrix} = 3 \begin{pmatrix} -4 \\ 1 \end{pmatrix}. \quad \checkmark$$

Eigenspaces

Definition

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . The λ -**eigenspace** of A is the set of all eigenvectors of A with eigenvalue λ , plus the zero vector:

$$\lambda\text{-eigenspace} = \{v \text{ in } \mathbf{R}^n \mid Av = \lambda v\}$$

Eigenspaces

Definition

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . The λ -**eigenspace** of A is the set of all eigenvectors of A with eigenvalue λ , plus the zero vector:

$$\begin{aligned}\lambda\text{-eigenspace} &= \{v \text{ in } \mathbf{R}^n \mid Av = \lambda v\} \\ &= \{v \text{ in } \mathbf{R}^n \mid (A - \lambda I)v = 0\}\end{aligned}$$

Eigenspaces

Definition

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . The λ -**eigenspace** of A is the set of all eigenvectors of A with eigenvalue λ , plus the zero vector:

$$\begin{aligned}\lambda\text{-eigenspace} &= \{v \text{ in } \mathbf{R}^n \mid Av = \lambda v\} \\ &= \{v \text{ in } \mathbf{R}^n \mid (A - \lambda I)v = 0\} \\ &= \text{Nul}(A - \lambda I).\end{aligned}$$

Eigenspaces

Definition

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . The λ -**eigenspace** of A is the set of all eigenvectors of A with eigenvalue λ , plus the zero vector:

$$\begin{aligned}\lambda\text{-eigenspace} &= \{v \text{ in } \mathbf{R}^n \mid Av = \lambda v\} \\ &= \{v \text{ in } \mathbf{R}^n \mid (A - \lambda I)v = 0\} \\ &= \text{Nul}(A - \lambda I).\end{aligned}$$

Since the λ -eigenspace is a null space, it is a *subspace* of \mathbf{R}^n .

Eigenspaces

Definition

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . The λ -**eigenspace** of A is the set of all eigenvectors of A with eigenvalue λ , plus the zero vector:

$$\begin{aligned}\lambda\text{-eigenspace} &= \{v \text{ in } \mathbf{R}^n \mid Av = \lambda v\} \\ &= \{v \text{ in } \mathbf{R}^n \mid (A - \lambda I)v = 0\} \\ &= \text{Nul}(A - \lambda I).\end{aligned}$$

Since the λ -eigenspace is a null space, it is a *subspace* of \mathbf{R}^n .

How do you find a basis for the λ -eigenspace?

Eigenspaces

Definition

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . The λ -**eigenspace** of A is the set of all eigenvectors of A with eigenvalue λ , plus the zero vector:

$$\begin{aligned}\lambda\text{-eigenspace} &= \{v \text{ in } \mathbf{R}^n \mid Av = \lambda v\} \\ &= \{v \text{ in } \mathbf{R}^n \mid (A - \lambda I)v = 0\} \\ &= \text{Nul}(A - \lambda I).\end{aligned}$$

Since the λ -eigenspace is a null space, it is a *subspace* of \mathbf{R}^n .

How do you find a basis for the λ -eigenspace? Parametric vector form!

Eigenspaces

Example

Find a basis for the 2-eigenspace of

$$A = \begin{pmatrix} 4 & -1 & 6 \\ 2 & 1 & 6 \\ 2 & -1 & 8 \end{pmatrix}.$$

Eigenspaces

Example

Find a basis for the 2-eigenspace of

λ 

$$A = \begin{pmatrix} 4 & -1 & 6 \\ 2 & 1 & 6 \\ 2 & -1 & 8 \end{pmatrix}.$$

$$A - 2I = \begin{pmatrix} 2 & -1 & 6 \\ 2 & -1 & 6 \\ 2 & -1 & 6 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & -\frac{1}{2} & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\begin{array}{c} \text{parametric} \\ \text{form} \end{array} \xrightarrow{\text{~~~~~}} x = \frac{1}{2}y - 3z$$

$$\begin{array}{c} \text{parametric vector} \\ \text{form} \end{array} \xrightarrow{\text{~~~~~}} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = y \begin{pmatrix} \frac{1}{2} \\ 1 \\ 0 \end{pmatrix} + z \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix}$$

$$\begin{array}{c} \text{basis} \end{array} \xrightarrow{\text{~~~~~}} \left\{ \begin{pmatrix} \frac{1}{2} \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix} \right\}$$

Eigenspaces

Picture

A basis for the 2-eigenspace of $\begin{pmatrix} 4 & -1 & 6 \\ 2 & 1 & 6 \\ 2 & -1 & 8 \end{pmatrix}$ is $\left\{ \begin{pmatrix} \frac{1}{2} \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix} \right\}$.

Eigenspaces


Picture

A basis for the 2-eigenspace of $\begin{pmatrix} 4 & -1 & 6 \\ 2 & 1 & 6 \\ 2 & -1 & 8 \end{pmatrix}$ is $\left\{ \begin{pmatrix} \frac{1}{2} \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix} \right\}$. What does this look like?

Eigenspaces

Picture


A basis for the 2-eigenspace of $\begin{pmatrix} 4 & -1 & 6 \\ 2 & 1 & 6 \\ 2 & -1 & 8 \end{pmatrix}$ is $\left\{ \begin{pmatrix} \frac{1}{2} \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix} \right\}$. What does this look like?



Eigenspaces

Picture

A basis for the 2-eigenspace of $\begin{pmatrix} 4 & -1 & 6 \\ 2 & 1 & 6 \\ 2 & -1 & 8 \end{pmatrix}$ is $\left\{ \begin{pmatrix} \frac{1}{2} \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix} \right\}$. What does this look like?




For any \mathbf{v} in the 2-eigenspace,

Eigenspaces

Picture

A basis for the 2-eigenspace of $\begin{pmatrix} 4 & -1 & 6 \\ 2 & 1 & 6 \\ 2 & -1 & 8 \end{pmatrix}$ is $\left\{ \begin{pmatrix} \frac{1}{2} \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix} \right\}$. What does this look like?




For any v in the 2-eigenspace, $Av = 2v$ by definition.

Eigenspaces

Picture

A basis for the 2-eigenspace of $\begin{pmatrix} 4 & -1 & 6 \\ 2 & 1 & 6 \\ 2 & -1 & 8 \end{pmatrix}$ is $\left\{ \begin{pmatrix} \frac{1}{2} \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix} \right\}$. What does this look like?




For any v in the 2-eigenspace, $Av = 2v$ by definition. So A acts by *scaling* by 2 on its 2-eigenspace.

Eigenspaces

Picture

A basis for the 2-eigenspace of $\begin{pmatrix} 4 & -1 & 6 \\ 2 & 1 & 6 \\ 2 & -1 & 8 \end{pmatrix}$ is $\left\{ \begin{pmatrix} \frac{1}{2} \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix} \right\}$. What does this look like?



For any v in the 2-eigenspace, $Av = 2v$ by definition. So A acts by *scaling by 2* on its 2-eigenspace. This is how eigenvalues and eigenvectors make matrices easier to understand.

Eigenspaces

Geometry

Eigenvectors, geometrically

An eigenvector of a matrix A is a nonzero vector v such that:

Eigenspaces

Geometry

Eigenvectors, geometrically

An eigenvector of a matrix A is a nonzero vector v such that:

- ▶ Av is a multiple of v , which means

Eigenspaces

Geometry

Eigenvectors, geometrically

An eigenvector of a matrix A is a nonzero vector v such that:

- ▶ Av is a multiple of v , which means
- ▶ Av is collinear with v , which means

Eigenspaces

Geometry

Eigenvectors, geometrically

An eigenvector of a matrix A is a nonzero vector v such that:

- ▶ Av is a multiple of v , which means
- ▶ Av is collinear with v , which means
- ▶ Av and v are *on the same line*.


Eigenspaces

Geometry

Eigenvectors, geometrically

An eigenvector of a matrix A is a nonzero vector v such that:

- ▶ Av is a multiple of v , which means
- ▶ Av is collinear with v , which means
- ▶ Av and v are *on the same line*.



v is an eigenvector

w is not an eigenvector

Eigenspaces

Geometry; example


Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!




Does anyone see any eigenvectors
(vectors that don't move off their line)?

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors
(vectors that don't move off their line)?


v is an eigenvector with eigenvalue ____.

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors (vectors that don't move off their line)?


v is an eigenvector with eigenvalue -1 .

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors
(vectors that don't move off their line)?


w is an eigenvector with eigenvalue _.

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors
(vectors that don't move off their line)?


w is an eigenvector with eigenvalue 1.

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors
(vectors that don't move off their line)?


u is *not* an eigenvector.

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors
(vectors that don't move off their line)?


Neither is z .

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors
(vectors that don't move off their line)?


The 1-eigenspace is _

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors
(vectors that don't move off their line)?


The 1-eigenspace is L
(all the vectors x where $Ax = x$).

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors
(vectors that don't move off their line)?


The (-1) -eigenspace is _____

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors (vectors that don't move off their line)?

The (-1) -eigenspace is **the line $y = x$** (all the vectors x where $Ax = -x$).

Eigenspaces

Summary

Let A be an $n \times n$ matrix and let λ be a number.

Eigenspaces

Summary

Let A be an $n \times n$ matrix and let λ be a number.

1. λ is an eigenvalue of A if and only if $(A - \lambda I)x = 0$ has a nontrivial solution, if and only if $\text{Nul}(A - \lambda I) \neq \{0\}$.

Eigenspaces

Summary

Let A be an $n \times n$ matrix and let λ be a number.

1. λ is an eigenvalue of A if and only if $(A - \lambda I)x = 0$ has a nontrivial solution, if and only if $\text{Nul}(A - \lambda I) \neq \{0\}$.
2. In this case, finding a basis for the λ -eigenspace of A means finding a basis for $\text{Nul}(A - \lambda I)$ as usual, i.e. by finding the parametric vector form for the general solution to $(A - \lambda I)x = 0$.

Eigenspaces

Summary

Let A be an $n \times n$ matrix and let λ be a number.

1. λ is an eigenvalue of A if and only if $(A - \lambda I)x = 0$ has a nontrivial solution, if and only if $\text{Nul}(A - \lambda I) \neq \{0\}$.
2. In this case, finding a basis for the λ -eigenspace of A means finding a basis for $\text{Nul}(A - \lambda I)$ as usual, i.e. by finding the parametric vector form for the general solution to $(A - \lambda I)x = 0$.
3. The eigenvectors with eigenvalue λ are the nonzero elements of $\text{Nul}(A - \lambda I)$, i.e. the nontrivial solutions to $(A - \lambda I)x = 0$.

The Eigenvalues of a Triangular Matrix are the Diagonal Entries

We've seen that finding eigenvectors for a given eigenvalue is a row reduction problem.

The Eigenvalues of a Triangular Matrix are the Diagonal Entries

We've seen that finding eigenvectors for a given eigenvalue is a row reduction problem.

Finding all of the eigenvalues of a matrix *is not a row reduction problem!*

The Eigenvalues of a Triangular Matrix are the Diagonal Entries

We've seen that finding eigenvectors for a given eigenvalue is a row reduction problem.

Finding all of the eigenvalues of a matrix *is not a row reduction problem!* We'll see how to do it in general next time.

The Eigenvalues of a Triangular Matrix are the Diagonal Entries

We've seen that finding eigenvectors for a given eigenvalue is a row reduction problem.

Finding all of the eigenvalues of a matrix *is not a row reduction problem!* We'll see how to do it in general next time. For now:

Fact: The eigenvalues of a triangular matrix are the diagonal entries.

Why? $\text{Nul}(A - \lambda I) \neq \{0\}$ if and only if $A - \lambda I$ is not invertible, if and only if $\det(A - \lambda I) = 0$.

$$\begin{pmatrix} 3 & 4 & 1 & 2 \\ 0 & -1 & -2 & 7 \\ 0 & 0 & 8 & 12 \\ 0 & 0 & 0 & -3 \end{pmatrix} - \lambda I_4 = \begin{pmatrix} 3 - \lambda & 4 & 1 & 2 \\ 0 & -1 - \lambda & -2 & 7 \\ 0 & 0 & 8 - \lambda & 12 \\ 0 & 0 & 0 & -3 - \lambda \end{pmatrix}.$$

The determinant is $(3 - \lambda)(-1 - \lambda)(8 - \lambda)(-3 - \lambda)$, which is zero exactly when $\lambda = 3, -1, 8$, or -3 .

A Matrix is Invertible if and only if Zero is not an Eigenvalue

Fact: A is invertible if and only if 0 is not an eigenvalue of A .

Why?

0 is an eigenvalue of $A \iff Ax = 0x$ has a nontrivial solution

$\iff Ax = 0$ has a nontrivial solution

$\iff A$ is not invertible.

invertible matrix theorem



Eigenvectors with Distinct Eigenvalues are Linearly Independent

Fact: If v_1, v_2, \dots, v_k are eigenvectors of A with *distinct* eigenvalues $\lambda_1, \dots, \lambda_k$, then $\{v_1, v_2, \dots, v_k\}$ is linearly independent.

Eigenvectors with Distinct Eigenvalues are Linearly Independent

Fact: If v_1, v_2, \dots, v_k are eigenvectors of A with *distinct* eigenvalues $\lambda_1, \dots, \lambda_k$, then $\{v_1, v_2, \dots, v_k\}$ is linearly independent.

Why? If $k = 2$, this says v_2 can't lie on the line through v_1 .

Eigenvectors with Distinct Eigenvalues are Linearly Independent

Fact: If v_1, v_2, \dots, v_k are eigenvectors of A with *distinct* eigenvalues $\lambda_1, \dots, \lambda_k$, then $\{v_1, v_2, \dots, v_k\}$ is linearly independent.

Why? If $k = 2$, this says v_2 can't lie on the line through v_1 .

But the line through v_1 is contained in the λ_1 -eigenspace, and v_2 does not have eigenvalue λ_1 .

Eigenvectors with Distinct Eigenvalues are Linearly Independent

Fact: If v_1, v_2, \dots, v_k are eigenvectors of A with *distinct* eigenvalues $\lambda_1, \dots, \lambda_k$, then $\{v_1, v_2, \dots, v_k\}$ is linearly independent.

Why? If $k = 2$, this says v_2 can't lie on the line through v_1 .

But the line through v_1 is contained in the λ_1 -eigenspace, and v_2 does not have eigenvalue λ_1 .

In general: see Lay, Theorem 2 in §5.1 (or work it out for yourself; it's not too hard).

Eigenvectors with Distinct Eigenvalues are Linearly Independent

Fact: If v_1, v_2, \dots, v_k are eigenvectors of A with *distinct* eigenvalues $\lambda_1, \dots, \lambda_k$, then $\{v_1, v_2, \dots, v_k\}$ is linearly independent.

Why? If $k = 2$, this says v_2 can't lie on the line through v_1 .

But the line through v_1 is contained in the λ_1 -eigenspace, and v_2 does not have eigenvalue λ_1 .

In general: see Lay, Theorem 2 in §5.1 (or work it out for yourself; it's not too hard).

Consequence: An $n \times n$ matrix has at most n distinct eigenvalues.

Eigenvectors with Distinct Eigenvalues are Linearly Independent

Fact: If v_1, v_2, \dots, v_k are eigenvectors of A with *distinct* eigenvalues $\lambda_1, \dots, \lambda_k$, then $\{v_1, v_2, \dots, v_k\}$ is linearly independent.

Why? If $k = 2$, this says v_2 can't lie on the line through v_1 .

But the line through v_1 is contained in the λ_1 -eigenspace, and v_2 does not have eigenvalue λ_1 .

In general: see Lay, Theorem 2 in §5.1 (or work it out for yourself; it's not too hard).

Consequence: An $n \times n$ matrix has at most n distinct eigenvalues.

Section 5.2

The Characteristic Equation

The Invertible Matrix Theorem

Addenda

We have a couple of new ways of saying “ A is invertible” now:

The Invertible Matrix Theorem

Addenda

We have a couple of new ways of saying “ A is invertible” now:

The Invertible Matrix Theorem

Let A be a square $n \times n$ matrix, and let $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$ be the linear transformation $T(x) = Ax$. The following statements are equivalent.

1. A is invertible.

The Invertible Matrix Theorem

Addenda

We have a couple of new ways of saying “ A is invertible” now:

The Invertible Matrix Theorem

Let A be a square $n \times n$ matrix, and let $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$ be the linear transformation $T(x) = Ax$. The following statements are equivalent.

1. A is invertible.
2. T is invertible.
3. A is row equivalent to I_n .
4. A has n pivots.
5. $Ax = 0$ has only the trivial solution.
6. The columns of A are linearly independent.
7. T is one-to-one.
8. $Ax = b$ is consistent for all b in \mathbf{R}^n .
9. The columns of A span \mathbf{R}^n .
10. T is onto.
11. A has a left inverse (there exists B such that $BA = I_n$).
12. A has a right inverse (there exists B such that $AB = I_n$).
13. A^T is invertible.
14. The columns of A form a basis for \mathbf{R}^n .
15. $\text{Col } A = \mathbf{R}^n$.
16. $\dim \text{Col } A = n$.
17. $\text{rank } A = n$.
18. $\text{Nul } A = \{0\}$.
19. $\dim \text{Nul } A = 0$.

The Invertible Matrix Theorem

Addenda

We have a couple of new ways of saying “ A is invertible” now:

The Invertible Matrix Theorem

Let A be a square $n \times n$ matrix, and let $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$ be the linear transformation $T(x) = Ax$. The following statements are equivalent.

1. A is invertible.

2. T is invertible.

3. A is row equivalent to I_n .

4. A has n pivots.

5. $Ax = 0$ has only the trivial solution.

6. The columns of A are linearly independent.

7. T is one-to-one.

8. $Ax = b$ is consistent for all b in \mathbf{R}^n .

9. The columns of A span \mathbf{R}^n .

10. T is onto.

11. A has a left inverse (there exists B such that $BA = I_n$).

12. A has a right inverse (there exists B such that $AB = I_n$).

13. A^T is invertible.

14. The columns of A form a basis for \mathbf{R}^n .

15. $\text{Col } A = \mathbf{R}^n$.

16. $\dim \text{Col } A = n$.

17. $\text{rank } A = n$.

18. $\text{Nul } A = \{0\}$.

19. $\dim \text{Nul } A = 0$.

19. The determinant of A is *not* equal to zero.

The Invertible Matrix Theorem

Addenda

We have a couple of new ways of saying “ A is invertible” now:

The Invertible Matrix Theorem

Let A be a square $n \times n$ matrix, and let $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$ be the linear transformation $T(x) = Ax$. The following statements are equivalent.

1. A is invertible.
2. T is invertible.
3. A is row equivalent to I_n .
4. A has n pivots.
5. $Ax = 0$ has only the trivial solution.
6. The columns of A are linearly independent.
7. T is one-to-one.
8. $Ax = b$ is consistent for all b in \mathbf{R}^n .
9. The columns of A span \mathbf{R}^n .
10. T is onto.
11. A has a left inverse (there exists B such that $BA = I_n$).
12. A has a right inverse (there exists B such that $AB = I_n$).
13. A^T is invertible.
14. The columns of A form a basis for \mathbf{R}^n .
15. $\text{Col } A = \mathbf{R}^n$.
16. $\dim \text{Col } A = n$.
17. $\text{rank } A = n$.
18. $\text{Nul } A = \{0\}$.
19. $\dim \text{Nul } A = 0$.
19. The determinant of A is *not* equal to zero.
20. The number 0 is *not* an eigenvalue of A .

The Characteristic Polynomial

Let A be a square matrix.

λ is an eigenvalue of $A \iff$

The Characteristic Polynomial

Let A be a square matrix.

λ is an eigenvalue of $A \iff Ax = \lambda x$ has a nontrivial solution

The Characteristic Polynomial

Let A be a square matrix.

λ is an eigenvalue of $A \iff Ax = \lambda x$ has a nontrivial solution

$\iff (A - \lambda I)x = 0$ has a nontrivial solution

The Characteristic Polynomial

Let A be a square matrix.

λ is an eigenvalue of $A \iff Ax = \lambda x$ has a nontrivial solution

$\iff (A - \lambda I)x = 0$ has a nontrivial solution

$\iff A - \lambda I$ is not invertible

The Characteristic Polynomial

Let A be a square matrix.

λ is an eigenvalue of $A \iff Ax = \lambda x$ has a nontrivial solution

$\iff (A - \lambda I)x = 0$ has a nontrivial solution

$\iff A - \lambda I$ is not invertible

$\iff \det(A - \lambda I) = 0.$

The Characteristic Polynomial

Let A be a square matrix.

λ is an eigenvalue of $A \iff Ax = \lambda x$ has a nontrivial solution

$\iff (A - \lambda I)x = 0$ has a nontrivial solution

$\iff A - \lambda I$ is not invertible

$\iff \det(A - \lambda I) = 0.$

This gives us a way to compute the eigenvalues of A .

The Characteristic Polynomial

Let A be a square matrix.

λ is an eigenvalue of $A \iff Ax = \lambda x$ has a nontrivial solution

$\iff (A - \lambda I)x = 0$ has a nontrivial solution

$\iff A - \lambda I$ is not invertible

$\iff \det(A - \lambda I) = 0$.

This gives us a way to compute the eigenvalues of A .

Definition

Let A be a square matrix. The **characteristic polynomial** of A is

$$f(\lambda) = \det(A - \lambda I).$$

The Characteristic Polynomial

Let A be a square matrix.

$$\begin{aligned}\lambda \text{ is an eigenvalue of } A &\iff Ax = \lambda x \text{ has a nontrivial solution} \\ &\iff (A - \lambda I)x = 0 \text{ has a nontrivial solution} \\ &\iff A - \lambda I \text{ is not invertible} \\ &\iff \det(A - \lambda I) = 0.\end{aligned}$$

This gives us a way to compute the eigenvalues of A .

Definition

Let A be a square matrix. The **characteristic polynomial** of A is

$$f(\lambda) = \det(A - \lambda I).$$

The **characteristic equation** of A is the equation

$$f(\lambda) = \det(A - \lambda I) = 0.$$

The Characteristic Polynomial

Let A be a square matrix.

$$\begin{aligned}\lambda \text{ is an eigenvalue of } A &\iff Ax = \lambda x \text{ has a nontrivial solution} \\ &\iff (A - \lambda I)x = 0 \text{ has a nontrivial solution} \\ &\iff A - \lambda I \text{ is not invertible} \\ &\iff \det(A - \lambda I) = 0.\end{aligned}$$

This gives us a way to compute the eigenvalues of A .

Definition

Let A be a square matrix. The **characteristic polynomial** of A is

$$f(\lambda) = \det(A - \lambda I).$$

The **characteristic equation** of A is the equation

$$f(\lambda) = \det(A - \lambda I) = 0.$$

Important

The eigenvalues of A are the roots of the characteristic polynomial $f(\lambda) = \det(A - \lambda I)$.

The Characteristic Polynomial

Example

Question: What are the eigenvalues of

$$A = \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}?$$

Answer: First we find the characteristic polynomial:

$$\begin{aligned} f(\lambda) &= \det(A - \lambda I) = \det \left[\begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix} - \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} \right] = \det \begin{pmatrix} 5 - \lambda & 2 \\ 2 & 1 - \lambda \end{pmatrix} \\ &= (5 - \lambda)(1 - \lambda) - 2 \cdot 2 \\ &= \lambda^2 - 6\lambda + 1. \end{aligned}$$

The eigenvalues are the roots of the characteristic polynomial, which we can find using the quadratic formula:

$$\lambda = \frac{6 \pm \sqrt{36 - 4}}{2} = 3 \pm 2\sqrt{2}.$$

The Characteristic Polynomial

Example

Question: What is the characteristic polynomial of

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}?$$

Answer:

$$\begin{aligned} f(\lambda) &= \det(A - \lambda I) = \det \begin{pmatrix} a - \lambda & b \\ c & d - \lambda \end{pmatrix} = (a - \lambda)(d - \lambda) - bc \\ &= \lambda^2 - (a + d)\lambda + (ad - bc) \end{aligned}$$

The Characteristic Polynomial

Example

Question: What is the characteristic polynomial of

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}?$$

Answer:

$$\begin{aligned} f(\lambda) &= \det(A - \lambda I) = \det \begin{pmatrix} a - \lambda & b \\ c & d - \lambda \end{pmatrix} = (a - \lambda)(d - \lambda) - bc \\ &= \lambda^2 - (a + d)\lambda + (ad - bc) \end{aligned}$$

What do you notice about $f(\lambda)$?

The Characteristic Polynomial

Example

Question: What is the characteristic polynomial of

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}?$$

Answer:

$$\begin{aligned} f(\lambda) &= \det(A - \lambda I) = \det \begin{pmatrix} a - \lambda & b \\ c & d - \lambda \end{pmatrix} = (a - \lambda)(d - \lambda) - bc \\ &= \lambda^2 - (a + d)\lambda + (ad - bc) \end{aligned}$$

What do you notice about $f(\lambda)$?

- The constant term is $\det(A)$, which is zero if and only if

The Characteristic Polynomial

Example

Question: What is the characteristic polynomial of

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}?$$

Answer:

$$\begin{aligned} f(\lambda) &= \det(A - \lambda I) = \det \begin{pmatrix} a - \lambda & b \\ c & d - \lambda \end{pmatrix} = (a - \lambda)(d - \lambda) - bc \\ &= \lambda^2 - (a + d)\lambda + (ad - bc) \end{aligned}$$

What do you notice about $f(\lambda)$?

- The constant term is $\det(A)$, which is zero if and only if $\lambda = 0$ is a root.

The Characteristic Polynomial

Example

Question: What is the characteristic polynomial of

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}?$$

Answer:

$$\begin{aligned} f(\lambda) &= \det(A - \lambda I) = \det \begin{pmatrix} a - \lambda & b \\ c & d - \lambda \end{pmatrix} = (a - \lambda)(d - \lambda) - bc \\ &= \lambda^2 - (a + d)\lambda + (ad - bc) \end{aligned}$$

What do you notice about $f(\lambda)$?

- ▶ The constant term is $\det(A)$, which is zero if and only if $\lambda = 0$ is a root.
- ▶ The linear term $-(a + d)$ is the negative of the sum of the diagonal entries of A .

The Characteristic Polynomial

Example

Question: What is the characteristic polynomial of

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}?$$

Answer:

$$\begin{aligned} f(\lambda) &= \det(A - \lambda I) = \det \begin{pmatrix} a - \lambda & b \\ c & d - \lambda \end{pmatrix} = (a - \lambda)(d - \lambda) - bc \\ &= \lambda^2 - (a + d)\lambda + (ad - bc) \end{aligned}$$

What do you notice about $f(\lambda)$?

- ▶ The constant term is $\det(A)$, which is zero if and only if $\lambda = 0$ is a root.
- ▶ The linear term $-(a + d)$ is the negative of the sum of the diagonal entries of A .

Definition

The **trace** of a square matrix A is $\text{Tr}(A) = \text{sum of the diagonal entries of } A$.

The Characteristic Polynomial

Example

Question: What is the characteristic polynomial of

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}?$$

Answer:

$$\begin{aligned} f(\lambda) &= \det(A - \lambda I) = \det \begin{pmatrix} a - \lambda & b \\ c & d - \lambda \end{pmatrix} = (a - \lambda)(d - \lambda) - bc \\ &= \lambda^2 - (a + d)\lambda + (ad - bc) \end{aligned}$$

What do you notice about $f(\lambda)$?

- ▶ The constant term is $\det(A)$, which is zero if and only if $\lambda = 0$ is a root.
- ▶ The linear term $-(a + d)$ is the negative of the sum of the diagonal entries of A .

Definition

The **trace** of a square matrix A is $\text{Tr}(A)$ = sum of the diagonal entries of A .

Shortcut

The characteristic polynomial of a 2×2 matrix A is

$$f(\lambda) = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

The Characteristic Polynomial

Example

Question: What are the eigenvalues of the following matrix

$$A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix}?$$

Answer: First we find the characteristic polynomial:

$$\begin{aligned} f(\lambda) &= \det(A - \lambda I) = \det \begin{pmatrix} -\lambda & 6 & 8 \\ \frac{1}{2} & -\lambda & 0 \\ 0 & \frac{1}{2} & -\lambda \end{pmatrix} \\ &= 8 \left(\frac{1}{4} - 0 \cdot -\lambda \right) - \lambda \left(\lambda^2 - 6 \cdot \frac{1}{2} \right) \\ &= -\lambda^3 + 3\lambda + 2. \end{aligned}$$

We know from before that one eigenvalue is $\lambda = 2$: indeed, $f(2) = -8 + 6 + 2 = 0$. Doing polynomial long division, we get:

$$\frac{-\lambda^3 + 3\lambda + 2}{\lambda - 2} = -\lambda^2 - 2\lambda - 1 = -(\lambda + 1)^2.$$

Hence $\lambda = -1$ is also an eigenvalue.

Algebraic Multiplicity

Definition

The **(algebraic) multiplicity** of an eigenvalue λ is its multiplicity as a root of the characteristic polynomial.

Algebraic Multiplicity

Definition

The **(algebraic) multiplicity** of an eigenvalue λ is its multiplicity as a root of the characteristic polynomial.

This is not a very interesting notion *yet*.

Algebraic Multiplicity

Definition

The **(algebraic) multiplicity** of an eigenvalue λ is its multiplicity as a root of the characteristic polynomial.

This is not a very interesting notion *yet*. It will become interesting when we also define *geometric* multiplicity later.

Algebraic Multiplicity

Definition

The **(algebraic) multiplicity** of an eigenvalue λ is its multiplicity as a root of the characteristic polynomial.

This is not a very interesting notion *yet*. It will become interesting when we also define *geometric* multiplicity later.

Example

In the previous example, $f(\lambda) = -(\lambda - 2)(\lambda + 1)^2$, so the algebraic multiplicity of the eigenvalue 2 is ,

Algebraic Multiplicity

Definition

The **(algebraic) multiplicity** of an eigenvalue λ is its multiplicity as a root of the characteristic polynomial.

This is not a very interesting notion *yet*. It will become interesting when we also define *geometric* multiplicity later.

Example

In the previous example, $f(\lambda) = -(\lambda - 2)(\lambda + 1)^2$, so the algebraic multiplicity of the eigenvalue 2 is 1,

Algebraic Multiplicity

Definition

The **(algebraic) multiplicity** of an eigenvalue λ is its multiplicity as a root of the characteristic polynomial.

This is not a very interesting notion *yet*. It will become interesting when we also define *geometric* multiplicity later.

Example

In the previous example, $f(\lambda) = -(\lambda - 2)(\lambda + 1)^2$, so the algebraic multiplicity of the eigenvalue 2 is 1, and the algebraic multiplicity of the eigenvalue -1 is .

Algebraic Multiplicity

Definition

The **(algebraic) multiplicity** of an eigenvalue λ is its multiplicity as a root of the characteristic polynomial.

This is not a very interesting notion *yet*. It will become interesting when we also define *geometric* multiplicity later.

Example

In the previous example, $f(\lambda) = -(\lambda - 2)(\lambda + 1)^2$, so the algebraic multiplicity of the eigenvalue 2 is 1, and the algebraic multiplicity of the eigenvalue -1 is 2.

Algebraic Multiplicity

Definition

The **(algebraic) multiplicity** of an eigenvalue λ is its multiplicity as a root of the characteristic polynomial.

This is not a very interesting notion *yet*. It will become interesting when we also define *geometric* multiplicity later.

Example

In the previous example, $f(\lambda) = -(\lambda - 2)(\lambda + 1)^2$, so the algebraic multiplicity of the eigenvalue 2 is 1, and the algebraic multiplicity of the eigenvalue -1 is 2.

Example

In the matrix $\begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$, $f(\lambda) = (\lambda - (3 - 2\sqrt{2}))(\lambda - (3 + 2\sqrt{2}))$, so the algebraic multiplicity of $3 + 2\sqrt{2}$ is $_$,

Algebraic Multiplicity

Definition

The **(algebraic) multiplicity** of an eigenvalue λ is its multiplicity as a root of the characteristic polynomial.

This is not a very interesting notion *yet*. It will become interesting when we also define *geometric* multiplicity later.

Example

In the previous example, $f(\lambda) = -(\lambda - 2)(\lambda + 1)^2$, so the algebraic multiplicity of the eigenvalue 2 is 1, and the algebraic multiplicity of the eigenvalue -1 is 2.

Example

In the matrix $\begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$, $f(\lambda) = (\lambda - (3 - 2\sqrt{2}))(\lambda - (3 + 2\sqrt{2}))$, so the algebraic multiplicity of $3 + 2\sqrt{2}$ is 1,

Algebraic Multiplicity

Definition

The **(algebraic) multiplicity** of an eigenvalue λ is its multiplicity as a root of the characteristic polynomial.

This is not a very interesting notion *yet*. It will become interesting when we also define *geometric* multiplicity later.

Example

In the previous example, $f(\lambda) = -(\lambda - 2)(\lambda + 1)^2$, so the algebraic multiplicity of the eigenvalue 2 is 1, and the algebraic multiplicity of the eigenvalue -1 is 2.

Example

In the matrix $\begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$, $f(\lambda) = (\lambda - (3 - 2\sqrt{2}))(\lambda - (3 + 2\sqrt{2}))$, so the algebraic multiplicity of $3 + 2\sqrt{2}$ is 1, and the algebraic multiplicity of $3 - 2\sqrt{2}$ is _.

Algebraic Multiplicity

Definition

The **(algebraic) multiplicity** of an eigenvalue λ is its multiplicity as a root of the characteristic polynomial.

This is not a very interesting notion *yet*. It will become interesting when we also define *geometric* multiplicity later.

Example

In the previous example, $f(\lambda) = -(\lambda - 2)(\lambda + 1)^2$, so the algebraic multiplicity of the eigenvalue 2 is 1, and the algebraic multiplicity of the eigenvalue -1 is 2.

Example

In the matrix $\begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$, $f(\lambda) = (\lambda - (3 - 2\sqrt{2}))(\lambda - (3 + 2\sqrt{2}))$, so the algebraic multiplicity of $3 + 2\sqrt{2}$ is 1, and the algebraic multiplicity of $3 - 2\sqrt{2}$ is 1.

The Characteristic Polynomial

Poll

Fact: If A is an $n \times n$ matrix, the characteristic polynomial

$$f(\lambda) = \det(A - \lambda I)$$

turns out to be a polynomial of degree n , and its roots are the eigenvalues of A :

$$f(\lambda) = (-1)^n \lambda^n + a_{n-1} \lambda^{n-1} + a_{n-2} \lambda^{n-2} + \cdots + a_1 \lambda + a_0.$$

The Characteristic Polynomial

Poll

Fact: If A is an $n \times n$ matrix, the characteristic polynomial

$$f(\lambda) = \det(A - \lambda I)$$

turns out to be a polynomial of degree n , and its roots are the eigenvalues of A :

$$f(\lambda) = (-1)^n \lambda^n + a_{n-1} \lambda^{n-1} + a_{n-2} \lambda^{n-2} + \cdots + a_1 \lambda + a_0.$$

Poll

If you count the eigenvalues of A , with their algebraic multiplicities, you will get:

- A. Always n .
- B. Always at most n , but sometimes less.
- C. Always at least n , but sometimes more.
- D. None of the above.

The Characteristic Polynomial

Poll

Fact: If A is an $n \times n$ matrix, the characteristic polynomial

$$f(\lambda) = \det(A - \lambda I)$$

turns out to be a polynomial of degree n , and its roots are the eigenvalues of A :

$$f(\lambda) = (-1)^n \lambda^n + a_{n-1} \lambda^{n-1} + a_{n-2} \lambda^{n-2} + \cdots + a_1 \lambda + a_0.$$

Poll

If you count the eigenvalues of A , with their algebraic multiplicities, you will get:

- A. Always n .
- B. Always at most n , but sometimes less.
- C. Always at least n , but sometimes more.
- D. None of the above.

The answer depends on whether you allow *complex* eigenvalues.

The Characteristic Polynomial

Poll

Fact: If A is an $n \times n$ matrix, the characteristic polynomial

$$f(\lambda) = \det(A - \lambda I)$$

turns out to be a polynomial of degree n , and its roots are the eigenvalues of A :

$$f(\lambda) = (-1)^n \lambda^n + a_{n-1} \lambda^{n-1} + a_{n-2} \lambda^{n-2} + \cdots + a_1 \lambda + a_0.$$

Poll

If you count the eigenvalues of A , with their algebraic multiplicities, you will get:

- A. Always n .
- B. Always at most n , but sometimes less.
- C. Always at least n , but sometimes more.
- D. None of the above.

The answer depends on whether you allow *complex* eigenvalues. If you only allow real eigenvalues, the answer is **B**.

The Characteristic Polynomial

Poll

Fact: If A is an $n \times n$ matrix, the characteristic polynomial

$$f(\lambda) = \det(A - \lambda I)$$

turns out to be a polynomial of degree n , and its roots are the eigenvalues of A :

$$f(\lambda) = (-1)^n \lambda^n + a_{n-1} \lambda^{n-1} + a_{n-2} \lambda^{n-2} + \cdots + a_1 \lambda + a_0.$$

Poll

If you count the eigenvalues of A , with their algebraic multiplicities, you will get:

- A. Always n .
- B. Always at most n , but sometimes less.
- C. Always at least n , but sometimes more.
- D. None of the above.

The answer depends on whether you allow *complex* eigenvalues. If you only allow real eigenvalues, the answer is **B**. Otherwise it is **A**, because any degree- n polynomial has exactly n *complex* roots, counted with multiplicity.

The Characteristic Polynomial

Poll

Fact: If A is an $n \times n$ matrix, the characteristic polynomial

$$f(\lambda) = \det(A - \lambda I)$$

turns out to be a polynomial of degree n , and its roots are the eigenvalues of A :

$$f(\lambda) = (-1)^n \lambda^n + a_{n-1} \lambda^{n-1} + a_{n-2} \lambda^{n-2} + \cdots + a_1 \lambda + a_0.$$

Poll

If you count the eigenvalues of A , with their algebraic multiplicities, you will get:

- A. Always n .
- B. Always at most n , but sometimes less.
- C. Always at least n , but sometimes more.
- D. None of the above.

The answer depends on whether you allow *complex* eigenvalues. If you only allow real eigenvalues, the answer is **B**. Otherwise it is **A**, because any degree- n polynomial has exactly n *complex* roots, counted with multiplicity. Stay tuned.

Section 5.3

Diagonalization

Motivation

Difference equations

Many real-word linear algebra problems have the form:

$$v_1 = Av_0, \quad v_2 = Av_1 = A^2v_0, \quad v_3 = Av_2 = A^3v_0, \quad \dots \quad v_n = Av_{n-1} = A^n v_0.$$

This is called a **difference equation**.

Motivation

Difference equations

Many real-world linear algebra problems have the form:

$$v_1 = Av_0, \quad v_2 = Av_1 = A^2v_0, \quad v_3 = Av_2 = A^3v_0, \quad \dots \quad v_n = Av_{n-1} = A^n v_0.$$

This is called a **difference equation**.

The question is, what happens to v_n as $n \rightarrow \infty$?

Motivation

Difference equations

Many real-word linear algebra problems have the form:

$$v_1 = Av_0, \quad v_2 = Av_1 = A^2v_0, \quad v_3 = Av_2 = A^3v_0, \quad \dots \quad v_n = Av_{n-1} = A^n v_0.$$

This is called a **difference equation**.

The question is, what happens to v_n as $n \rightarrow \infty$?

- ▶ Taking powers of diagonal matrices is easy!

Motivation

Difference equations

Many real-word linear algebra problems have the form:

$$v_1 = Av_0, \quad v_2 = Av_1 = A^2v_0, \quad v_3 = Av_2 = A^3v_0, \quad \dots \quad v_n = Av_{n-1} = A^n v_0.$$

This is called a **difference equation**.

The question is, what happens to v_n as $n \rightarrow \infty$?

- ▶ Taking powers of diagonal matrices is easy!
- ▶ Taking powers of *diagonalizable* matrices is still easy!

Motivation

Difference equations

Many real-word linear algebra problems have the form:

$$v_1 = Av_0, \quad v_2 = Av_1 = A^2v_0, \quad v_3 = Av_2 = A^3v_0, \quad \dots \quad v_n = Av_{n-1} = A^n v_0.$$

This is called a **difference equation**.

The question is, what happens to v_n as $n \rightarrow \infty$?

- ▶ Taking powers of diagonal matrices is easy!
- ▶ Taking powers of *diagonalizable* matrices is still easy!
- ▶ Diagonalizing a matrix is an eigenvalue problem.

Powers of Diagonal Matrices

If D is diagonal, then D^n is also diagonal; its diagonal entries are the n th powers of the diagonal entries of D :

$$D = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}, \quad D^2 = \begin{pmatrix} 4 & 0 \\ 0 & 9 \end{pmatrix}, \quad D^3 = \begin{pmatrix} 8 & 0 \\ 0 & 27 \end{pmatrix}, \quad \dots \quad D^n = \begin{pmatrix} 2^n & 0 \\ 0 & 3^n \end{pmatrix}.$$

$$D = \begin{pmatrix} -1 & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & \frac{1}{3} \end{pmatrix}, \quad D^2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{4} & 0 \\ 0 & 0 & \frac{1}{9} \end{pmatrix}, \quad D^3 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & \frac{1}{8} & 0 \\ 0 & 0 & \frac{1}{27} \end{pmatrix},$$
$$\dots \quad D^n = \begin{pmatrix} (-1)^n & 0 & 0 \\ 0 & \frac{1}{2^n} & 0 \\ 0 & 0 & \frac{1}{3^n} \end{pmatrix}$$

Similarity

Definition

Two $n \times n$ matrices A and B are **similar** if there is an invertible $n \times n$ matrix P such that

$$A = PBP^{-1}.$$

Similarity

Definition

Two $n \times n$ matrices A and B are **similar** if there is an invertible $n \times n$ matrix P such that

$$A = PBP^{-1}.$$

Important Facts:

Similarity

Definition

Two $n \times n$ matrices A and B are **similar** if there is an invertible $n \times n$ matrix P such that

$$A = PBP^{-1}.$$

Important Facts:

1. Similar matrices have the same characteristic polynomial.

Similarity

Definition

Two $n \times n$ matrices A and B are **similar** if there is an invertible $n \times n$ matrix P such that

$$A = PBP^{-1}.$$

Important Facts:

1. Similar matrices have the same characteristic polynomial.
2. It follows that similar matrices have the same eigenvalues.

Similarity

Definition

Two $n \times n$ matrices A and B are **similar** if there is an invertible $n \times n$ matrix P such that

$$A = PBP^{-1}.$$

Important Facts:

1. Similar matrices have the same characteristic polynomial.
2. It follows that similar matrices have the same eigenvalues.
3. If A is similar to B and B is similar to C , then A is similar to C .
4. Matrices with the same characteristic polynomial need not be similar.

Similarity

Definition

Two $n \times n$ matrices A and B are **similar** if there is an invertible $n \times n$ matrix P such that

$$A = PBP^{-1}.$$

Important Facts:

1. Similar matrices have the same characteristic polynomial.
2. It follows that similar matrices have the same eigenvalues.
3. If A is similar to B and B is similar to C , then A is similar to C .
4. Matrices with the same characteristic polynomial need not be similar.
5. Similarity has nothing to do with row equivalence.

Similarity

Definition

Two $n \times n$ matrices A and B are **similar** if there is an invertible $n \times n$ matrix P such that

$$A = PBP^{-1}.$$

Important Facts:

1. Similar matrices have the same characteristic polynomial.
2. It follows that similar matrices have the same eigenvalues.
3. If A is similar to B and B is similar to C , then A is similar to C .
4. Matrices with the same characteristic polynomial need not be similar.
5. Similarity has nothing to do with row equivalence.
6. Similar matrices usually do not have the same eigenvectors.

Powers of Matrices that are Similar to Diagonal Ones

What if A is not diagonal?

Powers of Matrices that are Similar to Diagonal Ones

What if A is not diagonal?

Example

Let $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$. Compute A^n .

Powers of Matrices that are Similar to Diagonal Ones

What if A is not diagonal?

Example

Let $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$. Compute A^n .

$$A = PDP^{-1} \quad \text{where} \quad P = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \quad \text{and} \quad D = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}.$$

Powers of Matrices that are Similar to Diagonal Ones

What if A is not diagonal?

Example

Let $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$. Compute A^n .

$$A = PDP^{-1} \quad \text{where} \quad P = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \quad \text{and} \quad D = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}.$$

Then

$$A^2 = (PDP^{-1})(PDP^{-1}) = PD(P^{-1}P)DP^{-1} = PDIDP^{-1} = PD^2P^{-1}$$

$$A^3 = (PDP^{-1})(PD^2P^{-1}) = PD(P^{-1}P)D^2P^{-1} = PDID^2P^{-1} = PD^3P^{-1}$$

$$\vdots$$

$$A^n = PD^nP^{-1}$$

Powers of Matrices that are Similar to Diagonal Ones

What if A is not diagonal?

Example

Let $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$. Compute A^n .

$$A = PDP^{-1} \quad \text{where} \quad P = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \quad \text{and} \quad D = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}.$$

Then

$$A^2 = (PDP^{-1})(PDP^{-1}) = PD(P^{-1}P)DP^{-1} = PDIDP^{-1} = PD^2P^{-1}$$

$$A^3 = (PDP^{-1})(PD^2P^{-1}) = PD(P^{-1}P)D^2P^{-1} = PDID^2P^{-1} = PD^3P^{-1}$$

$$\vdots$$

$$A^n = PD^nP^{-1}$$

Closed formula in terms of n :
easy to compute



Therefore

$$A^n = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2^n & 0 \\ 0 & 3^n \end{pmatrix} \begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix} = \begin{pmatrix} 2^{n+1} - 3^n & -2^{n+1} + 2 \cdot 3^n \\ 2^n - 3^n & -2^n + 2 \cdot 3^n \end{pmatrix}.$$

Diagonalizable Matrices

Definition

An $n \times n$ matrix A is **diagonalizable** if it is similar to a diagonal matrix:

$$A = PDP^{-1} \quad \text{for } D \text{ diagonal.}$$

Diagonalizable Matrices

Definition

An $n \times n$ matrix A is **diagonalizable** if it is similar to a diagonal matrix:

$$A = PDP^{-1} \quad \text{for } D \text{ diagonal.}$$

Important

If $A = PDP^{-1}$ for $D = \begin{pmatrix} d_{11} & 0 & \cdots & 0 \\ 0 & d_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_{nn} \end{pmatrix}$ then

$$A^k = PD^k P^{-1} = P \begin{pmatrix} d_{11}^k & 0 & \cdots & 0 \\ 0 & d_{22}^k & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_{nn}^k \end{pmatrix} P^{-1}.$$

Diagonalizable Matrices

Definition

An $n \times n$ matrix A is **diagonalizable** if it is similar to a diagonal matrix:

$$A = PDP^{-1} \quad \text{for } D \text{ diagonal.}$$

Important

If $A = PDP^{-1}$ for $D = \begin{pmatrix} d_{11} & 0 & \cdots & 0 \\ 0 & d_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_{nn} \end{pmatrix}$ then

$$A^k = PD^k P^{-1} = P \begin{pmatrix} d_{11}^k & 0 & \cdots & 0 \\ 0 & d_{22}^k & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_{nn}^k \end{pmatrix} P^{-1}.$$

So diagonalizable matrices are easy to raise to any power.

Diagonalization

The Diagonalization Theorem

An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors.

Diagonalization

The Diagonalization Theorem

An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors.

In this case, $A = PDP^{-1}$ for

$$P = \left(\begin{array}{c|c|c|c} | & | & \cdots & | \\ v_1 & v_2 & & v_n \\ | & | & & | \end{array} \right) \quad D = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix},$$

where v_1, v_2, \dots, v_n are linearly independent eigenvectors, and $\lambda_1, \lambda_2, \dots, \lambda_n$ are the corresponding eigenvalues (in the same order).

Diagonalization

The Diagonalization Theorem

An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors.

In this case, $A = PDP^{-1}$ for

$$P = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad D = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix},$$

where v_1, v_2, \dots, v_n are linearly independent eigenvectors, and $\lambda_1, \lambda_2, \dots, \lambda_n$ are the corresponding eigenvalues (in the same order).

Diagonalization

The Diagonalization Theorem

An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors.

In this case, $A = PDP^{-1}$ for

$$P = \left(\begin{array}{c|c|c|c} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{array} \right) \quad D = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix},$$

where v_1, v_2, \dots, v_n are linearly independent eigenvectors, and $\lambda_1, \lambda_2, \dots, \lambda_n$ are the corresponding eigenvalues (in the same order).

Corollary  a theorem that follows easily from another theorem

An $n \times n$ matrix with n distinct eigenvalues is diagonalizable.

Diagonalization

The Diagonalization Theorem

An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors.

In this case, $A = PDP^{-1}$ for

$$P = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad D = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix},$$

where v_1, v_2, \dots, v_n are linearly independent eigenvectors, and $\lambda_1, \lambda_2, \dots, \lambda_n$ are the corresponding eigenvalues (in the same order).

Corollary  a theorem that follows easily from another theorem

An $n \times n$ matrix with n distinct eigenvalues is diagonalizable.

The Corollary is true because eigenvectors with distinct eigenvalues are always linearly independent.

Diagonalization

The Diagonalization Theorem

An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors.

In this case, $A = PDP^{-1}$ for

$$P = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad D = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix},$$

where v_1, v_2, \dots, v_n are linearly independent eigenvectors, and $\lambda_1, \lambda_2, \dots, \lambda_n$ are the corresponding eigenvalues (in the same order).

Corollary  a theorem that follows easily from another theorem

An $n \times n$ matrix with n distinct eigenvalues is diagonalizable.

The Corollary is true because eigenvectors with distinct eigenvalues are always linearly independent. We will see later that a diagonalizable matrix need not have n distinct eigenvalues though.

Diagonalization

Example

Problem: Diagonalize $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$.

The characteristic polynomial is

$$f(\lambda) = \lambda^2 - \text{Tr}(A)\lambda + \det(A) = \lambda^2 - 5\lambda + 6 = (\lambda - 2)(\lambda - 3).$$

Therefore the eigenvalues are 2 and 3. Let's compute some eigenvectors:

$$(A - 2I)x = 0 \iff \begin{pmatrix} -1 & 2 \\ -1 & 2 \end{pmatrix} x = 0 \xrightarrow{\text{rref}} \begin{pmatrix} 1 & -2 \\ 0 & 0 \end{pmatrix} x = 0$$

The parametric form is $x = 2y$, so $v_1 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ is an eigenvector with eigenvalue 2.

$$(A - 3I)x = 0 \iff \begin{pmatrix} -2 & 2 \\ -1 & 1 \end{pmatrix} x = 0 \xrightarrow{\text{rref}} \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix} x = 0$$

The parametric form is $x = y$, so $v_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ is an eigenvector with eigenvalue 3.

The eigenvectors v_1, v_2 are linearly independent, so the Diagonalization Theorem says

$$A = PDP^{-1} \quad \text{for} \quad P = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \quad D = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}.$$

Diagonalization

Another example

Problem: Diagonalize $A = \begin{pmatrix} 4 & -3 & 0 \\ 2 & -1 & 0 \\ 1 & -1 & 1 \end{pmatrix}$.

The characteristic polynomial is

$$f(\lambda) = \det(A - \lambda I) = -\lambda^3 + 4\lambda^2 - 5\lambda + 2 = -(\lambda - 1)^2(\lambda - 2).$$

Therefore the eigenvalues are 1 and 2, with respective multiplicities 2 and 1.

Let's compute the 1-eigenspace:

$$(A - I)x = 0 \iff \begin{pmatrix} 3 & -3 & 0 \\ 2 & -2 & 0 \\ 1 & -1 & 0 \end{pmatrix} x = 0 \xrightarrow{\text{rref}} \begin{pmatrix} 1 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} x = 0$$

The parametric vector form is

$$\begin{array}{rcl} x & = & y \\ y & = & y \\ z & = & z \end{array} \implies \begin{pmatrix} x \\ y \\ z \end{pmatrix} = y \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + z \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

Hence a basis for the 1-eigenspace is

$$\mathcal{B}_1 = \{v_1, v_2\} \quad \text{where} \quad v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad v_2 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}.$$

Diagonalization

Another example, continued

Problem: Diagonalize $A = \begin{pmatrix} 4 & -3 & 0 \\ 2 & -1 & 0 \\ 1 & -1 & 1 \end{pmatrix}$.

Now let's compute the 2-eigenspace:

$$(A - 2I)x = 0 \iff \begin{pmatrix} 2 & -3 & 0 \\ 2 & -3 & 0 \\ 1 & -1 & -1 \end{pmatrix} x = 0 \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & -3 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{pmatrix} x = 0$$

The parametric form is $x = 3z, y = 2z$, so an eigenvector with eigenvalue 2 is

$$v_3 = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}.$$

The eigenvectors v_1, v_2, v_3 are linearly independent: v_1, v_2 form a basis for the 1-eigenspace, and v_3 is not contained in the 1-eigenspace. Therefore the Diagonalization Theorem says

$$A = PDP^{-1} \quad \text{for} \quad P = \begin{pmatrix} 1 & 0 & 3 \\ 1 & 0 & 2 \\ 0 & 1 & 1 \end{pmatrix} \quad D = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}.$$

Diagonalization

Another example, continued

Problem: Diagonalize $A = \begin{pmatrix} 4 & -3 & 0 \\ 2 & -1 & 0 \\ 1 & -1 & 1 \end{pmatrix}$.

Now let's compute the 2-eigenspace:

$$(A - 2I)x = 0 \iff \begin{pmatrix} 2 & -3 & 0 \\ 2 & -3 & 0 \\ 1 & -1 & -1 \end{pmatrix} x = 0 \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & -3 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{pmatrix} x = 0$$

The parametric form is $x = 3z, y = 2z$, so an eigenvector with eigenvalue 2 is

$$v_3 = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}.$$

The eigenvectors v_1, v_2, v_3 are linearly independent: v_1, v_2 form a basis for the 1-eigenspace, and v_3 is not contained in the 1-eigenspace. Therefore the Diagonalization Theorem says

$$A = PDP^{-1} \quad \text{for} \quad P = \begin{pmatrix} 1 & 0 & 3 \\ 1 & 0 & 2 \\ 0 & 1 & 1 \end{pmatrix} \quad D = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}.$$

Note: In this case, there are three linearly independent eigenvectors, but only two distinct eigenvalues.

Diagonalization

A non-diagonalizable matrix

Problem: Show that $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ is not diagonalizable.

This is an upper-triangular matrix, so the only eigenvalue is 1. Let's compute the 1-eigenspace:

$$(A - I)x = 0 \iff \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} x = 0.$$

This is row reduced, but has only one free variable x ; a basis for the 1-eigenspace is $\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\}$. So *all eigenvectors* of A are multiples of $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

Diagonalization

A non-diagonalizable matrix

Problem: Show that $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ is not diagonalizable.

This is an upper-triangular matrix, so the only eigenvalue is 1. Let's compute the 1-eigenspace:

$$(A - I)x = 0 \iff \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} x = 0.$$

This is row reduced, but has only one free variable x ; a basis for the 1-eigenspace is $\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\}$. So *all eigenvectors* of A are multiples of $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

Conclusion: A has only one linearly independent eigenvector, so by the “only if” part of the diagonalization theorem, A is not diagonalizable.

Poll

Which of the following matrices are diagonalizable, and why?

A. $\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ B. $\begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix}$ C. $\begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}$ D. $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$

Poll

Which of the following matrices are diagonalizable, and why?

A. $\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ B. $\begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix}$ C. $\begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}$ D. $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$

Matrix **A** is not diagonalizable: its only eigenvalue is 1, and its 1-eigenspace is spanned by $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

Poll

Which of the following matrices are diagonalizable, and why?

A. $\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ B. $\begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix}$ C. $\begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}$ D. $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$

Matrix **A** is not diagonalizable: its only eigenvalue is 1, and its 1-eigenspace is spanned by $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

Similarly, matrix **C** is not diagonalizable.

Poll

Which of the following matrices are diagonalizable, and why?

A. $\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ B. $\begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix}$ C. $\begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}$ D. $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$

Matrix **A** is not diagonalizable: its only eigenvalue is 1, and its 1-eigenspace is spanned by $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

Similarly, matrix **C** is not diagonalizable.

Matrix **B** is diagonalizable because it is a 2×2 matrix with distinct eigenvalues.

Poll

Which of the following matrices are diagonalizable, and why?

A. $\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ B. $\begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix}$ C. $\begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}$ D. $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$

Matrix **A** is not diagonalizable: its only eigenvalue is 1, and its 1-eigenspace is spanned by $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

Similarly, matrix **C** is not diagonalizable.

Matrix **B** is diagonalizable because it is a 2×2 matrix with distinct eigenvalues.

Matrix **D** is already diagonal!

Diagonalization

Procedure

How to diagonalize a matrix A :

Diagonalization

Procedure

How to diagonalize a matrix A :

1. Find the eigenvalues of A using the characteristic polynomial.

Diagonalization

Procedure

How to diagonalize a matrix A :

1. Find the eigenvalues of A using the characteristic polynomial.
2. For each eigenvalue λ of A , compute a basis \mathcal{B}_λ for the λ -eigenspace.

Diagonalization

Procedure

How to diagonalize a matrix A :

1. Find the eigenvalues of A using the characteristic polynomial.
2. For each eigenvalue λ of A , compute a basis \mathcal{B}_λ for the λ -eigenspace.
3. If there are fewer than n total vectors in the union of all of the eigenspace bases \mathcal{B}_λ , then the matrix is not diagonalizable.

Diagonalization

Procedure

How to diagonalize a matrix A :

1. Find the eigenvalues of A using the characteristic polynomial.
2. For each eigenvalue λ of A , compute a basis \mathcal{B}_λ for the λ -eigenspace.
3. If there are fewer than n total vectors in the union of all of the eigenspace bases \mathcal{B}_λ , then the matrix is not diagonalizable.
4. Otherwise, the n vectors v_1, v_2, \dots, v_n in your eigenspace bases are linearly independent, and $A = PDP^{-1}$ for

$$P = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad \text{and} \quad D = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix},$$

where λ_i is the eigenvalue for v_i .

Diagonalization

Proof

Why is the Diagonalization Theorem true?

A diagonalizable implies A has n linearly independent eigenvectors: Suppose $A = PDP^{-1}$, where D is diagonal with diagonal entries $\lambda_1, \lambda_2, \dots, \lambda_n$. Let v_1, v_2, \dots, v_n be the columns of P . They are linearly independent because P is invertible.

$$\begin{aligned} AP &= A \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} = \begin{pmatrix} | & | & \cdots & | \\ Av_1 & Av_2 & \cdots & Av_n \\ | & | & \cdots & | \end{pmatrix} \\ &= \begin{pmatrix} | & | & \cdots & | \\ \lambda_1 v_1 & \lambda_1 v_2 & \cdots & \lambda_1 v_n \\ | & | & \cdots & | \end{pmatrix} = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} \end{aligned}$$

Hence $AP = PD$.

Diagonalization

Proof

Why is the Diagonalization Theorem true?

A diagonalizable implies A has n linearly independent eigenvectors: Suppose $A = PDP^{-1}$, where D is diagonal with diagonal entries $\lambda_1, \lambda_2, \dots, \lambda_n$. Let v_1, v_2, \dots, v_n be the columns of P . They are linearly independent because P is invertible. So $Pe_i = v_i$, hence $P^{-1}v_i = e_i$.

$$Av_i = PDP^{-1}v_i = PDe_i = P(\lambda_i e_i) = \lambda_i Pe_i = \lambda_i v_i.$$

Hence v_i is an eigenvector of A with eigenvalue λ_i . So the columns of P form n linearly independent eigenvectors of A , and the diagonal entries of D are the eigenvalues.

A has n linearly independent eigenvectors implies A is diagonalizable: Suppose A has n linearly independent eigenvectors v_1, v_2, \dots, v_n , with eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$. Let P be the invertible matrix with columns v_1, v_2, \dots, v_n . Let $D = P^{-1}AP$.

$$De_i = P^{-1}APe_i = P^{-1}Av_i = P^{-1}(\lambda_i v_i) = \lambda_i P^{-1}v_i = \lambda_i e_i.$$

Hence D is diagonal, with diagonal entries $\lambda_1, \lambda_2, \dots, \lambda_n$. Solving $D = P^{-1}AP$ for A gives $A = PDP^{-1}$.

Non-Distinct Eigenvalues

Definition

Let λ be an eigenvalue of a square matrix A . The **geometric multiplicity** of λ is the dimension of the λ -eigenspace.

Non-Distinct Eigenvalues

Definition

Let λ be an eigenvalue of a square matrix A . The **geometric multiplicity** of λ is the dimension of the λ -eigenspace.

Theorem

Let λ be an eigenvalue of a square matrix A . Then

$$1 \leq (\text{the geometric multiplicity of } \lambda) \leq (\text{the algebraic multiplicity of } \lambda).$$

Non-Distinct Eigenvalues

Definition

Let λ be an eigenvalue of a square matrix A . The **geometric multiplicity** of λ is the dimension of the λ -eigenspace.

Theorem

Let λ be an eigenvalue of a square matrix A . Then

$$1 \leq (\text{the geometric multiplicity of } \lambda) \leq (\text{the algebraic multiplicity of } \lambda).$$

The proof is beyond the scope of this course.

Non-Distinct Eigenvalues

Definition

Let λ be an eigenvalue of a square matrix A . The **geometric multiplicity** of λ is the dimension of the λ -eigenspace.

Theorem

Let λ be an eigenvalue of a square matrix A . Then

$$1 \leq (\text{the geometric multiplicity of } \lambda) \leq (\text{the algebraic multiplicity of } \lambda).$$

The proof is beyond the scope of this course.

Corollary

Let λ be an eigenvalue of a square matrix A . If the algebraic multiplicity of λ is 1, then the geometric multiplicity is also 1.

Non-Distinct Eigenvalues

Definition

Let λ be an eigenvalue of a square matrix A . The **geometric multiplicity** of λ is the dimension of the λ -eigenspace.

Theorem

Let λ be an eigenvalue of a square matrix A . Then

$$1 \leq (\text{the geometric multiplicity of } \lambda) \leq (\text{the algebraic multiplicity of } \lambda).$$

The proof is beyond the scope of this course.

Corollary

Let λ be an eigenvalue of a square matrix A . If the algebraic multiplicity of λ is 1, then the geometric multiplicity is also 1.

The Diagonalization Theorem (Alternate Form)

Let A be an $n \times n$ matrix. The following are equivalent:

Non-Distinct Eigenvalues

Definition

Let λ be an eigenvalue of a square matrix A . The **geometric multiplicity** of λ is the dimension of the λ -eigenspace.

Theorem

Let λ be an eigenvalue of a square matrix A . Then

$$1 \leq (\text{the geometric multiplicity of } \lambda) \leq (\text{the algebraic multiplicity of } \lambda).$$

The proof is beyond the scope of this course.

Corollary

Let λ be an eigenvalue of a square matrix A . If the algebraic multiplicity of λ is 1, then the geometric multiplicity is also 1.

The Diagonalization Theorem (Alternate Form)

Let A be an $n \times n$ matrix. The following are equivalent:

1. A is diagonalizable.

Non-Distinct Eigenvalues

Definition

Let λ be an eigenvalue of a square matrix A . The **geometric multiplicity** of λ is the dimension of the λ -eigenspace.

Theorem

Let λ be an eigenvalue of a square matrix A . Then

$$1 \leq (\text{the geometric multiplicity of } \lambda) \leq (\text{the algebraic multiplicity of } \lambda).$$

The proof is beyond the scope of this course.

Corollary

Let λ be an eigenvalue of a square matrix A . If the algebraic multiplicity of λ is 1, then the geometric multiplicity is also 1.

The Diagonalization Theorem (Alternate Form)

Let A be an $n \times n$ matrix. The following are equivalent:

1. A is diagonalizable.
2. The sum of the geometric multiplicities of the eigenvalues of A equals n .

Non-Distinct Eigenvalues

Definition

Let λ be an eigenvalue of a square matrix A . The **geometric multiplicity** of λ is the dimension of the λ -eigenspace.

Theorem

Let λ be an eigenvalue of a square matrix A . Then

$$1 \leq (\text{the geometric multiplicity of } \lambda) \leq (\text{the algebraic multiplicity of } \lambda).$$

The proof is beyond the scope of this course.

Corollary

Let λ be an eigenvalue of a square matrix A . If the algebraic multiplicity of λ is 1, then the geometric multiplicity is also 1.

The Diagonalization Theorem (Alternate Form)

Let A be an $n \times n$ matrix. The following are equivalent:

1. A is diagonalizable.
2. The sum of the geometric multiplicities of the eigenvalues of A equals n .
3. The sum of the algebraic multiplicities of the eigenvalues of A equals n , and *the geometric multiplicity equals the algebraic multiplicity* of each eigenvalue.

Non-Distinct Eigenvalues

Examples

Example

If A has n distinct eigenvalues, then the algebraic multiplicity of each equals 1, hence so does the geometric multiplicity, and therefore A is diagonalizable.

Non-Distinct Eigenvalues

Examples

Example

If A has n distinct eigenvalues, then the algebraic multiplicity of each equals 1, hence so does the geometric multiplicity, and therefore A is diagonalizable.

For example, $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$ has eigenvalues 2 and 3, so it is diagonalizable.

Non-Distinct Eigenvalues

Examples

Example

If A has n distinct eigenvalues, then the algebraic multiplicity of each equals 1, hence so does the geometric multiplicity, and therefore A is diagonalizable.

For example, $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$ has eigenvalues 2 and 3, so it is diagonalizable.

Example

The matrix $A = \begin{pmatrix} 4 & -3 & 0 \\ 2 & -1 & 0 \\ 1 & -1 & 1 \end{pmatrix}$ has characteristic polynomial

$$f(\lambda) = -(\lambda - 1)^2(\lambda - 2).$$

Non-Distinct Eigenvalues

Examples

Example

If A has n distinct eigenvalues, then the algebraic multiplicity of each equals 1, hence so does the geometric multiplicity, and therefore A is diagonalizable.

For example, $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$ has eigenvalues 2 and 3, so it is diagonalizable.

Example

The matrix $A = \begin{pmatrix} 4 & -3 & 0 \\ 2 & -1 & 0 \\ 1 & -1 & 1 \end{pmatrix}$ has characteristic polynomial

$$f(\lambda) = -(\lambda - 1)^2(\lambda - 2).$$

The algebraic multiplicities of 1 and 2 are 2 and 1, respectively.

Non-Distinct Eigenvalues

Examples

Example

If A has n distinct eigenvalues, then the algebraic multiplicity of each equals 1, hence so does the geometric multiplicity, and therefore A is diagonalizable.

For example, $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$ has eigenvalues 2 and 3, so it is diagonalizable.

Example

The matrix $A = \begin{pmatrix} 4 & -3 & 0 \\ 2 & -1 & 0 \\ 1 & -1 & 1 \end{pmatrix}$ has characteristic polynomial

$$f(\lambda) = -(\lambda - 1)^2(\lambda - 2).$$

The algebraic multiplicities of 1 and 2 are 2 and 1, respectively. They sum to 3.

Non-Distinct Eigenvalues

Examples

Example

If A has n distinct eigenvalues, then the algebraic multiplicity of each equals 1, hence so does the geometric multiplicity, and therefore A is diagonalizable.

For example, $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$ has eigenvalues 2 and 3, so it is diagonalizable.

Example

The matrix $A = \begin{pmatrix} 4 & -3 & 0 \\ 2 & -1 & 0 \\ 1 & -1 & 1 \end{pmatrix}$ has characteristic polynomial

$$f(\lambda) = -(\lambda - 1)^2(\lambda - 2).$$

The algebraic multiplicities of 1 and 2 are 2 and 1, respectively. They sum to 3. We showed before that the geometric multiplicity of 1 is 2 (the 1-eigenspace has dimension 2).

Non-Distinct Eigenvalues

Examples

Example

If A has n distinct eigenvalues, then the algebraic multiplicity of each equals 1, hence so does the geometric multiplicity, and therefore A is diagonalizable.

For example, $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$ has eigenvalues 2 and 3, so it is diagonalizable.

Example

The matrix $A = \begin{pmatrix} 4 & -3 & 0 \\ 2 & -1 & 0 \\ 1 & -1 & 1 \end{pmatrix}$ has characteristic polynomial

$$f(\lambda) = -(\lambda - 1)^2(\lambda - 2).$$

The algebraic multiplicities of 1 and 2 are 2 and 1, respectively. They sum to 3. We showed before that the geometric multiplicity of 1 is 2 (the 1-eigenspace has dimension 2). The eigenvalue 2 automatically has geometric multiplicity 1.

Non-Distinct Eigenvalues

Examples

Example

If A has n distinct eigenvalues, then the algebraic multiplicity of each equals 1, hence so does the geometric multiplicity, and therefore A is diagonalizable.

For example, $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$ has eigenvalues 2 and 3, so it is diagonalizable.

Example

The matrix $A = \begin{pmatrix} 4 & -3 & 0 \\ 2 & -1 & 0 \\ 1 & -1 & 1 \end{pmatrix}$ has characteristic polynomial

$$f(\lambda) = -(\lambda - 1)^2(\lambda - 2).$$

The algebraic multiplicities of 1 and 2 are 2 and 1, respectively. They sum to 3. We showed before that the geometric multiplicity of 1 is 2 (the 1-eigenspace has dimension 2). The eigenvalue 2 automatically has geometric multiplicity 1. Hence the geometric multiplicities add up to 3, so A is diagonalizable.

Non-Distinct Eigenvalues

Another example

Example

The matrix $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ has characteristic polynomial $f(\lambda) = (\lambda - 1)^2$.

Non-Distinct Eigenvalues

Another example

Example

The matrix $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ has characteristic polynomial $f(\lambda) = (\lambda - 1)^2$.

It has one eigenvalue 1 of algebraic multiplicity 2.

Non-Distinct Eigenvalues

Another example

Example

The matrix $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ has characteristic polynomial $f(\lambda) = (\lambda - 1)^2$.

It has one eigenvalue 1 of algebraic multiplicity 2.

We showed before that the geometric multiplicity of 1 is 1 (the 1-eigenspace has dimension 1).

Non-Distinct Eigenvalues

Another example

Example

The matrix $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ has characteristic polynomial $f(\lambda) = (\lambda - 1)^2$.

It has one eigenvalue 1 of algebraic multiplicity 2.

We showed before that the geometric multiplicity of 1 is 1 (the 1-eigenspace has dimension 1).

Since the geometric multiplicity is smaller than the algebraic multiplicity, the matrix is *not* diagonalizable.

Linear Recurrence Relation

Linear Recurrence Relation

Let $(x_n) = (x_0, x_1, x_2, \dots)$ be a sequence of numbers that is defined as follows:

1. $x_0 = a_0, x_1 = a_1, \dots, x_{k-1} = a_{k-1}$, where a_0, a_1, \dots, a_{k-1} are scalars.
2. For all $n \geq k$, $x_n = c_1 x_{n-1} + c_2 x_{n-2} + \dots + c_k x_{n-k}$ where c_1, c_2, \dots, c_k are scalars.

If $c_k \neq 0$, the equation in (2) is called a linear recurrence relation of order k .
The equations in (1) are referred to as the initial conditions of the recurrence.

Examples

- ▶ $x_{n+2} = x_{n+1} + x_n, \quad x_0 = 1, x_1 = 1.$
- ▶ $x_{n+1} = 2x_n, \quad x_0 = 3.$

Linear Recurrence in Matrix Form

I am going to explain it using an example of second order linear recurrence relation Consider the following linear recurrence relation

$$x_{n+2} = ax_{n+1} + bx_n, \quad x_1 = c_1, x_0 = c_0,$$

where c_0 and c_1 are known constants. We can write it as

$$\begin{aligned}x_{n+2} &= ax_{n+1} + bx_n \\x_{n+1} &= x_{n+1}.\end{aligned}$$

In Matrix form, we can write

$$\begin{pmatrix} x_{n+2} \\ x_{n+1} \end{pmatrix} = \begin{pmatrix} a & b \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x_{n+1} \\ x_n \end{pmatrix}$$

$$\boxed{X_{n+1} = AX_n, \forall n \geq 0.}$$

where $X_{n+1} = \begin{pmatrix} x_{n+2} \\ x_{n+1} \end{pmatrix}$, and $A = \begin{pmatrix} a & b \\ 1 & 0 \end{pmatrix}$

Linear Recurrence in Matrix Form; contr'n'd...

For $n = 0$, we have

$$X_1 = AX_0,$$

where $X_0 = \begin{pmatrix} c_1 \\ c_0 \end{pmatrix}$

For $n = 1$, we can write

$$X_2 = AX_1 = A(AX_0) = A^2X_0.$$

$n = 2$, gives us

$$X_3 = AX_2 = A(A^2X_0) = A^3X_0.$$

Continuing in the same manner, we have

$$X_{n+1} = A^{n+1}X_0.$$

Examples

Suppose each "Gibonacci" number G_{k+2} is the average of the two previous numbers G_{k+1} and G_k . If $G_0 = 0$ and $G_1 = 1$. Find the k th term of the sequence only depending upon k . *Aim :* We want to find the general term of the sequence. *Steps :*

- ▶ Matrix Form
- ▶ Eigenvalues and Eigenvectors
- ▶ Diagonalize

Examples; Contn'd...

$$G_{k+2} = \frac{1}{2}G_{k+1} + \frac{1}{2}G_k$$
$$G_{k+1} = G_{k+1}.$$

In Matrix Form

$$\mathbf{G}_{k+1} = \mathbf{A}\mathbf{G}_k, \quad \forall k \geq 0.$$

$$\text{where } \mathbf{G}_{k+1} = \begin{pmatrix} G_{k+2} \\ G_{k+1} \end{pmatrix}, \mathbf{A} = \begin{pmatrix} 1/2 & 1/2 \\ 1 & 0 \end{pmatrix}, \mathbf{G}_0 = \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

$$\mathbf{G}_k = \mathbf{A}^k \mathbf{G}_0, \quad \forall k \geq 0.$$

Eigenvalues Characteristic Equation

$$\lambda^2 - \frac{1}{2}\lambda - \frac{1}{2} = 0$$

Eigenvalues are: $1, -\frac{1}{2}$

Examples; Contn'd...

Eigenvectors $\lambda = 1$

$$(A - 1I)X = 0$$

Augmented matrix

$$\begin{pmatrix} -\frac{1}{2} & \frac{1}{2} & 0 \\ 1 & -1 & 0 \end{pmatrix}$$

Eigenvector: All non-zero multiples of $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$

Eigenvectors $\lambda = \frac{-1}{2}$

$$(A + \frac{1}{2}I)X = 0$$

Augmented matrix

$$\begin{pmatrix} 1 & \frac{1}{2} & 0 \\ 1 & \frac{1}{2} & 0 \end{pmatrix}$$

Eigenvector: All non-zero multiples of $\begin{pmatrix} -1 \\ 2 \end{pmatrix}$

Examples; Contn'd...

$$\begin{pmatrix} 1 & -1 \\ 1 & 2 \end{pmatrix}, \quad D = \begin{pmatrix} 1 & 0 \\ 0 & \frac{-1}{2} \end{pmatrix}$$

As $A^k = PD^kP^{-1}$, so we need to calculate P^{-1} .

$$P^{-1} = \frac{\text{adj}P}{\det P} = \frac{1}{3} \begin{pmatrix} 2 & 1 \\ -1 & 1 \end{pmatrix}.$$

So,

$$A^k = \frac{1}{3} \begin{pmatrix} 1 & -1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & \frac{(-1)^k}{(2)^k} \end{pmatrix} \begin{pmatrix} 2 & 1 \\ -1 & 1 \end{pmatrix}$$

Simplification gives us

$$A^k = \frac{1}{3} \begin{pmatrix} \frac{(-1)^k}{(2)^k} + 2 & 1 - \frac{(-1)^k}{(2)^k} \\ 2 - \frac{2(-1)^k}{(2)^k} & \frac{(-1)^k}{(2)^k} + 1 \end{pmatrix}$$

Examples; Contn'd...

$$\mathbf{G}_k = \mathbf{A}^k \mathbf{G}_0$$

$$\begin{pmatrix} G_{k+2} \\ G_{k+1} \end{pmatrix} = \frac{1}{3} \begin{pmatrix} \frac{(-1)^k}{(2)^k} + 2 & 1 - \frac{(-1)^k}{(2)^k} \\ 2 - \frac{2(-1)^k}{(2)^k} & \frac{(-1)^k}{(2)^k} + 1 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

Hence

$$G_k = \frac{2}{3} - \frac{2}{3} \frac{(-1)^k}{(2)^k}$$

Important Result

Theorem

Let $x_n = ax_{n-1} + bx_{n-2}$ be a recurrence relation. Let λ_1 and λ_2 be the eigenvalues of the associated characteristic equation $\lambda^2 - a\lambda - b = 0$.

1. If $\lambda_1 \neq \lambda_2$, then

$$x_n = c_1 \lambda_1^n + c_2 \lambda_2^n$$

for some scalars c_1 and c_2 .

2. If $\lambda_1 = \lambda_2$, then

$$x_n = (c_1 + c_2 n) \lambda^n$$

for some scalars c_1 and c_2 .

Examples

Solve the following recurrence relation with the given initial conditions.

$$y_1 = 1, y_2 = 6, y_k = 4y_{k-1} - 4y_{k-2}, \quad k \geq 3.$$

Characteristics equation $\lambda^2 - 4\lambda + 4 = 0$.

Solution of the quadratic equation is Eigenvalues: $\lambda_1 = 2, \lambda_1 = 2$. So,

$$y_k = c_1(2)^k + c_2 k 2^k.$$

As, $y_1 = 1$, so, $2c_1 + 2c_2 = 1$, $y_2 = 6$, so, $4c_1 + 8c_2 = 6$.

Solution of above system is

$$c_1 = -\frac{1}{2}, \quad c_2 = 1.$$

Hence,

$$y_k = -\frac{1}{2}2^k + k2^k$$

Practice Problems

1. Solve the recurrence relation with the given initial conditions.

▶ $a_0 = 4, a_1 = 1, a_n = a_{n-1} - \frac{a_{n-2}}{4}, \text{ for } n \geq 2.$

▶ $a_n = 2a_{n-1} + a_{n-2} - 2a_{n-3}, \text{ subject to } a_0 = 2, a_1 = 2, a_2 = 4, \text{ for } n \geq 3.$

2. Find the limiting values of y_k and z_k , ($k \rightarrow \infty$) if

$$y_{k+1} = .8y_k + .3z_k, \quad y_0 = 0$$

$$z_{k+1} = .2y_k + .7z_k, \quad z_0 = 5.$$

Systems of Linear Differential Equations

First order linear homogenous differential equation

First order differential Equation

$$x' = kx, \quad k \text{ is a constant.}$$

Solution: $x(t) = x_0 e^{kt}$.

First order differential system

$$x' = 4x$$

$$y' = 9y$$

In Matrix form of above system can be written as

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 4 & 0 \\ 0 & 9 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

- ▶ Matrix is diagonal
- ▶ System is uncoupled

First order linear system of differential equations

General first order linear system

$$x' = ax + by$$

$$y' = cx + dy$$

In Matrix form

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

We can write the above system as

$$X' = AX,$$

where $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and $X = \begin{pmatrix} x \\ y \end{pmatrix}$.

- ▶ A is not a diagonal matrix.
- ▶ Can we diagonalize A ?

Diagonalization

Aim :

To solve the system $X' = AX$.

Challenges

Matrix is not diagonal.

Possible Solution

Transform the matrix into a diagonal matrix i.e., diagonalize it.

HOW?

We want to transform

$$X' = AX \xrightarrow{\text{to}} Y' = DY.$$

Diagonalization

As $PDP^{-1} = A$, so we can write

$$X' = AX = PDP^{-1}X$$

Pre multiplying by P^{-1} we get

$$P^{-1}X' = DP^{-1}X$$

Since, P is a constant matrix, so

$$\left(P^{-1}X\right)' = D\left(P^{-1}X\right).$$

Put $(P^{-1}X) = Y$ to get

$$\boxed{Y' = DY}.$$

Uncoupling system of differential equations

Summary

Coupled system of differential equation

$$X' = AX$$

can be transformed (uncoupled) to

$$Y' = DY$$

by using the transformation

$$X = PY.$$

Example

Find a solution to the system

$$x' = x + 3y$$

$$y' = 2x + 2y$$

subject to initial conditions $x(0) = 0$, $y(0) = 5$.

Solution: In Matrix form, we can write it as

$$X' = AX,$$

where $A = \begin{pmatrix} 1 & 3 \\ 2 & 2 \end{pmatrix}$ and $X = \begin{pmatrix} x \\ y \end{pmatrix}$.

We can uncoupled the system by using the transformation

$$X = PY.$$

Example

Eigenvalues:

Characteristic Equation:

$$\lambda^2 - 3\lambda - 4 = 0.$$

Eigenvalues are: $-1, 4$.

Corresponding eigenvectors are

$$\begin{pmatrix} 3 \\ -2 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

Hence,

$$P = \begin{pmatrix} 3 & 1 \\ -2 & 1 \end{pmatrix}, D = \begin{pmatrix} -1 & 0 \\ 0 & 4 \end{pmatrix}.$$

By using the transformation $X = PY$, we get

$$Y' = DY,$$

where $Y = \begin{pmatrix} u \\ v \end{pmatrix}$.

Example

$$\begin{pmatrix} u' \\ v' \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 4 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}.$$

Above equations can be written as

$$u' = -u$$

$$v' = 4v.$$

Solving, we get

$$u = c_1 e^{-t}, \quad v = c_2 e^{4t}.$$

As $X = PY$, so, we can write

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 3 & 1 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}.$$

$$x = 3u + v$$

$$y = -2u + v.$$

Substituting values of u and v , we get

$$x = 3c_1 e^{-t} + c_2 e^{4t}$$

$$y = -2c_1 e^{-t} + c_2 e^{4t}.$$

Example

Since, $x(0) = 0$ and $y(0) = 5$, we get

$$0 = 3c_1 + c_2$$

$$5 = -2c_1 + c_2.$$

Solving, above system we get

$$c_1 = -1, c_2 = 3.$$

In matrix form we can the solution as

$$X = -x_1 e^{-t} + 3x_2 e^{4t}$$

where $x_1 = \begin{pmatrix} 3 \\ -2 \end{pmatrix}$ and $x_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ are the eigenvectors corresponding to eigenvalues -1 and 4 respectively.

Example

Find a solution to the system

$$r'(t) = w(t) - 12$$

$$w'(t) = -r(t) + 10$$

Solution:

Issue :

Presence of -12 and 10 .

How to resolve it :

Put $w(t) - 12 = y(t)$ and $-r(t) + 10 = x(t)$, we get

$$-x'(t) = y(t)$$

$$y'(t) = x(t).$$

In Matrix form, we can write it as

$$X' = AX,$$

where $A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $X = \begin{pmatrix} x \\ y \end{pmatrix}$.

By using the substitution $X = PY$ we get

$$Y' = DY$$

$$\text{where } P = \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix}, D = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}.$$

So, solution is

$$u = c_1 e^{-it}$$

$$v = c_2 e^{it}$$

By using the relation $X = PY$, we get

$$x = c_1 e^{-it} + c_2 e^{it}$$

$$y = c_1 i e^{-it} - i c_2 e^{it}.$$

Solution of the system is

$$r(t) = 10 - c_1 e^{-it} - c_2 e^{it}$$

$$w(t) = 12 + c_1 i e^{-it} - i c_2 e^{it}$$

- In case of single linear differential equation, we have

$$x' = kx, \text{ } k \text{ is a constant.}$$

Solution of the differential equation is

$$x = ce^{kt}.$$

- In case of system of coupled differential equations, we have

$$X' = AX, \text{ } A \text{ is a constant matrix.}$$

Solution of the linear differential system should be

$$X = c e^{At}.$$

Exponential of a Matrix

Compute e^{Dt} where $D = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}$.

Since, $e^x = 1 + x + \frac{x^2}{2!} + \dots = \sum_{n=0}^{\infty} \frac{x^n}{n!}$. so,

$$e^{Dt} = I + Dt + \dots = \sum_{n=0}^{\infty} \frac{D^n t^n}{n!}$$

$$e^{Dt} = \sum_{n=0}^{\infty} \frac{1}{n!} \begin{pmatrix} 4^n t^n & 0 \\ 0 & t^n \end{pmatrix}$$

$$e^{Dt} = \begin{pmatrix} \sum_{n=0}^{\infty} \frac{4^n}{n!} & 0 \\ 0 & \sum_{n=0}^{\infty} \frac{1^n}{n!} \end{pmatrix}$$

$$e^{Dt} = \begin{pmatrix} e^{4t} & 0 \\ 0 & e^t \end{pmatrix}$$

Example

Compute e^{At} where $A = \begin{pmatrix} 1 & 3 \\ 2 & 2 \end{pmatrix}$

For given matrix, we have

$$P = \begin{pmatrix} 3 & 1 \\ -1 & 1 \end{pmatrix}, \quad D = \begin{pmatrix} -1 & 0 \\ 0 & 4 \end{pmatrix}.$$

$$e^{At} = \sum_{n=0}^{\infty} \frac{A^n t^n}{n!}$$

$$e^{At} = \sum_{n=0}^{\infty} \frac{PD^n P^{-1}}{n!}$$

$$e^{At} = P \sum_{n=0}^{\infty} \frac{D^n t^n}{n!} P^{-1}$$

$$e^{At} = P e^{Dt} P^{-1}$$

$$e^{At} = \begin{pmatrix} 3 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} e^{-t} & 0 \\ 0 & e^{4t} \end{pmatrix} \left(\begin{pmatrix} 3 & 1 \\ -1 & 1 \end{pmatrix} \right)^{-1}.$$