

# MT-1004

## Linear Algebra

Fall 2023

**Week # 3**

National University of Computer and Emerging Sciences


September 6, 2023

# Section 1.3

## Vector Equations


## Plan For Today

Today we will learn to describe and draw the solution set of an arbitrary system of linear equations  $Ax = b$ , using spans.



## Plan For Today


Today we will learn to describe and draw the solution set of an arbitrary system of linear equations  $Ax = b$ , using spans.



**Recall:** the **solution set** is the collection of all vectors  $x$  such that  $Ax = b$  is true.

## Plan For Today

Today we will learn to describe and draw the solution set of an arbitrary system of linear equations  $Ax = b$ , using spans.




**Recall:** the **solution set** is the collection of all vectors  $x$  such that  $Ax = b$  is true.

Last time we discussed the set of vectors  $b$  for which  $Ax = b$  has a solution.

## Plan For Today

Today we will learn to describe and draw the solution set of an arbitrary system of linear equations  $Ax = b$ , using spans.



**Recall:** the **solution set** is the collection of all vectors  $x$  such that  $Ax = b$  is true.

Last time we discussed the set of vectors  $b$  for which  $Ax = b$  has a solution.

We also described this set using spans, but it was a *different problem*.

# Homogeneous Systems

Everything is easier when  $b = 0$ , so we start with this case.

Dr Ali and Dr Sara

# Homogeneous Systems

Everything is easier when  $b = 0$ , so we start with this case.

## Definition

A system of linear equations of the form  $Ax = 0$  is called **homogeneous**.



# Homogeneous Systems

Everything is easier when  $b = 0$ , so we start with this case.

## Definition

A system of linear equations of the form  $Ax = 0$  is called **homogeneous**.

These are linear equations where everything to the right of the  $=$  is zero.

# Homogeneous Systems

Everything is easier when  $b = 0$ , so we start with this case.

## Definition

A system of linear equations of the form  $Ax = 0$  is called **homogeneous**.

These are linear equations where everything to the right of the  $=$  is zero.  
The opposite is:

## Definition

A system of linear equations of the form  $Ax = b$  with  $b \neq 0$  is called **nonhomogeneous** or **inhomogeneous**.

# Homogeneous Systems

Everything is easier when  $b = 0$ , so we start with this case.

## Definition

A system of linear equations of the form  $Ax = 0$  is called **homogeneous**.

These are linear equations where everything to the right of the  $=$  is zero.  
The opposite is:

## Definition

A system of linear equations of the form  $Ax = b$  with  $b \neq 0$  is called **nonhomogeneous** or **inhomogeneous**.

A homogeneous system always has the solution  $x = \underline{\quad}$ .

# Homogeneous Systems

Everything is easier when  $b = 0$ , so we start with this case.

## Definition

A system of linear equations of the form  $Ax = 0$  is called **homogeneous**.

These are linear equations where everything to the right of the  $=$  is zero.  
The opposite is:

## Definition

A system of linear equations of the form  $Ax = b$  with  $b \neq 0$  is called **nonhomogeneous** or **inhomogeneous**.

A homogeneous system always has the solution  $x = 0$ . This is called the **trivial solution**.

# Homogeneous Systems

Everything is easier when  $b = 0$ , so we start with this case.

## Definition

A system of linear equations of the form  $Ax = 0$  is called **homogeneous**.

These are linear equations where everything to the right of the  $=$  is zero.  
The opposite is:

## Definition

A system of linear equations of the form  $Ax = b$  with  $b \neq 0$  is called **nonhomogeneous** or **inhomogeneous**.

A homogeneous system always has the solution  $x = 0$ . This is called the **trivial solution**. The nonzero solutions are called **nontrivial**.

# Homogeneous Systems

Everything is easier when  $b = 0$ , so we start with this case.

## Definition

A system of linear equations of the form  $Ax = 0$  is called **homogeneous**.

These are linear equations where everything to the right of the  $=$  is zero.  
The opposite is:

## Definition

A system of linear equations of the form  $Ax = b$  with  $b \neq 0$  is called **nonhomogeneous** or **inhomogeneous**.

A homogeneous system always has the solution  $x = 0$ . This is called the **trivial solution**. The nonzero solutions are called **nontrivial**.

### Observation

$Ax = 0$  has a nontrivial solution



# Homogeneous Systems

Everything is easier when  $b = 0$ , so we start with this case.

## Definition

A system of linear equations of the form  $Ax = 0$  is called **homogeneous**.

These are linear equations where everything to the right of the  $=$  is zero.  
The opposite is:

## Definition

A system of linear equations of the form  $Ax = b$  with  $b \neq 0$  is called **nonhomogeneous** or **inhomogeneous**.

A homogeneous system always has the solution  $x = 0$ . This is called the **trivial solution**. The nonzero solutions are called **nontrivial**.

### Observation

$Ax = 0$  has a nontrivial solution

$\iff$  there is a free variable

$\iff$

# Homogeneous Systems

Everything is easier when  $b = 0$ , so we start with this case.

## Definition

A system of linear equations of the form  $Ax = 0$  is called **homogeneous**.

These are linear equations where everything to the right of the  $=$  is zero. The opposite is:

## Definition

A system of linear equations of the form  $Ax = b$  with  $b \neq 0$  is called **nonhomogeneous** or **inhomogeneous**.

A homogeneous system always has the solution  $x = 0$ . This is called the **trivial solution**. The nonzero solutions are called **nontrivial**.

### Observation

$Ax = 0$  has a nontrivial solution

$\iff$  there is a free variable

$\iff A$  has a column with no pivot.



# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 3 & 4 \\ 2 & -1 & 2 \\ 1 & 0 & 1 \end{pmatrix}?$$

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 3 & 4 \\ 2 & -1 & 2 \\ 1 & 0 & 1 \end{pmatrix}?$$

We know how to do this: first form an augmented matrix and row reduce.

$$\left( \begin{array}{ccc|c} 1 & 3 & 4 & 0 \\ 2 & -1 & 2 & 0 \\ 1 & 0 & 1 & 0 \end{array} \right) \xrightarrow{\text{row reduce}} \left( \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right).$$

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 3 & 4 \\ 2 & -1 & 2 \\ 1 & 0 & 1 \end{pmatrix}?$$

We know how to do this: first form an augmented matrix and row reduce.

$$\left( \begin{array}{ccc|c} 1 & 3 & 4 & 0 \\ 2 & -1 & 2 & 0 \\ 1 & 0 & 1 & 0 \end{array} \right) \xrightarrow{\text{row reduce}} \left( \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right).$$

The only solution is the trivial solution  $x = 0$ .

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 3 & 4 \\ 2 & -1 & 2 \\ 1 & 0 & 1 \end{pmatrix}?$$

We know how to do this: first form an augmented matrix and row reduce.

$$\left( \begin{array}{ccc|c} 1 & 3 & 4 & 0 \\ 2 & -1 & 2 & 0 \\ 1 & 0 & 1 & 0 \end{array} \right) \xrightarrow{\text{row reduce}} \left( \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right).$$

The only solution is the trivial solution  $x = 0$ .

### Observation

Since the last column (everything to the right of the  $=$ ) was zero to begin, it will always stay zero!

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 3 & 4 \\ 2 & -1 & 2 \\ 1 & 0 & 1 \end{pmatrix}?$$

We know how to do this: first form an augmented matrix and row reduce.

$$\left( \begin{array}{ccc|c} 1 & 3 & 4 & 0 \\ 2 & -1 & 2 & 0 \\ 1 & 0 & 1 & 0 \end{array} \right) \xrightarrow{\text{row reduce}} \left( \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right).$$

The only solution is the trivial solution  $x = 0$ .

### Observation

Since the last column (everything to the right of the  $=$ ) was zero to begin, it will always stay zero! So it's not really necessary to write augmented matrices in the homogeneous case.

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}?$$

$$\begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & -3 \\ 0 & 0 \end{pmatrix}$$

$$\xrightarrow{\text{equation}} x_1 - 3x_2 = 0$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 3x_2 \\ x_2 = x_2 \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}.$$

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}?$$

$$\begin{aligned} \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} &\xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & -3 \\ 0 & 0 \end{pmatrix} \\ &\xrightarrow{\text{equation}} x_1 - 3x_2 = 0 \\ &\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 3x_2 \\ x_2 = x_2 \end{cases} \\ &\xrightarrow{\text{parametric vector form}} x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}. \end{aligned}$$

This last equation is called the **parametric vector form** of the solution.

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}?$$

$$\begin{aligned} \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} &\xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & -3 \\ 0 & 0 \end{pmatrix} \\ &\xrightarrow{\text{equation}} x_1 - 3x_2 = 0 \\ &\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 3x_2 \\ x_2 = x_2 \end{cases} \\ &\xrightarrow{\text{parametric vector form}} x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}. \end{aligned}$$

This last equation is called the **parametric vector form** of the solution.

It is obtained by listing equations for all the variables, in order, including the free ones, and making a vector equation.



# Homogeneous Systems

## Example, continued

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}?$$

**Answer:**  $x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$  for any  $x_2$  in  $\mathbb{R}$ .

# Homogeneous Systems

## Example, continued

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}?$$

**Answer:**  $x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$  for any  $x_2$  in  $\mathbb{R}$ . The solution set is  $\text{Span}\left\{ \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\}$

# Homogeneous Systems

## Example, continued

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}?$$

**Answer:**  $x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$  for any  $x_2$  in  $\mathbb{R}$ . The solution set is  $\text{Span}\left\{\begin{pmatrix} 3 \\ 1 \end{pmatrix}\right\}$ .

# Homogeneous Systems


Example, continued

## Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}?$$

**Answer:**  $x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$  for any  $x_2$  in  $\mathbb{R}$ . The solution set is  $\text{Span}\left\{ \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\}$ .



# Homogeneous Systems


## Example, continued

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}?$$

**Answer:**  $x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$  for any  $x_2$  in  $\mathbb{R}$ . The solution set is  $\text{Span}\left\{\begin{pmatrix} 3 \\ 1 \end{pmatrix}\right\}$ .



**Note:** one free variable means the solution set is a *line* in  $\mathbb{R}^2$  ( $2 = \#$  variables  $= \#$  columns).

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix}?$$

$$\begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & -2 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\xrightarrow{\text{equations}} \begin{cases} x_1 - 2x_3 = 0 \\ x_2 + x_3 = 0 \end{cases}$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 2x_3 \\ x_2 = -x_3 \\ x_3 = x_3 \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = x_3 \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}.$$

# Homogeneous Systems

Example, continued

## Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix}?$$

Answer:  $\text{Span} \left\{ \right.$

# Homogeneous Systems

Example, continued

## Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix}?$$

**Answer:**  $\text{Span} \left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\}.$



# Homogeneous Systems


Example, continued

## Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix}?$$

Answer:  $\text{Span} \left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\}$ .



# Homogeneous Systems


## Example, continued

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix}?$$

**Answer:**  $\text{Span} \left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\}$ .



**Note:** one free variable means the solution set is a *line* in  $\mathbb{R}^3$  ( $3 = \#$  variables  $= \#$  columns).

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}?$$

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}$$

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$
$$\xrightarrow{\text{equations}} \begin{cases} x_1 - 8x_3 - 7x_4 = 0 \\ x_2 + 4x_3 + 3x_4 = 0 \end{cases}$$

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$
$$\xrightarrow{\text{equations}} \begin{cases} x_1 - 8x_3 - 7x_4 = 0 \\ x_2 + 4x_3 + 3x_4 = 0 \end{cases}$$
$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 8x_3 + 7x_4 \\ x_2 = -4x_3 - 3x_4 \\ x_3 = x_3 \\ x_4 = x_4 \end{cases}$$

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$
$$\xrightarrow{\text{equations}} \begin{cases} x_1 - 8x_3 - 7x_4 = 0 \\ x_2 + 4x_3 + 3x_4 = 0 \end{cases}$$
$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 8x_3 + 7x_4 \\ x_2 = -4x_3 - 3x_4 \\ x_3 = x_3 \\ x_4 = x_4 \end{cases}$$
$$\xrightarrow{\text{parametric vector form}} x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} =$$



# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\xrightarrow{\text{equations}} \begin{cases} x_1 - 8x_3 - 7x_4 = 0 \\ x_2 + 4x_3 + 3x_4 = 0 \end{cases}$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 8x_3 + 7x_4 \\ x_2 = -4x_3 - 3x_4 \\ x_3 = x_3 \\ x_4 = x_4 \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = x_3 \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix} +$$

# Homogeneous Systems

## Example

### Question

What is the solution set of  $Ax = 0$ , where

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\xrightarrow{\text{equations}} \begin{cases} x_1 - 8x_3 - 7x_4 = 0 \\ x_2 + 4x_3 + 3x_4 = 0 \end{cases}$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 8x_3 + 7x_4 \\ x_2 = -4x_3 - 3x_4 \\ x_3 = x_3 \\ x_4 = x_4 \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = x_3 \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix}.$$

# Homogeneous Systems

## Example, continued

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}?$$

Answer: Span {

# Homogeneous Systems

## Example, continued

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}?$$

**Answer:**  $\text{Span} \left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix} \right\},$

# Homogeneous Systems

## Example, continued

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}?$$

**Answer:**  $\text{Span} \left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \right\}.$

# Homogeneous Systems

## Example, continued

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}?$$

**Answer:**  $\text{Span} \left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \right\}.$

[not pictured here]

# Homogeneous Systems

## Example, continued

### Question

What is the solution set of  $Ax = 0$ , where

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}?$$

**Answer:**  $\text{Span} \left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \right\}.$

[not pictured here]

**Note:** two free variables means the solution set is a *plane* in  $\mathbb{R}^4$  ( $4 = \#$  variables  $= \#$  columns).

# Parametric Vector Form

## Homogeneous systems

Let  $A$  be an  $m \times n$  matrix.

Dr Ali and Dr Sara



# Parametric Vector Form

## Homogeneous systems

Let  $A$  be an  $m \times n$  matrix. Suppose that the free variables in the homogeneous equation  $Ax = 0$  are  $x_i, x_j, x_k, \dots$

# Parametric Vector Form

## Homogeneous systems

Let  $A$  be an  $m \times n$  matrix. Suppose that the free variables in the homogeneous equation  $Ax = 0$  are  $x_i, x_j, x_k, \dots$

Then the solutions to  $Ax = 0$  can be written in the form

$$x = x_i v_i + x_j v_j + x_k v_k + \dots$$

for some vectors  $v_i, v_j, v_k, \dots$  in  $\mathbb{R}^n$ ,

# Parametric Vector Form

## Homogeneous systems

Let  $A$  be an  $m \times n$  matrix. Suppose that the free variables in the homogeneous equation  $Ax = 0$  are  $x_i, x_j, x_k, \dots$

Then the solutions to  $Ax = 0$  can be written in the form

$$x = x_i v_i + x_j v_j + x_k v_k + \dots$$

for some vectors  $v_i, v_j, v_k, \dots$  in  $\mathbb{R}^n$ , and any scalars  $x_i, x_j, x_k, \dots$

# Parametric Vector Form

## Homogeneous systems

Let  $A$  be an  $m \times n$  matrix. Suppose that the free variables in the homogeneous equation  $Ax = 0$  are  $x_i, x_j, x_k, \dots$

Then the solutions to  $Ax = 0$  can be written in the form

$$x = x_i v_i + x_j v_j + x_k v_k + \dots$$

for some vectors  $v_i, v_j, v_k, \dots$  in  $\mathbb{R}^n$ , and any scalars  $x_i, x_j, x_k, \dots$

The solution set is

Span{

# Parametric Vector Form

## Homogeneous systems

Let  $A$  be an  $m \times n$  matrix. Suppose that the free variables in the homogeneous equation  $Ax = 0$  are  $x_i, x_j, x_k, \dots$

Then the solutions to  $Ax = 0$  can be written in the form

$$x = x_i v_i + x_j v_j + x_k v_k + \dots$$

for some vectors  $v_i, v_j, v_k, \dots$  in  $\mathbb{R}^n$ , and any scalars  $x_i, x_j, x_k, \dots$

The solution set is

$$\text{Span}\{v_i, v_j, v_k, \dots\}.$$

# Parametric Vector Form

## Homogeneous systems

Let  $A$  be an  $m \times n$  matrix. Suppose that the free variables in the homogeneous equation  $Ax = 0$  are  $x_i, x_j, x_k, \dots$

Then the solutions to  $Ax = 0$  can be written in the form

$$x = x_i v_i + x_j v_j + x_k v_k + \dots$$

for some vectors  $v_i, v_j, v_k, \dots$  in  $\mathbb{R}^n$ , and any scalars  $x_i, x_j, x_k, \dots$

The solution set is

$$\text{Span}\{v_i, v_j, v_k, \dots\}.$$

The **equation** above is called the **parametric vector form** of the solution.

## Poll

How many solutions can there be to a homogeneous system with more equations than variables?

- A. 0
- B. 1
- C.  $\infty$

## Poll

How many solutions can there be to a homogeneous system with more equations than variables?

- A. 0
- B. 1
- C.  $\infty$

The trivial solution is always a solution to a homogeneous system, so answer A is impossible.



## Poll

How many solutions can there be to a homogeneous system with more equations than variables?

- A. 0
- B. 1
- C.  $\infty$

The trivial solution is always a solution to a homogeneous system, so answer A is impossible.

This matrix has only one solution to  $Ax = 0$ :

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

## Poll

How many solutions can there be to a homogeneous system with more equations than variables?

- A. 0
- B. 1
- C.  $\infty$

The trivial solution is always a solution to a homogeneous system, so answer A is impossible.

This matrix has only one solution to  $Ax = 0$ :

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

This matrix has infinitely many solutions to  $Ax = 0$ :

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}$$

# Nonhomogeneous Systems

## Example

### Question

What is the solution set of  $Ax = b$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

$$\left( \begin{array}{cc|c} 1 & -3 & -3 \\ 2 & -6 & -6 \end{array} \right) \xrightarrow{\text{row reduce}} \left( \begin{array}{cc|c} 1 & -3 & -3 \\ 0 & 0 & 0 \end{array} \right)$$

$$\xrightarrow{\text{equation}} x_1 - 3x_2 = -3$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 3x_2 - 3 \\ x_2 = x_2 + 0 \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}.$$

# Nonhomogeneous Systems

## Example

### Question

What is the solution set of  $Ax = b$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

$$\left( \begin{array}{cc|c} 1 & -3 & -3 \\ 2 & -6 & -6 \end{array} \right) \xrightarrow{\text{row reduce}} \left( \begin{array}{cc|c} 1 & -3 & -3 \\ 0 & 0 & 0 \end{array} \right)$$

$$\xrightarrow{\text{equation}} x_1 - 3x_2 = -3$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 3x_2 - 3 \\ x_2 = x_2 + 0 \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}.$$

The only difference from the homogeneous case is the constant vector  $p = \begin{pmatrix} -3 \\ 0 \end{pmatrix}$ .

# Nonhomogeneous Systems

## Example

### Question

What is the solution set of  $Ax = b$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

$$\left( \begin{array}{cc|c} 1 & -3 & -3 \\ 2 & -6 & -6 \end{array} \right) \xrightarrow{\text{row reduce}} \left( \begin{array}{cc|c} 1 & -3 & -3 \\ 0 & 0 & 0 \end{array} \right)$$

$$\xrightarrow{\text{equation}} x_1 - 3x_2 = -3$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 3x_2 - 3 \\ x_2 = x_2 + 0 \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}.$$

The only difference from the homogeneous case is the constant vector  $p = \begin{pmatrix} -3 \\ 0 \end{pmatrix}$ .

Note that  $p$  is itself a solution: take  $x_2 = 0$ .

# Nonhomogeneous Systems

Example, continued

## Question

What is the solution set of  $Ax = b$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

**Answer:**  $x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}$  for any  $x_2$  in  $\mathbb{R}$ .

# Nonhomogeneous Systems

## Example, continued

### Question

What is the solution set of  $Ax = b$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

**Answer:**  $x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}$  for any  $x_2$  in  $\mathbb{R}$ .

This is a *translate* of  $\text{Span}\left\{\begin{pmatrix} 3 \\ 1 \end{pmatrix}\right\}$ : it is the parallel line through  $p = \begin{pmatrix} -3 \\ 0 \end{pmatrix}$ .

# Nonhomogeneous Systems

Example, continued


## Question

What is the solution set of  $Ax = b$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

**Answer:**  $x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}$  for any  $x_2$  in  $\mathbb{R}$ .

This is a *translate* of  $\text{Span}\left\{\begin{pmatrix} 3 \\ 1 \end{pmatrix}\right\}$ : it is the parallel line through  $p = \begin{pmatrix} -3 \\ 0 \end{pmatrix}$ .





# Nonhomogeneous Systems

## Example, continued


### Question

What is the solution set of  $Ax = b$ , where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

**Answer:**  $x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}$  for any  $x_2$  in  $\mathbb{R}$ .

This is a *translate* of  $\text{Span}\left\{\begin{pmatrix} 3 \\ 1 \end{pmatrix}\right\}$ : it is the parallel line through  $p = \begin{pmatrix} -3 \\ 0 \end{pmatrix}$ .



It can be written

$$\text{Span}\left\{\begin{pmatrix} 3 \\ 1 \end{pmatrix}\right\} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}.$$

# Nonhomogeneous Systems

## Example

### Question

What is the solution set of  $Ax = b$ , where

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -5 \\ -3 \\ -2 \end{pmatrix}?$$

$$\left( \begin{array}{ccc|c} 1 & 3 & 1 & -5 \\ 2 & -1 & -5 & -3 \\ 1 & 0 & -2 & -2 \end{array} \right) \xrightarrow{\text{row reduce}} \left( \begin{array}{ccc|c} 1 & 0 & -2 & -2 \\ 0 & 1 & 1 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

$$\xrightarrow{\text{equations}} \begin{cases} x_1 - 2x_3 = -2 \\ x_2 + x_3 = -1 \end{cases}$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 2x_3 - 2 \\ x_2 = -x_3 - 1 \\ x_3 = x_3 \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = x_3 \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} + \begin{pmatrix} -2 \\ -1 \\ 0 \end{pmatrix}.$$

# Nonhomogeneous Systems

Example, continued

## Question

What is the solution set of  $Ax = b$ , where

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -5 \\ -3 \\ -2 \end{pmatrix}?$$

**Answer:**  $\text{Span} \left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\} +$

# Nonhomogeneous Systems

Example, continued

## Question

What is the solution set of  $Ax = b$ , where

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -5 \\ -3 \\ -2 \end{pmatrix}?$$

**Answer:**  $\text{Span} \left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\} + \begin{pmatrix} -2 \\ -1 \\ 0 \end{pmatrix}.$

# Nonhomogeneous Systems


Example, continued

## Question

What is the solution set of  $Ax = b$ , where

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -5 \\ -3 \\ -2 \end{pmatrix}?$$

**Answer:**  $\text{Span} \left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\} + \begin{pmatrix} -2 \\ -1 \\ 0 \end{pmatrix}.$



The solution set is a *translate* of

$$\text{Span} \left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\} :$$

it is the parallel line through

$$p = \begin{pmatrix} -2 \\ -1 \\ 0 \end{pmatrix}.$$

## Homogeneous vs. Nonhomogeneous Systems

### Key Observation

The set of solutions to  $Ax = b$ , if it is nonempty, is obtained by taking one **specific** or **particular solution**  $p$  to  $Ax = b$ , and adding all solutions to  $Ax = 0$ .

# Homogeneous vs. Nonhomogeneous Systems

## Key Observation

The set of solutions to  $Ax = b$ , if it is nonempty, is obtained by taking one **specific** or **particular solution**  $p$  to  $Ax = b$ , and adding all solutions to  $Ax = 0$ .

**Why?** If  $Ap = b$  and  $Ax = 0$ , then

$$A(p + x) = Ap + Ax = b + 0 = b,$$

## Homogeneous vs. Nonhomogeneous Systems

### Key Observation

The set of solutions to  $Ax = b$ , if it is nonempty, is obtained by taking one **specific** or **particular solution**  $p$  to  $Ax = b$ , and adding all solutions to  $Ax = 0$ .

**Why?** If  $Ap = b$  and  $Ax = 0$ , then

$$A(p + x) = Ap + Ax = b + 0 = b,$$

so  $p + x$  is also a solution to  $Ax = b$ .



## Homogeneous vs. Nonhomogeneous Systems

### Key Observation

The set of solutions to  $Ax = b$ , if it is nonempty, is obtained by taking one **specific** or **particular solution**  $p$  to  $Ax = b$ , and adding all solutions to  $Ax = 0$ .

**Why?** If  $Ap = b$  and  $Ax = 0$ , then

$$A(p + x) = Ap + Ax = b + 0 = b,$$

so  $p + x$  is also a solution to  $Ax = b$ .

We know the solution set of  $Ax = 0$  is a span.

## Homogeneous vs. Nonhomogeneous Systems

### Key Observation

The set of solutions to  $Ax = b$ , if it is nonempty, is obtained by taking one **specific** or **particular solution**  $p$  to  $Ax = b$ , and adding all solutions to  $Ax = 0$ .

**Why?** If  $Ap = b$  and  $Ax = 0$ , then

$$A(p + x) = Ap + Ax = b + 0 = b,$$

so  $p + x$  is also a solution to  $Ax = b$ .

We know the solution set of  $Ax = 0$  is a span. So the solution set of  $Ax = b$  is a *translate* of a span: it is *parallel* to a span.

## Homogeneous vs. Nonhomogeneous Systems

### Key Observation

The set of solutions to  $Ax = b$ , if it is nonempty, is obtained by taking one **specific** or **particular solution**  $p$  to  $Ax = b$ , and adding all solutions to  $Ax = 0$ .

**Why?** If  $Ap = b$  and  $Ax = 0$ , then

$$A(p + x) = Ap + Ax = b + 0 = b,$$

so  $p + x$  is also a solution to  $Ax = b$ .

We know the solution set of  $Ax = 0$  is a span. So the solution set of  $Ax = b$  is a *translate* of a span: it is *parallel* to a span. (Or it is empty.)

## Homogeneous vs. Nonhomogeneous Systems

### Key Observation


The set of solutions to  $Ax = b$ , if it is nonempty, is obtained by taking one **specific** or **particular solution**  $p$  to  $Ax = b$ , and adding all solutions to  $Ax = 0$ .

**Why?** If  $Ap = b$  and  $Ax = 0$ , then

$$A(p + x) = Ap + Ax = b + 0 = b,$$

so  $p + x$  is also a solution to  $Ax = b$ .

We know the solution set of  $Ax = 0$  is a span. So the solution set of  $Ax = b$  is a *translate* of a span: it is *parallel* to a span. (Or it is empty.)



This works for *any* specific solution  $p$ : it doesn't have to be the one produced by finding the parametric vector form and setting the free variables all to zero, as we did before.

# Homogeneous vs. Nonhomogeneous Systems

Varying  $b$

If we understand the solution set of  $Ax = 0$ , then we understand the solution set of  $Ax = b$  for all  $b$ : they are all translates (or empty).


Dr Ali and Dr Sara

# Homogeneous vs. Nonhomogeneous Systems

## Varying $b$

If we understand the solution set of  $Ax = 0$ , then we understand the solution set of  $Ax = b$  for all  $b$ : they are all translates (or empty).

For instance, if  $A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$ , then the solution sets for varying  $b$  look like this:




# Homogeneous vs. Nonhomogeneous Systems

## Varying $b$

If we understand the solution set of  $Ax = 0$ , then we understand the solution set of  $Ax = b$  for all  $b$ : they are all translates (or empty).

For instance, if  $A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$ , then the solution sets for varying  $b$  look like this:




Which  $b$  gives the solution set  $Ax = b$  in red in the picture?

# Homogeneous vs. Nonhomogeneous Systems

Varying  $b$

If we understand the solution set of  $Ax = 0$ , then we understand the solution set of  $Ax = b$  for all  $b$ : they are all translates (or empty).

For instance, if  $A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$ , then the solution sets for varying  $b$  look like this:



Which  $b$  gives the solution set  $Ax = b$  in red in the picture?

Choose  $p$  on the red line, and set  $b = Ap$ . Then  $p$  is a specific solution to  $Ax = b$ , so the solution set of  $Ax = b$  is the red line.




# Homogeneous vs. Nonhomogeneous Systems

Varying  $b$

If we understand the solution set of  $Ax = 0$ , then we understand the solution set of  $Ax = b$  for all  $b$ : they are all translates (or empty).

For instance, if  $A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$ , then the solution sets for varying  $b$  look like this:



Which  $b$  gives the solution set  $Ax = b$  in red in the picture?

Choose  $p$  on the red line, and set  $b = Ap$ . Then  $p$  is a specific solution to  $Ax = b$ , so the solution set of  $Ax = b$  is the red line.


Note the cool optical illusion!

# Homogeneous vs. Nonhomogeneous Systems

## Varying $b$

If we understand the solution set of  $Ax = 0$ , then we understand the solution set of  $Ax = b$  for all  $b$ : they are all translates (or empty).

For instance, if  $A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$ , then the solution sets for varying  $b$  look like this:



Which  $b$  gives the solution set  $Ax = b$  in red in the picture?

Choose  $p$  on the red line, and set  $b = Ap$ . Then  $p$  is a specific solution to  $Ax = b$ , so the solution set of  $Ax = b$  is the red line.

Note the cool optical illusion!

For a matrix equation  $Ax = b$ , you now know how to find which  $b$ 's are possible, and what the solution set looks like for all  $b$ , both using spans.

# Section 1.7

## Linear Independence


## Motivation

Sometimes the span of a set of vectors is “smaller” than you expect from the number of vectors.

Dr Ali and Dr Sara



## Motivation

Sometimes the span of a set of vectors is “smaller” than you expect from the number of vectors.





## Motivation

Sometimes the span of a set of vectors is “smaller” than you expect from the number of vectors.



## Motivation



Sometimes the span of a set of vectors is “smaller” than you expect from the number of vectors.



This can mean many things. For example, it can mean you're using too many vectors to write your solution set.

## Motivation

Sometimes the span of a set of vectors is “smaller” than you expect from the number of vectors.





This can mean many things. For example, it can mean you're using too many vectors to write your solution set.

Notice in each case that one vector in the set is already in the span of the others—so it doesn't make the span bigger.



## Motivation

Sometimes the span of a set of vectors is “smaller” than you expect from the number of vectors.



This can mean many things. For example, it can mean you're using too many vectors to write your solution set.

Notice in each case that one vector in the set is already in the span of the others—so it doesn't make the span bigger.

Today we will formalize this idea in the concept of *linear (in)dependence*.

# Linear Independence

## Definition

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  in  $\mathbb{R}^n$  is **linearly independent** if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution  $x_1 = x_2 = \dots = x_p = 0$ .

# Linear Independence

## Definition

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  in  $\mathbb{R}^n$  is **linearly independent** if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution  $x_1 = x_2 = \dots = x_p = 0$ . The set  $\{v_1, v_2, \dots, v_p\}$  is **linearly dependent** otherwise.

# Linear Independence

## Definition

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  in  $\mathbb{R}^n$  is **linearly independent** if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution  $x_1 = x_2 = \dots = x_p = 0$ . The set  $\{v_1, v_2, \dots, v_p\}$  is **linearly dependent** otherwise.

In other words,  $\{v_1, v_2, \dots, v_p\}$  is linearly dependent if

# Linear Independence

## Definition

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  in  $\mathbb{R}^n$  is **linearly independent** if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution  $x_1 = x_2 = \dots = x_p = 0$ . The set  $\{v_1, v_2, \dots, v_p\}$  is **linearly dependent** otherwise.

In other words,  $\{v_1, v_2, \dots, v_p\}$  is linearly dependent if there exist numbers  $x_1, x_2, \dots, x_p$ , not all equal to zero, such that

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0.$$

This is called a **linear dependence relation**.

# Linear Independence

## Definition

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  in  $\mathbb{R}^n$  is **linearly independent** if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution  $x_1 = x_2 = \dots = x_p = 0$ . The set  $\{v_1, v_2, \dots, v_p\}$  is **linearly dependent** otherwise.

In other words,  $\{v_1, v_2, \dots, v_p\}$  is linearly dependent if there exist numbers  $x_1, x_2, \dots, x_p$ , not all equal to zero, such that

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0.$$

This is called a **linear dependence relation**.

Like span, linear (in)dependence is another one of those big vocabulary words that you absolutely need to learn. Much of the rest of the course will be built on these concepts, and you need to know exactly what they mean in order to be able to answer questions on quizzes and exams (and solve real-world problems later on).

# Linear Independence

## Definition

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  in  $\mathbb{R}^n$  is **linearly independent** if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution  $x_1 = x_2 = \dots = x_p = 0$ . The set  $\{v_1, v_2, \dots, v_p\}$  is **linearly dependent** otherwise.

Note that linear (in)dependence is a notion that applies to a *collection of vectors*, not to a single vector, or to one vector in the presence of some others.

## Checking Linear Independence

**Question:** Is  $\left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} \right\}$  linearly independent?

Equivalently, does the (homogeneous) the vector equation

$$x \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + y \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} + z \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

have a nontrivial solution?



## Checking Linear Independence

**Question:** Is  $\left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} \right\}$  linearly independent?

Equivalently, does the (homogeneous) the vector equation

$$x \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + y \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} + z \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

have a nontrivial solution?

How do we solve this kind of vector equation?

$$\begin{pmatrix} 1 & 1 & 3 \\ 1 & -1 & 1 \\ 1 & 2 & 4 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

## Checking Linear Independence

**Question:** Is  $\left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} \right\}$  linearly independent?

Equivalently, does the (homogeneous) the vector equation

$$x \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + y \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} + z \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

have a nontrivial solution?

How do we solve this kind of vector equation?

$$\begin{pmatrix} 1 & 1 & 3 \\ 1 & -1 & 1 \\ 1 & 2 & 4 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

So  $x = -2z$  and  $y = -z$ . So the vectors are linearly *dependent*, and an equation of linear dependence is (taking  $z = 1$ )

## Checking Linear Independence

**Question:** Is  $\left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} \right\}$  linearly independent?

Equivalently, does the (homogeneous) the vector equation

$$x \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + y \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} + z \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

have a nontrivial solution?

How do we solve this kind of vector equation?

$$\begin{pmatrix} 1 & 1 & 3 \\ 1 & -1 & 1 \\ 1 & 2 & 4 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

So  $x = -2z$  and  $y = -z$ . So the vectors are linearly dependent, and an equation of linear dependence is (taking  $z = 1$ )

$$-2 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} - \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} + \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

## Checking Linear Independence

**Question:** Is  $\left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} \right\}$  linearly independent?

Equivalently, does the (homogeneous) the vector equation

$$x \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + y \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} + z \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

have a nontrivial solution?

## Checking Linear Independence

**Question:** Is  $\left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} \right\}$  linearly independent?

Equivalently, does the (homogeneous) the vector equation

$$x \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + y \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} + z \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

have a nontrivial solution?

$$\begin{pmatrix} 1 & 1 & 3 \\ 1 & -1 & 1 \\ 0 & 2 & 4 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

## Checking Linear Independence

**Question:** Is  $\left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} \right\}$  linearly independent?

Equivalently, does the (homogeneous) the vector equation

$$x \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + y \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} + z \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

have a nontrivial solution?

$$\begin{pmatrix} 1 & 1 & 3 \\ 1 & -1 & 1 \\ 0 & 2 & 4 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

The trivial solution  $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$  is the unique solution. So the vectors are linearly *independent*.

## Linear Independence and Matrix Columns

In general,  $\{v_1, v_2, \dots, v_p\}$  is linearly independent

Dr Ali and Dr Sara

## Linear Independence and Matrix Columns

In general,  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution,



## Linear Independence and Matrix Columns

In general,  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution, if and only if the matrix equation

$$Ax = 0$$

has only the trivial solution, where  $A$  is the matrix

## Linear Independence and Matrix Columns

In general,  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution, if and only if the matrix equation

$$Ax = 0$$

has only the trivial solution, where  $A$  is the matrix with columns  $v_1, v_2, \dots, v_p$ :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & \cdots & | \end{pmatrix}.$$

## Linear Independence and Matrix Columns

In general,  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution, if and only if the matrix equation

$$Ax = 0$$

has only the trivial solution, where  $A$  is the matrix with columns  $v_1, v_2, \dots, v_p$ :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & \cdots & | \end{pmatrix}.$$

This is true if and only if the matrix  $A$  has a pivot in each \_\_\_\_\_.

## Linear Independence and Matrix Columns

In general,  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution, if and only if the matrix equation

$$Ax = 0$$

has only the trivial solution, where  $A$  is the matrix with columns  $v_1, v_2, \dots, v_p$ :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & \cdots & | \end{pmatrix}.$$

This is true if and only if the matrix  $A$  has a pivot in each column.

## Linear Independence and Matrix Columns

In general,  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution, if and only if the matrix equation

$$Ax = 0$$

has only the trivial solution, where  $A$  is the matrix with columns  $v_1, v_2, \dots, v_p$ :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & & | \end{pmatrix}.$$

This is true if and only if the matrix  $A$  has a pivot in each column.

### Important

- ▶ The vectors  $v_1, v_2, \dots, v_p$  are linearly independent if and only if the matrix with columns  $v_1, v_2, \dots, v_p$  has a pivot in each column.

## Linear Independence and Matrix Columns

In general,  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution, if and only if the matrix equation

$$Ax = 0$$

has only the trivial solution, where  $A$  is the matrix with columns  $v_1, v_2, \dots, v_p$ :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & \cdots & | \end{pmatrix}.$$

This is true if and only if the matrix  $A$  has a pivot in each column.

### Important

- ▶ The vectors  $v_1, v_2, \dots, v_p$  are linearly independent if and only if the matrix with columns  $v_1, v_2, \dots, v_p$  has a pivot in each column.
- ▶ Solving the matrix equation  $Ax = 0$  will either verify that the columns  $v_1, v_2, \dots, v_p$  of  $A$  are linearly independent, or will produce a linear dependence relation.

# Linear Independence

## Criterion

Suppose that one of the vectors  $\{v_1, v_2, \dots, v_p\}$  is a linear combination of the other ones (that is, it is in the span of the other ones):

# Linear Independence

## Criterion

Suppose that one of the vectors  $\{v_1, v_2, \dots, v_p\}$  is a linear combination of the other ones (that is, it is in the span of the other ones):

$$v_3 = 2v_1 - \frac{1}{2}v_2 + 6v_4$$

Then the vectors are linearly *dependent*:

$$2v_1 - \frac{1}{2}v_2 - v_3 + 6v_4 = 0.$$



# Linear Independence

## Criterion

Suppose that one of the vectors  $\{v_1, v_2, \dots, v_p\}$  is a linear combination of the other ones (that is, it is in the span of the other ones):

$$v_3 = 2v_1 - \frac{1}{2}v_2 + 6v_4$$

Then the vectors are linearly *dependent*:

$$2v_1 - \frac{1}{2}v_2 - v_3 + 6v_4 = 0.$$

Conversely, if the vectors are linearly dependent

$$2v_1 - \frac{1}{2}v_2 + 6v_4 = 0.$$

# Linear Independence

## Criterion

Suppose that one of the vectors  $\{v_1, v_2, \dots, v_p\}$  is a linear combination of the other ones (that is, it is in the span of the other ones):

$$v_3 = 2v_1 - \frac{1}{2}v_2 + 6v_4$$

Then the vectors are linearly *dependent*:

$$2v_1 - \frac{1}{2}v_2 - v_3 + 6v_4 = 0.$$

Conversely, if the vectors are linearly dependent

$$2v_1 - \frac{1}{2}v_2 + 6v_4 = 0.$$

then one vector is a linear combination of (in the span of) the other ones:

$$v_2 = 4v_1 + 12v_4.$$

# Linear Independence

## Criterion

Suppose that one of the vectors  $\{v_1, v_2, \dots, v_p\}$  is a linear combination of the other ones (that is, it is in the span of the other ones):

$$v_3 = 2v_1 - \frac{1}{2}v_2 + 6v_4$$

Then the vectors are linearly *dependent*:

$$2v_1 - \frac{1}{2}v_2 - v_3 + 6v_4 = 0.$$

Conversely, if the vectors are linearly dependent

$$2v_1 - \frac{1}{2}v_2 + 6v_4 = 0.$$

then one vector is a linear combination of (in the span of) the other ones:


$$v_2 = 4v_1 + 12v_4.$$

## Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly *dependent* if and only if one of the vectors is in the span of the other ones.

# Linear Independence

Pictures in  $\mathbb{R}^2$




In this picture

---

One vector  $\{v\}$ :

# Linear Independence

Pictures in  $\mathbb{R}^2$



In this picture


---

One vector  $\{v\}$ :

Linearly independent if  $v \neq 0$ .

# Linear Independence

Pictures in  $\mathbb{R}^2$



In this picture

---


One vector  $\{v\}$ :

Linearly independent if  $v \neq 0$ .

Two vectors  $\{v, w\}$ :

# Linear Independence

Pictures in  $\mathbb{R}^2$



In this picture

---

One vector  $\{v\}$ :


Linearly independent if  $v \neq 0$ .

Two vectors  $\{v, w\}$ :

Linearly independent: neither is in the span of the other.

# Linear Independence

Pictures in  $\mathbb{R}^2$



In this picture

---

One vector  $\{v\}$ :

Linearly independent if  $v \neq 0$ .

Two vectors  $\{v, w\}$ :


Linearly independent: neither is in the span of the other.

Three vectors  $\{v, w, u\}$ :



# Linear Independence

Pictures in  $\mathbb{R}^2$



In this picture

---

**One vector  $\{v\}$ :**

Linearly independent if  $v \neq 0$ .

**Two vectors  $\{v, w\}$ :**


Linearly independent: neither is in the span of the other.

**Three vectors  $\{v, w, u\}$ :**

Linearly dependent:  $u$  is in  $\text{Span}\{v, w\}$ .

# Linear Independence

Pictures in  $\mathbb{R}^2$



In this picture

**One vector  $\{v\}$ :**

Linearly independent if  $v \neq 0$ .

**Two vectors  $\{v, w\}$ :**

Linearly independent: neither is in the span of the other.


**Three vectors  $\{v, w, u\}$ :**

Linearly dependent:  $u$  is in  $\text{Span}\{v, w\}$ .

Also  $v$  is in  $\text{Span}\{u, w\}$  and  $w$  is in  $\text{Span}\{u, v\}$ .

# Linear Independence


Pictures in  $\mathbb{R}^2$



Two collinear vectors  $\{v, w\}$ :

# Linear Independence


Pictures in  $\mathbb{R}^2$



Two collinear vectors  $\{v, w\}$ :  
Linearly dependent:  $w$  is in  
 $\text{Span}\{v\}$  (and vice-versa).

# Linear Independence

Pictures in  $\mathbb{R}^2$




Two collinear vectors  $\{v, w\}$ :

Linearly dependent:  $w$  is in  $\text{Span}\{v\}$  (and vice-versa).

**Observe:** Two vectors are linearly dependent if and only if

# Linear Independence

Pictures in  $\mathbb{R}^2$




Two collinear vectors  $\{v, w\}$ :

Linearly dependent:  $w$  is in  $\text{Span}\{v\}$  (and vice-versa).

**Observe:** Two vectors are linearly dependent if and only if they are *collinear*.

# Linear Independence

Pictures in  $\mathbb{R}^2$



Two collinear vectors  $\{v, w\}$ :


Linearly dependent:  $w$  is in  $\text{Span}\{v\}$  (and vice-versa).

**Observe:** Two vectors are linearly dependent if and only if they are *collinear*.

Three vectors  $\{v, w, u\}$ :

# Linear Independence

Pictures in  $\mathbb{R}^2$



**Two collinear vectors  $\{v, w\}$ :**

Linearly dependent:  $w$  is in  $\text{Span}\{v\}$  (and vice-versa).

**Observe:** Two vectors are linearly dependent if and only if they are *collinear*.


**Three vectors  $\{v, w, u\}$ :**

Linearly dependent:  $w$  is in  $\text{Span}\{v\}$  (and vice-versa).



# Linear Independence

Pictures in  $\mathbb{R}^2$



**Two collinear vectors  $\{v, w\}$ :**

Linearly dependent:  $w$  is in  $\text{Span}\{v\}$  (and vice-versa).

**Observe:** Two vectors are linearly dependent if and only if they are *collinear*.


**Three vectors  $\{v, w, u\}$ :**

Linearly dependent:  $w$  is in  $\text{Span}\{v\}$  (and vice-versa).

**Observe:** If a set of vectors is linearly dependent, then so is any larger set of vectors!

# Linear Independence

Pictures in  $\mathbb{R}^3$




In this picture

---

Two vectors  $\{v, w\}$ :

# Linear Independence

Pictures in  $\mathbb{R}^3$




In this picture

Two vectors  $\{v, w\}$ :

Linearly independent: neither is in the span of the other.

# Linear Independence

Pictures in  $\mathbb{R}^3$



In this picture

---


Two vectors  $\{v, w\}$ :

Linearly independent: neither is in the span of the other.

Three vectors  $\{v, w, u\}$ :

# Linear Independence

Pictures in  $\mathbb{R}^3$



In this picture

Two vectors  $\{v, w\}$ :


Linearly independent: neither is in the span of the other.

Three vectors  $\{v, w, u\}$ :

Linearly independent: no one is in the span of the other two.

# Linear Independence

Pictures in  $\mathbb{R}^3$



In this picture

---


Two vectors  $\{v, w\}$ :

Linearly independent: neither is in the span of the other.

Three vectors  $\{v, w, x\}$ :

# Linear Independence

Pictures in  $\mathbb{R}^3$



In this picture

---

Two vectors  $\{v, w\}$ :

Linearly independent: neither is in the span of the other.

Three vectors  $\{v, w, x\}$ :

Linearly dependent:  $x$  is in  $\text{Span}\{v, w\}$ .

## Poll

Are there four vectors  $u, v, w, x$  in  $\mathbb{R}^3$  which are linearly dependent, but such that  $u$  is *not* a linear combination of  $v, w, x$ ? If so, draw a picture; if not, give an argument.



## Poll

Are there four vectors  $u, v, w, x$  in  $\mathbb{R}^3$  which are linearly dependent, but such that  $u$  is *not* a linear combination of  $v, w, x$ ? If so, draw a picture; if not, give an argument.

**Yes:** actually the pictures on the previous slides provide such an example.

## Poll

Are there four vectors  $u, v, w, x$  in  $\mathbb{R}^3$  which are linearly dependent, but such that  $u$  is *not* a linear combination of  $v, w, x$ ? If so, draw a picture; if not, give an argument.

**Yes:** actually the pictures on the previous slides provide such an example.

Linear dependence of  $\{v_1, \dots, v_p\}$  means *some*  $v_i$  is a linear combination of the others, not *any*.

# Linear Independence

## Stronger criterion

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly *dependent* if and only if one of the vectors is in the span of the other ones.

# Linear Independence

## Stronger criterion

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly *dependent* if and only if one of the vectors is in the span of the other ones.

Take the largest  $j$  such that  $v_j$  is in the span of the others.

# Linear Independence

## Stronger criterion

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly *dependent* if and only if one of the vectors is in the span of the other ones.

Take the largest  $j$  such that  $v_j$  is in the span of the others. Then  $v_j$  is in the span of  $v_1, v_2, \dots, v_{j-1}$ . Why?

# Linear Independence

## Stronger criterion

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly *dependent* if and only if one of the vectors is in the span of the other ones.

Take the largest  $j$  such that  $v_j$  is in the span of the others. Then  $v_j$  is in the span of  $v_1, v_2, \dots, v_{j-1}$ . Why? If not ( $j = 3$ ):

$$v_3 = 2v_1 - \frac{1}{2}v_2 + 6v_4$$

# Linear Independence

## Stronger criterion

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly *dependent* if and only if one of the vectors is in the span of the other ones.

Take the largest  $j$  such that  $v_j$  is in the span of the others. Then  $v_j$  is in the span of  $v_1, v_2, \dots, v_{j-1}$ . Why? If not ( $j = 3$ ):

$$v_3 = 2v_1 - \frac{1}{2}v_2 + 6v_4$$

Rearrange:

$$v_4 = -\frac{1}{6} \left( 2v_1 - \frac{1}{2}v_2 - v_3 \right)$$

# Linear Independence

## Stronger criterion

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly dependent if and only if one of the vectors is in the span of the other ones.

Take the largest  $j$  such that  $v_j$  is in the span of the others. Then  $v_j$  is in the span of  $v_1, v_2, \dots, v_{j-1}$ . Why? If not ( $j = 3$ ):

$$v_3 = 2v_1 - \frac{1}{2}v_2 + 6v_4$$

Rearrange:

$$v_4 = -\frac{1}{6} \left( 2v_1 - \frac{1}{2}v_2 - v_3 \right)$$

so  $v_4$  works as well, but  $v_3$  was supposed to be the last one that was in the span of the others.



# Linear Independence

## Stronger criterion

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly dependent if and only if one of the vectors is in the span of the other ones.

Take the largest  $j$  such that  $v_j$  is in the span of the others. Then  $v_j$  is in the span of  $v_1, v_2, \dots, v_{j-1}$ . Why? If not ( $j = 3$ ):

$$v_3 = 2v_1 - \frac{1}{2}v_2 + 6v_4$$

Rearrange:

$$v_4 = -\frac{1}{6} \left( 2v_1 - \frac{1}{2}v_2 - v_3 \right)$$

so  $v_4$  works as well, but  $v_3$  was supposed to be the last one that was in the span of the others.

### Better Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly dependent if and only if there is some  $j$  such that  $v_j$  is in  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

# Linear Independence

## Increasing span criterion

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly dependent if and only if there is some  $j$  such that  $v_j$  is in  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

# Linear Independence

## Increasing span criterion

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly dependent if and only if there is some  $j$  such that  $v_j$  is in  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

Equivalently,  $\{v_1, v_2, \dots, v_p\}$  is linearly *independent* if for every  $j$ , the vector  $v_j$  is not in  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

# Linear Independence

## Increasing span criterion

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly dependent if and only if there is some  $j$  such that  $v_j$  is in  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

Equivalently,  $\{v_1, v_2, \dots, v_p\}$  is linearly *independent* if for every  $j$ , the vector  $v_j$  is not in  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

This means  $\text{Span}\{v_1, v_2, \dots, v_j\}$  is *bigger* than  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

# Linear Independence

## Increasing span criterion

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly dependent if and only if there is some  $j$  such that  $v_j$  is in  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

Equivalently,  $\{v_1, v_2, \dots, v_p\}$  is linearly *independent* if for every  $j$ , the vector  $v_j$  is not in  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

This means  $\text{Span}\{v_1, v_2, \dots, v_j\}$  is *bigger* than  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if, for every  $j$ , the span of  $v_1, v_2, \dots, v_j$  is strictly larger than the span of  $v_1, v_2, \dots, v_{j-1}$ .

# Linear Independence

## Increasing span criterion

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly dependent if and only if there is some  $j$  such that  $v_j$  is in  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

Equivalently,  $\{v_1, v_2, \dots, v_p\}$  is linearly *independent* if for every  $j$ , the vector  $v_j$  is not in  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

This means  $\text{Span}\{v_1, v_2, \dots, v_j\}$  is *bigger* than  $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$ .

### Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if, for every  $j$ , the span of  $v_1, v_2, \dots, v_j$  is strictly larger than the span of  $v_1, v_2, \dots, v_{j-1}$ .

#### Translation


A set of vectors is linearly independent if and only if, every time you add another vector to the set, the span gets bigger.

# Linear Independence

Increasing span criterion: pictures

## Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if, for every  $j$ , the span of  $v_1, v_2, \dots, v_j$  is strictly larger than the span of  $v_1, v_2, \dots, v_{j-1}$ .



One vector  $\{v\}$ :


Linearly independent: span got bigger (than  $\{0\}$ ).

# Linear Independence

Increasing span criterion: pictures

## Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if, for every  $j$ , the span of  $v_1, v_2, \dots, v_j$  is strictly larger than the span of  $v_1, v_2, \dots, v_{j-1}$ .



One vector  $\{v\}$ :

Linearly independent: span got bigger (than  $\{0\}$ ).

Two vectors  $\{v, w\}$ :

Linearly independent: span got bigger.




# Linear Independence

Increasing span criterion: pictures

## Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if, for every  $j$ , the span of  $v_1, v_2, \dots, v_j$  is strictly larger than the span of  $v_1, v_2, \dots, v_{j-1}$ .



**One vector  $\{v\}$ :**

Linearly independent: span got bigger (than  $\{0\}$ ).

**Two vectors  $\{v, w\}$ :**

Linearly independent: span got bigger.

**Three vectors  $\{v, w, u\}$ :**


Linearly independent: span got bigger.

# Linear Independence

Increasing span criterion: pictures

## Theorem

A set of vectors  $\{v_1, v_2, \dots, v_p\}$  is linearly independent if and only if, for every  $j$ , the span of  $v_1, v_2, \dots, v_j$  is strictly larger than the span of  $v_1, v_2, \dots, v_{j-1}$ .



**One vector  $\{v\}$ :**

Linearly independent: span got bigger (than  $\{0\}$ ).

**Two vectors  $\{v, w\}$ :**

Linearly independent: span got bigger.

**Three vectors  $\{v, w, x\}$ :**

Linearly dependent: span didn't get bigger.

# Linear Independence

## Two more facts

**Fact 1:** Say  $v_1, v_2, \dots, v_n$  are in  $\mathbb{R}^m$ . If  $n > m$  then  $\{v_1, v_2, \dots, v_n\}$  is linearly

# Linear Independence

## Two more facts

**Fact 1:** Say  $v_1, v_2, \dots, v_n$  are in  $\mathbb{R}^m$ . If  $n > m$  then  $\{v_1, v_2, \dots, v_n\}$  is linearly dependent: the matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}.$$

cannot have a pivot in each column

# Linear Independence

## Two more facts

**Fact 1:** Say  $v_1, v_2, \dots, v_n$  are in  $\mathbb{R}^m$ . If  $n > m$  then  $\{v_1, v_2, \dots, v_n\}$  is linearly dependent: the matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}.$$

cannot have a pivot in each column (it is too wide).

# Linear Independence

## Two more facts

**Fact 1:** Say  $v_1, v_2, \dots, v_n$  are in  $\mathbb{R}^m$ . If  $n > m$  then  $\{v_1, v_2, \dots, v_n\}$  is linearly dependent: the matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}.$$

cannot have a pivot in each column (it is too wide).

This says you can't have 4 linearly independent vectors in  $\mathbb{R}^3$ , for instance.

# Linear Independence

## Two more facts

**Fact 1:** Say  $v_1, v_2, \dots, v_n$  are in  $\mathbb{R}^m$ . If  $n > m$  then  $\{v_1, v_2, \dots, v_n\}$  is linearly dependent: the matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}.$$

cannot have a pivot in each column (it is too wide).

This says you can't have 4 linearly independent vectors in  $\mathbb{R}^3$ , for instance.

A wide matrix can't have linearly independent columns.

# Linear Independence

## Two more facts

**Fact 1:** Say  $v_1, v_2, \dots, v_n$  are in  $\mathbb{R}^m$ . If  $n > m$  then  $\{v_1, v_2, \dots, v_n\}$  is linearly dependent: the matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}.$$

cannot have a pivot in each column (it is too wide).

This says you can't have 4 linearly independent vectors in  $\mathbb{R}^3$ , for instance.

A wide matrix can't have linearly independent columns.

**Fact 2:** If one of  $v_1, v_2, \dots, v_n$  is zero, then  $\{v_1, v_2, \dots, v_n\}$  is linearly



# Linear Independence

## Two more facts

**Fact 1:** Say  $v_1, v_2, \dots, v_n$  are in  $\mathbb{R}^m$ . If  $n > m$  then  $\{v_1, v_2, \dots, v_n\}$  is linearly dependent: the matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}.$$

cannot have a pivot in each column (it is too wide).

This says you can't have 4 linearly independent vectors in  $\mathbb{R}^3$ , for instance.

A wide matrix can't have linearly independent columns.

**Fact 2:** If one of  $v_1, v_2, \dots, v_n$  is zero, then  $\{v_1, v_2, \dots, v_n\}$  is linearly dependent. For instance, if  $v_1 = 0$ , then

# Linear Independence

## Two more facts

**Fact 1:** Say  $v_1, v_2, \dots, v_n$  are in  $\mathbb{R}^m$ . If  $n > m$  then  $\{v_1, v_2, \dots, v_n\}$  is linearly dependent: the matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}.$$

cannot have a pivot in each column (it is too wide).

This says you can't have 4 linearly independent vectors in  $\mathbb{R}^3$ , for instance.

A wide matrix can't have linearly independent columns.

**Fact 2:** If one of  $v_1, v_2, \dots, v_n$  is zero, then  $\{v_1, v_2, \dots, v_n\}$  is linearly dependent. For instance, if  $v_1 = 0$ , then

$$1 \cdot v_1 + 0 \cdot v_2 + 0 \cdot v_3 + \cdots + 0 \cdot v_n = 0$$

is a linear dependence relation.

# Linear Independence

## Two more facts

**Fact 1:** Say  $v_1, v_2, \dots, v_n$  are in  $\mathbb{R}^m$ . If  $n > m$  then  $\{v_1, v_2, \dots, v_n\}$  is linearly dependent: the matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}.$$

cannot have a pivot in each column (it is too wide).

This says you can't have 4 linearly independent vectors in  $\mathbb{R}^3$ , for instance.

A wide matrix can't have linearly independent columns.

**Fact 2:** If one of  $v_1, v_2, \dots, v_n$  is zero, then  $\{v_1, v_2, \dots, v_n\}$  is linearly dependent. For instance, if  $v_1 = 0$ , then

$$1 \cdot v_1 + 0 \cdot v_2 + 0 \cdot v_3 + \cdots + 0 \cdot v_n = 0$$

is a linear dependence relation.

A set containing the zero vector is linearly dependent.

- ▶ The columns of a matrix  $A$  are linearly independent if the equation  $Ax = 0$  has the trivial solution.
- ▶ If  $S$  is a linearly dependent set, then each vector is a linear combination of the other vectors in  $S$ .
- ▶ The columns of any  $4 \times 5$  matrix are linearly dependent. If  $x$  and  $y$  are linearly independent, and if  $\{x; y; z\}$  is linearly dependent, then  $z$  is in  $\text{Span}\{x; y\}$
- ▶ If  $\{v_1, \dots, v_5\}$  are in  $R^5$  and  $v_3 = 0$ , then  $\{v_1, \dots, v_5\}$  is linearly dependent.
- ▶ Suppose  $A$  is an  $m \times n$  matrix with the property that for all  $b$  in  $R^m$  the equation  $Ax = b$  has at most one solution. Explain why the columns of  $A$  must be linearly independent.

- (a). Find the value(s) of  $h$  for which the vectors are linearly dependent. Is

$$\left\{ \begin{pmatrix} 2 \\ -2 \\ 4 \end{pmatrix}, \begin{pmatrix} 4 \\ -6 \\ 7 \end{pmatrix}, \begin{pmatrix} -2 \\ 2 \\ h \end{pmatrix} \right\} \text{ linearly independent?}$$

- (b). (i) For what values of  $h$  is  $v_3$  in  $\text{Span}\{v_1; v_2\}$ , and (ii) for what values of  $h$  is  $\{v_1; v_2; v_3\}$  linearly dependent?

$$\left\{ \begin{pmatrix} 1 \\ -3 \\ 2 \end{pmatrix}, \begin{pmatrix} -3 \\ 9 \\ -6 \end{pmatrix}, \begin{pmatrix} 5 \\ -7 \\ h \end{pmatrix} \right\}$$

- (c). Determine by inspection whether the vectors are linearly independent

(i).  $\left\{ \begin{pmatrix} 5 \\ 8 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 3 \\ 7 \end{pmatrix} \right\}$

(ii).  $\left\{ \begin{pmatrix} 5 \\ 3 \\ -1 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} -7 \\ 2 \\ 4 \end{pmatrix} \right\}$

(iii).  $\left\{ \begin{pmatrix} 3 \\ 4 \end{pmatrix}, \begin{pmatrix} -1 \\ 5 \end{pmatrix}, \begin{pmatrix} 3 \\ 5 \end{pmatrix}, \begin{pmatrix} 7 \\ 1 \end{pmatrix} \right\}$

### Test your understanding:

- (i) The columns of a matrix  $A$  are linearly independent if the equation  $Ax = 0$  has the trivial solution.
- (ii) If  $S$  is a linearly dependent set, then each vector is a linear combination of the other vectors in  $S$ .
- (iii) The columns of any  $4 \times 5$  matrix are linearly dependent. If  $x$  and  $y$  are linearly independent, and if  $\{x; y; z\}$  is linearly dependent, then  $z$  is in  $\text{Span}\{x; y\}$ .
- (iv) How many pivot columns must a  $6 \times 4$  matrix have if its columns are linearly independent? Why?
- (v) How many pivot columns must a  $4 \times 6$  matrix have if its columns span  $\mathbb{R}^4$ ? Why?
- (vi) If  $\{v_1, \dots, v_5\}$  are in  $\mathbb{R}^5$  and  $v_3 = 0$ , then  $\{v_1, \dots, v_5\}$  is linearly dependent.
- (vii) Suppose  $A$  is an  $m \times n$  matrix with the property that for all  $b$  in  $\mathbb{R}^m$  the equation  $Ax = b$  has at most one solution. Explain why the columns of  $A$  must be linearly independent.

# Section: 2.8

## Subspaces

## Motivation

Now, we will discuss **subspaces** of  $\mathbb{R}^n$ .

Dr Ali and Dr Sara



## Motivation

Now, we will discuss **subspaces** of  $\mathbb{R}^n$ .


A subspace turns out to be the same as a span, except we don't know *which* vectors it's the span of.

## Motivation

Now, we will discuss **subspaces** of  $\mathbb{R}^n$ .

A subspace turns out to be the same as a span, except we don't know *which* vectors it's the span of.

This arises naturally when you have, say, a plane through the origin in  $\mathbb{R}^3$  which is *not* defined (a priori) as a span, but you still want to say something about it.



# Definition of Subspace

## Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ .

# Definition of Subspace

## Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ .

“not empty”

# Definition of Subspace

## Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ .
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ .

“not empty”

# Definition of Subspace

## Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ .
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ .

“not empty”

“closed under addition”

# Definition of Subspace

## Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ .
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ .
3. If  $u$  is in  $V$  and  $c$  is in  $\mathbb{R}$ , then  $cu$  is in  $V$ .

"not empty"

"closed under addition"

# Definition of Subspace

## Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ . "not empty"
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ . "closed under addition"
3. If  $u$  is in  $V$  and  $c$  is in  $\mathbb{R}$ , then  $cu$  is in  $V$ . "closed under  $\times$  scalars"



# Definition of Subspace

## Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ . "not empty"
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ . "closed under addition"
3. If  $u$  is in  $V$  and  $c$  is in  $\mathbb{R}$ , then  $cu$  is in  $V$ . "closed under  $\times$  scalars"

What does this mean?

- ▶ If  $v$  is in  $V$ , then all scalar multiples of  $v$  are in  $V$  by (3).

# Definition of Subspace

## Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ . "not empty"
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ . "closed under addition"
3. If  $u$  is in  $V$  and  $c$  is in  $\mathbb{R}$ , then  $cu$  is in  $V$ . "closed under  $\times$  scalars"

## What does this mean?

- ▶ If  $v$  is in  $V$ , then all scalar multiples of  $v$  are in  $V$  by (3). That is, the line through  $v$  is in  $V$ .

# Definition of Subspace

## Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ . "not empty"
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ . "closed under addition"
3. If  $u$  is in  $V$  and  $c$  is in  $\mathbb{R}$ , then  $cu$  is in  $V$ . "closed under  $\times$  scalars"

## What does this mean?

- ▶ If  $v$  is in  $V$ , then all scalar multiples of  $v$  are in  $V$  by (3). That is, the line through  $v$  is in  $V$ .
- ▶ If  $u, v$  are in  $V$ , then  $xu$  and  $yv$  are in  $V$  for scalars  $x, y$  by (3).

# Definition of Subspace

## Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ . "not empty"
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ . "closed under addition"
3. If  $u$  is in  $V$  and  $c$  is in  $\mathbb{R}$ , then  $cu$  is in  $V$ . "closed under  $\times$  scalars"

## What does this mean?

- ▶ If  $v$  is in  $V$ , then all scalar multiples of  $v$  are in  $V$  by (3). That is, the line through  $v$  is in  $V$ .
- ▶ If  $u, v$  are in  $V$ , then  $xu$  and  $yv$  are in  $V$  for scalars  $x, y$  by (3). So  $xu + yv$  is in  $V$  by (2).

# Definition of Subspace

## Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ . "not empty"
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ . "closed under addition"
3. If  $u$  is in  $V$  and  $c$  is in  $\mathbb{R}$ , then  $cu$  is in  $V$ . "closed under  $\times$  scalars"

## What does this mean?

- ▶ If  $v$  is in  $V$ , then all scalar multiples of  $v$  are in  $V$  by (3). That is, the line through  $v$  is in  $V$ .
- ▶ If  $u, v$  are in  $V$ , then  $xu$  and  $yv$  are in  $V$  for scalars  $x, y$  by (3). So  $xu + yv$  is in  $V$  by (2). So  $\text{Span}\{u, v\}$  is contained in  $V$ .

# Definition of Subspace

## Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ . "not empty"
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ . "closed under addition"
3. If  $u$  is in  $V$  and  $c$  is in  $\mathbb{R}$ , then  $cu$  is in  $V$ . "closed under  $\times$  scalars"

## What does this mean?

- ▶ If  $v$  is in  $V$ , then all scalar multiples of  $v$  are in  $V$  by (3). That is, the line through  $v$  is in  $V$ .
- ▶ If  $u, v$  are in  $V$ , then  $xu$  and  $yv$  are in  $V$  for scalars  $x, y$  by (3). So  $xu + yv$  is in  $V$  by (2). So  $\text{Span}\{u, v\}$  is contained in  $V$ .
- ▶ Likewise, if  $v_1, v_2, \dots, v_n$  are all in  $V$ , then  $\text{Span}\{v_1, v_2, \dots, v_n\}$  is contained in  $V$ .

## Definition of Subspace

### Definition

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  satisfying:

1. The zero vector is in  $V$ . "not empty"
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ . "closed under addition"
3. If  $u$  is in  $V$  and  $c$  is in  $\mathbb{R}$ , then  $cu$  is in  $V$ . "closed under  $\times$  scalars"

### What does this mean?


- ▶ If  $v$  is in  $V$ , then all scalar multiples of  $v$  are in  $V$  by (3). That is, the line through  $v$  is in  $V$ .
- ▶ If  $u, v$  are in  $V$ , then  $xu$  and  $yv$  are in  $V$  for scalars  $x, y$  by (3). So  $xu + yv$  is in  $V$  by (2). So  $\text{Span}\{u, v\}$  is contained in  $V$ .
- ▶ Likewise, if  $v_1, v_2, \dots, v_n$  are all in  $V$ , then  $\text{Span}\{v_1, v_2, \dots, v_n\}$  is contained in  $V$ .

A subspace  $V$  contains the span of any set of vectors in  $V$ .

# Examples

## Example

A line  $L$  through the origin: this contains the span of any vector in  $L$ .






# Examples

## Example

A line  $L$  through the origin: this contains the span of any vector in  $L$ .

## Example


A plane  $P$  through the origin: this contains the span of any vectors in  $P$ .



## Examples


### Example

A line  $L$  through the origin: this contains the span of any vector in  $L$ .



### Example

A plane  $P$  through the origin: this contains the span of any vectors in  $P$ .




### Example

All of  $\mathbb{R}^n$ : this contains 0, and is closed under addition and scalar multiplication.

## Examples


### Example

A line  $L$  through the origin: this contains the span of any vector in  $L$ .



### Example

A plane  $P$  through the origin: this contains the span of any vectors in  $P$ .



### Example

All of  $\mathbb{R}^n$ : this contains 0, and is closed under addition and scalar multiplication.


### Example

The subset  $\{0\}$ : this subspace contains only one vector.

## Examples


### Example

A line  $L$  through the origin: this contains the span of any vector in  $L$ .



### Example

A plane  $P$  through the origin: this contains the span of any vectors in  $P$ .



### Example

All of  $\mathbb{R}^n$ : this contains 0, and is closed under addition and scalar multiplication.

### Example

The subset  $\{0\}$ : this subspace contains only one vector.


Note these are all pictures of spans! (Line, plane, space, etc.)

# Non-Examples

## Non-Example

A line  $L$  (or any other set) that doesn't contain the origin is not a subspace.

Fails:




# Non-Examples

## Non-Example

A line  $L$  (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.



# Non-Examples


## Non-Example

A line  $L$  (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.

## Non-Example

A circle  $C$  is not a subspace. Fails:



# Non-Examples

## Non-Example


A line  $L$  (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.

## Non-Example

A circle  $C$  is not a subspace. Fails:

1,2,3.





# Non-Examples


## Non-Example

A line  $L$  (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.

## Non-Example

A circle  $C$  is not a subspace. Fails: 1,2,3. Think: a circle isn't a "linear space."




# Non-Examples

## Non-Example


A line  $L$  (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.




## Non-Example

A circle  $C$  is not a subspace. Fails: 1,2,3. Think: a circle isn't a "linear space."



## Non-Example

The first quadrant in  $\mathbb{R}^2$  is not a subspace. Fails:




# Non-Examples

## Non-Example


A line  $L$  (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.




## Non-Example

A circle  $C$  is not a subspace. Fails: 1,2,3. Think: a circle isn't a "linear space."



## Non-Example

The first quadrant in  $\mathbb{R}^2$  is not a subspace. Fails: 3 only.




# Non-Examples

## Non-Example


A line  $L$  (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.




## Non-Example

A circle  $C$  is not a subspace. Fails: 1,2,3. Think: a circle isn't a "linear space."




## Non-Example

The first quadrant in  $\mathbb{R}^2$  is not a subspace. Fails: 3 only.



## Non-Example

A line union a plane in  $\mathbb{R}^3$  is not a subspace. Fails:




# Non-Examples

## Non-Example


A line  $L$  (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.




## Non-Example

A circle  $C$  is not a subspace. Fails: 1,2,3. Think: a circle isn't a "linear space."




## Non-Example

The first quadrant in  $\mathbb{R}^2$  is not a subspace. Fails: 3 only.



## Non-Example

A line union a plane in  $\mathbb{R}^3$  is not a subspace. Fails: 2 only.



# Subsets and Subspaces

They aren't the same thing

A **subset** of  $\mathbb{R}^n$  is any collection of vectors whatsoever.

# Subsets and Subspaces

They aren't the same thing

A **subset** of  $\mathbb{R}^n$  is any collection of vectors whatsoever.

All of the non-examples are still subsets.

# Subsets and Subspaces

They aren't the same thing

A **subset** of  $\mathbb{R}^n$  is any collection of vectors whatsoever.

All of the non-examples are still subsets.

A **subspace** is a special kind of subset, which satisfies the three defining properties.




# Subsets and Subspaces

They aren't the same thing

A **subset** of  $\mathbb{R}^n$  is any collection of vectors whatsoever.

All of the non-examples are still subsets.

A **subspace** is a special kind of subset, which satisfies the three defining properties.



Subset: *yes*

Subspace: *no*

# Spans are Subspaces

## Theorem

Any  $\text{Span}\{v_1, v_2, \dots, v_n\}$  is a subspace.

Dr Ali and Dr Sara

# Spans are Subspaces

## Theorem

Any  $\text{Span}\{v_1, v_2, \dots, v_n\}$  is a subspace.

!!!

Every subspace is a span, and every span is a subspace.

# Spans are Subspaces

## Theorem

Any  $\text{Span}\{v_1, v_2, \dots, v_n\}$  is a subspace.

!!!

Every subspace is a span, and every span is a subspace.

## Definition

If  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , we say that  $V$  is the subspace **generated by** or **spanned by** the vectors  $v_1, v_2, \dots, v_n$ .

## Check:

1.  $0 = 0v_1 + 0v_2 + \dots + 0v_n$  is in the span.
2. If, say,  $u = 3v_1 + 4v_2$  and  $v = -v_1 - 2v_2$ , then

$$u + v = 3v_1 + 4v_2 - v_1 - 2v_2 = 2v_1 + 2v_2$$

is also in the span.

3. Similarly, if  $u$  is in the span, then so is  $cu$  for any scalar  $c$ .

Poll

Is the empty set  $\{\}$  a subspace? If not, which property(ies) does it fail?

Poll

Is the empty set  $\{\}$  a subspace? If not, which property(ies) does it fail?

The zero vector is not contained in the empty set, so it is *not* a subspace.

Poll

Is the empty set  $\{\}$  a subspace? If not, which property(ies) does it fail?


The zero vector is not contained in the empty set, so it is *not* a subspace.

**Question:** What is the difference between  $\{\}$  and  $\{0\}$ ?

# Subspaces

## Verification

Let  $V = \left\{ \begin{pmatrix} a \\ b \end{pmatrix} \text{ in } \mathbb{R}^2 \mid ab = 0 \right\}$ . Let's check if  $V$  is a subspace or not.

1. Does  $V$  contain the zero vector?  $\begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \implies ab = 0$  


3. Is  $V$  closed under scalar multiplication?

▶ Let  $\begin{pmatrix} a \\ b \end{pmatrix}$  be in  $V$ .

▶ *This means:*  $a$  and  $b$  are numbers such that  $ab = 0$ .

▶ Let  $c$  be a scalar. Is  $c \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} ca \\ cb \end{pmatrix}$  in  $V$ ?

▶ *This means:*  $(ca)(cb) = 0$ .

▶ Well,  $(ca)(cb) = c^2(ab) = c^2(0) = 0$  


2. Is  $V$  closed under addition?

▶ Let  $\begin{pmatrix} a \\ b \end{pmatrix}$  and  $\begin{pmatrix} a' \\ b' \end{pmatrix}$  be in  $V$ .

▶ *This means:*  $ab = 0$  and  $a'b' = 0$ .

▶ Is  $\begin{pmatrix} a \\ b \end{pmatrix} + \begin{pmatrix} a' \\ b' \end{pmatrix} = \begin{pmatrix} a+a' \\ b+b' \end{pmatrix}$  in  $V$ ?

▶ *This means:*  $(a+a')(b+b') = 0$ .


▶ This is not true for all such  $a, a', b, b'$ : for instance,  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$  are in  $V$ , but their sum  $\begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$  is not in  $V$ , because  $1 \cdot 1 \neq 0$ . 



# Subspaces

## Verification

Let  $V = \left\{ \begin{pmatrix} a \\ b \end{pmatrix} \text{ in } \mathbb{R}^2 \mid ab = 0 \right\}$ . Let's check if  $V$  is a subspace or not.

1. Does  $V$  contain the zero vector?  $\begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \implies ab = 0$  


3. Is  $V$  closed under scalar multiplication?

▶ Let  $\begin{pmatrix} a \\ b \end{pmatrix}$  be in  $V$ .

▶ *This means:*  $a$  and  $b$  are numbers such that  $ab = 0$ .

▶ Let  $c$  be a scalar. Is  $c \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} ca \\ cb \end{pmatrix}$  in  $V$ ?

▶ *This means:*  $(ca)(cb) = 0$ .

▶ Well,  $(ca)(cb) = c^2(ab) = c^2(0) = 0$  


2. Is  $V$  closed under addition?

▶ Let  $\begin{pmatrix} a \\ b \end{pmatrix}$  and  $\begin{pmatrix} a' \\ b' \end{pmatrix}$  be in  $V$ .

▶ *This means:*  $ab = 0$  and  $a'b' = 0$ .

▶ Is  $\begin{pmatrix} a \\ b \end{pmatrix} + \begin{pmatrix} a' \\ b' \end{pmatrix} = \begin{pmatrix} a+a' \\ b+b' \end{pmatrix}$  in  $V$ ?

▶ *This means:*  $(a+a')(b+b') = 0$ .

▶ This is not true for all such  $a, a', b, b'$ : for instance,  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$  are in  $V$ , but their sum  $\begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$  is not in  $V$ , because  $1 \cdot 1 \neq 0$ . 

We conclude that  $V$  is *not* a subspace.

# Subspaces

## Verification

Let  $V = \left\{ \begin{pmatrix} a \\ b \end{pmatrix} \text{ in } \mathbb{R}^2 \mid ab = 0 \right\}$ . Let's check if  $V$  is a subspace or not.

1. Does  $V$  contain the zero vector?  $\begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \implies ab = 0$  ✓

3. Is  $V$  closed under scalar multiplication?

▶ Let  $\begin{pmatrix} a \\ b \end{pmatrix}$  be in  $V$ .

▶ *This means:*  $a$  and  $b$  are numbers such that  $ab = 0$ .

▶ Let  $c$  be a scalar. Is  $c \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} ca \\ cb \end{pmatrix}$  in  $V$ ?

▶ *This means:*  $(ca)(cb) = 0$ .

▶ Well,  $(ca)(cb) = c^2(ab) = c^2(0) = 0$  ✓

2. Is  $V$  closed under addition?


▶ Let  $\begin{pmatrix} a \\ b \end{pmatrix}$  and  $\begin{pmatrix} a' \\ b' \end{pmatrix}$  be in  $V$ .

▶ *This means:*  $ab = 0$  and  $a'b' = 0$ .

▶ Is  $\begin{pmatrix} a \\ b \end{pmatrix} + \begin{pmatrix} a' \\ b' \end{pmatrix} = \begin{pmatrix} a+a' \\ b+b' \end{pmatrix}$  in  $V$ ?

▶ *This means:*  $(a+a')(b+b') = 0$ .

▶ This is not true for all such  $a, a', b, b'$ : for instance,  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$  are in  $V$ , but their sum  $\begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$  is not in  $V$ , because  $1 \cdot 1 \neq 0$ . ✗



We conclude that  $V$  is *not* a subspace. A picture is above. (It doesn't look like a span.)

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to two subspaces.

Dr Ali and Dr Sara

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to two subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$ —spanned by the columns of  $A$ . It is written  $\text{Col } A$ .

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to two subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . It is written  $\text{Col } A$ .

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to two subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . It is written  $\text{Col } A$ .
- ▶ The **null space** of  $A$  is the set of all solutions of the homogeneous equation  $Ax = 0$ :

$$\text{Nul } A = \{x \text{ in } \mathbb{R}^n \mid Ax = 0\}.$$

This is a subspace of  $\mathbb{R}^n$ .

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to two subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . It is written  $\text{Col } A$ .
- ▶ The **null space** of  $A$  is the set of all solutions of the homogeneous equation  $Ax = 0$ :

$$\text{Nul } A = \{x \text{ in } \mathbb{R}^n \mid Ax = 0\}.$$

This is a subspace of  $\mathbb{R}^n$ .

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to *two* subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . It is written  $\text{Col } A$ .
- ▶ The **null space** of  $A$  is the set of all solutions of the homogeneous equation  $Ax = 0$ :

$$\text{Nul } A = \{x \text{ in } \mathbb{R}^n \mid Ax = 0\}.$$

This is a subspace of  $\mathbb{R}^n$ .

The column space is defined as a span, so we know it is a subspace.



## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to *two* subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . It is written  $\text{Col } A$ .
- ▶ The **null space** of  $A$  is the set of all solutions of the homogeneous equation  $Ax = 0$ :

$$\text{Nul } A = \{x \text{ in } \mathbb{R}^n \mid Ax = 0\}.$$

This is a subspace of  $\mathbb{R}^n$ .

The column space is defined as a span, so we know it is a subspace. It is the range (as opposed to the codomain) of the transformation  $T(x) = Ax$ .

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to *two* subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . It is written  $\text{Col } A$ .
- ▶ The **null space** of  $A$  is the set of all solutions of the homogeneous equation  $Ax = 0$ :

$$\text{Nul } A = \{x \text{ in } \mathbb{R}^n \mid Ax = 0\}.$$

This is a subspace of  $\mathbb{R}^n$ .

The column space is defined as a span, so we know it is a subspace. It is the range (as opposed to the codomain) of the transformation  $T(x) = Ax$ .

**Check** that the null space is a subspace:

1.  $0$  is in  $\text{Nul } A$  because  $A0 = 0$ .
2. If  $u$  and  $v$  are in  $\text{Nul } A$ , then  $Au = 0$  and  $Av = 0$ . Hence
$$A(u + v) = Au + Av = 0,$$
so  $u + v$  is in  $\text{Nul } A$ .
3. If  $u$  is in  $\text{Nul } A$ , then  $Au = 0$ . For any scalar  $c$ ,  $A(cu) = cAu = 0$ . So  $cu$  is in  $\text{Nul } A$ .

## Column Space and Null Space

### Example

$$\text{Let } A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Let's compute the column space:

$$\text{Col } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\} = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}.$$

This is a line in  $\mathbb{R}^3$ .

# Column Space and Null Space


## Example

$$\text{Let } A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Let's compute the column space:

$$\text{Col } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\} = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}.$$

This is a line in  $\mathbb{R}^3$ .




# Column Space and Null Space

## Example

$$\text{Let } A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Let's compute the column space:

$$\text{Col } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\} = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}.$$



This is a line in  $\mathbb{R}^3$ .

Let's compute the null space:

$$A \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x+y \\ x+y \\ x+y \end{pmatrix}.$$

This zero if and only if  $x = -y$ . So

$$\text{Nul } A = \left\{ \begin{pmatrix} x \\ y \end{pmatrix} \text{ in } \mathbb{R}^2 \mid y = -x \right\}.$$

This defines a line in  $\mathbb{R}^2$ :


# Column Space and Null Space

## Example

$$\text{Let } A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Let's compute the column space:

$$\text{Col } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\} = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}.$$



This is a line in  $\mathbb{R}^3$ .


Let's compute the null space:

$$A \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x+y \\ x+y \\ x+y \end{pmatrix}.$$

This is zero if and only if  $x = -y$ . So

$$\text{Nul } A = \left\{ \begin{pmatrix} x \\ y \end{pmatrix} \text{ in } \mathbb{R}^2 \mid y = -x \right\}.$$

This defines a line in  $\mathbb{R}^2$ :



## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

Dr Ali and Dr Sara

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ .



## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

### Question

How to find vectors which span the null space?

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

### Question

How to find vectors which span the null space?

**Answer:** Parametric vector form!

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

### Question

How to find vectors which span the null space?

**Answer:** Parametric vector form! We know that the solution set to  $Ax = 0$  has a parametric form that looks like

$$x_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \quad \text{if, say, } x_3 \text{ and } x_4 \\ \text{are the free} \\ \text{variables.}$$

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

### Question

How to find vectors which span the null space?

**Answer:** Parametric vector form! We know that the solution set to  $Ax = 0$  has a parametric form that looks like

$$x_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \quad \text{if, say, } x_3 \text{ and } x_4 \text{ are the free variables. So} \quad \text{Nul } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

### Question

How to find vectors which span the null space?

**Answer:** Parametric vector form! We know that the solution set to  $Ax = 0$  has a parametric form that looks like

$$x_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \quad \begin{array}{l} \text{if, say, } x_3 \text{ and } x_4 \\ \text{are the free} \\ \text{variables. So} \end{array} \quad \text{Nul } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

Refer back to the slides for §1.5 (Solution Sets).

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

### Question

How to find vectors which span the null space?

**Answer:** Parametric vector form! We know that the solution set to  $Ax = 0$  has a parametric form that looks like

$$x_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \quad \begin{array}{l} \text{if, say, } x_3 \text{ and } x_4 \\ \text{are the free} \\ \text{variables. So} \end{array} \quad \text{Nul } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

Refer back to the slides for §1.5 (Solution Sets).

**Note:** It is much easier to define the null space first as a subspace, then find spanning vectors *later*, if we need them. This is one reason subspaces are so useful.

# The Null Space is a Span

Example, revisited

Find vector(s) that span the null space of  $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}$ .

The reduced row echelon form is  $\begin{pmatrix} 1 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}$ .

This gives the equation  $x + y = 0$ , or

$$\begin{array}{l} x = -y \\ y = y \end{array} \xrightarrow{\text{parametric vector form}} \begin{pmatrix} x \\ y \end{pmatrix} = y \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

The null space is

$$\text{Nul } A = \text{Span} \left\{ \begin{pmatrix} -1 \\ 1 \end{pmatrix} \right\}.$$



# The Null Space is a Span

Example, revisited

Find vector(s) that span the null space of  $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}$ .


The reduced row echelon form is  $\begin{pmatrix} 1 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}$ .

This gives the equation  $x + y = 0$ , or

$$\begin{array}{l} x = -y \\ y = y \end{array} \xrightarrow{\text{parametric vector form}} \begin{pmatrix} x \\ y \end{pmatrix} = y \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

The null space is

$$\text{Nul } A = \text{Span} \left\{ \begin{pmatrix} -1 \\ 1 \end{pmatrix} \right\}.$$



# Subspaces

## Summary

How do you check if a subset is a subspace?

- Is it a span? Can it be written as a span?

# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?

# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?

# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of  $\mathbb{R}^n$  or the zero subspace  $\{0\}$ ?

# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of  $\mathbb{R}^n$  or the zero subspace  $\{0\}$ ?
- ▶ Can it be written as a type of subspace that we'll learn about later (eigenspaces, ...)?

# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of  $\mathbb{R}^n$  or the zero subspace  $\{0\}$ ?
- ▶ Can it be written as a type of subspace that we'll learn about later (eigenspaces, ...)?

If so, then it's automatically a subspace.

# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of  $\mathbb{R}^n$  or the zero subspace  $\{0\}$ ?
- ▶ Can it be written as a type of subspace that we'll learn about later (eigenspaces, ...)?

If so, then it's automatically a subspace.

If all else fails:



# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of  $\mathbb{R}^n$  or the zero subspace  $\{0\}$ ?
- ▶ Can it be written as a type of subspace that we'll learn about later (eigenspaces, ...)?

If so, then it's automatically a subspace.

If all else fails:

- ▶ Can you verify directly that it satisfies the three defining properties?

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

Dr Ali and Dr Sara

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

The number of vectors in a basis is the **dimension** of  $V$ , and is written  $\dim V$ .

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

The number of vectors in a basis is the **dimension** of  $V$ , and is written  $\dim V$ .

Note the big  
red border here

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

The number of vectors in a basis is the **dimension** of  $V$ , and is written  $\dim V$ .

Note the big  
red border here

**Why** is a basis the smallest number of vectors needed to span?



## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

The number of vectors in a basis is the **dimension** of  $V$ , and is written  $\dim V$ .

Note the big  
red border here

**Why** is a basis the smallest number of vectors needed to span?

Recall: *linearly independent* means that every time you add another vector, the span gets bigger.

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

The number of vectors in a basis is the **dimension** of  $V$ , and is written  $\dim V$ .

Note the big  
red border here

**Why** is a basis the smallest number of vectors needed to span?

Recall: *linearly independent* means that every time you add another vector, the span gets bigger.

Hence, if we remove any vector, the span gets *smaller*: so any smaller set can't span  $V$ .

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

The number of vectors in a basis is the **dimension** of  $V$ , and is written  $\dim V$ .

Note the big  
red border here

**Why** is a basis the smallest number of vectors needed to span?

Recall: *linearly independent* means that every time you add another vector, the span gets bigger.

Hence, if we remove any vector, the span gets *smaller*: so any smaller set can't span  $V$ .

### Important

A subspace has *many different* bases, but they all have the same number of vectors (see the exercises in §2.9).

## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

Dr Ali and Dr Sara

# Bases of $\mathbb{R}^2$

## Question

What is a basis for  $\mathbb{R}^2$ ?


We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.

## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.




# Bases of $\mathbb{R}^2$

## Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} =$




## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .






## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .
2. They are linearly independent because they are not collinear.




## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .
2. They are linearly independent because they are not collinear.



### Question

What is another basis for  $\mathbb{R}^2$ ?


## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .
2. They are linearly independent because they are not collinear.



### Question

What is another basis for  $\mathbb{R}^2$ ?

Any two nonzero vectors that are not collinear.


## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.


1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .
2. They are linearly independent because they are not collinear.



### Question

What is another basis for  $\mathbb{R}^2$ ?

Any two nonzero vectors that are not collinear.  $\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$  is also a basis.




## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .
2. They are linearly independent because they are not collinear.




### Question

What is another basis for  $\mathbb{R}^2$ ?

Any two nonzero vectors that are not collinear.  $\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$  is also a basis.

1. They span:  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  has a pivot in every row.




## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .
2. They are linearly independent because they are not collinear.




### Question

What is another basis for  $\mathbb{R}^2$ ?

Any two nonzero vectors that are not collinear.  $\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$  is also a basis.

1. They span:  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  has a pivot in every row.
2. They are linearly independent:  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  has a pivot in every column.



## Bases of $\mathbb{R}^n$

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for  $\mathbb{R}^n$ .

## Bases of $\mathbb{R}^n$

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for  $\mathbb{R}^n$ .

1. They span:  $I_n$  has a pivot in every row.



## Bases of $\mathbb{R}^n$

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for  $\mathbb{R}^n$ . The identity matrix has columns  $e_1, e_2, \dots, e_n$ .

1. They span:  $I_n$  has a pivot in every row.

## Bases of $\mathbb{R}^n$

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for  $\mathbb{R}^n$ .  The identity matrix has columns  $e_1, e_2, \dots, e_n$ .

1. They span:  $I_n$  has a pivot in every row.
2. They are linearly independent:  $I_n$  has a pivot in every column.

## Bases of $\mathbb{R}^n$

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for  $\mathbb{R}^n$ .  The identity matrix has columns  $e_1, e_2, \dots, e_n$ .

1. They span:  $I_n$  has a pivot in every row.
2. They are linearly independent:  $I_n$  has a pivot in every column.

**In general:**  $\{v_1, v_2, \dots, v_n\}$  is a basis for  $\mathbb{R}^n$  if and only if the matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}$$

has a pivot in every row and every column, i.e. if  $A$  is \_\_\_\_\_.

## Bases of $\mathbb{R}^n$

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for  $\mathbb{R}^n$ .  The identity matrix has columns  $e_1, e_2, \dots, e_n$ .

1. They span:  $I_n$  has a pivot in every row.
2. They are linearly independent:  $I_n$  has a pivot in every column.

**In general:**  $\{v_1, v_2, \dots, v_n\}$  is a basis for  $\mathbb{R}^n$  if and only if the matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}$$

has a pivot in every row and every column, i.e. if  $A$  is *invertible*.

## Basis of a Subspace

### Example

#### Example

Let

$$V = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \text{ in } \mathbb{R}^3 \mid x + 3y + z = 0 \right\} \quad \mathcal{B} = \left\{ \begin{pmatrix} -3 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ -3 \end{pmatrix} \right\}.$$

Verify that  $\mathcal{B}$  is a basis for  $V$ .

0. In  $V$ : both vectors are in  $V$  because

$$-3 + 3(1) + 0 = 0 \quad \text{and} \quad 0 + 3(1) + (-3) = 0.$$

1. Span: If  $\begin{pmatrix} x \\ y \\ z \end{pmatrix}$  is in  $V$ , then  $y = -\frac{1}{3}(x + z)$ , so

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = -\frac{x}{3} \begin{pmatrix} -3 \\ 1 \\ 0 \end{pmatrix} - \frac{z}{3} \begin{pmatrix} 0 \\ 1 \\ -3 \end{pmatrix}.$$

2. Linearly independent:

$$c_1 \begin{pmatrix} -3 \\ 1 \\ 0 \end{pmatrix} + c_2 \begin{pmatrix} 0 \\ 1 \\ -3 \end{pmatrix} = 0 \implies \begin{pmatrix} -3c_1 \\ c_1 + c_2 \\ -3c_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \implies c_1 = c_2 = 0.$$

## Basis for $\text{Nul } A$

### Fact

The vectors in the parametric vector form of the general solution to  $Ax = 0$  always form a basis for  $\text{Nul } A$ .

## Basis for Nul $A$


### Fact

The vectors in the parametric vector form of the general solution to  $Ax = 0$  always form a basis for  $\text{Nul } A$ .

### Example

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

parametric vector form  $\xrightarrow{\text{~~~~~}}$

$$x = x_3 \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \xrightarrow{\text{basis of Nul } A} \left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \right\}$$


1. The vectors span  $\text{Nul } A$  by construction (every solution to  $Ax = 0$  has this form).
2. Can you see why they are linearly independent? (Look at the last two rows.)

## Basis for Col $A$

Fact

The *pivot columns* of  $A$  always form a basis for Col  $A$ .



## Basis for Col $A$

### Fact

The *pivot columns* of  $A$  always form a basis for Col  $A$ .

**Warning:** I mean the pivot columns of the *original* matrix  $A$ , not the row-reduced form.

## Basis for Col $A$

### Fact

The *pivot columns* of  $A$  always form a basis for Col  $A$ .

**Warning:** I mean the pivot columns of the *original* matrix  $A$ , not the row-reduced form. (Row reduction changes the column space.)

## Basis for Col A

### Fact

The *pivot columns* of  $A$  always form a basis for Col  $A$ .

**Warning:** I mean the pivot columns of the *original* matrix  $A$ , not the row-reduced form. (Row reduction changes the column space.)

### Example

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

pivot columns = basis  $\longleftrightarrow$  pivot columns in rref

So a basis for Col  $A$  is

$$\left\{ \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ -3 \\ 4 \end{pmatrix} \right\}.$$

## Basis for Col A

### Fact

The *pivot columns* of  $A$  always form a basis for Col  $A$ .

**Warning:** I mean the pivot columns of the *original* matrix  $A$ , not the row-reduced form. (Row reduction changes the column space.)

### Example

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

pivot columns = basis  $\longleftrightarrow$  pivot columns in rref

So a basis for Col  $A$  is

$$\left\{ \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ -3 \\ 4 \end{pmatrix} \right\}.$$

**Why?**

# Subspaces: Dimension & Ranks

Ex 2.8 & 2.9

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to two subspaces.

Dr Ali and Dr Sara

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to two subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$ —spanned by the columns of  $A$ . It is written  $\text{Col } A$ .

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to two subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . It is written  $\text{Col } A$ .



## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to two subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . It is written  $\text{Col } A$ .
- ▶ The **null space** of  $A$  is the set of all solutions of the homogeneous equation  $Ax = 0$ :

$$\text{Nul } A = \{x \text{ in } \mathbb{R}^n \mid Ax = 0\}.$$

This is a subspace of  $\mathbb{R}^n$ .

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to two subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . It is written  $\text{Col } A$ .
- ▶ The **null space** of  $A$  is the set of all solutions of the homogeneous equation  $Ax = 0$ :

$$\text{Nul } A = \{x \text{ in } \mathbb{R}^n \mid Ax = 0\}.$$

This is a subspace of  $\mathbb{R}^n$ .

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to *two* subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . It is written  $\text{Col } A$ .
- ▶ The **null space** of  $A$  is the set of all solutions of the homogeneous equation  $Ax = 0$ :

$$\text{Nul } A = \{x \text{ in } \mathbb{R}^n \mid Ax = 0\}.$$

This is a subspace of  $\mathbb{R}^n$ .

The column space is defined as a span, so we know it is a subspace.

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to *two* subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . It is written  $\text{Col } A$ .
- ▶ The **null space** of  $A$  is the set of all solutions of the homogeneous equation  $Ax = 0$ :

$$\text{Nul } A = \{x \text{ in } \mathbb{R}^n \mid Ax = 0\}.$$

This is a subspace of  $\mathbb{R}^n$ .

The column space is defined as a span, so we know it is a subspace. It is the range (as opposed to the codomain) of the transformation  $T(x) = Ax$ .

## Column Space and Null Space

An  $m \times n$  matrix  $A$  naturally gives rise to *two* subspaces.

### Definition

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . It is written  $\text{Col } A$ .
- ▶ The **null space** of  $A$  is the set of all solutions of the homogeneous equation  $Ax = 0$ :

$$\text{Nul } A = \{x \text{ in } \mathbb{R}^n \mid Ax = 0\}.$$

This is a subspace of  $\mathbb{R}^n$ .

The column space is defined as a span, so we know it is a subspace. It is the range (as opposed to the codomain) of the transformation  $T(x) = Ax$ .

**Check** that the null space is a subspace:

1.  $0$  is in  $\text{Nul } A$  because  $A0 = 0$ .
2. If  $u$  and  $v$  are in  $\text{Nul } A$ , then  $Au = 0$  and  $Av = 0$ . Hence
$$A(u + v) = Au + Av = 0,$$
so  $u + v$  is in  $\text{Nul } A$ .
3. If  $u$  is in  $\text{Nul } A$ , then  $Au = 0$ . For any scalar  $c$ ,  $A(cu) = cAu = 0$ . So  $cu$  is in  $\text{Nul } A$ .

## Column Space and Null Space

### Example

$$\text{Let } A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Let's compute the column space:

$$\text{Col } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\} = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}.$$

This is a line in  $\mathbb{R}^3$ .

# Column Space and Null Space


## Example

$$\text{Let } A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Let's compute the column space:

$$\text{Col } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\} = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}.$$

This is a line in  $\mathbb{R}^3$ .




# Column Space and Null Space

## Example

$$\text{Let } A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Let's compute the column space:

$$\text{Col } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\} = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}.$$



This is a line in  $\mathbb{R}^3$ .

Let's compute the null space:

$$A \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x+y \\ x+y \\ x+y \end{pmatrix}.$$

This zero if and only if  $x = -y$ . So

$$\text{Nul } A = \left\{ \begin{pmatrix} x \\ y \end{pmatrix} \text{ in } \mathbb{R}^2 \mid y = -x \right\}.$$

This defines a line in  $\mathbb{R}^2$ :




# Column Space and Null Space

## Example

$$\text{Let } A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Let's compute the column space:

$$\text{Col } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\} = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}.$$



This is a line in  $\mathbb{R}^3$ .


Let's compute the null space:

$$A \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x+y \\ x+y \\ x+y \end{pmatrix}.$$

This is zero if and only if  $x = -y$ . So

$$\text{Nul } A = \left\{ \begin{pmatrix} x \\ y \end{pmatrix} \text{ in } \mathbb{R}^2 \mid y = -x \right\}.$$

This defines a line in  $\mathbb{R}^2$ :



## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

Dr Ali and Dr Sara

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ .

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

### Question

How to find vectors which span the null space?

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

### Question

How to find vectors which span the null space?

**Answer:** Parametric vector form!

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

### Question

How to find vectors which span the null space?

**Answer:** Parametric vector form! We know that the solution set to  $Ax = 0$  has a parametric form that looks like

$$x_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \quad \text{if, say, } x_3 \text{ and } x_4 \\ \text{are the free} \\ \text{variables.}$$

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

### Question

How to find vectors which span the null space?

**Answer:** Parametric vector form! We know that the solution set to  $Ax = 0$  has a parametric form that looks like

$$x_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \quad \text{if, say, } x_3 \text{ and } x_4 \text{ are the free variables. So} \quad \text{Nul } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$



## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

### Question

How to find vectors which span the null space?

**Answer:** Parametric vector form! We know that the solution set to  $Ax = 0$  has a parametric form that looks like

$$x_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \quad \begin{array}{l} \text{if, say, } x_3 \text{ and } x_4 \\ \text{are the free} \\ \text{variables. So} \end{array} \quad \text{Nul } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

Refer back to the slides for §1.5 (Solution Sets).

## The Null Space is a Span

The column space of a matrix  $A$  is defined to be a span (of the columns).

The null space is defined to be the solution set to  $Ax = 0$ . It is a subspace, so it is a span.

### Question

How to find vectors which span the null space?

**Answer:** Parametric vector form! We know that the solution set to  $Ax = 0$  has a parametric form that looks like

$$x_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \quad \begin{array}{l} \text{if, say, } x_3 \text{ and } x_4 \\ \text{are the free} \\ \text{variables. So} \end{array} \quad \text{Nul } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

Refer back to the slides for §1.5 (Solution Sets).

**Note:** It is much easier to define the null space first as a subspace, then find spanning vectors *later*, if we need them. This is one reason subspaces are so useful.

# The Null Space is a Span

Example, revisited

Find vector(s) that span the null space of  $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}$ .

The reduced row echelon form is  $\begin{pmatrix} 1 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}$ .

This gives the equation  $x + y = 0$ , or

$$\begin{array}{l} x = -y \\ y = y \end{array} \xrightarrow{\text{parametric vector form}} \begin{pmatrix} x \\ y \end{pmatrix} = y \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

The null space is

$$\text{Nul } A = \text{Span} \left\{ \begin{pmatrix} -1 \\ 1 \end{pmatrix} \right\}.$$

# The Null Space is a Span

Example, revisited

Find vector(s) that span the null space of  $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}$ .


The reduced row echelon form is  $\begin{pmatrix} 1 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}$ .

This gives the equation  $x + y = 0$ , or

$$\begin{array}{l} x = -y \\ y = y \end{array} \xrightarrow{\text{parametric vector form}} \begin{pmatrix} x \\ y \end{pmatrix} = y \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

The null space is

$$\text{Nul } A = \text{Span} \left\{ \begin{pmatrix} -1 \\ 1 \end{pmatrix} \right\}.$$



# Subspaces

## Summary

How do you check if a subset is a subspace?

- Is it a span? Can it be written as a span?

# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?

# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?

# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of  $\mathbb{R}^n$  or the zero subspace  $\{0\}$ ?



# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of  $\mathbb{R}^n$  or the zero subspace  $\{0\}$ ?
- ▶ Can it be written as a type of subspace that we'll learn about later (eigenspaces, ...)?

# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of  $\mathbb{R}^n$  or the zero subspace  $\{0\}$ ?
- ▶ Can it be written as a type of subspace that we'll learn about later (eigenspaces, ...)?

If so, then it's automatically a subspace.

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of  $\mathbb{R}^n$  or the zero subspace  $\{0\}$ ?
- ▶ Can it be written as a type of subspace that we'll learn about later (eigenspaces, ...)?

If so, then it's automatically a subspace.

If all else fails:

# Subspaces

## Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of  $\mathbb{R}^n$  or the zero subspace  $\{0\}$ ?
- ▶ Can it be written as a type of subspace that we'll learn about later (eigenspaces, ...)?

If so, then it's automatically a subspace.

If all else fails:

- ▶ Can you verify directly that it satisfies the three defining properties?

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

Dr Ali and Dr Sara

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.



## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

The number of vectors in a basis is the **dimension** of  $V$ , and is written  $\dim V$ .

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

The number of vectors in a basis is the **dimension** of  $V$ , and is written  $\dim V$ .

Note the big  
red border here

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

The number of vectors in a basis is the **dimension** of  $V$ , and is written  $\dim V$ .

Note the big  
red border here

**Why** is a basis the smallest number of vectors needed to span?

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

The number of vectors in a basis is the **dimension** of  $V$ , and is written  $\dim V$ .

Note the big  
red border here

**Why** is a basis the smallest number of vectors needed to span?

Recall: *linearly independent* means that every time you add another vector, the span gets bigger.

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

The number of vectors in a basis is the **dimension** of  $V$ , and is written  $\dim V$ .

Note the big  
red border here

**Why** is a basis the smallest number of vectors needed to span?

Recall: *linearly independent* means that every time you add another vector, the span gets bigger.

Hence, if we remove any vector, the span gets *smaller*: so any smaller set can't span  $V$ .

## Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

### Definition

Let  $V$  be a subspace of  $\mathbb{R}^n$ . A **basis** of  $V$  is a set of vectors  $\{v_1, v_2, \dots, v_m\}$  in  $V$  such that:

1.  $V = \text{Span}\{v_1, v_2, \dots, v_m\}$ , and
2.  $\{v_1, v_2, \dots, v_m\}$  is linearly independent.

The number of vectors in a basis is the **dimension** of  $V$ , and is written  $\dim V$ .

Note the big  
red border here

**Why** is a basis the smallest number of vectors needed to span?

Recall: *linearly independent* means that every time you add another vector, the span gets bigger.

Hence, if we remove any vector, the span gets *smaller*: so any smaller set can't span  $V$ .

### Important

A subspace has *many different* bases, but they all have the same number of vectors (see the exercises in §2.9).

# Bases of $\mathbb{R}^2$

## Question

What is a basis for  $\mathbb{R}^2$ ?

Dr Ali and Dr Sara

## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.




## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.




# Bases of $\mathbb{R}^2$

## Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} =$




## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .




## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .
2. They are linearly independent because they are not collinear.




## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .
2. They are linearly independent because they are not collinear.



### Question

What is another basis for  $\mathbb{R}^2$ ?


## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .
2. They are linearly independent because they are not collinear.



### Question

What is another basis for  $\mathbb{R}^2$ ?

Any two nonzero vectors that are not collinear.


## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.


1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .
2. They are linearly independent because they are not collinear.



### Question

What is another basis for  $\mathbb{R}^2$ ?

Any two nonzero vectors that are not collinear.  $\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$  is also a basis.




## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .
2. They are linearly independent because they are not collinear.




### Question

What is another basis for  $\mathbb{R}^2$ ?

Any two nonzero vectors that are not collinear.  $\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$  is also a basis.

1. They span:  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  has a pivot in every row.






## Bases of $\mathbb{R}^2$

### Question

What is a basis for  $\mathbb{R}^2$ ?

We need two vectors that *span*  $\mathbb{R}^2$  and are *linearly independent*.  $\{e_1, e_2\}$  is one basis.

1. They span:  $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$ .
2. They are linearly independent because they are not collinear.




### Question

What is another basis for  $\mathbb{R}^2$ ?

Any two nonzero vectors that are not collinear.  $\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$  is also a basis.

1. They span:  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  has a pivot in every row.
2. They are linearly independent:  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  has a pivot in every column.



## Bases of $\mathbb{R}^n$

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for  $\mathbb{R}^n$ .

## Bases of $\mathbb{R}^n$

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for  $\mathbb{R}^n$ .

1. They span:  $I_n$  has a pivot in every row.

## Bases of $\mathbb{R}^n$

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for  $\mathbb{R}^n$ .  The identity matrix has columns  $e_1, e_2, \dots, e_n$ .

1. They span:  $I_n$  has a pivot in every row.

## Bases of $\mathbb{R}^n$

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for  $\mathbb{R}^n$ .  The identity matrix has columns  $e_1, e_2, \dots, e_n$ .

1. They span:  $I_n$  has a pivot in every row.
2. They are linearly independent:  $I_n$  has a pivot in every column.

## Bases of $\mathbb{R}^n$

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for  $\mathbb{R}^n$ .  The identity matrix has columns  $e_1, e_2, \dots, e_n$ .

1. They span:  $I_n$  has a pivot in every row.
2. They are linearly independent:  $I_n$  has a pivot in every column.

**In general:**  $\{v_1, v_2, \dots, v_n\}$  is a basis for  $\mathbb{R}^n$  if and only if the matrix

$$A = \left( \begin{array}{c|c|c|c} | & | & \cdots & | \\ v_1 & v_2 & & v_n \\ | & | & & | \end{array} \right)$$

has a pivot in every row and every column, i.e. if  $A$  is \_\_\_\_\_.

## Bases of $\mathbb{R}^n$

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for  $\mathbb{R}^n$ .  The identity matrix has columns  $e_1, e_2, \dots, e_n$ .

1. They span:  $I_n$  has a pivot in every row.
2. They are linearly independent:  $I_n$  has a pivot in every column.

**In general:**  $\{v_1, v_2, \dots, v_n\}$  is a basis for  $\mathbb{R}^n$  if and only if the matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}$$

has a pivot in every row and every column, i.e. if  $A$  is *invertible*.

## Basis of a Subspace

### Example

#### Example

Let

$$V = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \text{ in } \mathbb{R}^3 \mid x + 3y + z = 0 \right\} \quad \mathcal{B} = \left\{ \begin{pmatrix} -3 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ -3 \end{pmatrix} \right\}.$$

Verify that  $\mathcal{B}$  is a basis for  $V$ .

0. In  $V$ : both vectors are in  $V$  because

$$-3 + 3(1) + 0 = 0 \quad \text{and} \quad 0 + 3(1) + (-3) = 0.$$

1. Span: If  $\begin{pmatrix} x \\ y \\ z \end{pmatrix}$  is in  $V$ , then  $y = -\frac{1}{3}(x + z)$ , so

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = -\frac{x}{3} \begin{pmatrix} -3 \\ 1 \\ 0 \end{pmatrix} - \frac{z}{3} \begin{pmatrix} 0 \\ 1 \\ -3 \end{pmatrix}.$$

2. Linearly independent:

$$c_1 \begin{pmatrix} -3 \\ 1 \\ 0 \end{pmatrix} + c_2 \begin{pmatrix} 0 \\ 1 \\ -3 \end{pmatrix} = 0 \implies \begin{pmatrix} -3c_1 \\ c_1 + c_2 \\ -3c_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \implies c_1 = c_2 = 0.$$



## Basis for $\text{Nul } A$

### Fact

The vectors in the parametric vector form of the general solution to  $Ax = 0$  always form a basis for  $\text{Nul } A$ .

## Basis for Nul $A$


### Fact

The vectors in the parametric vector form of the general solution to  $Ax = 0$  always form a basis for Nul  $A$ .

### Example

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

parametric vector form  $\xrightarrow{\text{~~~~~}}$   $x = x_3 \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \xrightarrow{\text{basis of Nul } A} \left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \right\}$



1. The vectors span Nul  $A$  by construction (every solution to  $Ax = 0$  has this form).
2. Can you see why they are linearly independent? (Look at the last two rows.)

## Basis for Col $A$

Fact

The *pivot columns* of  $A$  always form a basis for Col  $A$ .

## Basis for Col $A$

### Fact

The *pivot columns* of  $A$  always form a basis for Col  $A$ .

**Warning:** I mean the pivot columns of the *original* matrix  $A$ , not the row-reduced form.

## Basis for Col $A$

### Fact

The *pivot columns* of  $A$  always form a basis for Col  $A$ .

**Warning:** I mean the pivot columns of the *original* matrix  $A$ , not the row-reduced form. (Row reduction changes the column space.)

## Basis for Col A

### Fact

The *pivot columns* of  $A$  always form a basis for Col  $A$ .

**Warning:** I mean the pivot columns of the *original* matrix  $A$ , not the row-reduced form. (Row reduction changes the column space.)

### Example

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

pivot columns = basis  $\longleftrightarrow$  pivot columns in rref

So a basis for Col  $A$  is

$$\left\{ \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ -3 \\ 4 \end{pmatrix} \right\}.$$

## Basis for Col A

### Fact

The *pivot columns* of  $A$  always form a basis for Col  $A$ .

**Warning:** I mean the pivot columns of the *original* matrix  $A$ , not the row-reduced form. (Row reduction changes the column space.)

### Example

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

pivot columns = basis  $\longleftrightarrow$  pivot columns in rref

So a basis for Col  $A$  is

$$\left\{ \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ -3 \\ 4 \end{pmatrix} \right\}.$$

**Why?**

## Coefficients of Basis Vectors

**Recall:** a **basis** of a subspace  $V$  is a set of vectors that *spans*  $V$  and is *linearly independent*.



## Coefficients of Basis Vectors

**Recall:** a **basis** of a subspace  $V$  is a set of vectors that *spans*  $V$  and is *linearly independent*.

Lemma

← like a theorem, but less important

## Coefficients of Basis Vectors

**Recall:** a **basis** of a subspace  $V$  is a set of vectors that *spans*  $V$  and is *linearly independent*.

### Lemma

If  $\mathcal{B} \leftarrow \{v_1, v_2, \dots, v_m\}$  <sup>like a theorem, but less important</sup> is a basis for a subspace  $V$ , then any vector  $x$  in  $V$  can be written as a linear combination

$$x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m$$

for *unique* coefficients  $c_1, c_2, \dots, c_m$ .

We know  $x$  is a linear combination of the  $v_i$  because they span  $V$ . Suppose that we can write  $x$  as a linear combination with different coefficients:

$$x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m$$

$$x = c'_1 v_1 + c'_2 v_2 + \cdots + c'_m v_m$$

Subtracting:

$$0 = x - x = (c_1 - c'_1)v_1 + (c_2 - c'_2)v_2 + \cdots + (c_m - c'_m)v_m$$

Since  $v_1, v_2, \dots, v_m$  are linearly independent, they only have the trivial linear dependence relation. That means each  $c_i - c'_i = 0$ , or  $c_i = c'_i$ .

## Bases as Coordinate Systems

The unit coordinate vectors  $e_1, e_2, \dots, e_n$  form a basis for  $\mathbb{R}^n$ . Any vector is a unique linear combination of the  $e_i$ :

$$v = \begin{pmatrix} 3 \\ 5 \\ -2 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + 5 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} - 2 \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = 3e_1 + 5e_2 - 2e_3.$$

## Bases as Coordinate Systems

The unit coordinate vectors  $e_1, e_2, \dots, e_n$  form a basis for  $\mathbb{R}^n$ . Any vector is a unique linear combination of the  $e_i$ :

$$v = \begin{pmatrix} 3 \\ 5 \\ -2 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + 5 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} - 2 \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = 3e_1 + 5e_2 - 2e_3.$$

**Observe:** the *coordinates* of  $v$  are exactly the *coefficients* of  $e_1, e_2, e_3$ .

## Bases as Coordinate Systems

The unit coordinate vectors  $e_1, e_2, \dots, e_n$  form a basis for  $\mathbb{R}^n$ . Any vector is a unique linear combination of the  $e_i$ :

$$v = \begin{pmatrix} 3 \\ 5 \\ -2 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + 5 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} - 2 \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = 3e_1 + 5e_2 - 2e_3.$$

**Observe:** the *coordinates* of  $v$  are exactly the *coefficients* of  $e_1, e_2, e_3$ .

We can go backwards: given any basis  $\mathcal{B}$ , we interpret the coefficients of a linear combination as “coordinates” with respect to  $\mathcal{B}$ .

## Bases as Coordinate Systems

The unit coordinate vectors  $e_1, e_2, \dots, e_n$  form a basis for  $\mathbb{R}^n$ . Any vector is a unique linear combination of the  $e_i$ :

$$v = \begin{pmatrix} 3 \\ 5 \\ -2 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + 5 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} - 2 \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = 3e_1 + 5e_2 - 2e_3.$$

**Observe:** the *coordinates* of  $v$  are exactly the *coefficients* of  $e_1, e_2, e_3$ .

We can go backwards: given any basis  $\mathcal{B}$ , we interpret the coefficients of a linear combination as “coordinates” with respect to  $\mathcal{B}$ .

### Definition

Let  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  be a basis of a subspace  $V$ . Any vector  $x$  in  $V$  can be written uniquely as a linear combination  $x = c_1 v_1 + c_2 v_2 + \dots + c_m v_m$ . The coefficients  $c_1, c_2, \dots, c_m$  are the **coordinates of  $x$  with respect to  $\mathcal{B}$** . The  **$\mathcal{B}$ -coordinate vector of  $x$**  is the vector

$$[x]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix} \quad \text{in } \mathbb{R}^m.$$

# Bases as Coordinate Systems

## Example 1

$$\text{Let } v_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \quad \mathcal{B} = \{v_1, v_2\}, \quad V = \text{Span}\{v_1, v_2\}.$$

## Bases as Coordinate Systems

### Example 1

$$\text{Let } v_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \quad \mathcal{B} = \{v_1, v_2\}, \quad V = \text{Span}\{v_1, v_2\}.$$

**Verify** that  $\mathcal{B}$  is a basis:

*Span*: by definition  $V = \text{Span}\{v_1, v_2\}$ .

*Linearly independent*: because they are not multiples of each other.



# Bases as Coordinate Systems

## Example 1

Let  $v_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$ ,  $v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ ,  $\mathcal{B} = \{v_1, v_2\}$ ,  $V = \text{Span}\{v_1, v_2\}$ .

**Verify** that  $\mathcal{B}$  is a basis:

*Span*: by definition  $V = \text{Span}\{v_1, v_2\}$ .

*Linearly independent*: because they are not multiples of each other.

**Question**: If  $[x]_{\mathcal{B}} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}$ , then what is  $x$ ?

$$[x]_{\mathcal{B}} = \begin{pmatrix} 5 \\ 2 \end{pmatrix} \quad \text{means} \quad x = 5v_1 + 2v_2 = 5 \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + 2 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 7 \\ 2 \\ 7 \end{pmatrix}.$$

## Bases as Coordinate Systems

### Example 1

$$\text{Let } v_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \quad \mathcal{B} = \{v_1, v_2\}, \quad V = \text{Span}\{v_1, v_2\}.$$

**Verify** that  $\mathcal{B}$  is a basis:

*Span*: by definition  $V = \text{Span}\{v_1, v_2\}$ .

*Linearly independent*: because they are not multiples of each other.

**Question:** If  $[x]_{\mathcal{B}} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}$ , then what is  $x$ ?

$$[x]_{\mathcal{B}} = \begin{pmatrix} 5 \\ 2 \end{pmatrix} \quad \text{means} \quad x = 5v_1 + 2v_2 = 5 \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + 2 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 7 \\ 2 \\ 7 \end{pmatrix}.$$

**Question:** Find the  $\mathcal{B}$ -coordinates of  $x = \begin{pmatrix} 5 \\ 3 \\ 5 \end{pmatrix}$ .

We have to solve the vector equation  $x = c_1 v_1 + c_2 v_2$  in the unknowns  $c_1, c_2$ .

$$\left( \begin{array}{cc|c} 1 & 1 & 5 \\ 0 & 1 & 3 \\ 1 & 1 & 5 \end{array} \right) \rightsquigarrow \left( \begin{array}{cc|c} 1 & 1 & 5 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{array} \right) \rightsquigarrow \left( \begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{array} \right)$$

So  $c_1 = 2$  and  $c_2 = 3$ , so  $x = 2v_1 + 3v_2$  and  $[x]_{\mathcal{B}} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$ .

# Bases as Coordinate Systems

## Example 2

$$\text{Let } v_1 = \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix}, v_2 = \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}, v_3 = \begin{pmatrix} 2 \\ 8 \\ 6 \end{pmatrix}, \quad V = \text{Span}\{v_1, v_2, v_3\}.$$

## Bases as Coordinate Systems

### Example 2

$$\text{Let } v_1 = \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix}, v_2 = \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}, v_3 = \begin{pmatrix} 2 \\ 8 \\ 6 \end{pmatrix}, \quad V = \text{Span}\{v_1, v_2, v_3\}.$$

**Question:** Find a basis for  $V$ .

$V$  is the column span of the matrix

$$A = \begin{pmatrix} 2 & -1 & 2 \\ 3 & 1 & 8 \\ 2 & 1 & 6 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}.$$

A basis for the column span is formed by the pivot columns:  $\mathcal{B} = \{v_1, v_2\}$ .

## Bases as Coordinate Systems

### Example 2

$$\text{Let } v_1 = \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix}, v_2 = \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}, v_3 = \begin{pmatrix} 2 \\ 8 \\ 6 \end{pmatrix}, \quad V = \text{Span}\{v_1, v_2, v_3\}.$$

**Question:** Find a basis for  $V$ .

$V$  is the column span of the matrix

$$A = \begin{pmatrix} 2 & -1 & 2 \\ 3 & 1 & 8 \\ 2 & 1 & 6 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}.$$

A basis for the column span is formed by the pivot columns:  $\mathcal{B} = \{v_1, v_2\}$ .

**Question:** Find the  $\mathcal{B}$ -coordinates of  $x = \begin{pmatrix} 4 \\ 11 \\ 8 \end{pmatrix}$ .

We have to solve  $x = c_1 v_1 + c_2 v_2$ .

$$\left( \begin{array}{cc|c} 2 & -1 & 4 \\ 3 & 1 & 11 \\ 2 & 1 & 8 \end{array} \right) \xrightarrow{\text{row reduce}} \left( \begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{array} \right)$$

So  $x = 3v_1 + 2v_2$  and  $[x]_{\mathcal{B}} = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$ .

# Bases as Coordinate Systems

## Summary

If  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  is a basis for a subspace  $V$  and  $x$  is in  $V$ , then

$$[x]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix} \quad \text{means} \quad x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m.$$

# Bases as Coordinate Systems

## Summary

If  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  is a basis for a subspace  $V$  and  $x$  is in  $V$ , then

$$[x]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix} \quad \text{means} \quad x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m.$$

Finding the  $\mathcal{B}$ -coordinates for  $x$  means solving the vector equation

$$x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m$$

in the unknowns  $c_1, c_2, \dots, c_m$ .

# Bases as Coordinate Systems

## Summary

If  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  is a basis for a subspace  $V$  and  $x$  is in  $V$ , then

$$[x]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix} \quad \text{means} \quad x = c_1 v_1 + c_2 v_2 + \dots + c_m v_m.$$

Finding the  $\mathcal{B}$ -coordinates for  $x$  means solving the vector equation

$$x = c_1 v_1 + c_2 v_2 + \dots + c_m v_m$$

in the unknowns  $c_1, c_2, \dots, c_m$ . This (usually) means row reducing the augmented matrix

$$\left( \begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_m & x \\ | & | & & | & | \end{array} \right).$$



# Bases as Coordinate Systems

## Summary

If  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  is a basis for a subspace  $V$  and  $x$  is in  $V$ , then

$$[x]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix} \quad \text{means} \quad x = c_1 v_1 + c_2 v_2 + \dots + c_m v_m.$$

Finding the  $\mathcal{B}$ -coordinates for  $x$  means solving the vector equation

$$x = c_1 v_1 + c_2 v_2 + \dots + c_m v_m$$

in the unknowns  $c_1, c_2, \dots, c_m$ . This (usually) means row reducing the augmented matrix

$$\left( \begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_m & x \\ | & | & & | & | \end{array} \right).$$

**Question:** What happens if you try to find the  $\mathcal{B}$ -coordinates of  $x$  *not* in  $V$ ? You end up with an inconsistent system:  $V$  is the span of  $v_1, v_2, \dots, v_m$ , and if  $x$  is not in the span, then  $x = c_1 v_1 + c_2 v_2 + \dots + c_m v_m$  has no solution.

# Bases as Coordinate Systems

Picture


Let

$$v_1 = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

These form a basis  $\mathcal{B}$  for the plane

$$V = \text{Span}\{v_1, v_2\}$$

in  $\mathbb{R}^3$ .



# Bases as Coordinate Systems

Picture


Let

$$v_1 = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

These form a basis  $\mathcal{B}$  for the plane

$$V = \text{Span}\{v_1, v_2\}$$

in  $\mathbb{R}^3$ .



**Question:** Estimate the  $\mathcal{B}$ -coordinates of these vectors:

$$[u_1]_{\mathcal{B}} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad [u_2]_{\mathcal{B}} = \begin{pmatrix} -1 \\ \frac{1}{2} \end{pmatrix} \quad [u_3]_{\mathcal{B}} = \begin{pmatrix} \frac{3}{2} \\ -\frac{1}{2} \end{pmatrix} \quad [u_4]_{\mathcal{B}} = \begin{pmatrix} 0 \\ \frac{3}{2} \end{pmatrix}$$

# Bases as Coordinate Systems

Picture


Let

$$v_1 = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

These form a basis  $\mathcal{B}$  for the plane

$$V = \text{Span}\{v_1, v_2\}$$

in  $\mathbb{R}^3$ .



**Question:** Estimate the  $\mathcal{B}$ -coordinates of these vectors:

$$[u_1]_{\mathcal{B}} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad [u_2]_{\mathcal{B}} = \begin{pmatrix} -1 \\ \frac{1}{2} \end{pmatrix} \quad [u_3]_{\mathcal{B}} = \begin{pmatrix} \frac{3}{2} \\ -\frac{1}{2} \end{pmatrix} \quad [u_4]_{\mathcal{B}} = \begin{pmatrix} 0 \\ \frac{3}{2} \end{pmatrix}$$

**Remark**

Many of you want to think of a plane in  $\mathbb{R}^3$  as “being”  $\mathbb{R}^2$ . Choosing a basis  $\mathcal{B}$  and using  $\mathcal{B}$ -coordinates is one way to make sense of that. But remember that the coordinates are the coefficients of a linear combination of the basis vectors.

# The Rank Theorem

Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the



# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$$\text{rank } A = \dim \text{Col } A =$$

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$\text{rank } A = \dim \text{Col } A = \text{the number of columns with pivots}$

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$$\begin{aligned}\text{rank } A &= \dim \text{Col } A = \text{the number of columns with pivots} \\ \dim \text{Nul } A &= \end{aligned}$$

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$$\begin{aligned}\text{rank } A &= \dim \text{Col } A = \text{the number of columns with pivots} \\ \dim \text{Nul } A &= \text{the number of free variables}\end{aligned}$$

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$$\begin{aligned}\text{rank } A &= \dim \text{Col } A = \text{the number of columns with pivots} \\ \dim \text{Nul } A &= \text{the number of free variables} \\ &= \text{the number of columns without pivots.}\end{aligned}$$

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$$\begin{aligned}\text{rank } A &= \dim \text{Col } A = \text{the number of columns with pivots} \\ \dim \text{Nul } A &= \text{the number of free variables} \\ &= \text{the number of columns without pivots.}\end{aligned}$$

## Rank Theorem

If  $A$  is an  $m \times n$  matrix, then

$$\text{rank } A + \dim \text{Nul } A =$$



# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$$\begin{aligned}\text{rank } A &= \dim \text{Col } A = \text{the number of columns with pivots} \\ \dim \text{Nul } A &= \text{the number of free variables} \\ &= \text{the number of columns without pivots.}\end{aligned}$$

## Rank Theorem

If  $A$  is an  $m \times n$  matrix, then

$$\text{rank } A + \dim \text{Nul } A = n = \text{the number of columns of } A.$$

# The Rank Theorem

## Example

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

A basis for Col  $A$  is

$$\left\{ \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ -3 \\ 4 \end{pmatrix} \right\},$$

so  $\text{rank } A = \dim \text{Col } A = 2$ .

Since there are two free variables  $x_3, x_4$ , the parametric vector form for the solutions to  $Ax = 0$  is

$$x = x_3 \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \xrightarrow{\text{basis for Nul } A} \left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

Thus  $\dim \text{Nul } A = 2$ .

The Rank Theorem says  $2 + 2 = 4$ .

# The Rank Theorem

## Example

$$A = \begin{pmatrix} \boxed{1} & \boxed{2} & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

basis of Col A

A basis for Col A is

$$\left\{ \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ -3 \\ 4 \end{pmatrix} \right\},$$

so  $\text{rank } A = \dim \text{Col } A = 2$ .

Since there are two free variables  $x_3, x_4$ , the parametric vector form for the solutions to  $Ax = 0$  is

$$x = x_3 \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \xrightarrow{\text{basis for Nul } A} \left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

Thus  $\dim \text{Nul } A = 2$ .

The Rank Theorem says  $2 + 2 = 4$ .

# The Rank Theorem

## Example

$$A = \begin{pmatrix} \boxed{1} & \boxed{2} & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & \boxed{-8} & \boxed{-7} \\ 0 & 1 & \boxed{4} & \boxed{3} \\ 0 & 0 & \boxed{0} & \boxed{0} \end{pmatrix}$$

basis of Col A free variables

A basis for Col A is

$$\left\{ \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ -3 \\ 4 \end{pmatrix} \right\},$$

so  $\text{rank } A = \dim \text{Col } A = 2$ .

Since there are two free variables  $x_3, x_4$ , the parametric vector form for the solutions to  $Ax = 0$  is

$$x = x_3 \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \xrightarrow{\text{basis for Nul } A} \left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

Thus  $\dim \text{Nul } A = 2$ .

The Rank Theorem says  $2 + 2 = 4$ .

## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?

## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?

If  $AB = 0$ , then  $ABx = 0$  for every  $x$  in  $\mathbb{R}^3$ .

## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?

If  $AB = 0$ , then  $ABx = 0$  for every  $x$  in  $\mathbb{R}^3$ .

This means  $A(Bx) = 0$ , so  $Bx$  is in  $\text{Nul } A$ .

## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?

If  $AB = 0$ , then  $ABx = 0$  for every  $x$  in  $\mathbb{R}^3$ .

This means  $A(Bx) = 0$ , so  $Bx$  is in  $\text{Nul } A$ .

This is true for every  $x$ , so  $\text{Col } B$  is contained in  $\text{Nul } A$ .



## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?

If  $AB = 0$ , then  $ABx = 0$  for every  $x$  in  $\mathbb{R}^3$ .

This means  $A(Bx) = 0$ , so  $Bx$  is in  $\text{Nul } A$ .

This is true for every  $x$ , so  $\text{Col } B$  is contained in  $\text{Nul } A$ .

But  $\dim \text{Nul } A = 1$  and  $\dim \text{Col } B = 2$ , and a 1-dimensional space can't contain a 2-dimensional space.

## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?

If  $AB = 0$ , then  $ABx = 0$  for every  $x$  in  $\mathbb{R}^3$ .

This means  $A(Bx) = 0$ , so  $Bx$  is in  $\text{Nul } A$ .

This is true for every  $x$ , so  $\text{Col } B$  is contained in  $\text{Nul } A$ .

But  $\dim \text{Nul } A = 1$  and  $\dim \text{Col } B = 2$ , and a 1-dimensional space can't contain a 2-dimensional space.

Hence it can't happen.

## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?


If  $AB = 0$ , then  $ABx = 0$  for every  $x$  in  $\mathbb{R}^3$ .

This means  $A(Bx) = 0$ , so  $Bx$  is in  $\text{Nul } A$ .


This is true for every  $x$ , so  $\text{Col } B$  is contained in  $\text{Nul } A$ .

But  $\dim \text{Nul } A = 1$  and  $\dim \text{Col } B = 2$ , and a 1-dimensional space can't contain a 2-dimensional space.

Hence it can't happen.



does not  
contain



# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

- ▶ Any  $m$  linearly independent vectors in  $V$  form a basis for  $V$ .

# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

- ▶ Any  $m$  linearly independent vectors in  $V$  form a basis for  $V$ .
- ▶ Any  $m$  vectors that span  $V$  form a basis for  $V$ .

# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

- ▶ Any  $m$  linearly independent vectors in  $V$  form a basis for  $V$ .
- ▶ Any  $m$  vectors that span  $V$  form a basis for  $V$ .

### Upshot

If you *already* know that  $\dim V = m$ , and you have  $m$  vectors  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  in  $V$ , then you only have to check *one* of

# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

- ▶ Any  $m$  linearly independent vectors in  $V$  form a basis for  $V$ .
- ▶ Any  $m$  vectors that span  $V$  form a basis for  $V$ .

### Upshot

If you *already* know that  $\dim V = m$ , and you have  $m$  vectors  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  in  $V$ , then you only have to check *one* of

1.  $\mathcal{B}$  is linearly independent, *or*



# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

- ▶ Any  $m$  linearly independent vectors in  $V$  form a basis for  $V$ .
- ▶ Any  $m$  vectors that span  $V$  form a basis for  $V$ .

### Upshot

If you *already* know that  $\dim V = m$ , and you have  $m$  vectors  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  in  $V$ , then you only have to check *one* of

1.  $\mathcal{B}$  is linearly independent, *or*
2.  $\mathcal{B}$  spans  $V$

# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

- ▶ Any  $m$  linearly independent vectors in  $V$  form a basis for  $V$ .
- ▶ Any  $m$  vectors that span  $V$  form a basis for  $V$ .

### Upshot

If you *already* know that  $\dim V = m$ , and you have  $m$  vectors  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  in  $V$ , then you only have to check *one* of

1.  $\mathcal{B}$  is linearly independent, *or*
2.  $\mathcal{B}$  spans  $V$

in order for  $\mathcal{B}$  to be a basis.