

BU5526 - Portfolio Analysis
Lecture Notes

Rodrigo Miguel

January 29, 2023

Contents

1	Lecture 1	2
1.1	Asset Pricing	2
1.1.1	Asset	2
1.1.2	Asset Pricing	2
1.1.3	How to price a stock:	2
1.2	Portfolio Management	3
1.3	Mathematical Concepts	3
1.3.1	Mean	3
1.3.2	Variance	3
1.3.3	Important Statistical Concept	4
1.3.4	Standard Deviation	4
1.3.5	Skewness	4
1.3.6	Kurtosis	4
1.4	Assumptions of Mean-Variance Analysis	4
1.4.1	Normal distribution	5
1.5	Returns on Financial Assets	5
1.5.1	Holding period return	5
1.5.2	Geometric Mean Return	5
1.5.3	Annualized Return	5
1.5.4	Portfolio Return	6
1.5.5	Historical and Expected Returns	6
1.5.6	Ways of Calculating Expected Returns	6
1.5.7	Standard deviation - Volatility	6
1.5.8	Portfolio Variance	6
1.5.9	Correlation and Portfolio Risk	7
2	Lecture 2	8

Chapter 1

Lecture 1

1.1 Asset Pricing

1.1.1 Asset

Something valuable that an entity owns, benefits from, or has use of, in generating income.

1.1.2 Asset Pricing

Needed to buy/sell at a fair price.

- Obtained by discovery process:
 - Demand and supply forces.
- The price of an asset is simply the current value of its cash-flows.

$$PV = \frac{FV_1}{(1+r)^1} + \frac{FV_2}{(1+r)^2} + \dots + \frac{FV_n}{(1+r)^n}$$

- What we need:
 - Prediction of cash flows;
 - Discount rate.

1.1.3 How to price a stock:

- Dividend Discount Model;
- FCF Model;
- Multipliers and Comparables approach.

Being able to estimate the required **rate of return** is everything you need to calculate the **value of any asset** - once you have predicted **future cash-flows**, including:

- Estimating the value of a stock;
- Estimating the value of a firm.

1.2 Portfolio Management

Prudent administration of **investable** (liquid) assets, aimed at achieving an **optimum risk-reward ratio**.

1.3 Mathematical Concepts

1.3.1 Mean

An average of different observations.

- Useful to describe a population;
- **Arithmetical**: sum of observations divided by the number of observations (if with equal weights).

$$\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n}$$

or

$$\bar{X} = W_1 \times X_1 + W_2 \times X_2 + \dots + W_n \times X_n$$

- **Geometrical**: product of observations divided by the number of observations.

$$\bar{X}_G = \sqrt[n]{X_1 \times X_2 \times \dots \times X_n}$$

or

$$\bar{X}_G = \sqrt[n]{X_1^{W_1} \times X_2^{W_2} \times \dots \times X_n^{W_n}}$$

There is a reason to use arithmetical or geometrical mean.

- **Equally weighted**: when all the observations have the same importance.
- **Unequally weighted**: different importance for different observations.

1.3.2 Variance

Seen as an extension of the mean.

- Dispersion to the mean (higher/lower):
 - Average of differences from the mean.

$$\sigma_x^2 = \frac{(X_1 - \mu_X)^2 + (X_2 - \mu_X)^2 + \dots + (X_n - \mu_X)^2}{n}$$

1.3.3 Important Statistical Concept

- The formula is correct if we possess all the data on the population;
- If we only have a sample, we need to reflect "impreciseness" by removing "one degree of freedom".

$$\sigma_x^2 = \frac{(X_1 - \mu_X)^2 + (X_2 - \mu_X)^2 + \dots + (X_n - \mu_X)^2}{n - 1}$$

This will **always** be the case in finance.

1.3.4 Standard Deviation

$$\sigma_X = \sqrt{\frac{(X_1 - \mu_n)^2 + (X_2 - \mu_n)^2 + \dots + (X_n - \mu_n)^2}{(n - 1)}}$$

1.3.5 Skewness

Brings back the sign. A positive skewness means more positive-value and reversely.

$$\sigma_X^3 = \frac{(X_1 - \mu_n)^3 + (X_2 - \mu_n)^3 + \dots + (X_n - \mu_n)^3}{n}$$

1.3.6 Kurtosis

Outweighs extremes - dropping the sign.

$$\sigma_X^4 = \frac{(X_1 - \mu_n)^4 + (X_2 - \mu_n)^4 + \dots + (X_n - \mu_n)^4}{n}$$

A large Kurtosis means a lot of extreme values.

- To get a **meaningful estimate: excess** kurtosis needs to be provided.
- Kurtosis of a normal distribution is 3.

Note: Unbiased equations of these two indicators are slightly more complex, but computer packages provide them automatically.

1.4 Assumptions of Mean-Variance Analysis

- Allows describing different assets simply.
- Assumes returns are normally distributed.

1.4.1 Normal distribution

- Its mean and median are equal;
- It's defined by two parameters, mean and variance;
- It's defined around its mean with:
 - 68% of observations within $\pm 1\sigma$ of the mean.
 - 95% of observations within $\pm 2\sigma$ of the mean.
 - 99% of observations within $\pm 3\sigma$ of the mean
- Returns are not normally distributed.
 - Skewed: not symmetric around the mean.
 - Characterized by high probability of extreme event.

1.5 Returns on Financial Assets

1.5.1 Holding period return

Return from holding an asset for a specific period of time.

$$R = \frac{P_t - P_{t-1}}{P_{t-1}} + \frac{D_t}{P_{t-1}}$$

Capital gain + Dividend yield

Holding period returns = compound returns

$$R = [(1 + r_1) \times (1 + r_2) \times (1 + r_3)] - 1$$

1.5.2 Geometric Mean Return

$$\bar{R} = \sqrt[T]{(1 + r_{i_1}) \times (1 + r_{i_2}) \times \dots \times (1 + r_{i_T})} - 1$$

1.5.3 Annualized Return

These returns should be capitalized.

- Such as:

$$r_{annual} = (1 + r_{period})^C - 1$$

with C being the number of periods in a year.

1.5.4 Portfolio Return

- When several assets are combined into a portfolio, we can compute the portfolio return.
- Weighted average of the returns of individual assets

$$R_p = W_1 \times R_1 + W_2 \times R_2$$

1.5.5 Historical and Expected Returns

- Computed from historical data.
- What the investor expects to earn.
- Historical returns are different from expected returns.

1.5.6 Ways of Calculating Expected Returns

- Gut feeling;
- Modelling - calculated from a formula.

1.5.7 Standard deviation - Volatility

- Historical standard deviation;
- Defined as **risk** of equity return.

1.5.8 Portfolio Variance

- Cannot simply add up two variances.
- Covariance:

$$cov(X, Y) = E(XY) - E(X) \times E(Y)$$

- The more the two assets move in the same way, the higher the covariance.
- A negative covariance means the assets move in opposite directions.

- Variance:

$$\sigma_v^2 = wCw^T$$

1.5.9 Correlation and Portfolio Risk

The correlation among assets determine the portfolio's risk.

- Is a measure of tendency for N investments to act similarly;
- Can range from -1 and $+1$.

$$\rho_{ij} = \frac{\text{cov}(R_i, R_j)}{\sigma_i \sigma_j}$$

Chapter 2

Lecture 2