Enhanced Expression of Glucose 2-Oxidase in *Phlebia tremellosa* by Addition of Phthalates

Baikjoong Kim, Hyewon Kim and Hyoung T. Choi*

Department of Biochemistry, Kangwon National University, Chuncheon 200-701, Korea (Received October 25, 2010. Accepted November 15, 2010)

Most fungi possess several hydrogen peroxide-generating enzymes, glucose oxidase and pyranose oxidase. Pyranose oxidase can use glucose as its substrate to generate hydrogen peroxide. White rot fungi, which degrade diverse recalcitrant compounds, contain lignin-degrading enzymes, and lignin peroxidase and manganese peroxidase require hydrogen peroxide for their enzymatic reactions. In this study, we isolated a cDNA fragment of pyranose oxidase from *Phlebia tremellosa* using PCR and examined its expression under the degradation conditions of diethylphthalate (DEP). Pyranose oxidase expression was enhanced up to 30% by the addition of DEP, and this result supports the possible involvement of pyranose oxidase in the degradation of recalcitrant compounds.

KEYWORDS: Hydrogen peroxide generation, Phlebia tremellosa, Pyranose oxidase

White rot fungi, which degrade lignins, possess related enzymes such as laccase, lignin peroxidase (LiP), and manganese peroxidase (MnP). Since these enzymes show relatively broad substrate specificities, white rot fungi can degrade polycyclic aromatic hydrocarbons, explosives, and endocrine-disrupting chemicals. We have previously reported the degradation of explosives [1] and endocrine-disrupting chemicals [2, 3]. We have also cloned the genes for laccase and MnP from *Phlebia tremellosa* and *Trametes versicolor* as well as constructed expression vectors. Genetic transformants using the expression vectors have been generated, and they show enhanced degrading activities against recalcitrant compounds [4-6].

Even though the original compounds of endocrine-disrupting chemicals in culture supernatants were removed up to 90% during the first 2~3 days, there were 30~40% residual estrogenic activities in the culture supernatants. This unbalanced degradation of endocrine-disrupting chemicals was observed in the wild type as well as in the genetic transformants of laccase or MnP [4-6]. This result implies that both of the original compounds and the degradation intermediates possessed estrogenic activities. Gartshore et al. [7] reported that the degradation intermediates of five different plasticizers have toxic effects, and phthalates can be changed to have strong estrogenic activities by UV irradiation [8]. These results also suggest that even though the original compounds were almost completely removed, the degradation intermediates still showed estrogenic activities in our previous experiments.

LiP and MnP require hydrogen peroxide as a co-substrate in their enzymatic reactions. Glucose oxidase (EC

1.1.3.4) catalyzes the oxidation of glucose to hydrogen peroxide and glucono-δ-lactone. Pyranose oxidase (EC 1.1.3.10; POx) is another enzyme that generates hydrogen peroxide; it produces hydrogen peroxide and 2-dehydro-glucose and therefore has another name, glucose 2oxidase. Since genetic transformants containing extra copies of laccase or MnP cDNA are not fully satisfactory as mentioned earlier, we tried to examine other enzymes that would efficiently remove the estrogenic activity of an endocrine-disrupting chemical. In the first step, we determined whether or not POx is involved in the degradation of endocrine-disrupting chemicals. Since we used a white rot basidiomycete, P. tremellosa, for the degradation of endocrine-disrupting chemicals, isolation of a cDNA fragment of pyranose oxidase from this fungus was performed.

POx cDNA (D73369) was found in a white rot fungus, *T. versicolor*, and the DNA sequence was used for construction of PCR primers as follows: forward primer 5'-ATGTCTACTAGCTCGAGCGA-3' and reverse primer 5'-GGGTGGTAACAGCCATACGC-3'. Total RNAs were isolated from the *P. tremellosa* liquid culture, and the cDNAs were synthesized by following a previous report [3]. PCR with the primers resulted in isolation of a cDNA fragment consisting of 1,872 bp, including the open reading frame and terminator. The nucleotide sequence showed 97% identity with that of POx of *T. versicolor*, and the deduced amino acid sequence showed 99% identity with the same enzyme of *T. versicolor* (Fig. 1). This was reported to the EMBL nucleotide sequence database with the accession number FR671428.

The degradation of phthalates by *P. tremellosa* has been reported [3], and genetic transformants containing extra-

^{*}Corresponding author <E-mail: htchoi@kangwon.ac.kr>





Fig. 1. Comparison of amino acid sequences of pyranose oxidase between *Phlebia tremellosa* and *Trametes versicolor*. Hyphens indicate where different amino acids from the two fungal species are encoded.

copies of the laccase gene show increased degrading activity against phthalates [6]. In order to determine whether or not isolated POx is involved in the degradation of phthalates, analysis of diethylphthalate (DEP) degradation was performed as follows. P. tremellosa was grown in liquid minimal medium (100 mL in 250 mL flask) at 30°C for 5 days with shaking, and the culture was homogenized five times for 1 sec each using a Waring blender [3]. The homogenized cells were transferred to 200 mL of the same medium containing DEP (200 mg/L) (inoculums size; 10% v/v) and incubated for 3 days. Quantitative determination of DEP from culture supernatant and culture medium with EDC was carried out using a Waters HPLC (HP 1,525 series, Gemini 5 µm C6-phenyl 110A 150 × 4.6 mm column; Waters, Milford, MA, USA) as reported previously [5]. The residual concentration of DEP was decreased to 65~70% level in the culture supernatant when compared to that of control (Table 1). The gene expression of POx was determined by reverse transcriptase-PCR using the total RNAs from the 3-day grown cultures and the POxspecific primers (forward primer 5'-GCACCCGGACTG-GTGGAACG-3'; reverse primer 5'-CGCAGCCACCGAG-GAACAGG-3'), with actin used as the control. The

Table 1. Removal of diethylphthalate by *Phlebia tremellosa* on day 3

Strain	Removal of diethylphthalate (%)
Control	0 ± 1.5
P. tremellosa	33 ± 2.5



POx Actin

Fig. 2. Determination of pyranose oxidase (POx) expression by reverse transcriptase-PCR using POx-specific primers. Left lane, MM culture; Right lane, MM + DEP culture. MM, minimal medium; DEP, diethylphthalate.

expected length (542 bp) of the PCR product was amplified in both the control and DEP-degrading cultures, and POx gene expression was enhanced about 30% by the addition of DEP (Fig. 2). It has been reported that POx from white rot fungi can supply H₂O₂ during the degradation of lignin and recalcitrant compounds [9, 10], and the enhanced expression of POx gene under the degradation conditions of DEP shows that POx is possibly involved during the degradation of various recalcitrant compounds in *P. tremellosa*.

Acknowledgements

This research was partly supported by a grant (# 2009-0073676) from the Ministry of Science, Education and Technology.

References

- Cheong S, Yeo S, Song HG, Choi HT. Determination of laccase gene expression during degradation of 2,4,6-trinitrotoluene and its catabolic intermediates in *Trametes versicolor*. Microbiol Res 2006;161:316-20.
- Kang AR, Choi HT, Song HG. Optimization of bisphenol A biodegradation by *Trametes versicolor*. Korean J Microbiol 2008;44:37-42.
- 3. Yeo S, Kim MK, Choi HT. Increased expression of laccase by the addition of phthalates in *Phlebia tremellosa*. FEMS Microbiol Lett 2008;278:72-7.
- Kum H, Kim M, Choi HT. Biodegradation of endocrine disrupting chemicals by genetic transformants of *Phlebia tremellosa* using manganese peroxidase gene from *Trametes versicolor*. Korean J Microbiol 2009;45:82-5.
- Kum H, Kim MK, Choi HT. Degradation of endocrine disrupting chemicals by genetic transformants in *Irpex lacteus* with an inducible laccase gene of *Phlebia tremellosa*. Biodegradation 2009;20:673-8.
- Yeo S, Kim M, Choi HT. Degradation of endocrine disrupting chemicals by laccase transformants of *Phlebia tremellosa*. Korean J Microbiol 2008;44:10-3.

66 Kim et al.

7. Gartshore J, Cooper DG, Nicell JA. Biodegradation of plast-cizers by *Rhodotorula rubra*. Environ Toxicol Chem 2003; 22:1244-51.

- 8. Okamoto Y, Hayashi T, Toda C, Ueda K, Hashizume K, Itoh K, Nishikawa J, Nishihara T, Kojima N. Formation of estrogenic products from environmental phthalate esters under light exposure. Chemosphere 2006;64:1785-92.
- 9. Ander P, Marzullo L. Sugar oxidoreductases and veratryl alcohol oxidase as related to lignin degradation. J Biotechnol 1997;53:115-31.
- 10. Giffhorn F. Fungal pyranose oxidases: occurrence, properties and biochemical applications in carbohydrate chemistry. Appl Microbiol Biotechnol 2000;54:727-40.