

Anti-hyperglycemic effect of the polysaccharides fraction from American ginseng berry extract in *ob/ob* mice

J.T. Xie^{1,2}, J.A. Wu^{1,2}, S. Mehendale^{1,2}, H.H. Aung^{1,2}, and C.-S. Yuan^{1,2,3}

¹Tang Center for Herbal Medicine Research

²Departments of Anesthesia & Critical Care

³Committee on Clinical Pharmacology and Pharmacogenomics, The Pritzker School of Medicine, The University of Chicago, USA

Summary

In this study, we evaluated the anti-hyperglycemic effect of a polysaccharides fraction from American ginseng berry extract in diabetic *ob/ob* mice. All animals received daily intraperitoneal injections of polysaccharides at 150 mg/kg body wt. ($n = 5$), polysaccharides at 50 mg/kg body wt. ($n = 5$), or vehicle ($n = 5$) for 10 consecutive days. On Day 5, as compared to the vehicle-treated mice (230.5 ± 13.5 mg/dl, mean \pm S.E), mice from both treated groups showed significantly lower fasting blood glucose levels (187.4 ± 20.5 mg/dl and 187.4 ± 17.1 mg/dl), respectively (both $P < 0.05$). On Day 10, compared to the vehicle group (240.1 ± 12.3 mg/dl), the 50 mg/kg dose group were at 188.4 ± 12.6 mg/dl ($P < 0.05$), and the 150 mg/kg dose group were normoglycemic (148.8 ± 17.6 mg/dl, $P < 0.01$). Those *ob/ob* mice treated with vehicle did not, however, show significant changes in fasting blood glucose levels. Data from the intraperitoneal glucose tolerance test (IPGTT) showed that, compared to Day 0, there was a significant improvement in glucose tolerance in animals who received the 50 and 150 mg/kg polysaccharide doses, and the area under the curve (AUC) decreased 15.5% ($P < 0.05$) and 28.2% ($P < 0.01$), respectively. Interestingly, after cessation of polysaccharide treatment, the fasting blood glucose levels stayed lower, and returned to control concentration on Day 30. We also observed that the polysaccharides fraction did not affect body weight changes in *ob/ob* mice. Our data suggest that the polysaccharides fraction from American ginseng berry extract has a potential clinical utility in treating diabetic patients.

Key words: American ginseng, ginseng berry, polysaccharides fraction, anti-hyperglycemic, diabetic *ob/ob* mice

Introduction

Ginseng root is a valuable herb in oriental medicine and has long been used as a remedy for a number of disorders. The root of ginseng has been utilized for over 2,000 years, in the belief that it is a panacea, tonic, roborant, and promotes longevity. The efficacy of ginseng root was known in the West by the eighteenth century, and the study of ginseng has a long history (Huang, 1999).

Historical records reveal that in traditional medical systems, a disease corresponding to type-2 diabetes was treated with plant extracts (Ackerknecht, 1982). Several reports have demonstrated that *Panax ginseng* C. A. Meyer (Asian ginseng) root has hypoglycemic effect both in normal and alloxan-induced hyperglycemic animals (Wang and Lei, 1957; Huang, 1999). *Panax quinquefolius* L. (American ginseng), another

common species of ginseng, is grown in the northern U.S. and Canada, but is also cultivated in France and northern China (Bensky and Gamble, 1993). Previous *in vitro* and *in vivo* animal studies (Kimura, 1980; Yokozawa et al. 1985; Kimura et al. 1999) and clinical trials (Sotaniemi et al. 1995; Vuksan et al. 2000a, 2000b) support the claim that the roots of Asian ginseng and American ginseng possess anti-hyperglycemic activities. Recently, we observed that extract of the berry (or fruit) of American ginseng has significant anti-hyperglycemic and anti-obesity effects in diabetic *ob/ob* mice (Xie et al. 2002a). In *ob/ob* mice, the *ob* gene was transferred from the stock of origin onto the B/6 genomic background and is located on chromosome 6 (Shafrir, 1992). Animals that are homozygous for the mutation exhibit morbid obesity and metabolic abnormalities that phenotypically resemble type 2 diabetes in humans.

Ginseng roots consist of a number of active constituents, such as ginsenosides, polysaccharides, peptides, polyacetylenic alcohols, and fatty acids (Li and Zhang, 1986; Cao, 1989; Lee, 1992; Attele et al. 1999). These constituents are also found in ginseng berry, in different proportions (Zhang and Jiang, 1981; Liu et al. 1988; Wang et al. 2000; Hao et al. 2000). It has been shown that polysaccharides from ginseng root possess a hypoglycemic effect in alloxan-induced hyperglycemic mice (Konno and Hikino, 1987). However, the effect of polysaccharides fraction from the berry of American ginseng in diabetic *ob/ob* mice has not been investigated. The present study was designed to evaluate anti-diabetic effect of polysaccharides fraction from American ginseng berry extract in diabetic *ob/ob* mice.

Materials and Methods

Animals

The study protocol was approved by the Institutional Animal Care and Use Committee of the University of Chicago. Male C57BL/6J *ob/ob* mice were obtained from Jackson Laboratory (Bar Harbor, ME). Mice were housed in environmentally controlled conditions with a 12-h light/dark cycle. All animals had free access to standard rodent pellet food (Zeigler Bros., Gardners, PA) and water *ad libitum*, except when fasted before experiments. Animals of 10–15 weeks of age were used in the study.

Preparation of polysaccharides fraction from American ginseng berry extract

Fresh American ginseng berry was obtained from a ginseng farm in Wausau, WI. In brief, 500 g of berry was mixed with 500 ml organic solvent. After removal

of the seeds, 356 g pulp was collected. An additional 500 ml organic solvent was added, and the solution was heated and refluxed. This procedure was repeated three times. The organic solvent was then evaporated and the solution was filtered. The filtered solution contained both polysaccharides and ginsenosides.

The remaining solution was loaded onto a Dialon HP-20 gel column (Supelco, VA), and the passed solution collected. Afterwards, the column was washed with distilled water several times, until the color of the solution disappeared. All collected solutions were mixed and filtered to obtain the polysaccharides fraction (Ma et al. 1998; Sung et al. 2000). Finally, the extract was lyophilized. The extraction rate was approximately 2% from the fresh ginseng berry.

Experimental protocol

Fifteen *ob/ob* mice received daily intraperitoneal (IP) injections, and were divided into three groups. The first drug group ($n = 5$) received 150 mg/kg body wt. of polysaccharides fraction of American ginseng berry extract. The second drug group ($n = 5$) received 50 mg/kg body wt. of polysaccharides fraction. The third vehicle group ($n = 5$) received vehicle (0.9% NaCl). No noticeable irritation or restlessness was observed following administration of the extract or vehicle solutions.

As our previous reports (Attele et al. 2002; Xie et al. 2002b), fasting blood glucose levels and body weight were measured after fasting the animals for 4 h (starting from 9:00 AM) on Day 0 (before treatment), Day 5, and Day 10. Blood glucose levels were determined from the tail vein blood samples at 1:00 PM using a Glucose Analyzer (Hemocue AB, Angelholm, Sweden). After cessation of consecutive 10-day administration, fasting blood glucose levels and body weights were measured every five days, until the effect of polysaccharides disappeared.

Intraperitoneal glucose tolerance test (IPGTT) was performed on Day 0 and Day 10. On days of the test, animals fasted for 4 h (starting from 9:00 AM), followed by an IP administration of glucose (2 g/kg body wt.). Blood glucose levels were determined in blood samples from the tail vein at 0 (prior to glucose administration), 30, 60 and 120 min after glucose administration.

Statistical Analysis

Data are expressed as mean \pm standard error. Statistical significance between vehicle-treated mice vs. drug-treated mice and between before treatment and after treatment were determined by paired Student's *t* test. A value of $P < 0.05$ was considered statistically significant.

Results

Effect of the polysaccharides fraction on fasting blood glucose levels

As shown in Fig. 1, diabetic *ob/ob* mice had remarkably high baseline fasting blood glucose levels. The polysaccharides fraction of American ginseng berry extract at doses of 50 and 150 mg/kg body wt. decreased fasting blood glucose levels significantly. On Day 5, as compared to the vehicle-treated mice (230.5 ± 13.5 mg/dl), polysaccharides-treated animals had significantly lower fasting blood glucose levels (187.4 ± 20.5 mg/dl in the 50 mg/kg group and 187.4 ± 17.1 mg/dl in the 150 mg/kg group; both $P < 0.05$). On Day 10, as compared to the vehicle group (240.1 ± 12.3 mg/dl), the 50 mg/kg group were 188.4 ± 12.6 mg/dl ($P < 0.05$), and the 150 mg/kg group were normoglycemic (148.8 ± 17.6 mg/dl, $P < 0.01$). However, those *ob/ob* mice treated with vehicle did not show significant changes in fasting blood glucose levels.

Effects of the polysaccharides fraction on the glucose tolerance test

Glucose tolerance was evaluated using the IPGTT, prior to and 10 days after treatment with the polysaccharides fraction. As shown in Fig. 2A (150 mg/kg




Fig. 1. Effect of the polysaccharides fraction from American ginseng berry extract on fasting blood glucose levels in diabetic *ob/ob* mice. The glucose levels decrease significantly in 50 mg/kg body wt. and 150 mg/kg body wt. polysaccharides-treated mice on Day 5 (both $P < 0.05$ as compared to vehicle-treated mice) and Day 10 ($P < 0.05$ and $P < 0.01$ as compared to vehicle-treated mice, respectively).

body wt. group) and 2B (50 mg/kg body wt. group), on Day 0, *ob/ob* mice demonstrated basal hyperglycemia, and this hyperglycemia was exacerbated by the IP glucose load, and failed to return to baseline after 120 min indicating glucose intolerance. After 10-day treatment with polysaccharides at 150 and 50 mg/kg body wt., overall glucose tolerance improved remarkably. After 150 and 50 mg/kg body wt. polysaccharides treatment, the area under the curve (AUC) decreased 28.2% (from 287.8 mg/ml · min of Day 0 to 206.7 mg/ml · min of Day 10, $P < 0.01$) and 15.5% (from 249.6 mg/ml · min of Day 0 to 211.0 mg/ml · min of Day 10, $P < 0.05$), respectively.




Fig. 2. Intraperitoneal glucose tolerance test in *ob/ob* mice before (Day 0) and after a 10-day treatment with polysaccharides fraction from American ginseng berry extract. (A) 150 mg/kg body wt. polysaccharides-treated mice ($n = 5$). (B) 50 mg/kg body wt. polysaccharides-treated mice ($n = 5$). There is a significantly higher rate of glucose disposal based on measurement of area under the plasma concentration curve (AUC) ($P < 0.01$ and $P < 0.05$ compared to Day 0, respectively).

Prolonged effect of the polysaccharides fraction on fasting blood glucose levels

To observe whether there was an effect on fasting blood glucose concentration after cessation of polysaccharides treatment, we measured the blood glucose




Fig. 3. Prolonged hypoglycemic effect on fasting blood glucose levels in *ob/ob* mice treated with 150 and 50 mg/kg body wt. polysaccharides fraction from American ginseng berry extract. After 5 days (Day 15), 10 days (Day 20), and 15 days (Day 25) cessation of treatment, fasting blood glucose levels were significantly lower as compared to the vehicle group (all $P < 0.01$).




Fig. 4. Effects of polysaccharides fraction on body weight changes in diabetic *ob/ob* mice. While there is a tendency to increase in body weight from Day 0 to Day 10 in mice receiving vehicle, polysaccharides treatment did not affect body weight changes.

every five days until the levels returned to those measured prior to the treatment. Fig. 3 shows a prolonged effect in animals who received polysaccharides treatment. Fasting blood glucose levels were 168.6 ± 17.7 (150 mg/kg) and 155.6 ± 7.4 (50 mg/kg) on Day 15, 176.8 ± 15.4 (150 mg/kg), 163.8 ± 15.7 (50 mg/kg) on Day 20, and 185.6 ± 7.9 (150 mg/kg) and 174.8 ± 10.4 (50 mg/kg) on Day 25, respectively (all $P < 0.01$ compared to the vehicle group).

Effects of the polysaccharides fraction on body weight changes

As shown in Fig. 4, the body weight of *ob/ob* mice in the vehicle-treated group had a tendency to increase from Day 0 to Day 10. This tendency of body weight increase was not affected by 150 or 50 mg/kg body wt. polysaccharides administration. After cessation of drug injection, a prolonged hypoglycemic effect was observed, as presented in Fig. 3. However, in our 30 days observation, body weight changes were not affected by polysaccharides treatment (Fig. 5).

Discussion

Type-2 diabetes is a major health problem, affecting approximately 5% of the total population of the U.S., and 3% of the population world-wide. Currently available drugs for type-2 diabetes have a number of limitations such as adverse effects and high rates of secondary failure (DeFronzo, 1999). These problems have led to the search for alternative therapies that may have




Fig. 5. Effect of polysaccharides on body weight changes in diabetic *ob/ob* mice. As compared to the vehicle group, polysaccharides administration did not affect body weight changes during and after treatment.

a similar degree of efficacy without the troublesome side-effects associated with the conventional drug treatment. The identification of compounds from medicinal plants with anti-hyperglycemic activity may also provide an opportunity to develop a new class of anti-diabetic agent.

In this study, we observed that the polysaccharides fraction of American ginseng berry extract had a significant anti-hyperglycemic effect. Glucose tolerance also improved significantly after polysaccharides treatment. This was shown by overall glucose excursion, calculated as area under the curve during the two-hour intraperitoneal glucose tolerance test, which decreased up to 28% as compared to pre-treatment levels. Whether this improvement of glucose tolerance is due to an increase in insulin sensitivity remains to be tested in future experiments using body-wide hyperinsulinemic-euglycemic clamp technique (DeFronzo, 1999).

Previously, we observed that extract of ginseng berry had significant anti-hyperglycemic and anti-obese effects in *ob/ob* and *db/db* mice (Attele et al. 2002; Xie et al. 2002b). However, unlike ginseng berry extract, the polysaccharides fraction of the extract did not affect body weight in the *ob/ob* mice. Thus, the anti-hyperglycemic action of the polysaccharides fraction is independent of body weight changes. This suggests that other constituents in the berry extract have distinct pharmacological mechanisms affecting energy metabolism.

Past anti-diabetic activity data suggested that ginseng root may affect glucose transport, which is mediated by nitric oxide (NO), and thus modulate NO-mediated insulin secretion (Hasegawa et al. 1994; Ohnishi et al. 1996; Roy et al. 1998). It was shown recently that NO stimulates glucose-dependent secretion of insulin in rat islet cells (Spinass et al. 1998). In these studies, however, the investigators did not test which fraction of ginseng extract, for example, ginsenoside or polysaccharide fraction, are responsible for the observed effects.

Another possible action site for ginseng berry extract to contribute its hypoglycemic effect is in the gastrointestinal tract. Gastric vagal afferents are the primary neuroanatomical link between the stomach and the central nervous system. In a previous study, we reported that ginseng extract, via gastric vagal afferents, inhibited brainstem neuronal activity (Yuan et al. 1998). Suzuki et al. (1991) reported that gastric secretion *in vitro* was inhibited by ginseng (Suzuki et al. 1991). These results suggest that ginseng may slow the digestion of food and decrease the rate of carbohydrate absorption. Since we observed a significant prolonged hypoglycemic effect after cessation of polysaccharides fraction treatment in this study, it is unlikely that the delay in digestion after the treatment would last several weeks.

Yang et al. reported that intraperitoneal or subcutaneous injection of polysaccharides from ginseng root, at dose of 50–200 mg/kg body wt., reduced blood glucose and liver glycogen levels significantly in mice (Yang et al. 1990; Yang and Wang, 1991). These investigators proposed that the reduction of blood glucose and liver glycogen by polysaccharides may be due primarily to the increase of carbohydrate utilization and the decrease of glycogenesis. They also observed that the polysaccharides stimulated the release of insulin. The exact mechanisms of the polysaccharides fraction from American ginseng berry extract in *ob/ob* mice are yet to be determined.

Various components from ginseng polysaccharides have been reported. For example, heteroglycans composed of six monosaccharides (Liu et al. 1988) and a number of biologically active panaxans were isolated (Konno and Hikino, 1987; Konno et al. 1984; 1985) in alloxan-induced type-1 diabetic mice. Whether these components have a similar effect on type-2 diabetic mice is still unknown. The activity-guided polysaccharides fractionation approach will be useful in our future studies to identify the particular constituents responsible for the pharmacological effect observed in this study.

The present study demonstrated that the polysaccharides fraction from American ginseng berry extract has a significant anti-hyperglycemic activity. Our observation suggested that the polysaccharides may have a potential clinical utility in treating type-2 diabetic patients. On the other hand, due to a prolonged effect of the polysaccharides on fasting blood glucose levels observed in this study, the pharmacokinetics of the compound warrant future investigation. Caution should be taken due to the undesirable additive hypoglycemic effect occurring when other anti-hyperglycemic drugs are added after cessation of polysaccharides administration.

Acknowledgement

This work was supported in part by the Tang Family Foundation, NIH/NCCAM grants AT00381 and AT00563, and an NIH Institutional Diabetes Research & Training Center grant. The authors are grateful for the assistance of the Ginseng Board of Wisconsin and Ms. Joan Eckes.

References

- Ackerknecht EHA (1982) Short History of Medicine, Johns Hopkins University Press, Baltimore/London
- Attele AS, Wu JA, Yuan CS (1999) Multiple pharmacological effects of ginseng. *Biochem Pharmacol* 58: 1685–1693
- Attele AS, Zhou YP, Xie JT, Wu JA, Zhang L, Dey L, Pugh W, Paul AR, Polonsky KS, Yuan CS (2002) Antidiabetic effects of *Panax ginseng* berry extract and the identification of an effective component. *Diabetes* 51: 1851–1858

- Bensky D, Gamble A (1993) Chinese Herbal Medicine Materia Medica, Eastland Press, Seattle, WA
- Cao LY (1989) Progress of chemical studies of polysaccharides from *Panax ginseng*. Chinese Traditional and Herbal Drugs 20: 36–41
- DeFronzo RA (1999) Pharmacologic therapy for type 2 diabetes mellitus. Ann Intern Med 131: 281–303
- Hao XH, Li PY, Li X, Liu YQ (2000) Saponins from the fruit of *Panax quinquefolius* collected in Canada. Chinese Traditional and Herbal Drugs 31: 801–803
- Hasegawa H, Matsumiya S, Murakami C (1994) Interaction of ginseng extract, ginseng separated fractions, and some triterpenoid saponins with glucose transporters in sheep erythrocytes. Planta Med 60: 153–157
- Huang KC (1999) The Pharmacology of Chinese Herbs. CRC Press, Boca Raton, FL
- Kim JK, Gavrilova O, Chen Y, Retiman ML, Shulman GL (2000) Mechanism of insulin resistance in A-2JP/F-1 fatless mice. J Biol Chem 275: 8456–8460
- Kimura I, Nakashima N, Sugihara Y, Fu-Jun C, Kimura M (1999) The antihyperglycemic blend effect of traditional Chinese medicine Byakko-ka-ninjin-to on alloxan and diabetic KK-CA^y mice. Phytother Res 13: 484–488
- Kimura M (1980) Hypoglycemic component in ginseng radix and its insulin release. Proceedings of the 3rd International Ginseng Symposium. Korean Ginseng Research Institute, Seoul
- Konno C, Sugiyama K, Kano M, Takahashi M, Hikino H (1984) Isolation and hypoglycemic activity of panaxans A, B, C D, and E: Glycans of *Panax ginseng* roots. Planta Med 50: 434–436
- Konno C, Mrakami Y, Oshima Y, Hikino H (1985) Isolation and hypoglycemic activity of panaxans Q, R, S, T and U glycans of *Panax ginseng* roots. J Ethnopharmacol 14: 69–74
- Konno C, Hikino H (1987) Isolation and hypoglycemic activity of panaxans M, N, O and P, glycans of *Panax ginseng* roots. Int J Crude Drug Res 25: 53–56
- Lee FC (1992) Facts about Ginseng, the Elixir of Life. Hollyn International Corp, Elizabeth, NJ
- Li RQ, Zhang YS (1986) Structural studies of *Panax Ginseng* C. A. Mey pectin. Acta Pharmaceutica Sinica 21: 912–916
- Liu CL, Zhang YS, Li RQ (1988) Studies on the water soluble polysaccharides from *Panax ginseng* fruit: Isolation, purification and structural studies on heteroglycan F. Acta Pharmaceutica Sinica 23: 863–867
- Ma XL, Hao CY, Lu SX, Sun YX, Lu JZ, Liu SY (1998) Isolation and characterization of a bioactive polysaccharide from *Panax quinquefolium* L. Chemical Research in Chinese Universities 14: 143–146
- Ohnishi Y, Takagi S, Miura T (1996) Effect of ginseng radix on GLUT2 protein content in mouse liver in normal and epinephrine-induced hyperglycemic mice. Biol Pharm Bull 19: 1238–1240
- Roy D, Perrault M, Marette A (1998) Insulin stimulation of glucose uptake in skeletal muscle and adipose tissue in vivo is NO dependent. Am J Physiol 274: E692–E699
- Shafir E (1992) Animal models of non-insulin-dependent diabetes. Diabetes-Metabolism Reviews 8: 180–208
- Sotaniemi EA, Haapakoski E, Rautio A (1995) Ginseng therapy in non-insulin-dependent diabetic patients. Diabetes Care 18: 1373–1375
- Spinas GA, Laffranchi R, Francoys I, David I, Richter C, Reinecke M (1998) The early phase of glucose-stimulated insulin secretion requires nitric oxide. Diabetologia 41: 292–299
- Sung JH, Chae HJ, Kim EY (2000) Development of separation method of ginseng saponins using ion exchange resin. Chromatography 21: 203–208
- Suzuki Y, Ito Y, Konno C, Furuya T (1991) Effects of tissue cultured of ginseng on gastric secretion and pepsin activity. (In Japanese) Yakugaku Zasshi 111: 770–774
- Vuksan V, Sievenpiper JL, Koo VYY, Francis T, Beljan-Zdravkovic U, Xu Z, Vidgen E (2000a) American Ginseng (*Panax quinquefolius* L) reduces postprandial glycemia in nondiabetic subjects and subjects with type II diabetes mellitus. Arch. Intern Med 160: 1009–1013
- Vuksan V, Leiter LA, Stavro MP, Josse RG, Sievenpiper JL, Xu Z, Beljan-Zdravkovic U, Vidgen E (2000b) Similar postprandial glycemic reductions with escalation of dose and administration time of American Ginseng in type II diabetes. Diabetes Care 23: 1221–1226
- Wang LJ, Li PY, Zhao CF, Li, X (2000) Studies on the chemical constituents in the fruit of *Panax quinquefolius*. Chinese Traditional and Herbal Drugs 31: 723–724
- Wang ZG, Lei HP (1957) Effect of ginseng in normal and alloxan-induced diabetic dogs. Chinese Medical Journal 11: 861–865
- Xie JT, Aung HH, Wu JA, Attele AS, Yuan CS (2002a) Effects of American ginseng berry extract on blood glucose levels in *ob/ob* mice. Am J Chin Med 30: 187–194
- Xie JT, Zhou YP, Dey L, Attele AS, Wu JA, Gu M, Polonnsky KS, Yuan CS (2002b) Ginseng berry reduces blood glucose and body weight in *db/db* mice. Phytomedicine 9: 254–258
- Yang M, Wang BX (1991) Effects of the ginseng polysaccharides on reducing liver glycogen. Traditional Medicine and Material Medica 12: 272–275
- Yang M, Wang BX, Jin YL, Wang Y, Cui ZY (1990) Effects of ginseng polysaccharides on reducing blood glucose and liver glycogen. Traditional Medicine and Material Medica 11: 520–524
- Yokozawa T, Kobayashi T, Oura H, Kawashima Y (1985) Studies on the mechanism of the hypoglycemic activity of ginsenoside-Rb₂ in streptozotocin-diabetic rats. Chem Pharm Bull 33: 869–872
- Yuan CS, Wu JA, Lowell T, Gu M (1998) Gut and brain effects of American ginseng root on brainstem neuronal activities in rats. Amer. J Chin Med 26: 47–55
- Zhang SC, Jiang XL (1981) The anti-stress effect of saponins extracted from *Panax ginseng* fruit and the hypophyseal-adrenal system. Acta Pharmaceutica Sinica 16: 860–863

Address

Chun-Su Yuan, Department of Anesthesia & Critical Care, The University of Chicago Medical Center, 5841 S Maryland Avenue, MC 4028, Chicago, Illinois 60637, USA
Tel.: (773) 702-1916; Fax: (773)834-0601;
e-mail: cyuan@airway.uchicago.edu