Digestion Pattern of Antihypertensive Angiotensin I-Converting Enzyme Inhibitory Peptides from *Saccharomyces cerevisiae* in a Successive Simulated Gastric-intestinal Bioreactor

Jeong-Hoon Jang, Seung-Chan Jeong, Jung-Kee Lee and Jong-Soo Lee*

Department of Life Science and Genetic Engineering, Paichai University, Daejeon 302-735, Korea (Received September 30, 2010. Accepted October 27, 2010)

A cell-free extract of *Saccharomyces cerevisiae* containing the angiotensin I-converting enzyme (ACE) inhibitory peptide was treated in a successive simulated gastric-intestinal bioreactor (step 1: amylase digestion, step 2: gastric fluid digestion, step 3: intestinal fluid digestion) to illustrate the absorption pattern of antihypertensive ACE inhibitory peptide, and the ACE inhibitory activities of each step were determined. Total ACE inhibitory activities of step 1, step 2, and step 3 were 55.96%, 80.09%, and 76.77%, respectively. The peptide sequence of each steps was analyzed by MS/MS spectrophotometry. Eleven kinds of representative peptide sequences were conserved in each step, and representative new peptides including RLPTES-VPEPK were identified in step 3.

KEYWORDS: Angiotensin I-converting enzyme inhibitory peptides, Digestion pattern, Simulated gastric-intestinal bioreactor

Angiotensin I-converting enzyme (dipeptidyl carboxy peptidase I, kinase II, E.C 3.4.15.1, ACE) is the key enzyme in the rennin-angiotensin system [1], which catalyzes production of the active hypertensive hormone angiotensin II (Asp-Arg-Val-Tyr-Ile-His-Pro-Phe) from the inactive prohormone angiotensin I (Asp-Arg-Val-Tyr-Ile-His-Pro-Phe-His-Leu) and also inactivates the hypotensive peptide bradykinin [2, 3]. This conversion endows ACE with a very important role in regulating blood pressure through the direct action of angiotensin II on blood vessels, sympathetic nerves, and adrenal glands [2, 3]. ACE inhibitors have long been very useful because they inhibit both ACE in the rennin-angiotensin system and the kallikrein-kinin system [4]. Although some ACE inhibitors have been produced and characterized from several natural sources [5], few are used commercially because of their low antihypertensive action and poor effectiveness in the gastric and intestinal systems. In a previous study [6], we purified and characterized an ACE inhibitor from Saccharomyces cerevisiae. The present study describes the absorption pattern of a novel ACE inhibitory peptide from alcohol fermented S. cerevisiae.

The purified ACE inhibitor from *S. cerevisiae* characterized in a previous paper with an IC_{∞} of 0.07 mg [6] was used in this study. ACE was extracted from rabbit lung acetone powder with 100 mM sodium borate buffer (pH 8.3) containing 300 mM NaCl. The extract was maintained overnight at 4°C and its activity was determined using hippuric acid-histidine-leucine (Hip-His-Leu) as a substrate [1]. The rabbit lung acetone powder, pepsin (4,150 unit/

mg), trypsin (1,200 units/mg), protease N (185 units/mg), and the Hip-His-Leu were obtained from Sigma Chemical Co. (St. Louis, MO, USA). Simulated gastric and intestinal fluid solutions were prepared according to United States Pharmacopeia specifications [7] as follows. For the simulated gastric fluid (SGF), 0.2% sodium chloride and 0.32% pepsin (from porcine stomach mucosa) were dissolved in 7.0 mL hydrochloric acid and a sufficient volume of water to make 1 L (pH 1.2). The purified peptide (1 mg/mL) was added to the SGF and incubated in a shaking incubator (37°C, 120 rpm) for 2 hr. Digestion was stopped by raising the pH to 7.5 using 1 N NaOH. For the simulated intestinal fluid (SIF), 0.68% monobasic potassium phosphate was dissolved in 250 mL water. To this, 77 mL of 0.2 N sodium hydroxide and 500 mL water were added and mixed along with 1% pancreatin (from porcine mucosa) (pH 6.8). The SIF solution was adjusted to pH 6.8 with 0.2 N HCl and then diluted with water to 1 L.

The SIF was added to the digests obtained from the SGF reaction and incubated for 2 hr. Then, these solutions were incubated in a shaking incubator (37°C, 120 rpm) for 10 hr. The final SGF and SIF digested samples were lyophilized immediately and used for measuring ACE inhibitory activity (IC₅₀). The stability of the purified peptide in SGF and SIF was analyzed by reverse phase high performance liquid chromatography. A simulated gastricintestinal (SGI) bioreactor was designed using the gastric and intestinal fluids as shown in Fig. 1.

ACE inhibitory activity was assayed by the modified method of Cushman and Cheung [8]. A mixture containing 100 mM sodium borate buffer (pH 8.3), 300 mM NaCl, 3 units of ACE from rabbit lung, and an appropri-

^{*}Corresponding author <E-mail: biotech8@pcu.ac.kr>

58 Jang et al.




Fig. 1. Flow chart of digestion in the simulated gastric-intestinal bioreactor.

Table 1. Angiotensin I-converting enzyme (ACE) inhibitory activity of the purified ACE inhibitor after digestion in a simulated gastric-intestinal bioreactor

Steps	Control	Step 1	Step 2	Step 3
ACE inhibitory activity (%)	67.6 ± 0.46	56.0 ± 0.66	80.1 ± 0.42	76.8 ± 0.19

Control, without treatment; Step 1, mouth digestion; Step 2, stomach digestion; Step 3, small intestinal digestion.

ate amount of the inhibitor solution was preincubated for 10 min at 37°C. The reaction was initiated by adding 50 μL of Hip-His-Leu at a final concentration of 5 mM and was terminated after a 30-min incubation by adding 250 µL of 1.0 M HCl. The hippuric acid liberated was extracted with 1 mL of ethyl acetate, and 0.8 mL of the extract was evaporated with a Speed Vac concentrator (Eyela Co., Tokyo, Japan). Then, the residue was dissolved in 1 mL of sodium borate buffer. Absorbance at 228 nm was measured to estimate ACE inhibitory activity. The ACE inhibitor concentration required to inhibit 50% of the ACE activity under the assay conditions was defined as the IC_{so}. The amino acid composition of the ACE inhibitor from S. cerevisiae was analyzed with a fluorometric analysis system (SLM-Aminoco, Urbana, IL, USA) after hydrolysis for 24 hr in 4 N methanesulfonic acid containing 0.2% 3-(2-aminoethyl) indole at 110°C [9]. The amino acid sequence was determined by the Edman method [4], using an Applied Biosystems 491A automatic protein sequencer (Applied Biosystems, Foster City, CA, USA) [10].

We determined the inhibitory activity of the purified ACE inhibitor from *S. cerevisiae* after a digestion with gastric and intestinal fluids in the SGI bioreactor (Table 1). The original ACE inhibitory activity (67.6%) of the purified ACE inhibitor decreased to 56.0% after mouth digestion (step 1) and then increased to 80.1% after stomach digestion (step 2) and again decreased to 76.8% after digestion by small intestinal fluids (step 3). These changes in inhibitory activity suggest that some ACE inhibitory

Table 2. Peptides sequence of the control (non-treatment), step 1 (mouth digestion), step 2 (stomach digestion), and step 3 (small intestine digestion) hydrolysates

step 3 (small intestine digestion) hydrolysates							
Peptide sequences							
Control	DDELDISWGIFTPLLK						
	EIKLQILLLEIIGLNDLDWNFR						
	SFYMILPYM*VADLSGVLHNPR						
	GQLGEM*AKC#LDNPDQGISDMCR						
	PYIAVGTSITKGEDIAVR						
	KYSESPWLLVDEAQLVISYAK						
	SC#LLPNVSSKMAVMM*AGLLGSLVIR						
	VGLVQYHPTAANVLASSSADNTIK						
	ITVEKALPQLEMLGGLVACTEK						
	LSGNPILGDGKSDNQNMSIPYIR						
	DDELDISWGIFTPLLK						
Step 1	LDVNTSVDEFFQGCFLNKLC#ISDK						
	LNGLQMPPIQTM*VRWGHQYAPR						
	YIAM*DC#EFVGVGPEGKESALAR						
	FGSQEEGLSLALSHYFDHNSGTSISK						
	GQLGEM*AKC#LDNPDQGISDMCR						
	FLVNAC#M*NPVTAILDC#VNGEMK						
	TDDDEDNSHLLVHHIPTIIRK						
	QLSMM*LM*LAQSNPQLFALMGTR						
	LCDFGVSGNLVASISK						
	SVSKDANIMVVINAAHCIQAM*AK						
Step 2	LDVNTSVDEFFQGCFLNKLC#ISDK MLTGSQIVTM*EPENERTM*CSSAHMK						
Step 2	ILKTELTTIGNHIESQK						
	FKFTPCGFC#FIIYSCPDEALNALK						
	ILC#NLC#NVVPGGM*VCFLPSYEYLR						
	GQLGEM*AKC#LDNPDQGISDMCR						
	DTHNAYSNDVYFLNLLSLKWYK						
	LVNEVENVLNKEGVLQK						
	LLLEELQRLSGLSHK						
	NPVYLTSQTSSSSDYVFDTSLTK						
	PAHPSTM*DENEIVHEEDVK						
	MLTGSQIVTM*EPENERTM*CSSAHMK						
Step 3	RLPTESVPEPK						
	KETLGNPLK						
	GLEFVRINNYFTTHDLQSFK						
	TRNPLTHSTPK						
	VGLVQYHPTAANVLASSSADNTIK						
	VVVDNQQPQPK						
	PNYLQYRIIGGIVDLYIYVGGTK						
	KYVSELLSNSWLSNLSETQVPYDPK						
	DLAQQFGINLHMFSQLSDM*EFVMK						
C1	LPNCIFLSDEMNHASM*INGIRNSR						
Conserved peptide	GLEFVRINNYFTTHDLQSFK						
sequences ^a	GQLGEM*AKC#LDNPDQGISDMCR LISHGKISSNNVPGHSGNLNHYHR						
sequences	DFTRDVSDLSNTVM*K						
	PNYLQYRIIGGIVDLYIYVGGTK						
	LSLAQYYGLTLEAADLC#LM*NFFQK						
	TNSMSSSGLGSPNRSSPSIICM*PK						
	IHARGPMGLDGLTIYK						
	NPDLGFSDYAAAQLR						
	EDNNDAVNSIMALK						
	GLEFVRINNYFTTHDLQSFK						

^aConserved peptides were present in every step including the control.

Table 3. Angiotensin I-converting enzyme (ACE) inhibitory activity of the 5-kDa cut-off filtrate extracts from the intestinal fluid hydrolysis following C18 solid phase chromatography (step 3)

C18 SPE									
0%	5%	25%	35%	45%	55%	65%	85%	100%	
n.d	4.43 ± 0.86	39.95 ± 0.12	69.93 ± 0.07	74.15 ± 0.37	63.29 ± 0.39	20.29 ± 0.34	9.22 ± 0.40	n.d	

n.d, not detected.

peptides were produced newly or decomposed by digestion with pepsin (stomach) and pancreatin (intestinal). Furthermore, the peptide sequences of the step 1, 2, and 3 hydrolysates were also analyzed by MS/MS spectrophotometry (Table 2). Eleven kinds of peptide sequences were conserved and ten kinds of new peptide sequences including RLPTESVPEPK were produced in step 3. It was assumed that these conserved and new peptides sequences were very important for antihypertensive action. Therefore, further determination of the ACE inhibitory activities of these conserved peptide sequences and new peptide sequences is required.

Several functional peptides are absorbed in the small intestine. Therefore, the step-3 hydrolysates were ultrafiltrated with a 5 kDa cut-off filter, and the filtrates were subjected to C18 solid phase extraction chromatography. As shown in Table 3, 45% of the C18 solid phase extract showed the highest ACE inhibitory activity of 74.2%, and 35% of the C18 solid phase extract also showed high ACE inhibitory activities of 69.9% and 63.3%, respectively. Further purification of the ACE inhibitor in the 45% fraction from the C18 solid phase extraction chromatography and a comparison of its inhibitory activities with the purified ACE inhibitor and the identified peptide sequences from SGI bioreactor are also required.

References

1. Fujita H, Yokoyama K, Yoshikawa M. Classification and

- antihypertensive activity of angiotensin I-converting enzyme inhibitory peptides derived from food proteins. J Food Sci 2000;65:564-9.
- Folkow B, Johansson B, Mellander S. The comparative effects of angiotensin and noradrenaline on consecutive vascular sections. Acta Physiol Scand 1961;53:99-104.
- Ondetti MA, Cushman DW. Enzymes of the rennin-angiotensin system and their inhibitors. Annu Rev Biochem 1982; 51:283-308.
- Ondetti MA, Rubin B, Cushman DW. Design of specific inhibitors of angiotensin converting enzyme: new class of orally active antihypertensive agent. Science 1977;196:441-4.
- Koo KC, Lee DH, Kim JH, Yu HE, Park JS, Lee JS. Production and characterization of antihypertensive angiotensin I-converting enzyme inhibitor from *Pholiota adipose*. J Microbiol Biotechnol 2006;16:757-63.
- Kim JH, Lee DH, Jeong SC, Chung KS, Lee JS. Characterization of antihypertensive angiotensin I-converting enzyme inhibitor from *Saccharomyces cerevisiae*. J Microbiol Biotechnol 2004;14:1318-23.
- Test solutions, United States Pharmacopeia 28/National Formulary 23, 2005, USP-NF [Internet]. Rockville: United States Pharmacoprial Convention [cited 2005 Feb 14]. Available from: http://www.usp.org.
- Cushman DW, Cheung HS. Spectrophotometric assay and properties of angiotensin-converting enzyme of rabbit lung. Biochem Pharmacol 1971;20:1637-48.
- 9. Bae EA, Han MJ, Song MJ, Kim DH. Purification of rotavirus infection-inhibitory protein from *Bifidobacterium breve* K-110. J Microbiol Biotechnol 2002;12:553-6.
- Walker JM. Basic protein and peptide protocols. Totowa: Humana Press; 1994.