

On a three-dimensional tumor-virus compartmental model, and two four-dimensional oncolytic virotherapy models

This Mathematica Notebook is a supplementary material to the paper which has the same title as this document. It contains some of the calculations and illustrations appearing in the paper.

1) Section 2 (in paper): Deterministic model with Logistic growth [Tian2011]

0) Definition of the model [Tian11]:

In[]:=

```

SetDirectory[NotebookDirectory[]];
AppendTo[$Path,Directory];
Clear["Global`*"];
(*Some aliases*)
Format[μv]:=Subscript[μ,v];Format[μy]:=Subscript[μ,y];
parT={β>0,λ>0,δ>0,b>1};
cparT={ μv→0,μy→0,γ→1, K→1};
cnTian={ μv→0, μy→0,K→1,γ→1,λ→0.36,β→0.11,δ→0.44}(*Numerical values of Tian*);

(***** Four dim Deterministic epidemic model with Logistic growth *****)
x1=λ x(1-(x+y)/K)- β x v ;
y1=β x v -μy y z- γ y;
v1=-β x v - μv v z+ b γ y - δ v;

dyn={x1,y1,v1}/.μy→0/.μv→0(*Tian case with K>0*);

Print["
      x'
      (y')=",dyn//FullSimplify//MatrixForm,
      v'

", and the reparametrized dynamics [Tian 2011] are
      x'
      (y')=",
      v'

dyn/.cparT//FullSimplify//MatrixForm]

```

$$\begin{pmatrix} x' \\ (y') \\ v' \end{pmatrix} = \begin{pmatrix} -v x \beta + x \left(1 - \frac{x+y}{K}\right) \lambda \\ v x \beta - y \gamma \\ b y \gamma - v (x \beta + \delta) \end{pmatrix}$$

$$, \text{ and the reparametrized dynamics [Tian 2011] are } \begin{pmatrix} x' \\ (y') \\ v' \end{pmatrix} = \begin{pmatrix} -x (v \beta + (-1 + x + y) \lambda) \\ -y + v x \beta \\ b y - v (x \beta + \delta) \end{pmatrix}$$

Fixed points and analysis of the Stability via Routh Hurwitz:

In[]:=

```

cfp=Solve[Thread[dyn=={0,0,0}],{x,y,v}]/FullSimplify;
fp={x,y,v}/.cfp;
Print[Length[fp]," fixed points, the third is E*="]
fp[[3]]//FullSimplify
(*"Jacobian is"*)
Jac=Grad[dyn,{x,y,v}]/FullSimplify;
J0=Jac/.cfp[[1]];J0//MatrixForm;
Print["J(E_K) is"]
J1=Jac/.cfp[[2]];J1//MatrixForm
Eigenvalues[J1]
R0=b β K/(β K+δ);bcrit=1+δ/(β K);(*Reduce[Join[{R0>1},pars],δ]*)
Print["J(E_*) is"]
Jst=Jac/.cfp[[3]]//FullSimplify;Jst//MatrixForm
Jstcr=Jst/.b->bcrit//FullSimplify;
Print["J(E*)/.b->b0 is",Jstcr//MatrixForm," eigvals are ",Eigenvalues[Jstcr]]

(*Routh Hurwitz conditions for the stability of E***)
pc=Collect[Det[ψ IdentityMatrix[3]-Jst],ψ];
coT=CoefficientList[pc,ψ]/FullSimplify;
Print["a1=",a1=Apart[coT[[3]]], ", a2=",a2=coT[[2]], ", a3=",a3=coT[[1]]]
H2=a1*a2-a3;
Print["H2(b0)=",H2/.b->bcrit//FullSimplify]
Print["Denominator of H2 is ",Denominator[Together[H2]]/.K->1//FullSimplify]
Together[H2//FullSimplify];
φb=Collect[Numerator[Together[H2]]/(δ λ),b]/.K->1//FullSimplify;
Print["Coefficients of φ(b)are:",cofi=CoefficientList[φb,b]/FullSimplify]
(*φb/.b->1//FullSimplify*)
Print["value at crit b is "]
φb/.b->bcrit/.K->1//FullSimplify

```

3 fixed points, the third is E*=

$$\text{Out[]}= \left\{ \frac{\delta}{(-1+b)\beta}, \frac{((-1+b)K\beta-\delta)\delta\lambda}{(-1+b)\beta((-1+b)K\beta\gamma+\delta\lambda)}, \frac{\gamma((-1+b)K\beta-\delta)\lambda}{\beta((-1+b)K\beta\gamma+\delta\lambda)} \right\}$$

J(E_K) is

Out[]//MatrixForm=

$$\begin{pmatrix} -\lambda & -\lambda & -K\beta \\ 0 & -\gamma & K\beta \\ 0 & b\gamma & -K\beta-\delta \end{pmatrix}$$

$$\text{Out[]}= \left\{ \frac{1}{2} \left(-K\beta-\gamma-\delta-\sqrt{(K\beta+\gamma+\delta)^2-4(K\beta\gamma-bK\beta\gamma+\gamma\delta)} \right), \right. \\ \left. \frac{1}{2} \left(-K\beta-\gamma-\delta+\sqrt{(K\beta+\gamma+\delta)^2-4(K\beta\gamma-bK\beta\gamma+\gamma\delta)} \right), -\lambda \right\}$$

J(E_*) is

Out[]//MatrixForm=

$$\begin{pmatrix} \frac{\delta\lambda}{K\beta-bK\beta} & \frac{\delta\lambda}{K\beta-bK\beta} & -\frac{\delta}{-1+b} \\ \frac{\gamma((-1+b)K\beta-\delta)\lambda}{(-1+b)K\beta\gamma+\delta\lambda} & -\gamma & \frac{\delta}{-1+b} \\ \frac{\gamma(K(\beta-b\beta)+\delta)\lambda}{(-1+b)K\beta\gamma+\delta\lambda} & b\gamma & \frac{b\delta}{1-b} \end{pmatrix}$$

$$J(E_*) /. b \rightarrow b_0 \text{ is } \begin{pmatrix} -\lambda & -\lambda & -K\beta \\ 0 & -\gamma & K\beta \\ 0 & \gamma + \frac{\gamma\delta}{K\beta} & -K\beta - \delta \end{pmatrix} \text{ eigvals are } \{0, -K\beta - \gamma - \delta, -\lambda\}$$

$$a_1 = \frac{-\gamma + b\gamma + b\delta}{-1 + b} + \frac{\delta\lambda}{(-1 + b)K\beta}, \quad a_2 = \frac{\delta\lambda \left((-1 + b)K\beta\gamma (K(\beta - b\beta) + (-1 + b)\gamma + \delta + b\delta) + ((-1 + b)^2 K\beta\gamma + b\delta^2)\lambda \right)}{(-1 + b)^2 K\beta (-1 + b)K\beta\gamma + \delta\lambda}$$

$$, \quad a_3 = \gamma\delta \left(1 + \frac{\delta}{K\beta - bK\beta} \right) \lambda$$

$$H_2(b_0) = (K\beta + \gamma + \delta)\lambda (K\beta + \gamma + \delta + \lambda)$$

$$\text{Denominator of } H_2 \text{ is } (-1 + b)^3 \beta^2 (-1 + b)\beta\gamma + \delta\lambda$$

Coefficients of $\phi(b)$ are:

$$\left\{ -\beta\gamma(\gamma + \lambda)(\beta\gamma - \delta\lambda), \beta^3\gamma(\gamma - \delta) + \delta^3\lambda^2 - \beta\gamma\delta\lambda(2\gamma + 3\delta + 2\lambda) + \beta^2\gamma(3\gamma^2 - \delta^2 + 3\gamma(\delta + \lambda)), \right. \\ \left. \beta(\beta^2\gamma(-3\gamma + 2\delta) - 3\beta\gamma^2(\gamma + 2\delta + \lambda) + \delta\lambda(\gamma^2 + \delta^2 + \gamma(3\delta + \lambda))), \right. \\ \left. \beta^2\gamma(3\beta\gamma + \gamma^2 - \beta\delta + 3\gamma\delta + \delta^2 + \gamma\lambda), -\beta^3\gamma^2 \right\}$$

value at crit b is

$$\text{Out[]} = \frac{\delta^3 (\beta + \gamma + \delta) (\gamma + \lambda) (\beta + \gamma + \delta + \lambda)}{\beta}$$

Computations of the Jacobians and Eigenvalues using EcoEvo package:

```
In[ ]:= <<EcoEvo`
(*EcoEvoDocs;*)
(*****Analysis of the Model, K=γ=1***)
dynKeq1=dyn/.cparT;
ClearParameters;
UnsetModel;
SetModel[{Pop[x]→{Equation→dynKeq1[[1]],Color→Red},Pop[y]→
{Equation→(dynKeq1[[2]]),Color→Green},
Pop[v]→{Equation→(dynKeq1[[3]]),Color→Purple},
Parameters→(cp=parT)}]

fpT=SolveEcoEq[]//FullSimplify;

J0T=EcoJacobian[fpT[[1]]]//FullSimplify;
J1T=EcoJacobian[fpT[[2]]]//FullSimplify;
Jst=EcoJacobian[fpT[[3]]]//FullSimplify;
Print["Jac (E0)=",J0T//MatrixForm]
Print["Jac (E1)=",J1T//MatrixForm]
Print["Jac (E*)=",Jst//MatrixForm]
Print["Eigenvalues of E1 are:",eiT=EcoEigenvalues[fpT[[2]]]//FullSimplify]

Print["b0=bs1=",bs1=Apart[Last[Last[Reduce[Join[{eiT[[2]]>0},parT],b]]]]]
```

Out[]:= EcoEvo Package Version 1.6.4 (November 5, 2021)

Christopher A. Klausmeier <christopher.klausmeier@gmail.com>

$$\text{Jac}(E_0) = \begin{pmatrix} \lambda & 0 & 0 \\ 0 & -1 & 0 \\ 0 & b & -\delta \end{pmatrix}$$

$$\text{Jac}(E_1) = \begin{pmatrix} -\lambda & -\lambda & -\beta \\ 0 & -1 & \beta \\ 0 & b & -\beta - \delta \end{pmatrix}$$

$$\text{Jac}(E_*) = \begin{pmatrix} \frac{\delta \lambda}{\beta - b \beta} & \frac{\delta \lambda}{\beta - b \beta} & -\frac{\delta}{-1 + b} \\ \frac{((-1 + b) \beta - \delta) \lambda}{(-1 + b) \beta + \delta \lambda} & -1 & \frac{\delta}{-1 + b} \\ \frac{(\beta - b \beta + \delta) \lambda}{(-1 + b) \beta + \delta \lambda} & b & \frac{b \delta}{1 - b} \end{pmatrix}$$

Eigenvalues of E_1 are:

$$\left\{ \frac{1}{2} \times \left(-1 - \beta - \delta - \sqrt{(1 + \beta + \delta)^2 - 4(\beta - b \beta + \delta)} \right), \frac{1}{2} \times \left(-1 - \beta - \delta + \sqrt{(1 + \beta + \delta)^2 - 4(\beta - b \beta + \delta)} \right), -\lambda \right\}$$

$$b_0 = b_{s1} = 1 + \frac{\delta}{\beta}$$

1) Numerical simulations:

Bifurcation Diagram:


In[]:=

```

ClearParameters;
λ=0.36;β=0.11;δ=0.44;bL=10;
Print["b0=",bs1/N]
fpT/N;
linb0=Line[{{bs1,0},{bs1,10}}];
lib0=Graphics[{Thick,Black,Dotted,linb0}];
px0=Plot[0,{b,0,bL},PlotStyle->{Blue},PlotRange->All,PlotLegends->{"E0 saddle point"}];
px1a=Plot[1,{b,0,bs1},PlotStyle->{Green},PlotRange->All,PlotLegends->{"E1 stable"}];
px1b=Plot[1,{b,bs1,bL},PlotStyle->{Red,Dashed},PlotRange->All,
PlotLegends->{"E1 unstable"}];
pxe1=Plot[{x/.fpT[[3]]},{b,0,bs1},PlotStyle->{Cyan,Dashed},PlotRange->All,
PlotLegends->{"E* unstable"}];
pxe2=Plot[{x/.fpT[[3]]},{b,bs1,bL},PlotStyle->{Purple},PlotRange->All,
PlotLegends->{"E* stable"}];
bif11T=Show[{px0,px1a,px1b,pxe1,pxe2,lib0},Epilog->{Text["b0",Offset[{-8,10},{bs1,0}],
{PointSize[Large],Style[Point[{bs1,0}],Purple]}},
PlotRange->{{0,10},{0,4}},AxesLabel->{"b","xee"}]
Export["bif11T.pdf",bif11T]

```

b0=5.



Out[]:= bif11T.pdf

Periodic x,y values when b=28:

In[]:=

```


ClearParameters;
λ=0.36;β=0.11;δ=0.44;b=28;
in={x→0.5,y→0.5,v→1.5};
fpT//N
EcoEigenvalues[fpT[[3]]] (*Eigenvalues corresponding to E* **)
solE3=EcoSim[RuleListAdd[fpT[[3]],in],20000];

Fig5T=PlotDynamics[{solE3[[1]],solE3[[2]]},PlotRange→{{0,2000},{0,0.7}}]
Export["Fig5T.pdf",Fig5T]

```

Out[]:= $\{\{x \rightarrow 0., y \rightarrow 0., v \rightarrow 0.\}, \{x \rightarrow 1., y \rightarrow 0., v \rightarrow 0.\},$
 $\{x \rightarrow 0.148148, y \rightarrow 0.0431317, v \rightarrow 2.64672\}\}$

Out[]:= $\{-1.51022, 0.000296187 + 0.298909 i, 0.000296187 - 0.298909 i\}$



Out[]:= Fig5T.pdf

Numerical illustrations (Parametric plot) when $b=28$:

```

In[ ]:= ClearParameters;
λ=0.36;β=0.11;δ=0.44;b=28;K=1;γ=1;
Print["E*",fp[[3]]/N]
x1=λ x[t] (1-(x[t]+y[t]))- β x[t]×v[t] ;
y1=β x[t]×v[t] - y[t];
v1=-β x[t]×v[t] + b y[t] - δ v[t];

ode={x'[t]==x1,y'[t]==y1,v'[t]==v1,x[0]==0.5,y[0]==0.5,v[0]==1.5};
sol=NDSolve[ode,{x,y,v},{t,0,400}];


x0=0.5; y0=0.5;v0=1.5;
ppb28=ParametricPlot[{ x[t],(y[t]) }/.sol,{t,0,400}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Blue}];
py=Plot[(y/.y→fp[[3,2]]),{t,0,400},PlotStyle→{Dashed,Green}];
pb28=Show[{ppb28,py, Graphics[{Green,Dashed,
Line[{ {x/.x→fp[[3,1]],0},{x/.x→fp[[3,1]],1}]}]}],
Epilog→{{Text["(x*,y*)",Offset[{10,10},{x/.x→fp[[3,1]],(y/.y→fp[[3,2]])}],
{PointSize[Large],Style[Point[{ {x/.x→fp[[3,1]],(y/.y→fp[[3,2]])}],Black}],
{PointSize[Large],Point[{x0,y0}],Text["(x0,y0)",Offset[{-10,8},{x0,y0}]]}}];
(*****b=28; different initial values *****)
ClearParameters;
λ=0.36;β=0.11;δ=0.44;b=28;K=1;γ=1;


ode={x'[t]==x1,y'[t]==y1,v'[t]==v1,x[0]==0.1,y[0]==0.04,v[0]==0.01};
sol=NDSolve[ode,{x,y,v},{t,0,400}];

(*****Parametric plot conditions***)
x0=0.1; y0=0.04;v0=0.01;
ppb28n=ParametricPlot[{ x[t],(y[t]) }/.sol,{t,0,400}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Red}];
pyn=Plot[(y/.y→fp[[3,2]]),{t,0,200},PlotStyle→{Dashed,Green}];
pb28n=Show[{ppb28n,pyn, Graphics[{Green,Dashed,
Line[{ {x/.x→fp[[3,1]],0},{x/.x→fp[[3,1]],1}]}]}],
Epilog→{{Text["(x*,y*)",Offset[{10,10},{x/.x→fp[[3,1]],(y/.y→fp[[3,2]])}],
{PointSize[Large],Style[Point[{ {x/.x→fp[[3,1]],(y/.y→fp[[3,2]])}],Black}],
{PointSize[Large],Point[{x0,y0}],Text["(x0,y0)",Offset[{-10,8},{x0,y0}]]}}];
cy11=Show[{pb28,pb28n},Epilog→{Text["(x*,y*)",
Offset[{10,10},{x/.x→fp[[3,1]],(y/.y→fp[[3,2]])}],
{PointSize[Large],Style[Point[{ {x/.x→fp[[3,1]],(y/.y→fp[[3,2]])}],Black}],
{PointSize[Large],Point[{0.1,0.04}],Text["(x0',y0')",Offset[{-10,8},{0.1,0.04}]]},
{PointSize[Large],Point[{0.5,0.5}],Text["(x0,y0)",Offset[{-10,8},{0.5,0.5}]]}}];
Export["cy11.pdf",cy11]

```

E*{0.148148, 0.0431317, 2.64672}





Out[]= cy11.pdf

2) Sections 3 and 4 (in paper): Deterministic model with Logistic growth (4 dim when $\epsilon=0$)

```
In[ ]:= SetDirectory[NotebookDirectory[]];
AppendTo[$Path, Directory];
Clear["Global`*"];
(*Some aliases*)
Format[μv] := Subscript[μ, v]; Format[μy] := Subscript[μ, y];
parE = {β > 0, λ > 0, γ > 0, δ > 0, μ > 0, μv > 0, b > 1, K > 0, s > 0, c > 0};
cKga1 = {ε → 0, K → 1, γ → 1};
cE = {ε → 1};
R0 = b β K / (β K + δ) (* Reproduction number*);
cnT17 = {μv → 0.16, μy → 0.48, K → 1, γ → 1, b → 9, β → 0.11,
λ → 0.36, δ → 0.2, s → 0.6, c → 0.036} (*Numerical values of Tian17*);
```

2-1) Description of the model when $\epsilon=0$ and analysis of the stability of the fixed point when $z \rightarrow 0$

```

In[ ]:= (***** Four dim Deterministic epidemic model with Logistic growth *****)
x1= $\lambda x(1-(x+y)/K) - \beta x v$  ;
y1= $\beta x v - \mu y y z - \gamma y$ ;
v1= $-\beta x v - \mu v v z + b \gamma y - \delta v$ ;
z1= $z(s y - c)$ ;
ye=c/s; vM= $\lambda(1-ye)/\beta$ ; vMN=vM/.cnT17;
dyn={x1,y1,v1,z1};
dyn3={x1,y1,v1}/.z->0; (*3dim case*)

Print[" $\begin{pmatrix} x' \\ y' \\ v' \\ z' \end{pmatrix} = "$ , dyn//FullSimplify//MatrixForm, " the reparametrized model is  $\begin{pmatrix} x' \\ y' \\ v' \\ z' \end{pmatrix} = "$ ,
dyn/.ckga1//FullSimplify//MatrixForm]
Print["b0=", b0=b/.Apart[Solve[R0==1,b][[1]]//FullSimplify]]
(*****Fixed points of Tian17 using the elimination when K=1,  $\gamma=1$ ****)

fv=(ye (b-1)-v  $\delta$ ); gv=(ye  $\mu y + v \mu v$ ); hv=(1-ye-v  $\beta/\lambda$ );

Print["xe= ",xe=hv," ze =", ze=fv/gv]

Pv=Numerator[Together[v  $\beta$  xe-ye (1+ $\mu y$  fv/gv)]]/(-s2  $\beta^2 \mu v$ );
Print["P(v)=",pc=Collect[Together[Pv],v]," coefs are"]
coP=CoefficientList[pc,v]//Simplify

(***Fixed point when z->0***)
eq=Thread[dyn3=={0,0,0}];
sol=Solve[eq,{x,y,v}]/FullSimplify;
Es={x,y,v}/.sol[[3]]; (*Endemic point with z=0*)
Print["when K= $\gamma=1$ , E*=",Est=Es/.ckga1//FullSimplify(* E* when K= $\gamma=1$ ****)]

bn=b/.Solve[Est[[2]]==ye,b]; bnn=bn/.cnT17;
bcn=b/.Solve[Est[[2]][[1]]==ye,b];
bcnn=bn/.cnT17;

Dis=Chop[Collect[Discriminant[Numerator[Pv],v],b]];
solb=Solve[Dis==0,b];
Jac=Grad[dyn/.ckga1,{x,y,v,z}]/FullSimplify;

```

$$\begin{pmatrix} x' \\ y' \\ v' \\ z' \end{pmatrix} = \begin{pmatrix} -v x \beta + x \left(1 - \frac{x+y}{K}\right) \lambda \\ v x \beta - y (\gamma + z \mu_y) \\ b y \gamma - v (x \beta + \delta + z \mu_v) \\ (-c + s y) z \end{pmatrix} \quad \text{the reparametrized model is} \quad \begin{pmatrix} x' \\ y' \\ v' \\ z' \end{pmatrix} = \begin{pmatrix} -x (v \beta + (-1 + x + y) \lambda) \\ v x \beta - y (1 + z \mu_y) \\ b y - v (x \beta + \delta + z \mu_v) \\ (-c + s y) z \end{pmatrix}$$

$$b\theta = 1 + \frac{\delta}{K \beta}$$

$$x_e = 1 - \frac{c}{s} - \frac{v \beta}{\lambda} \quad z_e = \frac{\frac{(-1+b)c}{s} - v \delta}{v \mu_v + \frac{c \mu_y}{s}}$$

$$P(v) = v^3 + \frac{b c^2 \lambda \mu_y}{s^2 \beta^2 \mu_v} + \frac{v^2 (c s \beta \lambda \mu_v - s^2 \beta \lambda \mu_v + c s \beta^2 \mu_y)}{s^2 \beta^2 \mu_v} + \frac{v (c s \lambda \mu_v + c^2 \beta \lambda \mu_y - c s \beta \lambda \mu_y - c s \delta \lambda \mu_y)}{s^2 \beta^2 \mu_v} \quad \text{coefs are}$$

$$Out[*] = \left\{ \frac{b c^2 \lambda \mu_y}{s^2 \beta^2 \mu_v}, \frac{c \lambda (c \beta \mu_y + s (\mu_v - (\beta + \delta) \mu_y))}{s^2 \beta^2 \mu_v}, \frac{c \lambda \mu_v - s \lambda \mu_v + c \beta \mu_y}{s \beta \mu_v}, 1 \right\}$$

$$\text{when } K=\gamma=1, E_* = \left\{ \frac{\delta}{(-1+b) \beta}, \frac{((-1+b) \beta - \delta) \delta \lambda}{(-1+b) \beta ((-1+b) \beta + \delta \lambda)}, \frac{((-1+b) \beta - \delta) \lambda}{\beta ((-1+b) \beta + \delta \lambda)} \right\}$$

2-2) Interior equilibrium

Analysis of the stability of the interior point E_* :

```
In[*]:= Jac3=Grad[dyn3/.cKga1,{x,y,v}]/FullSimplify;
bcrit=1+δ/(β);(*Reduce[Join[{R0>1},pars],δ]*)
Print["J(E_*) is"]
Jst=(Jac3/.x→Est[[1]].y→Est[[2]].v→Est[[3]])/FullSimplify;Jst//MatrixForm
Trs=Tr[Jst];
pc=Collect[Det[ψ IdentityMatrix[3]-Jst],ψ];
coT=CoefficientList[pc,ψ]/FullSimplify;
Print["a1=",a1=coT[[3]],",",a2=",a2=coT[[2]],",",a3=",a3=coT[[1]]]
H2=a1*a2-a3;
Print["H2(bθ)=",H2/.b→bcrit/FullSimplify]
Print["Denominator of H2 is ",Denominator[Together[H2]]/.cKga1/FullSimplify]
φb=Collect[Numerator[Together[H2]]/(δ λ),b]/.cKga1;
cofi=CoefficientList[φb,b](*Coefficients of φ(b)*);
Print["value of φ(b) at crit b is "]
φb/.b→bcrit/.cKga1/FullSimplify
```

J(E_*) is

Out[*]//MatrixForm=

$$\begin{pmatrix} \frac{\delta \lambda}{\beta - b \beta} & \frac{\delta \lambda}{\beta - b \beta} & -\frac{\delta}{-1+b} \\ \frac{((-1+b) \beta - \delta) \lambda}{(-1+b) \beta + \delta \lambda} & -1 & \frac{\delta}{-1+b} \\ \frac{(\beta - b \beta + \delta) \lambda}{(-1+b) \beta + \delta \lambda} & b & \frac{b \delta}{1-b} \end{pmatrix}$$

$$a_1 = \frac{\beta(-1+b+b\delta) + \delta\lambda}{(-1+b)\beta}, \quad a_2 = \frac{\delta\lambda((-1+b)\beta(-1+\beta+\delta+b(1-\beta+\delta)) + ((-1+b)^2\beta + b\delta^2)\lambda)}{(-1+b)^2\beta((-1+b)\beta + \delta\lambda)}, \quad a_3 = \delta \left(1 + \frac{\delta}{\beta - b\beta}\right)\lambda$$

$$H2(b0) = (1 + \beta + \delta)\lambda(1 + \beta + \delta + \lambda)$$

$$\text{Denominator of } H2 \text{ is } (-1+b)^3\beta^2((-1+b)\beta + \delta\lambda)$$


value of $\phi(b)$ at crit b is

$$\text{Out}[*]= \frac{\delta^3(1 + \beta + \delta) \times (1 + \lambda) \times (1 + \beta + \delta + \lambda)}{\beta}$$

Numerical values:

```
In[*]:=
cn=cnT17;
cb=NSolve[(ϕb//.Drop[cnT17,{5}])==0,b,WorkingPrecision->20]
bM=Max[Table[Re[b/.cb[[i]]],{i,Length[cb]}]];
Print["bH=",bH=N[bM,30]]
PR=Solve[Pv==0,v];
vn=v/.PR[[2]];vi=v/.PR[[3]];vp=v/.PR[[1]];
Chop[{vn,vi,vp}/.cnT17];
PR=Solve[Pv==0,v];
vn=v/.PR[[2]];vp=v/.PR[[1]];vi=v/.PR[[3]];
Print["b0=",b0=.cn//N," ",b1=", bnn[[1]], ", b2=", bnn[[2]], " ",bH=",bH]
PRN=Chop[PR//.cn//N](*values of the roots v*);
Print["E*",Es=Es/.cn//N]
Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v->vp/.cn//N]]
Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v->vi/.cn//N]]

Jiv=Jac/{x->xe,y->ye,z->ze};
JEi=Jiv/.v->vi//.cn//N//FullSimplify;
JEp=Jiv/.v->vp//.cn//N//FullSimplify;
JEif=Jac/.x->1/.y->0/.z->0/.v->0//.cn//N//FullSimplify;
Print["Eigv of E*:",Append[Eigenvalues[Jst]//.cn//N,Es[[2]]-ye/.cnT17]," Eigv of E+:",
Re[Eigenvalues[JEp]//N], " Eigv of Ei:",Re[Eigenvalues[JEi]//N],
" ", Eigv of Eif:",Eigenvalues[JEif]//N]
```

 **NSolve:** The precision of the argument function $\{-0.0056848 + 0.0319512b - 0.0515086b^2 + 0.0279268b^3 - 0.001331b^4\}$ is less than WorkingPrecision (20).

Out[*]= {{b → 0.29905192792090222818}, {b → 0.83532939126460210381 - 0.23115561298178191540 i}, {b → 0.83532939126460210381 + 0.23115561298178191540 i}, {b → 19.012107471368075382}}

bH=19.012107471368075382

b0=2.81818 , b1=3.58676, b2=8.66779 , bH=19.012107471368075382

E*={0.227273, 0.0584416, 2.33766}

E+={0.249944, 0.06, 2.25837, 0.072607}

Ei={0.483284, 0.06, 1.49471, 0.675711}

Eigv of E*:{-1.25056, -0.0281268 - 0.20904 i, -0.0281268 + 0.20904 i, -0.00155844}

Eigv of E+:{-1.29833, -0.0332218, -0.0332218, 0.000834552}

Eigv of Ei:{-1.69849, -0.076539, -0.076539, -0.00803072}

, Eigv of Eif:{-1.7081, 0.398103, -0.36, -0.036}

3D- Plot of the dynamic:


In[]:=

```
cn=cnT17;
Print["E*",Es=Es/.cn//N]
Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp/.cn//N]]
Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi/.cn//N]]
epi={Text["E*",Offset[{-10,10},{Es[[1]],Es[[2]]}]]//.cn,
{PointSize[Large],Style[Point[{Es[[1]],Es[[2]]}],Blue]//.cn},Text["E_i",Offset[{0,10},
{Ei[[1]],Ei[[2]]}],{PointSize[Large],Style[Point[{Ei[[1]],Ei[[2]]}],Purple}],
Text["E+",Offset[{0,10},{Ep[[1]],Ep[[2]]}],{PointSize[Large],
Style[Point[{Ep[[1]],Ep[[2]]}],Orange]}};
sp3=StreamPlot3D[dyn3//.cnT17,{x,0,1},{y,0,1},{v,0,3},AxesLabel→{"x","y","v"},
StreamColorFunction→Hue,PlotRange→All];
sp3D=Show[{sp3},Graphics3D[Text[Style["E*",Black,Thick],Es//.cn],
{PointSize[0.06],Style[Point[Es],Black]}],Graphics3D[Text[Style["E_i",Black,Thick],
Drop[Ei,{4}]//.cn],{PointSize[0.06],Style[Point[Drop[Ei,{4}]],Black]}]]
Export["sp3D.pdf",sp3D]
```

E*={0.227273, 0.0584416, 2.33766}

E+={0.249944, 0.06, 2.25837, 0.072607}

Ei={0.483284, 0.06, 1.49471, 0.675711}



Out[]:= sp3D.pdf

3)Sections 3 and 4(in paper):Figures used in the manuscript (*Run the previous cell*)

Numerical illustrations when $\epsilon=0$ (Bifurcations diagrams, parametric plots,

and 3D plot)

Bifurcation diagram when b varies:


In[]:=

```
ClearParameters;
μv=0.16; μy=0.48;K=1;b=9;γ=1;λ=0.36;β=0.11;δ=0.2;s=0.6; c=0.036;ε=0;
Print["{b2*,b1*}=",{b/.solb[[1]],b/.solb[[2]]} ,"", and b0=",b0, " ", bH=",bH,
    ",b1=", bnn[[1]], " ", b2=", bnn[[2]]]
Clear["b"];
vs=
$$\frac{((-1+b) \beta - \delta) \lambda}{\beta ((-1+b) \beta + \delta \lambda)}$$
 (*v of E* * ****);
bL=28; max=3.5;
bc2=b/.solb[[1]];
bc1=b/.solb[[2]];
lin1=Line[{{ bc1,0},{ bc1,max}}];
li1=Graphics[{Thick,Black,Dashed,lin1}];
lin2=Line[{{ bc2,0},{ bc2,max}}];
li2=Graphics[{Thick,Black,Dashed,lin2}];
lin3=Line[{{ b0,0},{ b0,max}}];
li3=Graphics[{Thick,Black,Dashed,lin3}];
linH=Line[{{ bH,0},{ bH,max}}];
liH=Graphics[{Thick,Black,Dashed,linH}];
linb1=Line[{{ bnn[[1]],0},{ bnn[[1]],max}}];
lib1=Graphics[{Thick,Black,Dashed,linb1}];
linb2=Line[{{ bnn[[2]],0},{ bnn[[2]],max}}];
lib2=Graphics[{Thick,Black,Dashed,linb2}];
linb9=Line[{{ 9,0},{ 9,max}}];
lib9=Graphics[{Thick,Black,Dashed,linb2}];
pn=Plot[{vn},{b,0,bL},PlotStyle→{Green,Dotted},PlotRange→All,PlotLegends→{"E- negative"}];
p0=Plot[0,{b,0,bL},PlotStyle→{Brown,Thick},PlotRange→All,PlotLegends→{"E0 saddle point"}];
ppa=Plot[{vp},{b,0, bnn[[2]]},PlotStyle→{Purple,Dotted},PlotRange→All,
PlotLegends→{"E+ outside the domain"}];
ppb=Plot[{vp},{b,bnn[[2]], bL},PlotStyle→{Purple,Thick,Dashed},
PlotRange→All,PlotLegends→{"E+ unstable"}];
pi1=Plot[{vi},{b,0,bnn[[1]]},PlotStyle→{Blue,Dotted},
PlotRange→All,PlotLegends→{"Ei outside the domain"}];
pi2=Plot[{vi},{b,bnn[[1]],bL},PlotStyle→{Red,Thick},
PlotRange→All,PlotLegends→{"Ei stable"}];
pi3=Plot[{vi},{b,bnn[[2]],bL},PlotStyle→{Blue,Thick,Dashed},
PlotRange→All,{PlotLegends→{"Ei unstable"}*}];
ps1=Plot[{vs},{b,b0,bnn[[1]]},PlotStyle→{Brown,Thick},
PlotRange→{{0,bL},{0,max}},PlotLegends→{"E+ stable"}];
ps2=Plot[{vs},{b,bnn[[1]],bnn[[2]]},PlotStyle→{Cyan,Thick,Dashed},
PlotRange→{{0,bL},{0,max}},PlotLegends→{"E+ unstable"}];
ps3=Plot[{vs},{b,bnn[[2]],bH},PlotStyle→{Brown,Thick},
PlotRange→{{0,bL},{0,max}}{*,PlotLegends→{"E+ stable"}*}];
ps4=Plot[{vs},{b,bH,bL},PlotStyle→{Cyan,Thick,Dashed},
PlotRange→{{0,bL},{0,max}}{*,PlotLegends→{"E+ unstable"}*}];
pdf1=Plot[{0},{b,0,b0},PlotStyle→{Magenta, Thick},
PlotRange→{{0,200},{0,max}},PlotLegends→{"E1 stable"}];
pdf2=Plot[{0},{b,b0,bL},PlotStyle→{Orange, Thick,Dashed},
PlotRange→{{0,200},{0,max}},PlotLegends→{"E1 unstable"}];
pvmax=Plot[{c (b-1)/(s δ),(λ(1-c/s))/β},{b,0,bL},PlotStyle→{Yellow, Pink},
PlotRange→{{0,200},{0,max}},
```

```
PlotLegends→{"c (b-1)/(s δ)", "(λ(1-c/s))/β"};
```

```
bifT=Show[{pn,ppa,pi1,ppb,pdf1,pdf2,pi2,ps1,ps2,ps3,ps4,li1,li2,li3,
lib2,lib1,lib9,pvmax,liH},
Epilog→{Text["b1*",Offset[{12,10},{bc1,0}]],{PointSize[Large],
Style[Point[{bc1,0}],Purple]},
Text["b2*",Offset[{11,10},{bc2,0}]],{PointSize[Large],Style[Point[{bc2,0}],Yellow]},
Text["b0",Offset[{-7,11},{b0,0}]],{PointSize[Large],Style[Point[{b0,0}],Black]},
Text["bH",Offset[{-10,11},{bH,0}]],{PointSize[Large],Style[Point[{bH,0}],Green]},
Text["b1",Offset[{8,11},{bnn[1],0}]],{PointSize[Large],Style[Point[{bnn[1],0}],Pink]},
Text["b2",Offset[{-7,11},{bnn[2],0}]],{PointSize[Large],Style[Point[{bnn[2],0}],Red]},
Text["b=9",Offset[{7,11},{9,0}]],{PointSize[Large],Style[Point[{9,0}],Brown]}},
PlotRange→All,AxesLabel→{"b","vee"}]
Export["BiifT17.pdf",bifT]
```

$\{b_{2*}, b_{1*}\} = \{10.2462, -0.00697038\}$, and $b_0 =$
 2.81818 , $b_H = 19.012107471368075382$, $b_1 = 3.58676$, $b_2 = 8.66779$



Out[]= BiifT17.pdf

Parametric plots at the intervals of stability:

When $b_0 < b < b_1$:

In[]:=

```

ClearParameters;
 $\mu v = 0.16$ ;  $\mu y = 0.48$ ;  $K = 1$ ;  $b = 3$ ;  $\gamma = 1$ ;  $\lambda = 0.36$ ;  $\beta = 0.11$ ;  $\delta = 0.2$ ;  $s = 0.6$ ;  $c = 0.036$ ;  $\epsilon = 0$ ;

Print["E*", Es = Est /. cn /. N]
Print["E+=", Ep = Chop[{xe, ye, v, ze} /. v -> vp /. cn /. N]
Print["Ei=", Ei = Chop[{xe, ye, v, ze} /. v -> vi /. cn /. N]
 $x_1 = \lambda x[t] (1 - (x[t] + y[t]) / K) - \beta x[t] \times v[t]$ ;
 $y_1 = \beta x[t] \times v[t] - \mu y y[t] \times z[t] - \gamma y[t]$ ;
 $v_1 = -\beta x[t] \times v[t] - \mu v v[t] \times z[t] + b \gamma y[t] - \delta v[t]$ ;
 $z_1 = z[t] (s y[t] - c)$ ;
ode1 = {x'[t] == x1, y'[t] == y1, v'[t] == v1, z'[t] == z1, x[0] == 0.9, y[0] == 0.01, v[0] == 0.01,
z[0] == 0.01};
sol01 = NDSolve[ode1, {x, y, v, z}, {t, 0, 500}];
pdy1 = Plot[{x[t] / 100 /. sol01, y[t] /. sol01, v[t] / 100 /. sol01, z[t] /. sol01},
{t, 0, 600}, PlotLegends -> {"x/100", "y", "v/100", "z"}];
pEs1 = Plot[{x / 100 /. x -> Es[[1]], y /. y -> Es[[2]], v / 100 /. v -> Es[[3]], z /. z -> 0}, {t, 0, 1000},
PlotStyle -> {Dashed}];
Dyn01 = Show[pdy1, pEs1]


(*****Parametric plot conditions****)
x0 = 0.9; y0 = 0.01; v0 = 0.01; z0 = 0.01;
ppb3 = ParametricPlot[{x[t], y[t]} /. sol01, {t, 0, 400}, AxesLabel -> {"x", "y"},
PlotRange -> Full, PlotStyle -> {Blue}];
py3 = Plot[y /. y -> Es[[2]], {t, 0, 400}, PlotStyle -> {Dashed, Green}];
pb3 = Show[{ppb3, py3, Graphics[{Green, Dashed, Line[{x /. x -> Es[[1]], 0}, {x /. x -> Es[[1]], 1}]}]},
Epilog -> {{Thick, Text["(x*, y*)", Offset[{10, -10}, {x /. x -> Es[[1]], y /. y -> Es[[2]]]}],
{PointSize[Large], Style[Point[{x /. x -> Es[[1]], y /. y -> Es[[2]]}], Black]}},
{PointSize[Large], Point[{x0, y0}]}, Text["(x0, y0)", Offset[{10, 10}, {x0, y0}]}]}]
Export["pb3.pdf", pb3]
Export["Dyn01.pdf", Dyn01]

```

E* {0.909091, 0.0224159, 0.224159}

E+ = {0.113348, 0.06, 2.70541, -0.912093}

Ei = {0.726116, 0.06, 0.699984, -0.142026}



Out[]= pb3.pdf

Out[]= Dyn01.pdf

When $b_1 < b = 6 < b_2$:



```
In[ ]:= ClearParameters;
μv=0.16; μy=0.48;K=1;b=6;γ=1;λ=0.36;β=0.11;δ=0.2;s=0.6; c=0.036;ε=0;
Print["E*",Es=Est/.cn//N]
Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp/.cn//N]]
Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi/.cn//N]]

x1=λ x[t] (1-(x[t]+y[t])/K)- β x[t]×v[t] ;
y1=β x[t]×v[t] -μy y[t]×z[t]- γ y[t];
v1=-β x[t]×v[t] - μv v[t]×z[t]+ b γ y[t] - δ v[t];
z1=z[t] (s y[t] - c );
ode2={x'[t]==x1,y'[t]==y1,v'[t]==v1,z'[t]==z1,x[0]==0.9,y[0]==0.01,v[0]==0.01,
z[0]==0.01};
sol12=NDSolve[ode2,{x,y,v,z},{t,0,1000}];
pdy1=Plot[{x[t]/100/. sol12,y[t]/100/. sol12,v[t]/100/. sol12,z[t]/100/. sol12},
{t,0,900},PlotLegends→{"x/100","y/100","v/100","z/100"}];
pEs1=Plot[{x/100/.x→Ei[[1]],y/100/.y→Ei[[2]],v/100/.v→Ei[[3]],z/100/.z→Ei[[4]]},{t,0,900},
PlotStyle→{Dashed}];
Dyn12=Show[pdy1,pEs1]
(*****Parametric plot conditions****)
x0=0.9; y0=0.01;v0=0.01;z0=0.01;
startP=Epilog[{{PointSize[Large],Point[{0.9,0.01}]}},Text["(x0,y0)",
Offset[{0,10},{0.9,0.01}]}];
bnd =Thread[{x[0],y[0],v[0],z[0]}=={x0,y0,v0,z0}]( *Starting point of the Paramateric Plot**);
ppb6=ParametricPlot[{ x[t],(y[t])}/.sol12,{t,0,900}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Blue}];
py6=Plot[{y/.y→Ei[[2]]},{t,0,1000},PlotRange→{{0,0.612},{0,0.08}},PlotStyle→{Dashed,Green}];
pb6=Show[{ppb6,py6, Graphics[{Green,Dashed,Line[{x/.x→Ei[[1]],0},{x/.x→Ei[[1]],1}]}]},
Epilog→{{Text["(xi,yi)",Offset[{10,10},{x/.x→Ei[[1]],(y/.y→Ei[[2]])}],
{PointSize[Large],Style[Point[{x/.x→Ei[[1]],(y/.y→Ei[[2]])}],Black]}},
{PointSize[Large],Point[{0.9,0.01}]}},Text["(x0,y0)",Offset[{-10,8},{0.9,0.01}]}]}]
Export["Dyn12.pdf",Dyn12]
Export["pb6.pdf",pb6]
```

E*{0.363636, 0.0736627, 1.84157}

E+={0.166196, 0.06, 2.53245, -0.475792}

Ei={0.612063, 0.06, 1.07325, 0.425644}



Out[*]= Dyn12.pdf

Out[*]= pb6.pdf

When $b_2 < b = 9 < b_2^*$:

```

ClearParameters;
μv=0.16; μy=0.48;K=1;b=9;γ=1;λ=0.36;β=0.11;δ=0.2;s=0.6; c=0.036;ε=0;
Print["E*=",Es=Est/.cn//N]
Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp/.cn//N]]
Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi/.cn//N]]

x1=λ x[t] (1-(x[t]+y[t])/K)- β x[t]×v[t] ;
y1=β x[t]×v[t] -μy y[t]×z[t]- γ y[t];
v1=-β x[t]×v[t] - μv v[t]×z[t]+ b γ y[t] - δ v[t];
z1=z[t] (s y[t] - c );
ode3={x'[t]==x1,y'[t]==y1,v'[t]==v1,z'[t]==z1,x[0]==0.9,y[0]==0.01,v[0]==0.01,
z[0]==0.01};
sol12=NDSolve[ode3,{x,y,v,z},{t,0,400}];
pdy2=Plot[{x[t]/100/. sol12,y[t]/. sol12,v[t]/100/. sol12,z[t]/. sol12},{t,0,400},
PlotLegends→{"x/100","y","v/100","z"}];
pEs2=Plot[{x/100/.x→Es[[1]],y/.y→Es[[2]],v/100/.v→Es[[3]],z/.z→0},{t,0,1000},
PlotStyle→{Dashed}];
pEi2=Plot[{x/100/.x→Ei[[1]],y/.y→Ei[[2]],v/100/.v→Ei[[3]],z/.z→Ei[[4]]},{t,0,1000},
PlotStyle→{Dashed}];
Dyni22=Show[pdy2,pEi2]
Dyns22=Show[pdy2,pEs2]


(*****Parametric plot conditions****)
x0=0.9; y0=0.01;v0=0.01;z0=0.01;
ppb9=ParametricPlot[{ x[t],(y[t])}/.sol12,{t,0,200}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Blue}];
py9=Plot[(y/.y→Es[[2]])},{t,0,800},PlotStyle→{Dashed,Green}];
pb9=Show[{ppb9,py9, Graphics[{Green,Dashed,Line[{x/.x→Es[[1]],0},{x/.x→Es[[1]],1}]}]},
Epilog→{{Text["(x*,y*)",Offset[{10,10},{x/.x→Es[[1]]},{y/.y→Es[[2]]}]}},{PointSize[Large],
Style[Point[{x/.x→Es[[1]]},{y/.y→Es[[2]]}],Black]}},{PointSize[Large],Point[{0.9,0.01}]},
Text["(x0,y0)",Offset[{-10,8},{0.9,0.01}]}]}]
Export["pb9.pdf",pb9]
Export["Dyni22.pdf",Dyni22]
Export["Dyns22.pdf",Dyns22]



```

E*={0.227273, 0.0584416, 2.33766}

E+={0.249944, 0.06, 2.25837, 0.072607}

Ei={0.483284, 0.06, 1.49471, 0.675711}





$Out[t]=$ pb9.pdf

$Out[t]=$ Dyni22.pdf

$Out[t]=$ Dyns22.pdf

When $b_2 < b = 9 < b_2^*$ and different initial values :

In[]:=

```

ClearParameters;
μv=0.16; μy=0.48;K=1;b=9;γ=1;λ=0.36;β=0.11;δ=0.2;s=0.6; c=0.036;ε=0;
Print["E*=",Es=Est/.cn//N]
Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp/.cn//N]
Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi/.cn//N]

x1=λ x[t] (1-(x[t]+y[t])/K)- β x[t]×v[t] ;
y1=β x[t]×v[t] -μy y[t]×z[t]- γ y[t];
v1=-β x[t]×v[t] - μv v[t]×z[t]+ b γ y[t] - δ v[t];
z1=z[t] (s y[t] - c );
ode3={x'[t]==x1,y'[t]==y1,v'[t]==v1,z'[t]==z1,x[0]==0.5,y[0]==0.01,v[0]==1.2,
z[0]==0.5};
sol22=NDSolve[ode3,{x,y,v,z},{t,0,1000}];
pdy2=Plot[{x[t]/100/. sol22,y[t]/. sol22,z[t]/100/. sol22},{t,0,600},
PlotLegends→{"x/100","y","v/100","z/100"}];
pEs2=Plot[{x/100/.x→Es[[1]],y/.y→Es[[2]],v/100/.v→Es[[3]],z/.z→0},{t,0,1000},
PlotStyle→{Dashed}];
pEi2=Plot[{x/100/.x→Ei[[1]],y/.y→Ei[[2]],v/100/.v→Ei[[3]],z/.z→Ei[[4]]},{t,0,1000},
PlotStyle→{Dashed}];
Dyni22b=Show[pdy2,pEi2]
DyNs22=Show[pdy2,pEs2]


(*****Parametric plot conditions****)
x0=0.5; y0=0.01;v0=1.2;z0=0.5;
ppb9=ParametricPlot[{ x[t],(y[t])}/.sol22,{t,0,500}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Blue}];
py9=Plot[(y/.y→Ei[[2]]),{t,0,800},PlotStyle→{Dashed,Green}];
pb9i=Show[{ppb9,py9, Graphics[{Green,Dashed,Line[{x/.x→Ei[[1]],0},{x/.x→Ei[[1]],1}]}]},
Epilog→{{Text["(x*,y*)",Offset[{10,10},{x/.x→Ei[[1]]},{y/.y→Ei[[2]]}]}},{PointSize[Large],
Style[Point[{x/.x→Ei[[1]]},{y/.y→Ei[[2]]}],Black]}},{PointSize[Large],Point[{0.9,0.01}]},
Text["(x0,y0)",Offset[{-10,8},{0.9,0.01}]}]}]
Export["pb9i.pdf",pb9i]
Export["DyNi22b.pdf",DyNi22b]


```

E*={0.227273, 0.0584416, 2.33766}

E+={0.249944, 0.06, 2.25837, 0.072607}

Ei={0.483284, 0.06, 1.49471, 0.675711}





Out[] = pb9i.pdf

Out[] = Dyni22b.pdf

When $b_2 < b = 10 < b_2^*$:

```
ClearParameters;
μv=0.16; μy=0.48;K=1;b=10;γ=1;λ=0.36;β=0.11;δ=0.2;s=0.6; c=0.036;ε=0;
Print["E*",Es=Est/.cn//N]
Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp/.cn//N]
Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi/.cn//N]

x1=λ x[t] (1-(x[t]+y[t])/K)- β x[t]×v[t] ;
y1=β x[t]×v[t] -μy y[t]×z[t]- γ y[t];
v1=-β x[t]×v[t] - μv v[t]×z[t]+ b γ y[t] - δ v[t];
z1=z[t] (s y[t] - c );

ode3={x'[t]==x1,y'[t]==y1,v'[t]==v1,z'[t]==z1,x[0]==0.9,y[0]==0.01,v[0]==0.01,
z[0]==0.01};
sol22=NDSolve[ode3,{x,y,v,z},{t,0,400}];


(*****Parametric plot conditions****)
x0=0.9; y0=0.01;v0=0.01;z0=0.01;
ppb10=ParametricPlot[{ x[t],(y[t]) }/.sol22,{t,0,200}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Blue}];
py10=Plot[(y/.y→Es[[2]]),{t,0,800},PlotStyle→{Dashed,Green}];
pb10=Show[{ppb10,Epilog→{{Text["(x*,y*)",Offset[{10,10},{(x/.x→Es[[1]]),
(y/.y→Es[[2]])}]}},{PointSize[Large],Style[Point[{(x/.x→Es[[1])},{(y/.y→Es[[2]])}],Black}}},
{PointSize[Large],Point[{0.9,0.01}],Text["(x0,y0)",Offset[{-10,8},{0.9,0.01}]}]}]

Export["pb10.pdf",pb10]
```

$E_* = \{0.20202, 0.0541003, 2.43451\}$

$E_+ = \{0.30876, 0.06, 2.06588, 0.352938\}$

$E_i = \{0.411479, 0.06, 1.72971, 0.635108\}$



Out[]:= pb10.pdf



```
ClearParameters;
μv=0.16; μy=0.48;K=1;b=10;γ=1;λ=0.36;β=0.11;δ=0.2;s=0.6; c=0.036;ε=0;
Print["E*",Es=Est/.cn//N]
Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp/.cn//N]]
Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi/.cn//N]]

x1=λ x[t] (1-(x[t]+y[t])/K)- β x[t]×v[t] ;
y1=β x[t]×v[t] -μy y[t]×z[t]- γ y[t];
v1=-β x[t]×v[t] - μv v[t]×z[t]+ b γ y[t] - δ v[t];
z1=z[t] (s y[t] - c );
ode3={x'[t]==x1,y'[t]==y1,v'[t]==v1,z'[t]==z1,x[0]==0.5,y[0]==0.01,v[0]==1.2,
z[0]==0.5};
sol22=NDSolve[ode3,{x,y,v,z},{t,0,1000}];
x0=0.5; y0=0.01;v0=1.2;z0=0.5;
ppb10i=ParametricPlot[{ x[t],(y[t])}/.sol22,{t,0,1900}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Red}];
py10i=Plot[(y/.y→Ei[[2]]),{t,0,800},PlotStyle→{Dashed,Green}];
pb10i=Show[{ppb10i },Epilog→{{Text["(xi,yi)",Offset[{10,10},{(x/.x→Ei[[1])],
(y/.y→Ei[[2])}]}],
{PointSize[Large],Style[Point[{(x/.x→Ei[[1])],(y/.y→Ei[[2])}],Black]}},
{PointSize[Large],Point[{x0,y0}],Text["(x0,y0)",Offset[{-10,8},{x0,y0}]]}}]
pp10si=Show[{pb10,pb10i},Epilog→{{Text["(xi,yi)",Offset[{10,10},{(x/.x→Ei[[1])],
(y/.y→Ei[[2])}]}],{PointSize[Large],Style[Point[{(x/.x→Ei[[1])],(y/.y→Ei[[2])}],Black]}},
{PointSize[Large],Point[{x0,y0}],Text["(x0i,y0i)",Offset[{-10,8},{x0,y0}]]},
{Text["(x*,y*)",Offset[{10,10},{(x/.x→Es[[1])],(y/.y→Es[[2])}]}],{PointSize[Large],
Style[Point[{(x/.x→Es[[1])],(y/.y→Es[[2])}],Black]}},{PointSize[Large],Point[{0.9,0.01}],
Text["(x0*,y0*)",Offset[{-10,8},{0.9,0.01}]]}}]
Export["pp10si.pdf",pp10si]
Export["pb10i.pdf",pb10i]
```

$E^* = \{0.20202, 0.0541003, 2.43451\}$

$E_+ = \{0.30876, 0.06, 2.06588, 0.352938\}$

$E_i = \{0.411479, 0.06, 1.72971, 0.635108\}$



Out[*]= pp10si.pdf

Out[*]= pb10i.pdf

When $b_2^* < b = 15 < b_H$:

```

In[*]:= ClearParameters;
μv=0.16; μy=0.48;K=1;b=15;γ=1;λ=0.36;β=0.11;δ=0.2;s=0.6; c=0.036;ε=0;
Print["E*",Es=Est/.cn//N]
Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp/.cn//N]]
Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi/.cn//N]]

x1=λ x[t] (1-(x[t]+y[t])/K)- β x[t]×v[t] ;
y1=β x[t]×v[t] -μy y[t]×z[t]- γ y[t];
v1=-β x[t]×v[t] - μv v[t]×z[t]+ b γ y[t] - δ v[t];
z1=z[t] (s y[t] - c);
ode4={x'[t]==x1,y'[t]==y1,v'[t]==v1,z'[t]==z1,x[0]==0.9,y[0]==0.01,v[0]==0.01,
z[0]==0.01};
sol2H=NDSolve[ode4,{x,y,v,z},{t,0,800}];
pdy2H=Plot[{x[t]/.sol2H,y[t]/.sol2H,v[t]/100/.sol2H,z[t]/.sol2H},{t,0,800},
PlotLegends→{"x","y","v/100","z"}];
pEs2H=Plot[{x/.x→Es[[1]],y/.y→Es[[2]],v/100/.v→Es[[3]],z/.z→0},{t,0,800},
PlotStyle→{Dashed}];
Dyn2H=Show[pdy2H,pEs2H,PlotRange→All]



(*****Parametric plot conditions****)
x0=0.9; y0=0.01;v0=0.01;z0=0.01;
ppb15=ParametricPlot[{ x[t],(y[t]) }/.sol2H,{t,0,500}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Blue}];
py15=Plot[(y/.y→Es[[2]]),{t,0,400},PlotStyle→{Dashed,Green}];
pb15=Show[{ppb15,py15, Graphics[{Green,Dashed,Line[{x/.x→Es[[1]],0},
{x/.x→Es[[1]],1}]}]},Epilog→{{Thick,Text["(x*,y*)",Offset[{10,10},{x/.x→Es[[1]],
(y/.y→Es[[2]])}]}},{PointSize[Large],Style[Point[{x/.x→Es[[1]],(y/.y→Es[[2]])}],Black]}},
{PointSize[Large],Point[{x0,y0}],Text["(x0,y0)",Offset[{-10,8},{x0,y0}]]}]
Export["pb15.pdf",pb15]
Export["Dyn2H.pdf",Dyn2H]

```


$E^* \{0.12987, 0.0388644, 2.72051\}$

$E_+ = \{0.332366 + 0.217051 i, 0.06, 1.98862 - 0.710347 i, 1.03002 + 0.746836 i\}$

$E_- = \{0.332366 - 0.217051 i, 0.06, 1.98862 + 0.710347 i, 1.03002 - 0.746836 i\}$



Out[]= pb15.pdf

Out[]= Dyn2H.pdf

When $b_H < b = 23 < b_\infty$:

In[]:=

```

ClearParameters;
μv=0.16; μy=0.48;K=1;b=23;γ=1;λ=0.36;β=0.11;δ=0.2;s=0.6; c=0.036;ε=0;
Print["E*",Es=Est/.cn//N]
Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp/.cn//N]]
Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi/.cn//N]]
x1=λ x[t] (1-(x[t]+y[t])/K)- β x[t]×v[t] ;
y1=β x[t]×v[t] -μy y[t]×z[t]- γ y[t];
v1=-β x[t]×v[t] - μv v[t]×z[t]+ b γ y[t] - δ v[t];
z1=z[t] (s y[t] - c );
ode5={x'[t]==x1,y'[t]==y1,v'[t]==v1,z'[t]==z1,x[0]==0.9,y[0]==0.01,v[0]==0.01,
z[0]==0.01};
solHI=NDSolve[ode5,{x,y,v,z},{t,0,10000}];
pdyHI=Plot[{x[t]/. solHI,y[t]/. solHI,v[t]/. solHI,z[t]/. solHI},{t,0,1000},
PlotLegends→{"x","y","v","z"}];
DynHI=Show[pdyHI,PlotRange→All]



(*****Parametric plot conditions****)
x0=0.9; y0=0.01;v0=0.01;z0=0.01;
ppb23=ParametricPlot[{ x[t],(y[t])}/.solHI,{t,0,200}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Blue}];
NDSolve[{y'[t]==y[t]×(x[t]-1),x'[t]==x[t] (2-y[t]),x[0]==1,y[0]==2.7},{x,y},{t,0,10}];
py23=Plot[(y/.y→Es[[2]]},{t,0,400},PlotStyle→{Dashed,Green}];
pb23=Show[{ppb23},Epilog→{{Thick,Text["(x*,y*)",Offset[{10,10},{(x/.x→Es[[1]]),
(y/.y→Es[[2]])}]}},{PointSize[Large],Style[Point[{(x/.x→Es[[1])},(y/.y→Es[[2]])}],Green}}},
{PointSize[Large],Point[{x0,y0}],Text["(x0,y0)",Offset[{-10,8},{x0,y0}]}]}]
Export["pb23.pdf",pb23]
Export["DynHI.pdf",DynHI]

```

E*{0.0826446, 0.0265046, 2.91551}

E+={0.297813 + 0.339863 i, 0.06, 2.1017 - 1.11228 i, 1.75115 + 1.463 i}

Ei={0.297813 - 0.339863 i, 0.06, 2.1017 + 1.11228 i, 1.75115 - 1.463 i}




Out[]= pb23.pdf

Out[]= DynHI.pdf

```


In[ ]:= ode5={x'[t]==x1,y'[t]==y1,v'[t]==v1,z'[t]==z1,x[0]==0.3,y[0]==0.05,v[0]==2,
z[0]==0.5};
solHI=NDSolve[ode5,{x,y,v,z},{t,0,10000}];
pdyHI=Plot[{x[t]/.solHI,y[t]/.solHI,v[t]/.solHI,z[t]/.solHI},{t,0,1000},
PlotLegends->{"x","y","v","z"}];
DynHIc=Show[pdyHI,PlotRange->All]
(*New initial conditions*)
x0=0.3; y0=0.05;v0=2;z0=0.5;
ppb23c=ParametricPlot[{x[t],(y[t])}/.solHI,{t,0,150}, AxesLabel->{"x","y"},
PlotRange->Full,PlotStyle->{Red}];
pb23c=Show[{ppb23c},Epilog->{{Thick,Text["(x*,y*)",Offset[{10,10},
{x/.x->Es[[1]],(y/.y->Es[[2]])}],{PointSize[Large],Style[Point[{x/.x->Es[[1]],
(y/.y->Es[[2]])}],Green}],{PointSize[Large],Point[{x0,y0}],Text["(x0,y0)",
Offset[{-10,8},{x0,y0}]]}}]
Export["pb23b.pdf",pb23c]
Export["DynHIb.pdf",DynHIc]
pp23cs=Show[{pb23,pb23c},Epilog->{{Thick,Text["(x*,y*)",Offset[{10,10},
{x/.x->Es[[1]],(y/.y->Es[[2]])}],{PointSize[Large],Style[Point[{x/.x->Es[[1]],(y/.y->Es[[2]])}],Green}],{
PointSize[Large],Point[{x0,y0}],Text["(x0',y0')",Offset[{16,6},{x0,y0}]],
{PointSize[Large],Point[{0.9,0.01}],Text["(x0*,y0*)",Offset[{-10,8},{0.9,0.01}]]}}]
Export["pp23s.pdf",pp23cs]

```



Out[]= pb23b.pdf

Out[]= DynHIb.pdf



Out[]= pp23s.pdf

4)Section 5(in paper): 4-Dim.Viro-therapy model when $\epsilon=1$

4-1)Definition of the model and fixed points when $\epsilon=1$

In[]:=

```

SetDirectory[NotebookDirectory[]];
AppendTo[$Path,Directory];
Clear["Global`*"];
Clear["K"];
Format[μv]:=Subscript[μ,v];Format[μy]:=Subscript[μ,y];Format[La]:=Δ;(*La >0*)
pars={β,λ,γ,δ,μy, μv,b,K,s,c};
cpos={β>0,λ>0,γ>0,δ>0,μy>0, μv>0,b>1,K>0,s>0,c>0};
La=0; bL=100;cnb={b→50};
cEri={μy→1/48,K→2139.258, β→.0002,λ→.2062,γ→1/18,δ→.025, μv→2*10^(-8),c→10^(-3),s→.027};
(***** Four dim Deterministic epidemic model with Logistic growth *****)
x1=La+λ x(1-(x+y)/K)- β x v ;
y1=β x v -μy y z- γ y;
v1=-β x v - μv v z+ b γ y - δ v;
z1=z(s y - c z);
x1s=λ (1-(x+y)/K)- β v ;
z1s=s y - c z;
dys={x1s,y1,v1,z1s};
dyn={x1,y1,v1,z1};
dyn3={x1,y1,v1}/.z→0;
Print["
      x'
      ( y' )= ",dyn//FullSimplify//MatrixForm]
      v'
      z'

(*Jacobian*)
Jac=Grad[(dyn),{x,y,v,z}]/FullSimplify;
det=Det[Jac]/FullSimplify; tr=Tr[Jac]/FullSimplify;
R0=b β K/(β K+δ);bcrit=1+δ/(β K);(*Reduce[Join[{R0>1},pars],δ]*)
(****Endemic points in 3dims **)
eqE=Thread[dyn3=={0,0,0}];
Print["Three Fixed points in 3-dim case:",solE=Solve[eqE,{x,y,v}]/FullSimplify]

el=Eliminate[Thread[dyns=={0,0,0,0}],{x,v,z}];
Qybyelim=Factor[el[[1,1]]-el[[1,2]]/y]/FullSimplify;
Print["Coefficients of Qy by elim polynomial are:"]
cof=CoefficientList[Qybyelim,y]/FullSimplify
so=y/.Solve[Qybyelim==0,y];(*Third order roots*)

(*****Fixed points of 4-dim model using P(y)*****)
fy=(c γ(b-1)-y μy s);
gy=( μv s y+c δ); hy=(γ +y s μy/c);
xe=hy gy/(β fy); ve=y fy/gy; ze= s y /c;
ys=y/.solE[[3]](* y of E* ****);

Py=λ(1-y/K)-β y fy/gy-λ hy gy/(β K fy); yb=c γ (b-1)/(μy s);
Qy=λ fy gy(1- y /K)- λ hy gy^2/(β K)-y β fy^2//FullSimplify;
Qycol=Collect[Qy,y];
Qycoef=CoefficientList[Qycol,y];
(*Print["Check Eric Qy -elim Qy=",Qybyelim+c K β Qy//FullSimplify]*)
Dis=Collect[Discriminant[Qy,y],b];
Discoef=CoefficientList[Dis,b];Length[Discoef];
DisE=Dis//.cEri//N;

```

$$\begin{pmatrix} x' \\ y' \\ v' \\ z' \end{pmatrix} = \begin{pmatrix} -v x \beta + x \left(1 - \frac{x+y}{K}\right) \lambda \\ v x \beta - y (\gamma + z \mu_y) \\ b y \gamma - v (x \beta + \delta + z \mu_v) \\ z (s y - c z) \end{pmatrix}$$

Three Fixed points in 3-dim case: $\{x \rightarrow 0, y \rightarrow 0, v \rightarrow 0\}, \{x \rightarrow K, y \rightarrow 0, v \rightarrow 0\},$

$$\left\{ x \rightarrow \frac{\delta}{(-1+b)\beta}, y \rightarrow \frac{((-1+b)K\beta - \delta)\delta\lambda}{(-1+b)\beta((-1+b)K\beta\gamma + \delta\lambda)}, v \rightarrow \frac{\gamma((-1+b)K\beta - \delta)\lambda}{\beta((-1+b)K\beta\gamma + \delta\lambda)} \right\}$$

Coefficients of Qy by elim polynomial are:

$$\begin{aligned} \text{Out}[*]= & \left\{ c^3 \gamma \delta (K(\beta - b\beta) + \delta) \lambda, \right. \\ & c^2 \left((-1+b) c \beta \gamma ((-1+b) K \beta \gamma + \delta \lambda) + s \lambda (\gamma (K(\beta - b\beta) + 2\delta) \mu_v + \delta (K\beta + \delta) \mu_y) \right), \\ & c s \left(s \lambda \mu_v (\gamma \mu_v + K \beta \mu_y + 2\delta \mu_y) + c \beta (-\delta \lambda \mu_y + (-1+b) \gamma (\lambda \mu_v - 2 K \beta \mu_y)) \right), \\ & \left. s^2 \mu_y (s \lambda \mu_v^2 + c \beta (-\lambda \mu_v + K \beta \mu_y)) \right\} \end{aligned}$$


```

In[*]:= Plot[{0,DisE},{b,0,2 bL},AxesLabel->{"b","Dis"}]
Print["Roots of Dis[b]=0 are: ",solbE=Solve[DisE==0,b]]
QR=Solve[Qy==0,y];
Print["critical b from R0=1 is:", bcrit//.cEri//N]
ym= y/.QR[[1]];
yp= y/.QR[[2]];yi= y/.QR[[3]];

{Chop[yp],Chop[ym]}//.cEri//Simplify;

jacEK=Jac/.x->K/.y->0/.v->0/.z->0;
Print["J(EK)=", jacEK//MatrixForm]
Print["Eig.val of J(EK) are:", Eigenvalues[jacEK]//FullSimplify]

```



Roots of Dis[b]=0 are:

$$\{b \rightarrow -468.749.\}, \{b \rightarrow -468.749.\}, \{b \rightarrow -159.797\}, \{b \rightarrow -132.005\}, \{b \rightarrow 133.421\}, \{b \rightarrow 159.122\}$$

critical b from R0=1 is:1.05843

$$J(EK) = \begin{pmatrix} -\lambda & -\lambda & -K\beta & 0 \\ 0 & -\gamma & K\beta & 0 \\ 0 & b\gamma & -K\beta - \delta & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Eig.val of J(EK) are: $\left\{0, \frac{1}{2} \left(-K\beta - \gamma - \delta - \sqrt{-4\gamma(K(\beta - b\beta) + \delta) + (K\beta + \gamma + \delta)^2} \right), \right.$

$$\left. \frac{1}{2} \left(-K\beta - \gamma - \delta + \sqrt{-4\gamma(K(\beta - b\beta) + \delta) + (K\beta + \gamma + \delta)^2} \right), -\lambda \right\}$$

4-2) Stability of the interior points using Routh Hurwitz:

In[]:=

```
jacE=Jac/.x->xe/.v->ve/.z->ze/.y->y;
jacE//MatrixForm;
Print["Det [Jac]=",Det[jacE]//FullSimplify, " , Trace [Jac]=",
Tr[jacE]//FullSimplify]
(*Reduce[Join[{Det [Jac]>0&&x>0&&y>0&&z>0&&v>0},cpos],β] (*take so long time***)
Reduce[Join[{Tr [Jac]<0&&x>0&&y>0&&z>0&&v>0},cpos],β]//FullSimplify;
poly=Collect[Together[Det [ψ IdentityMatrix[4]-Jac]],ψ]//FullSimplify;
coe=CoefficientList[poly,ψ]//FullSimplify
```

Det [Jac] =

$$\begin{aligned}
 & -\frac{1}{c^3 K \beta \left((-1+b) c \gamma - s y \mu_y \right)^2} s y^2 \left(-(-1+b)^2 c^5 \beta \gamma^3 \left((-1+b) K \beta \gamma + \delta \lambda \right) + 2 s^5 y^4 \lambda \mu_v^2 \mu_y^3 + c s^4 \right. \\
 & \quad y^3 \mu_y^2 \left(2 \times (3-2b) \gamma \lambda \mu_v^2 + 2 (-y \beta + \delta) \lambda \mu_v \mu_y + K \beta \mu_y \left(\lambda \mu_v + 2 y \beta \mu_y \right) \right) - \\
 & \quad c^2 s^3 y^2 \mu_y \left(6 \times (-1+b) \gamma^2 \lambda \mu_v^2 + y \beta \delta \lambda \mu_y^2 + \right. \\
 & \quad \gamma \mu_y \left((3 \times (-1+b) K \beta + (6-5b) y \beta + 6 \times (-1+b) \delta) \lambda \mu_v + (-7+6b) K y \beta^2 \mu_y \right) \left. \right) - \\
 & \quad c^3 s^2 y \gamma \left(2 \times (-1+b) \gamma^2 \lambda \mu_v^2 + \delta (3 y \beta + b (K \beta - 3 y \beta + 2 \delta)) \lambda \mu_y^2 + \right. \\
 & \quad \gamma \mu_y \left((3 \times (-2+b) \times (-1+b) y \beta - 6 \delta + 8 b \delta) \lambda \mu_v - (-1+b) \times (-3+2b) K \beta \left(\lambda \mu_v + 3 y \beta \mu_y \right) \right) \left. \right) + \\
 & \quad c^4 s \gamma^2 \left(-2 \times (-1+b) \gamma \left((-1+b) y \beta + \delta \right) \lambda \mu_v - \delta \left((-1+b) \times (-3+2b) y \beta + 2 b \delta \right) \lambda \mu_y + \right. \\
 & \quad \left. (-1+b) K \beta \left(b \delta \lambda \mu_y + (-1+b) \gamma \left(\lambda \mu_v + (5-2b) y \beta \mu_y \right) \right) \right) \left. \right) \\
 & , \text{Trace [Jac]} = -s y - \gamma - \delta + \lambda - \frac{s y \mu_v}{c} - \frac{s y \mu_y}{c} + \frac{y \beta \left(c \left(\gamma - b \gamma \right) + s y \mu_y \right)}{c \delta + s y \mu_v} - \\
 & \frac{(c \delta + s y \mu_v) \left(\gamma + \frac{s y \mu_y}{c} \right)}{(-1+b) c \gamma - s y \mu_y} - \\
 & \frac{\lambda \left(y + \frac{2 (c \delta + s y \mu_v) \left(\gamma + \frac{s y \mu_y}{c} \right)}{\beta \left((-1+b) c \gamma - s y \mu_y \right)} \right)}{K}
 \end{aligned}$$

$$\begin{aligned}
\text{Out}[*]= & \left\{ -s \left(y \gamma \lambda \left((-1+b) x \beta - \delta - z \mu_v \right) + v \beta \left(x z \lambda \mu_v + y \gamma \left(\delta + z \mu_v \right) \right) \right) + \right. \\
& 2 c z \left(x \beta \lambda \left((-1+b) \gamma - z \mu_y \right) + v \beta \left(\delta + z \mu_v \right) \left(\gamma + z \mu_y \right) - \lambda \left(\delta + z \mu_v \right) \left(\gamma + z \mu_y \right) \right) + \\
& \frac{1}{K} \lambda \left(s y \left((-1+b) x \left(2 x + y \right) \beta \gamma - \left(v x \beta + \left(2 x + y \right) \gamma \right) \delta \right) + \right. \\
& s z \left(2 v x^2 \beta - y \left(2 x + y \right) \gamma \right) \mu_v + 2 c z \left(y \left(\delta + z \mu_v \right) \left(\gamma + z \mu_y \right) + \right. \\
& \left. 2 x^2 \beta \left(\gamma - b \gamma + z \mu_y \right) + x y \beta \left(\gamma - b \gamma + z \mu_y \right) + x \left(\delta + z \mu_v \right) \left(v \beta + 2 \left(\gamma + z \mu_y \right) \right) \right) \left. \right), \\
& s y \left(-\gamma \delta - v \beta \left(\gamma + \delta \right) + \left(\gamma + \delta \right) \lambda + x \beta \left((-1+b) \gamma + \lambda \right) \right) + s z \left(v \left(x - y \right) \beta + y \left(-\gamma + \lambda \right) \right) \mu_v + \\
& \gamma \left(\lambda \left((-1+b) x \beta - \delta - z \mu_v \right) + v \beta \left(\delta + z \mu_v \right) \right) + \\
& z \left(v \beta \left(\delta + z \mu_v \right) - \lambda \left(x \beta + \delta + z \mu_v \right) \right) \mu_y + \\
& 2 c z \left(\gamma \delta - \gamma \lambda - \delta \lambda + z \gamma \mu_v - z \lambda \mu_v + z \left(\delta - \lambda + z \mu_v \right) \mu_y + \right. \\
& \left. x \beta \left(\gamma - b \gamma - \lambda + z \mu_y \right) + v \beta \left(\gamma + \delta + z \left(\mu_v + \mu_y \right) \right) \right) + \\
& \frac{1}{K} \lambda \left(2 x^2 \beta \gamma - 2 b x^2 \beta \gamma + x y \beta \gamma - b x y \beta \gamma + v x \beta \delta + 2 x \gamma \delta + y \gamma \delta + v x z \beta \mu_v + \right. \\
& 2 x z \gamma \mu_v + y z \gamma \mu_v - s y \left(v x \beta + \left(2 x + y \right) \left(x \beta + \gamma + \delta + z \mu_v \right) \right) + \\
& \left. \left(2 x + y \right) z \left(x \beta + \delta + z \mu_v \right) \mu_y + 2 c z \left(v x \beta + \left(2 x + y \right) \left(x \beta + \gamma + \delta + z \left(\mu_v + \mu_y \right) \right) \right) \right) \left. \right), \\
& v \beta \gamma + x \beta \gamma - b x \beta \gamma + v \beta \delta + \gamma \delta - x \beta \lambda - \gamma \lambda - \delta \lambda + v z \beta \mu_v + z \gamma \mu_v - z \lambda \mu_v - \\
& s y \left(\left(v + x \right) \beta + \gamma + \delta - \lambda + z \mu_v \right) + z \left(\left(v + x \right) \beta + \delta - \lambda + z \mu_v \right) \mu_y + \\
& \frac{1}{K} \lambda \left(-s y \left(2 x + y \right) + 2 c \left(2 x + y \right) z + v x \beta + 2 x^2 \beta + x y \beta + 2 x \gamma + y \gamma + 2 x \delta + y \delta + \right. \\
& \left. 2 x z \mu_v + y z \mu_v + \left(2 x + y \right) z \mu_y \right) + 2 c z \left(\left(v + x \right) \beta + \gamma + \delta - \lambda + z \left(\mu_v + \mu_y \right) \right), \\
& -s y + \left(v + x \right) \beta + \gamma + \delta - \lambda + \frac{\left(2 x + y \right) \lambda}{K} + z \left(2 c + \mu_v + \mu_y \right), \\
& 1 \}
\end{aligned}$$

4-3) Stability of the interior points numerically :

Routh Hurwitz conditions for the stability of E_- (4 dim)

```

In[*]:= Jac4=Grad[dyn//.cEri,{x,y,v,z}]/FullSimplify;
bcrit=1+δ/(β K);
Jst=(Jac4/.x→xe/.v→ve/.z→ze/.y→ym)//.cEri;Jst//N//MatrixForm;
Trs=Tr[Jst];
pc=Collect[Det[ψ IdentityMatrix[4]-Jst],ψ];
coT=CoefficientList[pc,ψ](*So long computations*);

```

Routh Hurwitz conditions for the stability of E_*

In[]:=

```

cF1={β→ $\frac{87}{2}$ , λ→1, γ→ $\frac{1}{128}$ , δ→1/2, μy→1, μv→1, K→1/2, s→1, c→1};
cEri=cF1;
Jac3=Grad[dyn3//.cEri,{x,y,v}]/FullSimplify;
bcrit=1+δ/(β-K);(*Reduce[Join[{R0>1},pars],δ]*)
Print["J(E_*) is"]
Jst=(Jac3/.x→x/.solE[[3]]/.y→ys/.v→v/.solE[[3]])//.cEri//FullSimplify;Jst//MatrixForm

Trs=Tr[Jst];
pc=Collect[Det[ψ IdentityMatrix[3]-Jst],ψ];
coT=CoefficientList[pc,ψ]//FullSimplify;
Print["a1=",a1=coT[[3]]//.cEri, ", a2=",a2=coT[[2]]//.cEri, ", a3=",a3=coT[[1]]//.cEri]
H2=a1*a2-a3;
Print["H2(b0)=",H2/.b→bcrit//FullSimplify]
Print["Denominator of H2 is ",Denominator[Together[H2]]/.cEri//FullSimplify]
φb=Collect[Numerator[Together[H2]]/(δ λ),b]//.cEri;
cofi=CoefficientList[φb,b](*Coefficients of φ(b)*);
Print["value of φ(b) at crit b is "]
φb/.b→bcrit//.cEri//N
cb=NSolve[(H2//.cEri)==0,b,WorkingPrecision→20]
bM=Max[Table[Re[b/.cb[[i]]],{i,Length[cb]}]];
Print["bH=",bH=N[bM,30]]

```

J(E_*) is

Out[]:=MatrixForm=

$$\begin{pmatrix} \frac{2}{87-87b} & \frac{2}{87-87b} & \frac{1}{2-2b} \\ 1-\frac{258}{169+87b} & -\frac{1}{128} & \frac{1}{2(-1+b)} \\ -1+\frac{258}{169+87b} & \frac{b}{128} & \frac{b}{2-2b} \end{pmatrix}$$

$$a_1 = \frac{65}{128} + \frac{91}{174(-1+b)}, \quad a_2 = \frac{-236553 + (478354 - 225417b)b}{5568(-1+b)^2(169+87b)}, \quad a_3 = \frac{87 - \frac{2}{-1+b}}{22272}$$

$$H2(b0) = -\frac{1}{256} + \frac{K\beta \left(-\frac{13695976827}{256K\beta+87\delta} + \frac{5963776K^2\beta^2+13781248K\beta\delta-76919571\delta^2}{\delta^3} \right)}{992083968}$$

Denominator of H2 is 62005248 (-1+b)³ (169+87b)

value of φ(b) at crit b is

$$\text{Out[]}= 2.01216 \times 10^8$$

$$\text{Out[]}= \{ \{b \rightarrow -61.53327023855142092\}, \{b \rightarrow 1.3349402618015729010\}, \\ \{b \rightarrow 0.7834975171549898868\}, \{b \rightarrow 0.0013985515488811205258\} \}$$

$$bH=1.3349402618015729010$$

Numerical solution of the stability (Bifurcation diagram)

In[]:=

```

cond=cF1;
Print["roots of Dis[b]=0:", bcE=NSolve[(Dis//.cond)==0,b]]
Print["b0=",bc=b/.Solve[R0==1,b][[1]]//.cond//N]
bL=100; max=2;
bc1=b/.bcE[[5]];
bc2=b/.bcE[[6]];
lin1=Line[{{bc1,0},{bc1,max}}];
li1=Graphics[{Thick,Black,Dashed,lin1}];
lin2=Line[{{bc2,0},{bc2,max}}];
li2=Graphics[{Thick,Black,Dashed,lin2}];
lin3=Line[{{bc,0},{bc,max}}];
li3=Graphics[{Thick,Black,Dashed,lin3}];
p1a=Plot[{ym} /. cond, {b,0,bc1},PlotStyle->{Dashed,Thick,Green},
PlotRange->All,PlotLegends->{"E_ unstable"}];
p1b=Plot[{ym} /. cond, {b,bc1,bL},PlotStyle->{Green},PlotRange->All,
PlotLegends->{"E_ stable"}];
p01=Plot[{0, {b,0,bc}},PlotStyle->{Magenta},PlotRange->All,PlotLegends->{"E_1 stable"}];
p02=Plot[{0, {b,bc,bL}},PlotStyle->{Blue},PlotRange->All,PlotLegends->{"E_1 unstable"}];
pp=Plot[{yi} /. cond, {b,0, bL},PlotStyle->{Purple},PlotRange->All,
PlotLegends->{"E_i unstable"}];
pm=Plot[{yp} /. cond, {b,0,bL},PlotStyle->{Red},PlotRange->All,
PlotLegends->{"E_+ unstable"}];
ps1=Plot[{ys} /. cond, {b,0,13.45},PlotStyle->{Orange, Dotted},
PlotRange->{{0,200},{0,max}},
PlotLegends->{"E_* outside the domain"}];
ps2=Plot[{ys} /. cond, {b,13.45,bL},PlotStyle->{Orange,Thick,Dashed},
PlotRange->{{0,200},{0,max}},
PlotLegends->{"E_* unstable"}];

pyb=Plot[{yb} /. cond, {b,0,bL},PlotStyle->{Dashed,Thick,Cyan},
PlotRange->{{0,200},{0,max}},PlotLegends->{"y_b "}]];
Print["y*'(b0)=",D[ys,b]/.b->bc//.cond//N//FullSimplify]
Chop[ys/.b->bc//.cond//N] (*Check*);
bifE2=Show[{p01,p02,ps1,ps2,pyb,li3},PlotRange->{{0,3},{0,1}},
Epilog->{Text["b0",Offset[{10,11},{bc} /. cond,0]],{PointSize[Large],
Style[Point[{bc} /. cond,0],Black]}},
PlotRange->All,AxesLabel->{"b","y_ee"}]
Export["bifEE.pdf",bifE2]


```

roots of Dis[b]=0:

{ {b → -152.24}, {b → -63.}, {b → -63.}, {b → -25.6124}, {b → 27.2273}, {b → 30.5559} }

b0=1.02299

y*'(b0)=21.5814




Out[]:= bifEE.pdf

In[]:=

```

p1a=Plot[{ym} /. cond, {b, 0, 13.45}, PlotStyle -> {Thick, Green}, PlotRange -> All];
p1c=Plot[{ym} /. cond, {b, 13.45, bc1}, PlotStyle -> {Thick, Green},
PlotRange -> All, PlotLegends -> {"E_ stable"}];
p1d=Plot[{ym} /. cond, {b, bc1, bL}, PlotStyle -> {Dashed, Thick, Green}, PlotRange -> All,
PlotLegends -> {"E_ unstable"}];
pi=Plot[{yi} /. cond, {b, 0, bL}, PlotStyle -> {Dashed, Thick, Red}, PlotRange -> All,
PlotLegends -> {"E_i unstable"}];
pp1=Plot[{yp} /. cond, {b, 0, bL}, PlotStyle -> {Purple}, PlotRange -> All,
PlotLegends -> {"E_+ stable"}];
shon=Show[{p1a, p1c, p1d, li1, li2}, PlotRange -> {{0, 60}, {0, 0.2}}, Epilog ->
{{Text["b1*", Offset[{-8, 10}, {bc1, 0}], {PointSize[Large],
Style[Point[{bc1, 0}], Purple]}, Text["b2*", Offset[{10, 10}, {bc2, 0}],
{PointSize[Large], Style[Point[{bc2, 0}], Yellow]}]
}, AxesLabel -> {"b", "yee"}];
shoip=Show[{pi, pp1}, PlotRange -> All, AxesLabel -> {"b", "yee"}];
Bnip=Show[shon, shoip];
EriB=Show[shon, shoip, bifE2, PlotRange -> {{0, 35}, {0, 0.3}}, Epilog -> {
Text["b0", Offset[{10, 11}, {bc /. cond, 0}], {PointSize[Large],
Style[Point[{bc /. cond, 0}], Black]},
Text["b1*", Offset[{-8, 10}, {bc1, 0}], {PointSize[Large],
Style[Point[{bc1, 0}], Purple]},
Text["b2*", Offset[{10, 10}, {bc2, 0}], {PointSize[Large],
Style[Point[{bc2, 0}], Yellow]}]}]
Export["Bnip.pdf", Bnip]
Export["EriB.pdf", EriB]

```



Out[]:= Bnip.pdf

Out[]:= EriB.pdf