On a four-dimensional oncolytic Virotherapy model (ϵ=1)

This Mathematica Notebook is a supplementary material to the paper "On a three-dimensional and two four-dimensional
oncolytic viro-therapy models". It contains some of the calculations and illustrations appearing in the paper.

Ep1)Section 3.5(in paper): 4-Dim.Viro-therapy model when ϵ=1

Ep1-1)Definition of the model and fixed points when ϵ=1

```
In[1]:= SetDirectory[NotebookDirectory[]];
        AppendTo[$Path, Directory];
        Clear["Global`*"];
        Clear["K"];
        (*Some aliases*)
        Format[βy] = Subscript[β, y];
        Format[βv] = Subscript[β, v];
        Format[βz] = Subscript[β, z];


        Unprotect[Power];
        Power[0, 0] = 1;
        Protect[Power];


        par = {b, β, λ, δ, βy, βv, βz, c, γ, K, ε};
        cp = Join[Thread[Drop[par, {1}]] > 0], {b > 1}];
        cKga1 = {K → 1, γ → 1};
        cep1 = {ε → 1};
        R0 = b β K / (β K + δ) (* Reproduction number*);


        (*cnb={b→50};
        cE1ri=Join[{βy→1/48,K→2139.258, β→.0002,λ→.2062,
            γ→1/18,δ→.025, βv→2*10^(-8),c→10^(-3),βz→.027},cep1];*)
        cF1 = {β → 87/2, λ → 1, γ → 1/128, δ → 1 / 2, βy → 1, βv → 1, K → 1, βz → 1, c → 1, ε → 1};


        (****** Four dim Deterministic  epidemic model with Logistic growth ****)
        x1 = λ x (1 - (x + y) / K) -  β x v ;
        y1 = β x v - βy y z - γ y;
        v1 = - β x v - βv v z + b γ y - δ v;
        z1 = z (βz y - c z^ε);
        dyn = {x1, y1, v1, z1};
        dyn3 = {x1, y1, v1} /. z → 0;(*3dim case used for E* *)
                    x'
        Print[" (y') =", dyn // FullSimplify // MatrixForm]
                    v'
                    z'
        Print["b0=", b0 = b /. Apart[Solve[R0 == 1, b][[1]] // FullSimplify]]


        (***Fixed point when z→0**)
        eq = Thread[dyn3 == {0, 0, 0}];
        sol = Solve[eq, {x, y, v}] // FullSimplify;
        Es = {x, y, v} /. sol[[3]]; (*Endemic point with z=0*);
        Print[" Endemic point with z=0 is E*=", Es // FullSimplify]
        jacD = Grad[dyn /. cep1, {x, y, v, z}];
        Print["J(x,y,v,z)="]
        jacD // MatrixForm
        Print["Det(J(x,y,v,z))="]
        Det[jacD] // FullSimplify
```

$$\left(\begin{matrix} x' \\ y' \\ v' \\ z' \end{matrix}\right) = \left(\begin{matrix} -v\,x\,\beta + x\left(1-\frac{x+y}{K}\right)\lambda \\ v\,x\,\beta - y\left(z\,\beta_y + \gamma\right) \\ b\,y\,\gamma - v\left(x\,\beta + z\,\beta_v + \delta\right) \\ z\left(-c\,z^\epsilon + y\,\beta_z\right) \end{matrix}\right)$$

$$b0 = 1 + \frac{\delta}{K\,\beta}$$

Endemic point with z=0 is E∗=

$$\left\{\frac{\delta}{(-1+b)\,\beta},\; \frac{\left((-1+b)\,K\,\beta - \delta\right)\delta\,\lambda}{(-1+b)\,\beta\left((-1+b)\,K\,\beta\,\gamma + \delta\,\lambda\right)},\; \frac{\gamma\left((-1+b)\,K\,\beta - \delta\right)\lambda}{\beta\left((-1+b)\,K\,\beta\,\gamma + \delta\,\lambda\right)}\right\}$$

J (x,y,v,z) =

Out[29]//MatrixForm=

$$\left(\begin{matrix} -v\,\beta - \frac{x\,\lambda}{K} + \left(1-\frac{x+y}{K}\right)\lambda & -\frac{x\,\lambda}{K} & -x\,\beta & 0 \\ v\,\beta & -z\,\beta_y - \gamma & x\,\beta & -y\,\beta_y \\ -v\,\beta & b\,\gamma & -x\,\beta - z\,\beta_v - \delta & -v\,\beta_v \\ 0 & z\,\beta_z & 0 & -2\,c\,z + y\,\beta_z \end{matrix}\right)$$

Det (J(x,y,v,z)) =

Out[31]= $\frac{1}{K}$

$\left(2\,c\,z\left(K\,v\,\beta\left(z\,\beta_y+\gamma\right)\left(z\,\beta_v+\delta\right)+v\,x\,\beta\left(z\,\beta_v+\delta\right)\lambda - K\left(x\,\beta\left(z\,\beta_y+\gamma-b\,\gamma\right)+\left(z\,\beta_y+\gamma\right)\left(z\,\beta_v+\delta\right)\right)\right.\right.$
$\left.\lambda + (2\,x+y)\left(x\,\beta\left(z\,\beta_y+\gamma-b\,\gamma\right)+\left(z\,\beta_y+\gamma\right)\left(z\,\beta_v+\delta\right)\right)\lambda\right) -$
$\beta_z\left(K\,y\,\gamma\left((-1+b)\,x\,\beta - z\,\beta_v-\delta\right)\lambda - y\,(2\,x+y)\,\gamma\left((-1+b)\,x\,\beta - z\,\beta_v-\delta\right)\lambda + \right.$
$\left.\left. v\,x\,\beta\left(-2\,x\,z\,\beta_v + y\,\delta\right)\lambda + K\,v\,\beta\left(y\,\gamma\left(z\,\beta_v+\delta\right)+x\,z\,\beta_v\,\lambda\right)\right)\right)$

In[32]:=
```
(*****Fixed points of 4-dim model using P(y)***)
fy=(c γ(b-1)-y βy βz);
gy=( βv βz y+c δ); hy=(γ +y βz βy/c);
xey=hy gy/(β fy); vey=y fy/gy; zey= βz y /c;
ys=y/.sol〚3〛(* y of E*      ****);

Py=λ(1-y/K)-β y fy/gy-λ hy gy/(β K fy); yb=c γ (b-1)/(βy βz);
Qy=λ fy gy(1-  y /K)- λ hy gy^2/(β K )-y β fy^2;
Qycol=Collect[Together[Qy],y];
Qycoef=CoefficientList[Qycol,y];
Print["f(y)=", fy, " ,g(y)=", gy, " , h(y)=", hy]
Print["p(y)=", Py//FullSimplify]
Print["The polynomial  Q(y) is of order  ", Length[Qycoef]-1]
Print["Coefficients of Q(y) are ", Qycoef//FullSimplify]

Dis=Collect[Discriminant[Qy,y],b];
Discoef=CoefficientList[Dis,b];Length[Discoef];
Disn=Dis//.cF1//N;
bL=50;
Plot[{0,Disn},{b,0,2 bL},AxesLabel→{"b","Dis"},PlotRange→{{0,bL},{-10,12}}]
Print["Roots of Dis[b]=0 are: ",solbE=Solve[Disn==0,b]]
```

$$f(y) = -y \, \beta_y \, \beta_z + (-1 + b) \, c \, \gamma \quad , g(y) = y \, \beta_v \, \beta_z + c \, \delta \quad , \quad h(y) = \frac{y \, \beta_y \, \beta_z}{c} + \gamma$$

$$p(y) = \frac{y \, \beta \, (y \, \beta_y \, \beta_z - (-1 + b) \, c \, \gamma)}{y \, \beta_v \, \beta_z + c \, \delta} + \lambda - \frac{y \, \lambda}{K} - \frac{\left(\frac{y \, \beta_y \, \beta_z}{c} + \gamma \right) (y \, \beta_v \, \beta_z + c \, \delta) \, \lambda}{K \, \beta \, \left(-y \, \beta_y \, \beta_z + (-1 + b) \, c \, \gamma \right)}$$

The polynomial Q(y) is of order 3

Coefficients of Q(y) are $\Big\{ \dfrac{c^2 \, \gamma \, ((-1 + b) \, K \, \beta - \delta) \, \delta \, \lambda}{K \, \beta}$,

$-\dfrac{c \, \left(\beta_z \, \left(\delta \, \left(2 \, \beta_v \, \gamma + \beta_y \, \delta \right) + K \, \beta \, \left(\beta_v \, \gamma - b \, \beta_v \, \gamma + \beta_y \, \delta \right) \right) \lambda + (-1 + b) \, c \, \beta \, \gamma \, ((-1 + b) \, K \, \beta \, \gamma + \delta \, \lambda) \right)}{K \, \beta}$,

$\dfrac{\beta_z \, \left(-\beta_v \, \beta_z \, \left(K \, \beta \, \beta_y + \beta_v \, \gamma + 2 \, \beta_y \, \delta \right) \lambda + c \, \beta \, \left(2 \times (-1 + b) \, K \, \beta \, \beta_y \, \gamma + \beta_v \, (\gamma - b \, \gamma) \, \lambda + \beta_y \, \delta \, \lambda \right) \right)}{K \, \beta}$,

$-\dfrac{\beta_y \, \beta_z^2 \, \left(\beta_v^2 \, \beta_z \, \lambda + c \, \beta \, \left(K \, \beta \, \beta_y - \beta_v \, \lambda \right) \right)}{c \, K \, \beta} \Big\}$

Out[48]=

Roots of Dis[b]=0 are:

$\{\{b \to -126.518\}, \{b \to -63.\}, \{b \to -63.\}, \{b \to -24.5518\}, \{b \to 29.361\}, \{b \to 45.9232\}\}$

In[50]:=
```
QR=Solve[Qy==0,y,Cubics→False]//ToRadicals(*casus irreducibilis*);
ym= y/.QR〚2〛;
yp=  y/.QR〚1〛;yi=  y/.QR〚3〛;
Em1={xey,ym,vey,zey}//.y→ym;
Ep1={xey,yp,vey,zey}//.y→yp;
Eim1={xey,yi,vey,zey}//.y→yi;
Es1=Join[{x,y,v}/.sol〚3〛,{0}];
{Chop[yp],Chop[ym]}//.cF1//N;

jacE1K=jacD/.x→K/.y→0/.v→0/.z→0;
Print["J(EK)=", jacE1K//MatrixForm]
(*Jacobians of the fixed points**)
jEs1=jacD/.sol〚3〛/.z→0;
jEm1=jacD/.x→xey/.v→vey/.z→zey/.y→ym;
jEim1=jacD/.x→xey/.v→vey/.z→zey/.y→yi;
jEp1=jacD/.x→xey/.v→vey/.z→zey/.y→yp;
Print["Eig.val of J(EK) are:", Eigenvalues[jacE1K]//FullSimplify]
jacE1=jacD/.x→xey/.v→vey/.z→zey/.y→y//FullSimplify;
jacE1 //MatrixForm;
Det[jacE1]//FullSimplify;
Print[" Trace of either Ei, E+ or E- is : ", Tr[jacE1]//FullSimplify]
Print["J(E_*) is"]
jEs1//FullSimplify//MatrixForm
bbs=b/.Solve[yb==(y/.sol〚3〛),b]〚1〛(*long expression*);
```

$$J(EK) = \begin{pmatrix} -\lambda & -\lambda & -K\,\beta & 0 \\ 0 & -\gamma & K\,\beta & 0 \\ 0 & b\,\gamma & -K\,\beta - \delta & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Eig.val of J(EK) are:$\left\{0, \dfrac{1}{2}\left(-K\,\beta - \gamma - \delta - \sqrt{-4\,\gamma\,(K\,(\beta - b\,\beta) + \delta) + (K\,\beta + \gamma + \delta)^2}\right),\right.$

$\left.\dfrac{1}{2}\left(-K\,\beta - \gamma - \delta + \sqrt{-4\,\gamma\,(K\,(\beta - b\,\beta) + \delta) + (K\,\beta + \gamma + \delta)^2}\right), -\lambda\right\}$

Trace of either Ei, E+ or E- is : $-y\,\beta_z - \dfrac{y\,\beta_y\,\beta_z + c\,\gamma}{c} + \dfrac{y^2\,\beta\,\beta_y\,\beta_z}{y\,\beta_v\,\beta_z + c\,\delta} -$

$\dfrac{(-1+b)\,c\,y\,\beta\,\gamma}{y\,\beta_v\,\beta_z + c\,\delta} + \dfrac{b\,\gamma\,(y\,\beta_v\,\beta_z + c\,\delta)}{y\,\beta_y\,\beta_z - (-1+b)\,c\,\gamma} + \lambda - \dfrac{y\,\lambda}{K} - \dfrac{2\left(\dfrac{y\,\beta_y\,\beta_z}{c} + \gamma\right)(y\,\beta_v\,\beta_z + c\,\delta)\,\lambda}{K\,\beta\,\left(-y\,\beta_y\,\beta_z + (-1+b)\,c\,\gamma\right)}$

J(E_*) is

Out[70]//MatrixForm=

$$\begin{pmatrix} \dfrac{\delta\,\lambda}{K\,\beta - b\,K\,\beta} & \dfrac{\delta\,\lambda}{K\,\beta - b\,K\,\beta} & -\dfrac{\delta}{-1+b} & 0 \\[2ex] \dfrac{\gamma\,((-1+b)\,K\,\beta - \delta)\,\lambda}{(-1+b)\,K\,\beta\,\gamma + \delta\,\lambda} & -\gamma & \dfrac{\delta}{-1+b} & \dfrac{\beta_y\,\delta\,(K\,(\beta - b\,\beta) + \delta)\,\lambda}{(-1+b)\,\beta\,((-1+b)\,K\,\beta\,\gamma + \delta\,\lambda)} \\[2ex] \dfrac{\gamma\,(K\,(\beta - b\,\beta) + \delta)\,\lambda}{(-1+b)\,K\,\beta\,\gamma + \delta\,\lambda} & b\,\gamma & \dfrac{b\,\delta}{1-b} & \dfrac{\beta_v\,\gamma\,(K\,(\beta - b\,\beta) + \delta)\,\lambda}{\beta\,((-1+b)\,K\,\beta\,\gamma + \delta\,\lambda)} \\[2ex] 0 & 0 & 0 & \dfrac{\beta_z\,((-1+b)\,K\,\beta - \delta)\,\delta\,\lambda}{(-1+b)\,\beta\,((-1+b)\,K\,\beta\,\gamma + \delta\,\lambda)} \end{pmatrix}$$

Hopf bifurcation point related to b- for E- :

```
In[ ]:=   (*it takes few minutes to run*)
          cn=Join[cF1];
          pcm=Collect[Det[ψ IdentityMatrix[4]-(jEm1)],ψ];
          coT=CoefficientList[pcm,ψ];
          Length[coT]
          a1=coT[[4]] ;a2=coT[[3]];a3=coT[[2]] ; a4=coT[[1]];
          H4m=a1*a2*a3-a3^2+a1^2 a4//.cF1;
          ϕb4m=Collect[Numerator[Together[H4m]],b];
          cbm=NSolve[(ϕb4m//.cn)==0,b,WorkingPrecision→10];
          bMm=Max[Table[Re[b/.cbm[[i]]],{i,Length[cbm]}]];
          Print["bH-=",bHm=N[bMm,30]]
```

Out[ ]= 5

```
bH-=29.19440675
```

Ep1-2)Trace, Det and third criterion of Routh Hurwitz applied to E*:

Det and Trace of of E* and Analysis of the stability of E* in 4 dim when ϵ=1:

```
In[72]:=   Print["Tr[J[E*]]="]
           trEs1=Tr[jacD//.Join[sol[[3]],{z→0}]]//FullSimplify
           Print["Det[J[E*]]="]
           detEs1=Det[jacD//.Join[sol[[3]],{z→0}]]//FullSimplify

           pc=Collect[Det[ψ IdentityMatrix[4]-(jEs1)],ψ];
           coT=CoefficientList[pc,ψ]//FullSimplify;
           Length[coT]
           a1=coT[[4]]//FullSimplify ;a2=coT[[3]]//FullSimplify;a3=coT[[2]]//FullSimplify ; a4=coT[[1]]//FullSim
           Print["a₁=",a1, ", a₂=",a2, ", a₃=",a3, " ,a₄=", a4 ]
           H4=a1*a2*a3-a3^2+a1^2 a4;
           Print["H2(b0)=",H4/.b→b0//FullSimplify]
           Print["Denominator of H2 is ",Denominator[Together[H4]]//FullSimplify]
           ϕb4=Collect[Numerator[Together[H4]],b];
           cofi=CoefficientList[ϕb4,b];
           Length[cofi]
           (*Print["value of ϕ(b) at crit b is "]
            ϕb4/.b→b0//FullSimplify; (*so long expression*)*)
```

Tr[J[E*]]=

Out[73]= $-\dfrac{K\,\delta\,(2\times(-1+b)\,\beta\,\gamma+b\,\beta\,\delta+\beta_z\,\delta)\,\lambda+\delta^2\,\lambda^2+(-1+b)\,K^2\,\beta\,(\beta\,\gamma\,((-1+b)\,\gamma+b\,\delta)-\beta_z\,\delta\,\lambda)}{(-1+b)\,K\,\beta\,((-1+b)\,K\,\beta\,\gamma+\delta\,\lambda)}$

Det[J[E*]]=

Out[75]= $-\dfrac{\beta_z\,\gamma\,\delta^2\,(K\,(\beta-b\,\beta)+\delta)^2\,\lambda^2}{(-1+b)^2\,K\,\beta^2\,((-1+b)\,K\,\beta\,\gamma+\delta\,\lambda)}$

Out[78]= 5

$$a_1 = \frac{K \delta (2 \times (-1+b) \beta \gamma + b \beta \delta + \beta_z \delta) \lambda + \delta^2 \lambda^2 + (-1+b) K^2 \beta (\beta \gamma ((-1+b) \gamma + b \delta) - \beta_z \delta \lambda)}{(-1+b) K \beta ((-1+b) K \beta \gamma + \delta \lambda)}$$

$$, a_2 = \Big(\big(\delta \lambda \big(- \big((-1+b) K^2 \beta^2 ((-1+b) (\beta + \beta_z) \gamma + b \beta_z \delta) \big) + (b \beta + \beta_z) \delta^2 \lambda + $$
$$K \beta (\beta_z \delta ((-1+b) \gamma + b \delta + \lambda - b \lambda) + (-1+b) \beta \gamma ((-1+b) \gamma + \delta - \lambda + b (\delta + \lambda))) \big) \big) \Big) /$$
$$\Big((-1+b)^2 K \beta^2 ((-1+b) K \beta \gamma + \delta \lambda) \Big) \Big), a_3 = $$
$$\Big(\big(\big(((-1+b) K \beta - \delta) \delta \lambda \big(\delta^2 \big((-1+b)^2 \beta \gamma - b \beta_z \delta \big) \lambda^2 + (-1+b)^2 K^2 \beta^2 \gamma \big((-1+b)^2 \beta \gamma^2 + \beta_z \delta \lambda \big) + $$
$$(-1+b) K \beta \gamma \delta \lambda \big(2 (-1+b)^2 \beta \gamma - \beta_z ((-1+b) \gamma + \delta - \lambda + b (\delta + \lambda)) \big) \big) \big) \Big) /$$
$$\Big((-1+b)^3 K \beta^2 ((-1+b) K \beta \gamma + \delta \lambda)^2 \Big) \Big), a_4 = - \frac{\beta_z \gamma \delta^2 (K (\beta - b \beta) + \delta)^2 \lambda^2}{(-1+b)^2 K \beta^2 ((-1+b) K \beta \gamma + \delta \lambda)}$$

H2 (b0) =0

Denominator of H2 is $(-1+b)^6 K^3 \beta^5 ((-1+b) K \beta \gamma + \delta \lambda)^4$

Out[86]= 11

Numerical values ϵ=1:

```
In[87]:=  cn=Join[cF1]; cnb={b→40};
          cb=NSolve[(ϕb4//.cn)==0,b,WorkingPrecision→10]
          bM=Max[Table[Re[b/.cb〚i〛],{i,Length[cb]}]];
          Print["bH=",bH=N[bM,30]]
          Print["b0=",b0/.cn//N]

          Print["E*",Es1//.cn/.cnb//N]
          Print["E+=",Em1//.cn/.cnb//N]
          Print["Eim=",Eim1//.cn/.cnb//N]
          Print["roots of Dis[b]=0:", bcE1=NSolve[(Dis//.cn)==0,b]]
          bc1=Chop[Evaluate[b/.bcE1〚5〛]];
          bc2=Chop[Evaluate[b/.bcE1〚6〛]];
          Print["b1*=", bc1]
          Print[" b2*=", bc2]
```

Out[88]= $\{\{b \to -258.7317419\}, \{b \to -0.01615999598\},$
$\{b \to 0.060848063 - 10.686990738\, i\}, \{b \to 0.060848063 + 10.686990738\, i\},$
$\{b \to 0.8448282668 - 0.9299250641\, i\}, \{b \to 0.8448282668 + 0.9299250641\, i\},$
$\{b \to 0.8878046288\}, \{b \to 1.011494253\}, \{b \to 1.022996712\}, \{b \to 1.152364263\}\}$

bH=1.152364263

b0=1.01149

E*$\{0.000294724, 0.0363426, 0.0221463, 0.\}$

E+=$\{0.00262408 + 5.51068 \times 10^{-19}\, i, 0.0462057 + 8.67362 \times 10^{-18}\, i,$
$0.021866 + 3.02367 \times 10^{-18}\, i, 0.0462057 + 8.67362 \times 10^{-18}\, i\}$

Eim=$\{0.114678 - 9.26732 \times 10^{-17}\, i, 0.263221 - 2.77556 \times 10^{-17}\, i,$
$0.0143012 + 8.58446 \times 10^{-18}\, i, 0.263221 - 2.77556 \times 10^{-17}\, i\}$

roots of Dis[b]=0:
$\{\{b \to -126.518\}, \{b \to -63.\}, \{b \to -63.\}, \{b \to -24.5518\}, \{b \to 29.361\}, \{b \to 45.9232\}\}$

b1*=29.361

b2*=45.9232

Ep1-3)Bifurcation diagrams :

Numerical solution of the stability (Bifurcation diagram) wrt y:

In[100]:= `(*`**Checks on the stability of the fixed points**`**)`
`Print["Eigenvalues of E* when b=20 and when b=40, respectively "]`
`Eigenvalues[jEs1] //. cF1 /. b → 20 // N`
`Eigenvalues[jEs1] //. cF1 /. b → 40 // N`
`Print["Eigenvalues of E+  when b=40"]`
`Chop[Eigenvalues[jEp1 //. cF1 /. b → 20 // N]]`

`Print["Eigenvalues of Eim  when b=40"]`
`Chop[Eigenvalues[jEim1 //. cF1 /. b → 40 // N]]`
`Print["Eigenvalues of E- between b1* and b2* "]`
`Chop[Eigenvalues[jEm1 //. cF1 /. b → 35 // N]]`
`Print["Eigenvalues of E- between 0 and b1*"]`
`Chop[Eigenvalues[jEm1 //. cF1 /. b → 20 // N]]`

Eigenvalues of E* when b=20 and when b=40, respectively

Out[101]= $\{0.0718263, -0.586224, 0.0257451 - 0.0774375\,\dot{\iota}, 0.0257451 + 0.0774375\,\dot{\iota}\}$

Out[102]= $\{0.0363426, -0.555066, 0.017069 - 0.082122\,\dot{\iota}, 0.017069 + 0.082122\,\dot{\iota}\}$

Eigenvalues of E+ when b=40

Out[104]= $\{-36.8126, -0.800277, -0.073135 + 0.126319\,\dot{\iota}, -0.073135 - 0.126319\,\dot{\iota}\}$

Eigenvalues of Eim when b=40

Out[106]= $\{-6.50192, 0.307571, -0.103153 + 0.233849\,\dot{\iota}, -0.103153 - 0.233849\,\dot{\iota}\}$

Eigenvalues of E- between b1* and b2*

Out[108]= $\{-0.999601, 0.0879326 + 0.166504\,\dot{\iota}, 0.0879326 - 0.166504\,\dot{\iota}, -0.0384878\}$

Eigenvalues of E- between 0 and b1*

Out[110]= $\{-0.840551 - 1.00512\,\dot{\iota}, 0.242338 + 0.343832\,\dot{\iota},$
$0.0933864 - 0.157103\,\dot{\iota}, -0.0937105 - 0.0144354\,\dot{\iota}\}$

```
In[ ]:=  cut=cF1;bL=50;max=0.37;
         b0n=b0//.cn;
         b1=Chop[bbs//.cF1//N];
         Print["b0=",b0n//N, " ,b_b*=", b1, ", b1*=", bc1, " , b2*=", bc2]
         lin1=Line[{{ bc1,0},{ bc1,max}}];
         li1=Graphics[{Thick,Black,Dashed,lin1}];
         lin2=Line[{{ bc2,0},{ bc2,max}}];
         li2=Graphics[{Thick,Black,Dashed,lin2}];
         lin3=Line[{{ b0n,0},{ b0n,max}}];
         li3=Graphics[{Thick,Black,Dashed,lin3}];
         lin4=Line[{{ b1,0},{ b1,max}}];
         li4=Graphics[{Thick,Black,Dashed,lin4}];
         pyb=Plot[{yb}//.cut,{b,0,bL},PlotStyle→{Dashed,Thick,Cyan},
         PlotRange→{{0,200},{0,max}},PlotLegends→{"y_b "}];
         (*pym1n=Plot[{ym}//.cut,{b,0,bL},PlotStyle→{Green,Thick},PlotRange→All,PlotPoints→200,
         PlotLegends→{"E_ unstable"}];*)

         pym=Plot[{ym}//.cut,{b,0,bL},PlotStyle→{Green,Dashed,Thick},PlotRange→All,PlotPoints→180,
         PlotLegends→{"E_ unstable"}];
         pyK1=Plot[0,{b,0,b0n},PlotStyle→{Magenta},PlotRange→All,PlotLegends→{"E_K stable"}];
         pyK2=Plot[0,{b,b0n,bL},PlotStyle→{Blue},PlotRange→All,PlotLegends→{"E_K unstable"}];
         pyi=Plot[{yi}//.cut,{b,0, bL},PlotStyle→{Red,Dashed,Thick},PlotRange→All,
         PlotLegends→{"E_im unstable"}];
         pyp=Plot[{yp}//.cut,{b,b0n,bL},PlotStyle→{Purple, Thick},PlotRange→All,
         PlotLegends→{"E_+ stable"}];
         pypn=Plot[{yp}//.cut,{b,0,b0n},PlotStyle→{Yellow, Thick},PlotRange→All,
         PlotLegends→{"E_+ (y<0)"}];
         N[{yp}//.cut/.b→40,20](*check*)
         Print["Q(yp) at b0 is "]
         Qy/.y→yp/.b→b0//FullSimplify
         Show[pyp,pyb,li3,li1,li2,li4];
         pys1=Plot[{ys}//.cut,{b,0,b1},PlotStyle→{Orange, Dotted},
         PlotRange→{{0,200},{0,max}},
         PlotLegends→{"E_* outside domain"}];
         pys2=Plot[{ys}//.cut,{b,b1,bL},PlotStyle→{Orange,Thick,Dashed},
         PlotRange→{{0,200},{0,max}},
         PlotLegends→{"E_* unstable"}];


         Print["y*'(b0)=",D[ys,b]/.b→b0n//.cut//N//FullSimplify]
         Chop[ys/.b→b0n//.cut//N](*Check*);
         bifep1=Show[{pys2,pys1,pyi,pyp,pypn,pyK1,pyK2,pym,pyb,li3,li1,li2,li4},
         Epilog→{Text["b_0",Offset[{-2,10},{ b0n//.cut,0}]],{PointSize[Large],
         Style[Point[{ b0n//.cut,0}],Black]},
         Text["b_1*",Offset[{-8,10},{ bc1,0}]],{PointSize[Large],
         Style[Point[{ bc1,0}],Purple]},
         Text["b_2*",Offset[{10,10},{ bc2,0}]],{PointSize[Large],
         Style[Point[{ bc2,0}],Yellow]},Text["b_b*",Offset[{10,10},{ b1,0}]],{PointSize[Large],
         Style[Point[{ b1,0}],Magenta]}},AxesLabel→{"b","y"},PlotRange→{{-0.2,bL},{-0.1,max}}]
         Export["EriB.pdf",bifep1]
```

b0=1.01149 ,b_b*=14.0011, b1*=29.361 , b2*=45.9232

Out[ ]= $\{0.29448140956728204616 + 0. \times 10^{-21} \, \mathbb{i}\}$

```
Q(yp) at b0 is
```

Out[ ]= `0`

```
y*'(b0)=86.3256
```


Out[ ]= `EriB.pdf`

(x,b)-Bifurcation diagram:

Determination of the endemic points with respect to x when $\epsilon=1$:

In[ ]:= **Solve[((y1) /. vex /. y → yex) == 0, z]**

Out[ ]=
$$\left\{\left\{z \to \frac{-c\,\gamma - \frac{c\,x\,\lambda}{K} - \beta_z\,\sqrt{\frac{4\,c\,\beta_y\,\left(x\,\lambda - \frac{x^2\,\lambda}{K}\right)}{\beta_z} + \left(-\frac{c\,\gamma}{\beta_z} - \frac{c\,x\,\lambda}{K\,\beta_z}\right)^2}}{2\,c\,\beta_y}\right\},\right.$$

$$\left.\left\{z \to \frac{-c\,\gamma - \frac{c\,x\,\lambda}{K} + \beta_z\,\sqrt{\frac{4\,c\,\beta_y\,\left(x\,\lambda - \frac{x^2\,\lambda}{K}\right)}{\beta_z} + \left(-\frac{c\,\gamma}{\beta_z} - \frac{c\,x\,\lambda}{K\,\beta_z}\right)^2}}{2\,c\,\beta_y}\right\}\right\}$$

In[372]:=
```
yex=c z/βz;(*Frome Solve[(z1/z/.cep1)==0,y]*)
Print["ye(x)=",yex]
Print["ve(x)="]
vex=Solve[(x1/x)==0,v]〚1〛
Print["ze(x)="]
zex=Solve[((y1/y)/.vex/.y→yex)==0,z]〚1〛//FullSimplify
(*the first solution of z above doesn't belong to the domain*)
vnn=(v1/v/.vex/.y→yex/.zex );
vnnn=vnn/.cF1
xex=Solve[vnnn==0,x,Cubics→False];
(*or // ComplexExpand[#, TargetFunctions → {Re, Im}] &*)
(*so long time when it's not numeric**)

Print["Number of endemic x"]
Length[xex]
Print["Numerical check"]
xex/.b→40//N
```

$$ye(x) = \frac{c\,z}{\beta_z}$$

$$ve(x) =$$

Out[375]= $$\left\{ v \to \frac{(K - x - y)\,\lambda}{K\,\beta} \right\}$$

$$ze(x) =$$

Out[377]= $$\left\{ z \to \frac{-c\,(K\,\gamma + x\,\lambda) + \sqrt{c\left(4\,K\,(K-x)\,x\,\beta_y\,\beta_z\,\lambda + c\,(K\,\gamma + x\,\lambda)^2\right)}}{2\,c\,K\,\beta_y} \right\}$$

Out[379]= $$\left(87 \times \left(\frac{1}{256}\,b\left(-\frac{1}{128} - x + \sqrt{4 \times (1-x)\,x + \left(\frac{1}{128} + x\right)^2} \right) - \right.\right.$$

$$x\left(1 - x + \frac{1}{2} \times \left(\frac{1}{128} + x - \sqrt{4 \times (1-x)\,x + \left(\frac{1}{128} + x\right)^2} \right) \right) -$$

$$\frac{1}{87} \times \left(-\frac{1}{128} - x + \sqrt{4 \times (1-x)\,x + \left(\frac{1}{128} + x\right)^2} \right) \times$$

$$\left(1 - x + \frac{1}{2} \times \left(\frac{1}{128} + x - \sqrt{4 \times (1-x)\,x + \left(\frac{1}{128} + x\right)^2} \right) \right) +$$

$$\left. \frac{1}{87} \times \left(-1 + x + \frac{1}{2} \times \left(-\frac{1}{128} - x + \sqrt{4 \times (1-x)\,x + \left(\frac{1}{128} + x\right)^2} \right) \right) \right) \right) \Bigg/$$

$$\left(2 \times \left(1 - x + \frac{1}{2} \times \left(\frac{1}{128} + x - \sqrt{4 \times (1-x)\,x + \left(\frac{1}{128} + x\right)^2} \right) \right) \right)$$

⋯ Solve: Solutions may not be valid for all values of parameters.

Number of endemic x

Out[382]= 3

Numerical check

Out[384]= $\{\{x \to 0.00262408\}, \{x \to 0.114678\}, \{x \to 0.540959\}\}$

In[385]:= **(∗Jacobians of the fixed points∗∗)**
jEsx = jacD /. cep1 /. sol⟦3⟧ /. z → 0;
jEmx = jacD /. cep1 /. vex /. y → yex /. zex /. xex⟦1⟧;
jEimx = jacD /. cep1 /. vex /. y → yex /. zex /. xex⟦2⟧;
jEpx = jacD /. cep1 /. vex /. y → yex /. zex /. xex⟦3⟧;
(∗Checks on the stability of the fixed points∗∗)
Print["Eigenvalues of E∗ : "]
Print[" between b0 and b1∗ "]
Eigenvalues[jEsx] //. cF1 /. b → 15 // N
Print[" between b1∗ and b2∗ "]
Eigenvalues[jEsx] //. cF1 /. b → 40 // N
Print["Eigenvalues of E+ between b1∗ and b2∗"]
Chop[Eigenvalues[jEpx //. cF1 /. b → 42 // N]]
Print["Eigenvalues of Eim between b1∗ and b2∗"]
Chop[Eigenvalues[jEimx //. cF1 /. b → 35 // N]]
Print["Eigenvalues of E− between b0 and b1∗ "]
Chop[Eigenvalues[jEmx //. cF1 /. b → 15 // N]]
Print["Eigenvalues of E− between b1∗ and b2∗"]
Chop[Eigenvalues[jEimx //. cF1 /. b → 40 // N]]
Print["Eigenvalues of E− after b2∗"]
Chop[Eigenvalues[jEimx //. cF1 /. b → 50 // N]]

Eigenvalues of E∗ :

 between b0 and b1∗

Out[391]= $\{0.0950185, -0.606269, 0.0309604 - 0.074022\, \mathbb{i}, 0.0309604 + 0.074022\, \mathbb{i}\}$

 between b1∗ and b2∗

Out[393]= $\{0.0363426, -0.555066, 0.017069 - 0.082122\, \mathbb{i}, 0.017069 + 0.082122\, \mathbb{i}\}$

Eigenvalues of E+ between b1∗ and b2∗

Out[395]= $\{-22.81, -0.157506 + 0.275785\, \mathbb{i}, -0.157506 - 0.275785\, \mathbb{i}, -0.301641\}$

Eigenvalues of Eim between b1∗ and b2∗

Out[397]= $\{-4.10357, 0.346228, -0.0664887 + 0.180685\, \mathbb{i}, -0.0664887 - 0.180685\, \mathbb{i}\}$

Eigenvalues of E− between b0 and b1∗

Out[399]= $\{-38.9444, -0.864225, -0.053898 + 0.0931984\, \mathbb{i}, -0.053898 - 0.0931984\, \mathbb{i}\}$

Eigenvalues of E− between b1∗ and b2∗

Out[401]= $\{-6.50192, 0.307571, -0.103153 + 0.233849\, \mathbb{i}, -0.103153 - 0.233849\, \mathbb{i}\}$

Eigenvalues of E− after b2∗

Out[403]= $\{-14.0734 + 7.77106\, \mathbb{i}, -0.136004 + 0.346012\, \mathbb{i},$
$-0.181515 - 0.313963\, \mathbb{i}, -0.00338735 + 0.318699\, \mathbb{i}\}$

In[477]:=
```
cut=cF1;bL=49;max=1.1;
b0n=b0//.cn;
b1=Chop[b/.Solve[(Es1[[1]])==(1-yb),b][[2]]//.cF1//N]
Print["bH=",bH, " ,b1*=",bc1, " ,b2*=",bc2]
lin1=Line[{{ bc1,0},{ bc1,max}}];
li1=Graphics[{Thick,Black,Dashed,lin1}];
lin2=Line[{{ bc2,0},{ bc2,max}}];
li2=Graphics[{Thick,Black,Dashed,lin2}];
lin3=Line[{{ b0n,0},{ b0n,max}}];
li3=Graphics[{Thick,Black,Dashed,lin3}];
lin4=Line[{{ b1,0},{ b1,max}}];
li4=Graphics[{Thick,Black,Dashed,lin4}];
pxp=Plot[{x/.xex[[3]]}//.cut,{b,0,bL},PlotStyle→{Purple,Thick},PlotRange→All,PlotPoints→180,
PlotLegends→{"E₊ stable"}]
pxK1=Plot[K//.cut,{b,0,b0n},PlotStyle→{Magenta},PlotRange→All,PlotLegends→{"E_K stable"}];
pxK2=Plot[K//.cut,{b,b0n,bL},PlotStyle→{Blue},PlotRange→All,PlotLegends→{"E_K unstable"}];
pxi=Plot[{x/.xex[[2]]}//.cut,{b,b0n, bL},PlotStyle→{Red, Dashed,Thick},PlotRange→All,
PlotLegends→{"E_im unstable"}]
pxm0=Plot[{x/.xex[[1]]}//.cut,{b,0,b0n},PlotStyle→{Yellow,  Thick},PlotRange→All,
PlotLegends→{"E₋ outside domain "}]
pxm=Plot[{x/.xex[[1]]}//.cut,{b,b0n,bc1},PlotStyle→{Green,  Thick},PlotRange→All,
PlotLegends→{"E₋ stable"}]
pxm1=Plot[{x/.xex[[1]]}//.cut,{b,bc2,bL},PlotStyle→{Green, Thick},PlotRange→All,PlotPoints→40]
pxm2=Plot[{x/.xex[[1]]}//.cut,{b,bc1,bc2},PlotStyle→{Green,Dashed,  Thick},PlotRange→All,
PlotLegends→{"E₋ unstable"}]

pxs=Plot[{x/.sol[[3]]}//.cut,{b,0,bL},PlotStyle→{Orange,Thick,Dashed},
PlotRange→{{0,bL},{0,max}},
PlotLegends→{"E* unstable"}]

pxb=Plot[{1-yb}//.cut,{b,0,bL},PlotStyle→{Dashed,Thick,Cyan},
PlotRange→{{0,bL},{0,max}},PlotLegends→{"1-y_b "}];

bifep1=Show[{pxs,pxi,pxp,pxK1,pxK2,pxm,pxm1,pxm2,(*pxm0,*)pxb,li3,li1,li2,li4},
Epilog→{Text["b₀",Offset[{10,10},{ b0n//.cut,0}]],{PointSize[Large],
Style[Point[{ b0n//.cut,0}],Black]},
Text["b₁*",Offset[{-8,10},{ bc1,0}]],{PointSize[Large],
Style[Point[{ bc1,0}],Purple]},
Text["b₂*",Offset[{10,10},{ bc2,0}]],{PointSize[Large],
Style[Point[{ bc2,0}],Yellow]},Text["b_b*",Offset[{0,10},{ b1,0}]],{PointSize[Large],
Style[Point[{ b1,0}],Magenta]}},AxesLabel→{"b","x"},PlotRange→{{-0.1,bL},{-0.1,max}}]
Export["Bif1x.pdf",bifep1]
```

Out[479]= 128.989

bH=1.152364263 ,b1*=29.361 ,b2*=45.9232

Out[489]=

E_+ stable

Out[492]=

E_{im} unstable

Out[493]=

E_- outside domain

Out[494]=

E_- stable

Out[495]=

Out[496]=

- - - - E_- unstable

Out[497]=

- - - - E_* unstable

Out[499]=

- - - - E_* unstable

- - - - E_{im} unstable

—— E_+ stable

—— E_K stable

—— E_K unstable

—— E_- stable

- - - - E_- unstable

- - - - $1-y_b$

Out[500]= `Bif1x.pdf`

```
In[ ]:= (*Some checks*)
     Print["between b0 and b1*; E-="]
     {x /. xex〚2〛, yex〚2〛 /. zex /. xex〚2〛,
         v /. vex /. y → yex /. (zex) /. xex〚2〛, z /. zex /. xex〚2〛} //. cut /. b → 15 // N
     Print["with yb="]
     yb //. cut /. b → 15 // N
     Print["between b1* and b2*; E-="]
     {x /. xex〚2〛, yex〚2〛 /. zex /. xex〚2〛,
         v /. vex /. y → yex /. (zex) /. xex〚2〛, z /. zex /. xex〚2〛} //. cut /. b → 35 // N
     Print[" after b2*; E-="]
     {x /. xex〚2〛, yex〚2〛 /. zex /. xex〚2〛,
         v /. vex /. y → yex /. (zex) /. xex〚2〛, z /. zex /. xex〚2〛} //. cut /. b → 47 // N
     Print["between  b0 and  b1*; E*="]
     sol〚3〛 //. cut /. b → 15 // N
     Print[" between b1* and  b2*; E*="]
     sol〚3〛 //. cut /. b → 35 // N
     Print["After  b2*; E*="]
     sol〚3〛 //. cut /. b → 47 // N
```

between b0 and b1*; E-=

Out[ ]= {0.878336, 0.107541, 0.000324678, 0.107541}

with yb=

Out[ ]= 0.109375

between b1* and b2*; E-=

Out[ ]= {0.00405931, 0.0579238, 0.0215636, 0.0579238}

 after b2*; E-=

Out[ ]= {0.0017057, 0.0367794, 0.0221038, 0.0367794}

between b0 and b1*; E*=

Out[ ]= {x → 0.000821018, y → 0.0950185, v → 0.0207853}

 between b1* and b2*; E*=

Out[ ]= {x → 0.000338066, y → 0.0414636, v → 0.0220275}

After b2*; E*=

Out[ ]= {x → 0.000249875, y → 0.030985, v → 0.0222705}

```
(*Endemic x using the elimination**)
elx = Eliminate[
    Thread[({x1 / x, y1 / y, v1, z1 / z} /. cep1 /. cKga1) == {0, 0, 0, 0}], {y, v, z}];
Qx = (elx[[1]] - elx[[2]]);
Print["Coefficients of  Qx by elim polynomial are:"]
cofx = CoefficientList[Qx, x] // FullSimplify;
Length[cofx]
solx = x /. Solve[Qx == 0, x, Cubics → False] (*Third order roots*)
solx //. cF1 /. b → 40 // N
```

Coefficients of Qx by elim polynomial are:

Out[○]= 4

Out[○]= $\{ \text{Root}[-b\,c^2\,\beta\,\beta_y\,\delta + c^2\,\beta_v\,\delta\,\lambda + c\,\beta_v\,\beta_y\,\beta_z\,\delta\,\lambda - c^2\,\beta_y\,\delta^2\,\lambda -$
$c\,\beta_y^2\,\beta_z\,\delta^2\,\lambda + (-b\,c^2\,\beta^2\,\beta_y + b^2\,c^2\,\beta^2\,\beta_y + c^2\,\beta\,\beta_v\,\lambda - b\,c^2\,\beta\,\beta_v\,\lambda + c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda -$
$2\,b\,c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda - 2\,c^2\,\beta\,\beta_y\,\delta\,\lambda + b\,c^2\,\beta\,\beta_y\,\delta\,\lambda - c\,\beta_v\,\beta_y\,\beta_z\,\delta\,\lambda - 2\,c\,\beta\,\beta_y^2\,\beta_z\,\delta\,\lambda +$
$c\,\beta_y^2\,\beta_z\,\delta^2\,\lambda + c\,\beta_v^2\,\beta_z\,\lambda^2 + \beta_v^2\,\beta_y\,\beta_z^2\,\lambda^2 + c^2\,\beta_v\,\delta\,\lambda^2 + c\,\beta_v\,\beta_y\,\beta_z\,\delta\,\lambda^2)\, \#1 +$
$(-c^2\,\beta^2\,\beta_y\,\lambda + b\,c^2\,\beta^2\,\beta_y\,\lambda - c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda + 2\,b\,c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda - c\,\beta^2\,\beta_y^2\,\beta_z\,\lambda + 2\,c\,\beta\,\beta_y^2\,\beta_z\,\delta\,\lambda +$
$c^2\,\beta\,\beta_v\,\lambda^2 - b\,c^2\,\beta\,\beta_v\,\lambda^2 - c\,\beta_v^2\,\beta_z\,\lambda^2 + c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda^2 - 2\,\beta_v^2\,\beta_y\,\beta_z^2\,\lambda^2 - c\,\beta_v\,\beta_y\,\beta_z\,\delta\,\lambda^2)\, \#1^2 +$
$(c\,\beta^2\,\beta_y^2\,\beta_z\,\lambda - c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda^2 + \beta_v^2\,\beta_y\,\beta_z^2\,\lambda^2)\, \#1^3\, \&, 1],$
$\text{Root}[-b\,c^2\,\beta\,\beta_y\,\delta + c^2\,\beta_v\,\delta\,\lambda + c\,\beta_v\,\beta_y\,\beta_z\,\delta\,\lambda - c^2\,\beta_y\,\delta^2\,\lambda - c\,\beta_y^2\,\beta_z\,\delta^2\,\lambda +$
$(-b\,c^2\,\beta^2\,\beta_y + b^2\,c^2\,\beta^2\,\beta_y + c^2\,\beta\,\beta_v\,\lambda - b\,c^2\,\beta\,\beta_v\,\lambda + c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda -$
$2\,b\,c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda - 2\,c^2\,\beta\,\beta_y\,\delta\,\lambda + b\,c^2\,\beta\,\beta_y\,\delta\,\lambda - c\,\beta_v\,\beta_y\,\beta_z\,\delta\,\lambda - 2\,c\,\beta\,\beta_y^2\,\beta_z\,\delta\,\lambda +$
$c\,\beta_y^2\,\beta_z\,\delta^2\,\lambda + c\,\beta_v^2\,\beta_z\,\lambda^2 + \beta_v^2\,\beta_y\,\beta_z^2\,\lambda^2 + c^2\,\beta_v\,\delta\,\lambda^2 + c\,\beta_v\,\beta_y\,\beta_z\,\delta\,\lambda^2)\, \#1 +$
$(-c^2\,\beta^2\,\beta_y\,\lambda + b\,c^2\,\beta^2\,\beta_y\,\lambda - c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda + 2\,b\,c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda - c\,\beta^2\,\beta_y^2\,\beta_z\,\lambda + 2\,c\,\beta\,\beta_y^2\,\beta_z\,\delta\,\lambda +$
$c^2\,\beta\,\beta_v\,\lambda^2 - b\,c^2\,\beta\,\beta_v\,\lambda^2 - c\,\beta_v^2\,\beta_z\,\lambda^2 + c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda^2 - 2\,\beta_v^2\,\beta_y\,\beta_z^2\,\lambda^2 - c\,\beta_v\,\beta_y\,\beta_z\,\delta\,\lambda^2)\, \#1^2 +$
$(c\,\beta^2\,\beta_y^2\,\beta_z\,\lambda - c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda^2 + \beta_v^2\,\beta_y\,\beta_z^2\,\lambda^2)\, \#1^3\, \&, 2],$
$\text{Root}[-b\,c^2\,\beta\,\beta_y\,\delta + c^2\,\beta_v\,\delta\,\lambda + c\,\beta_v\,\beta_y\,\beta_z\,\delta\,\lambda - c^2\,\beta_y\,\delta^2\,\lambda - c\,\beta_y^2\,\beta_z\,\delta^2\,\lambda +$
$(-b\,c^2\,\beta^2\,\beta_y + b^2\,c^2\,\beta^2\,\beta_y + c^2\,\beta\,\beta_v\,\lambda - b\,c^2\,\beta\,\beta_v\,\lambda + c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda -$
$2\,b\,c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda - 2\,c^2\,\beta\,\beta_y\,\delta\,\lambda + b\,c^2\,\beta\,\beta_y\,\delta\,\lambda - c\,\beta_v\,\beta_y\,\beta_z\,\delta\,\lambda - 2\,c\,\beta\,\beta_y^2\,\beta_z\,\delta\,\lambda +$
$c\,\beta_y^2\,\beta_z\,\delta^2\,\lambda + c\,\beta_v^2\,\beta_z\,\lambda^2 + \beta_v^2\,\beta_y\,\beta_z^2\,\lambda^2 + c^2\,\beta_v\,\delta\,\lambda^2 + c\,\beta_v\,\beta_y\,\beta_z\,\delta\,\lambda^2)\, \#1 +$
$(-c^2\,\beta^2\,\beta_y\,\lambda + b\,c^2\,\beta^2\,\beta_y\,\lambda - c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda + 2\,b\,c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda - c\,\beta^2\,\beta_y^2\,\beta_z\,\lambda + 2\,c\,\beta\,\beta_y^2\,\beta_z\,\delta\,\lambda +$
$c^2\,\beta\,\beta_v\,\lambda^2 - b\,c^2\,\beta\,\beta_v\,\lambda^2 - c\,\beta_v^2\,\beta_z\,\lambda^2 + c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda^2 - 2\,\beta_v^2\,\beta_y\,\beta_z^2\,\lambda^2 - c\,\beta_v\,\beta_y\,\beta_z\,\delta\,\lambda^2)\, \#1^2 +$
$(c\,\beta^2\,\beta_y^2\,\beta_z\,\lambda - c\,\beta\,\beta_v\,\beta_y\,\beta_z\,\lambda^2 + \beta_v^2\,\beta_y\,\beta_z^2\,\lambda^2)\, \#1^3\, \&, 3] \}$

Out[○]= $\{0.000294987, -19.9296 - 34.5878\,\mathbb{i}, -19.9296 + 34.5878\,\mathbb{i}\}$