On a four-dimensional oncolytic Virotherapy model (ϵ=0)

This Mathematica Notebook is a supplementary material to the paper "On a three-dimensional and two four-dimensional
oncolytic viro-therapy models". It contains some of the calculations and illustrations appearing in the paper concerning the section on 4 dimensional model.

Section 3 : Four-dim viro-therapy model with Logistic growth when ϵ>0 and ϵ=0

```
In[72]:= SetDirectory[NotebookDirectory[]];
AppendTo[$Path, Directory];
Clear["Global`*"];
Clear["K"];
(*Some aliases*)
Format[βy] = Subscript[β, y];
Format[βv] = Subscript[β, v];
Format[βz] = Subscript[β, z];


Unprotect[Power];
Power[0, 0] = 1;
Protect[Power];

par = {b, β, λ, δ, βy, βv, βz, c, γ, K, ϵ};
cp = Join[Thread[Drop[par, {1}] > 0], {b > 1}];
cKga1 = {K → 1, γ → 1};
cep1 = {ϵ → 1}; cep0 = {ϵ → 0};
R0 = b β K / (β K + δ) (* Reproduction number*);
cnT17 = {βv → 0.16, βy → 0.48, K → 1,
    γ → 1, b → 9, β → 0.11, λ → 0.36, δ → 0.2, βz → 0.6, c → 0.036};
(*Numerical values of Tian 17 after dimensionalization*)

(****** Four dim Deterministic  epidemic model with Logistic growth ****)
x1 = λ x (1 - (x + y) / K) -  β x v ;
y1 = β x v - βy y z - γ y;
v1 = -β x v - βv v z + b γ y - δ v;
z1 = z (βz y - c z^ϵ);
ye = c / βz; vM = λ (1 - ye) / β; vMN = vM /. cnT17;
dyn = {x1, y1, v1, z1};
dynS = dyn //. cKga1;
```

```mathematica
dyn3 = {x1, y1, v1} /. z → 0; (*3dim case used for E* *)
dyn3S = dyn3 //. cKga1;
          x'
          y'
Print[" (   )=", dyn // FullSimplify // MatrixForm,
          v'
          z'

                                    x'
                                    y'
   "  the reparametrized model is (   )=",
                                    v'
                                    z'

 dynS // FullSimplify // MatrixForm]
Print["b0=", b0 = b /. Apart[Solve[R0 == 1, b]〚1〛 // FullSimplify]]
b0S = b0 //. cKga1;
(*****Fixed points of Tian17 using the scaling when K=1, γ=1***)

fv = (ye (b - 1) - v δ); gv = (ye βy + v βv); hv = (1 - ye - v β / λ);

Print["xe= ", xe = hv, " , ye=", ye, " , ze =", ze = fv / gv]

Pv = Numerator[Together[v β xe - ye (1 + βy fv / gv)]] / (-βz² β² βv);
Print["P(v)=", pc = Collect[Together[Pv], v], " coefs are"]
coP = CoefficientList[pc, v] // Simplify

(***Fixed point when z→0**)
eq = Thread[dyn3 == {0, 0, 0}];
sol = Solve[eq, {x, y, v}] // FullSimplify;
Es = {x, y, v} /. sol〚3〛; (*Endemic point with z=0*);
Print[" Endemic point with z=0 is E*=", Es // FullSimplify]
EsS = Es /. cKga1 // FullSimplify(* E* when K=γ=1***)
Print["Check: when K=γ=1, E*=", EsS]

bn = b /. Solve[EsS〚2〛 == ye, b]; bnn = bn /. cnT17;
Print["y*/ye="]
EsS〚2〛 / ye // FullSimplify
Print["(b1,b2)="]
bn // FullSimplify

bcn = b /. Solve[EsS〚2〛〚1〛 == ye, b];
bcnn = bn /. cnT17;

Dis = Chop[Collect[Discriminant[Numerator[Pv], v], b]];
solb = Solve[Dis == 0, b];
jacS = Grad[dynS, {x, y, v, z}] // FullSimplify;
Print["Jacobian"]
jacS // MatrixForm
Print["Jacobian at E_K "]
jacK = FullSimplify[jacS /. Join[sol〚2〛, {z → 0}], Assumptions → ε ≥ 0];
jacK // MatrixForm
Print["Jacobian at E_* when K>1"]
```

```
jacEs = FullSimplify[jacS /. Join[sol[[3]], {z → 0}], Assumptions → ϵ ≥ 0] (*when K>0*);
jacEs // MatrixForm

PR = Solve[Pv == 0, v, Cubics → False]
  (*//ComplexExpand[#,TargetFunctions→{Re,Im}]&*);
vn = v /. PR[[1]]; vp = v /. PR[[3]]; vi = v /. PR[[2]];
Ep = {xe, ye, v, ze} /. v → vp;
Ei = {xe, ye, v, ze} /. v → vi;

JEi = jacS //. {x → xe, y → ye, v → vi, z → ze};
JEp = jacS //. {x → xe, y → ye, v → vp, z → ze};
JEm = jacS //. {x → xe, y → ye, v → vn, z → ze};
jacEs1 = jacEs /. cKga1;
Print["Tr[J[E*]] corresponding to 4 dim model when K=γ=1 is "]
Tr[jacEs1]
Print["Tr[J[E*]] >0 iff"]
Reduce[Join[{(Tr[jacEs1] /. ϵ → 0) > 0}, Drop[cp, {9, 10}]],
  Drop[par, {10, 11}]] // FullSimplify
bc2 = b /. solb[[1]] // FullSimplify;
bc1 = b /. solb[[2]] // FullSimplify;
Print["b1*=", bc1]
Print[" b2*=", bc2]
```

$$\left(*\left(-1 + \frac{b\,\delta}{1-b} - c - \frac{\delta\,\lambda}{(-1+b)\,\beta} + \frac{\delta}{(-1+b)\,\beta}\left(\frac{b\,\lambda(Rz-1)\,\beta z}{\lambda(b-Rz) + (-1+b)\,\gamma\,Rz}\right)\right)*\right)$$

$$\begin{pmatrix} x' \\ y' \\ v' \\ z' \end{pmatrix} = \begin{pmatrix} -v\,x\,\beta + x\left(1 - \frac{x+y}{K}\right)\lambda \\ v\,x\,\beta - y\left(z\,\beta_y + \gamma\right) \\ b\,y\,\gamma - v\left(x\,\beta + z\,\beta_v + \delta\right) \\ z\left(-c\,z^\epsilon + y\,\beta_z\right) \end{pmatrix}$$

the reparametrized model is

$$\begin{pmatrix} x' \\ y' \\ v' \\ z' \end{pmatrix} = \begin{pmatrix} -x\left(v\,\beta + (-1+x+y)\,\lambda\right) \\ v\,x\,\beta - y\left(1 + z\,\beta_y\right) \\ b\,y - v\left(x\,\beta + z\,\beta_v + \delta\right) \\ z\left(-c\,z^\epsilon + y\,\beta_z\right) \end{pmatrix}$$

$$b0 = 1 + \frac{\delta}{K\,\beta}$$

$$xe = 1 - \frac{c}{\beta_z} - \frac{v\,\beta}{\lambda} \quad,\quad ye = \frac{c}{\beta_z} \quad,\quad ze = \frac{\frac{(-1+b)\,c}{\beta_z} - v\,\delta}{v\,\beta_v + \frac{c\,\beta_y}{\beta_z}}$$

$$P(v) = v^3 + \frac{b\,c^2\,\beta_y\,\lambda}{\beta^2\,\beta_v\,\beta_z^2} + \frac{v^2\left(c\,\beta^2\,\beta_y\,\beta_z + c\,\beta\,\beta_v\,\beta_z\,\lambda - \beta\,\beta_v\,\beta_z^2\,\lambda\right)}{\beta^2\,\beta_v\,\beta_z^2} +$$

$$\frac{v\left(c^2\,\beta\,\beta_y\,\lambda + c\,\beta_v\,\beta_z\,\lambda - c\,\beta\,\beta_y\,\beta_z\,\lambda - c\,\beta_y\,\beta_z\,\delta\,\lambda\right)}{\beta^2\,\beta_v\,\beta_z^2} \quad \text{coefs are}$$

Out[102]= $\left\{\dfrac{b\,c^2\,\beta_y\,\lambda}{\beta^2\,\beta_v\,\beta_z^2} \,,\, \dfrac{c\left(c\,\beta\,\beta_y + \beta_z\left(\beta_v - \beta_y\,(\beta+\delta)\right)\right)\lambda}{\beta^2\,\beta_v\,\beta_z^2} \,,\, \dfrac{c\,\beta\,\beta_y + c\,\beta_v\,\lambda - \beta_v\,\beta_z\,\lambda}{\beta\,\beta_v\,\beta_z} \,,\, 1\right\}$

Endemic point with z=0 is E*=

$\left\{\dfrac{\delta}{(-1+b)\,\beta} \,,\, \dfrac{((-1+b)\,K\,\beta - \delta)\,\delta\,\lambda}{(-1+b)\,\beta\,((-1+b)\,K\,\beta\,\gamma + \delta\,\lambda)} \,,\, \dfrac{\gamma\left((-1+b)\,K\,\beta - \delta\right)\lambda}{\beta\,((-1+b)\,K\,\beta\,\gamma + \delta\,\lambda)}\right\}$

Out[107]= $\left\{\dfrac{\delta}{(-1+b)\,\beta} \,,\, \dfrac{((-1+b)\,\beta - \delta)\,\delta\,\lambda}{(-1+b)\,\beta\,((-1+b)\,\beta + \delta\,\lambda)} \,,\, \dfrac{((-1+b)\,\beta - \delta)\,\lambda}{\beta\,((-1+b)\,\beta + \delta\,\lambda)}\right\}$

Check: when K=γ=1, E$*$=$\left\{ \dfrac{\delta}{(-1+b)\,\beta}, \dfrac{((-1+b)\,\beta-\delta)\,\delta\,\lambda}{(-1+b)\,\beta\,((-1+b)\,\beta+\delta\,\lambda)}, \dfrac{((-1+b)\,\beta-\delta)\,\lambda}{\beta\,((-1+b)\,\beta+\delta\,\lambda)} \right\}$

y$*$/ye=

Out[111]= $\dfrac{\beta_z\,((-1+b)\,\beta-\delta)\,\delta\,\lambda}{(-1+b)\,c\,\beta\,((-1+b)\,\beta+\delta\,\lambda)}$

$\langle$b1,b2$\rangle$ =

Out[113]= $\left\{ \dfrac{2\,c\,\beta - c\,\delta\,\lambda + \beta_z\,\delta\,\lambda - \delta\,\sqrt{\lambda}\,\sqrt{-4\,c\,\beta_z+(c-\beta_z)^2\,\lambda}}{2\,c\,\beta}, \right.$

$\left. \dfrac{2\,c\,\beta - c\,\delta\,\lambda + \beta_z\,\delta\,\lambda + \delta\,\sqrt{\lambda}\,\sqrt{-4\,c\,\beta_z+(c-\beta_z)^2\,\lambda}}{2\,c\,\beta} \right\}$

Jacobian

Out[120]//MatrixForm=
$$\begin{pmatrix} -v\,\beta - (-1+2\,x+y)\,\lambda & -x\,\lambda & -x\,\beta & 0 \\ v\,\beta & -1-z\,\beta_y & x\,\beta & -y\,\beta_y \\ -v\,\beta & b & -x\,\beta - z\,\beta_v - \delta & -v\,\beta_v \\ 0 & z\,\beta_z & 0 & y\,\beta_z - c\,z^\in\,(1+\in) \end{pmatrix}$$

Jacobian at E_K

Out[123]//MatrixForm=
$$\begin{pmatrix} \lambda - 2\,K\,\lambda & -K\,\lambda & -K\,\beta & 0 \\ 0 & -1 & K\,\beta & 0 \\ 0 & b & -K\,\beta - \delta & 0 \\ 0 & 0 & 0 & -0^\in\,c\,(1+\in) \end{pmatrix}$$

Jacobian at E_$*$ when K>1

Out[126]//MatrixForm=
$$\begin{pmatrix} -\dfrac{\delta\,\lambda\,(\delta\,\lambda+(-1+b)\,\beta\,((-1+2\,K)\,\gamma+(-1+K)\,\lambda))}{(-1+b)\,\beta\,((-1+b)\,K\,\beta\,\gamma+\delta\,\lambda)} & \dfrac{\delta\,\lambda}{\beta-b\,\beta} & -\dfrac{\delta}{-1+b} & 0 \\ \dfrac{\gamma\,((-1+b)\,K\,\beta-\delta)\,\lambda}{(-1+b)\,K\,\beta\,\gamma+\delta\,\lambda} & -1 & \dfrac{\delta}{-1+b} & \dfrac{\beta_y\,\delta\,(K\,(\beta-b\,\beta)+\delta)\,\lambda}{(-1+b)\,\beta\,((-1+b)\,K\,\beta\,\gamma+\delta\,\lambda)} \\ \dfrac{\gamma\,(K\,(\beta-b\,\beta)+\delta)\,\lambda}{(-1+b)\,K\,\beta\,\gamma+\delta\,\lambda} & b & \dfrac{b\,\delta}{1-b} & \dfrac{\beta_v\,\gamma\,(K\,(\beta-b\,\beta)+\delta)\,\lambda}{\beta\,((-1+b)\,K\,\beta\,\gamma+\delta\,\lambda)} \\ 0 & 0 & 0 & -0^\in\,c\,(1+\in) + \dfrac{\beta_z\,((-1+b)\,K\,\beta-\delta)\,\delta\,\lambda}{(-1+b)\,\beta\,((-1+b)\,K\,\beta\,\gamma+\delta\,\lambda)} \end{pmatrix}$$

Tr[J[E$*$]] corresponding to 4 dim model when K=γ=1 is

Out[136]= $-1 + \dfrac{b\,\delta}{1-b} - 0^\in\,c\,(1+\in) - \dfrac{\delta\,\lambda}{(-1+b)\,\beta} + \dfrac{\beta_z\,((-1+b)\,\beta-\delta)\,\delta\,\lambda}{(-1+b)\,\beta\,((-1+b)\,\beta+\delta\,\lambda)}$

Tr[J[E$*$]] >0 iff

Out[138]= $\beta>0$ && $\lambda>0$ && $0<\delta<(-1+b)\,\beta$ && $\beta_y>0$ && $\beta_v>0$ &&

$((-1+b)\,\beta-\delta)\,\delta\,\lambda\,\left((-1+b)\,\beta^2\,(-1+b+b\,\delta)+\beta\,\delta\,(-2+\beta_z+b\,(2-\beta_z+\delta))\,\lambda+\delta^2\,\lambda\,(\beta_z+\lambda)\right) <$

0 && $0<c< -\dfrac{(-1+b)\,\beta^2\,(-1+b+b\,\delta)+\beta\,\delta\,(-2+\beta_z+b\,(2-\beta_z+\delta))\,\lambda+\delta^2\,\lambda\,(\beta_z+\lambda)}{(-1+b)\,\beta\,((-1+b)\,\beta+\delta\,\lambda)}$ && $\gamma>0$

$$b1_* = \frac{1}{27\, c^2\, \beta^2\, \beta_v^2\, \beta_y^2\, \beta_z^2\, \lambda}$$

$$\left(-2\, \sqrt{ \beta^2\, \beta_y^2\, \beta_z^2\, \left(c^2\, \beta^2\, \beta_y^2 + c\, \beta_v\, \left(-c\, \beta\, \beta_y - 3\, \beta_v\, \beta_z + \beta_y\, \beta_z\, (\beta + 3\, \delta) \right)\, \lambda + \beta_v^2\, (c - \beta_z)^2\, \lambda^2 \right)^3 } \right. +$$

$$\beta\, \beta_y\, \beta_z\, \left(2\, c^2\, \beta^2\, \beta_y^2 + c\, \beta_v\, \left(-5\, c\, \beta\, \beta_y + \beta_z\, \left(-9\, \beta_v + 5\, \beta\, \beta_y + 9\, \beta_y\, \delta \right) \right) \right)\, \lambda + 2\, \beta_v^2\, (c - \beta_z)^2\, \lambda^2 \right)$$

$$\left. \left(\beta_v\, \beta_z\, \lambda - c\, \left(\beta\, \beta_y + \beta_v\, \lambda \right) \right) \right)$$

$$b2_* = \frac{1}{27\, c^2\, \beta^2\, \beta_v^2\, \beta_y^2\, \beta_z^2\, \lambda}$$

$$\left(2\, \sqrt{ \beta^2\, \beta_y^2\, \beta_z^2\, \left(c^2\, \beta^2\, \beta_y^2 + c\, \beta_v\, \left(-c\, \beta\, \beta_y - 3\, \beta_v\, \beta_z + \beta_y\, \beta_z\, (\beta + 3\, \delta) \right)\, \lambda + \beta_v^2\, (c - \beta_z)^2\, \lambda^2 \right)^3 } \right. +$$

$$\beta\, \beta_y\, \beta_z\, \left(2\, c^2\, \beta^2\, \beta_y^2 + c\, \beta_v\, \left(-5\, c\, \beta\, \beta_y + \beta_z\, \left(-9\, \beta_v + 5\, \beta\, \beta_y + 9\, \beta_y\, \delta \right) \right) \right)\, \lambda + 2\, \beta_v^2\, (c - \beta_z)^2\, \lambda^2 \right)$$

$$\left. \left(\beta_v\, \beta_z\, \lambda - c\, \left(\beta\, \beta_y + \beta_v\, \lambda \right) \right) \right)$$

2-2)Interior equilibrium

Analysis of the stability of the interior point E_* when $K=\gamma=1$:

In[143]:=
```
(*Since z=0, we reduce our analysis for this point to 3 dimensions*)
jac3=Grad[dyn3/.cKga1,{x,y,v}]//FullSimplify;
Print["J(E_*) is when K=1"](*in three dimension*)
JsS=(jac3/.sol〚3〛/.cKga1)//FullSimplify;
JsS//MatrixForm
Trs=Tr[JsS](*the trace of J[E*] in 3 dimension*);
Print["Tr(J(E*))=",Trs//FullSimplify]
Print["Det(J(E*))=",Det[JsS]//FullSimplify]


pc=Collect[Det[ψ IdentityMatrix[3]-JsS],ψ];
coT=CoefficientList[pc,ψ]//FullSimplify;
Length[coT]
Print["a₁=",a1=coT〚3〛, ", a₂=",a2=coT〚2〛, ", a₃=",a3=coT〚1〛]
H2=a1*a2-a3;
Print["H2(b0)=",H2/.b→b0S//FullSimplify]
Print["Denominator of H2 is ",Denominator[Together[H2]]/.cKga1//FullSimplify]
ϕb=Collect[Numerator[Together[H2]]/(δ λ),b]/.cKga1;
cofi=CoefficientList[ϕb,b](*Coefficients of ϕ(b)*);
Print["value of ϕ(b) at crit b is "]
ϕb/.b→b0S/.cKga1//FullSimplify
```

J(E_*) is when K=1

Out[146]//MatrixForm=

$$\begin{pmatrix} \dfrac{\delta\, \lambda}{\beta - b\, \beta} & \dfrac{\delta\, \lambda}{\beta - b\, \beta} & -\dfrac{\delta}{-1+b} \\[2ex] \dfrac{\left((-1+b)\, \beta - \delta \right)\, \lambda}{(-1+b)\, \beta + \delta\, \lambda} & -1 & \dfrac{\delta}{-1+b} \\[2ex] \dfrac{(\beta - b\, \beta + \delta)\, \lambda}{(-1+b)\, \beta + \delta\, \lambda} & b & \dfrac{b\, \delta}{1-b} \end{pmatrix}$$

$$\text{Tr}(J(E*)) = -1 + \frac{\delta \ (b \ \beta + \lambda)}{\beta - b \ \beta}$$

$$\text{Det}(J(E*)) = \frac{\delta \ (\beta - b \ \beta + \delta) \ \lambda}{(-1 + b) \ \beta}$$

Out[152]= 4

$$a_1 = \frac{\beta \ (-1 + b + b \ \delta) + \delta \ \lambda}{(-1 + b) \ \beta}, \quad a_2 =$$

$$\frac{\delta \ \lambda \ \left((-1 + b) \ \beta \ (-1 + \beta + \delta + b \ (1 - \beta + \delta)) + \left((-1 + b)^2 \ \beta + b \ \delta^2 \right) \ \lambda \right)}{(-1 + b)^2 \ \beta \ ((-1 + b) \ \beta + \delta \ \lambda)}, \quad a_3 = \delta \left(1 + \frac{\delta}{\beta - b \ \beta} \right) \lambda$$

H2(b0) = $(1 + \beta + \delta) \ \lambda \ (1 + \beta + \delta + \lambda)$

Denominator of H2 is $(-1 + b)^3 \ \beta^2 \ ((-1 + b) \ \beta + \delta \ \lambda)$

value of $\phi(b)$ at crit b is

Out[160]=
$$\frac{\delta^3 \ (1 + \beta + \delta) \times (1 + \lambda) \times (1 + \beta + \delta + \lambda)}{\beta}$$

In[161]:=
```
(*Analysis of the stability of E* in 4 dim when ϵ=0*)

pc=Collect[Det[ψ IdentityMatrix[4]-(jacEs1/.cep0)],ψ];
coT=CoefficientList[pc,ψ]//FullSimplify;
Length[coT]
a1=coT〚4〛//FullSimplify ;a2=coT〚3〛//FullSimplify;a3=coT〚2〛//FullSimplify ; a4=coT〚1〛//FullSi
Print["a₁=",a1, ", a₂=",a2, ", a₃=",a3, " ,a₄=", a4 ]
H4=a1*a2*a3-a3^2+a1^2 a4;
Print["H2(b0)=",H4/.b→b0S//FullSimplify]
Print["Denominator of H2 is ",Denominator[Together[H4]]//FullSimplify]
ϕb4=Collect[Numerator[Together[H4]],b];
cofi=CoefficientList[ϕb4,b];
Length[cofi]
Print["value of ϕ(b) at crit b is "]
ϕb4/.b→b0S//FullSimplify
```

Out[163]= 5

$$a_1 = c + \frac{(-1+b)\,\beta^2\,(-1+b+b\,\delta) + \beta\,\delta\,(-2+\beta_z + b\,(2-\beta_z+\delta))\,\lambda + \delta^2\,\lambda\,(\beta_z+\lambda)}{(-1+b)\,\beta\,((-1+b)\,\beta+\delta\,\lambda)}$$

$$, \quad a_2 = \frac{1}{(-1+b)^2\,\beta^2}\left((-1+b)\,c\,\beta\,(\beta\,(-1+b+b\,\delta)+\delta\,\lambda) + \frac{1}{(-1+b)\,\beta+\delta\,\lambda}\right.$$
$$\delta\,\lambda\,\left(-(-1+b)^2\,\beta^3+\beta_z\,\delta^2\,\lambda - (-1+b)\,\beta^2\,(1-\beta_z-\delta+b\,(-1+\beta_z)\times(1+\delta)+\lambda - b\,\lambda) + \right.$$
$$\left.\left.\beta\,\delta\,(b\,\delta\,\lambda+\beta_z\,(-1+b+b\,\delta+\lambda-b\,\lambda)))\right)\right), \quad a_3 = \frac{1}{(-1+b)^3\,\beta^2\,((-1+b)\,\beta+\delta\,\lambda)^2}$$
$$\delta\,\lambda\,\left(((-1+b)\,\beta-\delta)\,\left((-1+b)^4\,\beta^3 + (-1+b)\,\beta\,\delta\,(\beta_z+(-1+b)\,\beta\,(-2+2\,b+\beta_z) - \beta_z\,(b+\delta+b\,\delta))\,\lambda - \right.\right.$$
$$\left.\delta\,\left((-1+b)^2\,\beta\,(\beta_z-\delta)+b\,\beta_z\,\delta^2\right)\lambda^2\right) -$$
$$\left.(-1+b)\,c\,\beta\,((-1+b)\,\beta+\delta\,\lambda)\,\left((-1+b)^2\,\beta^2 - b\,\delta^2\,\lambda - (-1+b)\,\beta\,(-1+b+\delta+b\,\delta+(-1+b)\,\lambda)\right)\right)$$

$$, a_4 = \frac{((-1+b)\,\beta-\delta)\,\delta\,\lambda\,(\beta_z\,\delta\,(\beta-b\,\beta+\delta)\,\lambda + (-1+b)\,c\,\beta\,((-1+b)\,\beta+\delta\,\lambda))}{(-1+b)^2\,\beta^2\,((-1+b)\,\beta+\delta\,\lambda)}$$

H2(b0) = $c\,(1+\beta+\delta)\times(1+c+\beta+\delta)\,\lambda\,(c+\lambda)\,(1+\beta+\delta+\lambda)$

Denominator of H2 is $(-1+b)^6\,\beta^5\,((-1+b)\,\beta+\delta\,\lambda)^4$

Out[171]= **11**

value of $\phi(b)$ at crit b is

Out[173]= $\dfrac{c\,\delta^{10}\,(1+\beta+\delta)\times(1+c+\beta+\delta)\,\lambda\,(1+\lambda)^4\,(c+\lambda)\,(1+\beta+\delta+\lambda)}{\beta}$

```
FindInstance[Join[{Tr[jacEs/.cKga1/.ϵ→0]>0},Drop[cp,{9,10}]],Drop[par,{10,11}]]
```

Out[∘]= $\left\{\left\{b\to 2,\,\beta\to 2,\,\lambda\to 1,\,\delta\to 1,\,\beta_y\to 1,\,\beta_v\to 1,\,\beta_z\to 28,\,c\to 1,\,\gamma\to 1\right\}\right\}$

Numerical values ϵ=0:

```
In[174]:=
cn=Join[cnT17,cep0];
cb=NSolve[(ϕb//.Drop[cnT17,{5}])==0,b,WorkingPrecision→10]
bM=Max[Table[Re[b/.cb〚i〛],{i,Length[cb]}]];
Print["bH=",bH=N[bM,30]]
Chop[{vn,vi,vp}/.cnT17];

Print["b0=",b0/.cn//N, " , b1=", bnn〚1〛,    ", b2=", bnn〚2〛, "  ,bH=",bH]
PRN=Chop[PR//.cn//N](*values of the roots v*);
Print["E*",Chop[Es/.cn]//N]
Print["E+=",Chop[Ep/.cn]//N]
Print["Ei=",Chop[Ei/.cn]//N]
Print["Eigv of E*:",Append[Eigenvalues[JsS],Es〚2〛-ye]/.cn," , Eigv of E+:",
Chop[Eigenvalues[JEp//.cn]]//N//FullSimplify, " Eigv of Eim:",Chop[Eigenvalues[JEi//.cn]]//N,
"   , Eigv of EK:",Eigenvalues[jacK//.cn]//N]

Print["b1*=",  bc1//.cn]
Print[" b2*=",  bc2//.cn]
```

Out[175]= $\{\{b\to 0.299052\},\,\{b\to 0.835329 - 0.231156\,i\},\,\{b\to 0.835329 + 0.231156\,i\},\,\{b\to 19.0121\}\}$

bH=19.0121

b0=2.81818 , b1=3.58676, b2=8.66779 ,bH=19.0121

E∗{0.227273, 0.0584416, 2.33766}

E+={0.249944, 0.06, 2.25837, 0.072607}

Ei={0.483284, 0.06, 1.49471, 0.675711}

Eigv of E∗:{-1.25056, -0.0281268 - 0.20904 i, -0.0281268 + 0.20904 i, -0.00155844}
    , Eigv of E+:{-1.29833, -0.0332218 + 0.21255 i, -0.0332218 - 0.21255 i, 0.000834552}
     Eigv of Eim:{-1.69849, -0.076539 + 0.209952 i, -0.076539 - 0.209952 i, -0.00803072}
        , Eigv of EK:{-1.7081, 0.398103, -0.36, -0.036}

b1∗=-0.00697038

 b2∗=10.2462

Determination of the endemic points with respect to x:

In[187]:=
```
Print["ve(x)="]
vee=Solve[(x1/x)==0,v]〚1〛/.cKga1
Print["ze(x)="]
zee=Solve[v1+(x1/x)==0,z]〚1〛/.cKga1//FullSimplify
(*(y1/.zee/.vee/.y→ye);*)
xee=Solve[Collect[Numerator[Together[(y1/.zee/.vee/.y→ye)]],x]==0,x](*The expressions are so
Print["Number of endemic x"]
Length[xee]
Chop[Evaluate[xee//.cnT17]](*Numerical check*)
```

ve(x)=

Out[188]= $\left\{ v \to \dfrac{(1-x-y)\ \lambda}{\beta} \right\}$

ze(x)=

Out[190]= $\left\{ z \to \dfrac{b\ y - v\ (\beta + x\ \beta + \delta)\ + \lambda - (x+y)\ \lambda}{v\ \beta_v} \right\}$

Number of endemic x

Out[193]= 3

Out[194]= $\{\{x \to 0.249944\},\ \{x \to 1.20177\},\ \{x \to 0.483284\}\}$

3)Sections 3.1---3.3(in paper):Figures used in the manuscript (*Run the previous cell*)

Numerical illustrations when ϵ=0 (Bifurcations diagrams, parametric plots, and 3D plot)

Bifurcation diagram when b varies:

In[195]:=
```
ClearParameters;
βv=0.16; βy=0.48;K=1;b=9;γ=1;λ=0.36;β=0.11;δ=0.2;βz=0.6; c=0.036;ϵ=0;
Print["{b2∗,b1∗}=",{b/.solb〚1〛,b/.solb〚2〛} ,", and b₀=",b0, "  , b_H=",bH,
 " ,b1=", bnn〚1〛,   ", b2=", bnn〚2〛]
Clear["b"];
```

```
vs= ((-1+b) β-δ) λ  (*v of E* *  ****);
     ─────────────────
     β ((-1+b) β+δ λ)
bL=28; max=3.5;bmin=-0.9;
lin1=Line[{{ bc1,0},{ bc1,max}}];
li1=Graphics[{Thick,Black,Dashed,lin1}];
lin2=Line[{{ bc2,0},{ bc2,max}}];
li2=Graphics[{Thick,Black,Dashed,lin2}];
lin3=Line[{{ b0,0},{ b0,max}}];
li3=Graphics[{Thick,Black,Dashed,lin3}];
linH=Line[{{ bH,0},{ bH,max}}];
liH=Graphics[{Thick,Black,Dashed,linH}];
linb1=Line[{{ bnn[[1]],0},{ bnn[[1]],max}}];
lib1=Graphics[{Thick,Black,Dashed,linb1}];
linb2=Line[{{ bnn[[2]],0},{ bnn[[2]],max}}];
lib2=Graphics[{Thick,Black,Dashed,linb2}];
linb9=Line[{{ 9,0},{9,max}}];
lib9=Graphics[{Thick,Black,Dashed,linb2}];
pn=Plot[{vn},{b,0,bL},PlotStyle→{Green,Dotted},PlotRange→{{bmin,bL},{-2,max}},PlotLegends→{"
p0=Plot[0,{b,0,bL},PlotStyle→{Brown,Thick},PlotRange→{{bmin,bL},{-2,max}},PlotLegends→{"E₀ s
ppa=Plot[{vp},{b,0, bnn[[2]]},PlotStyle→{Purple,Dotted},PlotRange→{{bmin,bL},{-2,max}},
PlotLegends→{"E₊ outside domain"}];
ppb=Plot[{vp},{b,bnn[[2]], bL},PlotStyle→{Purple,Thick,Dashed},
PlotRange→{{bmin,bL},{-2,max}},PlotLegends→{"E₊ unstable"}];
pi1=Plot[{vi},{b,0,bnn[[1]]},PlotStyle→{Blue,Dotted},
PlotRange→{{bmin,bL},{-2,max}},PlotLegends→{"Eᵢₘ outside domain"}];
pi2=Plot[{vi},{b,bnn[[1]],bL},PlotStyle→{Red,Thick},
PlotRange→{{bmin,bL},{-2,max}},PlotLegends→{"Eᵢₘ stable"}];
pi3=Plot[{vi},{b,bnn[[2]],bL},PlotStyle→{Blue,Thick,Dashed},
PlotRange→{{bmin,bL},{-2,max}}(*,PlotLegends→{"Eᵢ unstable"}*)];
ps1=Plot[{vs},{b,b0,bnn[[1]]},PlotStyle→{Brown,Thick},
PlotRange→{{bmin,bL},{-2,max}},PlotLegends→{"E∗ stable"}];
ps2=Plot[{vs},{b,bnn[[1]],bnn[[2]]},PlotStyle→{Cyan,Thick,Dashed},
PlotRange→{{bmin,bL},{-2,max}},PlotLegends→{"E∗ unstable"}];
ps3=Plot[{vs},{b,bnn[[2]],bH},PlotStyle→{Brown,Thick},
PlotRange→{{bmin,bL},{0,max}}(*,PlotLegends→{"E∗ stable"}*)];
ps4=Plot[{vs},{b,bH,bL},PlotStyle→{Cyan,Thick,Dashed},
PlotRange→{{bmin,bL},{-2,max}}(*,PlotLegends→{"E∗ unstable"}*)];
pdf1=Plot[{0},{b,0,b0},PlotStyle→{Magenta, Thick},
PlotRange→{{bmin,bL},{0,max}},PlotLegends→{"Eₖ stable"}];
pdf2=Plot[{0},{b,b0,bL},PlotStyle→{Orange, Thick,Dashed},
PlotRange→{{bmin,bL},{0,max}},PlotLegends→{"Eₖ unstable"}];
pvmax=Plot[{c (b-1)/(βz δ)(*,(λ(1-c/βz))/β*)},{b,0,bL},PlotStyle→{Yellow},
PlotRange→{{0,bL},{0,max}},
PlotLegends→{"c (b-1)/(s δ)"}];


bifT=Show[{ppa,ppb,pi1,pi2,ps1,ps2,ps3,ps4,pdf1,pdf2,pn,li1,li2,li3,
lib2,lib1,lib9,pvmax,liH},AxesLabel→{"b","v"},PlotRange→{{bmin,bL},{-3/2,max}},
Epilog→{Text["b₁∗",Offset[{-2,10},{ bc1,0}]],{PointSize[Large],
Style[Point[{ bc1,0}],Purple]},
Text["b₂∗",Offset[{11,10},{ bc2,0}]],{PointSize[Large],Style[Point[{ bc2,0}],Yellow]},
Text["b₀",Offset[{-7,11},{ b0,0}]],{PointSize[Large],Style[Point[{ b0,0}],Black]},
Text["bH",Offset[{-10,11},{ bH,0}]],{PointSize[Large],Style[Point[{ bH,0}],Green]},
Text["b₁",Offset[{8,11},{ bnn[[1]],0}]],{PointSize[Large],Style[Point[{ bnn[[1]],0}],Pink]},
Text["b₂",Offset[{-7,11},{ bnn[[2]],0}]],{PointSize[Large],Style[Point[{ bnn[[2]],0}],Red]},
```

```
Text["b=9",Offset[{7,11},{9,0}]],{PointSize[Large],Style[Point[{ 9,0}],Brown]}}]
Export["BiifT17.pdf",bifT]
```

$\{b_{2*},b_{1*}\}=\{10.2462, -0.00697038\}$, and b_0=2.81818 , b_H=19.0121 ,b1=3.58676, b2=8.66779

Out[230]= BiifT17.pdf

Numerical solution of the stability (Bifurcation diagram) wrt v:

```
In[ ]:= (*Checks on the stability of the fixed points**)
Print["Eigenvalues of E*:"]
Print["E* between b0 and b1"]
Eigenvalues[jacEs1] /. b → 3 // N
Print["E* between b1 and b2"]
Eigenvalues[jacEs1] /. b → 5 // N
Print["E* between b2 and bH"]
Eigenvalues[jacEs1] /. b → 15 // N
Print["E* after bH"]
Eigenvalues[jacEs1] /. b → 25 // N
Print["Eigenvalues of Eim:"]
Print["Eim between 0 and b1"]
Chop[Eigenvalues[JEi] /. b → 2 // N]
Print["E* between b1 and b2*"]
Chop[Eigenvalues[JEi] /. b → 9 // N]
Print["Eigenvalues of E+:"]
Print["Eim between 0 and b2"]
Chop[Eigenvalues[JEp] /. b → 2 // N]
Print["E* between b2 and b2*"]
Chop[Eigenvalues[JEp] /. b → 9 // N]
```

Eigenvalues of E∗:

E∗ between b0 and b1

Out[ ]= $\{-0.0225504, -1.29942, -0.311695, -0.0161607\}$

E∗ between b1 and b2

Out[ ]= $\{0.0100227, -1.26091, -0.0763636 - 0.159107\,\mathbb{i}, -0.0763636 + 0.159107\,\mathbb{i}\}$

E∗ between b2 and bH

Out[ ]= $\{-0.0126814, -1.24799, -0.00652554 - 0.223959\,\mathbb{i}, -0.00652554 + 0.223959\,\mathbb{i}\}$

E∗ after bH

Out[ ]= $\{-0.0212776, -1.24705, 0.00572283 - 0.230932\,\mathbb{i}, 0.00572283 + 0.230932\,\mathbb{i}\}$

Eigenvalues of Eim:

Eim between 0 and b1

Out[ ]= $\{-1.00272, -0.249728, -0.0735624, 0.0456119\}$

E∗ between b1 and b2∗

Out[ ]= $\{-1.69849, -0.00803072, -0.076539 - 0.209952\,\mathbb{i}, -0.076539 + 0.209952\,\mathbb{i}\}$

Eigenvalues of E+:

Eim between 0 and b2

Out[ ]= $\{-0.559828, -0.123746, 0.053141 - 0.0959514\,\mathbb{i}, 0.053141 + 0.0959514\,\mathbb{i}\}$

E∗ between b2 and b2∗

Out[ ]= $\{-1.29833, 0.000834552, -0.0332218 - 0.21255\,\mathbb{i}, -0.0332218 + 0.21255\,\mathbb{i}\}$

(x-b)-Bifurcation diagram :

In[ ]:=
```
Clear["K"];
K=1;
Print["{b₂∗,b₁∗}=",{b/.solb[[1]],b/.solb[[2]]} ,", and b₀=",b0, "   , bₕ=",bH,
 " ,b1=", bnn[[1]],    ", b2=", bnn[[2]]]
Clear["b"];
xs=EsS[[1]](*x of E∗ ∗  ∗∗∗∗);
bL=22; max=1.4;bmin=-0.9;
lin1=Line[{{ bc1,0},{ bc1,max}}];
li1=Graphics[{Thick,Black,Dashed,lin1}];
lin2=Line[{{ bc2,0},{ bc2,max}}];
li2=Graphics[{Thick,Black,Dashed,lin2}];
lin3=Line[{{ b0,0},{ b0,max}}];
li3=Graphics[{Thick,Black,Dashed,lin3}];
linH=Line[{{ bH,0},{ bH,max}}];
liH=Graphics[{Thick,Black,Dashed,linH}];
linb1=Line[{{ bnn[[1]],0},{ bnn[[1]],max}}];
lib1=Graphics[{Thick,Black,Dashed,linb1}];
linb2=Line[{{ bnn[[2]],0},{ bnn[[2]],max}}];
lib2=Graphics[{Thick,Black,Dashed,linb2}];
linb9=Line[{{ 9,0},{9,max}}];
lib9=Graphics[{Thick,Black,Dashed,linb2}];
(*pn=Plot[{x/.xee[[1]]},{b,0,b0},PlotStyle→{Purple,Dotted,Thick},PlotRange→{{bmin,bL},{-2,max}
pn1=Plot[{x/.xee[[1]]},{b,0,bnn[[2]]},PlotStyle→{Purple,Dotted,Thick},PlotRange→{{bmin,bL},{-2,m
PlotLegends→{"E₊ (z<0)"}];
```

```
pn2=Plot[{x/.xee[[1]]},{b,bnn[[2]],bL},PlotStyle→{Purple,Dashed,Thick},PlotRange→{{bmin,bL},{-2,
PlotLegends→{"E₊ unstable"}];
p0=Plot[0,{b,0,bL},PlotStyle→{Brown,Thick},PlotRange→{{bmin,bL},{-2,max}},PlotLegends→{"E₀ s
ppa=Plot[{x/.xee[[2]]},{b,0, bL},PlotStyle→{Green,Dotted},PlotRange→{{bmin,bL},{-2,max}},
PlotLegends→{"E₋ (v<0)"}];
pi1=Plot[{x/.xee[[3]]},{b,0,bnn[[1]]},PlotStyle→{Red,Dotted},
PlotRange→{{bmin,bL},{-2,max}},PlotLegends→{"E_im (z<0)"}];
pi2=Plot[{x/.xee[[3]]},{b,bnn[[1]],bL},PlotStyle→{Red,Thick},
PlotRange→{{bmin,bL},{-2,max}},PlotLegends→{"E_im stable"}];
ps1=Plot[{xs},{b,b0,bnn[[1]]},PlotStyle→{Brown,Thick},
PlotRange→{{bmin,bL},{-2,max}},PlotLegends→{"E_* stable"}];
ps2=Plot[{xs},{b,bnn[[1]],bnn[[2]]},PlotStyle→{Cyan,Thick,Dashed},
PlotRange→{{bmin,bL},{-2,max}},PlotLegends→{"E_* unstable"}];
ps3=Plot[{xs},{b,bnn[[2]],bH},PlotStyle→{Brown,Thick},
PlotRange→{{bmin,bL},{0,max}}(*,PlotLegends→{"E_* stable"}*)];
ps4=Plot[{xs},{b,bH,bL},PlotStyle→{Cyan,Thick,Dashed},
PlotRange→{{bmin,bL},{-2,max}}(*,PlotLegends→{"E_* unstable"}*)];
pdf1=Plot[K,{b,0,b0},PlotStyle→{Magenta, Thick},
PlotRange→{{bmin,bL},{0,max}},PlotLegends→{"E_K stable"}];
pdf2=Plot[K,{b,b0,bL},PlotStyle→{Orange, Thick,Dashed},
PlotRange→{{bmin,bL},{0,max}},PlotLegends→{"E_K unstable"}];
pxmax=Plot[{1-ye-β/λ (ye (b-1)/δ)},{b,0,bL},PlotStyle→{Yellow},
PlotRange→{{0,bL},{0,max}},
PlotLegends→{"1-y_e ( 1- β (b-1) / δ λ )"}];


bifx=Show[{pn1,pn2,pi1,pi2,ps1,ps2,ps3,ps4,pdf1,pdf2,ppa,pxmax,li1,li2,li3,
lib2,lib1,lib9,liH},AxesLabel→{"b","x"},PlotRange→{{bmin,bL},{-0.1,max}},
Epilog→{Text["b₁*",Offset[{-2,10},{ bc1,0}]],{PointSize[Large],
Style[Point[{ bc1,0}],Purple]},
Text["b₂*",Offset[{11,10},{ bc2,0}]],{PointSize[Large],Style[Point[{ bc2,0}],Yellow]},
 Text["b₀",Offset[{-7,11},{ b0,0}]],{PointSize[Large],Style[Point[{ b0,0}],Black]},
Text["b_H",Offset[{-10,11},{ bH,0}]],{PointSize[Large],Style[Point[{ bH,0}],Green]},
 Text["b₁",Offset[{8,11},{ bnn[[1]],0}]],{PointSize[Large],Style[Point[{ bnn[[1]],0}],Pink]},
Text["b₂",Offset[{-7,11},{ bnn[[2]],0}]],{PointSize[Large],Style[Point[{ bnn[[2]],0}],Red]},
Text["b=9",Offset[{7,11},{9,0}]],{PointSize[Large],Style[Point[{ 9,0}],Brown]}}]
Export["bifx.pdf",bifx]
```

$\{b_{2*},b_{1*}\}=\{10.2462, -0.00697038\}$, and $b_0=2.81818$, $b_H=19.0121$,$b1=3.58676$, $b2=8.66779$

Out[ ]= `bifx.pdf`

Bifurcation diagram of codimension 2:

In[231]:=
```
Clear["b"];
Clear["β"];
(*Here we modified a slightly the values from FindInstance when Tr[jacEs]>0*)
(*λ=1;δ=0.2;βy=0.48;βv=0.16;βz=28;c=1;γ=1;K=1; *)

bm=1; bM=25; ym=0; yM=0.38;
trEp=Tr[JEp];
trEim=Tr[JEi];
detEs=Det[jacEs1];

R1=Style[Text["R₀>1,H∗<0,Δ<0",{16,0.30}]],13];
R1a=Style[Text["Attractor cycle",{17,0.25}]],13];
R2=Style[Text["R₀>1,H∗>0,Δ<0",{13,0.13}]],10];
R2a=Style[Text["E∗ Stable",{15,0.09}]],10];
RIII=Style[Text["R₀>1,H∗>0,Δ>0",{10,0.06}]],10];
RIIIa=Style[Text["Eim stable,E∗ unstable",{15,0.04}]],10];
RII=Style[Text["E∗",{4,0.08}]],10];
RI=Style[Text["EK Stable",{6,0.01}]],10];
RIa=Style[Text["R₀<1,H∗>0,Δ>0",{4,0.03}]],10];
Rbi=Style[Text["Eim",{11,0.09}]],9];
Rbia=Style[Text["E∗",{13,0.075}]],9];

pt0s=Text["b₀",Offset[{-10,6},{b0/.β→0.11,0.11}]];
pt0={PointSize[Medium],Style[Point[{b0/.β→0.11,0.11}],Black]};
ptHs=Text["bH",Offset[{10,6},{bH,0.11}]];
ptH1={PointSize[Medium],Style[Point[{bH,0.11}],Black]};
ptb2as=Text["b2∗",Offset[{14,-3},{bc2/.β→0.11,0.11}]];
ptb2a={PointSize[Medium],Style[Point[{bc2/.β→0.11,0.11}],Black]};
ptb2bs=Text["b2",Offset[{-6,-5},{bnn〚2〛,0.11}]];
ptb2b={PointSize[Medium],Style[Point[{bnn〚2〛,0.11}],Black]};
ptb1s=Text["b1",Offset[{8,6},{bnn〚1〛,0.11}]];
```

```
ptb1={PointSize[Medium],Style[Point[{bnn〚1〛,0.11}],Black]};
ptbas=Text["b₇",Offset[{-2,6},{9.5,0.11}]];
ptba={PointSize[Medium],Style[Point[{9.5,0.11}],Yellow]};
ptbbs=Text["b₈=23",Offset[{10,6},{23,0.11}]];
ptbb={PointSize[Medium],Style[Point[{23,0.11}],Red]};
epi={R1,R1a,R2,R2a,RIII,RIIIa,RII,RI,RIa,Rbi,Rbia,pt0s,pt0,ptHs,ptH1,ptb2as,ptb2a,ptb2bs,ptb2b
ptbas,ptba,ptbbs,ptbb,ptb1s,ptb1};

pR0=ContourPlot[R0==1,{b,bm,bM},{β,ym,yM},ContourStyle→{Black,Dotted},
 FrameLabel→{"b","β"},LabelStyle→{Black,Bold},Frame→True,PlotLegends→{"R₀=1"}];
(*ptrEs=ContourPlot[Tr[jacEs]==0,{b,bm,bM},{β,ym,yM},ContourStyle→{Green,Bold},
 FrameLabel→{"b","β"},LabelStyle→{Black,Bold},Frame→True,PlotLegends→{"Tr[J(E*)]=0"}];*)
 pphi=ContourPlot[H2(*or ϕb*)==0,{b,bm,bM},{β,ym,yM},ContourStyle→{Blue,Bold},PlotPoints → 18
 FrameLabel→{"b","β"},LabelStyle→{Black,Bold},Frame→True,PlotLegends→{"H*=0"}];
 pdetEs=ContourPlot[detEs==0,{b,bm,bM},{β,ym,yM},ContourStyle→{Green,Bold},
 FrameLabel→{"b","β"},LabelStyle→{Black,Bold},Frame→True,PlotLegends→{"Det[J(E*)]=0"}];
  pdis=ContourPlot[Dis==0,{b,bm,bM},{β,ym,yM},ContourStyle→{Red,Thick},PlotPoints→180,
 FrameLabel→{"b","β"},LabelStyle→{Black,Bold},Frame→True,PlotLegends→{"Δ=0"}];
(*ptrEi=ContourPlot[trEim==0,{b,bm,bM},{β,ym,yM},ContourStyle→{Red,Dashed},PlotPoints → 180,
 FrameLabel→{"b","β"},LabelStyle→{Black,Bold},Frame→True,PlotLegends→{"Tr[J(Eim)]=0"}];*)
 pbeta=ContourPlot[β==0.11,{b,bm,bM},{β,ym,yM},ContourStyle→{Cyan,Dashed},
 FrameLabel→{"b","β"},LabelStyle→{Black,Bold},Frame→True,PlotLegends→{"β=0.11"}];
 ppar=ContourPlot[ EsS〚2〛/ye==1,{b,bm,bM},{β,ym,yM},ContourStyle→{Green,Thick},
 FrameLabel→{"b","β"},LabelStyle→{Black,Bold},Frame→True,PlotLegends→{"y*/ye=1"}];



 peq={pR0,pdis,pphi,ppar(*pdetEs,ptrEi*)(*ptrEs,*),pbeta};
pcut=Show[peq,Epilog→epi]
Export["map4.pdf",pcut]
```

Out[270]=

Out[271]= map4.pdf

```
In[ ]:= (***Some checks***)
    (*For Ein**)
    Chop[Evaluate[xee[[3]] /. b → 2]] // N
    Chop[Evaluate[zee /. vee /. xee[[3]] /. y → ye /. b → 2]] // N
    Chop[vee /. zee /. xee[[3]] /. y → ye /. b → 2] // N
    (*For E+**)
    Chop[Evaluate[xee[[1]] /. b → 3]] // N
    Chop[zee /. vee /. xee[[1]] /. y → ye /. b → 3] // N
    Chop[vee /. zee /. xee[[1]] /. y → ye /. b → 3] // N
```

$Out[ ]=$ $\{x \to 0.768927\}$

$Out[ ]=$ $\{z \to -0.439052\}$

$Out[ ]=$ $\{v \to 0.559875\}$

$Out[ ]=$ $\{x \to 0.113348\}$

$Out[ ]=$ $\{z \to -0.912093\}$

$Out[ ]=$ $\{v \to 2.70541\}$

```
In[ ]:= Chop[Evaluate[xee[[2]] /. b → 6]] // N
    zee /. vee /. xee[[2]] /. y → ye /. b → 6 // N
    vee /. zee /. xee[[2]] /. y → ye /. b → 6 // N
```

$Out[ ]=$ $\{x \to 1.15674\}$

$Out[ ]=$ $\left\{z \to -5.21725 + 3.87119 \times 10^{-16} \, \mathbb{i}\right\}$

$Out[ ]=$ $\left\{v \to -0.709334 - 9.08364 \times 10^{-17} \, \mathbb{i}\right\}$

Parametric plots at the intervals of stability:

When b2<b=9<b2*:

In[ ]:=
```
b=9;
Print["E*",Es=EsS//N]
Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp//N]]
Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi//N]]

X={x,y,v,z};
Xt=Map[#[t]&,X ];
Thread[X→Xt];
dynt=dyn/.Thread[X→Xt];
x1=dynt〚1〛 ;
y1=dynt〚2〛;
v1=dynt〚3〛;
z1=dynt〚4〛;
x0=0.9; y0=0.01;v0=0.01;z0=0.01;
ode3={x'[t]⩵x1,y'[t]⩵y1,v'[t]⩵v1,z'[t]⩵z1,x[0]⩵x0,y[0]⩵y0,v[0]⩵v0,z[0]⩵z0};
sol22=NDSolve[ode3,{x,y,v,z},{t,0,400}];
pdy2=Plot[{x[t]/100/. sol22,y[t]/. sol22,v[t]/100/. sol22,z[t]/. sol22},{t,0,400},
PlotLegends→{"x/100","y","v/100","z"}];
pEs2=Plot[{x/100/.x→Es〚1〛,y/.y→Es〚2〛,v/100/.v→Es〚3〛,z/.z→0},{t,0,1000},
PlotStyle→{Dashed}];
pEi2=Plot[{x/100/.x→Ei〚1〛,y/.y→Ei〚2〛,v/100/.v→Ei〚3〛,z/.z→Ei〚4〛},{t,0,1000},
PlotStyle→{Dashed}];
Dyni22=Show[pdy2,pEi2];
Dyns22=Show[pdy2,pEs2]

(*****…Parametric plot conditions***)
ppb9=ParametricPlot[{  x[t],(y[t])}/.sol22,{t,0,200}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Blue}];
py9=Plot[(y/.y→Es〚2〛),{t,0,800},PlotStyle→{Dashed,Green}];
pb9=Show[{ppb9,py9, Graphics[{Green,Dashed,Line[{{x/.x→Es〚1〛,0},{x/.x→Es〚1〛,1}}]}] },
Epilog→{{Text["(x*,y*)",Offset[{10,10},{(x/.x→Es〚1〛),(y/.y→Es〚2〛)}]]},{PointSize[Large],
Style[Point[{(x/.x→Es〚1〛),(y/.y→Es〚2〛)}],Black]}},{PointSize[Large],Point[{x0,y0}]},
Text["(x0,y0)",Offset[{-10,8},{x0,y0}]]}]
Export["pb9.pdf",pb9]
Export["Dyni22.pdf",Dyni22]
Export["Dyns22.pdf",Dyns22]
```

E*{0.227273, 0.0584416, 2.33766}

E+={0.249944, 0.06, 2.25837, 0.072607}

Ei={0.483284, 0.06, 1.49471, 0.675711}

Out[ ]=

Out[•]= pb9.pdf

Out[•]= Dyni22.pdf

Out[•]= Dyns22.pdf

When b2<b=9<b2* and different initial values :

```
In[•]:=  b=9;
         Print["E*",Es=EsS//N]
         Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp//N]]
         Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi//N]]

         X={x,y,v,z};
         Xt=Map[#[t]&,X ];
         Thread[X→Xt];
         dynt=dyn/.Thread[X→Xt];
         x1=dynt〚1〛 ;
         y1=dynt〚2〛;
         v1=dynt〚3〛;
         z1=dynt〚4〛;
         x0=0.5; y0=0.01;v0=1.2;z0=0.5;
         ode3={x'[t]⩵x1,y'[t]⩵y1,v'[t]⩵v1,z'[t]⩵z1,x[0]⩵x0,y[0]⩵y0,v[0]⩵v0,z[0]⩵z0};
         sol22=NDSolve[ode3,{x,y,v,z},{t,0,1000}];
         pdy2=Plot[{x[t]/100/. sol22,y[t]/. sol22,v[t]/100/. sol22,z[t]/100/. sol22},{t,0,600},
         PlotLegends→{"x/100","y","v/100","z/100"}];
         pEs2=Plot[{x/100/.x→Es〚1〛,y/.y→Es〚2〛,v/100/.v→Es〚3〛,z/.z→0},{t,0,1000},
         PlotStyle→{Dashed}];
         pEi2=Plot[{x/100/.x→Ei〚1〛,y/.y→Ei〚2〛,v/100/.v→Ei〚3〛,z/.z→Ei〚4〛},{t,0,1000},
         PlotStyle→{Dashed}];
         Dyni22b=Show[pdy2,pEi2]
         Dyns22=Show[pdy2,pEs2]

         (*****…Parametric plot conditions***)
         ppb9=ParametricPlot[{  x[t],(y[t])}/.sol22,{t,0,500}, AxesLabel→{"x","y"},
         PlotRange→Full,PlotStyle→{Blue}];
         py9=Plot[(y/.y→Ei〚2〛),{t,0,800},PlotStyle→{Dashed,Green}];
         pb9i=Show[{ppb9,py9, Graphics[{Green,Dashed,Line[{{x/.x→Ei〚1〛,0},{x/.x→Ei〚1〛,1}}]}] },
         Epilog→{{Text["(x*,y*)",Offset[{10,10},{(x/.x→Ei〚1〛),(y/.y→Ei〚2〛)}]],{PointSize[Large],
         Style[Point[{(x/.x→Ei〚1〛),(y/.y→Ei〚2〛)}],Black]}},{PointSize[Large],Point[{x0,y0}]},
         Text["(x0,y0)",Offset[{-10,8},{x0,y0}]]}]
         Export["pb9i.pdf",pb9i]
         Export["Dyni22b.pdf",Dyni22b]
```

E*{0.227273, 0.0584416, 2.33766}

E+={0.249944, 0.06, 2.25837, 0.072607}

Ei={0.483284, 0.06, 1.49471, 0.675711}

Out[●]= pb9i.pdf

Out[●]= Dyni22b.pdf

When b2<b=10<b2*:

In[ ]:=
```
Clear["b"];
b=9.5;
Print["E*",Es=EsS//N]
Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp//N]]
Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi//N]]

X={x,y,v,z};
Xt=Map[#[t]&,X ];
Thread[X→Xt];
dynt=dyn/.Thread[X→Xt];
x1=dynt〚1〛 ;
y1=dynt〚2〛;
v1=dynt〚3〛;
z1=dynt〚4〛;
x0=0.9; y0=0.01;v0=0.01;z0=0.01;
ode3={x'[t]⩵x1,y'[t]⩵y1,v'[t]⩵v1,z'[t]⩵z1,x[0]⩵x0,y[0]⩵y0,v[0]⩵v0,z[0]⩵z0};
sol22=NDSolve[ode3,{x,y,v,z},{t,0,400}];
pdy2=Plot[{x[t]/100/. sol22,y[t]/. sol22,v[t]/100/. sol22,z[t]/. sol22},{t,0,400},
PlotLegends→{"x/100","y","v/100","z"}];
pEs2=Plot[{x/100/.x→Es〚1〛,y/.y→Es〚2〛,v/100/.v→Es〚3〛,z/.z→0},{t,0,1000},
PlotStyle→{Dashed}];
Dyns22=Show[pdy2,pEs2]

(*****…Parametric plot conditions***)
ppb10=ParametricPlot[{  x[t],(y[t])}/.sol22,{t,0,200}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Blue}];
py10=Plot[(y/.y→Es〚2〛),{t,0,800},PlotStyle→{Dashed,Green}];
pb10=Show[{ppb10},Epilog→{{Text["(x*,y*)",Offset[{10,10},{(x/.x→Es〚1〛),
(y/.y→Es〚2〛)}]],{PointSize[Large],Style[Point[{(x/.x→Es〚1〛),(y/.y→Es〚2〛)}],Black]}},
{PointSize[Large],Point[{x0,y0}]},Text["(x₀,y₀)",Offset[{-10,8},{x0,y0}]]}]]

Export["pb10.pdf",pb10]
Export["Dyns22T.pdf",Dyns22]
```

E*{0.213904, 0.0562055, 2.38873}

E+={0.273477, 0.06, 2.18135, 0.195148}

Ei={0.453156, 0.06, 1.59331, 0.67437}

Out[ ]=

Out[ ]= pb10.pdf

Out[ ]= Dyns22T.pdf

In[ ]:=

```
Print["E*",Es=EsS//N]
Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp//N]]
Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi//N]]

X={x,y,v,z};
Xt=Map[#[t]&,X ];
Thread[X→Xt];
dynt=dyn/.Thread[X→Xt];
x1=dynt[[1]] ;
y1=dynt[[2]];
v1=dynt[[3]];
z1=dynt[[4]];
x0=0.5; y0=0.01;v0=1.2;z0=0.5;
ode3={x'[t]==x1,y'[t]==y1,v'[t]==v1,z'[t]==z1,x[0]==x0,y[0]==y0,v[0]==v0,
z[0]==z0};
sol22=NDSolve[ode3,{x,y,v,z},{t,0,1000}];
pdy2=Plot[{x[t]/100/. sol22,y[t]/. sol22,v[t]/100/. sol22,z[t]/100/. sol22},{t,0,400},
PlotLegends→{"x/100","y","v/100","z/100"}];
pEi2=Plot[{x/100/.x→Ei[[1]],y/.y→Ei[[2]],v/100/.v→Ei[[3]],z/.z→Ei[[4]]},{t,0,1000},
PlotStyle→{Dashed}];
Dyni22=Show[pdy2,pEi2]

ppb10i=ParametricPlot[{  x[t],(y[t])}/.sol22,{t,0,1900}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Red}];
py10i=Plot[(y/.y→Ei[[2]]),{t,0,800},PlotStyle→{Dashed,Green}];
pb10i=Show[{ppb10i },Epilog→{{Text["(xᵢ,yᵢ)",Offset[{10,10},{(x/.x→Ei[[1]]),
(y/.y→Ei[[2]])}]],
{PointSize[Large],Style[Point[{(x/.x→Ei[[1]]),(y/.y→Ei[[2]])}],Black]}},
{PointSize[Large],Point[{x0,y0}]},Text["(x₀,y₀)",Offset[{-10,8},{x0,y0}]]}]
pp10si=Show[{pb10,pb10i},Epilog→{{Text["(xᵢ,yᵢ)",Offset[{10,10},{(x/.x→Ei[[1]]),
(y/.y→Ei[[2]])}]]},{PointSize[Large],Style[Point[{(x/.x→Ei[[1]]),(y/.y→Ei[[2]])}],Black]}},
{PointSize[Large],Point[{x0,y0}]},Text["(x₀ᵢ,y₀ᵢ)",Offset[{-10,8},{x0,y0}]],
{Text["(x*,y*)",Offset[{10,10},{(x/.x→Es[[1]]),(y/.y→Es[[2]])}]]},{PointSize[Large],
Style[Point[{(x/.x→Es[[1]]),(y/.y→Es[[2]])}],Black]}},{PointSize[Large],Point[{0.9,0.01}]},
Text["(x₀*,y₀*)",Offset[{-10,8},{0.9,0.01}]]}]
Export["pp10si.pdf",pp10si]
Export["pb10i.pdf",pb10i]
Export["Dyni22T.pdf",Dyni22]
```

E*{0.213904, 0.0562055, 2.38873}

E+={0.273477, 0.06, 2.18135, 0.195148}

Ei={0.453156, 0.06, 1.59331, 0.67437}

Out[●]= `pp10si.pdf`

Out[●]= `pb10i.pdf`

Out[●]= `Dyni22T.pdf`

When bH<b=23<b∞:

```
In[ ]:=   Clear["b"];
          b=23;T=2000;
          Print["E*",Es=EsS//N]
          Print["E+=",Ep=Chop[{xe,ye,v,ze}/.v→vp//N]]
          Print["Ei=",Ei=Chop[{xe,ye,v,ze}/.v→vi//N]]
          X={x,y,v,z};
          Xt=Map[#[t]&,X ];
          Thread[X→Xt];
          dynt=dyn/.Thread[X→Xt];
          x1=dynt〚1〛 ;
          y1=dynt〚2〛;
          v1=dynt〚3〛;
          z1=dynt〚4〛;
          x01=0.25; y01=0.05;v01=0.01;z01=0.01;
          ode5={x'[t]⩵x1,y'[t]⩵y1,v'[t]⩵v1,z'[t]⩵z1,x[0]⩵x01,y[0]⩵y01,v[0]⩵v01,
          z[0]⩵z01};
          solHI=NDSolve[ode5,{x,y,v,z},{t,0,T}];
          pdyHI1=Plot[{x[t]/. solHI(*,y[t]/. solHI,v[t]/. solHI,z[t]/. solHI*)},{t,0,T/2},PlotStyle→Or
          DynHI1=Show[pdyHI1,PlotRange→All]

          (*****…Parametric plot conditions***)
          ppb23=ParametricPlot[{  x[t],(y[t])}/.solHI,{t,0,T}, AxesLabel→{"x","y"},
          PlotRange→Full,PlotStyle→{Blue,Dashed}];
          NDSolve[{y'[t]⩵y[t]×(x[t]-1),x'[t]⩵x[t]  (2-y[t]),x[0]⩵1,y[0]⩵2.7},{x,y},{t,0,10}];
          py23=Plot[(y/.y→Es〚2〛),{t,0,400},PlotStyle→{Dashed,Green}];
          pb23=Show[{ppb23},Epilog→{{Thick,Text["(x*,y*)",Offset[{10,10},{(x/.x→Es〚1〛),
          (y/.y→Es〚2〛)}]],{PointSize[Large],Style[Point[{(x/.x→Es〚1〛),(y/.y→Es〚2〛)}],Green]}},
          {PointSize[Large],Point[{x01,y01}]},Text["(x₀,y₀)",Offset[{-10,8},{x01,y01}]]}}]
          Export["pb23.pdf",pb23]
          Export["DynHI.pdf",DynHI]
```

E*{0.0826446, 0.0265046, 2.91551}

E+={0.297813 + 0.339863 $\mathbb{i}$, 0.06, 2.1017 - 1.11228 $\mathbb{i}$, 1.75115 + 1.463 $\mathbb{i}$}

Ei={0.297813 - 0.339863 $\mathbb{i}$, 0.06, 2.1017 + 1.11228 $\mathbb{i}$, 1.75115 - 1.463 $\mathbb{i}$}

Out[ ]=

Out[●]=


Out[●]= pb23.pdf

Out[●]= DynHI.pdf

In[●]:=
```
x0=0.3; y0=0.05;v0=2;z0=0.5;T=2000;
ode5={x'[t]==x1,y'[t]==y1,v'[t]==v1,z'[t]==z1,x[0]==x0,y[0]==y0,v[0]==v0,z[0]==z0};
solHI=NDSolve[ode5,{x,y,v,z},{t,0,10000}];
pdyHI=Plot[{x[t]/. solHI(*,y[t]/. solHI,v[t]/. solHI,z[t]/. solHI*)},{t,0,T/2},PlotStyle→Gre
AxesLabel→{"t","x"}];
DynHIc=Show[pdyHI,PlotRange→All];
(*New initial conditions**)
ppb23c=ParametricPlot[{  x[t],(y[t])}/.solHI,{t,0,150}, AxesLabel→{"x","y"},
PlotRange→Full,PlotStyle→{Red}];
pb23c=Show[{ppb23c },Epilog→{{Thick,Text["(x*,y*)",Offset[{10,10},
{(x/.x→Es〚1〛),(y/.y→Es〚2〛)}]],{PointSize[Large],Style[Point[{(x/.x→Es〚1〛),
(y/.y→Es〚2〛)}],Green]}},{PointSize[Large],Point[{x0,y0}]},Text["(x₀,y₀)",
Offset[{-10,8},{x0,y0}]]}]
Export["pb23b.pdf",pb23c]
Export["DynHIb.pdf",DynHIc]
pp23cs=Show[{pb23,pb23c},Epilog→{{Thick,Text["(x*,y*)",Offset[{10,10},{(x/.x→Es〚1〛),
(y/.y→Es〚2〛)}]],{PointSize[Large],Style[Point[{(x/.x→Es〚1〛),(y/.y→Es〚2〛)}],Green]}},
{PointSize[Large],Point[{x0,y0}]},Text["(x₀',y₀')",Offset[{16,6},{x0,y0}]],
{PointSize[Large],Point[{x01,y01}]},Text["(x₀*,y₀*)",Offset[{-10,8},{x01,y01}]]}]
xtH=Show[{DynHI1,DynHIc},AxesLabel→{"t","x"}]
Export["xtH.pdf",xtH]
Export["pp23s.pdf",pp23cs]
```

Out[●]=

Out[●]= pb23b.pdf

Out[●]= DynHIb.pdf

Out[●]=

Out[●]= xtH.pdf

Out[●]= pp23s.pdf

Illustration of the limit cycle by combining EcoEvo package with Parametric plot:

```
In[●]:=   <<EcoEvo`
         (*EcoEvoDocs;*)
         UnsetModel; T=10000;x0=0.9;y0=0.01;
         SetModel[{Pop[x]→{Equation:→dyn〚1〛,Color→Red},Pop[y]→
         {Equation:→(dyn〚2〛),Color→Green},
         Pop[v]→{Equation:→(dyn〚3〛),Color→Purple},Pop[z]→{Equation:→(dyn〚4〛),Color→Blue}}]

         fpT=SolveEcoEq[]//FullSimplify
         Print["Eigenvalues of E∗:",EcoEigenvalues[fpT〚3〛]]
         Print["We simulate around "]
         Esn=RuleListTweak[fpT〚3〛,{x,y},{0.1,0.01}]
         sol=EcoSim[Esn,T];
         psol=PlotDynamics[sol](*Time plot around E∗ *);

         lc=FindEcoCycle[FinalSlice[sol]](*Finding the cycle using time slice*);
         Print["Final time slice of the cycle is"]
         FinalTime[lc]

         pyn=Plot[(y/.fpT〚3,2〛),{t,0,T},PlotStyle→{Dashed,Green}];
         cyH=Show[{ppb23,pyn, Graphics[{Green,Dashed,
         Line[{{x/.fpT〚3,1〛,0},{x/.fpT〚3,1〛,1}}]}] },RuleListPlot[lc,{x,y},PlotStyle→{Red}],
         Epilog→{{Text["(x∗,y∗)",Offset[{10,10},{(x/.fpT〚3,1〛),(y/.fpT〚3,2〛)}]],
         {PointSize[Large],Style[Point[{(x/.fpT〚3,1〛),(y/.fpT〚3,2〛)}],Black]}},
         {PointSize[Large],Point[{x01,y01}]},Text["(x₀,y₀)",Offset[{10,8},{x01,y01}]]}]
         Export["cyH.pdf",cyH]
```

Out[●]= EcoEvo Package Version 1.6.4 (November 5, 2021)
Christopher A. Klausmeier <christopher.klausmeier@gmail.com>

••• Solve: Solve was unable to solve the system with inexact coefficients. The answer was obtained by solving a corresponding exact system and numericizing the result.

Out[ ]= {{x → 0, y → 0, v → 0, z → 0}, {x → 1., y → 0, v → 0, z → 0},
    {x → 0.0826446, y → 0.0265046, v → 2.91551, z → 0},
    {x → 0, y → 0.06, v → -10.35, z → -2.08333},
    {x → 1.33937, y → 0.06, v → -1.30704, z → -8.7697},
    {x → 0.297813 + 0.339863 𝕚, y → 0.06, v → 2.1017 - 1.11228 𝕚, z → 1.75115 + 1.463 𝕚},
    {x → 0.297813 - 0.339863 𝕚, y → 0.06, v → 2.1017 + 1.11228 𝕚, z → 1.75115 - 1.463 𝕚}}

Eigenvalues of E∗:{-1.24715, 0.00415397 + 0.230094 𝕚, 0.00415397 - 0.230094 𝕚, -0.0200972}

We simulate around

Out[ ]= {v → 2.91551, z → 0, x → 0.182645, y → 0.0365046}

Final time slice of the cycle is

Out[ ]= 32.613

Out[ ]= cyH.pdf