

MEDLINE- acetaminophen tablet
Medline Industries, LP

196 OTCM00023 500mg Acetaminophen Cap

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

for the temporary relief of minor aches and pains associated with

- the common cold
- the flu
- headache
- toothache
- muscular aches
- backache
- the premenstrual and menstrual periods
- for the minor pain from arthritis
- to reduce fever

Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 4,000 milligrams in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Ask a doctor before use if the user

- has liver disease

- is a child with pain of arthritis

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor or pharmacist before use if

- the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- symptoms do not improve
- any new symptoms appear

If pregnant or breast-feeding,

ask a professional before use

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

adults and children 12 years and over	• take 1 to 2 tablets every 6 hours while symptoms persist • do not take more than 6 tablets in 24 hours, unless directed by a doctor
children under 12 years	• ask a doctor


Other information

- store between 20–25°C (68–77°F)
- not made with Natural Rubber Latex
- **do not use if inner seal is broken or missing**

Inactive Ingredients

hydroxy propyl methyl cellulose, polyethylene glycol, pregelatinised Starch, providone K-30, purified Water, sodium starch glycolate,

Package Label



acetaminophen tablet

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53329-196
Route of Administration	ORAL		

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients	
Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z 65AP)	
WATER (UNII: 059QF0KO0R)	
HYDROXYPROPYL METHYLCELLULOSE (UNII: 3NXW29V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE K30 (UNII: U725QWY32X)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	

Product Characteristics			
Color	white ((white to off-white))	Score	score with uneven pieces
Shape	CAPSULE ((Caplet Shape))	Size	17mm
Flavor		Imprint Code	G551
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53329-196-30	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/29/2024	08/31/2026



Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/29/2024	08/31/2026

Labeler - Medline Industries, LP (025460908)

Registrant - Medline Industries, LP (025460908)

Revised: 6/2026

Medline Industries, LP