

# MATH 320/321 (Real Analysis) Notes

Rio Weil

*This document was typeset on July 18, 2021*

## Introduction:

This set of notes is transcribed from UBC's MATH 320/321 (Real Variables I/II) sequence. The course covers the first 9 chapters of Rudin's "Principles of Mathematical Analysis" with occasional omissions & additions. The numbering of the definitions/theorems/examples will follow that used in Rudin for convenience. The structure of these notes is such that they are split into main text (the boxed elements) and side text (everything else). It is possible to solely read the main text for all of the material, but the additional discussion provided by the side text may be useful.

## Contents

<b>1</b>	<b>The Real and Complex Number Systems</b>	<b>3</b>
1.1	The Naturals, Integers, and Rationals . . . . .	3
1.2	Ordered Sets . . . . .	5
1.3	The Least Upper Bound Property . . . . .	6
1.4	Fields and Ordered Fields . . . . .	8
1.5	Consequences of the LUB Property . . . . .	11
1.6	Integer Roots of the Reals . . . . .	12
1.7	Construction of the Reals . . . . .	13
1.8	The Complex Field . . . . .	16
1.9	The Cauchy-Schwartz Inequality . . . . .	20
1.10	Euclidean Space . . . . .	22
<b>2</b>	<b>Basic Topology</b>	<b>25</b>
2.1	Finite and Countable Sets . . . . .	25
2.2	Uncountable Sets . . . . .	28
2.3	Topology of Metric Spaces . . . . .	30
2.4	Closure and Relative Topology . . . . .	35
2.5	Compactness . . . . .	37
2.6	Compactness in $\mathbb{R}^k$ and the Cantor Set . . . . .	40
2.7	Connected Sets . . . . .	44
<b>3</b>	<b>Numerical Sequences and Series</b>	<b>46</b>
3.1	Sequences . . . . .	46
3.2	Subsequences . . . . .	50
3.3	Cauchy Sequences and Completeness . . . . .	51
3.4	Limit Supremum and Limit Infimum . . . . .	54
3.5	Series . . . . .	56
3.6	p-Series and Euler's Number . . . . .	59
3.7	The Ratio and Root Tests . . . . .	61
3.8	Power Series . . . . .	63
3.9	Absolute Convergence . . . . .	66
3.10	Addition and Multiplication of Series . . . . .	67

<b>4</b>	<b>Continuity</b>	<b>69</b>
4.1	Limits and Continuity . . . . .	69
4.2	Topological Characterization of Continuity . . . . .	71
4.3	Continuity and Compactness . . . . .	74
4.4	Uniform Continuity, Connectedness, and IVT . . . . .	76
4.5	Topological Spaces . . . . .	78
<b>5</b>	<b>Differentiation</b>	<b>80</b>
5.1	Derivatives . . . . .	80
5.2	MVT . . . . .	83
5.3	Taylor's Theorem . . . . .	85
5.4	Local Behavior of Functions . . . . .	89
<b>6</b>	<b>The Riemann-Stieltjes Integral</b>	<b>92</b>
6.1	Definition of the Integral . . . . .	92
6.2	Criterion for Integrability . . . . .	95
6.3	Properties of the Integral . . . . .	101
6.4	The Fundamental Theorem of Calculus . . . . .	108
<b>7</b>	<b>Sequences and Series of Functions</b>	<b>113</b>
7.1	Motivating Examples . . . . .	113
7.2	Uniform Convergence . . . . .	116
7.3	Uniform Convergence and Integration . . . . .	123
7.4	Uniform Convergence and Differentiation . . . . .	125
7.5	Equicontinuous Families of Functions . . . . .	128
7.6	The Stone-Weierstrass Theorem . . . . .	132
<b>8</b>	<b>Some Special Functions</b>	<b>136</b>
8.1	Power Series, Revisited . . . . .	136
8.2	The Exponential Function . . . . .	136
8.3	The Logarithm . . . . .	136
8.4	Sine and Cosine . . . . .	136
8.5	Fourier Series . . . . .	136
<b>9</b>	<b>Functions of Several Variables</b>	<b>136</b>
9.1	Banach Fixed Point Theorem . . . . .	136
9.2	Differentiation of Functions of Several Variables . . . . .	136
9.3	The Inverse Function Theorem . . . . .	136

# 1 The Real and Complex Number Systems

## 1.1 The Naturals, Integers, and Rationals

We begin by a review of number systems which are already familiar.

### Definition: The Natural Numbers

The **Naturals**, denoted by  $\mathbb{N}$ , is the set  $\{1, 2, 3, \dots\}$ .

For  $x, y \in \mathbb{N}$ , we have that  $x + y \in \mathbb{N}$  and  $xy \in \mathbb{N}$ , so the naturals are closed under addition and multiplication. However, we note that it is not closed under subtraction; take for example  $2 - 4 = -2 \notin \mathbb{N}$ .

### Definition: The Integers

The **Integers**, denoted by  $\mathbb{Z}$ , is the set  $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ .

The integers are closed under addition, multiplication, and subtraction. However, it is not closed under division; for example,  $1/2 \notin \mathbb{Z}$ .

### Definition: The Rationals (informal)

The **Rationals**, denoted by  $\mathbb{Q}$ , can be defined as  $\left\{\frac{m}{n} : m \in \mathbb{Z}, n \in \mathbb{N}\right\}$ , where  $\frac{m_1}{n_1}$  and  $\frac{m_2}{n_2}$  are identified if  $m_1 n_2 = m_2 n_1$ .

We note that unlike the naturals/integers, the rationals do not have as obvious of a denumeration. This above is a good definition if we already have the same rigorous idea of what a rational number is in our mind; i.e. it works because we have a shared preconceived understanding of a rational number.

If this is not the case, it may help to define the rational numbers more rigorously/formally (even if the definition may be slightly harder to parse). As a second attempt at a definition, we can say that  $\mathbb{Q}$  is the set of ordered pairs  $\{(m, n) : m \in \mathbb{Z}, n \in \mathbb{N}\}$ . However, this is not quite enough as we need a notion of equivalence between two rational numbers (e.g.  $(1, 2) = (2, 4)$ ). Hence, a complete and rigorous definition would be:

### Definition: The Rationals (formal)

The **Rationals**, denoted by  $\mathbb{Q}$ , is the set  $\{(m, n) : m \in \mathbb{Z}, n \in \mathbb{N}\} / \sim$  where  $(m_1, n_1) \sim (m_2, n_2)$  if  $m_1 n_2 = m_2 n_1$ .

Under the formal definition, the rationals are a set of equivalence classes of ordered pairs, under the equivalence relation  $\sim$ . We note that the rationals are closed under addition, subtraction, multiplication, and division.

This formal definition might be slightly harder to parse, so it might be useful to consider an example with a similar flavour. Consider the set  $X = \{m \in \mathbb{Z}\} / \sim$  such that  $m_1 \sim m_2$  if  $m_1 - m_2$  is divisible by 12. This is "clock arithmetic", with equivalence classes  $[0], [1], [2], \dots$  for each hour on an analog clock. A fun side note: If instead of 12 we picked a prime number, we would get a field (we will discuss what this is in a later lecture)!

Note that under this definition,  $(1, 2)$  and  $(2, 4)$  are different representations of the same rational number. With this definition, we would define addition such that  $(m_1, n_1) + (m_2, n_2) = (m_1 n_2 + m_2 n_1, n_1 n_2)$ . Note that  $(2m_1, 2n_2) + (m_2, n_2) = (2m_1 n_2 + 2m_2 n_1, 2n_1 n_2)$  and we can identify  $(m_1 n_2 + m_2 n_1, n_1 n_2)$  with  $(2m_1 n_2 + 2m_2 n_1, 2n_1 n_2)$ . If we choose different representations when we do addition, we might get a different representation in our result, but it will represent the same rational number regardless of the choice of representations we originally chose to do the addition.

A natural question then becomes if the rationals are sufficient for doing all of real analysis. Certainly, it seems as we have a number system that is closed under all our basic arithmetic operations; but is this enough? For example, are we able to take limits just using the rationals? The answer turns out to be no (they are insufficient!) and the following example will serve as one illustration of this fact.

### Example 1.1: Incompleteness of the Rationals

There exists no  $p \in \mathbb{Q}$  such that  $p^2 = 2$ .

We proceed via proof by contradiction. Recall in that these types of proof, we start with a certain wrong assumption, follow a correct/true line of reasoning, reach an eventual absurdity, and therefore conclude that the original assumption was mistaken.

#### Proof

Let us then suppose for the contradiction that there exists  $p = \frac{m}{n}$  with  $p^2 = 2$ . We then have that not both  $m, n$  are even, and hence at least one is odd. Then, we have that  $2 = p^2 = \frac{m^2}{n^2}$  and hence  $m^2 = 2n^2$ , so  $m^2$  is even, implying  $m$  is even. So, let us write  $m = 2k$  for  $k \in \mathbb{Z}$ . Then,  $(2k)^2 = 4k^2 = 2n^2$ , and hence  $2k^2 = n^2$ . Therefore,  $n^2$  is even and hence  $n$  is even.  $m$  and  $n$  are therefore both even, a contradiction. We conclude that no such  $p$  exists.  $\square$

Why can we conclude that not both  $m, n$  are even in the above proof? This is the case as if  $m, n$  we both even, then we could write  $m = 2m', n = 2n'$  for some  $m', n'$ , and then  $p = \frac{m}{n} = \frac{2m'}{2n'} = \frac{m'}{n'}$  which we can continue until either the numerator or denominator is odd. A natural question to consider is how to prove that this process of reducing fractions will eventually conclude. The resolution is to invoke the fundamental theorem of arithmetic, and write  $m, n$  in terms of their unique prime factorization. We are then able to cancel out factors of 2 from the numerator/denominator until at least one is odd.

We note that this example leads us to conclude that the rationals have certain “holes” in them. This is concerning, as there are sequences of rational numbers that tend to  $\sqrt{2}$ . Conversely, it's not as concerning that there is no rational number  $x$  such that  $x^2 = -1$ , as there is no such sequence of rational numbers that is “close to”  $i$  (note that both  $\sqrt{2}$  and  $i$  have not yet been defined, but this will come shortly).

### Example 1.1: Incompleteness of the Rationals

Let  $A = \{p \in \mathbb{Q} : p > 0, p^2 < 2\}$ , and  $B = \{p \in \mathbb{Q} : p > 0, p^2 > 2\}$ . Then,  $\forall p \in A, \exists q \in A$  such that  $p < q$ , and  $\forall p \in B, \exists q \in B$  such that  $q < p$ .



Figure 1: Visualization of sets  $A$  and  $B$ . We note that  $\sqrt{2}$  has not been defined in our formalism yet, but from our prior mathematical intuition it would be what goes in the “hole” of the rationals.

For the proof of this statement, we consider playing a 2 person game. One person is  $\forall$ , one person is  $\exists$ , and we consider if one person has a winning strategy.  $\forall$  goes first, and then  $\exists$  goes next, having seen the choice that  $\forall$  has made. Then, we check if indeed  $p < q$ . If  $p < q$ , then  $\exists$  wins. If  $p \not< q$ , then  $\forall$  wins.

### Proof

Let  $p \in A$ . Then, let  $q = \frac{2p+2}{2+p}$ . Since  $p \in \mathbb{Q}$ , it follows that  $2p+2 \in \mathbb{Q}$  and  $2+p \in \mathbb{Q}$  so  $q \in \mathbb{Q}$ . Furthermore, we have that  $2p+2 > 0$  and  $2+p > 0$ , so  $q > 0$ . We also have that:

$$q^2 = \frac{(2p+2)^2}{(2+p)^2} = 2 + \frac{2(p^2-2)}{(p+2)^2} < 2$$

Where the inequality follows from the fact that  $p^2 < 2$  and hence  $(p^2 - 2) < 0$ . It therefore follows that  $q \in A$ . Finally, we have that:

$$q = p + \frac{2-p^2}{2+p} > p$$

so  $q > p$ , completing the proof of the first part of the claim. The second part is left as an exercise (we note that the same  $q$  can be used).  $\square$

The number  $q = \frac{2p+2}{2+p}$  seems to be pulled out of a hat, but actually comes from a fairly geometric picture (the secant method of approximating roots). Discussion on this topic can be found here: <https://math.stackexchange.com/questions/141774/choice-of-q-in-baby-rudins-example-1-1>.

## 1.2 Ordered Sets

Over the next couple sections, we will be discussing certain properties of sets that will give us a better understanding of the real numbers, and allow us to construct them.

### Definition 1.5: Order

An **order**  $<$  on a set  $S$  is a relation with the following properties:

- (i) For every pair  $x, y \in S$ , exactly one of  $x < y$ ,  $x = y$ , or  $y < x$  is true.
- (ii) For  $x, y, z \in S$ , if  $x < y$  and  $y < z$ , then  $x < z$ .

A point on notation; We note that  $x > y$  means  $y < x$ , and  $x \leq y$  means  $x < y$  or  $x = y$ .

### Definition 1.6: Ordered Sets

An **ordered set** is a pair  $(S, <)$ . We may write just  $S$  if the order can be inferred by the context.

A familiar (and useful) set of examples is  $S = \mathbb{N}$  or  $S = \mathbb{Z}$  or  $S = \mathbb{Q}$ . For these three sets, we have that  $x < y$  if  $y - x$  is positive. For another example, consider the set  $S$  of english words; then the order  $<$  can be the dictionary/lexographic order.

### Definition 1.7: Upper & Lower Bounds

Let  $S$  be an ordered set and  $E \subset S$  (for the duration of these notes, we will follow Rudin's notation, with  $E \subset S$  as a non-strict subset, and  $E \subsetneq S$  as a strict subset).  $E$  is **bounded above** if there exists an element  $\beta \in S$  such that  $\forall x \in E, x \leq \beta$ . Any such  $\beta$  is an **upper bound** of  $E$ . Similarly, we say that  $E$  is **bounded below** if there exists an element  $\alpha \in S$  such that  $\forall x \in E, \alpha \leq x$ . In this case,  $\alpha$  is a **lower bound** of  $E$ .

As an example, one can take  $S = \mathbb{Q}$ ,  $E = A = \{p \in \mathbb{Q} : p > 0, p^2 > 2\}$  (as in Example 1.1(b)). Here,  $E$  is

bounded above, with  $\beta = 2$  as one possible upper bound. to see this is the case, consider that if  $p \in E$ :

$$2 - p = \frac{4 - p^2}{2 + p} > \frac{4 - 2}{2 + p} > 0$$

However, if we take  $S = A$ ,  $E = A$ , then  $E$  is not bounded above as we saw in the example. There is no upper bound of  $A$  in  $A$ . In general, this example reveals the subtle point that "the upper bound of a set" is ill-defined; we need to specify  $E \subset S$ .

### 1.3 The Least Upper Bound Property

#### Definition 1.8: Least Upper Bound & Greatest Lower Bound

Let  $S$  be an ordered set, and let  $E \subset S$  with  $E$  bounded above. If  $\exists \alpha \in S$  such that:

- (i)  $\alpha$  is an upper bound for  $E$
- (ii) If  $\gamma < \alpha$ , then  $\gamma$  is not an upper bound for  $E$

The  $\alpha$  is the **least upper bound**, or **supremum** of  $E$ . This can be denoted as  $\alpha = \sup(E)$ . Analogously, the **greatest lower bound**, or **infimum** of  $E$  (denoted  $\alpha = \inf(E)$ ) is an element  $\alpha \in S$  (if it exists) such that:

- (i)  $\alpha$  is a lower bound for  $E$
- (ii) If  $\gamma > \alpha$ , then  $\gamma$  is not an upper bound of  $E$ .

#### Theorem

If the supremum/infimum of  $E \subset S$  exist, they are unique.

#### Proof

Let  $E \subset S$ . Suppose that there exist  $\alpha_1, \alpha_2$  such that  $\alpha_1 = \sup(E)$  and  $\alpha_2 = \sup(E)$ . If  $\alpha_1 < \alpha_2$ , as  $\alpha_1$  is an upper bound of  $E$ , this contradicts the fact that  $\alpha_2$  is the least upper bound of  $E$ . We reach an identical contradiction if  $\alpha_2 < \alpha_1$ . Therefore we conclude that  $\alpha_1 = \alpha_2$  and the supremum of  $E$  is unique (if it exists). The proof for the infimum is analogous.  $\square$

#### Theorem

If  $E \subset S$  has a maximum element  $\alpha$  (that is, an element such that  $x < \alpha$  for all  $x \in E$ ) then  $\alpha = \sup(E)$ . Similarly, if  $E$  has a minimum element  $\alpha$ , then  $\alpha = \inf(E)$ .

#### Proof

Let  $E \subset S$  and  $\alpha = \max(E)$ . By definition  $\alpha$  is an upper bound of  $E$ , and if  $x < \alpha$  for some  $x \in E$  then  $x$  is not an upper bound of  $E$  as it is not greater than  $\alpha \in E$ . The claim follows (with an identical proof for the minimum).  $\square$

### Example 1.9

- (a) Consider again the sets  $A, B \subset \mathbb{Q}$  from example 1.1.  $A$  is bounded above by any element in  $B$ , and the upper bounds of  $A$  are exactly the elements of  $B$ . Since  $B$  has no smallest member,  $A$  does not have a least upper bound in  $\mathbb{Q}$ .
- (b) Let  $E_1, E_2 \subset \mathbb{Q}$  such that  $E_1 = \{r : \mathbb{Q}, r < 0\}$  and  $E_2 = \{r : \mathbb{Q}, r \leq 0\}$ . Then  $\sup(E_1) = \sup(E_2) = 0$ . Note that this example shows that the supremum can either be contained or not contained in the set;  $0 \notin E_1$  but  $0 \in E_2$ .
- (c) Let  $E \subset \mathbb{Q}$  such that  $E = \{\frac{1}{n} : n \in \mathbb{N}\}$ . Then  $\sup(E) = 1$  and  $\inf(E) = 0$ . This is proven below.

### Proof

$\sup(E) = 1$  immediately follows from the equivalence of the maximum and supremum as proven above. To see that  $\inf(E) = 0$ , first note that 0 is a lower bound for  $E$  as all of the elements of  $E$  are positive. To see that it is the lower bound, take any  $x > 0$ . Then, we have that for any  $n > \frac{1}{x}$ ,  $\frac{1}{n} < x$  and hence  $x$  is not an upper bound of  $E$ . This proves the claim.  $\square$

### Definition 1.10: The LUB/GUB Property

An ordered set  $S$  has the **least upper bound property** if for every  $E \subset S$ , if  $E \neq \emptyset$  and  $E$  is bounded above, then  $E$  has a least upper bound (that is,  $\sup(E)$  exists in  $S$ ). Similarly, an ordered set  $S$  has the **greatest lower bound property** if for every  $E \subset S$ , if  $E \neq \emptyset$  and  $E$  is bounded below, then  $E$  has a greatest lower bound.

We will show in the next theorem that these properties are actually equivalent; before then, we briefly consider two examples.

### Example

$\mathbb{Z}$  has the least upper bound property, while  $\mathbb{Q}$  does not.

### Proof

For the first claim, consider any nonempty  $E \subset \mathbb{Z}$  that is bounded above. Choose any  $x \in E$ . Since  $\mathbb{Z}$  is bounded above, there exist finitely many elements that are greater than  $x$ . Take the maximum of these finitely many elements. This maximum is also the maximum of  $E$ , so it is the supremum of  $E$ . Therefore  $\mathbb{Z}$  has the LUB property as claimed. The second claim immediately follows from Example 1.9(a).  $\square$

### Theorem 1.11

Let  $S$  be an ordered set. Then  $S$  has the LUB property if and only if it has the GUB property.

### Proof

$\Rightarrow$  Let  $S$  be an ordered set with the LUB property. Let  $E \subset S$  with  $E \neq \emptyset$ , with  $E$  bounded below. Let  $L = \{x \in S : x \text{ is a lower bound of } E\}$ .  $L \neq \emptyset$  as  $E$  is bounded below (and hence has at least one lower bound). If  $y \in E$ , then  $y$  is an upper bound for  $L$ . Since  $E$  is nonempty,  $L$  is therefore bounded above. Since  $S$  has the LUB property, then  $\sup(L)$  must exist. Let us call this  $\alpha$ . Then,  $\alpha \leq x \forall x \in E$  (as if  $\gamma < \alpha$ , then  $\gamma$  is not an upper bound of  $L$  and hence  $\gamma \notin E$ ). Hence,  $\alpha$  is a lower bound for  $E$  and hence  $\alpha \in L$ . Since  $\alpha = \sup(L)$  and  $\alpha$  is an upper bound for  $L$ , we have that  $\alpha \geq \gamma \forall \gamma \in L$ . Thus,  $\alpha = \inf(E)$ .

$\Leftarrow$  Left as an exercise. □

## 1.4 Fields and Ordered Fields

### Definition 1.12: Fields

A **field**  $F$  is a set with two binary operations,  $+$  and  $\cdot$  (addition and multiplication) such that the following axioms are satisfied:

- (A1): If  $x, y \in F$ , then  $x + y \in F$ . (Closure under addition)
- (A2):  $x + y = y + x$  for all  $x, y \in F$ . (Commutativity of addition)
- (A3):  $(x + y) + z = x + (y + z)$  for all  $x, y, z \in F$ . (Associativity of addition)
- (A4):  $\exists 0 \in F$  such that  $\forall x \in F, 0 + x = x$ . (Additive identity)
- (A5):  $\forall x \in F, \exists y$  such that  $x + y = 0$ . We can denote  $y = -x$ . (Additive inverse)
- (M1): If  $x, y \in F$ , then  $x \cdot y \in F$ . (Closure under multiplication)
- (M2):  $x \cdot y = y \cdot x$  for all  $x, y \in F$ .
- (M3):  $(x \cdot y) \cdot z = x \cdot (y \cdot z)$  for all  $x, y, z \in F$ . (Associativity under multiplication)
- (M4):  $\exists 1 \in F$  such that  $1 \neq 0$  and  $\forall x \in F, 1 \cdot x = x$ . (Multiplicative identity)
- (M5):  $\forall x \in F$ , exists  $y \in F$  such that  $x \cdot y = 1$ . We can denote  $y = \frac{1}{x}$ . (Multiplicative inverse)
- (D):  $x \cdot (y + z) = x \cdot y + x \cdot z, \forall x, y, z \in F$ . (Distributive law)

Note that A3/M3 show that  $x + y + z$  and  $x \cdot y \cdot z$  are well defined in a mathematical sense; however, associativity may not hold for computers that do math with finite precision!

### Theorem

The additive/multiplicative identities given by (A4)/(M4) and the additive/multiplicative inverses given by (A5)/(M5) are unique.



### Proof

Let  $F$  be an ordered field. Suppose that there exist  $0_1, 0_2 \in F$  such that  $0_1 + x = x$  and  $0_2 + x = x$  for all  $x \in F$ . We then have that:

$$\begin{aligned} 0_1 + 0_2 &= 0_1 + 0_2 \\ 0_1 + 0_2 &= 0_2 + 0_1 & (A2) \\ 0_2 &= 0_1 & (\text{Property of additive identity}) \end{aligned}$$

Which shows that the additive identity is unique. The remaining proofs are left as an exercise.  $\square$

Some easy (and familiar) consequences of the field axioms can be found in Rudin 1.14-1.16. Instead of repeating those here, we will discuss some examples.

The rationals form a field (under the usual notions of addition/multiplication), but the integers do not, as there are no multiplicative inverses (e.g. there exists no integer  $x \in \mathbb{Z}$  such that  $2 \cdot x = 1$ ). The simplest example of a field is  $F = \{0, 1\}$ , with the relations:

$$\begin{aligned} 0 + 0 &= 0 & 0 \cdot 0 &= 1 \\ 0 + 1 &= 0 & 0 \cdot 1 &= 0 \\ 1 + 1 &= 0 & 1 \cdot 1 &= 1 \end{aligned}$$

This field is often called  $\mathbb{F}_2$  or  $F_2$ , and is useful in computer science (where bits can take on two states, 0 or 1). As a slight tangent, a byte (8 bits) can be considered an element of an 8-dimensional vector space over the field  $\mathbb{F}_2$ , where  $+$  would be the XOR operator and  $\cdot$  would be the AND operation.

A generalization of the above example is  $\mathbb{F}_p$  or  $F_p$ , for a prime number  $p$ . This field would consist of the elements  $0, 1, \dots, p-1$ . The addition and multiplication are carried out mod  $p$ . An interesting result is that in general, finite fields must have cardinality of some prime power.

Note that a field cannot have a single element; the field axioms (A4) and (M4) require the existence of distinct additive and multiplicative identities, which a singleton set cannot satisfy.

Although algebra is not the focus of this course, it may be interesting to briefly think about sets with less structure than a field. We start by considering a group.

A **group**  $G$  is a set with a binary operation  $(a, b) \mapsto a \cdot b$  such that the following axioms are satisfied:

- (M1): If  $a, b \in G$ , then  $a \cdot b \in G$  (Closure)
- (M3): For  $a, b, c \in G$ ,  $(a \cdot b) \cdot c = a \cdot (b \cdot c)$  (Associativity)
- (M4): There exists  $1 \in G$  such that  $\forall x \in G, 1 \cdot x = x$ . (Identity)
- (M5):  $\forall x \in G$ , there exists  $y \in G$  such that  $x \cdot y = 1$ . (Inverse)

We note that  $\mathbb{Z}$  is a group under addition, but not under multiplication (due to lack of multiplicative inverses). We can also consider the set of  $2 \times 2$  matrices with integer entries:

$$G = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{Z} \right\}$$

$G$  is again a group under matrix addition, but not under matrix multiplication (as not every matrix in  $G$  is invertible). If we restricted  $G$  to be the set of  $2 \times 2$  invertible matrices, in this case it could form a group under matrix multiplication. A set with slightly more structure than a group (though not quite as structured as a field) is a ring:

A **ring**  $R$  is a set with two binary operations  $(a, b) \mapsto a + b$  and  $(a, b) \mapsto a \cdot b$  such that the following axioms are satisfied:

- (A1): If  $x, y \in R$ , then  $x + y \in R$ . (Closure under addition)
- (A2):  $x + y = y + x$  for all  $x, y \in R$ . (Commutativity of addition)
- (A3):  $(x + y) + z = x + (y + z)$  for all  $x, y, z \in R$ . (Associativity of addition)
- (A4):  $\exists 0 \in R$  such that  $\forall x \in R, 0 + x = x$ . (Additive identity)
- (A5):  $\forall x \in R, \exists y$  such that  $x + y = 0$ . We can denote  $y = -x$ . (Additive inverse)
- (M1): If  $x, y \in R$ , then  $x \cdot y \in R$ . (Closure under multiplication)
- (M3):  $(x \cdot y) \cdot z = x \cdot (y \cdot z)$  for all  $x, y, z \in R$ . (Associativity under multiplication)
- (M4):  $\exists 1 \in R$  such that  $1 \neq 0$  and  $\forall x \in R, 1 \cdot x = x$ . (Multiplicative identity)
- (D1):  $x \cdot (y + z) = x \cdot y + x \cdot z, \forall x, y, z \in R$ . (Left distributivity)
- (D2):  $(y + z) \cdot x = y \cdot x + z \cdot x, \forall x, y, z \in R$ . (Right distributivity)

Rings have the same axioms as fields under addition, but multiplication is not necessarily commutative (this is why an additional distributivity axiom is added), and multiplicative inverses are not required. We note that  $\mathbb{Z}$  and  $G$  are both rings under their respective operations of addition and multiplication.

For the remainder of this course, we will really only be discussing fields; however, they will be the objects of interest in abstract algebra courses!

#### Definition 1.17: Ordered Field

An **Ordered field** is a field  $F$  that is also an ordered set, such that the following axioms are satisfied:

- (i) If  $x, y, z \in F$  and  $y < z$ , then  $x + y < x + z$ .
- (ii) If  $x, y \in F$  and  $x > 0, y > 0$ , then  $x \cdot y > 0$ .

Some properties of ordered fields are discussed in Rudin 1.18. We will again refer the reader to the discussion in the textbook for these properties, and here consider some examples.

$\mathbb{Q}$  is an ordered field, with the familiar order of  $a > b$  if  $a - b > 0$ . A question may arise if  $\mathbb{F}_2$  is an ordered field. A priori fields do not have order, but is it possible to impose an order on this set such that it is an ordered field? The answer turns out to be no.

*Proof.* It suffices to show that both possible orderings leads to a contradiction. Suppose  $0 < 1$ . Then,  $1 = 0 + 1 < 1 + 1 = 0$  which is a contradiction. Suppose instead that  $1 < 0$ . Then,  $0 = 1 + 1 < 1 + 0 = 1$  which again is a contradiction.  $\square$

#### Theorem 1.19: Existence of $\mathbb{R}$

There exists an ordered field  $\mathbb{R}$  which has the LUB property and contains  $\mathbb{Q}$  as a subfield.

What does it mean for  $\mathbb{Q}$  to be a subfield? It means that there exists an injective function  $\mathbb{Q} \mapsto \mathbb{R}$  that respects the properties of an ordered field.

This field  $\mathbb{R}$  happens to be exactly the set of real numbers we are familiar with. However, a natural question is “what does it mean that there exists a field?” It turns out that we can define the reals based on the definitions we have made already. One further question might be that could there not exist several fields with the above property; however, taking the appropriate view, we will find that there is a unique such field.

## 1.5 Consequences of the LUB Property

We will use the least upper bound property and the fact that  $\mathbb{R}$  has  $\mathbb{Q}$  as a subfield to derive its properties.

### Theorem 1.20: Archimedean Property, Density of Rationals/Irrationals in $\mathbb{R}$

- (a) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (b) If  $x, y \in \mathbb{R}$ , and  $x < y$ , then  $\exists p \in \mathbb{Q}$  such that  $x < p < y$ . ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ )
- (c) If  $x, y \in \mathbb{R}$ , and  $x < y$ , then  $\exists \alpha \in \mathbb{R} \setminus \mathbb{Q}$  such that  $x < \alpha < y$ . ( $\mathbb{R} \setminus \mathbb{Q}$  is dense in  $\mathbb{R}$ )

#### Proof

(a) Let  $A = \{nx : n \in \mathbb{N}\}$ . Suppose for the sake of contradiction that the conclusion was false; then  $y$  is an upper bound of  $A$ . Then,  $\alpha = \sup(A)$  exists by the LUB property of  $\mathbb{R}$ . Since  $x > 0$ , we then have that  $\alpha - x < \alpha$  by the property of an ordered field. Hence,  $\alpha - x$  is not an upper bound for  $A$ . Therefore, there exists some  $m \in \mathbb{N}$  such that  $mx > \alpha - x$ . It then follows that  $(m+1)x > \alpha$ . We therefore have found  $m+1 = k \in \mathbb{N}$  such that  $kx > \alpha$ , contradicting  $\alpha$  being the least upper bound of  $A$ .  $\square$

In order to prove (b) and (c), we first prove a stronger version of 1.20(a):

#### Lemma

If  $x, y \in \mathbb{R}$  and  $x > 0$ , then there exists  $n \in \mathbb{Z}$  such that  $(n-1)x \leq y < nx$ .

#### Proof

Suppose  $y \geq 0$ . Let  $A = \{m \in \mathbb{N} : y < mx\} \subset \mathbb{N}$ . By Theorem 1.20 (a), we have that  $A \neq \emptyset$ . Every non-empty subset of  $\mathbb{N}$  has a smallest element (to see this, let  $x \in A$ , and define  $A' = \{y \in A : y \leq x\}$ . This is finite and nonempty and so has a smallest element, and the minimum element of this set will also be a lower bound and hence the minimum element of all of  $A$ ), so let  $n = \min(A)$ . The claim holds for this  $n$ . The case for  $y < 0$  is left as an exercise.  $\square$

#### Proof

(b) Since  $y - x > 0$ , by (a),  $\exists n \in \mathbb{N}$  such that  $1 < n(y - x)$ . Furthermore, by the Lemma we have that  $\exists m \in \mathbb{Z}$  such that  $m - 1 \leq nx < m$  and hence  $m \leq nx + 1$ . From these inequalities we obtain that  $nx < m \leq nx + 1 < ny$ , and therefore  $x < \frac{m}{n} < y$  for some  $m \in \mathbb{Z}$ ,  $n \in \mathbb{N}$ .  $\square$

For the proof of part (c), we will use the result of Theorem 1.21 from the next section, specifically that there exists  $s \in \mathbb{R} \setminus \mathbb{Q}$  such that  $s > 0$  and  $s^2 = 2$ . We will call this  $\sqrt{2}$ .

### Proof

(c) First, we have that  $\sqrt{2} < 2$  as if  $\sqrt{2} = 2$  then  $(\sqrt{2})^2 = 2 = 2^2 = 4$  which is a contradiction, and if  $\sqrt{2} > 2$  then  $2 = \sqrt{2} \cdot \sqrt{2} > 2 \cdot 2 = 4$  by Rudin 1.18 which is yet again a contradiction. Thus,  $\frac{\sqrt{2}}{2} < 1$ .

Let  $x, y \in \mathbb{R}$  such that  $x < y$ . By Theorem 1.20(b), there exists  $p, q \in \mathbb{Q}$  such that  $x < p < q < y$ . Let  $\alpha = p + \frac{\sqrt{2}}{2}(q - p)$ . Then, we have that  $p < \alpha < p + 1(q - p) < q$  and hence  $x < p < \alpha < q < y$ .

If  $\alpha \in \mathbb{Q}$ , then  $\sqrt{2} = 2 \left( \frac{\alpha - p}{q - p} \right) \in \mathbb{Q}$ , which is a contradiction, so it follows that  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ .  $\square$

## 1.6 Integer Roots of the Reals

In this section, we will prove that  $\sqrt{2}$  exists and is an irrational number, but we will not use the fact that  $\mathbb{R} \setminus \mathbb{Q}$  is dense in  $\mathbb{R}$ ; this would of course be circular reasoning. The more general idea will be to prove that for any  $n \in \mathbb{N}$ , there exists  $y \in \mathbb{R}$  such that  $y = x^{1/n}$ . Before this, we prove a lemma.

### Lemma

If  $0 < a < b$  and  $n \in \mathbb{N}$ , then  $0 < b^n - a^n \leq nb^{n-1}(b - a)$

Note that a “Calculus proof” of this Lemma would be to let  $f(x) = x^n$ , and then

$$f(b) - f(a) = f'(c)(b - a) = nc^{n-1}(b - a) \leq nb^{n-1}(b - a)$$

Where we invoke the mean value theorem. But this obviously doesn’t work as we have neither defined a derivative nor proven the mean value theorem. A proper proof would be:

### Proof

Let  $0 < a < b$ . Then, we may factor  $b^n - a^n$  such that:

$$b^n - a^n = (b - a)(b^{n-1} + ab^{n-2} + a^2b^{n-3} + \dots + a^{n-2}b + a^{n-1})$$

The second factor is a sum of  $n$  terms, each positive, and in between 0 and  $b^{n-1}$ . Therefore:

$$b^n - a^n \leq nb^{n-1}(b - a)$$

which proves the claim.  $\square$

We will now state the theorem formally:

### Theorem 1.21: Integer Roots of the Reals

Let  $x \in \mathbb{R}$ ,  $x > 0$ , and  $n \in \mathbb{N}$ . Then, there exists a unique  $y \in \mathbb{R}$  such that  $y > 0$  and  $y^n = x$ .

Note that somewhere in the proof, we will use the fact that  $y \in \mathbb{R}$ ; this statement doesn’t hold for rationals (see Example 1.1) so some property of the reals must come into play somewhere.

## Proof

If  $n = 1$ , then the unique solution is  $y = x$ ; we may therefore assume that  $n \geq 2$ .

**Uniqueness:** Suppose there exist two distinct numbers  $y_1, y_2$  with  $y_1 > 0, y_2 > 0$ , and  $y_1^n = y_2^n = x$ . WLOG, suppose  $0 < y_1 < y_2$ . We then have that  $0 < y_1^n < y_2^n$  which is a contradiction.

**Existence:** We prove existence in three steps.

1. We show that  $E \neq \emptyset$ . Let  $E = \{t \in \mathbb{R} : t > 0, t^n < x\}$ . If  $x < 1$ , then  $x^n < x$ , so  $x \in E$ . If  $x \geq 1$ , then  $\left(\frac{1}{2}\right)^n < \frac{1}{2} < x$ , so  $\frac{1}{2} \in E$ . Therefore,  $E \neq \emptyset$ .
2. We show that  $E$  is bounded above and has a supremum in  $\mathbb{R}$ . If  $t > 1 + x$ , then it follows that  $t^n > t > x$ , so  $t \notin E$ . Hence,  $1 + x$  is an upper bound of  $E$ . By Theorem 1.19 (the LUB property of  $\mathbb{R}$ ), it follows that  $\sup(E) \in \mathbb{R}$  exists.
3. We show that  $y = \sup(E)$  satisfies  $y^n = x$ . As  $\mathbb{R}$  is an ordered field, one of  $y^n < x$ ,  $y^n = x$ , or  $y^n > x$  must be true; we show that the first and third are impossible.
  - (a) Suppose  $y^n < x$ . We will obtain a contradiction by finding  $h > 0$  such that  $(y + h)^n < x$ . (Why is this a contradiction?  $y + h > y$ , so if  $(y + h)^n < x$ , then  $y + h \in E$ , contradicting the fact that  $y$  would be an upper bound of  $E$ ). WLOG, suppose that  $h < 1$ . By the above Lemma, we have that:

$$(y + h)^n - y^n \leq n(y + h)^{n-1}h \leq n(y + 1)^{n-1}h$$

By choosing  $h$  sufficiently small, that is:

$$h < \min \left\{ 1, \frac{x - y^n}{n(y + 1)^{n-1}} \right\}$$

Then  $n(y + 1)^{n-1}h < x^n - y^n$  from which it follows that  $(y + h)^n - y^n < x^n - y^n$  and so  $y + h < x$ , which is the desired contradiction.

- (b) Suppose  $y^n > x$ . We will obtain a contradiction by finding  $h > 0$  such that  $(y - h)^n > x$ . If this is true, then  $y - h$  is an upper bound for  $E$ , contradicting the fact that  $y$  is the least upper bound for  $E$ . WLOG suppose that  $h < 0$ . Again applying the Lemma, we have that:

$$y^n - (y - h)^n \leq ny^{n-1}h$$

By choosing  $h$  sufficiently small, that is:

$$h < \min \left\{ 1, \frac{y^n - x}{ny^{n-1}} \right\}$$

It then follows that:

$$y^n - (y - h)^n \leq ny^{n-1}h < y^n - x$$

and hence  $(y - h)^n > x$ , which is the desired contradiction. □

## 1.7 Construction of the Reals

Theorem 1.19 says that there exists an ordered field that contains  $\mathbb{Q}$  as a subfield. We now go about proving this statement. The construction is fairly technical and hence will be carried out in multiple steps. Some of the steps are left as exercises (one can refer to Rudin for the fully complete construction).

### Step 1: Defining the elements of $\mathbb{R}$

The members of  $\mathbb{R}$  will be proper subsets of  $\mathbb{Q}$ , called cuts.  $\mathbb{R} = \{\text{all cuts}\}$ .

#### Definition: Cuts

A **cut** is a proper subset  $\alpha \subsetneq \mathbb{Q}$  with the three properties:

- (I)  $\alpha \neq \emptyset$
- (II) If  $p \in \alpha$ , then  $q \in \alpha \forall q < p$ .
- (III) If  $p \in \alpha$ , then  $\exists r \in \alpha$  such that  $p < r$ .



Figure 2: Visualization of a cut  $\alpha$ . The real number being described of this cut can be thought of as the number at the right boundary (the arrow).

In a sense, a cut gives us a way of discussing the real numbers (in the way we are familiar with them already) without referring to them directly; much like we could formally define/refer to rationals as equivalence classes of ordered pairs.

As a note, we could very well define cuts to be bounded below rather than above, and the following construction would still work out.

### Step 2: $\mathbb{R}$ is an ordered set

We define  $\alpha < \beta$  to mean  $\alpha \subsetneq \beta$ . We show that this makes  $\mathbb{R}$  into an ordered set. First checking transitivity, we have that if  $\alpha < \beta$  and  $\beta < \gamma$  then  $\alpha < \gamma$  by the fact that set inclusion is transitive. Furthermore, at most one of  $\alpha < \beta$ ,  $\alpha = \beta$ , and  $\beta < \alpha$  hold; to see this is the case, suppose the first two fail. Then,  $\alpha \not\subsetneq \beta$ . Hence,  $\exists p \in \alpha$  with  $p \notin \beta$ . If  $q \in \beta$ ,  $q < p$  and hence  $q \in \alpha$  by (II), so  $\beta \subset \alpha$ , and since  $\beta \neq \alpha$  it follows that  $\beta \subsetneq \alpha$ .

### Step 3: $\mathbb{R}$ has the LUB property

We show that  $\mathbb{R}$  has the LUB property. To see this is the case, let  $A \subset \mathbb{R}$  with  $A \neq \emptyset$ , and suppose that there exists  $\beta \in \mathbb{R}$  that is an upper bound for  $A$ . We will now define  $\gamma = \bigcup_{\alpha \in A} \alpha$  and prove that  $\gamma \in \mathbb{R}$  and  $\gamma = \sup A$  (hence  $A$  has a supremum and  $\mathbb{R}$  has the LUB property).

Since  $A \neq \emptyset$ ,  $\exists \alpha_0 \in A$ , and since  $\alpha_0 \neq \emptyset$  (as it is a cut) and  $\alpha_0 \subset \gamma$ , it follows that  $\gamma \neq \emptyset$ . Next, we have that  $\gamma \subset \beta$ , since  $\alpha \subset \beta$  for every  $\alpha \in A$ , and hence  $\gamma \neq \mathbb{Q}$ , that is,  $\gamma \subsetneq \mathbb{Q}$ . Hence  $\gamma$  satisfies property (I) of a cut.

Take  $p \in \gamma$ . Then  $p \in \alpha_1$  for some  $\alpha_1 \in A$ . If  $q < p$ , then  $q \in \alpha_1$  (as  $\alpha_1$  is a cut) so  $q \in \gamma$ , satisfying property (II).

Next, choose  $r \in \alpha_1$  such that  $r > p$ , then  $r \in \gamma$  (as  $\alpha_1 \subset \gamma$ ) and hence  $\gamma$  satisfies property (III). Hence  $\gamma$  is a cut, and  $\gamma \in \mathbb{R}$ .

Finally, we show that  $\gamma = \sup A$ . Clearly,  $\alpha \leq \gamma$  for all  $\alpha \in A$ , as  $\gamma = \bigcup_{\alpha \in A} \alpha$ , so  $\gamma$  is an upper bound of  $A$ . To show that it is the least upper bound, let  $\delta < \gamma$  be a cut. Then,  $\exists s \in \gamma$  such that  $s \notin \delta$ . Therefore,  $\exists \alpha_2 \in A$  such that  $s \in \alpha_2$ ; hence  $\delta < \alpha_2$ , so  $\delta$  is not an upper bound for  $A$ , giving the desired result.

#### Step 4: Addition on $\mathbb{R}$

##### Definition: Addition

If  $\alpha, \beta \in \mathbb{R}$ , we define  $\alpha + \beta = \{s + t : s \in \alpha, t \in \beta\}$ . Showing that this is a cut is left as an exercise.

##### Definition: Zero

$0^* = \{s \in \mathbb{Q}\}$ . Showing that this is a cut is left as an exercise.

We leave it as an exercise to show that the addition axioms (A1)-(A5) of a field are satisfied under this definition of addition on  $\mathbb{R}$ , with the 0 element as  $0^*$  defined above.

#### Step 5: $\mathbb{R}$ satisfies the Ordered Field Property (i)

We verify that if  $\alpha, \beta, \gamma \in \mathbb{R}$  and  $\beta < \gamma$ , then  $\alpha + \beta < \alpha + \gamma$ .

For every  $s \in \alpha, t \in \beta$ , we have that  $t \in \gamma$  as  $\beta$  is a subset of  $\gamma$  by the definition of order on  $\mathbb{R}$ . Hence,  $s + t \in \alpha + \beta$  implies  $s + t \in \alpha + \gamma$ . Therefore,  $\alpha + \beta \subset \alpha + \gamma$  and hence  $\alpha + \beta \leq \alpha + \gamma$ .

We are then left to check that  $\alpha + \beta \neq \alpha + \gamma$ . To see that this is the case, if  $\alpha + \beta = \alpha + \gamma$ , then  $\beta = \alpha + \beta - \alpha = \alpha + \gamma - \alpha = \gamma$  by the field axioms for addition. Therefore we obtain that  $\beta = \gamma$ , contradicting that  $\beta < \gamma$ . Hence the claim is proven.

As a remark, note that  $0^* < \alpha \iff -\alpha < 0^*$ .

Next we will define multiplication on  $\mathbb{R}$ . A first attempt would be  $\alpha \cdot \beta = \{s \cdot t : s \in \alpha, t \in \beta\}$ . However, this definition is inconsistent with negative numbers from what we require multiplication to accomplish.  $-1 \cdot -1$  would fail to be a cut (it would not contain any negative numbers and hence fail criteria (II)) and  $-1 \cdot 1$  would yield the entirety of the rationals (again not a cut!)

#### Step 6: Positive Multiplication on $\mathbb{R}$

##### Definition: Positive Reals

We define  $\mathbb{R}^+ = \{\alpha \in \mathbb{R} : \alpha > 0^*\}$

##### Definition: Multiplication of Positive Reals

If  $\alpha, \beta \in \mathbb{R}^+$ , we define  $\alpha \cdot \beta = \{r \cdot s : r \in \alpha, r > 0, s \in \beta, s > 0\} \cup \{t \in \mathbb{Q}, t \leq 0\}$ . Equivalently,  $\alpha \cdot \beta = \{p \in \mathbb{Q} : \exists r \cdot s : r \in \alpha, r > 0, s \in \beta, s > 0\}$ . We leave it as an exercise to show that  $\alpha \cdot \beta \in \mathbb{R}$ , and moreover,  $\alpha \cdot \beta \in \mathbb{R}^+$ . Showing this second fact proves ordered field property (ii).

##### Definition: One

$1^* = \{r \in \mathbb{Q} : r < 1\}$ . We again leave showing  $1^* \in \mathbb{R}^+$  as an exercise.

### Step 7: Multiplication on all of $\mathbb{R}$

#### Definition: Multiplication by zero

$$\alpha \cdot 0^* = 0^* = 0^* \cdot \alpha$$

#### Definition: Multiplication

We define general multiplication as below, where the  $\cdot$  on the RHS represents the multiplication of positive reals as outlined in Step 5.

$$\alpha \cdot \beta = \begin{cases} (-\alpha) \cdot (-\beta) & \text{if } \alpha < 0^* \text{ and } \beta < 0^* \\ -((-\alpha) \cdot \beta) & \text{if } \alpha < 0^* \text{ and } \beta > 0^* \\ -(\alpha \cdot (-\beta)) & \text{if } \alpha > 0^* \text{ and } \beta < 0^* \end{cases}$$

We leave it as an exercise to show that the multiplicative axioms (M1)-(M5), as well as the distributive law (D) of a field are satisfied under this definition of multiplication on  $\mathbb{R}$ .

Up until this point, we have shown  $\mathbb{R}$  is an ordered field with the LUB property; we last check that it contains  $\mathbb{Q}$  as a subfield. Note that we do have to be a bit careful with what we mean here;  $\mathbb{R}$  does not literally contain  $\mathbb{Q}$ ;  $\mathbb{R}$  is indeed a set of proper subsets of  $\mathbb{Q}$ . What we really mean is to associate every element of  $\mathbb{Q}$  to an element of  $\mathbb{R}$  such that the field structure is preserved.

### Step 8: $\mathbb{R}$ contains $\mathbb{Q}$ as a subfield

For each  $r \in \mathbb{Q}$ , associate the cut  $r^* = \{p \in \mathbb{Q}, p < r\}$ . We then leave as an easy exercise to verify that  $r^* < s^* \iff r < s$ ,  $r^* + s^* = r + s$ , and  $r^* \cdot s^* = r \cdot s$ . This concludes the construction of the reals.  $\square$

Note that later on in the course, we will construct the real numbers in a different fashion; by considering Cauchy sequences modulo an equivalence relation. Also note that from here on out, it will suffice to have the standard/traditional picture of a "real number" in mind (i.e. infinite decimal expansions) and we will not have to really think about the real numbers as cuts; this was just necessary for the formal construction.

## 1.8 The Complex Field

### Definition 1.24: The Complex Numbers

We define the set of **complex numbers** to be  $\{(a, b) : a, b \in \mathbb{R}\}$ , denoted by  $\mathbb{C}$ . For  $x = (a, b) \in \mathbb{C}$  and  $y = (c, d) \in \mathbb{C}$ , we write  $x = y$  if and only if  $a = c$  and  $b = d$  (note that this is a very different notion of equality compared to the rationals). We define the zero element to be  $(0, 0)$  and the one element to be  $(1, 0)$ . We define addition of complex numbers such that:

$$x + y = (a, b) + (c, d) = (a + c, b + d)$$

And multiplication of complex numbers such that:

$$x \cdot y = (a, b) \cdot (c, d) = (ac - bd, ad + bc)$$



**Theorem 1.25**

The operations of  $+$  and  $\cdot$ , as well as the zero/one elements defined above turn  $\mathbb{C}$  into a field.

**Proof**

It suffices to verify the field axioms (A1)-(A5), (M1)-(M5), and (D) as discussed in 1.12. We will here show (M3), (M4), and (M5) and leave the rest as exercises.

(M3): Let  $x, y, z \in \mathbb{C}$ . We show that  $(x \cdot y) \cdot z = x \cdot (y \cdot z)$ . Let  $x = (a, b)$ ,  $y = (c, d)$ , and  $z = (e, f)$ . We then have that:

$$\begin{aligned}(x \cdot y) \cdot z &= (ac - bd, ad + bc) \cdot (e, f) \\ &= ((ac - bd)e - (ad + bc)f, (ac - bd)f + (ad + bc)e)\end{aligned}$$

We also have that:

$$\begin{aligned}x \cdot (y \cdot z) &= (a, b) \cdot (ce - df, cf + de) \\ &= (a(ce - df) - b(cf + de), a(cf + de) + b(ce - df)) \\ &= (ace - adf - bcf - bde, acf + ade + bce - bdf) \\ &= ((ac - bd)e - (ad + bc)f, (ac - bd)f + (ad + bc)e)\end{aligned}$$

So the claim is proven.

(M4):  $(a, b)(1, 0) = (a \cdot 1 - b \cdot 0, a \cdot 0 + b \cdot 1) = (a, b)$

(M5): Let  $x \in \mathbb{C}$  such that  $x \neq 0$ . Then,  $x = (a, b)$  where either  $a \neq 0$  or  $b \neq 0$  or both. Hence,  $a^2 + b^2 > 0$ . Then, let  $\frac{1}{x} = (\frac{a}{a^2 + b^2}, -\frac{b}{a^2 + b^2})$ . We then have that:

$$\begin{aligned}x \frac{1}{x} &= (a, b) \left( \frac{a}{a^2 + b^2}, -\frac{b}{a^2 + b^2} \right) \\ &= \left( a \frac{a}{a^2 + b^2} - b \left( -\frac{b}{a^2 + b^2} \right), a \left( -\frac{b}{a^2 + b^2} \right) + b \left( \frac{a}{a^2 + b^2} \right) \right) \\ &= \left( \frac{a^2 + b^2}{a^2 + b^2}, -\frac{ab}{a^2 + b^2} + \frac{ab}{a^2 + b^2} \right) \\ &= (1, 0)\end{aligned}$$

Which proves the claim. □

Much like  $\mathbb{Q}$  was a subfield of  $\mathbb{R}$ ,  $\mathbb{R}$  is a subfield of  $\mathbb{C}$ , and there exists a map  $\phi$  from  $\mathbb{R}$  to  $\mathbb{C}$  that respects the field axioms, namely:

$$\begin{aligned}\phi &: \mathbb{R} \longrightarrow \mathbb{C} \\ x &\longmapsto (x, 0)\end{aligned}$$

The theorem below shows that  $\phi$  preserves the field structure:

**Theorem 1.26**

For  $a, b \in \mathbb{R}$  we have that  $(a, 0) + (b, 0) = (a + b, 0)$  and  $(a, 0)(b, 0) = (ab, 0)$ .

**Definition 1.27:  $i$** 

$$i = (0, 1).$$

**Theorem 1.28**

$$i^2 = -1.$$

**Theorem 1.29**

If  $a, b \in \mathbb{R}$ , then  $(a, b) = a + bi$ .

**Proof**

Below are the trivial proofs for the above three theorems.

$$\begin{aligned}(a, 0) + (b, 0) &= (a + b, 0 + 0) = (a + b, 0) \\ (a, 0) \cdot (b, 0) &= (a \cdot b - 0 \cdot 0, a \cdot 0 + 0 \cdot b) = (ab, 0) \\ i^2 &= i \cdot i = (0, 1) \cdot (0, 1) = (-1, 0) = -1 \\ a + bi &= (a, 0) + b(0, 1) = (a, 0) + (0, b) = (a, b)\end{aligned}$$

A slightly odd question may be to ask whether  $\mathbb{C}$  is a subfield of  $\mathbb{R}$ , i.e. does there exist  $\psi : \mathbb{C} \mapsto \mathbb{R}$  such that  $\psi(a + b) = \psi(a) + \psi(b)$  and  $\psi(a \cdot b) = \psi(a) \cdot \psi(b)$ . As we will prove in Chapter 2, we do have that  $|\mathbb{C}| = |\mathbb{R}^2| = |\mathbb{R}|$  (where  $||$  denotes cardinality of the set, to be defined shortly), so there does exist a bijection (i.e. a function that is both injective/one-to-one and surjective/onto; we will define these terms precisely in the next chapter) between the two sets.

As a Lemma, we have that the only injective function  $f : \mathbb{Q} \mapsto \mathbb{R}$  that satisfies  $f(a + b) = f(a) + f(b)$  and  $f(a \cdot b) = f(a) \cdot f(b)$  is  $f(x) = x$ . The proof of this is left as a homework problem (HW2). Therefore, it follows that the only injective function  $g : \mathbb{Q} \times \{0\} \mapsto \mathbb{R}$  (where  $\times$  denotes the Cartesian product) is given by  $g((x, 0)) = x$ . We now give a proof that  $\mathbb{C}$  is not a subfield of  $\mathbb{R}$ .

*Proof.* Suppose then for the sake of contradiction that there exists an injective function  $\psi : \mathbb{Q} \times \{0, 1\} \mapsto \mathbb{R}$ . Such a function then must satisfy  $\psi(i \cdot i) = \psi(-1) = -1$ , and  $\psi(i \cdot i) = \psi(i) \cdot \psi(i) = \psi((0, 1)) \cdot \psi((0, 1)) = 0 \cdot 0 = 0$  which is a contradiction. Hence, no such injection exists from  $\mathbb{Q} \times \{0, 1\}$  to  $\mathbb{R}$  and hence no such injection could exist from  $\mathbb{C}$  ( $\mathbb{R}^2$ ) to  $\mathbb{R}$ . Hence  $\mathbb{C}$  is not a subfield of  $\mathbb{R}$ .  $\square$

**Definition 1.30: Real/Imaginary Parts and Complex Conjugates**

Let  $z = a + bi \in \mathbb{C}$ . Then,  $\text{Re}(z) = a$  is the **real part** of  $z$  and  $\text{Im}(z) = b$  is the **imaginary part** of  $z$ . The **complex conjugate** of  $z$ , denoted by  $\bar{z}$ , is defined as  $\bar{z} = a - bi$ .

### Theorem 1.31

Let  $z, w \in \mathbb{C}$ . It then follows that:

- (a)  $\overline{z + w} = \bar{z} + \bar{w}$ .
- (b)  $\overline{zw} = \bar{z} \cdot \bar{w}$ .
- (c)  $z + \bar{z} = 2\operatorname{Re}(z)$ ,  $z - \bar{z} = 2i\operatorname{Im}(z)$ .
- (d)  $z\bar{z}$  is real and positive (except when  $z = 0$ ).

### Proof

We prove (d). We have that:

$$z\bar{z} = (a + bi)(a - bi) = a^2 + b^2$$

$a^2 + b^2 \geq 0$ , and  $a^2 + b^2 = 0 \iff a = 0, b = 0$  which proves the claim.  $\square$

### Definition 1.32: Absolute Value

We define the **absolute value**  $|z|$  of a complex number  $z$  as  $|z| = \sqrt{z\bar{z}}$ . Note that if  $a \in \mathbb{R}$  and  $z = (a, 0)$ , then

$$|z| = \sqrt{a^2} = \begin{cases} a & \text{if } a \geq 0 \\ -a & \text{if } a < 0 \end{cases}$$

Hence if  $a \in \mathbb{R}$ , we can define  $|a| = |(a, 0)|$ .

### Theorem 1.33

Let  $z, w \in \mathbb{C}$ .

- (a)  $|z| \geq 0$ ,  $|z| = 0 \iff z = 0$ .
- (b)  $|\bar{z}| = |z|$ .
- (c)  $|z||w| = |zw|$ .
- (d)  $|\operatorname{Re}(z)| \leq |z|$ ,  $|\operatorname{Im}(z)| \leq |z|$ .
- (e)  $|z + w| \leq |z| + |w|$ .

### Proof

We prove (d) and (e). Let  $z, w \in \mathbb{C}$ , with  $z = a + bi$ . For (d) we have that  $\operatorname{Re}(z) = a$ , so

$$|\operatorname{Re}(a)| = \sqrt{a^2} \leq \sqrt{a^2 + b^2} = |z|$$

And an equivalent proof follows for  $\operatorname{Im}(z)$ . For (e), we have that:

$$\begin{aligned} |z + w|^2 &= (z + w)(\overline{z + w}) \\ &= z\bar{z} + z\bar{w} + w\bar{z} + w\bar{w} \\ &= |z|^2 + 2\operatorname{Re}(z\bar{w}) + |w|^2 \\ &\leq |z|^2 + 2|\operatorname{Re}(z\bar{w})| + |w|^2 && (|x| \geq x) \\ &= |z|^2 + 2|z\bar{w}| + |w|^2 && (1.33(d)) \\ &= |z|^2 + 2|z||\bar{w}| + |w|^2 && (1.33(c)) \\ &= |z|^2 + 2|z||w| + |w|^2 && (1.33(b)) \\ &= (|z| + |w|)^2 \end{aligned}$$

The claim follows by taking square roots on both sides. □

## 1.9 The Cauchy-Schwartz Inequality

Recall the summation notation:

$$x_1 + x_2 + \dots + x_n = \sum_{j=1}^n x_j$$

### Theorem 1.35: Cauchy-Schwartz Inequality

Let  $a_1, \dots, a_n, b_1, \dots, b_n \in \mathbb{C}$ . We then have that:

$$\left| \sum_{j=1}^n a_j \bar{b}_j \right|^2 \leq \left( \sum_{j=1}^n |a_j|^2 \right) \left( \sum_{j=1}^n |b_j|^2 \right)$$

Note that in the above theorem, both the RHS and the LHS are real numbers (check!) so the equality makes sense (recall that there is no order on  $\mathbb{C}$ ; in fact, it is impossible to define one).

A geometric interpretation of the above inequality is as follows. Let  $\mathbf{a}, \mathbf{b}$  be vectors in  $\mathbb{C}^n$ . Then,  $\langle \mathbf{a}, \mathbf{b} \rangle = \sum_{j=1}^n a_j \bar{b}_j$  is the inner product of  $\mathbf{a}$  and  $\mathbf{b}$ . Then, the inequality says that  $|\langle \mathbf{a}, \mathbf{b} \rangle|^2 \leq \langle \mathbf{a}, \mathbf{a} \rangle \cdot \langle \mathbf{b}, \mathbf{b} \rangle$ .

### Proof

Define  $A = \sum_{j=1}^n |a_j|^2$ ,  $B = \sum_{j=1}^n |b_j|^2$ , and  $C = \sum_{j=1}^n a_j \bar{b}_j$ . If  $B = 0$  (that is, all of the  $b_j$ s are zero) then the LHS/RHS are both zero and we are done. So, let us assume that  $B > 0$ . Let  $\lambda \in \mathbb{C}$ , and we then have that:

$$\begin{aligned} 0 &\leq \sum_{j=1}^n |a_j + \lambda b_j|^2 \\ &= \sum_{j=1}^n (a_j + \lambda b_j)(\bar{a}_j + \bar{\lambda} \bar{b}_j) \\ &= \sum_{j=1}^n |a_j|^2 + \bar{\lambda} \sum_{j=1}^n a_j \bar{b}_j + \lambda \sum_{j=1}^n \bar{a}_j b_j + |\lambda|^2 \sum_{j=1}^n |b_j|^2 \\ &= A + \bar{\lambda} C + \lambda \bar{C} + |\lambda|^2 B \end{aligned}$$

This inequality holds for any  $\lambda$ ; it therefore holds for  $\lambda = -\frac{C}{B}$ , so:

$$\begin{aligned} 0 &\leq A - \frac{\bar{C}}{B} C - \frac{C}{B} \bar{C} + \frac{C \bar{C}}{B^2} B \\ &= A - \frac{|C|^2}{B} \end{aligned}$$

So we therefore obtain that  $|C|^2 \leq AB$  which is the desired inequality. □

A natural question given any inequality is when does equality hold; the answer turns out to be if the vectors are linearly independent, that is, at least one of  $\mathbf{a} = \alpha \mathbf{b}$  and  $\mathbf{b} = \beta \mathbf{a}$  ( $\alpha, \beta \in \mathbb{C}$ ) hold. Note that we only require one of the two relations to hold; in the case that one of  $\mathbf{a}, \mathbf{b}$  are  $\mathbf{0}$  (the vector of all zeros) both equalities cannot be true. It is left as a homework problem to verify equality in the Cauchy-Schwartz inequality if and only if at least one of the two conditions holds (HW3).

## 1.10 Euclidean Space

### Definition 1.36: Euclidean k-space

If  $k \in \mathbb{N}$ , define  $\mathbb{R}^k$  as the set of  $k$ -tuples of real numbers:

$$\mathbb{R}^k = \{\mathbf{x} = (x_1, x_2, \dots, x_k) : x_1, x_2, \dots, x_k \in \mathbb{R}\}$$

We can then define vector addition as:

$$\mathbf{x} + \mathbf{y} = (x_1 + y_1, x_2 + y_2, \dots, x_k + y_k)$$

And scalar multiplication (for  $\alpha \in \mathbb{R}$ ) to be:

$$\alpha \mathbf{x} = (\alpha x_1, \alpha x_2, \dots, \alpha x_k)$$

These operations make  $\mathbb{R}^k$  into a vector space over the real field. We can define the inner product over  $\mathbb{R}^k$  to be:

$$(\mathbf{x}, \mathbf{y}) = \mathbf{x} \cdot \mathbf{y} = \sum_{j=1}^k x_j y_j$$

This allows us to define the norm of  $\mathbf{x}$  to be:

$$|\mathbf{x}| = \sqrt{\mathbf{x} \cdot \mathbf{x}} = \left( \sum_{j=1}^n x_j^2 \right)^{1/2}$$

$\mathbb{R}^k$  with the above inner product and norm is called **Euclidean k-space**.

We briefly remark that the above inner product we defined agrees with the inner product we defined over  $\mathbb{C}^k$ ; we can identify  $r \in \mathbb{R}$  with  $(r, 0) \in \mathbb{C}$ , and hence recognize that  $\mathbb{R}^k \subset \mathbb{C}^k$  where the imaginary part of each coordinate is zero. Then, for the inner product we get the exact same result, as  $\bar{b}_j = b_j$  for any complex numbers with imaginary part zero. From this we can conclude that the Cauchy-Schwartz inequality also holds in  $\mathbb{R}^k$ .

Note that although the field  $\mathbb{C}$  is  $\mathbb{R}^2$  with multiplication defined as in Definition 1.24, in general vector multiplication on  $\mathbb{R}^n$  is not well defined. That is, we cannot make  $\mathbb{R}^n$  into a field in general; though we can make it into a vector space, which has slightly less structure.

One possibly familiar notion of vector multiplication in  $\mathbb{R}^3$  is the cross product. For  $\mathbf{x} = (x_1, x_2, x_3)$  and  $\mathbf{y} = (y_1, y_2, y_3)$ , the cross product is defined as:

$$\mathbf{x} \times \mathbf{y} = (x_2 y_3 - x_3 y_2, x_3 y_1 - x_1 y_3, x_1 y_2 - x_2 y_1)$$

However, the cross product does not satisfy properties that would be necessary to make  $\mathbb{R}^3$  a field. For one, it is not commutative, but anticommutative;  $\mathbf{x} \times \mathbf{y} = -\mathbf{y} \times \mathbf{x}$ . One might ask whether vectors in  $\mathbb{R}^3$  have well-defined inverses, but even before that, there does not exist an identity vector in  $\mathbb{R}^3$  under the cross product! In fact,  $\mathbb{R}^3$  under vector addition and cross product multiplication can be viewed as a noncommutative ring without an identity.

Note that there is a more general notion of a “wedge product” between vectors in  $\mathbb{R}^n$ . We are in a sense very “lucky” that in  $\mathbb{R}^3$ , the wedge product of two vectors returns another vector in  $\mathbb{R}^3$ .

**Theorem 1.37**

Let  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}^k$ , and  $\alpha \in \mathbb{R}$ . Then:

- (a)  $|\mathbf{x}| \geq 0$
- (b)  $|\mathbf{x}| = 0 \iff \mathbf{x} = (0, \dots, 0)$ . This is often denoted as  $\mathbf{0}$ , the “zero vector”.
- (c)  $|\alpha \mathbf{x}| = |\alpha| |\mathbf{x}|$
- (d)  $|\mathbf{x} \cdot \mathbf{y}| \leq |\mathbf{x}| |\mathbf{y}|$
- (e)  $|\mathbf{x} + \mathbf{y}| \leq |\mathbf{x}| + |\mathbf{y}|$
- (f)  $|\mathbf{x} - \mathbf{z}| \leq |\mathbf{x} - \mathbf{y}| + |\mathbf{y} - \mathbf{z}|$

(e) and (f) are often called “triangle inequalities”; a visual intuition for these inequalities is given in the following figure:



Figure 3: Visual picture for Theorem 1.37(f), drawn in  $\mathbb{R}^2$ . Suppose we started at  $\mathbf{x}$  and wanted the shortest path to  $\mathbf{z}$ ; we could try walking directly to  $\mathbf{z}$ , or we could try walking somewhere else first ( $\mathbf{y}$ ) and then to  $\mathbf{z}$ . However, the theorem tells us that the direct path will always be shorter in Euclidean space.

Note that equality in part (f) arises if and only if  $\mathbf{y}$  lies on the line segment between  $\mathbf{x}$  and  $\mathbf{z}$ .

**Proof**

(a)-(c) are immediate, and (d) immediately follows from Theorem 1.35 (Cauchy-Schwartz). For (e), we have that:

$$\begin{aligned}
 |\mathbf{x} + \mathbf{y}|^2 &= (\mathbf{x} + \mathbf{y}) \cdot (\mathbf{x} + \mathbf{y}) \\
 &= |\mathbf{x}|^2 + 2\mathbf{x} \cdot \mathbf{y} + |\mathbf{y}|^2 \\
 &\leq |\mathbf{x}|^2 + 2|\mathbf{x}| |\mathbf{y}| + |\mathbf{y}|^2 \\
 &\leq |\mathbf{x}|^2 + 2|\mathbf{x}| |\mathbf{y}| + |\mathbf{y}|^2 \quad (1.37(d)) \\
 &= (|\mathbf{x}| + |\mathbf{y}|)^2
 \end{aligned}$$

And the claim follows by taking square roots on both sides. For (f), substitute  $\mathbf{x} \mapsto \mathbf{x} - \mathbf{y}$  and  $\mathbf{y} \mapsto \mathbf{y} - \mathbf{z}$  into (e).  $\square$

Though we discuss the Euclidean norm here, it may also be of interest to consider/discuss other norms. One example is the  $L_1$  norm (c.f. the norm discussed in Definition 1.36, which is the  $L_2$  norm), which is the sum of the absolute values of each of the components. For  $\mathbf{x} = (x_1, x_2, \dots, x_n)$  and  $\mathbf{y} = (y_1, y_2, \dots, y_n)$  we have that:

$$|\mathbf{x}|_1 = |x_1| + |x_2| + \dots + |x_n|, \quad |\mathbf{x} - \mathbf{y}|_1 = |x_1 - y_1| + |x_2 - y_2| + \dots + |x_n - y_n|$$

The  $L_1$  norm is often called the “Taxicab norm” or the “Manhattan norm” as the way it quantifies distance is akin to walking in discrete NSEW chunks; much like a taxi running through a grid-like New York City!



Figure 4: Visual comparison of the  $L_1$  and  $L_2$  norms in  $\mathbb{R}^2$ .

We are free to generalize this notion to the  $L_n$  norm, and we may also define the  $L_\infty$  norm, which for  $\mathbf{x} \in \mathbb{R}^n$  is defined as:

$$|\mathbf{x}|_\infty = \max_i |x_i|$$

In general for any  $\mathbf{x} \in \mathbb{R}^n$ , we have that  $|x|_1 \geq |x|_2 \geq |x|_3 \geq \dots \geq |x|_\infty$ . We note that we can generalize these norms to the cases where we have infinite components:

$$\|\mathbf{x}\|_p = \left( \sum_{i=1}^{\infty} |x_i|^p \right)^{1/p} < \infty \quad \|f\|_p \equiv \left( \int_S |f|^p d\mu \right)^{1/p} < \infty$$

Which allow us to define norms for function spaces. However, a detailed discussion of these are beyond the scope of this course (to be covered in a later course in functional analysis!) Moreover, we haven't even defined what an infinite sum or integral are yet, which we will get to in later chapters.



## 2 Basic Topology

### 2.1 Finite and Countable Sets

This chapter is split into two portions; the first looks at counting, what it means for us to say that two sets have the same number of elements, and concludes with a classic theorem of Cantor concerning uncountable sets. The second part looks at the topology of metric spaces, before moving onto the topology of the real numbers.

Let us then begin with our discussion of counting. If we consider counting how many bananas there are on a table (say there are 10 bananas), then what we are formally doing is establishing a correspondence between each ball on the table with an element in the set  $\{1, \dots, 10\}$ . When we refer to the number of elements in a set, it will be good to keep in mind that we are establishing functions between sets. Although we have been discussing functions with some frequency in the course already, we give a definition below for completeness.

#### Definition 2.1: Functions

Let  $A, B$  be sets. Then, a map that associates each element  $x \in A$  with a unique element denoted as  $f(x) \in B$  is a **function**  $f : A \rightarrow B$ . We then define  $A$  as the **domain** of  $f$  and the set  $\{f(x) : x \in A\}$  as the **range**. For  $E \subset A$ , we call  $f(E) = \{f(x) : x \in E\}$  the **image** of  $E$  under  $f$ . For  $F \subset B$ , we call  $f^{-1}(F) = \{x \in A : f(x) \in F\}$  the **preimage** of  $F$ .

#### Definition 2.2: Injective/Surjective Functions

Let  $f : A \mapsto B$  be a function. If for  $x_1, x_2 \in A$  we have that  $f(x_1) = f(x_2) \implies x_1 = x_2$  (or equivalently,  $x_1 \neq x_2 \implies f(x_1) \neq f(x_2)$ ), then we say that  $f$  is **injective**, or **one-to-one**. If for all  $y \in B$  there exists  $x \in A$  such that  $y = f(x)$ , then we say that  $f$  is **surjective**, or **onto**. If a function is both injective and surjective, it is **bijective**.

Intuitively, we can think of injectivity as implying each element in  $B$  being reached at most once, and surjectivity implying that each element in  $B$  is reached at least once.

#### Definition 2.3: Cardinality & Equivalence

Let  $A, B$  be sets. We say that  $A, B$  have the same **cardinality** if there exists  $f : A \mapsto B$  such that  $f$  is bijective. We can denote this as  $A \sim B$  where  $\sim$  indicates an **equivalence relation**. An equivalence relation has three properties:

- (a) Reflexivity:  $A \sim A$ .
- (b) Symmetry: If  $A \sim B$  then  $B \sim A$ .
- (c) Transitivity: If  $A \sim B$  and  $B \sim C$  then  $A \sim C$ .

As a point of notation,  $|S|$  denotes the cardinality of the set  $S$ .

We get (a) from each set having a bijection to itself (i.e. the identity function), (b) from the fact that if there exists a bijection  $f : A \mapsto B$ , then there must exist an inverse  $f^{-1} : B \mapsto A$ , and (c) from if there exist bijections  $f : A \mapsto B$  and  $g : B \mapsto C$  then the composition  $g \circ f : A \mapsto C$  will also be a bijection.

### Definition 2.4: Countability

First, we denote  $J_n = \{1, 2, 3, \dots, n\}$  and  $J = \mathbb{N} = \{1, 2, 3, \dots\}$ . Let  $A$  be a set. We say that  $A$  is **finite** if it has a finite number of elements, that is, there exists  $n \in \mathbb{N}$  such that  $A \sim J_n$ . A set  $A$  is **infinite** if it is not finite, and we cannot put  $A$  in bijection with  $J_n$  for any  $n \in \mathbb{N}$ . We say that  $A$  is **countable** if  $A \sim \mathbb{N}$ , and **uncountable** otherwise.

Note that the above definition gives us a useful notion for what sets we can consider countable; if we can enumerate a set with the naturals, this yields a bijection with  $\mathbb{N}$  and hence the set must be countable.

We here give some additional properties concerning cardinalities of sets, which may be useful:

- $|A| \leq |B| \iff \exists f : A \mapsto B$  such that  $f$  is injective
- $|A| \geq |B| \iff \exists f : A \mapsto B$  such that  $f$  is surjective
- $|A| \leq |B|$  and  $|A| \geq |B| \implies |A| = |B|$

### Example 2.5

$\mathbb{Z}$  is countable. To see this, consider the function:

$$f = \begin{cases} \frac{n}{2} & n \text{ is even} \\ -\frac{n-1}{2} & n \text{ is odd} \end{cases}$$

$f$  is a bijection (check!) and hence  $\mathbb{N} \sim \mathbb{Z}$ .

The above example serves as a bit of a warning sign. Even though  $\mathbb{N} \subsetneq \mathbb{Z}$  and  $\mathbb{Z}$  has “more elements”, we still find that the two sets have the same cardinality. A similar example is given by  $\mathbb{N}$  and the set of all even natural numbers (which we may denote  $2\mathbb{N}$ ); the bijection  $f(n) : n \mapsto 2n$  between these two sets shows that  $\mathbb{N} \sim 2\mathbb{N}$ , even though  $2\mathbb{N}$  is a strict subset of  $\mathbb{N}$ .

### Theorem 2.8

A subset of a countable set is either finite or countable.

### Proof

(Sketch) The countability of  $A$  implies that  $A = \{a_1, a_2, a_3, a_4, a_5, \dots\}$  (in other words, we can enumerate the elements using  $\mathbb{N}$ ). Let  $S \subset A$ . Then,  $S = \{a_1, \cancel{a_2}, a_3, a_4, \cancel{a_5}, \dots\}$ , that is,  $A$  with some (or none) of the elements removed. Now, we can rename all the elements with  $a_1, a_2, \dots$ ; what we have left is again an enumeration, so it is yet again (at most) countable.

One potentially useful fact is that if we have a set  $S$  and a function  $f : \mathbb{N} \mapsto S$  such that  $f$  is surjective, then  $S$  is at most countable.

*Proof.* Let  $T = \{n \in \mathbb{N} : f(n) \neq f(m), \forall m = 1, 2, \dots, n\}$ . We restrict  $f : T \mapsto S$ , then  $f$  is injective by constructive. It is still surjective, hence  $T \sim S$ . Since  $T \subset \mathbb{N}$ , by Theorem 2.8,  $S$  is finite or countable.  $\square$

### Theorem 2.12

Let  $E_1, E_2, \dots$  be countable sets (i.e. we have a countable number of countable sets). Define  $S = \bigcup_{n=1}^{\infty} E_n$ . Then,  $S$  is countable.

### Proof

Write  $E_n = \{x_{n1}, x_{n2}, x_{n3}, \dots\}$  (which we can do as each of the  $E_n$ s are countable). Then, we form an array:

$$\begin{array}{rcll} E_1 & = & \cancel{x_{11}} & x_{12}^{\nearrow} & x_{13}^{\nearrow} & \cdots \\ E_2 & = & x_{21} & \cancel{x_{22}} & x_{23}^{\nearrow} & \cdots \\ E_3 & = & x_{31} & x_{32} & \cancel{x_{33}} & \cdots \\ & \vdots & & & & \end{array}$$

Then, we can re-number the elements along the diagonal lines (i.e.  $x_{11}, x_{21}, x_{12}, x_{31}, x_{22}, x_{13}, \dots$ ). This new enumeration corresponds to a countable set. From there, we let  $T \subset \mathbb{N}$  be the remaining labels in the enumeration after removing the repeated elements from the sequence. Then,  $T \sim S$ , and hence  $S$  is at most countable.  $S$  cannot be finite as  $E_1 \subset S$  and  $E_1$  is not finite. Hence  $S$  is countable.  $\square$

### Corollary 2.13: $\mathbb{Q}$ is Countable.

- If  $A$  is countable, the set of  $n$ -tuples of  $(a_1, \dots, a_n)$  is also countable for any  $n \in \mathbb{N}$ .
- $\mathbb{Q}$  is countable.

We defined  $\mathbb{Q}$  as pairs of integers, but by the first part of the corollary (which follows immediately by application of Theorem 2.12)  $\mathbb{Z}^2$  (the set of pairs of integers) has equal cardinality to  $\mathbb{Z}$ , and since  $\mathbb{Q}$  is a subset of the set of pairs of integers,  $\mathbb{Q}$  is countable.

Another way we can formalize this argument: if we let  $X_t = \left\{ \frac{m}{n} : m \in \mathbb{Z}, n \in \mathbb{N}, |m| \leq t, n \leq t \right\}$ , then we have that  $|X_t|$  is at most  $t \cdot (2t + 1)$  (the first term being the choices for  $n$ , the second term being the choices for  $m$ ). Of course, the cardinality of  $X_t$  is actually less than that as we would have to remove repeats. Nonetheless, we have that  $\mathbb{Q} = \bigcap_{t=1}^{\infty} X_t$  and hence  $\mathbb{Q}$  would be countable by Theorem 2.12.

From the discussion of today, we have established that  $|\mathbb{N}| = |\mathbb{Z}| = |\mathbb{Q}|$ . Does  $\mathbb{R}$  also have equal cardinality to these sets? Do infinite sets in general have the same cardinality? The answer turns out to be no for both of these questions. We will answer the first question in the next lecture (when we discuss Cantor diagonalization, a highlight of the course), but we can discuss the second statement now. First, we make a definition:

### Definition: Power Sets

Let  $A$  be a set. Then, the **power set** of  $A$ , denoted  $\mathcal{P}(A)$ , is the set of all subsets of  $A$ .

An interesting theorem then follows:

### Theorem: Power Set Cardinality

Let  $A$  be a set. Then,  $|A| < |\mathcal{P}(A)|$ .

### Proof

Suppose for the sake of contradiction that there exists a surjection  $f : A \rightarrow \mathcal{P}(A)$  (this would imply that  $|A| \geq |\mathcal{P}(A)|$ , so by showing this is false, we obtain the desired result). Then, each element  $x \in A$  gets mapped to some subset of  $A$ . We either have that  $x$  belongs to the subset that it gets mapped to, or it doesn't. Therefore, we can define a new subset  $B \subset A$ , such that:

$$B = \{x \in A : x \notin f(x)\}$$

In other words, the set of all  $x$ s that are not in the subset that they get mapped to by  $f$ . Since  $f$  is surjective, there must be an element  $y \in A$  such that  $f(y) = B$ . One of  $y \in B, y \notin B$  must be true. If  $y \in B$ , by construction of  $B$  we have that  $y$  is not in the subset that it gets mapped to by  $f$ , which is a contradiction. If  $y \notin B$ , by definition of  $B$ ,  $y \in B$  as it is not in the subset that it gets mapped to, yet again a contradiction. Therefore, we conclude that no  $y \in A$  exists such that  $f(y) = B$ , and hence, no surjective  $f$  exists such that  $f : A \rightarrow \mathcal{P}(A)$ . Hence,  $|A| < |\mathcal{P}(A)|$ .  $\square$

An interesting consequence of this theorem is that for a countable set  $A$ , we then have that  $\mathcal{P}(A)$  is an infinite set which has greater cardinality! For example,  $|\mathbb{N}| < |\mathcal{P}(\mathbb{N})|$ . Moreover, this gives rise to an infinite number of cardinalities in ascending order;  $|\mathbb{N}| < |\mathcal{P}(\mathbb{N})| < |\mathcal{P}(\mathcal{P}(\mathbb{N}))| < \dots$  and so on.

## 2.2 Uncountable Sets

### Theorem 2.14: Existence of Uncountable Sets

Let  $A = \{(b_1, b_2, \dots), b_n \in \{0, 1\}\}$  be the set of binary sequences. Then,  $A$  is uncountable.

### Proof

It suffices to show that every countable subset of  $A$  is a proper subset of  $A$ . Let  $E \subset A$  be countable, and let  $E = \{S^{(1)}, S^{(2)}, S^{(3)}, \dots\}$ . To show that  $E$  is a proper subset, we show that there exists a sequence  $S \in A \setminus E$ . To construct such an  $S$ , let us put the elements of  $E$  in an array.

$$\begin{array}{ccccccc} S^{(1)} & = & \boxed{b_1^1} & b_2^1 & b_3^1 & \dots \\ S^{(2)} & = & b_1^2 & \boxed{b_2^2} & b_3^2 & \dots \\ S^{(3)} & = & b_1^3 & b_2^3 & \boxed{b_3^3} & \dots \end{array}$$

Then, define:

$$\tilde{b}_n^n = \begin{cases} 1 & \text{if } b_n^n = 0 \\ 0 & \text{if } b_n^n = 1 \end{cases}$$

I.e.  $\tilde{b}_n^n$  is the bit flip of  $b_n^n$ . Then, let  $S = (\tilde{b}_1^1, \tilde{b}_2^2, \tilde{b}_3^3, \dots)$ , that is,  $S$  is the sequence of bit-flipped diagonal elements of the original array. By construction,  $S \neq S^{(k)}$  as for any  $S^{(k)} \in E$  as  $S$  differs at the  $k$ th position. Hence,  $S \notin E$  and therefore  $E \subsetneq A$ .  $\square$

The above proof is a very famous argument, invented by the mathematician George Cantor. The discovery that there exist sets with greater cardinality than  $\mathbb{N}$  was initially quite controversial in the math community!

### Corollary: $\mathbb{R}$ is Uncountable

$\mathcal{P}(\mathbb{N})$  (the power set of  $\mathbb{N}$ ) is uncountable.  $\mathbb{R}$  is uncountable.

#### Proof

Although we showed that  $\mathcal{P}(\mathbb{N})$  was uncountable last lecture, we show this in an alternative way by considering a bijection between  $\mathcal{P}(\mathbb{N})$  and the set  $A$  of binary sequences. To do this, consider that we can associate a subset  $T \subset \mathbb{N}$ ,  $T \in \mathcal{P}(\mathbb{N})$  with the sequence corresponding to:

$$b_n = \begin{cases} 1 & \text{if } n \in T \\ 0 & \text{if } n \notin T \end{cases}$$

Since  $A$  is uncountable, it follows that  $\mathcal{P}(\mathbb{N})$  is uncountable. The second statement in the corollary follows (roughly) by considering  $\mathbb{R}$  represented in binary, though this requires more justification than what we present here (we will show below that a subset of  $\mathbb{R}$  is uncountable).  $\square$

### Theorem

$[0, 1] \subset \mathbb{R}$  is uncountable.

#### Proof

(Sketch) We construct a bijection from  $[0, 1]$  to  $A$ . Let  $x \in [0, 1]$ , and let  $b_1$  be the largest integer such that  $N_1 = \frac{b_1}{2} \leq x$ . Then, let  $b_2 \in \{0, 1\}$  be the largest integer such that  $N_2 = \frac{b_1}{2} + \frac{b_2}{2^2} \leq x$ . We can continue dividing  $[0, 1]$  in half in this way, approximating  $x$  by powers of 2 (a decimal expansion in binary). Then, let  $E(x) = \{N_1, N_2, N_3, \dots\}$ . By construction,  $E(x)$  is bounded above by  $x$  and nonempty. Hence,  $\sup(E(x))$  exists and is unique, and in fact is equal to  $x$  (note that we are in essence constructing an "infinite series" where the sequence of partial sums is increasing and bounded, approaching  $x$  from the left). Doing this we can associate  $(b_1, b_2, b_3, \dots) \in A$  with every number  $x \in [0, 1]$  and therefore  $[0, 1] \sim A$ . Hence  $[0, 1]$  is uncountable.  $\square$

It might be worth investigating why a more naive approach to the above argument may fail. Let  $D$  be the sets of all decanary sequences, i.e. all sequences of the form  $(d_1, d_2, d_3, \dots)$  where  $d_n \in \{n \in \mathbb{Z}, 0 \leq n \leq 9\}$ . Then, it might be tempting to say that the function:

$$\begin{aligned} f : D &\longrightarrow [0, 1] \\ (d_1, d_2, d_3, \dots) &\longmapsto 0.d_1d_2d_3\dots \end{aligned}$$

Is a bijection; however, it is actually not the case! To see this, consider that  $0.1000\dots = 0.0999\dots$ . The sequences  $(1, 0, 0, 0, \dots)$  and  $(0, 9, 9, 9, \dots)$  are distinct, but the real number they would map to would be the same. This map is not an injection but a surjection. To show that  $\mathbb{R}$  is uncountable, we require an injection from  $D$  to  $[0, 1]$ . So instead, what we have done with the above proof is map the set of binary sequences  $B$  to  $[0, 1]$ :

$$\begin{aligned} g : B &\longrightarrow [0, 1] \\ (b_1, b_2, b_3, \dots) &\longmapsto \sum_{j=1}^{\infty} b_j 3^{-j} = \sup \left\{ \sum_{j=1}^N b_j 3^{-j} : N \in \mathbb{N} \right\} \end{aligned}$$

Note that in general, an infinite sum is not equal to the supremum of its partial sums (as a counterexample, consider the sum with first term 1, the second term -1, and all the remaining terms 0; evidently the value

of the infinite sum is 0, but the supremum of the partial sums would be 1.) We will define an infinite sum more generally in chapter 3, but in our case here, this definition suffices as in the case where all terms of the sum are non-negative, the supremum of the partial sums is indeed the value of the infinite sum. We leave it as an exercise to show that  $g$  is an injection. Also, one neat remark; the above set is equivalent to the Cantor set, which we will more formally define/discuss in Chapter 2.

We take note a theorem that if  $Y$  is a set, and there exists  $X \subset Y$  such that  $X$  is in bijective correspondence with an uncountable set, then  $Y$  is also uncountable. Thus we can use the above theorem to conclude that  $\mathbb{R}$  is uncountable.

## 2.3 Topology of Metric Spaces

In our investigation of topology, we will try to better understand distances between and neighbourhoods of points. To do so, we first introduce the notion of a metric space.

### Definition 2.15: Metric Spaces

A set  $X$  is a **metric space** (whose elements we call points) with a **metric**  $d : X \times X \mapsto \mathbb{R}$  such that for  $x, y, z \in X$ :

- (a)  $d(x, y) > 0$  if  $x \neq y$ ;  $d(x, x) = 0$ ;
- (b)  $d(x, y) = d(y, x)$  ( $d$  is symmetric);
- (c)  $d(x, z) \leq d(x, y) + d(y, z)$  (triangle inequality).

### Example 2.16

$\mathbb{N}, \mathbb{Q}, \mathbb{R}, \mathbb{R}^n, \mathbb{C}$  are all metric spaces with  $d(x, y) = |x - y|$ . Any subset  $Y \subset X$  of a metric space is also a metric space, with the same metric.

Another example of a metric space is given below; note that this example can be generalized, in that any connected graph can be made into a metric space.



Figure 5: A graph-theoretic example of a metric space. Let  $X = \{1, 2, 3, 4, 5, 6\}$ . Then, for  $x, y \in X$ , let  $d(x, y)$  be the number of edges in the shortest path between  $x$  and  $y$ . Properties (a), (b) of a metric are immediately satisfied, and (c) follows from the property of the shortest path.

It may be of interest to consider other possible metrics. The discrete metric is defined such that  $d(x, y) = 1$  if  $x \neq y$  and  $d(x, x) = 0$ . The L1 Norm is defined such that  $d(f, g) = \int_0^1 |f(x) - g(x)| dx$  where  $f, g$  are functions. In an inner product space, we have that  $d(v, w) = \|v - w\| = \langle v - w, v - w \rangle^{1/2}$ ; if  $v, w$  are functions, then we have that  $d(v, w) = \left( \int_0^1 |v(x) - w(x)|^2 dx \right)^{1/2}$ . In analysis, we are often interested in the notion of things being "close to other things" (e.g. with limits, continuity) so a notion of distance is extremely important to define.

### Definition 2.18: Neighbourhoods

A **neighbourhood** in a metric space  $X$  is a set  $N_r(p) = \{q \in X : d(p, q) < r\}$  with  $r > 0$ .

### Example

- In  $\mathbb{R}$ ,  $N_r(p)$  is the interval  $(p - r, p + r)$  about midpoint  $p$ .
- In  $\mathbb{R}^2$ ,  $N_r(p)$  is the open disk about center  $p$ .
- In  $\mathbb{R}^3$ ,  $N_r(p)$  is the open ball about center  $p$ .
- In  $\mathbb{R}^n$ ,  $N_r(p)$  is the open hyperball about center  $p$ .

As another example, consider that in  $\mathbb{Z}$ ,  $N_1(0) = \{0\}$ ,  $N_{3/2} = \{-1, 0, 1\}$  and  $N_2(0) = \{-1, 0, 1\}$ .

### Definition 2.18: Interior Points

Let  $E \subset X$ . Then,  $p$  is an **interior point** of  $S$  if there is a neighbourhood  $N_r(p)$  such that  $N \subset E$ .

Intuitively, an interior point of  $E$  is a point that is not on the boundary of  $E$ . As an example, in  $\mathbb{R}^n$ , if  $E = \{y : |x - y| \leq 1\}$ , then the interior points of  $E$  (which we can denote as  $E^\circ$ ) are  $E^\circ = \{y : |x - y| < 1\}$ . The idea is that there is always some finite distance to the boundary, so we can always fit a (perhaps small) open ball in. But this doesn't hold at the boundary!



Figure 6: A visualization of an interior point. A set  $E \subset X$  is pictured.  $p_1$  is an interior point as there exists  $N_r(p_1) \subset E$ .  $p_2$  is not an interior point as there does not exist a neighbourhood of  $p_2$  that is entirely contained in  $E$  (it is on the boundary).

**Definition 2.18: Open Sets**

A set  $E \subset X$  is **open** if every point of  $E$  is an interior point of  $E$ .

**Theorem 2.19**

Every neighbourhood is an open set.

**Proof**

Consider a neighbourhood  $E = N_r(p) \subset X$ . Let  $q \in E$ . We will show that  $q$  is an interior point of  $E$ . Choose  $s < r - d(p, q)$ . Then, let  $x \in N_s(q)$ . By the triangle inequality:

$$d(x, p) \leq d(x, q) + d(q, p) < s + d(q, p) < r - d(p, q) + d(p, q) = r.$$

Hence,  $d(x, p) < r$  and it follows that  $x \in N_r(p)$ . Hence,  $N_s(q) \subset N_r(p)$  and  $q$  is an interior point of  $E$ .  $\square$




Figure 7: Visualization of the Sets/Points in Theorem 2.19

**Definition 2.18: Limit Points/Isolated Points**

Let  $E \subset X$  and  $p \in X$ . Then,  $p$  is a **limit point** of  $E$  if every neighbourhood of  $p$  contains  $q \in E$ ,  $q \neq p$ . If  $p \in E$  and  $p$  is not a limit point of  $E$ , then  $p$  is an **isolated point** of  $E$ .

**Example**

Let  $E = \left\{ \frac{1}{n} : n \in \mathbb{N} \right\}$ . Then,  $\frac{1}{2}$  is not a limit point of  $E$ , as for  $r < \frac{1}{4}$   $N_r(\frac{1}{2})$  does not contain any other points of  $E$ . On the other hand, 0 is a limit point of  $E$ . For any neighbourhood  $N_r(0)$  of 0,  $\frac{1}{N} \in N_r(0)$  for  $N > \frac{1}{r}$ . Note that 0 is the only limit point of  $E$ , and is not contained in  $E$  (indeed, there is no requirement that a limit point be contained in the set).

**Theorem 2.20**

If  $p$  is a limit point of  $E$ , then every neighbourhood of  $p$  contains an infinite number of  $q \in E$ .



### Proof

(Sketch) Let  $r_1 = 1$ . Then, there exists  $q_1 \in N_{r_1}(p)$  such that  $q_1 \in E$  and  $q_1 \neq p$  as  $p$  is a limit point of  $E$  by assumption. Let  $r_2 = d(q_1, p)$ . Then, there exists  $q_2 \in N_{r_2}(p)$  such that  $q_2 \in E$  and  $q_2 \neq p$ . We can repeat this process to get a (countably infinite) sequence of distinct points  $q \in N_{r_1}(p)$ , which proves the claim.  $\square$

### Corollary

If  $E \subset X$  is finite, then  $E$  has no limit points.

### Definition 2.18: Dense Sets

$E \subset X$  is dense in  $X$  if every point of  $X$  is a limit point of  $E$ , or a point of  $E$ .

By consequence of Theorem 1.20, we have that  $\mathbb{Q}$  is dense in  $\mathbb{R}$  and  $\mathbb{R} \setminus \mathbb{Q}$  is dense in  $\mathbb{R}$ . As a general method to show that a set is dense in another set, take  $x \in X \setminus E$ , and show  $x$  must be a limit point of  $E$ .

### Definition 2.18: Bounded Sets

$E \subset X$  is bounded if there exists  $M \in \mathbb{R}$  and a point  $q \in X$  such that  $d(p, q) < M$  for all  $p \in E$ .

For example,  $(0, 1)$ ,  $[0, 1]$ ,  $[0, 1] \times [0, 1]$  are bounded, and  $(0, \infty)$ ,  $\mathbb{N}$ ,  $\mathbb{R}$  are unbounded (with the usual metric on  $\mathbb{R}$ ).

### Definition 2.18: Closed Sets

A set  $E \subset X$  is closed if every limit point of  $E$  is in  $E$ .

Note that with the above Corollary, we find that every finite set is (trivially) closed.

### Definition 2.18: Complement

Let  $E \subset X$ . Then, the complement of  $E$ , denoted  $E^c$  is  $E^c = \{x \in X : x \notin E\}$ .



Figure 8: Visualization of a set  $E$  and its complement.

### Theorem 2.23

A set  $E \subset X$  is open if and only if  $E^c$  is closed.

Note that this theorem does not imply that all sets are closed or open; it is possible to have a set that is neither closed or open (such as  $[0, 1) \subset \mathbb{R}$ ) and then its complement (with the former example,  $(-\infty, 0) \cup [1, \infty)$ ) which is also neither closed nor open (indeed, most sets are neither closed nor open). As an

additional note, we have that  $X$  (the entire metric space) and  $\emptyset$  are both open and closed (which we may affectionately label as “clopen”).

#### Proof

$\Rightarrow$  Assume  $E$  is open. If  $E^c$  has no limit points, it is trivially closed, so suppose that there exists a limit point  $x$  of  $E^c$ . Suppose for the sake of contradiction that  $x \notin E^c$ . Then,  $x \in E$ . As  $E$  is open,  $x$  is an interior point of  $E$ , so there exists a neighbourhood  $N_r(x) \subset E$ . In particular,  $N_r(x) \cap E^c = \emptyset$ , contradicting the fact that  $x$  is a limit point of  $E^c$ . Hence,  $x \in E^c$  and  $E^c$  is closed.

$\Leftarrow$  Assume  $E^c$  is closed. Let  $x \in E$ . In particular,  $x \notin E^c$ , so  $x$  is not a limit point of  $E^c$ . So, there exists a neighbourhood  $N_r(x)$  which contains no point of  $E^c$ , i.e.  $N_r(x) \cap E^c = \emptyset$ . It follows that  $N_r(x) \subset E$ , and hence  $x$  is an interior point of  $E$ . This argument applies to all points of  $E$ , hence  $E$  is open.  $\square$

#### Corollary

A set  $F \subset X$  is closed if and only if  $F^c$  is open.

Let  $F = E^c$  in Theorem 2.23 to realize the above Corollary.

#### Theorem 2.24

- (a) For any collection  $\{E_\alpha\}$  of open sets,  $\bigcup_\alpha E_\alpha$  is open.
- (b) For any collection  $\{F_\alpha\}$  of closed sets,  $\bigcap_\alpha F_\alpha$  is closed.
- (c) For any finite collection  $E_1, \dots, E_n$  of open sets,  $\bigcap_{i=1}^n E_i$  is open.
- (d) For any finite collection  $F_1, \dots, F_n$  of closed sets,  $\bigcup_{i=1}^n F_i$  is closed.

A point of notation;  $\{E_\alpha\}$  can be finite, countable, or uncountable; the indices  $\alpha$  are taken from an index set  $A$  which can be chosen to be of any cardinality.

#### Proof

- (a) Suppose all sets in  $\{E_\alpha\}$  are closed. Let  $x \in \bigcup_\alpha E_\alpha$ . Then, there exists  $\alpha_0$  such that  $x \in E_{\alpha_0}$ . Since  $E_{\alpha_0}$  is open, there exists a neighbourhood  $N_r(x)$  of  $x$  such that  $N_r(x) \subset E_{\alpha_0} \subset \bigcup_\alpha E_\alpha$ . Hence,  $\bigcup_\alpha E_\alpha$  is open.
- (b) Suppose all sets in  $\{F_\alpha\}$  are open. To show that  $\bigcap_\alpha F_\alpha$  is closed, we show that  $(\bigcap_\alpha F_\alpha)^c$  is open (by Theorem 2.23). We have that  $(\bigcap_\alpha F_\alpha)^c = \bigcup_\alpha F_\alpha^c$ . As all  $F_\alpha^c$  are open, by part (a) we have that  $\bigcup_\alpha F_\alpha^c$  is also open. Hence  $\bigcap_\alpha F_\alpha$  is closed.
- (c) Suppose  $E_1, \dots, E_n$  are open. Let  $x \in \bigcap_{i=1}^n E_i$ , and then we have that  $x \in E_i$  for all  $i \in \{1, \dots, n\}$ . Hence, there exists  $r_i$  such that  $N_{r_i}(x) \subset E_i$  as each of the  $E_i$ s are open. Let  $r = \min\{r_1, \dots, r_n\}$  and then we have that  $N_r(x) \subset N_{r_i}(x) \subset E_i$  for all  $E_i$ . Therefore,  $N_r(x) \subset \bigcap_{i=1}^n E_i$  and  $\bigcap_{i=1}^n E_i$  is open.
- (d) Suppose  $F_1, \dots, F_n$  are closed. By Theorem 2.23 we have that  $\bigcup_{i=1}^n F_i$  is closed if and only if  $(\bigcup_{i=1}^n F_i)^c = \bigcap_{i=1}^n F_i^c$  is open. Since all  $F_i^c$ s are open, by part (c)  $\bigcap_{i=1}^n F_i^c$  is open, and hence  $\bigcup_{i=1}^n F_i$  is closed.  $\square$

### Example 2.25

We consider some examples to see why the finiteness of the collections in parts (c)/(d) of the theorem are essential. Suppose  $E_n = \left(-\frac{1}{n}, \frac{1}{n}\right) \subset \mathbb{R}$ . These sets form a countably infinite collection of subsets of  $\mathbb{R}$ . We then consider that  $\bigcap_{n=1}^{\infty} E_n = \{0\}$ , which is not open; showing that openness is not preserved under infinite intersections. Next, consider  $F_n = [0, 1 - \frac{1}{n}] \subset \mathbb{R}$ , which form a countably infinite collection of closed sets in  $\mathbb{R}$ . We then have that  $\bigcup_{n=1}^{\infty} F_n = [0, 1)$  which is not closed as 0 is not an interior point of the set. Hence, closedness is not preserved under infinite unions.

## 2.4 Closure and Relative Topology

As a review of some definitions of the previous section, we call a set open if every point of the set is an interior point, and closed if it contains all of its limit points. A complement of an open set is closed and vice versa. We also note that openness is preserved under infinite unions and finite intersections, and closedness is preserved under infinite intersections and finite unions.

### Definition 2.26: Closure

Let  $X$  be a metric space. Let  $E \subset X$ , and denote  $E'$  as the set of all limit points of  $E$ . Then, the set  $\bar{E} = E \cup E'$  is the **closure** of  $E$ .

### Theorem 2.27

- (a)  $\bar{E}$  is closed.
- (b)  $E = \bar{E}$  if and only if  $E$  is closed.
- (c)  $\bar{E} \subset F$  for every closed set  $F \subset X$  such that  $E \subset F$ .

### Proof

- (a) Let  $p \in \bar{E}^c = E^c \cap (E')^c$ . Then,  $p \notin E$ , and furthermore  $p \notin E'$  so  $p$  is not a limit point of  $E$ . Hence, there exists a neighbourhood  $N_r(p)$  such that  $N_r(p) \cap E = \emptyset$ . Moreover, no point of  $N_r(p)$  is in  $E'$  (if there existed  $q \in E'$  such that  $q \in N_r(p)$ , then there would exist some  $N_{r'}(q)$  (which contains points of  $E$ ) such that  $N_{r'}(q) \subset N_r(p)$  which contradicts the fact that  $N_r(p)$  contains no points of  $E$ ). Hence,  $N_r(p) \subset E^c \cap (E')^c = \bar{E}^c$ . Hence,  $\bar{E}^c$  is open and  $\bar{E}$  is closed.
- (b)  $\implies$  If  $E = \bar{E}$ , then  $E$  is closed by (a).  
 $\impliedby$  If  $E$  is closed, then  $E$  contains all of its limit points, so  $\bar{E} = E \cup E' \subset E$ . By definition  $\bar{E} \supseteq E$ , so  $E = \bar{E}$ .
- (c) Let  $E \subset F$  with  $F$  closed. Then,  $F' \subset F$ . Also,  $E' \subset F'$ . So, by (b) we have  $F = \bar{F} = F \cup F' \supseteq E \cup E' = \bar{E}$ . □

We will soon define the notion of being relatively open, but before doing so, we consider a motivating example.

In  $\mathbb{R}$ , we know  $(a, b)$  to be in an open set. However, embedded in  $\mathbb{R}^2$ ,  $(a, b)$  has no interior points; any neighbourhood about any  $x \in (a, b)$  extends into the plane and will inevitably contain points  $y \in \mathbb{R}^2$ ,  $y \notin (a, b)$ . Hence,  $(a, b)$  is open as a subset of  $\mathbb{R}$ , but not as a subset of  $\mathbb{R}^2$ .



Figure 9: The interval  $(a, b)$  embedded in  $\mathbb{R}^2$ .

### Definition 2.29: Relative Openness

Let  $E \subset Y \subset X$ . Then,  $E$  is **relatively open** with respect to  $Y$  if it is an open set in the metric space  $Y$ .

However, we note that the above definition is not very useful. Is there a better notion of relative openness? Going back to our prior example, consider that  $(a, b)$  can be viewed as the intersection of an open disk with  $\mathbb{R}$ . This gives an equivalent definition!



Figure 10: The interval  $(a, b)$  can be viewed as the intersection of  $\mathbb{R}$  with the open disk  $N = \left\{ (x, y) \in \mathbb{R}^2 : \sqrt{\left(x - \frac{b+a}{2}\right)^2 + y^2} < \frac{b-a}{2} \right\}$ , giving rise to a more useful notion of relative openness.

### Theorem 2.30

Let  $E \subset Y \subset X$ . Then,  $E$  is open relative to  $Y$  if and only if there exists an open set  $G \subset X$  such that  $E = G \cap Y$ .

#### Proof

$\Rightarrow$  Let  $p \in E$ . Then, there exists a neighbourhood  $N_r^Y(p) \subset E$  (Note that here,  $N_r^Y(p) = \{y \in Y : d(p, y) < r\}$ ). By Theorem 2.24, we have that  $G = \bigcup_{p \in E} N_{r_p}^X(p)$  is open. Then, we have that  $G \cap Y = \bigcup_{p \in E} N_{r_p}^X(p) \cap Y = \bigcup_{p \in E} N_{r_p}^Y(p) = E$ .

$\Leftarrow$  Suppose  $G \subset X$  is open and  $E = G \cap Y$ . Let  $p \in E$ . Then, there exists a neighbourhood  $N_r^X(p) \subset G$  as  $G$  is open, so  $N_r^Y(p) = N_r^X(p) \cap Y \subset G \cap Y = E$ . Hence,  $p$  is an interior point of  $E$  in the metric space of  $Y$ , and hence  $E$  is relatively open in  $Y$ .  $\square$

## 2.5 Compactness

### Definition 2.31: Open Covers

An **open cover** of a subset  $E \subset X$  is a collection  $\{G_\alpha\}$  of open sets of  $X$  such that  $E \subset \bigcup_\alpha G_\alpha$ .

Note that the open cover can be either a finite or infinite (countable or uncountable) collection.




Figure 11: Visualization of a set  $E$  and a collection of open disks that form a (finite) open cover of  $E$ .

### Definition 2.32: Compactness

A subset  $K \subset X$  is **compact** if every open cover of  $K$  has a finite subcover. Explicitly, if  $\{G_\alpha\}$  is an open cover of  $K$ , then there exist indices  $\alpha_1, \dots, \alpha_n$  such that  $K \subset G_{\alpha_1} \cup \dots \cup G_{\alpha_n}$ .

While the above definition might seem strange/esoteric, it turns out to be very useful and important to analysis. We saw in the last lecture that open sets in a sense do not behave “nicely”; a given set could be open with respect to a subspace  $Y$  but not to  $X$  (and we had to introduce the notion of relative openness to take care of this fact). Compactness behaves more nicely, as we will see with the following theorem.

### Theorem 2.33

Suppose  $K \subset Y \subset X$ . Then,  $K$  is compact with respect to  $X$  if and only if  $K$  is compact with respect to  $Y$ .

While openness depends on the choice of subspace, this theorem states that compactness is independent of what subspace we choose to look at. It is in a sense an intrinsic property of the set.

### Proof

$\Rightarrow$  Suppose  $K$  is compact with respect to  $X$ . Let  $\{V_\alpha\}$  be an open (in  $Y$ ) cover of  $K$ . Then, by Theorem 2.30 we have that there exists  $\{G_\alpha\}$  of open sets in  $X$  such that  $V_\alpha = G_\alpha \cap Y$ . Then,  $\{G_\alpha\}$  is an open (in  $X$ ) cover of  $K$ , and hence there exists a finite subcover  $K \subset \bigcup_{i=1}^n G_{\alpha_i}$ . Then, we have that  $K \subset \bigcup_{i=1}^n (G_{\alpha_i} \cap Y) = \bigcup_{i=1}^n V_{\alpha_i}$  which is a finite (sub)cover of  $K$  in  $Y$ . Hence,  $K$  is compact with respect to  $Y$ .

$\Leftarrow$  Suppose  $K$  is compact with respect to  $Y$ . Let  $\{G_\alpha\}$  be an open (in  $X$ ) cover of  $K$ . Let  $V_\alpha = G_\alpha \cap Y$ . These are relatively open in  $Y$  by Theorem 2.30, and  $\{V_\alpha\}$  still cover  $K$ . Hence, as  $K$  is compact in  $Y$ , there exists a finite subcover  $K \subset \bigcup_{i=1}^n V_{\alpha_i} \subset \bigcup_{i=1}^n G_{\alpha_i}$ . Therefore, we have found a finite subcover in  $X$ , so  $K$  is compact in  $X$ .  $\square$

### Theorem 2.34

Let  $K \subset X$  be compact. Then,  $K$  is closed.

A priori, it might seem like the notions of compactness and closedness are quite different, but this theorem shows that they are indeed quite closely related.

#### Proof

We show that  $K^c$  is open. Let  $p \in K^c$ . For any  $q \in K$ . Let  $r_q = \frac{1}{2}d(p, q)$ , and define  $W_q = N_{r_q}(q)$ . So, taking the collection  $\{W_q\}$  we have that  $\bigcup_{q \in K} W_q$  is an open cover (as it clearly contains every  $q \in K$ ).  $K$  is compact, so there exists a finite subcover  $\bigcup_{i=1}^n W_{q_i}$ . Then, let  $r = \min\{r_{q_1}, \dots, r_{q_n}\}$ . Then,  $N_r(p) \subset K^c$ , hence  $p$  is an interior point of  $K^c$ . Therefore  $K^c$  is open, and  $K$  is closed by Theorem 2.23.  $\square$

### Theorem 2.35

If  $F \subset X$  is closed,  $K \subset X$  is compact, and  $F \subset K$ , then  $F$  is compact.

#### Proof

Let  $\{V_\alpha\}$  be an open cover of  $F$ . Then,  $\{V_\alpha\} \cup F^c$  is an open cover of  $K$  ( $F^c$  is open as  $F$  is closed). Hence, as  $K$  is compact, there exists a finite subcover. Dropping  $F^c$  (if it is part of the finite subcover of  $K$ ), we obtain a finite subcover of  $F$ . Hence,  $F$  is compact.  $\square$

### Corollary

If  $F \subset X$  is closed and  $K \subset X$  is compact, then  $F \cap K$  is compact. In other words, compactness is preserved under intersection (with closed sets).

#### Proof

If  $K$  is compact, it is closed by Theorem 2.34.  $F \cap K$  is closed by Theorem 2.24, and since  $F \cap K \subset K$ , by Theorem 2.35  $F \cap K$  is compact.  $\square$

In a previous course, we may have introduced the definition that compact sets are closed and bounded. Here, we started with a significantly different definition, and work up to prove the Heine-Borel Theorem (coming soon!) which tells us that compactness is equivalent to being closed and bounded in  $\mathbb{R}^k$ . Before we move onto such further properties and theorems of compact sets, we may find it useful to consider some examples of sets that are and are not compact.

The singleton set  $\{x\}$  is compact in any metric space (out of any open cover, simply pick a single open set that contains  $x$ ; this yields the desired subcover). Moreover, any finite set is compact in any topology (consider that for any open cover of a set of  $n$  elements, we can just choose at most  $n$  open sets corresponding to each element to form our finite subcover).

Next, consider any infinite set  $Y$  in a metric space  $X$  with the discrete metric:

$$d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$$

Then is  $Y \subset X$  compact? The answer is no; consider the open cover of singleton sets,  $U = \{\{x\} : x \in Y\}$ . First, note that this is indeed an open cover, as  $\{x\}$  is open in  $X$ ; consider that  $N_r(x) \subset \{x\}$  if  $r < 1$ , and hence  $x$  is an interior point of  $\{x\}$ , making it open. However, there exists no finite subcover of this open cover as we need to pick an infinite number of singletons to cover each point in  $Y$ .

$[1, \infty) \subset \mathbb{R}$  is not compact as we could consider the open cover  $\{N_1(n) : n \in \mathbb{N}\}$ , which has no finite subcover. For the sake of contradiction, suppose such a finite subcover existed. Then, there exists some maximum  $m$  for which there would be a neighbourhood around in this subcover. Then, all real numbers greater than  $m + 1$  would not be in this subcover, which is a contradiction. In a similar way, we can show that  $\mathbb{R} \subset \mathbb{R}, \mathbb{R}^2 \subset \mathbb{R}^2$  are not compact.

Another interesting example to consider is  $(0, 1)$ . To see that this interval is not compact, consider the open cover  $\{(\frac{1}{n}, 1) : n \in \mathbb{N}\}$ . This open cover has no finite subcover. Suppose for the sake of contradiction that a finite subcover existed. Then, there would be some minimum  $\frac{1}{N}$  such that  $(\frac{1}{N}, 1)$  would be in the subcover. But then no  $0 < x < \frac{1}{N}$  would be contained in the subcover, which is a contradiction.

From the Heine-Borel theorem, it is clear here that  $[1, \infty), \mathbb{R}, \mathbb{R}^2$  fail due to the sets being unbounded, and the  $(0, 1)$  fails due to the set not being closed. We will now move onto a sequence of Theorems that will eventually lead to Heine-Borel (the climax of this chapter)!

### Theorem 2.36

Let  $\{K_\alpha\}$  is a collection of compact sets such that  $\bigcap_{i=1}^n K_{\alpha_i} \neq \emptyset$  for any subcollection/any choice of indices  $\alpha_1, \dots, \alpha_n$ . Then,  $\bigcap_\alpha K_\alpha \neq \emptyset$ .

### Proof

Suppose for the sake of contradiction that the assumptions hold but  $\bigcap_\alpha K_\alpha \neq \emptyset$ . Pick  $\alpha_0$ . Then,  $K_{\alpha_0} \cap (\bigcap_{\alpha \neq \alpha_0} K_\alpha) = \emptyset$ . Hence,  $K_{\alpha_0} \subset (\bigcap_{\alpha \neq \alpha_0} K_\alpha)^c = \bigcup_{\alpha \neq \alpha_0} K_\alpha^c$ . Hence,  $\{K_\alpha^c\}$  is an open cover of  $K_{\alpha_0}$ .  $K_{\alpha_0}$  is compact, so there exists a finite subcover  $\bigcup_{i=1}^n K_{\alpha_i}^c = (\bigcap_{i=1}^n K_{\alpha_i})^c$ . But then we have that  $K_{\alpha_0} \cap (\bigcap_{i=1}^n K_{\alpha_i}) = \emptyset$ , which is a contradiction as a finite intersection should be nonempty by assumption.  $\square$

### Corollary

Let  $\{K_1, K_2, \dots\}$  be a collection of nonempty and compact sets such that  $K_{i+1} \subset K_i$ . Then,  $\bigcap_{i=1}^\infty K_i \neq \emptyset$ .

### Proof

If  $n_1 < \dots < n_m$ , then  $\bigcap_{i=1}^m K_{n_i} = K_{n_m} \neq \emptyset$ . Then,  $\bigcap_{i=1}^\infty K_{n_i} \neq \emptyset$  by the above theorem.  $\square$

### Theorem 2.37

Let  $K$  be compact and  $E \subset K$  be an infinite set. Then,  $E$  has a limit point in  $K$ .

### Proof

Suppose for the sake of contradiction that no point of  $K$  is a limit point of  $E$ . Then, for all  $q \in K$ , then there exists  $V_q = N_{r_q}(q)$  such that:

$$V_q \cap E = \begin{cases} \{q\} & \text{if } q \in E \\ \emptyset & \text{if } q \notin E \end{cases}$$

Trivially,  $\{V_q\}$  is an open cover of  $K$ . So, there exists a finite subcover  $\{V_{q_i}\}_{i=1}^n$  of  $K$ . So:

$$E \subset K \cap E \subset \left( \bigcup_{i=1}^n V_{q_i} \right) \cap E = \bigcup_{i=1}^n (V_{q_i} \cap E)$$

But the RHS is a finite set, contradicting the assumption that  $E$  is infinite.  $\square$

## 2.6 Compactness in $\mathbb{R}^k$ and the Cantor Set

### Theorem 2.38

Let  $I_n = [a_n, b_n] \subset \mathbb{R}$  such that  $I_{n+1} \subset I_n$  for all  $n$ . Then,  $\bigcap_{i=1}^{\infty} I_i \neq \emptyset$ .

This theorem is very reminiscent of Theorem 2.37. However, we cannot apply it directly as we do not know if  $[a, b]$  is compact (though we will show that this is indeed the case by the end of the lecture, we want to avoid circular reasoning)!



Figure 12: Visualization of the first few sets  $I_n$  in Theorem 2.38. Note that this is just an example, and the sets do not need to be “symmetrically shrinking” around a point as pictured.

### Proof

We have that  $a_n \leq a_{n+m} \leq a_{n+m} \leq b_n$  for all  $n, m \in \mathbb{N}$ . Let  $E = \{a_1, a_2, \dots\}$ .  $E$  is nonempty and bounded by any  $b_n$ , so by the least upper bound property of  $\mathbb{R}$ , there exists  $x = \sup E \in \mathbb{R}$ . We then have that  $a_n \leq x \leq b_n$  for all  $n$  as  $x$  is the least upper bound. Hence,  $x \in I_n$  for all  $n$ , and hence  $x \in \bigcap_{i=1}^{\infty} I_i$ . We conclude that  $\bigcap_{i=1}^{\infty} I_i \neq \emptyset$ .  $\square$

### Definition: k-cells

A **k-cell** is  $I \subset \mathbb{R}^k$  such that  $I = \{(x_1, x_2, \dots, x_k) \in \mathbb{R}^k : a_j \leq x_j \leq b_j\}$  for some  $\{a_1, \dots, a_k, b_1, \dots, b_k\}$ .

A  $k$ -cell can be viewed as the generalization of a rectangle for  $\mathbb{R}^k$ .



**Theorem 2.39**

Theorem 2.38 holds for general  $k$ -cells.

The above theorem follows naturally by applying Theorem 2.38 to each coordinate.

**Theorem 2.40**

Let  $I \subset \mathbb{R}^k$  be a  $k$ -cell. Then,  $I$  is compact.

**Proof**

Let  $\{G_\alpha\}$  be an open cover of  $I = I_0$ . Suppose for the sake of contradiction that  $\{G_\alpha\}$  has no finite subcover. Let  $c_j = \frac{1}{2}(a_j + b_j)$ . Splitting up  $I_0$  along each coordinate (i.e.  $[a_j, c_j], [c_j, b_j]$ ) we obtain  $2^k$   $k$ -cells  $Q_i$ , with  $I_0 = \bigcup_{i=1}^{2^k} Q_i$ . At least one of these  $Q_n$ s has no finite subcover by assumption. Call this  $I_1$ . Then, repeat the same division process using  $I_1$  (and so on). This yields a sequence of  $k$ -cells  $I_0, I_1, I_2, \dots$ . This sequence has the following properties:

- (a)  $I_0 \supset I_1 \supset I_2 \supset \dots$
- (b) None of  $I_n$ s have a finite subcover from  $\{G_\alpha\}$  by construction.
- (c) Let  $\delta = \sqrt{\sum_{j=1}^k (b_j - a_j)^2}$  be the diameter of  $I$ . Then, for  $x, y \in I_n$ ,  $|x - y| < 2^{-n}\delta$ .

By (a) and Theorem 2.39, we have that there exists  $\mathbf{x}^*$  such that  $\mathbf{x}^* \in \bigcap_{n=1}^{\infty} I_n \subset I$ . There exists  $\alpha_0$  such that  $\mathbf{x} \in G_{\alpha_0}$ , and as  $G_{\alpha_0}$  is open, there exists some  $r > 0$  such that  $N_r(\mathbf{x}^*) \subset G_{\alpha_0}$ . By (c), we have that  $I_n \subset N_{2^{-n+1}\delta}(\mathbf{x}^*)$ . For sufficiently large  $n$ ,  $I_n \subset N_{2^{-n+1}\delta}(\mathbf{x}^*) \subset N_r(\mathbf{x}^*) \subset G_{\alpha_0}$ , but this contradicts (b). Hence, there must exist a finite subcover of  $\{G_\alpha\}$  for  $I$ .  $\square$



Figure 13: Visualization of the division process (first three iterations shown) in the proof of Theorem 2.40, for  $I \subset \mathbb{R}^2$ .

### Theorem 2.41: Heine-Borel

If a set in  $\mathbb{R}^k$  has the following three properties, then it has the other two.

- (a)  $E$  is bounded and closed.
- (b)  $E$  is compact.
- (c) Every infinite subset of  $E$  has at least one limit point in  $E$

In particular, the equivalence of (a) and (b) is what is commonly referred to as the Heine-Borel theorem.

### Proof

(a)  $\implies$  (b) If  $E$  is bounded, then there exists a  $k$ -cell  $I$  such that  $E \subset I$ .  $I$  is compact by Theorem 2.40, and since  $E$  is closed, by Theorem 2.35 we have that  $E$  is compact.

(b)  $\implies$  (c) See Theorem 2.37.

(c)  $\implies$  (a) Suppose first for the sake of contradiction that  $E$  is not bounded. Then,  $E$  has an infinite subset  $S = \{x_1, x_2, \dots\}$  with  $|x_n| > n$  for all  $n$ . Hence,  $S$  has no limit points in  $\mathbb{R}^k$ , and therefore no limit points in  $E$ . But this contradicts (c).

Suppose next that  $E$  is not closed. Then, there exists  $x_0 \in \mathbb{R}^k$  which is a limit point of  $E$  but is not in  $E$ . Form  $S = \{x_1, x_2, \dots\} \subset E$  with  $|x_n - x_0| < \frac{1}{n}$  for all  $n$ . We now show that  $S$  has no limit point in  $\mathbb{R}^k$  except for  $x_0$ . To see this, let  $y \in \mathbb{R}^k, y \neq x_0$ . Then:

$$|x_n - y| \geq |x_0 - y| - |x_n - x_0| \geq |x_0 - y| - \frac{1}{n} \geq |x_0 - y| - \frac{n}{2}|x_0 - y| = \frac{1}{2}|x_0 - y|$$

Where the last inequality holds for sufficiently large  $n$ . As every neighbourhood of  $y$  must contain an infinite number of points in  $S$  if  $y$  is to be a limit point of  $S$  (Theorem 2.20) we find that  $y$  is not a limit point of  $S$ . Hence,  $S$  has no limit point in  $E$ , again contradicting (c). We conclude that  $E$  must be bounded and closed.  $\square$

We have already shown that compactness implies closed in any metric space (see Theorem 2.34). It also turns out that compactness implies boundedness in any metric space, as well!

*Proof.* Let  $E \subset X$  be compact. Suppose for the sake of contradiction that  $E$  was unbounded. Pick any  $x_0 \in E$ , then  $E$  is unbounded, so  $(N_n(x_0))^c \cap E \neq \emptyset$  for all  $n \in \mathbb{N}$ . But,  $E \subset \bigcup_{n \in \mathbb{N}} N_n(x_0)$ , and hence  $\{N_n(x_0)\}$  forms an (infinite) subcover of  $E$ . By the compactness of  $E$ , there exists a finite subcover; but then, there is a maximal radius neighbourhood of  $x_0$  that is contained in this subcover. However, this neighbourhood would not contain all  $x \in E$  as  $E$  is unbounded, which is a contradiction.  $\square$

From this, we have shown that compact implies closed and bounded in any metric space (i.e. the forwards direction of Heine-Borel holds in general). The converse is not always true, however. For one example, consider any infinite set  $E$  in a metric space equipped with the discrete metric.  $E$  is closed, since  $E$  has no limit points (the neighbourhood of radius  $r \leq 1$  around any point contains no other point of  $E$ ).  $E$  is bounded as a neighbourhood of radius  $r > 1$  around any point contains all points of  $E$ . However, as we have discussed previously,  $E$  is not compact. Another example would be  $\mathbb{R}^\infty$  (i.e. the set of all infinite sequences of real numbers), but the argument for this is left as homework (HW5).

**Theorem 2.42: Weierstrauss**

Suppose  $E \subset \mathbb{R}^k$  is bounded and infinite. Then,  $E$  has a limit point in  $\mathbb{R}^k$ .

**Proof**

Since  $E$  is bounded,  $E \subset I \subset \mathbb{R}^k$  for some  $k$ -cell  $I$ . By Theorem 2.40  $E$  is compact, so by Theorem 2.37  $E$  has a limit point in  $I$  (and hence in  $\mathbb{R}^k$ ).  $\square$

Having proven the landmark (Heine-Borel) theorem of this section, we close this chapter with discussion of perfect sets and the Cantor set.

**Definition 2.18: Perfect Sets**

A set  $P \subset X$  is **perfect** if  $P$  is closed and every point of  $P$  is a limit point of  $P$ .

**Theorem 2.43**

Let  $P \subset \mathbb{R}^k$  be nonempty and perfect. Then,  $P$  is uncountable.

**Proof**

Since  $P$  has limit points, it is not finite by Theorem 2.20. Assume for the sake of contradiction that  $P$  is countable. Then,  $P = \{x_1, x_2, \dots\}$ . Let  $V_1 = N_r(x_1)$  be any neighbourhood of  $x_1$ . Suppose we construct  $B_n$  such that  $V_n \cap P \neq \emptyset$  (note that we don't assume that  $x_n \in V_n$ , just that some point  $p \in P$  is in  $V_n$ ). Since  $P$  is perfect, we may construct  $V_{n+1}$  such that:

- (i)  $\overline{V}_{n+1} \subset V_n$
- (ii)  $x_n \notin \overline{V}_{n+1}$
- (iii)  $V_{n+1} \cap P \neq \emptyset$

Note that  $\overline{V}_n$  is closed and bounded, and hence is compact by Theorem 2.41. Hence, by the corollary to Theorem 2.35 we have that  $K_n = \overline{V}_n \cap P$  is also compact. By (ii), we have that  $x_n \notin \bigcap_{n=1}^{\infty} K_n$ , and this holds for all  $n \in \mathbb{N}$ . Hence,  $P \cap \bigcap_{n=1}^{\infty} K_n = \emptyset$ , but  $K_n \subset P$ , so it follows that  $\bigcap_{n=1}^{\infty} K_n = \emptyset$ . But  $K_n$  is non-empty, compact, and  $K_{n+1} \subset K_n$ , so this contradicts the corollary to Theorem 2.36. We conclude that  $P$  is uncountable.  $\square$

**Corollary**

Every interval  $[a, b]$  is uncountable. It follows that  $\mathbb{R}$  is uncountable.

**Definition 2.44: The Cantor Set**

Let  $E_0 = [0, 1]$ . Then, remove the middle third (that is,  $(\frac{1}{3}, \frac{2}{3})$ ) to obtain  $E_1 = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$ . Then remove the middle thirds of each of these parts to get  $E_2 = [0, \frac{1}{9}] \cup [\frac{2}{9}, \frac{1}{3}] \cup [\frac{2}{3}, \frac{7}{9}] \cup [\frac{8}{9}, 1]$ . This yields a sequence of sets  $E_0 \supset E_1 \supset E_2 \supset E_3 \supset \dots$ . Then, we may define the **Cantor set** as  $P = \bigcap_{n=1}^{\infty} E_n$ .

Although we will rigorously define a measure in this course, each  $E_n$  in the above construction is the union of  $2^n$  intervals of measure  $\frac{1}{3^n}$ , for a total measure of  $\frac{2^n}{3^n}$  for each  $E_n$ . Taking the limit of  $n \rightarrow \infty$ , we

can therefore see that the Cantor set has measure zero. However, the Cantor set is uncountable (as we will see shortly), making it an example of an uncountable set of measure zero.



Figure 14: Visualization of the first few sets  $E_0, E_1, E_2$  used in the construction of the Cantor set.

### Theorem

The Cantor set contains no interval  $(a, b)$ .

### Proof

(Sketch) Any interval  $(a, b)$  contains some interval  $\left(\frac{3k+1}{3^m}, \frac{3k+2}{3^m}\right)$  which are all removed in the construction of the set.  $\square$

If the Cantor set contains no intervals, then how is it uncountable? It might help to consider that by construction, the endpoints of each  $E_n$  belong to  $P$ ; in the limit, this yields an uncountable number of points contained in  $P$ .

### Theorem

The Cantor set is uncountable. Moreover, it is perfect.

### Proof

Let  $x \in P$  and  $S$  be an interval such that  $x \in S$ . Let  $I_n$  be the interval of  $E_n$  containing  $x$ . The length of  $I_n$  is given by  $\frac{1}{3^n}$ . For sufficiently large  $n$ ,  $I_n \subset S$ . Let  $x_n$  be an endpoint of  $I_n$  such that  $x_n \neq x$ . By construction,  $x_n \in P$ , and hence  $x$  is a limit point of  $P$  by construction. Additionally,  $P$  is closed by Theorem 2.24 as it is an infinite intersection of closed sets. Hence  $P$  is perfect, and by Theorem 2.43 it is uncountable.  $\square$

## 2.7 Connected Sets

Though not covered in lecture, we here briefly discuss connectedness as it comes up later in Chapter 4.

### Definition 2.45: Connected Sets

Two subsets  $A, B$  of a metric space  $X$  are separated if  $A \cap \bar{B} = \bar{A} \cap B = \emptyset$ . A set  $E \subset X$  is connected if  $E$  is not a union of two nonempty separated sets.

Separated sets are disjoint, but disjoint sets are not necessarily separated; for example,  $A = [0, 1]$  and  $B = (1, 2)$  are not separated as  $\bar{B} = [1, 2]$  and hence  $A \cap \bar{B} = \{1\}$ . However,  $A = (0, 1)$  and  $B = (1, 2)$  are separated. The following theorem characterizes connected subsets of  $\mathbb{R}$ .

**Theorem 2.47**

A subset  $E \subset \mathbb{R}$  is connected if and only if it has the property that if  $x, y \in E$  and  $x < z < y$ , then  $z \in E$ .

**Proof**

Not covered in lecture, see Rudin.

### 3 Numerical Sequences and Series

#### 3.1 Sequences

We begin by formally defining a sequence.

##### Definition: Sequences

Let  $X$  be a metric space. A **sequence** is a function  $f : \mathbb{N} \mapsto X$ . We can denote a term in the sequence as  $f(n) = x_n$ , or the entire sequence as  $\{x_n\}_{n=1}^{\infty}$ ,  $\{x_n\}$ ,  $(x_n)$ , or  $\{x_1, x_2, x_3 \dots\}$ .

We now discuss the notion of convergence of a sequence. Intuitively, we can equate convergence with the notion of points getting closer together.

##### Definition 3.1: Convergence of Sequences

A sequence  $\{p_n\}_{n=1}^{\infty}$  **converges** to  $p \in X$  if for all  $\epsilon > 0$ , there exists  $N \in \mathbb{N}$  such that  $n \geq N$  implies  $d(p_n, p) < \epsilon$ . In this case, we say that  $\{p_n\}$  converges to  $p$ , or that  $p$  is the limit of  $\{p_n\}$ , and denote this as  $p_n \rightarrow p$  or  $\lim_{n \rightarrow \infty} p_n = p$ . If  $\{p_n\}$  does not converge, we say it **diverges**.

To phrase this definition in another way, we fix some  $\epsilon > 0$ , and then we have that all points in the sequence past some  $N \in \mathbb{N}$  are contained in the neighbourhood  $N_{\epsilon}(p)$ . In practice, it can be difficult to apply this definition of convergence if we don't know what the limiting  $p$  is, as the definition implicitly uses the value of the limit. We will later discuss another definition of convergence (in  $\mathbb{R}^k$ ) that does not use the value of the limit.




Figure 15: Visualization of a sequence  $\{p_n\} \subset \mathbb{R}^2$  converging to a point  $p$ . For the  $\epsilon > 0$  shown in the picture, we have that all points of the sequence past  $N = 5$  lie in the open disk of radius  $\epsilon$  around  $p$ .

As a remark, consider that convergence can depend on our choice of metric space; for example,  $\{\frac{1}{n}\}$  as a sequence in  $\mathbb{R}$  converges to 0, but the same sequence in the strictly positive reals ( $\mathbb{R}^+ = \{x \in \mathbb{R} : x > 0\}$ ) does not converge.

Another interesting example (that again shows us the importance of the choice of metric space). Is  $\mathbb{R}$  equipped with the discrete metric. A question we can ask is “given some points  $p \in \mathbb{R}$ , what sequences converge to  $p$ ?” The answer turns out to be eventually constant sequences only; that is, sequences for which  $p_n = p$  for  $n \geq N$  for some  $N$ .

*Proof.* If  $p_n \rightarrow p$ , then setting  $\epsilon = \frac{1}{2}$ , we have that there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$ ,  $d(p_n, p) < \epsilon = \frac{1}{2}$ . Under the discrete metric, this is only possible if  $p_n = p$ .  $\square$

This of course is a strikingly different picture for  $\mathbb{R}$  with the standard metric of  $d(x, y) = |x - y|$ . For example, the sequence  $p_n = \frac{1}{n}$  has no term equal to zero, but converges to  $p = 0$ . The takeaway message here can be that in the Euclidean metric, points can “get closer” but in the discrete metric, they cannot.

### Theorem 3.3

Suppose  $\{s_n\}, \{t_n\}$  are complex sequences that converge, with  $\lim_{n \rightarrow \infty} s_n \rightarrow s$  and  $\lim_{n \rightarrow \infty} t_n \rightarrow t$ . Then:

- (a)  $\lim_{n \rightarrow \infty} (s_n + t_n) = s + t$ .
- (b)  $\lim_{n \rightarrow \infty} cs_n = cs$  and  $\lim_{n \rightarrow \infty} (c + s_n) = c + s$  for all  $c \in \mathbb{C}$ .
- (c)  $\lim_{n \rightarrow \infty} s_n t_n = st$
- (d)  $\lim_{n \rightarrow \infty} \frac{1}{s_n} = \frac{1}{s}$  provided  $s \neq 0$  and  $s_n \neq 0$  for all  $n$ .

### Proof

- (a) Let  $\epsilon > 0$ . There exist  $N_1, N_2 \in \mathbb{N}$  such that  $|s_{n_1} - s| < \frac{\epsilon}{2}$  for  $n_1 \geq N_1$  and  $|t_{n_2} - t| < \frac{\epsilon}{2}$  for  $n_2 \geq N_2$ . Take  $N = \max N_1, N_2$ , and using the triangle inequality, it follows that for  $n \geq N$ :

$$|(s_n + t_n) - (s + t)| \leq |s_n - s| + |t_n - t| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

We conclude that  $\lim_{n \rightarrow \infty} (s_n + t_n) = s + t$ .

- (b) Let  $\epsilon > 0$ . If  $c = 0$  then the first sequence trivially converges to 0, so suppose that  $c \neq 0$ . There exists  $N$  such that  $|s_n - s| < \frac{\epsilon}{|c|}$  for  $n \geq N$ , so it follows that:

$$|cs_n - cs| = |c||s_n - s| < |c| \frac{\epsilon}{|c|} = \epsilon.$$

For the second identity, we have that  $c_n \rightarrow c$  for any constant sequence  $c_n = c$  so we may apply (a).

- (c) Let  $\epsilon > 0$ . There exist  $N_1, N_2$  such that  $|s_{n_1} - s| < \sqrt{\epsilon}$  for  $n_1 \geq N_1$  and  $|t_{n_2} - t| < \sqrt{\epsilon}$  for  $n_2 \geq N_2$ . We then consider that:

$$s_n t_n - st = (s_n - s)(t_n - t) + s(t_n - t) + t(s_n - s)$$

For  $n \geq N = \max N_1, N_2$ , we have that:

$$(s_n - s)(t_n - t) < \epsilon$$

And we hence observe that  $\lim_{n \rightarrow \infty} (s_n - s)(t_n - t) = 0$ . We can then use (a) and (b) to find that:

$$\lim_{n \rightarrow \infty} s(t_n - t) = 0, \quad \lim_{n \rightarrow \infty} t(s_n - s) = 0$$

So we conclude that  $\lim_{n \rightarrow \infty} (s_n t_n - st) = 0$  and hence  $s_n t_n \rightarrow st$ .

**Proof (Continued)**

(d) Choose  $m$  such that  $|s_n - s| < \frac{1}{2}|s|$  if  $n \geq m$ . We then have that  $|s_n| > \frac{1}{2}|s|$  for  $n \geq m$ . Let  $\epsilon > 0$ . Then, there exists  $N$  with  $N > m$  such that for  $n \geq N$ :

$$|s_n - s| < \frac{1}{2}|s|^2\epsilon$$

Hence, for  $n \geq N$ :

$$\left| \frac{1}{s_n} - \frac{1}{s} \right| = \left| \frac{s_n - s}{s_n s} \right| < \frac{2}{|s|^2} |s_n - s| < \epsilon$$

□

**Lemma: Squeeze Lemma**

Let  $\{x_n\}, \{s_n\}$  be real-valued sequences. Then, if  $0 \leq x_n \leq s_n$  for all  $n$ , and  $\lim_{n \rightarrow \infty} s_n = 0$ , then  $\lim_{n \rightarrow \infty} x_n = 0$ .

**Proof**

Let  $\epsilon > 0$ . Choose  $N \in \mathbb{N}$  such that  $n \geq N$  implies  $0 \leq s_n < \epsilon$ . Then, we have that for  $n \geq N$ ,  $0 \leq x_n \leq s_n < \epsilon$  and hence  $x_n \rightarrow 0$  as claimed. □

Note that we can prove a more generalized version of the Squeeze Lemma.

**Lemma: Generalized Squeeze Lemma**

Suppose we have sequences  $\{l_n\}, \{x_n\}, \{u_n\}$  such that  $l_n \leq x_n \leq u_n$  for all  $n$  and  $\lim_{n \rightarrow \infty} l_n = \lim_{n \rightarrow \infty} u_n = L \in \mathbb{R}$ . Then,  $\lim_{n \rightarrow \infty} x_n = L$ .

**Proof**

We have that  $0 \leq u_n - l_n \leq u_n - l_n$ . We have that  $\lim_{n \rightarrow \infty} u_n - l_n = 0$  by Theorem 3.3(a), so by the (original) Squeeze Lemma we have that  $\lim_{n \rightarrow \infty} u_n - l_n = 0$ . It then follows that  $\lim_{n \rightarrow \infty} u_n = \lim_{n \rightarrow \infty} l_n = L$  as claimed. □

**Theorem 3.20**

- (a) Let  $p > 0$ . Then,  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 0$ .
- (b) Let  $p > 0$ . Then,  $\lim_{n \rightarrow \infty} \sqrt[n]{p} = 1$ .
- (c)  $\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1$ .
- (d) Let  $p > 0$  and  $\alpha \in \mathbb{R}$ . Then,  $\lim_{n \rightarrow \infty} \frac{n^\alpha}{(1+p)^n} = 0$ .
- (e) Let  $|x| < 1$ . Then,  $\lim_{n \rightarrow \infty} x^n = 0$ .



## Proof

(a) Let  $\epsilon > 0$ . Choose  $N$  such that  $\frac{1}{N^p} < \epsilon$ , namely  $N > \left(\frac{1}{\epsilon}\right)^{1/p}$ . Then, for  $n \geq N$ ,  $\frac{1}{n^p} < \frac{1}{N^p} < \epsilon$ .

(b) If  $p = 1$ , the sequence is constant and the conclusion immediate.

If  $p > 1$ , then let  $x_n = \sqrt[p]{p} - 1$ . We then have that:

$$p = (x_n + 1)^n = \sum_{k=0}^n \binom{n}{k} x_n^k \geq nx_n$$

Where the second equality follows from the binomial theorem (where  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$ ), and the inequality follows by considering that we just keep the  $k = 1$  term (and the series is non-negative). Hence, we have that  $x_n \leq \frac{p}{n}$ , and  $x_n \rightarrow 0$  by (a).

If  $p < 1$ , then let  $q = \frac{1}{p} > 1$ . Then,  $\sqrt[q]{q} \rightarrow 1$  by the argument above. By Theorem 3.3(d), we then have that  $\sqrt[p]{p} = \frac{1}{\sqrt[q]{q}} \rightarrow \frac{1}{1} = 1$ .

(c) Let  $x_n = \sqrt[n]{n} - 1$ . Then, we have that:

$$n = (x_n + 1)^n = \sum_{k=0}^n \binom{n}{k} x_n^k \geq \frac{n(n-1)}{2} x_n^2$$

Where the inequality follows from keeping the  $k = 2$  term only. We then have that  $x_n \leq \sqrt{\frac{2}{n-1}}$  and hence  $x_n \rightarrow 0$  by the Squeeze Lemma.

(d) We want to show  $\frac{n^\alpha}{(1+p)^n} \rightarrow 0$ ; we therefore want an upper bound on the expression, and hence a lower bound on  $(1+p)^n$ . Applying the Binomial Theorem we have that:

$$(1+p)^n = \sum_{k=0}^n \binom{n}{k} p^k = ((n)(n-1)(n-2) \cdots (n-k+1)) \frac{p^k}{k!}$$

Now, we pick  $k > \alpha$ . For  $2n > k$ , we then have that:

$$(1+p)^n \geq \left(\frac{n}{2}\right)^k \frac{p^k}{k!}$$

We therefore have that:

$$\frac{n^\alpha}{(1+p)^k} \leq \frac{2^k k!}{p^k} n^{\alpha-k} \rightarrow 0$$

And the claim follows by the Squeeze Lemma.

(e) Taking  $\alpha = 0$  in (d), the claim follows by setting  $|x| = \frac{1}{1+p} < 1$  (as  $p > 0$ ) and recognizing that  $x_n \rightarrow 0 \iff |x^n| = |x|^n \rightarrow 0$ .  $\square$

### 3.2 Subsequences

#### Theorem 3.2

Let  $\{p_n\}$  be a sequence in  $X$ .

- (a)  $p_n \rightarrow p$  in  $X$  if and only if for all  $r > 0$ ,  $N_r(p)$  contains all but finitely many points of  $\{p_n\}$ .
- (b) If  $p_n \rightarrow p$  and  $p_n \rightarrow p'$  then  $p = p'$ . In other words, the limit is unique.
- (c) If  $\{p_n\}$  is convergent, then it is bounded (that is, for any  $q \in X$  there exists  $M \in \mathbb{R}$  such that  $d(q, p_n) \leq M$  for all  $n \in \mathbb{N}$ ).
- (d) If  $E \subset X$  has a limit point  $p$ , then there exists  $\{p_n\}$  in  $E$  such that  $p_n \rightarrow p$ .

#### Proof

- (a) The claim follows immediately from the definition of convergence; for any  $r = \epsilon > 0$ , there exists  $N \in \mathbb{N}$  such that  $N_r(p)$  contains  $\{p_n : n \geq N\}$ .
- (b) There exist  $N_1, N_2$  such that  $d(p, p_{n_1}) < \frac{\epsilon}{2}$  if  $n_1 \geq N_1$  and  $d(p, p_{n_2}) < \frac{\epsilon}{2}$  if  $n_2 \geq N_2$ . Then for  $n \geq N = \max N_1, N_2$  we have (using the triangle inequality) that:

$$d(p, p') \leq d(p, p_n) + d(p_n, p') < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

Since  $\epsilon$  is arbitrary,  $d(p, p') = 0$  and hence  $p = p'$ .

- (c) If  $p_n \rightarrow p$ , there exists  $N$  such that  $d(p_n, p) < 1$  for all  $n \geq N$ . Set:

$$r = \max \{1, d(p_1, p), d(p_2, p), \dots, d(p_{N-1}, p)\}$$

For any  $q \in X$ , we then have that:

$$d(q, p_n) \leq d(q, p) + d(p, p_n) \leq d(q, p) + r$$

so the claim follows with  $M = r + d(q, p) + 1$ .

- (d) Pick  $p_n \in E$  such that  $d(p_n, p) < \frac{1}{n}$ . Let  $\epsilon > 0$ , and  $N > \frac{1}{\epsilon}$ . Then,  $n \geq N$  implies  $\frac{1}{n} \leq \frac{1}{N} < \epsilon$  and hence  $d(p_n, p) < \epsilon$  for all  $n \geq N$ , and hence  $p_n \rightarrow p$  as desired.  $\square$

#### Definition 3.5: Subsequences

Given  $\{p_n\}$  and  $n_1 < n_2 < n_3 < \dots$ , we say that  $\{p_{n_j}\}$  is a **subsequence** of  $\{p_n\}$ .

We first consider some examples. Let  $p_n = n$ . Then some valid subsequences of  $\{p_n\}$  are  $\{1, 2, 3, 4, 5, \dots\}$  (the original sequence),  $\{1, 3, 5, 7, \dots\}$  (the odds),  $\{2, 3, 5, 7, 11, 13, \dots\}$  (the primes). Next, let  $p_n = i^n$ . We have that  $\{p_n\} = \{i, -1, -i, 1, i, -1, -i, 1, \dots\}$  which is clearly divergent. However, the subsequences  $\{i, i, i, \dots\}$ ,  $\{-1, -1, -1, \dots\}$ ,  $\{-i, -i, -i, \dots\}$  and  $\{1, 1, 1, \dots\}$  are all convergent! It is hence possible for a divergent sequence to have a convergent subsequence.

### Lemma

If  $p_n \rightarrow p$ , then every subsequence of  $\{p_n\}$  converges to  $p$ .

### Proof

Suppose  $p_n \rightarrow p$  and let  $\{p_{n_j}\}$  be a subsequence of  $p_n$ . Let  $\epsilon > 0$ . Then, there exists some  $N \in \mathbb{N}$  such that  $d(p, p_n) < \epsilon$  if  $n \geq N$ . Hence,  $d(p, p_{n_j}) < \epsilon$  if  $n_j \geq N$  and hence  $p_{n_j} \rightarrow p$ .  $\square$

### Theorem 3.6: Bolzano–Weierstrass

- (a) If  $\{p_n\} \subset X$  with  $X$  compact, then  $\{p_n\}$  has a convergent subsequence.
- (b) If  $\{p_n\} \subset \mathbb{R}^k$  and  $\{p_n\}$  is bounded, then  $\{p_n\}$  has a convergent subsequence.

### Proof

- (a) Let  $E$  be the range of  $\{p_n\}$ . If  $E$  is finite, then there exists  $x \in X$  and  $n_1 < n_2 < n_3 < \dots$  such that  $p_{n_j} = x$  for all  $j$ . Therefore  $p_{n_j} \rightarrow x$  and we are done. If  $E$  is infinite, then by compactness,  $E \subset X$  has a limit point in  $X$  by Theorem 2.37. By Theorem 3.2(d) there exists a sequence  $\{p_{n_j}\}$  in  $E$  such that  $p_{n_j} \rightarrow p$ .
- (b) By Theorem 2.41,  $E$  (being bounded) lies in a compact subset of  $\mathbb{R}^k$ . We then apply (a).  $\square$

## 3.3 Cauchy Sequences and Completeness

### Definition 3.8: Cauchy Sequences

A sequence  $\{p_n\} \subset X$  is a **Cauchy sequence** if for all  $\epsilon > 0$ , there exists  $N \in \mathbb{N}$  such that for all  $n, m \geq N$ ,  $d(p_n, p_m) < \epsilon$ .

Note the fact that this definition does not refer to a particular  $p$  that the sequence may converge to! It instead formalizes the notion of the points of a sequence getting “closer together” as the sequence goes on. It is therefore easier to check if a sequence is Cauchy than if it converges, as we don’t need to know the value of the limit. To this end, it is useful to know in what situations a sequence being Cauchy implies that the sequence is convergent. We will soon arrive at a theorem that addresses this question, but first we establish a little more machinery.

### Definition 3.9: Diameter

Let  $E \subset X$ . Then the **diameter** of  $E$ , denoted  $\text{diam } E$  is defined as  $\text{diam } E = \sup \{d(p, q) : p, q \in E\}$ . It follows from the definition that a sequence  $\{p_n\}$  is Cauchy if and only if  $\lim_{N \rightarrow \infty} \text{diam } E_n = 0$  where  $E_n = \{p_n\}_{n=N}^{\infty}$  (the tail of the sequence).

### Example

- (a) If  $E = (a, b) \subset \mathbb{R}$  or  $E = [a, b] \subset \mathbb{R}$ , then  $\text{diam } E = b - a$ .
- (b) If  $E = (0, 1) \times (0, 1) \subset \mathbb{R}^2$ , then  $\text{diam } E = \sqrt{2}$  (the diagonal of the open square).

### Theorem 3.10

- (a) Let  $E \subset X$ . Then,  $\text{diam } \bar{E} = \text{diam } E$ .
- (b) If  $K_n \subset X$  are compact,  $K_{n+1} \subset K_n$  for all  $n$ , and  $\lim_{n \rightarrow \infty} \text{diam } K_n = 0$ , then  $\bigcap_{n=1}^{\infty} K_n$  consists of exactly one point.

### Proof

- (a) Since  $E \subset \bar{E}$ , it is clear that  $\text{diam } \bar{E} \geq \text{diam } E$ . Next, let  $\epsilon > 0$  and  $p, q \in \bar{E}$ . Choose  $p', q' \in E$  such that  $d(p, p') < \frac{\epsilon}{2}$ ,  $d(q, q') < \frac{\epsilon}{2}$  (this choice is possible as either  $p, q$  are in  $E$ , or  $p, q$  are limit points of  $E$ ). Then, we have that:

$$d(p, q) \leq d(p, p') + d(p', q) \leq d(p, p') + d(p', q') + d(q', q) < \frac{\epsilon}{2} + \text{diam } E + \frac{\epsilon}{2} = \text{diam } E + \epsilon$$

$\epsilon, p$ , and  $q$  are arbitrary, so it follows that  $\text{diam } \bar{E} \leq \text{diam } E + \epsilon$  from the definition of the diameter. It then follows that  $\text{diam } \bar{E} \leq \text{diam } E$ . We conclude that  $\text{diam } \bar{E} = \text{diam } E$ .

- (b) Let  $K = \bigcap_{n=1}^{\infty} K_n$ . By the corollary to Theorem 2.36, we have that  $K \neq \emptyset$ , so  $K$  contains at least one point. Since  $K \subset K_n$ , it follows that  $\text{diam } K \leq \text{diam } K_n$  for any  $n$ , and since  $\text{diam } K_n \rightarrow 0$ ,  $\text{diam } K = 0$ . If there were  $p, q \in K$  such that  $p \neq q$ , then  $\text{diam } K \neq 0$ , so it must follow that  $K$  has at most one point. We conclude that  $K$  has exactly one point.  $\square$

### Lemma

If a sequence  $\{p_n\}$  is Cauchy, then it is bounded.

### Proof

If  $\{p_n\}$  is Cauchy, then we have that  $\lim_{N \rightarrow \infty} \text{diam } E_N = \lim_{N \rightarrow \infty} \text{diam } \{p_n\}_{n=N}^{\infty} = 0$ . Then for some  $N \in \mathbb{N}$ ,  $\text{diam } E_N < 1$ . The range of  $\{p_n\}$  is the union of  $E_N$  and the finite set  $\{p_1, \dots, p_{N-1}\}$  and hence  $\{p_n\}$  is bounded.  $\square$

### Theorem 3.11

- (a) If a sequence  $\{p_n\} \subset X$  converges, then it is Cauchy.
- (b) If a sequence  $\{p_n\} \subset X$  is Cauchy and  $X$  is compact, then  $\{p_n\}$  converges to some  $p \in X$ .
- (c) In  $\mathbb{R}^k$ , every Cauchy sequence is convergent.

### Proof

- (a) Let  $p_n \rightarrow p$  and let  $\epsilon > 0$ . There exists  $N \in \mathbb{N}$  such that  $d(p_n, p) < \frac{\epsilon}{2}$  if  $n \geq N$ . Then, for  $n, m \geq N$ , we have that:

$$d(p_n, p_m) \leq d(p_n, p) + d(p, p_m) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

so  $\{p_n\}$  is Cauchy.

- (b) Let  $E_N = \{p_n\}_{n=N}^{\infty}$ . Then,  $\bar{E}_N \subset X$  is closed, so by the compactness of  $X$  we have that  $\bar{E}_N$  is compact by Theorem 2.35. Since  $E_{N+1} \subset E_N$ , we have that  $\bar{E}_{N+1} \subset \bar{E}_N$ , and additionally we have that  $\lim_{N \rightarrow \infty} \bar{E}_N = \lim_{N \rightarrow \infty} E_N = 0$  where the first equality follows from Theorem 3.10(a) and the second equality follows from the fact that  $\{p_n\}$  is Cauchy and Definition 3.9. Thus, Theorem 3.10(b) says that there exists a unique point  $p \in \bigcap_{n=1}^{\infty} \bar{E}_N$ . Next, let  $\epsilon > 0$ . Then, there exists  $N_0$  such that  $\text{diam } \bar{E}_N < \epsilon$  for all  $N \geq N_0$ . So,  $d(p, q) < \epsilon$  for all  $q \in \bar{E}_N$ , so the same holds for all  $q \in E_N$ . Hence,  $d(p, p_n) < \epsilon$  for all  $n \geq N_0$ , which shows that  $p_n \rightarrow p$  and proves the claim.

- (c) By the above Lemma, Cauchy sequences are bounded. Hence,  $\{p_n\} \subset I$  for some  $k$ -cell  $I \subset \mathbb{R}^k$ . Since  $I$  is compact in  $\mathbb{R}^k$ , the claim follows from (b).  $\square$

### Definition 3.12: Completeness

A metric space  $X$  is called **complete** if every Cauchy sequence converges in  $X$ .

It might be tempting at first to think that every space would be complete, but this is not the case. For example, something that can go wrong is a sequence can be Cauchy, but the limit can lie “outside” of the space. To see this, consider again the sequence  $\left\{\frac{1}{n}\right\}$  in the metric space  $\mathbb{R}^+ = \mathbb{R} \setminus \{x \in \mathbb{R} : x \leq 0\}$ . The sequence is Cauchy, but does not converge in  $\mathbb{R}^+$  (as it converges to 0, which lies outside of the space).

### Example

- (i) Compact sets are complete by Theorem 3.11(b).
- (ii)  $\mathbb{R}^k$  (and  $\mathbb{C}$ ) are complete by Theorem 3.11(c).
- (iii)  $\mathbb{Q}$  is not complete. We can make a sequence of rational points that converges to an irrational number in  $\mathbb{R}$  which is Cauchy, but does not converge in  $\mathbb{Q}$  (Example 1.1 gives a way one might construct such a sequence).

Note that  $\mathbb{Q}$  can be completed to  $\mathbb{R}$ , and in general for any  $(X, d)$  which is not complete, there exists  $(X^*, d^*)$  that is complete such that  $|X| = X^*$ . Indeed, this is another way we can construct the real numbers!  $\mathbb{R}$  can be viewed as equivalence classes of Cauchy sequences in  $\mathbb{Q}$ . The idea is to define an equivalence relation  $\sim$  such that  $p_n \sim q_n$  if  $\lim_{n \rightarrow \infty} d(p_n, q_n) = 0$ .  $X^*$  is then defined as the set of equivalence classes under that equivalence relation, equipped with the metric  $d^*([p], [q]) = \lim_{n \rightarrow \infty} d(p_n, q_n)$ . It can then be checked that  $d^*$  is a valid metric and that  $X^*$  is complete. For the full proof, see HW7, or exercises 3.23-3.25 in Rudin (note: this proof is quite technical/difficult).

To motivate the next theorem, consider that all convergent sequences in  $\mathbb{R}$  (and in general) are bounded (as we saw in Theorem 3.2(c)). However, this is not always true; for example consider  $p_n = (-1)^n$  which is clearly bounded but divergent. What then are conditions that a bounded sequence may converge?

### Definition 3.13: Monotonic Sequences

A sequence  $\{p_n\} \subset \mathbb{R}$  is **monotonically increasing** if  $p_{n+1} \geq p_n$  for all  $n$ , and **monotonically decreasing** if  $p_{n+1} \leq p_n$ .

### Theorem 3.14

Suppose  $\{p_n\} \subset \mathbb{R}$  is monotonic. Then,  $\{p_n\}$  is convergent if and only if it is bounded.

### Proof

$\Rightarrow$  See Theorem 3.2(c).

$\Leftarrow$  We show the proof for the increasing case as the decreasing case is analogous. Let  $p = \sup p_n : n \in \mathbb{N}$  which exists as  $\{p_n\}$  is bounded and  $\mathbb{R}$  has the LUB property. Then,  $p_n \leq p$  for all  $n$ . Let  $\epsilon > 0$ . Then, there exists  $N \in \mathbb{N}$  such that  $p - \epsilon < p_N < p_{N+1}$ . By the monotonicity of  $\{p_n\}$ , it follows that  $|p_n - p| < \epsilon$  for all  $n \geq N$ . Hence,  $p_n \rightarrow p$ .  $\square$

### Definition 3.15: Limits to Infinity

Let  $\{p_n\} \subset \mathbb{R}$ . If for all  $M \in \mathbb{R}$ , there exists  $N \in \mathbb{N}$  such that  $p_n > M$  for all  $n \geq N$ , then we write  $p_n \rightarrow \infty$ . If instead for all  $M \in \mathbb{R}$  there exists  $N \in \mathbb{N}$  such that  $p_n < M$  for all  $n \geq N$ , then we write  $p_n \rightarrow -\infty$ .

## 3.4 Limit Supremum and Limit Infimum

As a motivating question, how would we say something about the largest and smallest accumulation points of a sequence? This leads us to the following definition.

### Definition 3.16: limsup and liminf

Let  $\{s_n\} \subset \mathbb{R}$ , then, we define the **limit supremum** as:

$$\limsup_{n \rightarrow \infty} s_n = \inf_{n \geq 1} \sup_{m \geq n} s_m = \lim_{n \rightarrow \infty} \sup_{m \geq n} s_m$$

And the **limit infimum** as:

$$\liminf_{n \rightarrow \infty} s_n = \sup_{n \geq 1} \inf_{m \geq n} s_m = \lim_{n \rightarrow \infty} \inf_{m \geq n} s_m$$

Note that unlike the limit of a real-valued sequence, the limsup and liminf always exist.

In the above definition, the equivalence of  $\inf_{n \geq 1} \sup_{m \geq n} s_m$  and  $\lim_{n \rightarrow \infty} \sup_{m \geq n} s_m$  may be slightly confusing. To see this, consider the fact that the sequence  $q_n = \sup \{p_n : n \geq 1\}$  is a strictly decreasing sequence in  $n$  (with increasing  $n$ , we take the supremum over less terms each time), so taking the limit of  $n \rightarrow \infty$  or taking the infimum over  $n$  are equivalent.

Note that Rudin defines the limsup/liminf differently, but perfectly equivalently. Namely, if  $\{p_n\} \subset \mathbb{R}$  is a sequence, then  $E$  is the set of all subsequential limits (i.e. there is a subsequence of  $\{p_n\}$  with a given limit). Then,  $\limsup^* p_n = \sup E$  (we use the  $*$  to denote Rudin's definition). The equivalence is not immediately obvious, so we here give a sketch to show that the two definitions coincide. We will use the

general technique of showing that the two expressions are  $\epsilon$  close to one another. Namely, for any  $\epsilon > 0$ , we show that the following two statements hold:

- 1)  $\limsup^* p_n \leq \limsup p_n + \epsilon$
- 2)  $\limsup^* p_n + \epsilon \geq \limsup p_n$

To show 1), for each  $N$ , we let  $n_N$  be an index such that  $p_{n_N}$  satisfies:

$$p_{n_N} \geq \sup \{p_n : n \geq N\} - \frac{\epsilon}{N}$$

And then we claim that  $\lim_{n \rightarrow \infty} p_{n_N} = \limsup p_n$  (Exercise). There is one slight technical issue in that  $\{p_{n_N}\}$  may not be a subsequence of  $\{p_n\}$ , in particular we don't know that  $n_{N_1} < n_{N_2} < n_{N_3} < \dots$  just by the above construction (but in order for this to be a valid subsequence, we need this to be the case). Fortunately, this is a fixable issue. We do know that  $n_{N_1} \geq 1$ ,  $n_{N_2} \geq 2$ ,  $n_{N_3} \geq 3$  by construction, so if it turns out to be the case that  $n_{N_1} < n_{N_2}$  doesn't hold, we can skip ahead into the sequence until we find the first  $j$  for which the equality holds. Concretely, if  $n_{N_1} = 1000$  (as an example), then if we skip ahead to  $n_{N_{1001}}$ , it is guaranteed that  $n_{N_1} < n_{N_{1001}}$  and we can from there construct a valid sequence. The sketch for 2) is left as an exercise.

### Example

- (a) Consider  $s_n = (-1)^n \left(1 + \frac{1}{n^2}\right)$ . Then, we have that  $1 \leq \sup_{m \geq n} s_m \leq 1 + \frac{1}{1^2} = 2$ , so  $\limsup_{n \rightarrow \infty} s_n = 1$ . Similarly,  $\limsup_{n \rightarrow \infty} s_n = -1$ . Note that this sequence has no limit (it oscillates indefinitely and does not converge) but these quantities are well defined. We notice that the limsup is greater than the liminf in this case, and indeed it is true in general that  $\liminf_{n \rightarrow \infty} s_n \leq \limsup_{n \rightarrow \infty} s_n$ .
- (b) If  $\{s_n\}$  is not bounded above, then  $\sup_{m \geq n} s_m = \infty$  for all  $n$  and we write  $\limsup_{n \rightarrow \infty} s_n = \infty$ . Similarly, if  $\{s_n\}$  is not bounded below, then  $\inf_{m \geq n} s_m = -\infty$  for all  $n$  and we write  $\limsup_{n \rightarrow \infty} s_n = -\infty$ .

One difficulty with discussing the convergence of a sequence is that the definition is difficult to apply; we need to know what the sequence converges to. The notion of a Cauchy sequence then begins helpful (as Cauchy and convergent are equivalent in complete metric spaces). In addition, it is helpful to consider the limsup and liminf, as we can bound the limit above and below with these quantities respectively. In particular, the limit of the supremum of the tail of the sequence equals the limit of the infimum of the tail of the sequence equals the limit of the sequence if the sequence is convergent.

### Theorem 3.18

Let  $\{s_n\} \subset \mathbb{R}$ . Then,  $\lim_{n \rightarrow \infty} s_n = L$  if and only if  $\limsup_{n \rightarrow \infty} s_n = \liminf_{n \rightarrow \infty} s_n = L$ .

### Proof

$\Rightarrow$  Let  $\epsilon > 0$ . Then, there exists  $N \in \mathbb{N}$  such that  $s_m \in (L - \epsilon, L + \epsilon)$  for all  $m \geq N$ . Then, we have that:

$$L - \epsilon \leq \inf_{m \geq N} s_m \leq \sup_{m \geq N} s_m \leq L + \epsilon$$

Taking limits we have:

$$L - \epsilon \leq \liminf_{n \rightarrow \infty} s_n \leq \limsup_{n \rightarrow \infty} s_n \leq L + \epsilon$$

$\epsilon$  is arbitrary, so we have that  $\limsup_{n \rightarrow \infty} s_n = \liminf_{n \rightarrow \infty} s_n = L$ .

$\Leftarrow$  We have that:

$$\inf_{m \geq n} s_m \leq s_n \leq \sup_{m \geq n} s_m$$

for all  $n \in \mathbb{N}$ . By assumption we have  $\limsup_{n \rightarrow \infty} s_n = \liminf_{n \rightarrow \infty} s_n = L$  so by the Generalized Squeeze Lemma we conclude that  $\lim_{n \rightarrow \infty} s_n = L$ .  $\square$

## 3.5 Series

### Definition 3.21: Infinite Series

Let  $\{a_n\} \subset \mathbb{C}$ . We then form a new sequence of  $s_n = \sum_{j=1}^n a_j$  (the sequence of partial sums). Then, if  $s_n \rightarrow s$ . We say that the series  $\sum_{j=1}^{\infty} a_j$  converges, and write  $\sum_{j=1}^{\infty} a_j = s$ . If  $s_n$  does not converge, we say that  $\sum_{j=1}^{\infty} a_j$  diverges. As a notational point, we will sometimes omit the bounds of summation and write  $\sum a_j$  to denote an infinite series, where the meaning is clear from context.

Note that series are just a specific subset of sequences. Although the above definition states that  $\{a_n\} \subset \mathbb{C}$ , it is in general possible to define series over general vector spaces, with  $\{a_n\} \in V$  and  $\{s_n\} \in V$ .

Note that this generalization allows us to state an equivalent notion of completeness for vector spaces, namely that  $V$  is complete if and only if for all sequences  $\{a_n\} \subset V$ , the sum  $\sum_{n=0}^{\infty} \|a_n\|$  converges to a point in  $V$ . As before,  $\mathbb{R}, \mathbb{R}^k$  (both over the field  $\mathbb{R}$ ), are complete vector spaces. An example of a vector space that is not complete is the set of functions  $f : \mathbb{R} \mapsto \mathbb{C}$  such that  $\int_{-\infty}^{\infty} |f(x)|^2 dx < \infty$  (where the integral is the familiar Riemann integral from first year calculus, to be defined more rigorously in Chapter 6). For example, the sequence  $f_n \subset V$  such that:

$$f_n = \begin{cases} 1 & \text{for the first } n \text{ rationals} \\ 0 & \text{elsewhere} \end{cases}$$

does not converge to a function in  $V$ . In fact, Lebesgue integration (which is not covered in this course, but will be the primary focus of a course in measure theory) deals with this issue.

We will now restate the Cauchy criterion (Theorem 3.11) for series.

### Theorem 3.22

A series  $\sum a_j$  converges if and only if for all  $\epsilon > 0$ , there exists  $N \in \mathbb{N}$  such that  $\left| \sum_{j=m}^n a_j \right| < \epsilon$  for all  $n \geq m \geq N$ .



**Proof**

$\sum a_j$  converges if and only if  $\{s_n\}$  converges if and only if  $\{s_n\}$  is Cauchy. So by definition there exists  $N$  such that for  $n \geq m - 1 \geq N$ :

$$|s_n - s_m| = \left| \sum_{j=1}^n a_j - \sum_{j=1}^{m-1} a_j \right| = \left| \sum_{j=m}^n a_j \right| < \epsilon$$

which proves the claim.  $\square$

**Theorem 3.23: Divergence Test**

If  $\sum a_n$  converges, then  $\lim_{n \rightarrow \infty} a_n = 0$ .

**Proof**

Choose  $n = m$  in Theorem 3.22.  $\square$

Note that this criteria gives us the ability to easily check if a series diverges; if  $a_j$  does not converge to 0, then  $\sum a_j$  diverges. However, note that the reverse implication does NOT hold!  $\sum \frac{1}{n}$  diverges (as we will show next lecture) even though clearly  $\frac{1}{n} \rightarrow 0$ .

**Theorem 3.24**

Let  $\{a_n\} \subset \mathbb{R}$  and  $a_n \geq 0$  for all  $n$ . Then, we have that  $\sum a_n$  converges if and only if the sequence of partial sums  $\{s_n\}$  is bounded.

**Proof**

If  $a_n \geq 0$ , then  $\{s_n\}$  is monotonically increasing. Then by Theorem 3.14,  $\{s_n\}$  converges if and only if it is bounded.  $\square$

**Theorem 3.25: Comparison Test**

- (a) Let  $\{a_n\} \subset \mathbb{C}$  and  $\{c_n\} \subset \mathbb{R}$ . Then, if  $|a_n| \leq c_n$  for all  $n \geq N_0$  for some  $N_0 \in \mathbb{N}$ , and  $\sum c_n$  converges, then  $\sum a_n$  converges.
- (b) Let  $\{a_n\} \subset \mathbb{R}$  and  $\{d_n\} \subset \mathbb{R}$ . If  $a_n \geq d_n \geq 0$  for all  $n \geq N_0$  for some  $N_0 \in \mathbb{N}$  and  $\sum d_n$  diverges, then  $\sum a_n$  diverges.

**Proof**

- (a) Let  $\epsilon > 0$ . Then, there exists  $N \in \mathbb{N}$  such that  $\sum_{j=m}^n c_j < \epsilon$  for all  $n \geq m \geq N$ . Then, take  $N \geq N_0$ , and we have that:

$$\left| \sum_{j=m}^n a_j \right| \leq \sum_{j=m}^n |a_j| \leq \sum_{j=m}^n c_j < \epsilon$$

Where in the first inequality we apply the triangle inequality (Theorem 1.37). We conclude that  $\sum a_j$  converges by the Cauchy criterion (Theorem 3.22).

- (b) The claim follows by considering the contrapositive of (a). □

**Theorem 3.26: The Geometric Series**

If  $0 \leq x < 1$ , then  $\sum_{j=0}^{\infty} x^j = \frac{1}{1-x}$ . If  $x \geq 1$ , then  $\sum_{j=0}^{\infty} x^j$  diverges.

**Proof**

Suppose  $x = 1$ . Then,  $x_n = 1$  does not converge to zero, so by Theorem 3.24  $\sum_{j=0}^{\infty} x^j$  diverges. Suppose then that  $x \neq 1$ . We have that  $s_n = 1 + x + x^2 + \dots + x^n$ . Hence,  $xs_n = x + x^2 + x^3 + \dots + x^{n+1}$ . Hence,  $(1-x)s_n = 1 + x^{n+1}$  and therefore a closed form expression for  $s_n$  is:

$$s_n = \frac{1 + x^{n+1}}{1 - x}$$

If  $x > 1$ , we have that  $s_n$  diverges (again) by Theorem 3.24. If  $x < 1$ , then  $x^{n+1} \rightarrow 0$  by 3.20(e), so  $s_n \rightarrow \frac{1}{1-x}$ . □

Note that by the comparison test, the above result can be generalized to see that  $\sum z^j$  for  $z \in \mathbb{C}$  converges for  $|z| < 1$  and diverges for  $|z| \geq 1$ .

**Theorem 3.27: Cauchy Condensation Test**

Suppose  $\{a_n\} \subset \mathbb{R}$  and  $a_1 \geq a_2 \geq a_3 \geq \dots \geq 0$ . Then,  $\sum a_j$  converges if and only if  $\sum_{n=1}^{\infty} 2^n a_{2^n}$  converges.

**Proof**

$\Rightarrow$  For  $2^k < n$ , using the fact that the sequence is decreasing, we have that:

$$\begin{aligned} a_1 + a_2 + \dots + a_n &\geq a_1 + a_2 + (a_3 + a_4) + \dots + (a_{2^{k-1}+1} + \dots + a_{2^k}) \\ &\geq \frac{1}{2}a_1 + a_2 + 2a_4 + \dots + 2^{k-1}a_{2^k} \\ &= \frac{1}{2}(a_1 + 2a_2 + 4a_4 + \dots + 2^k a_{2^k}) \end{aligned}$$

Hence by the comparison test (Theorem 3.25) we have that  $\sum 2^n a_{2^n}$  converges if  $\sum a_j$  converges.

$\Leftarrow$  We show the contrapositive. For  $2^k > n$ , We have that:

$$\begin{aligned} a_1 + a_2 + \dots + a_n &\leq a_1 + (a_2 + a_3) + \dots + (a_{2^k} + \dots + a_{2^{k+1}-1}) \\ &\leq a_1 + 2a_2 + 4a_4 + \dots + 2^k a_{2^k} \end{aligned}$$

Hence by the comparison test (Theorem 3.25) we have that  $\sum 2^n a_{2^n}$  diverges if  $\sum a_j$  diverges.  $\square$

**3.6 p-Series and Euler's Number**

We now use the result of Theorem 3.27 to prove a result about a familiar subset of series.

**Theorem 3.28: p-Series**

$\sum_{n=1}^{\infty} \frac{1}{n^p}$  converges if  $p > 1$  and diverges if  $p \leq 1$ .

**Proof**

If  $p \leq 0$ , then  $n^p$  does not converge to 0 and hence  $\sum \frac{1}{n^p}$  diverges by Theorem 3.23. If  $p > 0$ , then  $\frac{1}{n^p}$  is a monotonically decreasing sequence of positive terms. Hence, we can apply the result of Theorem 3.27.  $\sum \frac{1}{n^p}$  converges if and only if  $\sum_k 2^k \frac{1}{2^{kp}} = \sum_k \left(\frac{1}{2^{p-1}}\right)^k$  converges. By Theorem 3.26, this last expression is convergent if and only if  $0 < \frac{1}{2^{p-1}} < 1$ , i.e. if  $p > 1$ , proving the claim.  $\square$

From the above result, we have that the  $p$ -series converges if  $p > 1$  and diverges otherwise. Is there something “in between” these two regions?

**Theorem 3.29**

$\sum_{n=2}^{\infty} \frac{1}{n(\log n)^p}$  converges if  $p > 1$  and diverges if  $p \leq 1$ .

**Proof**

$\log n$  is monotonically increasing for  $p > 0$ . So,  $\frac{1}{n(\log n)^p}$  is monotonically decreasing. Hence,  $\sum \frac{1}{n(\log n)^p}$  converges if and only if  $\sum_k 2^k \frac{1}{2^k (\log 2^k)^p} = \sum_k \frac{1}{k^p}$  converges, and hence the claim follows from Theorem 3.28.  $\square$

**Definition 3.30: Euler's Number**

$e = \sum_{n=0}^{\infty} \frac{1}{n!}$ , where  $0! = 1$  and  $n! = n \cdot (n-1)! = n \cdot (n-1) \cdot \dots \cdot 2 \cdot 1$  for  $n \geq 1$ .

We should check that this expression is well defined (namely, that the series converges). We first observe that  $\frac{1}{n!} \leq \frac{1}{2^{n-1}}$  for  $n \geq 1$ . Therefore, we have that:

$$s_n = \sum_{j=0}^n \frac{1}{j!} \leq 1 + \sum_{j=1}^n \frac{1}{2^{j-1}} \leq 1 + \sum_{j=0}^{\infty} \frac{1}{2^j} = 1 + \frac{1}{1 - \frac{1}{2}} = 3$$

Where we use Theorem 3.26 in the second last equality. Hence, we conclude that  $\sum_{j=0}^{\infty} \frac{1}{j!}$  converges by comparison, and moreover, that  $0 < e < 3$ .

It will also be of interest to investigate the rate of convergence of this series. To this end, we observe:

$$\begin{aligned} 0 < e - s_q &= \frac{1}{(q+1)!} + \frac{1}{(q+2)!} + \dots \leq \frac{1}{(q+1)!} \left( 1 + \frac{1}{q+1} + \frac{1}{(q+1)^2} + \dots \right) \\ &= \frac{1}{(q+1)!} \frac{1}{1 - \frac{1}{q+1}} = \frac{1}{(q+1)!} \frac{q+1}{q} = \frac{1}{q!q} \end{aligned}$$

Hence we have that the error goes to zero extremely quickly! Moreover, we can use this fact to show that  $e$  is irrational.

**Theorem 3.32: Irrationality of  $e$** 

$e \notin \mathbb{Q}$ .

**Proof**

Suppose  $e = \frac{p}{q}$  for  $p, q \in \mathbb{N}$ . Then, by the argument above, we have that  $0 < e - s_q < \frac{1}{q!q}$ . Hence, we have that  $0 < q!e - q!s_q < \frac{1}{q} \leq 1$ . But,  $q!e = q! \frac{p}{q} = (q-1)!p \in \mathbb{N}$ , and  $q!s_q = \sum_{j=0}^q q! \frac{1}{j!} \in \mathbb{N}$ . Hence,  $q!e - q!s_q \in \mathbb{Z}$ , but this is a contradiction as  $0 < q!e - q!s_q < 1$ .  $\square$

There is another familiar definition of  $e$  involving a limit that one may recall from first year calculus. These definitions are equivalent, as we will show in the next theorem.

**Theorem 3.31**

$$e = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right)^n.$$

### Proof

Let  $t_n = \left(1 + \frac{1}{n}\right)^n$ . By the Binomial theorem, we have that:

$$t_n = \sum_{j=0}^n \binom{n}{j} \left(\frac{1}{n}\right)^j = \sum_{j=0}^n \frac{1}{j!} \left(\frac{n}{n} \cdot \frac{(n-1)}{n} \cdots \frac{(n-j+1)}{n}\right) \leq s_n$$

Where the last inequality follows from the fact that the term in brackets is less than (or equal to) 1. Hence, we have that  $\limsup_{n \rightarrow \infty} t_n \leq \limsup_{n \rightarrow \infty} s_n = \lim_{n \rightarrow \infty} s_n = e$ . On the other hand, fix  $m \in \mathbb{N}$  and let  $n \geq m$ . Then, we have that:

$$t_n \geq \sum_{j=0}^m \binom{n}{j} \frac{1}{n^j} = \sum_{j=0}^m \frac{1}{j!} \left( \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{j-1}{n}\right) \right)$$

The infimum of the term in the brackets is just 1, so we therefore have that:

$$\inf_{n \geq m} t_n = \sum_{j=0}^m \frac{1}{j!} = s_m$$

Now, we take  $m \rightarrow \infty$  to find that  $\liminf_{m \rightarrow \infty} t_m \geq \liminf_{m \rightarrow \infty} s_m = \lim_{m \rightarrow \infty} s_m = e$ .

Having shown that  $\liminf_{n \rightarrow \infty} t_n \geq e \geq \limsup_{n \rightarrow \infty} t_n$ , we conclude that  $\limsup_{n \rightarrow \infty} = \liminf_{n \rightarrow \infty} = e$  and hence  $\lim_{n \rightarrow \infty} t_n = e$ .  $\square$

## 3.7 The Ratio and Root Tests

### Theorem 3.33: The Root Test

Let  $\sum a_n$  be a series, and put  $\alpha = \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}$ . Then,

- (i)  $\sum a_n$  converges if  $\alpha < 1$ .
- (ii)  $\sum a_n$  diverges if  $\alpha > 1$ .
- (iii) If  $\alpha = 1$ , the test is inconclusive.

### Proof

- (i) Suppose  $\limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|} = \alpha < 1$ . Take  $\beta$  such that  $\alpha < \beta < 1$  and  $N \in \mathbb{N}$  such that  $\sqrt[n]{|a_n|} < \beta$  for all  $n \geq N$ . Hence, for  $n \geq N$ ,  $|a_n| < \beta^n$ , and  $\beta < 1$ . The result follows by using the comparison Test with the geometric series.
- (ii) Suppose  $\limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|} = \alpha > 1$ . Then, there exists a subsequence such that  $\sqrt[n_j]{|a_{n_j}|} \rightarrow \alpha$ . Therefore, there exists  $N$  such that for  $j \geq N$ ,  $\sqrt[n_j]{|a_{n_j}|} > 1$ , that is to say,  $\sqrt[n_j]{|a_{n_j}|} > 1$  for infinitely many terms. Hence,  $\sqrt[n]{|a_n|}$  does not converge to zero, and hence the series does not converge by the divergence test.
- (iii) Consider  $\sum \frac{1}{n}$  and  $\sum \frac{1}{n^2}$ .  $\alpha = 1$  for both sums, but by Theorem 3.28 the former diverges and the latter converges.  $\square$

### Theorem 3.34: The Ratio Test

Let  $\sum a_n$  be a series such that  $a_n \neq 0$  for all  $n$ . Then,

- (i)  $\sum a_n$  converges if  $\limsup_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$ .
- (ii) Diverges if there exists  $N_0$  such that  $\left| \frac{a_{n+1}}{a_n} \right| \geq 1$  for all  $n \geq N_0$ .

In other cases, the test is inconclusive.

### Proof

- (i) By assumption, there exists  $\beta < 1$  such that for some  $N$ ,  $\left| \frac{a_{n+1}}{a_n} \right| < \beta$  for all  $n \geq N$ . We then have that  $|a_{N+1}| < \beta|a_N|$ , that  $|a_{N+1}| < \beta|a_{N+1}| < \beta^2|a_N|$  and inductively we obtain that  $|a_{N+p}| < \beta^p|a_N|$ . In other words, we have that for  $n \geq N$ ,  $|a_n| < |a_N|\beta^{-N}\beta^n$ . Since  $\sum \beta^n$  converges (convergent geometric series),  $\sum a_n$  converges by the comparison test.
- (ii) For  $n \geq N_0$ , we have that  $|a_n| \leq |a_{N+1}|$ . Hence,  $a_n$  does not converge to 0, and the claim follows by the divergence test.  $\square$

As a remark, the the ratio test is less powerful than the root test. For any series for which the ratio test is conclusive, the root test is also conclusive. But the converse is not true. However, the ratio test is easier to apply in practice. We also note that the above ratio test implies the (perhaps more familiar) version from first year calculus:

### Corollary

Let  $\sum a_n$  be a series such that  $a_n \neq 0$  for all  $n$ . Then:

- (i)  $\sum a_n$  converges if  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$ .
- (ii)  $\sum a_n$  diverges if  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| > 1$ .
- (iii)  $\sum a_n$  diverges if  $\liminf \left| \frac{a_{n+1}}{a_n} \right| > 1$ .

### Example 3.35

Consider the series:

$$\frac{1}{2} + 1 + \frac{1}{8} + \frac{1}{4} + \frac{1}{32} + \frac{1}{16} + \dots$$

We then have that the ratio  $\frac{a_{n+1}}{a_n}$  is the sequence  $2, \frac{1}{8}, 2, \frac{1}{8}, \dots$ . Therefore,  $\limsup_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 2 > 1$  and  $\liminf_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{8} < 1$  and the ratio test is inconclusive/tells us nothing. We then consider the root test. For  $n \geq 3$ , we have that:

$$a_n = \begin{cases} \left(\frac{1}{4}\right)^k = \left(\frac{1}{4}\right)^{\frac{n-1}{2}} = 2\left(\frac{1}{4}\right)^{\frac{n}{2}} & n = 2k + 1 \\ \frac{1}{2} \left(\frac{1}{4}\right)^k = \frac{1}{2} \left(\frac{1}{4}\right)^{\frac{n}{2}} & n = 2k \end{cases}$$

Since  $\lim_{n \rightarrow \infty} \sqrt[n]{p} = 1$  for  $p > 0$  (Theorem 3.20(b)), we have that:

$$\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \lim_{n \rightarrow \infty} \sqrt[n]{c \left(\frac{1}{4}\right)^{\frac{n}{2}}} = \lim_{n \rightarrow \infty} \sqrt[n]{c} \frac{1}{2} = \frac{1}{2} < 1$$

where the above limit holds for either  $c = \frac{1}{2}$  or  $c = 2$ . Hence, we conclude that the series converges by the root test. This example demonstrates how the root test is “sharper” than the ratio test (though harder to apply).

## 3.8 Power Series

### Definition 3.38: Power Series

For  $z \in \mathbb{C}$  and a sequence  $\{c_n\}$ ,  $\sum_{n=0}^{\infty} c_n z^n$  is called a **power series**.

### Theorem 3.39: Radius of Convergence

Let  $R = \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{|c_n|}}$ , with the convention  $R = \infty$  if  $\limsup_{n \rightarrow \infty} \sqrt[n]{|c_n|} = 0$  and  $R = 0$  if  $\limsup_{n \rightarrow \infty} \sqrt[n]{|c_n|} = \infty$ . Then,  $\sum_{n=0}^{\infty} c_n z^n$  converges if  $|z| < R$  and diverges if  $|z| > R$ .  $R$  is called the **radius of convergence** of  $\sum c_n z^n$ . We note that on the circle  $|z| = R$ , the behavior is varied; the series can be divergent or convergent, and it can also depend on the particular choice of  $z$  on the circle.

### Proof

We have that  $\limsup_{n \rightarrow \infty} \sqrt[n]{c_n z^n} = \limsup_{n \rightarrow \infty} \sqrt[n]{c_n} |z| = \frac{|z|}{R}$ . Therefore, by the root test (Theorem 3.33) the series converges if  $|z| < R$  and diverges if  $|z| > R$  (and nothing can be said if  $|z| = R$ ).  $\square$

Note that we can use the ratio test to determine  $R$  as well, as we will see in the next few examples.

### Example 3.40

(a)  $\sum n! z^n$ . By the ratio test, we have that  $\lim_{n \rightarrow \infty} \left| \frac{(n+1)! z^{n+1}}{n! z^n} \right| = \lim_{n \rightarrow \infty} (n+1) |z| = \infty$  for all  $z \neq 0$ . Hence the series diverges for all  $z \in \mathbb{C} \neq \{0\}$ , and we conclude that  $R = 0$ .

(b)  $\sum \frac{z^n}{n^n}$ . By Theorem 3.39, we have that:

$$R = \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{\frac{1}{n^n}}} = \frac{1}{\limsup_{n \rightarrow \infty} \frac{1}{n}} = \frac{1}{\lim_{n \rightarrow \infty} \frac{1}{n}} = \infty$$

(c)  $\sum \frac{z^n}{n!}$  also has  $R = \infty$  (as can be checked easily with the ratio test). For  $R = 1$ , the series is equal to  $e$ . As we will define later in Chapter 8, this series is equal to  $e^z$ .

(d)  $\sum \frac{z^n}{n^p}$  with  $p > 1$ . By Theorem 3.39, we have that:

$$R = \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{\frac{1}{n^p}}} = \frac{1}{\limsup_{n \rightarrow \infty} \left( \frac{1}{n} \right)^{\frac{p}{n}}} = \frac{1}{1^p} = 1$$

where for the second last equality we apply Theorem 3.20(c). Note that this series converges for all  $|z| \leq 1$  (although the above calculation does not show convergence on the boundary).

Before moving on, let us consider some further examples. Suppose  $a_n = 1$  for all  $n$ . Then, our power series is just  $\sum_{n=0}^{\infty} z^n$ , which is just the Geometric series. By Theorem 3.26, we have that the series converges if  $|z| < 1$  to  $\frac{1}{1-z}$ .

As another example, consider the series  $\sum_{n=1}^{\infty} \frac{1}{n} z^n$ . By the ratio test, we find that  $R = 1$ , as:

$$\limsup_{n \rightarrow \infty} \left| \frac{\frac{1}{n+1} z^{n+1}}{\frac{1}{n} z^n} \right| = \limsup_{n \rightarrow \infty} \left| \frac{n}{n+1} z \right| = |z| \limsup_{n \rightarrow \infty} \left| 1 - \frac{1}{n+1} \right| = |z|$$

So this series converges if  $|z| < 1$ , diverges if  $|z| > 1$ . What happens for  $|z| = 1$ ? At  $z = 1$ , we have that the series is just the standard harmonic series and diverges (by Theorem 3.28). At  $z = -1$ , we have an alternating series (a series whose terms are decreasing and tend to zero, and alternate in sign with





Figure 16: Visualization of the radius of convergence of the Geometric Series. The series converges in the shaded region, and diverges outside of it (and on the boundary).

each term), so we can apply the Alternating series test (below) to conclude that it converges. What about elsewhere on the circle? Let's look at  $z = i$ . We then have that the terms look like:

$$1 + \frac{i}{2} - \frac{1}{3} - \frac{i}{4} + \frac{1}{5} + \frac{i}{6} - \dots$$

we then have that the real and imaginary parts of the series are separately alternating series that decrease in magnitude; hence both parts are convergent, and the series as a whole is convergent at  $z = i$ . The same argument can be applied to conclude convergence of the series at  $z = -i$ . In fact, this series converges everywhere on the unit circle except at  $z = 1$ , which is a conclusion that follows from Theorem 3.44 (not covered in lecture, but feel free to refer to Rudin)!

#### Theorem 3.43: The Alternating Series Test

Let  $\{a_n\} \subset \mathbb{C}$  and suppose that:

- (a)  $|a_1| \geq |a_2| \geq |a_3| \dots$
- (b)  $a_{2m-1} \geq 0, a_{2m} \leq 1$  for  $m \in \mathbb{N}$
- (c)  $\lim_{n \rightarrow \infty} a_n = 0$ .

Then,  $\sum a_n$  converges.

#### Proof

Rudin establishes a partial summation formula (Theorem 3.41) and proves a more general theorem (Theorem 3.42) to prove this claim. However, an alternative proof in the case where  $\{a_n\}$  is real is left as homework (HW7).  $\square$

### 3.9 Absolute Convergence

#### Definition: Absolute Convergence

$\sum a_n$  is **absolutely convergent** if  $\sum |a_n|$  converges. Note that if  $\sum a_n$  is convergent but  $\sum |a_n|$  diverges, then  $\sum a_n$  is **conditionally convergent**. Note that for real series with strictly positive terms, absolute convergence and conditional convergence are equivalent. Also, note that the root and ratio tests test for absolute convergence, and hence do not yield any information for conditional convergence.

As an example, consider that  $\sum \frac{(-1)^n}{n}$  converges, but  $\sum \frac{1}{n}$  diverges, so  $\sum \frac{(-1)^n}{n}$  is conditionally convergent.

#### Theorem 3.45

If  $\sum a_n$  converges absolutely, then  $\sum a_n$  converges.

#### Proof

We have that:

$$\left| \sum_{j=m}^n a_j \right| \leq \sum_{j=m}^n |a_j| < \epsilon$$

For all  $n \geq m \geq N$  for some  $N$  by the fact that  $\sum |a_j|$  converges. Hence,  $\sum a_j$  is convergent by the Cauchy Criterion.  $\square$

For absolutely convergent series, we can freely change the order of the additions without affecting the value of the sum (as we will soon see). However, for series that are not absolutely convergent, this turns out to not be the case!

#### Definition 3.52: Rearrangements

Given a bijection  $K : \mathbb{N} \rightarrow \mathbb{N}$ , the series:

$$\sum_n a'_n = \sum_n a_{K(n)}$$

is called a rearrangement of  $\sum_n a_n$ .

#### Theorem 3.55

If  $\sum a_n$  is absolutely convergent, every rearrangement  $\sum a'_n$  converges to the same limit.

#### Proof

Let  $\{s'_n\}$  be the sequence of partial sums of the rearrangement  $\sum a'_n$ . Let  $\epsilon > 0$ . By the absolute convergence of the original series, there exists  $N \in \mathbb{N}$  such that for all  $n \geq m \geq N$ ,  $\sum_{j=m}^n |a_j| < \epsilon$ . Then, pick  $p$  such that  $\{1, 2, \dots, N\} \subset \{K(1), K(2), K(3), \dots, K(p)\}$ . Then, the summands  $a_1, a_2, \dots, a_N$  cancel out in  $s_n - s'_n$  for  $n \geq p$ , leaving only terms  $a_{K(j)}$  past  $a_N$ . Hence,  $|s_n - s'_n| < \epsilon$  for  $n \geq p \geq N$ , and we conclude that  $\sum a'_n$  converges to the same limit as  $\sum a_n$ .  $\square$

### Theorem 3.54: Riemann Rearrangement Theorem

If  $\sum a_n$  is a conditionally convergent series of real numbers, and  $-\infty \leq \alpha \leq \beta \leq \infty$ , then there is a rearrangement  $\sum a'_n$  such that  $\liminf_{n \rightarrow \infty} s'_n = \alpha$  and  $\limsup_{n \rightarrow \infty} s'_n = \beta$ . Taking  $\alpha = \beta$ , we have that for any real number, there exists a rearrangement that converges to it.

#### Proof

Not covered in lecture, see Rudin. □

For example, given  $\sum (-1)^n a_n$  with  $a_n \geq 0$  and  $\lim_{n \rightarrow \infty} a_n = 0$ , we can rearrange this series to converge to any point in  $\mathbb{R}$  that we like (for example,  $\pi$ ). The idea is to select positive terms from the series until we overshoot  $\pi$ , then choose a sequence of alternating negative/positive terms of decreasing magnitude until the  $\epsilon$  distance from  $\pi$  decreases to zero.

## 3.10 Addition and Multiplication of Series

### Theorem 3.47: Series Addition

Let  $\sum a_n = A$  and  $\sum b_n = B$ . Then,  $\sum (a_n + b_n) = A + B$  and  $\sum ca_n = cA$  for any fixed  $c \in \mathbb{C}$ .

#### Proof

Let  $A_n = \sum_{j=0}^n a_j$  and  $B_n = \sum_{j=0}^n b_j$ . Then,  $A_n + B_n = \sum_{j=0}^n a_j + b_j$  and since  $\lim_{n \rightarrow \infty} A_n = A$  and  $\lim_{n \rightarrow \infty} B_n = B$ , it follows that:

$$\lim_{n \rightarrow \infty} (A_n + B_n) = A + B.$$

For the second assertion, we have that  $\lim_{n \rightarrow \infty} cA_n = c \lim_{n \rightarrow \infty} A_n = cA$ . □

### Definition 3.48: Series Multiplication

Let  $\sum a_n$  and  $\sum b_n$  be two series. Then, the **product** of the two series is the series  $\sum c_n$  where:

$$c_n = \sum_{j=0}^n a_j b_{n-j} = \sum_{j=0}^n a_{n-j} b_j$$

Any student who has studied Fourier Series prior to this course will notice the similarity of the above definition to the convolution of two functions.

### Theorem 3.50

Suppose that  $\sum a_n$  converges absolutely and  $\sum b_n$  converges. Let  $\sum a_n = A$  and  $\sum b_n = B$ . Then,  $\sum c_n$  converges and  $\sum c_n = AB$ .

### Proof

Let  $A_n = \sum_{j=0}^n a_j$  and  $B_n = \sum_{j=0}^n b_j$ . Let  $\beta_n = B_n - B$  (note that  $\beta_n \rightarrow 0$ ). Now, we have that:

$$C_n = \sum_{k=0}^n c_k = \sum_{k=0}^n \sum_{j=0}^k a_j b_{k-j} = \sum_{j=0}^n \sum_{k=j}^n a_j a_{k-j} = \sum_{j=0}^n a_j \sum_{k=j}^n b_{k-j} = \sum_{j=0}^n a_j B_{n-j} = \sum_{j=0}^n a_j (B + \beta_{n-j})$$

From here, we split the sum and then we have that:

$$C_n = \sum_{j=0}^n a_j B + \sum_{j=0}^n a_j \beta_{n-j}$$

Defining  $\gamma_n = \sum_{j=0}^n a_j \beta_{n-j}$  and taking the limit of  $n \rightarrow \infty$ , we have:

$$\lim_{n \rightarrow \infty} C_n = C = \lim_{n \rightarrow \infty} \left( \sum_{j=0}^n a_j B + \gamma_n \right) = \lim_{n \rightarrow \infty} \sum_{j=0}^n a_j B + \lim_{n \rightarrow \infty} \gamma_n = AB + \lim_{n \rightarrow \infty} \gamma_n$$

So the claim is proven if  $\lim_{n \rightarrow \infty} \gamma_n = 0$ . Let  $\alpha = \sum |a_n| < \infty$  (by assumption of absolute convergence). Let  $\epsilon > 0$ . Then, there exists  $N \in \mathbb{N}$  such that  $|\beta_j| < \frac{\epsilon}{\alpha}$  for all  $j \geq N$  (as  $\beta_n \rightarrow 0$ ). Hence,

$$|\gamma_n| \leq \left| \sum_{j=0}^n a_{n-j} \beta_j \right| \leq \left| \sum_{j=0}^N a_{n-j} \beta_j \right| + \left| \sum_{j=N+1}^n a_{n-j} \beta_j \right| < \left| \sum_{j=0}^N a_{n-j} \beta_j \right| + \sum_{j=N+1}^n |a_{n-j}| \frac{\epsilon}{\alpha} \leq \left| \sum_{j=0}^N a_{n-j} \beta_j \right| + \epsilon$$

Letting  $n \rightarrow \infty$  with  $N$  fixed, we have the first term goes to 0 as  $a_n \rightarrow 0$  as  $n \rightarrow \infty$ . Hence, We have that:

$$\lim_{n \rightarrow \infty} |\gamma_n| < \epsilon$$

And as  $\epsilon$  is arbitrary,  $\lim_{n \rightarrow \infty} |\gamma_n| = 0$  and the claim follows.  $\square$

## 4 Continuity

### 4.1 Limits and Continuity

#### Definition 4.1: Limits

Let  $X, Y$  be metric spaces. Let  $E \subset X$ , and let  $f : E \mapsto Y$ . Let  $p \in X$  be a limit point of  $E$ . Then, we say that  $\lim_{x \rightarrow p} f(x) = q$  or  $f(x) \rightarrow q$  as  $x \rightarrow p$  if there exists  $q \in Y$  such that for all  $\epsilon > 0$ , there exists  $\delta > 0$  such that for all  $x \in E$  with  $0 < d_X(x, p) < \delta$  we have that  $d_Y(f(x), q) < \epsilon$ .



Figure 17: Visualization of the limit  $\lim_{x \rightarrow 1} f(x) = 1$  for  $f(x) = x$ . For any  $\epsilon > 0$ , we can take  $\delta = \epsilon$  and then we have that  $|f(x) - 1| < \epsilon$  if  $|x - 1| < \delta$ .

Note in the above definition that we do not care about  $f(p)$ , that is, the actual value of  $f$  at  $p$ . In particular, if  $p \notin E$ , then  $f(p)$  is not even necessarily defined. This distinction between the limit and the actual value of a function at a point becomes crucial later on when we want to define a derivative. Although we will discuss this in more detail in Chapter 5, the definition of a derivative of a function  $g$  at a point  $p \in \mathbb{R}$  involves the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that:

$$f(x) = \frac{g(x) - g(p)}{x - p}$$

Evidently, the domain of  $f$  does not contain the point  $p$ , but we are interested in the value of  $f$  in the limit of  $x \rightarrow p$  (which, if it exists, is the value of the derivative).



Figure 18: Visualization of the function  $f(x) = x$  for  $x \in \mathbb{R} \setminus \{1\}$ ,  $f(x) = 0$  for  $x = 1$ . In this case, we have that  $f(1) = 0$  but  $\lim_{x \rightarrow 1} f(x) = 1$ , demonstrating that the actual value of the function is irrelevant when defining the limit.

### Theorem 4.2

Let  $X, Y$  be metric spaces. Let  $E \subset X$  and  $f : E \mapsto Y$ . Suppose that for all sequences  $\{p_n\} \subset E$  with  $p_n \rightarrow p$  and  $p_n \neq p$ , we have that  $f(p_n) \rightarrow q \in Y$ . Then, this is equivalent to saying that  $\lim_{x \rightarrow p} f(x) = q$ .

### Proof

$\Rightarrow$  Suppose that  $\lim_{x \rightarrow p} f(x) = q$ , and let  $\{p_n\}$  be a sequence in  $E$  with  $p_n \rightarrow p$  and  $p_n \neq p$  for all  $n$ . We wish to show that  $f(p_n) \rightarrow q$ . Let  $\epsilon > 0$ . We show that there exists  $N \in \mathbb{N}$  such that  $d_Y(f(p_n), q) < \epsilon$  for all  $n \geq N$ . Since  $\lim_{x \rightarrow p} f(x) = q$ , there exists  $\delta > 0$  such that for all  $x \in E$  with  $d_X(p, x) < \delta$ ,  $d_Y(f(x), q) < \epsilon$ . Since we know that  $p_n \rightarrow p$ , there exists some  $N$  such that  $0 < d(p_n, p) < \delta$  for all  $n \geq N$ , so we have that  $d_Y(f(p_n), q) < \epsilon$  as required.

$\Leftarrow$  We show the contrapositive. Suppose that  $\lim_{x \rightarrow p} f(x) \neq q$ . We wish to find a sequence  $\{p_n\} \subset E$  with  $p_n \rightarrow p$  and  $p_n \neq p$  for all  $n$  such that  $f(p_n)$  does not converge to  $q$ . Since  $\lim_{x \rightarrow p} f(x) \neq q$ , then there exists  $\epsilon > 0$  such that for all  $\delta > 0$ , there exists  $x \in E$  such that  $0 < d_X(x, p) < \delta$  but  $d_Y(f(x), q) \geq \epsilon$ . For each  $\delta$  of the form  $\frac{1}{n}$ , let  $p_n \in E$  be the corresponding value of  $x$ . Then,  $p_n \rightarrow p$ ,  $p_n \neq p$  for all  $n$ , and  $f(p_n)$  does not converge to  $q$  as  $d(f(p_n), q) \geq \epsilon$  for all  $n$ .  $\square$

### Theorem 4.4

When  $Y = \mathbb{C}$  (i.e. the functions we consider are complex), then limits respect sums, differences, products, and functions. That is, let  $X$  is a metric space,  $E \subset X$ , and  $f, g : E \mapsto \mathbb{C}$  with  $p$  a limit point of  $E$ . If  $\lim_{x \rightarrow p} f(x) = q$  and  $\lim_{x \rightarrow p} g(x) = r$ , then  $\lim_{x \rightarrow p} (f + g)(x) = q + r$ . The same holds for subtraction, multiplication, and division (provided we do not divide by zero).

### Proof

By Theorem 4.2, these properties of limits follow from the analogous properties of sequences (Theorem 3.3).

### Definition 4.5: Continuity

Let  $X, Y$  be metric spaces, and  $E \subset X$ . Let  $p \in E$ , and define  $f : E \mapsto Y$ . We say that  $f$  is **continuous** at  $p$  if for all  $\epsilon > 0$ , there exists  $\delta > 0$  such that for all  $x \in E$  with  $d_X(x, p) < \delta$ , we have that  $d_Y(f(x), f(p)) < \epsilon$ . Equivalently,  $f(N_\delta^E(p)) \subset N_\epsilon^Y(f(p))$ . If  $f$  is continuous at  $p$  for all  $p \in E$ , we say that  $f$  is continuous.

Note that this definition of continuity is heavily reliant on the particular metric of  $X$  and  $Y$ ; in particular, there can be functions that are continuous for some choices of metric but not others.

Let us consider some examples of continuous functions (while thinking about different possible metric spaces).

First, let us take consider  $X = E = \mathbb{Z}$  and  $Y = \mathbb{R}$ . What functions  $f : E \rightarrow Y$  are continuous at  $p = 0$ ? The answer turns out to be all functions! To see this, fix  $n \in \mathbb{Z}$  and let  $\epsilon > 0$ . We then have that if  $|n - m| < \frac{1}{2}$ , then  $|f(n) - f(m)| < \epsilon$  as the only point  $m$  contained in  $N_{1/2}^{\mathbb{Z}}(n)$  is  $n$  itself (and hence  $|f(n) - f(m)| = |f(n) - f(n)| = 0$ ). This argument applies to every  $n \in \mathbb{Z}$  and hence all functions  $f : \mathbb{Z} \mapsto \mathbb{R}$  are continuous.

As further examples (that work for arbitrary metric spaces), If we have that  $f : X \mapsto X$ ,  $f(x) = x$ , we have that  $f$  is continuous (pick  $\delta = \epsilon$  in the definition of continuity). If we have that  $f : X \mapsto Y$ ,  $f(x) = c$

for some  $c \in Y$ , then  $f$  is also continuous (pick any  $\delta > 0$  in the definition).

Finally, the above definition doesn't make a distinction between limit points and isolated points. However, it turns out that according to the definition, if  $p \in E$  is isolated, then every function  $f$  with  $E$  as its domain is continuous. To see this, consider that for any  $\epsilon > 0$ , we can pick  $\delta > 0$  such that the only point  $x \in E$  for which  $d_X(x, p) < \delta$  is  $x = p$  (such a choice of  $\delta$  is possible as  $p$  is isolated). It then follows that  $d_Y(f(x), f(p)) = 0 < \epsilon$ .

We now consider a Theorem which gives us a familiar notion of continuity (that may have been encountered in first year calculus).

#### Theorem 4.6

Suppose that  $p$  is a limit point of  $E$  in Definition 4.5. Then,  $f$  is continuous at  $p$  if and only if  $\lim_{x \rightarrow p} f(x) = f(p)$ .

#### Proof

The claim immediately follows by comparing Definitions 4.1 and 4.5. □

#### Theorem 4.7

Let  $X, Y, Z$  be metric spaces, and  $E \subset X, F \subset Y$ . Let  $f : E \mapsto Y, g : F \mapsto Z$ , and suppose  $f(E) \subset F$ . Let  $p \in E$ . If  $f$  is continuous at  $p$  and  $g$  is continuous at  $f(p)$ , then  $g \circ f : E \mapsto Z$  is continuous at  $p$ .

#### Proof

Let  $\epsilon > 0$ . Since  $g$  is continuous at  $f(p)$ , there exists  $\gamma > 0$  such that  $d_Z(g(y), g(f(p))) < \epsilon$  if  $d_Y(y, f(p)) < \gamma$ . Since  $f$  is continuous at  $p$ , there exists  $\delta > 0$  such that  $d_Y(f(x), f(p)) < \gamma$  if  $d_X(x, p) < \delta$ . Hence, we have that  $d_Z(h(x), h(p)) = d_Z(g(f(x)), g(f(p))) < \epsilon$  if  $d_X(x, p) < \delta$  and  $x \in E$ . We conclude that  $h$  is continuous at  $p$ . □

## 4.2 Topological Characterization of Continuity

#### Theorem 4.8

Let  $X, Y$  be metric spaces, and  $f : X \mapsto Y$ . Then,  $f$  is continuous if and only if  $f^{-1}(V) \subset X$  is open for every open set  $V \subset Y$ .

#### Proof

$\Rightarrow$  Suppose  $f$  is continuous, and  $V \subset Y$  is open. Let  $p \in f^{-1}(V)$ , so  $f(p) \in V$ .  $V$  is open, so it follows that  $f(p)$  is an interior point of  $V$ . So, there exists  $r > 0$  such that  $N_r^Y(f(p)) \subset V$ . Next,  $f$  is continuous, so there exists  $\delta > 0$  such that for all  $x \in X$  with  $d_X(x, p) < \delta$ ,  $d_Y(f(x), f(p)) < r$ . Hence, we obtain that  $f(x) \in N_r^Y(f(p)) \subset V$ . In particular,  $N_\delta^X(p) \subset f^{-1}(V)$ , so  $p$  is an interior point of  $f^{-1}(V)$ . Hence every point of  $f^{-1}(V)$  is an interior point, and  $f^{-1}(V)$  is open.

$\Leftarrow$  Suppose  $f^{-1}(V)$  is open for every open set  $V \subset Y$ . Let  $p \in X$  and  $\epsilon > 0$ . Let  $V = N_\epsilon^Y(f(p))$  which is open, so by assumption  $f^{-1}(V)$  is open.  $p \in f^{-1}(V)$ , so  $p$  is an interior point. Hence, there exists  $\delta > 0$  such that  $N_\delta^X(p) \subset f^{-1}(V)$ . In other words, if  $d_X(x, p) < \delta$ , then  $f(x) \in V$ , so  $d_Y(f(x), f(p)) < \epsilon$ .  $f$  is then continuous by definition. □

### Corollary

$f : X \mapsto Y$  is continuous if and only if  $f^{-1}(F) \subset X$  is closed for every closed set  $F \subset Y$ .

### Proof

Let  $V = F^c$  with  $V$  open. Then, the above statement is equivalent to saying that a function  $f : X \mapsto Y$  is continuous if and only if  $f^{-1}(V^c) = \left(f^{-1}(V)\right)^c$  is closed for every open set  $V \subset Y$ . This is equivalent to the statement that  $f^{-1}(V)$  is open for every open set  $V \subset Y$ , and the claim follows by the previous theorem.  $\square$

Note that the above topological characterization says properties of sets are preserved under taking the preimage; it does not say anything about preserving properties under the image. That is to say, images of open/closed sets are not necessarily open/closed.

### Example

Consider  $X = \mathbb{R}^+ = (0, \infty)$  and  $Y = \mathbb{R}$ . Then, the function:

$$\begin{array}{ccc} f & : & X \longrightarrow Y \\ & & x \longmapsto \frac{1}{x} \end{array}$$

is continuous. Then defining  $A = [1, \infty)$  we have that  $A$  is closed, but  $f(A) = (0, 1]$  is not closed.

### Theorem 4.9

Let  $f : X \mapsto \mathbb{C}$  and  $g : X \mapsto \mathbb{C}$  be continuous functions. Then,  $f + g$ ,  $fg$  are continuous, and  $\frac{f}{g}$  is continuous if  $g(x) \neq 0$ .

### Proof

At isolated points there is nothing to prove (as any choice of function that is defined at an isolated  $p$  will be continuous there). For limit points, the claim follows from Theorems 4.4 and 4.6.  $\square$

### Theorem 4.10

(a) Let  $f_1, f_2, \dots, f_k : X \mapsto \mathbb{R}$ , and define  $\mathbf{f} : X \mapsto \mathbb{R}^k$  by:

$$\mathbf{f}(x) = (f_1(x), f_2(x), \dots, f_k(x))$$

Then,  $\mathbf{f}$  is continuous if and only if every  $f_i$  is continuous.

(b) If  $\mathbf{f} = (f_1, \dots, f_k) : X \mapsto \mathbb{R}^k$  and  $\mathbf{g} = (g_1, \dots, g_k) : X \mapsto \mathbb{R}^k$  are continuous, then the functions:

$$\mathbf{f} + \mathbf{g} = (f_1 + g_1, \dots, f_k + g_k)$$

$$\mathbf{f} \cdot \mathbf{g} = f_1 g_1 + \dots + f_k g_k$$

are continuous.



### Proof

- (a)  $\Rightarrow$  Suppose  $\mathbf{f}$  is continuous. Then, let  $\epsilon > 0$ . For each  $p \in X$ , there exists  $\delta > 0$  such that  $d_X(x, p) < \delta$  implies  $|\mathbf{f}(x) - \mathbf{f}(p)| < \epsilon$ . We then observe that for any  $i \in \{1, \dots, k\}$ :

$$|\mathbf{f}(x) - \mathbf{f}(p)| = \left( \sum_{j=1}^k |f_j(x) - f_j(p)|^2 \right)^{1/2} \geq |f_i(x) - f_i(p)|$$

Where the inequality follows by just keeping one term from the sum of non-negative terms. So for any  $d_X(x, p) < \delta$  we therefore have that  $|f_i(x) - f_i(p)| < \epsilon$ , and hence each  $f_i$  is continuous.

$\Leftarrow$  Suppose each of  $f_1, \dots, f_k$  is continuous. Then let  $\frac{\epsilon}{\sqrt{k}} > 0$ . For each  $p \in X$ , there exists  $\delta_i$  such that  $d_X(x, p) < \delta_i$  implies that  $|f_i(x) - f_i(p)| < \frac{\epsilon}{\sqrt{k}}$ . Take  $\delta = \min \{\delta_1, \dots, \delta_k\}$ . Then, if  $d_X(x, p) < \delta$ , we have that:

$$|\mathbf{f}(x) - \mathbf{f}(p)| = \left( \sum_{j=1}^k |f_j(x) - f_j(p)|^2 \right)^{1/2} < \left( \sum_{j=1}^k \left( \frac{\epsilon}{\sqrt{k}} \right)^2 \right)^{1/2} = \epsilon$$

So it follows that  $\mathbf{f}$  is continuous.

- (b) The claim follows from (a) and Theorem 4.9.  $\square$

### Example 4.11

We will now explore some interesting examples of continuous functions.

- (a) For each index  $i = 1, \dots, k$ , define  $\phi_i : \mathbb{R}^k \mapsto \mathbb{R}$  by  $\phi_i(\mathbf{x}) = x_i$ . Then,  $\phi_i$  is continuous.

*Proof.* Let  $\epsilon > 0$ . Then, for  $\mathbf{p} \in \mathbb{R}^k$ , if  $|\mathbf{x} - \mathbf{p}| < \delta$  with  $\delta = \epsilon$ , we have that:

$$\epsilon > |\mathbf{x} - \mathbf{p}| = \left( \sum_{j=1}^k |x_j - p_j|^2 \right)^{1/2} \geq (|x_i - p_i|^2)^{1/2} = |x_i - p_i|$$

So for  $|\mathbf{x} - \mathbf{p}| < \delta$ , we have that  $|\phi_i(\mathbf{x}) - \phi_i(\mathbf{p})| < \epsilon$ . We conclude that  $\phi_i$  is continuous.  $\square$

We could also use the topological characterization of continuity to prove this claim. If  $V \subset \mathbb{R}$  is open, then  $\mathbb{R} \times \mathbb{R} \times \dots \times V \times \mathbb{R} \times \dots \times \mathbb{R}$  is open, showing again that  $\phi_i$  is continuous (a much easier proof)!

- (b) Let  $f : \mathbb{R}^k \mapsto \mathbb{R}$  be given by  $f(\mathbf{x}) = x_1^{n_1} x_2^{n_2} \dots x_k^{n_k}$  with  $n_i \in \mathbb{N} \cup \{0\}$ .  $f$  is continuous, and hence so is any polynomial  $P : \mathbb{R}^k \mapsto \mathbb{R}$ .
- (c) Rational functions  $P/Q$  where  $P, Q$  are polynomials are continuous everywhere except where  $Q$  is zero.
- (d)  $f : \mathbb{R}^k \mapsto \mathbb{R}$  given by  $f(\mathbf{x}) = |\mathbf{x}|$  is continuous.
- (e) Suppose  $f : X \mapsto \mathbb{R}^k$  is continuous. Then so is  $g : X \mapsto \mathbb{R}^k$  with  $g(x) = |f(x)|$ .

### 4.3 Continuity and Compactness

#### Theorem 4.14

Let  $f : X \mapsto Y$  be continuous, and suppose  $X$  is compact. Then, the image  $f(X)$  is compact.

#### Proof

Let  $\{V_\alpha\}$  be an open cover of  $f(X)$ . For each  $\alpha$ , Let  $O_\alpha = f^{-1}(V_\alpha)$ . Then,  $\{O_\alpha\}$  is an open cover of  $X$ . to see this, we recognize that each  $O_\alpha$  is open due to the continuity of  $f$  (Theorem 4.8) and each  $p \in X$  satisfies  $V_\beta$  for some  $\beta$ , so  $p \in O_\beta$ .  $X$  is compact, so there exists a finite subcover  $\{O_{\alpha_1}, \dots, O_{\alpha_n}\}$ . But it then follows that  $\{V_{\alpha_1}, \dots, V_{\alpha_n}\}$  is a finite subcover of  $f(X)$ . Hence  $f(X)$  is compact.  $\square$

We previously discussed how an image of a closed set (under a continuous function) does not have to be closed. However, the above Theorem shows that the image of a compact set (under a continuous function) must be compact.

Taking a slight tangent, a question of interest may be to consider the topology of the extended real numbers; namely,  $\mathbb{R} \cup \{\infty, -\infty\}$  (where  $-\infty < x < \infty$  for all  $x \in \mathbb{R}$ ). In particular, a natural question to ask is whether  $[1, \infty) \cup \{\infty\}$  would be closed/compact.

We start then by defining a reasonable metric on  $X = \mathbb{R} \cup \{\infty, -\infty\}$ . A “reasonable” definition of a metric will satisfy the requirement that  $[1, \infty) \cup \{\infty\}$  is compact or not. We then define the metric:

$$d(x, y) = |\arctan(x) - \arctan(y)|$$

where  $\arctan(\infty) = \frac{\pi}{2}$  and  $\arctan(-\infty) = -\frac{\pi}{2}$ . We then have that  $(X, d)$  is a metric space (check!)

Note that the extended reals have been made into a metric space here, but it is no longer a field; in particular, we have that  $(-\infty + \infty) + \infty = 0 + \infty = \infty$  but  $-\infty + (\infty + \infty) = -\infty + \infty = 0$  so associativity no longer holds.

With a reasonable metric defined on the extended reals, let us now return to our previous example in Section 4.2 where we considered the function  $f(x) = \frac{1}{x}$ . We now extend the domain of the function such that:

$$\begin{aligned} f : X \setminus \{0\} &\longrightarrow \mathbb{R} \\ x &\longmapsto \frac{1}{x} \end{aligned}$$

With  $f(\infty) = f(-\infty) = 0$ . We leave it as an exercise to verify that  $f$  is continuous on  $X \setminus \{0\}$  using the continuity of the arctan function.

Now, consider  $A = [1, \infty) \subset X$ .  $A$  is not closed, as  $\infty$  is a limit point of  $A$  (this can be verified by checking that for all  $\epsilon > 0$ , there exists  $x \in [1, \infty)$  such that  $|\arctan x - \arctan \infty| = \frac{\pi}{2} - \arctan x < \epsilon$ ). However, we do have that  $\overline{A} = [1, \infty]$  is closed and compact, and that  $f(\overline{A}) = [0, 1]$  is compact as Theorem 4.14 states. In a sense, we have “compactified” the reals through our construction, as  $X$  is compact while  $\mathbb{R}$  was not. We also have that  $X$  is bounded (unlike  $\mathbb{R}$ ), as all distances between points are bounded by at most  $\pi$ . In particular,  $\text{diam } X = \pi$ .

#### Definition 4.13: Bounded functions

Let  $X$  be a metric space, and  $E \subset X$ . We say that  $f : E \mapsto Y$  is **bounded** if  $f(E)$  is a bounded set. In particular, if  $Y = \mathbb{R}^k$ , then  $f$  is bounded if and only if there exists  $M \in \mathbb{R}$  such that  $|f(x)| \leq M$  for all  $x \in E$ .

**Theorem 4.15**

Let  $X$  be a compact metric space, and  $f : X \mapsto Y$  be continuous. Then,  $f(X)$  is bounded.

**Proof**

By Theorem 4.14, we have that  $f(X)$  is compact.  $f(X)$  is therefore bounded (see discussion preceding Theorem 2.41).  $\square$

An interesting special case in the above theorem to think about is when  $Y = \mathbb{R}$ ; this yields the Extreme Value Theorem, below.

**Theorem 4.16: Extreme Value Theorem**

Let  $X$  be a compact metric space, and let  $f : X \mapsto \mathbb{R}$  be continuous. Then, there exist  $p, q \in X$  such that  $f(p) \leq f(x) \leq f(q)$  for all  $x \in X$ . In other words,  $f$  attains its minimum (infimum) and maximum (supremum) on  $X$ .

**Proof**

By Theorems 4.15 and 4.16,  $f(X)$  is closed and bounded. In particular,  $m = \inf f(X)$  and  $M = \sup f(X)$  exist (as the set is bounded), and  $m, M \in f(X)$  as  $f(X)$  contains all of its limit points. Hence, there exist  $p, q \in X$  such that  $f(p) = m, f(q) = M$ .  $\square$

We consider some examples which demonstrate the necessity of the compactness of the domain. Let  $E = (0, 1) \subset \mathbb{R}$  and consider  $f : (0, 1) \mapsto \mathbb{R}$ . Let  $f(x) = \frac{1}{x^2}$ .  $(0, 1)$  is bounded but not closed, and we observe that  $f((0, 1)) = [1, \infty)$  which is unbounded (and hence  $f$  does not attain a maximum). Instead let  $f(x) = x$ . Then, we have that  $f((0, 1)) = (0, 1)$  which is bounded, but  $f$  does not attain a maximum/minimum on its domain.

**Theorem 4.17**

Let  $X$  be a compact metric space, and  $f : X \mapsto Y$  be continuous and a bijection. Define  $f^{-1} : Y \mapsto X$  by  $f^{-1}(y) = x$  if  $f(x) = y$  (this is well-defined due to the bijectivity of  $f$ ). Then,  $f^{-1}$  is continuous.

**Proof**

By Theorem 4.8, it suffices to show that for every open set  $V \subset X$ ,  $(f^{-1})^{-1}(V) = f(V)$  is open. By the openness of  $V$ ,  $V^c$  is closed, and hence compact (as closed subsets of compact sets are compact by Theorem 2.35). So,  $f(V^c) \subset Y$  is compact by Theorem 4.14. Therefore,  $f(V^c) = (f(V))^c$  is closed (as compact sets are closed by Theorem 2.34). Therefore,  $((f(V))^c)^c = f(V)$  is open as desired. Hence  $f^{-1}$  is continuous.  $\square$

### Example 4.21

This example shows the necessity of compactness of  $X$  in Theorem 4.17. Let  $S = [0, 2\pi)$  which is not closed and hence not compact (Heine Borel/Theorem 2.41). Let  $Y = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$ . Define  $f : X \mapsto Y$  where  $f(\theta) = (\cos \theta, \sin \theta)$ .  $f$  is continuous and a bijection (check!) but  $f^{-1}$  is not continuous. To see this is the case, consider first that  $[0, \frac{\pi}{2}) \subset X$  is open in  $X$ . At everywhere except 0 it should be evident that all points of  $[0, \frac{\pi}{2})$  are interior to  $X$ , and at 0, we have that  $N_{1/2}(0) \subset [0, \frac{\pi}{2})$  showing that 0 is also an interior point. But,  $(f^{-1})^{-1}([0, \frac{\pi}{2})) = f([0, \frac{\pi}{2}))$  is not open, because not every point is an interior point; namely,  $(1, 0) = f(0) \in f([0, \frac{\pi}{2}))$  is not an interior point. So,  $f^{-1}$  is not continuous.



Figure 19: Visual of the map  $f$  which maps a point  $\theta \in [0, 2\pi)$  to the point  $(\cos \theta, \sin \theta)$  on the unit circle. Pictured is the map from  $[0, \frac{\pi}{2})$  to  $f([0, \frac{\pi}{2}))$ , the former which is open in  $[0, 2\pi)$ , and the latter which is not open in  $Y = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$ .

## 4.4 Uniform Continuity, Connectedness, and IVT

### Definition 4.18: Uniform Continuity

Let  $X, Y$  be metric spaces and  $f : X \mapsto Y$ . Then,  $f$  is **uniformly continuous** if for all  $\epsilon > 0$ , there exists  $\delta > 0$  such that for all  $p, q \in X$  with  $d_X(p, q) < \delta$ , we have that  $d_Y(f(p), f(q)) < \epsilon$ . It is clear from the definition that every uniformly continuous function is continuous.

Note that this definition is very similar to the definition of continuity, but with a difference in the quantifiers. With continuity, for each  $\epsilon$  there is a  $\delta$  that we can find for a given point  $p \in X$ . For uniform continuity, we have that for each  $\epsilon$ , there exists a  $\delta$  that works uniformly for all  $p$ .

We now consider some examples to see the difference concretely. Take  $X = (0, 1)$ ,  $Y = \mathbb{R}$ , and  $f(x) = \frac{1}{x}$ . We then have that  $f$  is continuous, but not uniformly continuous, showing that in general continuity does not imply uniform continuity.

Next, consider  $X = Y = \mathbb{R}$  and  $f(x) = \sin x$ . Then,  $f$  is uniformly continuous. Given any  $\epsilon$ , we can pick  $\delta = \epsilon$  and this proof will work. The “calculus-inspired” proof would use that  $\frac{d}{dx} \sin x = \cos x$  and  $|\cos x| \leq 1$ , so we can always pick  $\delta = \epsilon$  with the MVT. However, we have yet to define the derivative or prove the mean value theorem, so an alternate approach would be to invoke trigonometric identities.

Finally, let  $X = [0, 10]$ ,  $Y = \mathbb{R}$ , and  $f(x) = x^2$ . Then,  $f$  is uniformly continuous. However, if  $X = Y = \mathbb{R}$  and  $f(x) = x^2$ , then  $f$  is not uniformly continuous. The uniform continuity of  $f$  depends on the domain; in particular,  $[0, 10]$  is closed/compact, while  $\mathbb{R}$  is not compact. This motivates our next theorem.

**Theorem 4.19**

Let  $X, Y$  be metric spaces with  $X$  compact. Let  $f : X \mapsto Y$  be continuous. Then,  $f$  is uniformly continuous.

**Proof**

Let  $\epsilon > 0$ . By the continuity of  $f$ , for each  $p \in X$ , there exists  $\delta_p > 0$  such that for all  $q \in X$  with  $d_X(p, q) < \delta_p$ , we have that  $d_Y(f(p), f(q)) < \frac{\epsilon}{2}$ . Define  $U_p = N_{\delta_p/2}(p)$ . We then have that  $\{U_p\}_{p \in X}$  is an open cover for  $X$ . By the compactness of  $X$ , it has a finite subcover. Let  $\{U_{p_1}, \dots, U_{p_n}\}$  be a finite subcover. Let  $\delta = \frac{1}{2} \min \{\delta_{p_1}, \dots, \delta_{p_n}\} > 0$  (note the importance of taking the minimum over a *finite* number of  $\delta_{p_i}$ s; if we took an infimum over an infinite number, it could be possible that the infimum could be zero). Then, take  $p, q \in X$  with  $d_X(p, q) < \delta$ . Then,  $p \in U_{p_i}$  for some  $i$ . So,  $d_X(p, p_i) < \frac{\delta_{p_i}}{2}$ , and then by the triangle inequality we have that:

$$d_X(q, p_i) \leq d_X(q, p) + d_X(p, p_i) < \delta + \frac{\delta_{p_i}}{2} \leq \frac{\delta_{p_i}}{2} + \frac{\delta_{p_i}}{2} = \delta_{p_i}$$

Therefore, we have that:

$$d_Y(f(p), f(q)) \leq d_Y(f(p), f(p_i)) + d_Y(f(p_i), f(q)) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

And we conclude that  $f$  is uniformly continuous. □

**Theorem 4.22**

Let  $X, Y$  be metric spaces, and let  $f : X \mapsto Y$  be continuous. If  $E \subset X$  is connected (see Definition 2.45) then its image  $f(E)$  is also connected.

**Proof**

Suppose for the sake of contradiction that we can write  $f(E) = A \cup B$  with  $A, B \neq \emptyset$  and  $A \cap \bar{B} = \bar{A} \cap B = \emptyset$  (i.e.  $f(E)$  is not connected). Then, define  $G = f^{-1}(A) \cap E$  and  $H = f^{-1}(B) \cap E$ . Then,  $E = G \cup H$ , with  $G \neq \emptyset$ ,  $H \neq \emptyset$ , and  $G \cap H = \emptyset$ . We wish to show that  $\bar{G} \cap H = G \cap \bar{H} = \emptyset$  as this will contradict the connectedness of  $E$ . Since  $A \subset \bar{A}$ , we have that  $G \subset f^{-1}(A) \subset f^{-1}(\bar{A})$ .  $f$  is continuous, so by Theorem 4.8, we have that  $f^{-1}(\bar{A})$  is closed. Hence by Theorem 2.27 we have that  $\bar{G} \subset f^{-1}(\bar{A})$ . Since  $f(H) = B$  and  $\bar{A} \cap B = \emptyset$ , we have that:

$$f(\bar{G}) \cap f(H) = \emptyset$$

And therefore  $\bar{G} \cap H = \emptyset$ . By an identical argument,  $G \cap \bar{H} = \emptyset$ . Hence,  $E = G \cup H$  is not connected, which is a contradiction. □

**Theorem 4.23: Intermediate Value Theorem**

Let  $f : [a, b] \mapsto \mathbb{R}$  be continuous and  $f(a) < f(b)$ . Then, for all  $\alpha \in (f(a), f(b))$ , there exists  $c \in (a, b)$  with  $f(c) = \alpha$ .

### Proof

$[a, b]$  is connected, so by Theorem 4.22,  $f([a, b])$  is connected. Hence by 2.47, it contains every point between  $f(a)$  and  $f(b)$ .  $\square$



Figure 20: Visualization of functions for which the IVT applies (left) and for which it doesn't (right).

As a closing remark for this chapter, consider that continuity over a closed interval implies the intermediate value theorem. Does this then imply that a connected space with the intermediate value theorem implies continuity? The answer turns out to be no. Consider discontinuities other than jumps (IVT only tells us a function does not “jump” on the domain of interest), that is, a discontinuity that arises from the non-existence of a limit. As an example, consider:

$$f : \mathbb{R} \longrightarrow \mathbb{R} \\ x \longmapsto \begin{cases} \frac{1}{x} & x \neq 0 \\ 0 & x = 0 \end{cases}$$

This function is not continuous at  $x = 0$  (the limit does not even exist there!) but satisfies the IVT.

## 4.5 Topological Spaces

Up until this point, we have been discussing metric spaces  $(X, d)$ , where  $U \subset X$  is open if every point of  $U$  is an interior point according to the metric  $d$ . In this picture,  $\emptyset$  is open,  $X$  is open, and if  $\{U_\alpha\}$  are open, then  $\bigcup_\alpha U_\alpha$  (arbitrary unions) and  $\bigcap_{i=1}^n U_{\alpha_i}$  (finite intersections) are open. Although this metric space picture is the one we have been using (and will be using going forwards in the course), it could be interesting to take a temporary tangent, and consider a generalization of these notions to topological spaces:

### Definition: Topological Spaces

A pair  $(X, \tau)$  is a **topological space** if  $\tau \subset \mathcal{P}(X)$  such that  $\emptyset \in \tau$ ,  $X \in \tau$ , and if  $\{U_\alpha\} \subset \tau$ , then  $\bigcup_\alpha U_\alpha \in \tau$  and  $\bigcap_{i=1}^n U_{\alpha_i} \in \tau$ . In this definition,  $\tau$  is the set of “open sets” of  $X$ .

In metric spaces, we defined openness in terms of a metric, but in topological spaces, we throw away the metric.

If  $(X, \tau)$  and  $(Y, \rho)$  are topological spaces, how do we define a continuous function  $f : X \mapsto Y$ ? Evidently, the  $\epsilon - \delta$  definition no longer makes sense (in the absence of a metric), but our topological characterization of continuity still holds:

### Definition: Continuity on Topological Spaces

Let  $(X, \tau)$  and  $(Y, \rho)$  be topological spaces. Then,  $f : X \mapsto Y$  is **continuous** if for every  $U \in \rho$ ,  $f^{-1}(U) \in \tau$ .

A natural question that arises given a topological space is whether it is possible or not to find a metric that gives us this topology.

### Definition: Metrizable

Let  $(X, \tau)$  be a topological space. If there exists a metric  $d : X \times X \mapsto \mathbb{R}$  such that every set  $U \in \tau$  is open with respect to the metric  $d$ , then we say that  $X$  is **metrizable**.

However, not all topological spaces are metrizable. For example, take  $(X, \tau)$  with  $\tau = \{\emptyset, X\}$ . If  $X$  has more than 1 element, then there is no metric that gives rise to this trivial topology.

*Proof.* We show the proof for the case of 2 elements (the proof for any finite  $|X|$  follows analogously, and the argument for the infinite case is left as an exercise). Let  $X = \{a, b\}$  and  $\tau = \{\emptyset, X\}$ . If  $d$  is a metric on  $X$ , then  $d(a, b) = r > 0$ . Then, let  $0 < s < r$ . Then,  $N_s(a)$  has to be open, as does  $N_s(b) = \{b\}$ . The set of open sets of  $X$  under  $d$  is therefore  $\{X, \emptyset, \{a\}, \{b\}\} \neq \tau$  and hence  $(X, \tau)$  is not metrizable.  $\square$

The original definition we had of convergent sequences (and Cauchy sequences) also does not generalize well to topological spaces (as both invoke the notion of a metric). We can define convergence in topological spaces (in a way that does not explicitly use a metric) as follows:

### Definition: Convergence in Topological Spaces

Let  $(X, \tau)$  be a topological space, and suppose  $\{x_n\} \subset X$ . We say that  $x_n$  **converges** to  $x$  if for every set  $U \in \tau$  such that  $x \in U$ , there exists  $N$  such that  $x_n \in U$  for all  $n \geq N$ .

Note that it is impossible to define the notion of a Cauchy sequence without a metric (it cannot be defined solely in terms of open sets). Convergence is topological, but the metric matters for Cauchy.

Recall that the compactness of a set  $K$  was defined in terms of every cover having a finite subcover. The sequential compactness of a set  $K$  can be defined by saying that every sequence in  $K$  has a convergent subsequence. These definitions have no explicit reference to a metric given our above two definitions of open sets/convergence, so the definition of compactness holds in the same way in the picture of topological spaces. Note that in metric spaces, compactness and sequentially compactness are equivalent, but they can be different in topological spaces (this is another fact that tells us that there exist topological spaces that are not metrizable)!

So reviewing what we can define meaningfully in the picture of topological spaces, we see that we can define continuity, convergence, compactness, and sequential compactness; unfortunately, Cauchy sequences are left on the cutting room floor.

## 5 Differentiation

### 5.1 Derivatives

#### Definition 5.1: Derivatives

Let  $f : [a, b] \mapsto \mathbb{R}$ , and  $x \in [a, b]$ . We then define the **derivative** of  $f$  at  $x$  as:

$$f'(x) = \lim_{t \rightarrow x} \frac{f(t) - f(x)}{t - x}$$

If the limit exists. Alternative notations for the derivative are given by:

$$\frac{\partial f}{\partial x}(x) \text{ or } \frac{d}{dx} f(x) \text{ or } \frac{d}{dy} f(x) \Big|_{y=x}$$

As an interpretation of the derivative, take  $[a, b]$  to be a metric space, with  $x$  a limit point of  $[a, b] \setminus \{x\}$ . Then,  $g(t) = \frac{f(t) - f(x)}{t - x}$  is a function from  $[a, b] \setminus \{x\} \mapsto \mathbb{R}$ . If  $x \in (a, b)$ , then the above definition of the derivative agrees with the definition of  $f'(x)$  from first year calculus. If  $x = a$  or  $x = b$ , then the above definition agrees with the definition of the one-sided derivative from first year calculus. Note that we will not discuss in this class cases where the domain gets more complicated (i.e. not just closed intervals of  $\mathbb{R}$ ).

#### Theorem 5.2

Let  $f : [a, b] \mapsto \mathbb{R}$ , let  $x \in [a, b]$ , and suppose  $f'(x)$  exists. Then,  $f$  is continuous at  $x$ .

#### Proof

For  $t \neq x$ , we can write:

$$f(t) = f(x) + (f(t) - f(x)) = f(x) + \frac{f(t) - f(x)}{t - x}(t - x)$$

Taking the limit of  $t \rightarrow x$ , we then have that:

$$\lim_{t \rightarrow x} f(t) = \lim_{t \rightarrow x} \left( f(x) + \frac{f(t) - f(x)}{t - x}(t - x) \right) = \lim_{t \rightarrow x} f(x) + \lim_{t \rightarrow x} \frac{f(t) - f(x)}{t - x} \lim_{t \rightarrow x} (t - x)$$

Where in the last line we invoke Theorem 4.4. Evaluating the limits on the RHS by using the existence of the derivative of  $f$  at  $x$ , we have

$$\lim_{t \rightarrow x} f(t) = f(x) + f'(x) \cdot (0) = f(x)$$

So we conclude that  $f$  is continuous at  $x$  by Theorem 4.6. □

The interpretation is that differentiability at  $x \in (a, b)$  implies continuity of  $f$  at  $x$ , and the left/right differentiability of  $f$  at  $a/b$  implies the left/right continuity of  $f$  at  $a/b$ . We have wrapped the proof of all these cases into one!

Note that the converse of the above theorem is not true. As a simple example, take  $f(x) = |x|$  on  $[-1, 1]$ , which is continuous at  $x = 0$  (it can be verified that  $\lim_{x \rightarrow 0} f(x) = f(0) = 0$ ) but is not differentiable there (the left/right handed limits of the difference quotient do not agree and hence the



derivative does not exist). In Chapter 7, we will construct a function that is continuous everywhere and differentiable nowhere!

NWe will now proceed to prove a series of theorems that have been seen in first year, but using our new/rigorous definitions.

### Theorem 5.3: Sum, Product, and Quotient Rules

Let  $f, g : [a, b] \mapsto \mathbb{R}$ . Let  $x \in [a, b]$  and suppose  $f$  and  $g$  are differentiable at  $x$ . Then,  $f + g$ ,  $f - g$ ,  $f \cdot g$  are differentiable at  $x$ , and so is  $\frac{f}{g}$  provided  $g(x) \neq 0$ . Furthermore:

$$(a) (f + g)'(x) = f'(x) + g'(x)$$

$$(b) (fg)'(x) = f'(x)g(x) + f(x)g'(x)$$

$$(c) \left(\frac{f}{g}\right)'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$$

### Proof

(a) Follows immediately from the additive property of limits (Theorem 4.4).

(b) Let  $h = fg$ . We then have that:

$$h(t) - h(x) = f(t)[g(t) - g(x)] + g(x)[f(t) - f(x)]$$

For  $t \neq x$ , we can divide both sides by  $t - x$  to obtain:

$$\frac{h(t) - h(x)}{t - x} = f(t)\frac{g(t) - g(x)}{t - x} + g(x)\frac{f(t) - f(x)}{t - x}$$

Taking the limit of  $t \rightarrow x$  on both sides, we obtain:

$$h'(x) = f(x)g'(x) + f'(x)g(x)$$

as desired.

(c) Let  $h(t) = \frac{f(t)}{g(t)}$ . Then:

$$\begin{aligned} h(t) - h(x) &= \frac{f(t)}{g(t)} - \frac{f(x)}{g(x)} \\ &= \frac{1}{g(t)g(x)} (f(t)g(x) - g(t)f(x)) \\ &= \frac{1}{g(t)g(x)} [g(x)(f(t) - f(x)) - f(x)(g(t) - g(x))] \end{aligned}$$

For  $t \neq x$ , we can divide both sides by  $t - x$  to get:

$$\frac{h(t) - h(x)}{t - x} = \frac{1}{g(t)g(x)} \left[ g(t)\frac{f(t) - f(x)}{t - x} - f(x)\frac{g(t) - g(x)}{t - x} \right]$$

Taking the limit as  $t \rightarrow x$  on both sides, we obtain the desired expression.  $\square$

As an exercise, one can prove by induction (applying 5.3(b)) that  $(f_1 f_2 f_3 \dots f_n)'(x)$  (where  $f_i : [a, b] \mapsto \mathbb{R}$  and each  $f_i'(x)$  exists) is given by:

$$f_1'(x)f_2(x) \dots f_n(x) + \dots + f_1(x)f_2(x) \dots f_n'(x).$$

Note that as a corollary of this, we get that if  $f(x) = x^n$ , then  $f'(x) = nx^{n-1}$  and we hence recover the familiar power rule from first year calculus!

### Theorem 5.5: Chain Rule

Let  $f : [a, b] \mapsto \mathbb{R}$ ,  $x \in [a, b]$ , and suppose  $f$  is differentiable at  $x$ . Suppose furthermore that  $f([a, b])$  is contained in some interval  $I$ . Let  $g : I \mapsto \mathbb{R}$  and suppose  $g$  is differentiable at  $f(x)$ . Then,  $g \circ f : [a, b] \mapsto \mathbb{R}$  is differentiable at  $x$ , and furthermore:

$$(g \circ f)'(x) = g'(f(x))f'(x)$$

### Proof

Define  $h(t) = g \circ f(t)$  for  $a \leq t \leq b$ ,  $t \neq x$ . We can then write:

$$f(t) - f(x) = (t - x) [f'(x) + u(t)]$$

For a function  $u(t)$  with  $\lim_{t \rightarrow x} u(t) = 0$ . Now defining  $y = f(x)$ , we write:

$$g(s) - g(y) = (s - y) [g'(y) + r(s)]$$

For a function  $r(s)$  with  $\lim_{s \rightarrow y} r(s) = 0$ . Hence, we have that:

$$\begin{aligned} h(t) - h(x) &= g(f(t)) - g(f(x)) \\ &= (f(t) - f(x)) (g'(y) + r(s)) \\ &= (t - x) [f'(x) + u(t)] (g'(y) + r(s)) \end{aligned}$$

Dividing both sides by  $t - x$ , we obtain:

$$\frac{h(t) - h(x)}{t - x} = [f'(x) + u(t)] (g'(y) + r(s))$$

We now take the limit of  $t \rightarrow x$  on both sides.  $\lim_{t \rightarrow x} u(t) = 0$ , and  $f$  is differentiable and hence continuous at  $x$ , so  $s = f(t) \rightarrow y$  as  $t \rightarrow x$ . Thus,  $r(s) \rightarrow 0$  as  $t \rightarrow x$ , and in conclusion:

$$h'(x) = (g \circ f)'(x) = f'(x)g'(y) = g'(f(x))f'(x)$$

as desired. □

## 5.2 MVT

### Definition 5.7: Local Maxima/Minima

Let  $X$  be a metric space. Let  $f : X \mapsto \mathbb{R}$ , and let  $x \in X$ . We say that  $x$  is a **local maximum** of  $f$  if there exists  $\delta > 0$  such that  $f(y) \leq f(x)$  for all  $y \in N_\delta(x)$ . A **local minimum** is defined similarly, with  $f(y) \geq f(x)$  instead.

For a metric space  $X$  equipped with the discrete metric, all points  $x \in X$  are simultaneously local maxima and minima. To see this, take any  $0 < \delta \leq 1$ .

### Theorem 5.8

Let  $f : [a, b] \mapsto \mathbb{R}$ . Let  $x \in [a, b]$  and suppose that  $f'(x)$  exists, and  $f$  is either a local maximum or local minimum of  $f$ . Then,  $f'(x) = 0$ .

#### Proof

Suppose  $x$  is a local minimum. Then, there exists  $\delta > 0$  such that  $N_\delta(x) \subset [a, b]$ , and  $f(y) \geq f(x)$  for all  $y \in N_\delta(x)$ . Thus, if  $x < y < x + \delta$ , then:

$$\frac{f(y) - f(x)}{y - x} \geq 0 \implies f'(x) \geq 0$$

Conversely, if  $x - \delta < y < x$ , then:

$$\frac{f(y) - f(x)}{y - x} \leq 0 \implies f'(x) \leq 0$$

So taken together we obtain that  $f'(x) = 0$ . An identical argument is used for the case of a local maximum.  $\square$

### Theorem: Rolle's Theorem

Let  $f : [a, b] \mapsto \mathbb{R}$  be continuous, and suppose  $f$  is differentiable on  $(a, b)$ . If  $f(a) = f(b)$ , then there exists  $x \in (a, b)$  such that  $f'(x) = 0$ .

#### Proof

Since  $[a, b]$  is compact and  $f$  is continuous, by the EVT (Theorem 4.16)  $f$  attains its maximum on  $[a, b]$ , that is, there exists  $c \in [a, b]$  such that  $f(y) \leq f(c)$  for all  $y \in [a, b]$ . If  $c \in (a, b)$ , then by Theorem 5.8,  $f'(c) = 0$  and we are done. Next, suppose  $c = a$  or  $c = b$ . Again by the EVT,  $f$  attains its minimum on  $[a, b]$ , that is, there exists  $d \in [a, b]$  such that  $f(y) \geq f(d)$  for all  $y \in [a, b]$ . If  $d \in (a, b)$ , then by Theorem 5.8,  $f'(d) = 0$  and we are done. Suppose then that  $d = a$  or  $d = b$ . Since  $f(a) = f(b)$ , we therefore obtain that  $f(a) = f(b) = f(c) = f(d)$  and the maximum/minimum values agree. Hence,  $f(y) = f(a)$  for all  $y \in [a, b]$ , so  $f'(y) = 0$  for all  $y \in [a, b]$ . So, the desired  $x$  may be any point in  $[a, b]$ .  $\square$



Figure 21: A simple parabolic function that demonstrates Rolle's Theorem.

#### Theorem 5.10: Mean Value Theorem

Let  $f : [a, b] \mapsto \mathbb{R}$  be continuous and differentiable on  $(a, b)$ . Then, there exists  $x \in (a, b)$  such that  $f(b) - f(a) = f'(x)(b - a)$ .

The visual interpretation of this theorem is that there exists  $x \in (a, b)$  such that the slope of the tangent line to  $f$  at  $x$  is equal to the secant line slope between  $(a, f(a))$  and  $(b, f(b))$ . The idea of the proof is to rotate one's head such that the secant line is horizontal; one is then able to apply Rolle's Theorem!

#### Proof

Define  $h(y) = f(y) - \frac{f(b)-f(a)}{b-a}(y-a)$ .  $h$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$  (being a sum of continuous/differentiable functions). We have that  $h(a) = f(a) - 0 = f(a)$ , and  $h(b) = f(b) - \frac{f(b)-f(a)}{b-a}(b-a) = f(a)$ . Applying Rolle's Theorem to  $h$ , there exists  $x \in (a, b)$  such that  $h'(x) = 0$ . Therefore,  $h'(x) = 0 = f'(x) - \frac{f(b)-f(a)}{b-a} = 0$ , and we conclude that  $f(b) - f(a) = f'(x)(b - a)$  for some  $x \in (a, b)$ .  $\square$



Figure 22: A simple continuous function that demonstrates the MVT.

### Theorem 5.11

Let  $f : [a, b] \mapsto \mathbb{R}$  be differentiable on  $(a, b)$ . Then:

- (a) If  $f'(x) \geq 0$  for all  $x \in (a, b)$ , then  $f$  is monotonically increasing.
- (b) If  $f'(x) = 0$  for all  $x \in (a, b)$ , then  $f$  is constant.
- (c) If  $f'(x) \leq 0$  for all  $x \in (a, b)$ , then  $f$  is monotonically decreasing.

### Proof

If  $a < x < y < b$ , by the mean value theorem, there exists  $z \in (x, y)$  such that:

$$f(y) - f(x) = f'(z)(y - x)$$

Note that  $y - x > 0$  by construction.

- (a) If  $f'(x) \geq 0$  for all  $x \in (a, b)$ , then  $f'(z) \geq 0$ , showing that  $f(y) - f(x) \geq 0$  and hence that  $f$  is monotonically increasing.
- (b) If  $f'(x) = 0$  for all  $x \in (a, b)$ , then  $f'(z) = 0$ , showing that  $f(y) - f(x) = 0$  and hence that  $f$  is constant on  $(a, b)$ .
- (c) If  $f'(x) \leq 0$  for all  $x \in (a, b)$ , then  $f'(z) \leq 0$ , showing that  $f(y) - f(x) \leq 0$  and hence that  $f$  is monotonically decreasing.  $\square$

## 5.3 Taylor's Theorem

### Definition 5.14: Higher Order Derivatives

If  $f$  is differentiable in a neighbourhood of  $x$ , then we may compute a second order derivative:

$$\lim_{t \rightarrow x} \frac{f'(t) - f'(x)}{t - x} = (f')'(x) = f''(x)$$

We can then continue this process to obtain  $f^{(3)}(x), f^{(4)}(x), \dots, f^{(n)}(x)$ .

### Definition: $C^n(I, \mathbb{R})$

If  $f$  is continuous in  $I$ , we can write  $f \in C^0(I, \mathbb{R})$ . If  $f$  is differentiable in a neighbourhood  $I$  and the derivative  $f'$  is continuous in  $I$ , then we write  $f \in C^1(I, \mathbb{R})$ . In general,  $f \in C^n(I, \mathbb{R})$  denotes the  $n$ th derivative of  $f$  is continuous in  $I$ . Note that where it is clear from context, we may drop the  $\mathbb{R}$  and just write  $C^n(I)$ .

Recall that for a function  $f$  continuous and differentiable on  $(x_0, x)$ , the Mean Value Theorem proved the existence of some  $\tilde{x}$  such that:

$$f(x) = f(x_0) + f'(\tilde{x})(x - x_0)$$

This gives us a natural method to build up approximations for functions; we can start with a constant approximation  $f(x_0)$ , then add a linear term  $f'(\tilde{x})(x - x_0)$ , then add on a quadratic term  $(x - x_0)^2$  and so on. The following theorem gives us a way to construct these approximations and bound their error.

### Theorem 5.15: Taylor's Theorem

Let  $I$  be a neighbourhood of  $x_0$ , and  $f \in C^p(I)$ . Then, for any  $n < p$ , we have that:

$$f(x) = \sum_{j=0}^n \frac{f^{(j)}(x_0)}{j!} (x - x_0)^j + \frac{f^{(n+1)}(\tilde{x})}{(n+1)!} (x - x_0)^{n+1}$$

Where  $\tilde{x} = x_0 + \lambda(x - x_0)$  for some  $\lambda \in (0, 1)$  (i.e.  $\tilde{x} \in (x_0, x)$ ). Note that  $\tilde{x}$  depends on  $x, x_0$ , and  $n$ .

#### Proof

For  $n = 0$ , the claim reduces to the Mean Value Theorem. Then, let  $n \geq 1$ . Let  $A$  be a constant that depends on  $x, x_0$ , and  $n$ , and let  $P_n(x) = \sum_{j=0}^n \frac{f^{(j)}(x_0)}{j!} (x - x_0)^j$ . Then, we can write:

$$f(x) = P_n(x) + A(x - x_0)^{n+1}$$

We need to show that we can express  $A$  as relating to the derivative, namely, that there exists  $\tilde{x}$  such that  $A = \frac{f^{(n+1)}(\tilde{x})}{(n+1)!}$ . Let  $g(t) = f(t) - P_n(t) - A(t - x_0)^{n+1}$  with  $t \in I$ . Then,  $g \in C^p(I)$ . For  $n < p$ , we then have that:

$$g^{(n+1)}(t) = f^{(n+1)}(t) - 0 - A(n+1)!(t - x_0)$$

We claim that there exists  $\tilde{x} \in (x_0, x)$  such that  $g^{(n+1)}(\tilde{x}) = 0$ . To see this, consider that  $P^{(j)}(x_0) = f^{(j)}(x_0)$  for  $j = 0, 1, \dots, n$ , so  $g(x_0) = 0$ , and furthermore:

$$g'(x_0) = g''(x_0) = g^{(3)}(x_0) = \dots = g^{(n)}(x_0) = 0$$

Moreover by the choice of  $n$ , we have that  $g(x) = 0$ . Hence, by Rolle's Theorem, there exists a point  $x_1$  between  $x_0$  and  $x$  such that  $g'(x_1) = 0$ . Similarly, repeating the argument above, there exists an  $x_2$  between  $x_0$  and  $x_1$  such that  $g''(x_2) = 0$ . Repeating this process up to  $g^{(n)}$ , we have that  $g^{(n)}(x_n) = 0$ , for some  $x_0 < x_n < x_{n-1} < \dots < x$  and in turn, there exists  $x_{n+1} \in (x_0, x_n)$  such that  $g^{(n+1)}(x_{n+1}) = 0$ . Setting  $\tilde{x} = x_{n+1}$ , the claim is shown.  $\square$

As an example, we consider the Taylor series of the function  $f(x) = \cos(x)$  (We will formally define this function later on, but for now, let us assume its familiar properties and derivatives). We then have that:

$$f^{(j)}(0) = \begin{cases} (-1)^m & \text{if } j = 2m \\ 0 & \text{if } j = 2m + 1 \end{cases}$$

If we have the sum run from  $j = 0$  to some  $j = n$ , let us then try to estimate the rest. Let  $\tilde{x} \in (0, x)$ . Then, letting the error term be represented by  $\epsilon$ , we have that:

$$\epsilon = \left| \frac{f^{(n+1)}(\tilde{x})}{(n+1)!} (x - x_0)^{n+1} \right| \leq \frac{|x|^{n+1}}{(n+1)!}$$

And we observe that  $\lim_{n \rightarrow \infty} \frac{|x|^{n+1}}{(n+1)!} = 0$  and hence the error  $\epsilon$  goes to zero in the  $n \rightarrow \infty$  limit. Therefore, the difference between  $\cos(x)$  and its Taylor polynomial vanishes quickly for any  $x$ , and the Taylor series

converges for all  $x$ . Taking the limit of the sum, we have that:

$$f(x) = \cos(x) = \lim_{n \rightarrow \infty} \left( P_n(x) + \frac{f^{(n+1)}(x_0)}{(n+1)!} (x - x_0)^{n+1} \right) = \sum_{j=0}^{\infty} \frac{(-1)^{2m}}{m!} x^{2m} = 1 - \frac{x^2}{2} + \frac{x^4}{4} - \frac{x^6}{6} + \dots$$

A question of interest might be how many terms do we need in the Polynomial such that our error is less than  $10^{-6}$ , say, for estimating the value of  $\cos\left(\frac{\pi}{12}\right)$ . In other words, we want to find the  $m$  such that:

$$\frac{1}{2m!} \left( \frac{\pi}{12} \right)^{2m} \leq 10^{-6}$$

Rearranging, we require:

$$(2m)! \left( \frac{12}{\pi} \right)^{2m} > 10^6$$

Making a table of the value of the LHS as a function of  $m$ , we have: So we see that three terms are sufficient

$m$	$(2m)! \left( \frac{12}{\pi} \right)^{2m}$
1	$\approx 29$
2	$\approx 5110$
3	$\approx 2.23 \times 10^6$

for a good approximation in this case (and as stated before, the series converges very quickly)!

A natural question of interest is the convergence of the sum in the  $N \rightarrow \infty$  limit, that is, the convergence of the power series  $\sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$ . We are also interested are interested for when  $f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$  holds (that is, when is a function equal to its Taylor series)? These turn out to be distinct questions; in particular, there are functions whose power series converge for all  $x$  but are equal to their power series nowhere (except at  $x_0$  where equality must hold). This motivates the following definition:

**Definition: Analyticity**

A function  $f$  is **analytic** if  $f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$  in a neighbourhood of  $x_0$ .

To motivate this definition, it will help to study a function which is *not* analytic. Consider the function:

$$f(x) = \begin{cases} \exp\left(-\frac{1}{x}\right) & x > 0 \\ 0 & x \leq 0 \end{cases}$$



Figure 23: Plot of  $f$ .

$f$  is continuous everywhere by construction. It is infinitely differentiable at  $x = 0$ , but it is only equal to its Taylor series around  $x_0 = 0$  for  $x \leq 0$ . To see this, we observe that:

$$f'(x) = \frac{1}{x^2} \exp\left(-\frac{1}{x}\right)$$

$$f^{(n)}(x) = \frac{P_n(x)}{x^{2n}} \exp\left(-\frac{1}{x}\right)$$

We have that  $f^{(n)}(x) \rightarrow 0$  as  $x \rightarrow 0$  for all  $n$  as the exponential dominates the polynomial singularity. We hence have that  $f \in C^\infty$ . The Taylor polynomial at  $x_0 = 0$  however, as we have that  $f(0) = 0$  and  $f^{(n)}(0) = 0$ , leading to:

$$\sum_{n=0}^N \frac{f^{(n)}(0)}{n!} x^n = 0$$

for all  $N$ . We therefore have that the series converges for all  $x$ , but is only equal to  $f$  for  $x \leq 0$ ; hence it is not analytic (as there exists no neighbourhood around  $x_0 = 0$  for which  $f$  is equal to its Taylor series).

We now consider a function  $\chi(x)$  defined as  $\chi(x) = \frac{f(x)}{f(x)+f(1-x)}$ . This function is also continuous, its denominator is never zero, and  $\chi \in C^\infty$ . We observe that  $\chi(x) = 0$  for  $x \leq 0$  and  $\chi(x) = 1$  for  $x \geq 1$ , and overall the function looks much like a step function:



Figure 24: Plot of  $\chi$ .

Indeed, this function can be used as a “cutoff”/“switch” function that behaves much like a step function (except it is infinitely differentiable).

A question becomes whether such a function could be analytic. The answer turns out to be no, and the proof we leave as an exercise. As a sketch, consider an analytic function  $g$  such that  $g(x) = 0$  for  $x \leq x_0$  and  $g(x) \neq 0$  for  $x > x_0$ . One can derive a contradiction by considering the Taylor series expansion about  $x_0$  and then using the assumed analyticity of  $g$ .

Note that in a sense, Taylor’s Theorem is the culmination of a sequence of theorems we have proven in the course. Roughly, the sequence was as follows:

- 1)  $f : X \mapsto Y$  and  $K \subset X$ , then  $f(K)$  compact (Theorem 4.14)
- 2) Extreme Value Theorem: If  $f : X \mapsto \mathbb{R}$  with  $f$  continuous and  $K$  compact, then  $f$  realizes its supremum and infimum on  $K$ . (Theorem 4.16)
- 3) Rolle’s Theorem
- 4) Mean Value Theorem (Theorem 5.10)
- 5) Taylor’s Theorem (Theorem 5.15)



A question that arises is could we have gone through this sequence of proofs with just the rational numbers ( $\mathbb{Q}$ )? The intuitive answer is no, but it may be interesting to see where along this chain the logic breaks down.

The first step that looks immediately questionable is step 2; the supremum/infimum is not well-defined for all subsets of  $\mathbb{Q}$ , so we might be able to find a breakdown there. To this end, we consider the set  $A = \{q \in \mathbb{Q} : q > 0, q^2 < 2\}$  that arises in Example 1.1 and try to find a function  $f : K \mapsto \mathbb{Q}$  with  $K \subset \mathbb{Q}$  compact such that  $f(K) = A$ . An idea would be to try  $f(x) = \sqrt{x}$ , with  $K = \{q \in \mathbb{Q} : \exists r : r^2 = q\} \cap [0, 2]$  but this doesn't work as  $K$  is not compact. Trying another attempt,  $K = [1, 2] \cap \mathbb{Q}$  with  $f(x) = \sin(x)$  does not work either as  $\sin(x)$  is not necessarily rational, and moreover,  $[1, 3] \cap \mathbb{Q}$  is not a compact set as not all Cauchy sequences in  $S$  converge! Another attempt would be  $K = [1, 2] \cap \mathbb{Q} = X$  with  $f(q) = |q^2 - 2|$  where it would seem as though  $f(K) = \{r \in \mathbb{Q} : 0 < r \leq 2\}$  provides a good counterexample, but this fails for the same reason as  $K$  is not compact. Finding a valid counterexample for the EVT is therefore difficult.

An easier break along the chain to find is with Rolle's Theorem. One can consider the function  $f(x) = x^2 - 2$  which can break the Intermediate value theorem as 0 is not contained in the image if the domain is  $\mathbb{Q}$ . We can dress this up to construct a counterexample for Rolle's Theorem.

## 5.4 Local Behavior of Functions

### Theorem: Second Derivative Test

Suppose  $f \in C^3(I)$  where  $I$  is a neighbourhood of  $x_0$ . Furthermore, suppose that  $f'(x_0) = 0$ . If  $f''(x_0) > 0$ , then  $x_0$  is a local minimum. Conversely, if  $f''(x_0) < 0$ , then  $x_0$  is a local maximum.

### Proof

By Taylor's Theorem (Theorem 5.15), if we let  $x = x_0 + h$  for  $h > 0$ , there exists some  $\tilde{x} = x_0 + \lambda h$  with  $\lambda \in (0, 1)$  such that:

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{1}{2}f''(x_0)(x - x_0)^2 + \frac{1}{6}f^{(3)}(\tilde{x})(x - x_0)^3$$

Using that  $f'(x_0) = 0$ , we have:

$$f(x) - f(x_0) = h^2 \left( \frac{1}{2}f''(x_0) + \frac{1}{6}f^{(3)}(x_0 + \lambda h)h \right)$$

Let  $0 < \epsilon < \left| \frac{1}{2}f''(x_0) \right|$ . Then, by the assumed continuity of  $f^{(3)}$ , we have that there exists  $\delta > 0$  such that  $|h| < \delta$  implies  $\left| \frac{1}{6}f^{(3)}(x_0 + \lambda h)h \right| < \epsilon$ . Hence, for sufficiently small  $h$ , we have that:

$$\text{sgn}(f(x) - f(x_0)) = \text{sgn}(f''(x_0))$$

So we conclude that if  $f'(x_0) = 0$  and  $f''(x_0) > 0$  then  $x_0$  is a local minimum, and if  $f''(x_0) < 0$ , then  $x_0$  is a local maximum.  $\square$

### Definition: Convex Functions

Let  $f : (a, b) \mapsto \mathbb{R}$  is **convex** if for all  $x, y \in (a, b)$  with  $a < x < y < b$  and for all  $\lambda \in (0, 1)$ ,

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$$

Note that an alternative definition of convexity is that for any  $x, y$  with  $x < y$ , the function evaluated at some point  $z \in (x, y)$  will always be below the average of the function at  $x, y$ . That is:

$$f(z) \leq \frac{f(x) + f(y)}{2}$$



Figure 25: Visualization of a convex and non-convex function. For the upwards facing parabola, the function (and hence all points  $(\lambda x + (1 - \lambda)y, f(\lambda x + (1 - \lambda)y))$  for  $\lambda \in (0, 1)$ ) always lies below the line connecting  $(x, f(x))$  and  $(y, f(y))$  for  $a < x < y < b$  and hence it is convex. For the downwards facing parabola, this is no longer true and the function is not convex (it is instead *concave*).

Another equivalent way of defining convexity is to say that  $f$  is convex if and only if for all  $a < s < t < y < b$ :

$$\frac{f(t) - f(s)}{t - s} \leq \frac{f(u) - f(t)}{u - t}$$



Figure 26: Visualization of the alternative definition of convexity. For any  $a < s < t < u < v$ , the slope of the line segment joining  $s$  and  $t$  is less than the slope of the line segment joining  $t$  and  $u$ .

### Theorem

- (a) Assume that  $f : (a, b) \mapsto \mathbb{R}$  is convex. Then,  $f$  is continuous.
- (b) Assume that  $f \in C^1(a, b)$ . Then, if  $f$  is convex,  $f'$  is increasing.
- (c) Assume that  $f \in C^2(a, b)$ . Then  $f$  convex implies  $f'' \geq 0$ .

### Proof

- (a) Let  $[c, d] \subset (a, b)$  and  $a < c_1 < c < x < y < d < d_1 < b$ . By convexity, we have that:

$$\frac{f(y) - f(x)}{y - x} \leq \frac{f(d) - f(y)}{d - y} \leq \frac{f(d_1) - f(d)}{d_1 - d}$$

and also that:

$$\frac{f(y) - f(x)}{y - x} \geq \frac{f(x) - f(c)}{x - c} \geq \frac{f(c) - f(c_1)}{c - c_1}$$

We therefore have that:

$$\left\{ \left| \frac{f(y) - f(x)}{y - x} \right| : c < x < y < b \right\} < M$$

for some  $M \in \mathbb{R}$ . Therefore,  $|f(y) - f(x)| < M|y - x|$  for all  $x, y \in (c, d)$ . This holds for all  $[c, d] \subset (a, b)$ , showing the continuity of  $f$ .

- (b) Let  $f$  be convex, and let  $a < c < x < y < d < b$ . Then, by convexity we have that:

$$\frac{f(x) - f(c)}{x - c} \leq \frac{f(y) - f(x)}{y - x} \leq \frac{f(d) - f(y)}{d - y}$$

Ignoring the central term in the inequality, and taking the limit as  $x \rightarrow c$  and  $y \rightarrow b$ , we have that:

$$f'(c) \leq f'(b)$$

So we conclude that  $f'$  is increasing on  $(a, b)$ .

- (c) If  $f$  is convex,  $f'$  is increasing by (b). Then, we have that for any  $a < x < y < b$ :

$$\frac{f(y) - f(x)}{y - x} \geq 0$$

And hence taking  $y \rightarrow x$  we have that  $f'(x) \geq 0$ . □

### Corollary

If  $f \in C^3(I)$  and  $f'(x_0) = 0$  and  $f''(x_0) \neq 0$ ,  $x_0$  is a local minimum if  $f$  is convex in a neighbourhood of  $x_0$ .

## 6 The Riemann-Stieltjes Integral

### 6.1 Definition of the Integral

#### Definition 6.1: Partition

A **partition** of  $[a, b] \subset \mathbb{R}$  is a set  $\{x_0, x_1, \dots, x_n\}$  (for some  $n \in \mathbb{N}$ ) such that:

$$a = x_0 \leq x_1 \leq x_2 \leq \dots \leq x_{n-1} \leq x_n = b$$

We can then write:

$$\Delta x_i = x_i - x_{i-1}$$



Figure 27: Visualization of a partition  $\{x_0, x_1, x_2, x_3, x_4\}$  of  $[a, b]$ . Note that the points in the partitions need not be equally spaced.

#### Definition 6.1: Upper and Lower Sums

Given  $f : [a, b] \mapsto \mathbb{R}$  and a partition  $P$  of  $[a, b]$  let:

$$M_i = \sup \{f(x) : x_{i-1} \leq x \leq x_i\}$$

$$m_i = \inf \{f(x) : x_{i-1} \leq x \leq x_i\}$$

Then, we can define the **upper** and **lower sums**:

$$U(P, f) = \sum_{i=1}^n M_i \Delta x_i$$

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i$$



Figure 28: Example of a function  $f$ , a partition  $P$  of  $[a, b]$ , and the  $M_i, m_i$ s for this choice of partition.

By construction, it should be evident that  $L(P, f) \leq U(P, f)$  for all  $P, f$ .

A natural question that arises from the form of the above expression is whether these are Riemann sums or not. Recall from first year calculus that we would choose the left endpoint, right endpoint, or some other arbitrary choice of a point in the subinterval. Here, we in a sense use a “special case” of the supremum/infimum. We will see that this choice is much easier to use in proofs due to monotonicity properties. Namely, if we have a partition and add another point, then  $U(P, f)$  can only decrease, and  $L(P, f)$  can only increase (we will see this in a theorem soon)!

### Definition 6.1: Upper/Lower Integrals and Riemann Integrability

We define the **upper Riemann integral** to be:

$$\overline{\int_a^b} = \inf_P U(P, f).$$

and the **lower Riemann integral** to be:

$$\underline{\int_a^b} = \sup_P L(P, f).$$

Here, the infimum/supremum is taken over all partitions  $P$ . We say that  $f$  is **Riemann integrable** on  $[a, b]$ , and write  $f \in \mathcal{R}[a, b]$  if:

$$\overline{\int_a^b} = \underline{\int_a^b}$$

which we can write as:

$$\int_a^b f dx \text{ or } \int_a^b f(x) dx$$

Note that the choice of variable in the above definition is totally arbitrary.

Also, note that while  $f$  is not required to be continuous in the above definition, it is required to be bounded; else,  $M_i$  and  $m_i$  may not exist. Since  $f$  is bounded,  $U(P, f)$ ,  $L(P, f)$  are bounded for all  $P, f$  and hence we have a set of real numbers for which we may consider the supremum/infimum of by the LUB/GLB property of the reals. Since the upper/lower sums lie in a bounded interval, there is no questions about whether the lower/upper integrals exist. The question becomes whether they are equal or not. Before getting into further discussion on this topic, we discuss a bound:

### Theorem 6.1: ML Bounds

Let  $m = \inf \{f(x) : a \leq x \leq b\}$  and  $M = \sup \{f(x) : a \leq x \leq b\}$  (which exist by the boundedness of  $f$ ). Then,

$$m(b - a) \leq L(P, f) \leq U(P, f) \leq M(b - a)$$

For any choice of partition  $P$ .

**Proof**

For any  $i$ , we have that:

$$m \leq m_i \leq M_i \leq M$$

Therefore:

$$\sum_{i=1}^n m \Delta x_i \leq \sum_{i=1}^n m_i \Delta x_i \leq \sum_{i=1}^n M_i \Delta x_i \leq \sum_{i=1}^n M \Delta x_i$$

So we conclude that:

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a)$$

□

Now that we have established the Riemann integral, a natural question is how can we extend this notion. In order to do so, we will use a monotonically increasing function  $\alpha : [a, b] \mapsto \mathbb{R}$  (that is,  $\alpha(x) \leq \alpha(y)$  for all  $x \leq y$ ). Note that  $\alpha$  need not be continuous. Indeed, compared to the Riemann integral where  $\alpha(x) = x$  and was continuous, in this general setting,  $\alpha$  is allowed to have jumps. This allows for certain benefits, as we will soon discuss. However, we note that  $\alpha$  can only have a finite number of jumps.

**Theorem 4.30**

Let  $\alpha : [a, b] \mapsto \mathbb{R}$  be monotonic. Then, it can only have finitely many discontinuities.

**Proof**

Assign a rational number  $r(x)$  to each of the discontinuities of  $\alpha$ . Then, we have that:

$$\lim_{x \rightarrow r(x)^-} \alpha(x) < \alpha(x) < \lim_{x \rightarrow r(x)^+} \alpha(x)$$

Since  $x_1 < x_2$  implies  $\lim_{x \rightarrow r(x_1)^+} \alpha(x) \leq \lim_{x \rightarrow r(x_2)^-} \alpha(x)$ , we have that  $r(x_1) \neq r(x_2)$  if  $x_1 \neq x_2$ . We therefore have established a function  $r$  from the set of discontinuities of  $\alpha$  to the rationals. As the rationals are countable, the set of discontinuities of  $\alpha$  are also countable. □

With this established, we now define the generalized Riemann-Stieltjes integral.

### Definition 6.2: Riemann-Stieltjes Integral

Let  $\alpha : [a, b] \mapsto \mathbb{R}$  be increasing, and given a partition  $P$  of  $[a, b]$ , define:

$$\Delta\alpha_i = \alpha(x_i) - \alpha(x_{i-1}) (\geq 0)$$

For bounded  $f : [a, b] \mapsto \mathbb{R}$ , Let:

$$U(P, f, \alpha) = \sum_{i=1}^n M_i \Delta\alpha_i$$

$$L(P, f, \alpha) = \sum_{i=1}^n m_i \Delta\alpha_i$$

We then take the infimum/supremum over partitions  $P$  to get:

$$\overline{\int_a^b f d\alpha} = \inf_P U(P, f, \alpha)$$

$$\underline{\int_a^b} = \sup_P L(P, f, \alpha)$$

If equal, we write their value as:

$$\int_a^b f d\alpha \text{ or } \int_a^b f(x) d\alpha(x)$$

and we write that  $f \in \mathcal{R}_\alpha[a, b]$ . In the case where  $\alpha(x) = x$ , we recover the Riemann integral.

Why is this definition useful? What does it accomplish for us that the original Riemann integral does not? We consider a physically motivating example. Suppose we have a thin wire with varying mass density  $\rho(x)$ . If we wanted to calculate the mass density of the wire, we would integrate the density  $\rho(x)$  over the length of the wire. Now, suppose our wire consists of steel of continuously varying mass density, as well as beads/point masses placed on certain locations of the wire. The Riemann integral cannot handle these point masses, but the Riemann-Stieltjes integral can deal with this case if we use an  $\alpha$  with discontinuities in it. Hence, the Riemann-Stieltjes integral allows us to handle cases where we both have continuous and discrete masses to integrate over. It acts as a bridge between Riemann and Lebesgue integration (the latter of which will be the subject of a later course in measure theory).

We now will answer the question: “for what choices of  $f, \alpha$  is  $f$  Riemann-Stieltjes integrable?”

## 6.2 Criterion for Integrability

### Definition 6.3: Refinements and Common Refinement

$P^*$  is a **refinement** of  $P$  if  $P \subset P^*$  and  $P, P^*$  are partitions. The common refinement of  $P_1, P_2$  is  $P^* = P_1 \cup P_2$ .

### Theorem 6.4

If  $P^*$  is a refinement of  $P$ , then:

$$L(P, f, \alpha) \leq L(P^*, f, \alpha) \leq U(P^*, f, \alpha) \leq U(P, f, \alpha)$$

As a remark, when we take infimums/supremums over partitions  $P$  to obtain the upper/lower Riemann-Stieltjes integrals, we are taking refinements.

Also, note that the above theorem does *not* apply to (right-hand, left-hand, midpoint, arbitrary) Riemann sums, and is a consequence of the choice of upper/lower sums with supremums/infimums taken over the subintervals.

### Proof

It suffices to consider  $P^*$  with a single extra point  $x_{i-1} < x^* < x_i$ , and then the genral case follows by induction.

For the case where  $\alpha(x) = x$ , the refinement adds  $(m^* - m_i)(x^* - x_{i-1}) \geq 0$ .

For the general case, we have that:

$$\begin{aligned} L(P^*, f, \alpha) - L(P, f, \alpha) &= (m^*(\alpha(x^*) - \alpha(x_{i-1})) + m_i(x^* - x_{i-1})) - m_i(\alpha(x_i) - \alpha(x_{i-1})) \\ &= (m^*(\alpha(x^*) - \alpha(x_{i-1})) + m_i(x^* - x_{i-1})) \\ &\quad - m_i[(\alpha(x^*) - \alpha(x_{i-1})) + (\alpha(x_i) - \alpha(x^*))] \\ &= (m^* - m_i)(\alpha(x^*) - \alpha(x_i)) \end{aligned}$$

$\alpha$  is monotonically increasing, so  $x^* \geq x_i$  implies that the second term is positive. Furthermore,  $m^* \geq m_i$  as  $\inf \{f(x) : x \in [x_{i-1}, x^*]\} \geq \inf \{f(x) : x \in [x_{i-1}, x_i]\}$ . It follows that:

$$L(P^*, f, \alpha) - L(P, f, \alpha) \geq 0$$

and the proof for  $U(P^*, f, \alpha) - U(P, f, \alpha) \leq 0$  follows analogously. □



Figure 29: Visualization of the effect of adding an extra point  $x^*$  to the partition  $P$ . We can see that this has the net effect of increasing  $L(P, f, \alpha)$  as  $m^* \geq m_i$ .



Note that as a point of notation, if  $f, \alpha$  are fixed, we sometimes can write  $L(P), U(P)$  in place of  $L(P, f, \alpha)$  and  $U(P, f, \alpha)$ . Sometimes where the context is clear, it is also common to write  $\mathcal{R}(\alpha)$  in place of  $\mathcal{R}_\alpha[a, b]$ .

### Theorem 6.5

$$\underline{\int_a^b} f d\alpha \leq \overline{\int_a^b} f d\alpha.$$

### Proof

For partitions  $P_1, P_2$ , let  $P^* = P_1 \cup P_2$  be the common refinement. By Theorem 6.4, we have that:

$$L(P_1) \leq L(P^*) \leq U(P^*) \leq U(P_2)$$

And in particular,  $L(P_1) \leq U(P_2)$ . Therefore, for any fixed  $P_2$ ,  $U(P_2)$  is an upper bound on the set of all lower sums. As the supremum is the least upper bound, we have that:

$$\sup_{P_1} L(P_1) \leq U(P_2)$$

Therefore, as  $\sup_{P_1} L(P_1)$  is a lower bound on the set of all upper sums, and the infimum is the greatest lower bound, we have that:

$$\sup_{P_1} L(P_1) \leq \inf_{P_2} U(P_2)$$

So therefore:

$$\underline{\int_a^b} f d\alpha \leq \overline{\int_a^b} f d\alpha$$

as claimed. □

### Theorem 6.6: $\epsilon$ -Criterion for Integrability

$f \in \mathcal{R}_\alpha[a, b]$  if and only if for all  $\epsilon > 0$ , there exists a partition  $P_\epsilon$  of  $[a, b]$  such that  $U(P_\epsilon) - L(P_\epsilon) < \epsilon$ .

### Proof

$\Rightarrow$  By hypothesis,  $\sup_P L(P) = \int_a^b f d\alpha = \inf_P U(P)$ . Let  $\epsilon > 0$ . Then by the property of sup/inf, there exist  $P_1, P_2$  such that:

$$\begin{aligned}\int_a^b f d\alpha - L(P_1) &< \frac{\epsilon}{2} \\ U(P_2) - \int_a^b f d\alpha &< \frac{\epsilon}{2}\end{aligned}$$

Adding the first inequality to the second, we get:

$$U(P_2) - L(P_1) < \epsilon$$

Letting  $P^* = P_1 \cup P_2$ , by Theorem 6.4 we have that:

$$U(P^*) - L(P^*) \leq U(P_2) - L(P_1) < \epsilon$$

which proves the claim.

$\Leftarrow$  Let  $\epsilon > 0$  Then by Theorem 6.5 we have that:

$$0 \leq \overline{\int_a^b f d\alpha} - \underline{\int_a^b f d\alpha}$$

and furthermore:

$$0 \leq \overline{\int_a^b f d\alpha} - \underline{\int_a^b f d\alpha} \leq U(P_\epsilon) - L(P_\epsilon) < \epsilon$$

Where the second inequality is true for any choice of partition.  $\epsilon$  is arbitrary, so we conclude that:

$$\overline{\int_a^b f d\alpha} - \underline{\int_a^b f d\alpha} = 0$$

And therefore  $\overline{\int_a^b f d\alpha} = \underline{\int_a^b f d\alpha}$ , and  $f \in \mathcal{R}_\alpha[a, b]$ . □

Theorem 6.7 is a little technical, so we shall skip it for now.

### Theorem 6.8: Continuity implies integrability

If  $f$  is continuous on  $[a, b]$ , then  $f \in \mathcal{R}_\alpha[a, b]$ .

Note in the above theorem that we make no assumptions on  $\alpha$ , only that (of course) it is monotonic.

**Proof**

By definition,  $U(P) - L(P) = \sum_{i=1}^n (M_i - m_i) \Delta \alpha_i$ . The idea will be to choose small intervals to make these differences small. Since  $[a, b]$  is compact,  $f$  is uniformly continuous by Theorem 4.19. So, for all  $\eta > 0$ , there exists  $\delta > 0$  such that  $|f(x) - f(t)| < \eta$  if  $|x - t| < \delta$ . Thus, if  $P$  is constructed such that  $\Delta x_i < \delta$  then  $M_i - m_i < \eta$ . We then have that:

$$U(P) - L(P) \leq \sum_{i=1}^n \eta \Delta \alpha_i = \eta \sum_{i=1}^n (\alpha(x_i) - \alpha(x_{i-1})) = \eta (\alpha(b) - \alpha(a))$$

Where in the last equality we use the fact that we have a telescoping sum. Given  $\epsilon > 0$ , we choose  $\eta < \frac{\epsilon}{\alpha(b) - \alpha(a)}$ . With this choice of partition with  $\Delta x_i < \delta = \delta(\eta)$ , we have that:

$$U(P) - L(P) < \epsilon$$

and we conclude that  $f \in \mathcal{R}_\alpha[a, b]$  by Theorem 6.6. □

A natural question is “what  $f$ s are Riemann integrable in general?” The answer turns out to be “if  $f$  is continuous almost everywhere”. This sounds handwavy, but has a precise definition; although it is not covered in this course, one can refer to Rudin 11.33(b) for details.

**Theorem 6.9**

If  $f$  is monotone on  $[a, b]$  and  $\alpha$  is continuous on  $[a, b]$  then  $f \in \mathcal{R}_\alpha[a, b]$ .

In the proof of Theorem 6.8, we used the continuity of  $f$  to bound the maximum/minimum on each subinterval. Here,  $f$  is no longer continuous, so we cannot control the maximum/minimum. Instead, we will use the continuity of  $\alpha$  to control the size of the  $\Delta \alpha$ s.

**Proof**

Given  $n \in \mathbb{N}$ , choose  $P$  such that  $\Delta \alpha_i = \frac{\alpha(b) - \alpha(a)}{n}$  for all  $i \in 1, \dots, n$ . Note that such a choice is possible by the continuity of  $\alpha$  and the IVT (Theorem 4.23). We then have that:

$$U(P) - L(P) = \sum_{i=1}^n (M_i - m_i) \Delta \alpha_i = \frac{\alpha(b) - \alpha(a)}{n} \sum_{i=1}^n (M_i - m_i)$$

Suppose (WLOG) that  $f$  is an increasing function. Then,  $M_i = f(x_i)$  and  $m_i = f(x_{i-1})$  due to the monotone increasing property. Hence:

$$U(P) - L(P) = \frac{\alpha(b) - \alpha(a)}{n} \sum_{i=1}^n f(x_i) - f(x_{i-1}) = \frac{\alpha(b) - \alpha(a)}{n} [f(b) - f(a)]$$

Where in the last equality we use the fact that the sum telescopes. If  $\alpha(b) = \alpha(a)$ , then  $U(P) - L(P) = 0$  and the claim immediately follows. If  $\alpha(b) > \alpha(a)$ , then the claim follows by choosing  $n > \frac{1}{\epsilon} \frac{f(b) - f(a)}{\alpha(b) - \alpha(a)}$  (that is, selecting a partition  $P$  with sufficiently large  $n$ ), from which it follows that:

$$U(P) - L(P) < \epsilon$$

and hence  $f \in \mathcal{R}_\alpha[a, b]$  by Theorem 6.6. □



Figure 30: Visualization of the idea of the proof of Theorem 6.9. We chop up  $[\alpha(a), \alpha(b)]$  into equal sized subintervals.

### Theorem 6.10

Suppose  $f : [a, b] \mapsto \mathbb{R}$  is bounded and has finitely many discontinuities. Suppose  $\alpha$  is continuous at every point where  $f$  is not. Then,  $f \in \mathcal{R}_\alpha[a, b]$ .

### Proof

As has become standard, we will be applying Theorem 6.6. We have that:

$$U(P) - L(P) = \sum_{i=1}^n (M_i - m_i) \Delta \alpha_i$$

Let  $\epsilon > 0$ , and let  $E = \{e_1, \dots, e_k\}$  be the set of points where  $f$  is not continuous.  $\alpha$  is continuous at each  $e_j$  by hypothesis. Therefore, there exists disjoint intervals  $(u_j, v_j)$  that cover points in  $E$ , such that  $u_j < e_j < v_j$  and  $\alpha(v_j) - \alpha(u_j) < \epsilon$  (by the continuity of  $\alpha$ ). Let  $K = [a, b] \cap \left( \bigcup_{j=1}^k (u_j, v_j) \right)^c$ .  $K$  is a compact set as it is a finite union of closed intervals.  $f$  is continuous on  $K$  by hypothesis, so  $f$  is uniformly continuous on  $K$  by Theorem 4.19. Hence, there exists  $\delta > 0$  such that for  $s, t \in K$ ,  $|s - t| < \delta \implies |f(s) - f(t)| < \epsilon$ . We then form a partition  $P = \{x_1, \dots, x_n\}$  to consist of  $\{u_1, v_1, \dots, u_k, v_k\}$  and additional points in  $K$  with  $\delta x_i < \delta$ . For such  $i$ ,  $f$  is continuous on the subinterval and hence  $M_i - m_i < \epsilon$ . For the other intervals  $[u_j, v_j]$ , we have that  $M_j - m_j \leq 2M$  where  $M = \sup \{ |f(x)| : x \in [a, b] \}$  and that  $\Delta \alpha_j < \epsilon$ . Therefore, we have that:

$$\begin{aligned} 0 \leq U(P) - L(P) &= \sum_{i=1}^n (M_i - m_i) \Delta \alpha_i \\ &\leq 2M\epsilon + \epsilon (\alpha(b) - \alpha(a)) \end{aligned}$$

Where the first term comes from the  $[u_j, v_j]$ s and the second term is the maximum possible value from the subintervals of  $K$ . By choosing  $\epsilon$  small enough, the RHS is small as desired for some choice of  $P$ . Hence,  $U(P) - L(P) < \epsilon$  for some  $P$  and hence  $f \in \mathcal{R}_\alpha[a, b]$ .  $\square$

A natural question that arises is “what if  $f$  and  $\alpha$  are discontinuous at the same point?” In this case, we can construct functions such that  $f \notin \mathcal{R}_\alpha[a, b]$  (see HW1Q5).



Figure 31: Visualization of how  $[a, b]$  gets split up in the proof of Theorem 6.10.  $K$  consists of the union of the closed intervals marked in blue.

### Theorem 6.11

If  $f \in \mathcal{R}_\alpha[a, b]$ ,  $m \leq f(x) \leq M$  for all  $x \in [a, b]$ , and  $\phi$  is continuous on  $[m, M]$ , then  $\phi \circ f \in \mathcal{R}_\alpha[a, b]$ .

### Proof

Not covered in lecture, see Rudin. □

As an example of the above theorem, suppose we have that  $f \in \mathcal{R}(\alpha)$ . Then, we have that  $f^2 \in \mathcal{R}(\alpha)$  and  $|f| \in \mathcal{R}(\alpha)$  as  $\phi(x) = x^2$  and  $\phi(x) = |x|$  are both continuous functions.

## 6.3 Properties of the Integral

### Theorem 6.12: Linearity and Related Properties

- (a) Suppose  $f_1 \in \mathcal{R}_\alpha[a, b]$  and  $f_2 \in \mathcal{R}_\alpha[a, b]$ . Then,  $f_1 + f_2 \in \mathcal{R}_\alpha[a, b]$  and  $cf_1 \in \mathcal{R}_{[\alpha]}(a)$  for all  $c \in \mathbb{R}$ . Furthermore:

$$\int_a^b (f_1 + f_2) d\alpha = \int_a^b f_1 d\alpha + \int_a^b f_2 d\alpha \text{ and } \int_a^b cf_1 d\alpha = c \int_a^b f_1 d\alpha$$

- (b) Suppose  $f_1, f_2 \in \mathcal{R}_\alpha[a, b]$  and  $f_1(x) \leq f_2(x)$  for all  $x \in [a, b]$ . Then,

$$\int_a^b f_1 d\alpha \leq \int_a^b f_2 d\alpha$$

- (c) If  $f \in \mathcal{R}_\alpha[a, b]$  and  $c \in (a, b)$ , then  $f \in \mathcal{R}_\alpha[a, c] \cap \mathcal{R}_\alpha[c, b]$  and furthermore:

$$\int_a^b f d\alpha = \int_a^c f d\alpha + \int_c^b f d\alpha$$

- (d) If  $f \in \mathcal{R}_\alpha[a, b]$  and  $|f(x)| \leq M$  for all  $x \in [a, b]$ , then:

$$\left| \int_a^b f d\alpha \right| \leq M(\alpha(b) - \alpha(a))$$

- (e) If  $f \in \mathcal{R}_{\alpha_1}[a, b]$  and  $f \in \mathcal{R}_{\alpha_2}[a, b]$ , then  $f \in \mathcal{R}_{\alpha_1 + \alpha_2}[a, b]$  and  $f \in \mathcal{R}_{c\alpha_1}[a, b]$  for all  $c \geq 0$ . Furthermore:

$$\int_a^b f d\alpha_1 + \int_a^b f d\alpha_2 = \int_a^b f d(\alpha_1 + \alpha_2) \text{ and } \int_a^b f d(c\alpha_1) = c \int_a^b f d\alpha_1$$

## Proof

(a) See Rudin and HW2Q1.

(b) We have that  $L(P, f_1) \leq L(P, f_2)$  for all partitions  $P$  as  $\inf \{f_1(x) : x \in [x_{i-1}, x_i]\} \leq \inf \{f_2(x) : x \in [x_{i-1}, x_i]\}$  for all  $i$  (as  $f_1(x) \leq f_2(x)$ ). Therefore, we have that:

$$\int_a^b f_1 d\alpha = \sup_P L(P, f_1) \leq \sup_P L(P, f_2) = \int_a^b f_2 d\alpha.$$

(c) Exercise (see HW2Q1 for the  $\overline{\int_a^b}$  case)

(d) Consider any partition  $P$  of  $[a, b]$ . Let  $M = \sup \{|f(x)| : x \in [a, b]\}$ . We then have that (using the ML bound from Theorem 6.1):

$$\begin{aligned} -M[\alpha(b) - \alpha(a)] &= \sum_{i=1}^n (-M)\Delta\alpha_i \leq \sum_{i=1}^n M_i\Delta\alpha_i = L(P, f, \alpha) \leq \int_a^b f d\alpha \\ &\leq U(P, f, \alpha) = \sum_{i=1}^n M_i\Delta\alpha_i \leq \sum_{i=1}^n M\Delta\alpha_i = M[\alpha(b) - \alpha(a)]. \end{aligned}$$

(e) We prove the first (additivity) statement. Let  $\epsilon > 0$ . Choose  $P_i, i \in \{1, 2\}$  (By Theorem 6.6) such that

$$U(P_i, f, \alpha_i) - L(P_i, f, \alpha_i) < \frac{\epsilon}{2}$$

Let  $P = P_1 \cup P_2$ . Then, by Theorem 6.4, we have that:

$$U(P^*, f, \alpha_i) - L(P^*, f, \alpha_i) < \frac{\epsilon}{2}$$

Adding the two expressions (for  $i = 1, 2$ ) together, we have:

$$U(P^*, f, \alpha_1 + \alpha_2) - L(P^*, f, \alpha_1 + \alpha_2) < \epsilon$$

So  $f \in \mathcal{R}_{\alpha_1 + \alpha_2}[a, b]$  by Theorem 6.6. Furthermore, We have that:

$$\int_a^b f d(\alpha_1 + \alpha_2) \leq U(P^*, f, \alpha_1 + \alpha_2) = U(P^*, f, \alpha_1) + U(P^*, f, \alpha_2) < \int_a^b f d\alpha_1 + \int_a^b f d\alpha_2 + \epsilon$$

where we apply the inequality of  $U(P^*, f, \alpha_i) < \int_a^b f d\alpha_i + \frac{\epsilon}{2}$  in the last line. Similarly, we have that:

$$\int_a^b f d(\alpha_1 + \alpha_2) \geq L(P^*, f, \alpha_1 + \alpha_2) = L(P^*, f, \alpha_1) + L(P, f, \alpha_2) > \int_a^b f d\alpha_1 + \int_a^b f d\alpha_2 - \epsilon$$

Since  $\epsilon$  is arbitrary, we therefore conclude that:

$$\int_a^b f d(\alpha_1 + \alpha_2) = \int_a^b f d\alpha_1 + \int_a^b f d\alpha_2$$

□

**Theorem 6.13**

- (a) Suppose  $f, g \in \mathcal{R}(\alpha)$ . Then,  $fg \in \mathcal{R}(\alpha)$ .
- (b) Suppose  $f \in \mathcal{R}(\alpha)$ . Then,  $|f| \in \mathcal{R}(\alpha)$  and  $\left| \int_a^b f d\alpha \right| \leq \int_a^b |f| d\alpha$ .

**Proof**

- (a) By Theorem 6.12(a),  $f \pm g \in \mathcal{R}(\alpha)$ , so by Theorem 6.11,  $(f \pm g)^2 \in \mathcal{R}(\alpha)$ . Since  $(f + g)^2 - (f - g)^2 = 4fg$ , we then have that:

$$\frac{(f + g)^2 + (f - g)^2}{4} = fg \in \mathcal{R}(\alpha)$$

- (b) By Theorem 6.11,  $|f| \in \mathcal{R}(\alpha)$  letting  $\phi(t) = |t|$ . Let  $c = \operatorname{sgn} \left( \int_a^b f d\alpha \right) \in \{-1, 0, 1\}$ . Then, we have that:

$$\left| \int_a^b f d\alpha \right| = c \int_a^b d\alpha = \int_a^b c f d\alpha \leq \int_a^b |f| d\alpha$$

Where we use Theorem 6.12(a) in the second equality, and Theorem 6.12(b) in the last inequality (as  $cf \leq |f|$ ).  $\square$

**Theorem 6.15**

Suppose  $f$  is bounded on  $[a, b]$ ,  $s \in (a, b)$ , and  $f$  is continuous at  $s$ . Let:

$$\alpha(x) = \begin{cases} 0 & x \leq s \\ 1 & x > s \end{cases}$$

Then,  $\int_a^b f d\alpha = f(s)$  (and in particular,  $f \in \mathcal{R}(\alpha)$ ).

This result is interesting, and some remarks on the nature of this theorem are in order. Firstly, by Theorem 4.29 (not covered in Lecture, see Rudin), since  $\alpha$  is monotonically increasing, we have that  $\lim_{t \rightarrow x^+} \alpha(t) = \alpha(x^+)$  and  $\lim_{t \rightarrow x^-} \alpha(t) = \alpha(x^-)$  exist at every  $x \in (a, b)$  and  $\alpha(x^-) \leq \alpha(x) \leq \alpha(x^+)$ . Secondly, note that Rudin defines  $\alpha$  in the above Theorem to be left continuous, but in probability theory, it is conventional to use a right continuous (i.e.  $\alpha(x) = \alpha(x^+)$  for all  $x \in (a, b)$ ). We leave it as an exercise to prove the theorem for the case of a right continuous  $\alpha$  (the strategy and result are identical).

Any physicists looking at the result of the Theorem will note that it looks very much like the “Dirac Delta” function; indeed, this choice of  $\alpha$  is making rigorous the notion of a  $\delta$  function where:

$$\int_a^b f(x) \delta(x - s) = f(s)$$

We leave it as a homework exercise (HW2Q2) to prove that no such function can actually exist. This step-function method is one way to make the  $\delta$  function well-defined; other methods include taking the limit of a bell curve, or bringing in the theory of distributions (the latter which is most definitely outside the scope of this course).

Finally, we note that this example really shows off something that the Riemann integral cannot reproduce; the incorporation of “discrete” points.



Figure 32: Visualization of Theorem 4.29. For any strictly increasing function, we have that  $\alpha(x^-) \leq \alpha(x) \leq \alpha(x^+)$  at a discontinuity  $x$ .

#### Proof

Choose  $P = \{x_0, x_1 = s, x_2, x_3 = b\}$ . We then have that:

$$U(P) = \sum_{i=1}^3 M_i \Delta \alpha_i = M_2 \Delta \alpha_2 = M_2 = \sup \{f(x) : x \in [x_1, x_2]\}$$

$$L(P) = \sum_{i=1}^3 m_i \Delta \alpha_i = m_2 \Delta \alpha_2 = m_2$$

So, we have that  $m_2 \leq \int_a^b f d\alpha \leq M_2$  (assuming the integral exists). Taking the limit as  $x_2 \rightarrow s^+$ , by the continuity of  $f$  at  $s$  we have that  $M_2 \rightarrow f(s)$  and  $m_2 \rightarrow f(s)$ . Therefore,  $\int_a^b f d\alpha$  exists and equals  $f(s)$ .  $\square$

Note that the above proof works exactly the same way if  $s = a$ . If  $s = b$ , then because we take  $\alpha$  to be left continuous,  $\alpha = 0$  everywhere on  $[a, b]$  and the integral just equals zero.

#### Definition 6.14: Unit Step Function

We define the (left continuous) **unit step function** as:

$$I(x) = \begin{cases} 0 & x \leq 0 \\ 1 & x > 0 \end{cases}$$

In Theorem 6.15, we have that  $\alpha(x) = I(x - s)$ .

#### Theorem 6.16

Let  $c_n \geq 0$  be such that  $\sum_{n=1}^{\infty} c_n < \infty$ . Take  $s_n \in (a, b)$  distinct (that is,  $s_n \neq s_m$  if  $n \neq m$ ). Define  $\alpha(x) = \sum_{n=1}^{\infty} c_n I(x - s_n)$  (which converges/exists by comparison, as  $\sum c_n$  converges). Let  $f$  be continuous on  $[a, b]$ . Then,

$$\int_a^b f d\alpha = \sum_{n=1}^{\infty} c_n f(s_n)$$





Figure 33: Plot of the (left continuous) unit step function. To obtain the  $\alpha$  used in Theorem 6.15, move the step to  $x = s$  instead of  $x = 0$ .

Note that the resultant series in the theorem above is convergent/well defined as  $f$  is bounded on  $[a, b]$  and hence  $\sum_{n=1}^{\infty} c_n f(s_n) \leq M \sum_{n=1}^{\infty} c_n$  where  $M = \max \{f(x)\}$ . Hence the series converges by comparison. If all the  $c_n$ s are zero except for one, this is just Theorem 6.15. Note that the above result holds for a finite sum as well (we would use induction to prove it in this case).

#### Proof

Let  $R_N = \sum_a^b f d\alpha - \sum_{n=1}^N c_n f(s_n)$ . We show that  $R_N \rightarrow 0$  as  $N \rightarrow \infty$  (i.e. given  $\epsilon > 0$ , we show there exists  $N_0$  such that  $|R_N| < \epsilon$  if  $N \geq N_0$ ). We define:

$$\alpha_1(x) = \sum_{n=1}^N c_n I(x - s_n) \quad \alpha_2(x) = \sum_{n=N+1}^{\infty} c_n I(x - s_n)$$

Then, by Theorem 6.12(e) we have that:

$$\int_a^b f d\alpha = \int_a^b f d\alpha_1 + \int_a^b f d\alpha_2$$

Then, we have that:

$$\begin{aligned} \int_a^b d\alpha_1 &= \sum_{i=1}^N \int_a^b f(x) d(c_n I(x - s_n)) \\ &= \sum_{n=1}^N c_n \int_a^b f(x) d(I(x - s_n)) \\ &= \sum_{n=1}^N c_n f(s_n) \end{aligned}$$

Wherein the first two equalities we again apply Theorem 6.12(e) and in the last equality we use Theorem 6.15. We therefore obtain that  $R_N = \int_a^b f d\alpha_2$ . We then have that:

$$|R_N| \leq M [\alpha_2(b) - \alpha_2(a)] = M \sum_{n=N+1}^{\infty} c_n$$

Then by the convergence of  $\sum c_n$ , we can choose  $N_0$  such that  $\sum_{n=N_0+1}^{\infty} c_n < \frac{\epsilon}{M}$ . We therefore have that  $R_N < \epsilon$  if  $N \geq N_0$ , proving the claim.  $\square$

**Theorem 6.17**

Suppose:

- (i)  $|f(x)| \leq M$  for all  $x \in [a, b]$ .
- (ii)  $\alpha$  is continuous and increasing on  $[a, b]$  and differentiable on  $(a, b)$ .
- (iii)  $\alpha' \in \mathcal{R}[a, b]$ .

Then,  $f \in \mathcal{R}_\alpha[a, b]$  if and only if  $f\alpha' \in \mathcal{R}[a, b]$  and in this case:

$$\int_a^b f d\alpha = \int_a^b f(x)\alpha'(x)dx$$

**Proof**

It suffices to show that:

$$\overline{\int_a^b f d\alpha} = \overline{\int_a^b f\alpha' dx} \text{ and } \underline{\int_a^b f d\alpha} = \underline{\int_a^b f\alpha' dx}$$

We prove the first equality and the second equality follows analogously. Let  $\epsilon > 0$ . Since  $\alpha' \in \mathcal{R}$ , there exists a partition  $P$  such that  $U(P, \alpha') - L(P, \alpha') < \epsilon$  (this also holds for any refinement of  $P$ ). Letting  $A_i = \sup \alpha', a_i = \inf \alpha'$  on  $[x_{i-1}, x_i]$  we have that:

$$\sum_{i=1}^n (A_i - a_i) \Delta x_i < \epsilon$$

By the Mean Value Theorem (Theorem 5.10) we have that there exists  $t_i \in [x_{i-1}, x_i]$  such that  $\Delta \alpha_i = \alpha'(t_i) \Delta x_i$ . We also have that for all  $s_i \in [x_{i-1}, x_i]$ ,  $|\alpha'(s_i) - \alpha'(t_i)| \leq A_i - a_i$  ( $\alpha'$  can only vary as much as the maximum minus the minimum on the interval). We therefore have that:

$$\sum_{i=1}^n |\alpha'(s_i) - \alpha'(t_i)| \Delta x_i \leq \sum_{i=1}^n (A_i - a_i) \Delta x_i < \epsilon$$

Recall that we skipped Theorem 6.7, but it might be worth comparing this result to Theorem 6.7(c). Let  $M = \sup f(x) : x \in [a, b]$ . Now, for any choice of  $s_i \in [x_{i-1}, x_i]$ , we have that:

$$\left| \sum_{i=1}^n f(s_i) \Delta \alpha_i - \sum_{i=1}^n \alpha'(s_i) \Delta x_i \right| \leq \sum_{i=1}^n M |\alpha'(t_i) - \alpha'(s_i)| \Delta x_i < M\epsilon \quad (*)$$

Therefore:

$$\sum_{i=1}^n f(s_i) \Delta \alpha_i \leq \sum_{i=1}^n f(s_i) \alpha'(s_i) \Delta x_i + M\epsilon \leq U(P, f\alpha') + M\epsilon$$

We now take the supremum over each  $[s_{i-1}, s_i]$ . Since  $s_i$  is arbitrary, we have that  $\overline{\int_a^b f d\alpha} \leq U(P, f, \alpha) \leq U(P, f\alpha') + M\epsilon$ , and hence  $\overline{\int_a^b f d\alpha} \leq \overline{\int_a^b f\alpha' dx} + M\epsilon$ . (\*) gives  $\overline{\int_a^b f\alpha' dx} \leq \overline{\int_a^b f d\alpha} + M\epsilon$ . Since  $\epsilon$  is arbitrary, we have that  $\overline{\int_a^b f d\alpha} \leq \overline{\int_a^b f\alpha' dx}$  and  $\overline{\int_a^b f\alpha' dx} \leq \overline{\int_a^b f d\alpha}$ , so  $\overline{\int_a^b f\alpha' dx} = \overline{\int_a^b f d\alpha}$ .  $\square$

This theorem tells us something that looks very familiar from Calculus... namely, a change of variables!

Recall the COV formula from first year:

$$\int_a^b f(x)dx = \int_{\phi^{-1}(a)}^{\phi^{-1}(b)} f(\phi(y))\phi'(y)dy$$

Where  $x = \phi(y), y = \phi^{-1}(y)$ . We now prove this notion rigorously.

**Theorem 6.19: Change of Variable**

Let  $f : [a, b] \mapsto \mathbb{R}$ . Suppose  $\phi : [A, B] \mapsto [a, b]$  is continuous and strictly increasing. Suppose  $\alpha$  is increasing on  $[a, b]$  and  $f \in \mathcal{R}_\alpha[a, b]$ . Let  $g = f \circ \phi : [A, B] \mapsto \mathbb{R}$  and  $\beta = \alpha \circ \phi : [A, B] \mapsto \mathbb{R}$ . Then, we have that  $g \in \mathcal{R}_\beta[A, B]$ , and:

$$\int_A^B g d\beta = \int_a^b f d\alpha$$

As an example, consider the integral  $\int_a^b \sin x^2 dx$  for  $0 \leq a < b$ . We then have that  $f(x) = \sin x^2$  and  $\alpha(x) = x$ . We make the substitution  $x^2 = y$ , so  $\phi(x) = \sqrt{x}$  and  $\phi^{-1}(y) = y^2$ . Then,  $A = \phi^{-1}(a) = a^2$ ,  $B = \phi^{-1}(b) = b^2$ . We then have that  $g(y) = f \circ \phi(y) = f(\sqrt{y}) = \sin(y)$ , and  $\beta(y) = \alpha \circ \phi(y) = \alpha(\sqrt{y}) = \sqrt{y}$ . We then have that:

$$\int_a^b \sin x^2 dx = \int_{a^2}^{b^2} \sin y d\beta = \int_{a^2}^{b^2} \frac{\sin y}{2\sqrt{y}} dy$$

Where the first equality follows by Theorem 6.19 and the second equality follows by Theorem 6.17.



Figure 34: Plot of a (monotonically increasing and continuous) function  $\phi$  and a demonstration of how it puts partitions of  $[a, b]$  and  $[A, B]$  in one-to-one correspondence. This gives the intuition for the proof of Theorem 6.19.

### Proof

We make the observation that partitions  $P$  of  $[a, b]$  and partitions  $Q$  of  $[A, B]$  can be put in 1-1 correspondence via  $x_i = \phi(y_i)$ . Additionally, we make the observation that the set of  $g$  values on  $[y_{i-1}, y_i]$  is equal to the set of  $f$  values on  $[x_{i-1}, x_i]$ . Finally, we observe that  $\alpha(x_i) = (\alpha \circ \phi)(y_i) = \beta(y_i)$ . With these three observations, we have that:

$$U(P, f, \alpha) = U(Q, g, \beta) \text{ and } L(P, f, \alpha) = L(Q, g, \beta)$$

Let  $\epsilon > 0$ . Since  $f \in \mathcal{R}_\alpha[a, b]$ , there exists a  $P$  such that  $U(P, f, \alpha) - L(P, f, \alpha) < \epsilon$  by Theorem 6.6. For this partition, we have that  $U(Q, g, \beta) - L(Q, g, \beta) < \epsilon$  for the corresponding partition  $Q$ . Hence,  $g \in \mathcal{R}_\beta[A, B]$ . Finally, we have that:

$$\int_A^B g d\beta = \inf_Q U(Q, g, \beta) = \inf_P U(P, f, \alpha) = \int_a^b f d\alpha$$

□

## 6.4 The Fundamental Theorem of Calculus

### Theorem 6.20: Fundamental Theorem of Calculus I

Let  $f \in \mathcal{R}[a, b]$  and for  $x \in [a, b]$ , define  $F(x) = \int_a^x f(t) dt$ . Then,  $F$  is continuous on  $[a, b]$ . If  $f$  is continuous at  $x_0 \in [a, b]$  then  $F'(x_0)$  exists and  $F'(x_0) = f(x_0)$ .




Figure 35: The Fundamental Theorem of Calculus/Theorem 6.20 gives us a way to relate a curve ( $f$ ) with the cumulative area beneath it ( $F$ ) by means of the derivative.

To get intuition for Theorem 6.20, we think about what happens when we “zoom in” to a function  $f$ . Over an interval  $(x_0, x_0 + h)$ , if  $x_0$  is continuous at  $x_0$  and  $h$  is small, then  $f$  will be roughly constant over the interval. Hence,  $F(x_0 + h) - F(x_0) \approx f(x_0)h$ . In the limit of  $h \rightarrow 0$ , this approximation becomes exact.



Figure 36: Visual Intuition for Theorem 6.20 and its proof. As  $f$  is continuous at  $x_0$ , for small  $h$ ,  $f$  is roughly constant on  $(x_0, x_0 + h)$ . Hence, the increase in cumulative area below the curve is roughly  $f(x_0) \cdot h$  (height of  $f(x_0)$  times width  $h$  of the approximate rectangle).

### Proof

We first show the continuity of  $F$ . Choose  $M$  such that  $|f(t)| \leq M$  for all  $t \in [a, b]$ . For  $a \leq x < y \leq b$ , we then have that:

$$|F(x) - F(y)| = \left| \int_a^x f(t)dt - \int_a^y f(t)dt \right| = \left| \int_x^y f(t)dt \right| \leq M(y - x)$$

Where we use Theorem 6.12(c) for the second equality and Theorem 6.12(d) for the inequality at the end. Let  $\epsilon > 0$ . If  $|x - y| < \delta = \frac{\epsilon}{M}$ , then  $|F(x) - F(y)| < \epsilon$  so  $F$  is continuous.

We next show the differentiability of  $F$ . We wish to show that:

$$\lim_{h \rightarrow 0} \frac{F(x_0 + h) - F(x_0)}{h} = f(x_0)$$

We show the case for  $h > 0$ , that is, taking the limit of  $h \rightarrow 0^+$ . We have that:

$$\frac{1}{h} [F(x_0 + h) - F(x_0)] - f(x_0) = \frac{1}{h} \int_{x_0}^{x_0+h} f(t)dt - f(x_0) = \frac{1}{h} \int_{x_0}^{x_0+h} [f(t) - f(x_0)] dt$$

Where in the last line we use the observation that  $\frac{1}{h} \int_{x_0}^{x_0+h} f(x_0)dt = f(x_0)$ . Since  $f$  is continuous at  $x_0$ , for any  $\epsilon > 0$ , there exists  $\delta > 0$  such that:

$$|t - x_0| < \delta \implies |f(t) - f(x_0)| < \epsilon$$

Thus, if  $h < \delta$ , then:

$$\left| \frac{1}{h} [F(x_0 + h) - F(x_0)] - f(x_0) \right| \leq \frac{1}{h} \int_{x_0}^{x_0+h} |f(t) - f(x_0)| dt < \frac{1}{h} \epsilon h = \epsilon$$

So we conclude that:

$$\lim_{h \rightarrow 0^+} \frac{F(x_0 + h) - F(x_0)}{h} = F'(x_0) = f(x_0)$$

□

**Theorem 6.21: Fundamental Theorem of Calculus II**

If  $f \in \mathcal{R}[a, b]$  and there exists  $F$  on  $[a, b]$  such that  $F' = f$ , then:

$$\int_a^b f(x)dx = F(b) - F(a)$$

**Proof**

For any partition  $P$ , we have that

$$\begin{aligned} F(b) - F(a) &= \sum_{i=1}^n [F(x_i) - F(x_{i-1})] \\ &= \sum_{i=1}^n F'(t_i) \Delta x_i \\ &= \sum_{i=1}^n f(t_i) \Delta x_i \end{aligned}$$

Where the first equality follows by the fact that the sum telescopes, the second equality follows from the Mean Value Theorem/Theorem 5.10 (By the continuity of  $F$ , there exists  $t_i \in [x_{i-1}, x_i]$  such that  $F(x_i) - F(x_{i-1}) = F'(t_i)(x_i - x_{i-1})$ ) and the third equality follows by Theorem 6.20. Now, we have that  $f(t_i) \in [m_i, M_i]$  for each  $i$ , so we therefore have that:

$$\sum_{i=1}^n m_i \Delta x_i \leq \sum_{i=1}^n f(t_i) \Delta x_i \leq \sum_{i=1}^n M_i \Delta x_i$$

Hence:

$$F(b) - F(a) \in [L(P, f), U(P, f)]$$

Additionally, by definition of the integral we have that:

$$\int_a^b f(x)dx \in [L(P, f), U(P, f)]$$

Let  $\epsilon > 0$ . Choose  $P$  such that  $U(P, f) - L(P, f) < \epsilon$ . Then, we have that:

$$\left| F(b) - F(a) - \int_a^b f(x)dx \right| < \epsilon$$

Since  $\epsilon$  is arbitrary, we have that:

$$F(b) - F(a) = \int_a^b f(x)dx$$

□

### Theorem 6.22: Integration By Parts

If  $F, G$  are differentiable on  $[a, b]$  with  $F' = f \in \mathcal{R}[a, b]$  and  $G' = g \in \mathcal{R}[a, b]$ , then:

$$\int_a^b F(x)g(x)dx = F(b)G(b) - F(a)G(a) - \int_a^b f(x)G(x)dx$$

Note that the above integration by parts formula generalizes to different  $\alpha$  (not just  $\alpha(x) = x$ ). The proof of this is left as an exercise (Rudin Chapter 6 Problem 17).

#### Proof

Let  $H(x) = F(x)G(x)$ . Then,  $H'(x) = F'(x)G(x) + F(x)G'(x) = f(x)G(x) + F(x)g(x) \in \mathcal{R}[a, b]$  by the product rule (Theorem 5.3) and the FTC I (Theorem 6.20). Therefore, we have that:

$$H(b) - H(a) = \int_a^b H'(x)dx = \int_a^b f(x)G(x)dx + \int_a^b F(x)g(x)dx$$

Where the first equality follows by FTC II (Theorem 6.21). We have that  $H(b) - H(a) = F(b)G(b) - F(a)G(a)$ , so rearranging the above expression, we find that:

$$\int_a^b F(x)g(x)dx = F(b)G(b) - F(a)G(a) - \int_a^b f(x)G(x)dx$$

Which is the desired expression. □

### Definition: Improper Integrals

If  $f \in \mathcal{R}[a, b]$  for all  $b > a$ , then we define:

$$\int_a^\infty f(x)dx = \lim_{b \rightarrow \infty} \int_a^b f(x)dx$$

if the integral exists in  $\mathbb{R}$ . Then, we say that the **improper integral**  $\int_a^\infty f(x)dx$  converges.

### Definition: Absolute Convergence of Integrals

If  $\int_a^\infty |f(x)|dx$  converges, then we say that  $\int_a^\infty f(x)dx$  **converges absolutely**.

To finish off this chapter, we will work through a comprehensive problem together; we prove that  $\int_0^\infty \sin t^2 dt$  converges but not absolutely.

*Proof.* The proof that the integral does not converge absolutely is left as a homework exercise (HW3Q1). We will prove that the integral converges here. Let  $p_n = \int_0^n \sin t^2 dt$  where  $n \in \mathbb{N}$ . We will show that:

- (i)  $\{p_n\}$  is a Cauchy sequence and hence has a limit in  $\mathbb{R}$ .
- (ii) This is enough to show that the improper integral converges.

We start by showing (i), namely that  $\{p_n\}$  is Cauchy. For  $x < y$ , we have that:

$$\int_x^y \sin t^2 dt = \int_{x^2}^{y^2} \sin u \frac{1}{2\sqrt{u}} du$$

by a change of variable (Theorem 6.19). Now performing an integration by parts (Theorem 6.22) with  $F(u) = \frac{1}{2\sqrt{u}}$  and  $G'(u)du = \sin u du$ , we have that  $F'(u) = -\frac{1}{4u^{3/2}}$  and  $G(u) = -\cos u$ , so:

$$\int_x^y \sin t^2 dt = \int_{x^2}^{y^2} \sin u \frac{1}{2\sqrt{u}} du = -\frac{\cos u}{2\sqrt{u}} \Big|_{x^2}^{y^2} - \int_{x^2}^{y^2} \frac{\cos u}{4u^{3/2}} du$$

Hence:

$$\int_x^y \sin t^2 dt = \frac{\cos x^2}{2x} - \frac{\cos y^2}{2y} - \int_{x^2}^{y^2} \frac{\cos u}{4u^{3/2}} du$$

Suppose  $n \geq m$ . Then, we have that:

$$\begin{aligned} |p_n - p_m| &= \left| \int_m^n \sin t^2 dt \right| \\ &= \left| \frac{\cos m^2}{2m} - \frac{\cos n^2}{2n} - \int_{m^2}^{n^2} \frac{\cos u}{4u^{3/2}} du \right| \end{aligned}$$

Using the fact that  $|\cos(x)| \leq 1$  and applying the Triangle inequality, we have that:

$$|p_n - p_m| \leq \frac{1}{2m} + \frac{1}{2n} - \int_{m^2}^{n^2} \frac{1}{4u^{3/2}} du = \frac{1}{2m} + \frac{1}{2n} + \frac{-1}{2u^{1/2}} \Big|_{m^2}^{n^2} = \frac{1}{2m} + \frac{1}{2n} - \frac{1}{2n} + \frac{1}{2m} = \frac{1}{m} \quad (*)$$

Let  $\epsilon > 0$ . Choose  $N_0$  such that  $\frac{1}{N_0} < \epsilon$ . Then, for  $m \geq n \geq N_0$  we have that:

$$|p_n - p_m| \leq \frac{1}{m} \leq \frac{1}{N_0} < \epsilon$$

and we conclude that  $\{p_n\}$  is Cauchy. Since  $\mathbb{R}$  is complete,  $p_n \rightarrow p$  for some  $p \in \mathbb{R}$ . To finish the proof, we show (ii); that this is sufficient. For  $b \geq N_0$ , choose  $N \geq N_0$  such that  $b \in [N, N+1)$ . Then, we have that:

$$\int_0^b \sin t^2 dt = p_N + \int_N^b \sin t^2 dt \leq p_N + \frac{1}{N}$$

Where the inequality follows by (\*). We therefore have that:

$$\left| p - \int_0^b \sin t^2 dt \right| \leq |p - p_N| + \left| \int_N^b \sin t^2 dt \right| < \epsilon + \frac{1}{N} \leq \epsilon + \frac{1}{N_0} < \epsilon + \epsilon = 2\epsilon$$

Since  $\epsilon$  is arbitrary, we conclude that  $\lim_{b \rightarrow \infty} \int_0^b \sin t^2 dt = p$  and hence the improper integral converges.  $\square$



## 7 Sequences and Series of Functions

### 7.1 Motivating Examples

#### Example

For  $m, n \in \mathbb{N}$ , let  $p_{n,m} = \frac{m}{n}$ . Then,

$$\lim_{m \rightarrow \infty} p_{m,n} = \infty, \quad \lim_{n \rightarrow \infty} p_{m,n} = 0$$

In particular,

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} p_{m,n} = 0, \quad \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} p_{m,n} = \infty.$$

Which demonstrates that the order of which limits are taken in can affect the value.

#### Example

Define the sequence of functions:

$$f_n(x) = \begin{cases} 1 & x \geq 0 \\ 1 + nx & -\frac{1}{n} < x < 0 \\ 0 & x \leq -\frac{1}{n} \end{cases}$$

Since  $f_n$  is piecewise linear, it is continuous. However, looking at the  $n \rightarrow \infty$  limit, we have:

$$\lim_{n \rightarrow \infty} f_n(x) = \begin{cases} 1 & x \geq 0 \\ 0 & x < 0 \end{cases}$$

Which is the right continuous step function, which is evidently discontinuous at  $x = 0$ . Hence, the limit of continuous functions can be discontinuous. Another way of viewing this problem is:

$$\lim_{n \rightarrow \infty} \lim_{x \rightarrow 0} f_n(x) = 0, \quad \lim_{x \rightarrow 0} \lim_{n \rightarrow \infty} f_n(x) = \text{D.N.E.}$$

so again we see the order of taking our limits can be important.



Figure 37: Plot of  $f_n$  in the above example.

### Example 7.4

For  $m \in \mathbb{N}$  and  $x \in \mathbb{R}$ , let  $f_m(x) = \lim_{n \rightarrow \infty} [\cos(m! \pi x)]^{2n}$ . Since  $|\cos(k\pi)| = 1$  if  $k \in \mathbb{Z}$ , we see that  $f_m(x) = 1$  when  $m!x \in \mathbb{Z}$ . Conversely, since  $|\cos(k\pi)| < 1$  if  $k \notin \mathbb{Z}$ ,  $f_m(x) = 0$  when  $m!x \notin \mathbb{Z}$ . Some plots of  $f_m(x)$  on  $[0, 1]$  for  $m = 1, 2, 3$  are below as a visualization. We now define  $f(x) = \lim_{m \rightarrow \infty} f_m(x)$ . If  $x = \frac{p}{q} \in \mathbb{Q}$ , then  $m!x = \frac{m!p}{q} \in \mathbb{Z}$  for  $m$  large enough (for  $m \geq q$ , as the denominator cancels). Therefore, we have that  $f(x) = 1$  for  $x \in \mathbb{Q}$ . Conversely, if  $x \notin \mathbb{Q}$ , then  $m!x \notin \mathbb{Z}$  for all  $m \in \mathbb{N}$ . So,  $f_m(x) = 0$  for all  $m$ , and  $f(x) = 0$ . Therefore, we have that:

$$f(x) = \lim_{m \rightarrow \infty} f_m(x) = \begin{cases} 1 & x \in \mathbb{Q} \\ 0 & x \notin \mathbb{Q} \end{cases}.$$

In other words,  $f$  is the Dirichlet function. The interesting part is that each of the  $f_m(x)$  are Riemann integrable on  $[0, 1]$  by Theorem 6.10 (as  $f$  has finitely many discontinuities for any  $m \in \mathbb{N}$ ). However, the limit is not Riemann integrable, as we prove below. Hence, the limit of Riemann integrable functions is not necessarily Riemann integrable.



Figure 38: Plot of  $f_m(x)$  over the interval  $[0, 1]$  for  $m = 1, 2, 3$ . For  $m = 1$ , only  $x = 0, 1$  satisfy  $m!x = x \in \mathbb{Z}$ . For  $m = 2$ , we have that  $x = 0, \frac{1}{2}, 1$  satisfy  $m!x = 2x \in \mathbb{Z}$ . Finally, for  $m = 3$ , we have that  $x = 0, \frac{1}{6}, \frac{2}{6}, \frac{3}{6}, \frac{4}{6}, \frac{5}{6}, 1$  satisfy  $m!x = 6x \in \mathbb{Z}$ .

We now show that  $f$  defined in the above example is not Riemann integrable on  $[0, 1]$ .

*Proof.* Consider any partition  $P$  of  $[0, 1]$ . Due to the density of rational and irrational numbers in  $\mathbb{R}$  (Theorem 1.20) we have that  $M_i = \sup \{f(x) : x \in [x_{i-1}, x_i]\} = 1$  and  $m_i = \inf \{f(x) : x \in [x_{i-1}, x_i]\} = 0$  for all  $i$ . Therefore, we have that  $U(P, f) = \sum_{i=1}^N M_i \Delta x_i = 1$  and  $L(P, f) = \sum_{i=1}^N m_i \Delta x_i = 0$  for all partitions  $P$ . Therefore,  $\sup_P U(P, f) = 1$  and  $\inf_P L(P, f) = 0$ , and we conclude that  $f$  is not Riemann integrable on  $[0, 1]$ .  $\square$

### Example

Define  $f_n$  such that:

$$f_n(x) = \begin{cases} 0 & |x| \geq \frac{1}{n} \\ n(nx+1) & -\frac{1}{n} < x < 0 \\ -n(nx+1) & 0 < x < \frac{1}{n} \\ 0 & x = 0 \end{cases}$$

Then, we have that  $f(x) = \lim_{n \rightarrow \infty} f_n(x) = 0$  for all  $x$ . Furthermore, we have that  $\int_{-1}^1 f_n(x) dx = 1$  for all  $n$ , but  $\int_{-1}^1 f(x) dx = 0$ . Hence, we have that:

$$\lim_{n \rightarrow \infty} \int_{-1}^1 f_n(x) dx = 1 \neq 0 = \int_{-1}^1 \lim_{n \rightarrow \infty} f_n(x) dx$$

showing that problems can arise when we interchange the order of an integral with a limit.



Figure 39: Plot of  $f_n$  in the above example.

### Example 7.5

Let  $f_n(x) = \frac{\sin nx}{\sqrt{n}}$  for  $n \in \mathbb{N}, x \in \mathbb{R}$ . Then, let  $f(x) = \lim_{n \rightarrow \infty} f_n(x) = 0$  for all  $x \in \mathbb{R}$ , so  $f'(x) = 0$ . However,  $f'_n(x) = \frac{1}{\sqrt{n}} n \cos nx = \sqrt{n} \cos nx$  and  $\lim_{n \rightarrow \infty} \sqrt{n} \cos nx$  does not exist. For example,  $f'_n(\pi) = \sqrt{n}(-1)^n$  which is a divergent sequence. So:

$$f'(\pi) = \left( \lim_{n \rightarrow \infty} f_n \right)'(\pi) = 0 \neq \lim_{n \rightarrow \infty} f'_n(\pi)$$

which shows us that problems can arise when interchanging a derivative (which is just a type of limit) with a limit.

With the above five examples, we have seen examples of bad behaviour that can occur under interchange of limits. Namely:

1. An interchange of the order of limits can change the limiting value for a double sequence.
2. The limit of a sequence of continuous functions is not necessarily continuous.
3. The limit of a sequence of Riemann integrable functions is not necessarily Riemann integrable.

4. The limit of a sequence of Riemann integrals can differ from the Riemann integral of the limit of a sequence.
5. The limit of a sequence of derivatives can differ from the derivative of a limit of a sequence.

The good news is that in all of these examples, the sequences we looked at had a “weak” form of convergence, where we fix  $x$  and then take the  $n \rightarrow \infty$  limit. We will now proceed to look at a stronger version of convergence, which looks at “all  $x$  at once”, ensuring that this bad behaviour does not (for the most part) occur.

## 7.2 Uniform Convergence

### Definition 7.7: Uniform Convergence

Let  $E$  be any set and  $f_n : E \mapsto \mathbb{R}$  or  $f_n : E \mapsto \mathbb{C}$  for  $n \in \mathbb{N}$ . Then,  $f_n$  **converges uniformly** to  $f$  on  $E$  if for all  $\epsilon > 0$ , there exists  $N$  such that  $n \geq N$  implies that  $|f_n(x) - f(x)| < \epsilon$  for all  $x \in E$ .

Note the lack of  $x$  dependence in the above definition. We give a useful visual intuition of uniform convergence below:



Figure 40: Visualization of the intuition behind uniform convergence. If  $f_n \rightarrow f$ , uniformly, for any  $\epsilon > 0$ , we can find  $N$  such that for  $n \geq N$ ,  $f_n(x)$  lies in the  $\epsilon$ -tube (pictured above) around  $f$ .

### Example

Let us return to Example 7.5. We have that:

$$|f_n(x) - f(x)| = \left| \frac{\sin nx}{\sqrt{n}} - 0 \right| \leq \frac{1}{\sqrt{n}}$$

So taking  $n$  large enough such that  $\frac{1}{\sqrt{n}} < \epsilon$ , we can see that  $f_n(x)$  converges uniformly to  $f(x) = 0$ . Note that this example does show that uniform convergence is *not* sufficient for:

$$\lim_{n \rightarrow \infty} f'_n = \left( \lim_{n \rightarrow \infty} f_n \right)'$$

to hold. We will return to the relation of uniform convergence and differentiation in a later theorem.

### Example

Let us return to our second example from our section on motivating examples. Recall we had:

$$f_n(x) = \begin{cases} 1 & x \geq 0 \\ 1 + nx & -\frac{1}{n} < x < 0 \\ 0 & x \leq -\frac{1}{n} \end{cases} \quad f(x) = \begin{cases} 1 & x \geq 0 \\ 0 & x < 0 \end{cases}$$

We then have that:

$$f_n(x) - f(x) = \begin{cases} 1 + nx & -\frac{1}{n} < x < 0 \\ 0 & \text{otherwise} \end{cases}$$

So for  $x = -\frac{1}{2n}$ , we have that:

$$f_n\left(-\frac{1}{2n}\right) - f\left(-\frac{1}{2n}\right) = 1 + n\left(-\frac{1}{2n}\right) - 0 = \frac{1}{2}$$

Which will never be less than  $\epsilon$  for  $\epsilon < \frac{1}{2}$ . Hence, we conclude that  $f_n$  does not converge uniformly to  $f$  on  $\mathbb{R}$ .



Figure 41: Visualization of why the convergence of  $f_n \rightarrow f$  in the above example is not uniform. We can see that if we draw a small enough  $\epsilon$  tube (i.e.  $\epsilon \leq 1$ ), there is no way to choose  $n$  large enough to make all of  $f_n(x)$  lie in the tube.

### Theorem 7.8: Cauchy Criterion for Uniform Convergence

$f_n$  converges uniformly on  $E$  if and only if for all  $\epsilon > 0$ , there exists  $N$  such that if  $m, n \geq N$ , then  $|f_m(x) - f_n(x)| < \epsilon$  for all  $x \in E$ .

Again, note the lack of  $x$  dependence in the above theorem.

### Proof

$\Rightarrow$  Suppose  $f_n \rightarrow f$  uniformly on  $E$ . Then, there exists some  $N$  such that for  $m, n \geq N$ :

$$|f_m(x) - f(x)| < \frac{\epsilon}{2}, \quad |f_n(x) - f(x)| < \frac{\epsilon}{2}$$

for all  $x \in E$ . Therefore by the triangle inequality, we have that:

$$|f_m(x) - f_n(x)| \leq |f_m(x) - f(x)| + |f(x) - f_n(x)| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

Hence  $|f_m(x) - f_n(x)| < \epsilon$  for all  $x \in E$ .

$\Leftarrow$  Let  $x \in E$ . By assumption,  $\{f_n(x)\}_{n \in \mathbb{N}}$  is a Cauchy sequence, and hence has a limit  $f(x)$  (as both  $\mathbb{R}$  and  $\mathbb{C}$ , the possible codomains of  $f$ , are complete). We then let  $f(x) = \lim_{n \rightarrow \infty} f_n(x)$ , so we have pointwise convergence. To see that the convergence is uniform, let  $\epsilon > 0$ . We know that  $|f_m(x) - f_n(x)| < \epsilon$  for  $m, n \geq N$  and for all  $x$ . Then, let  $m \rightarrow \infty$ . Then,  $|f(x) - f_n(x)| \leq \epsilon$  for all  $n \geq N$  and all  $x \in E$ , so the convergence is uniform.  $\square$

### Theorem 7.9

Suppose  $\lim_{n \rightarrow \infty} f_n(x) = f(x)$  for  $x \in E$ , and let:

$$M_n = \sup_{x \in E} |f_n(x) - f(x)|$$

Then,  $f_n \rightarrow f$  uniformly on  $E$  if and only if  $M_n \rightarrow 0$  as  $n \rightarrow \infty$ .

### Proof

$\Rightarrow$  suppose  $f_n \rightarrow f$  uniformly. Then, for any  $\epsilon > 0$ , there exists some  $N \in \mathbb{N}$  such that for all  $n \geq N$  and all  $x \in E$ :

$$|f_n(x) - f(x)| < \epsilon$$

Since this holds for all  $x \in E$ , taking the supremum of  $|f_n(x) - f(x)|$  we have that:

$$\sup_{x \in E} |f_n(x) - f(x)| = M_n \leq \epsilon$$

We then have that  $M_n \leq \epsilon$  for  $n \geq N$  for some  $N$ , and hence  $M_n \rightarrow 0$ .

$\Leftarrow$  Suppose that  $M_n \rightarrow 0$ . Then, for any  $\epsilon > 0$ , there exists some  $N \in \mathbb{N}$  such that for all  $n \geq N$ :

$$\sup_{x \in E} |f_n(x) - f(x)| = M_n < \epsilon$$

We then have that for any  $x \in E$ :

$$|f_n(x) - f(x)| \leq \sup_{x \in E} |f_n(x) - f(x)| < \epsilon$$

so we conclude that  $f_n \rightarrow f$  uniformly.  $\square$

### Definition: Uniform Convergence of Series

We say that  $\sum_{n=1}^{\infty} f_n(x)$  **converges uniformly** on  $E$  if  $S_n(x) = \sum_{i=1}^n f_i(x)$  is a uniformly convergent sequence of functions.

### Theorem 7.10: Weierstrass M-Test

Suppose  $|f_n(x)| < M_n$  for all  $n \geq N_0$  and for all  $x \in E$ . Suppose also that  $\sum_{n=N_0}^{\infty} M_n < \infty$ . Then,  $\sum_{n=1}^{\infty} f_n(x)$  converges uniformly on  $E$ .

### Proof

Let  $S_n(x) = \sum_{i=1}^n f_i(x)$ . For  $n > m \geq N_0$ , we have that:

$$|S_n(x) - S_m(x)| = \left| \sum_{i=m+1}^n f_i(x) \right| \leq \sum_{i=m+1}^n |f_i(x)| \leq \sum_{i=m+1}^n M_i$$

Let  $\epsilon > 0$ . Choose  $N \geq N_0$  such that  $\sum_{i=N+1}^{\infty} M_i < \epsilon$  (which we can choose as the series converges by assumption). We then have that  $|S_n(x) - S_m(x)| < \epsilon$  for all  $n > m \geq N$  for all  $x \in E$ . Hence,  $S_n(x)$  converges uniformly on  $E$ .  $\square$

### Theorem 7.11

Let  $E \subset X$  and  $f_n : E \mapsto \mathbb{R}$  or  $\mathbb{C}$ ,  $n \in \mathbb{N}$ . Suppose  $f_n \rightarrow f$  uniformly on  $E$ , and let  $x \in E$  (where  $x$  is a limit point of  $E$ ). Suppose  $\lim_{t \rightarrow x} f_n(t) = A_n$  exists for each  $n \in \mathbb{N}$ . Then,  $A_n \rightarrow A$  for some  $A$ . And  $\lim_{t \rightarrow x} f(t) = A$ . In other words:

$$\lim_{t \rightarrow x} \lim_{n \rightarrow \infty} f_n(t) = \lim_{n \rightarrow \infty} \lim_{t \rightarrow x} f_n(t)$$

showing that the interchange of limits is valid when we have uniform convergence.



Figure 42: Visualization of Theorem 7.11, with  $E = (0, \infty)$  and  $x = 0$ . Eventually, the graph of  $f_n$  lies in the  $\epsilon$  tube around  $f$  (no matter how skinny the tube is). But,  $A_n$  is being determined by  $f_n$  near 0, so there is nowhere for  $A_n$  to go except to the limiting value. That is,  $A_n \rightarrow A$  as the  $\epsilon$  tube gets compressed.

### Proof

We first show that  $A_n \rightarrow A$  for some  $A$ . Since  $\mathbb{R}, \mathbb{C}$  are complete metric spaces, it suffices to show that  $\{A_n\}$  is Cauchy. Given  $\epsilon > 0$ , choose  $N$  such that for  $m, n \geq N$ ,  $|f_n(t) - f_m(t)| < \epsilon$  for all  $t$  (such an  $N$  exists by Theorem 7.8). Letting  $t \rightarrow x$ , we therefore obtain that  $|A_n - A_m| \leq \epsilon$  for all  $m, n \geq N$ , showing that  $\{A_n\}$  is Cauchy. Hence, the sequence converges to some limit  $A$ .

Now, we show that  $\lim_{t \rightarrow x} f(t) = A$ . We show this by the common “ $\epsilon/3$  argument”. For all  $t \in E$  and  $n \in \mathbb{N}$ , we have by the triangle inequality that:

$$|f(t) - A| \leq |f(t) - f_n(t)| + |f_n(t) - A_n| + |A_n - A| (*)$$

Which is a good move, as we know that we can make each of the three terms on the RHS arbitrarily small (they are “close”). Let  $\epsilon > 0$ . Since  $f_n \rightarrow f$  uniformly, there exists  $N_1$  such that  $|f(t) - f_n(t)| < \frac{\epsilon}{3}$  for all  $n \geq N_1$  and all  $t \in E$ . Since  $A_n \rightarrow A$ , there exists some  $N_2$  such that  $|A_n - A| < \frac{\epsilon}{3}$  for all  $n \geq N_2$ . Letting  $N = \max\{N_1, N_2\}$  and taking  $n = N$  in  $(*)$ , we have that:

$$|f(t) - A| < \frac{\epsilon}{3} + |f_N(t) - A_N| + \frac{\epsilon}{3}$$

Since  $\lim_{t \rightarrow x} f_N(t) = A_N$ , we can choose  $\delta > 0$  such that  $t \in N_\delta(x)$  implies  $|f_N(t) - A_N| < \frac{\epsilon}{3}$  (Note a subtle point here that this choice of  $\delta$  depends on  $N$ !). Therefore, if  $t \in N_\delta(x)$ , we have that:

$$|f(t) - A| < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon$$

Hence, as  $t \rightarrow x$ ,  $f(t) \rightarrow A$ . □

### Theorem 7.12

Suppose  $f_n$  is continuous on  $E$  for all  $n \in \mathbb{N}$ , and  $f_n \rightarrow f$  uniformly on  $E$ . Then,  $f$  is continuous.

### Proof

Every  $f_n$  is continuous at isolated points of  $E$ , so it suffices to consider limit points  $x \in E' \cap E$ . For these points, we have that:

$$f(x) = \lim_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \lim_{t \rightarrow x} f_n(t) = \lim_{t \rightarrow x} \lim_{n \rightarrow \infty} f_n(t) = \lim_{t \rightarrow x} f(t)$$

Where the third equality (the interchange of the two limits) follows from Theorem 7.11. We conclude that  $f$  is continuous by Theorem 4.6 (as  $f(x) = \lim_{t \rightarrow x} f(t)$ ). □

### Theorem 7.13

Suppose  $K$  is compact, and:

- (a)  $f_n$  is continuous on  $K$  for each  $n \in \mathbb{N}$
- (b)  $f_n \rightarrow f$  pointwise (that is, for each  $x \in K$ ,  $f_n(x) \rightarrow f(x)$ ) and  $f$  is continuous
- (c)  $f_n(x) \geq f_{n+1}(x)$  for all  $x \in K$  and all  $n \in \mathbb{N}$  (note that the opposite inequality also works, just multiply by  $-1$ ).

Then,  $f_n \rightarrow f$  uniformly on  $K$ .



Note that this theorem is not super useful, being that it requires so many specific assumptions; however, we will find that it does have an interesting proof. Before we move to that, let us show some counterexamples for when the assumptions do not hold.

### Example

Let  $K = [-1, 0)$ , and define:

$$f_n(x) = \begin{cases} 0 & -1 \leq x \leq -\frac{1}{n} \\ 1 + nx & -\frac{1}{n} < x < 0 \end{cases}$$

We then have that  $f_n$  is continuous on  $K$ , that  $f_n \rightarrow 0$  pointwise on  $K$ , that  $f$  is continuous (the zero function), and  $f_n$  is decreasing with  $n$ . However, we note that  $f_n$  does not converge uniformly to  $f$  on  $K$ , with points close to zero being problem points (for example, take  $x = -\frac{1}{2n}$ , and then  $f_n(x) - f(x) = \frac{1}{2}$  for all  $n$ ). We note that  $K$  is *not* compact, showing the importance of compactness of the domain in the above Theorem.



Figure 43: Plot of  $f_n$  on  $K = [-1, 0)$  from the above example.

### Example

Let  $K = [0, 1]$ , and define:

$$f_n(x) = \begin{cases} 2nx & 0 \leq x < \frac{1}{2n} \\ 2 - 2nx & \frac{1}{n} \leq x \leq \frac{1}{n} \\ 0 & \frac{1}{n} < x \leq 1 \end{cases}$$

We then have that  $f_n$  is continuous,  $f_n \rightarrow f = 0$  pointwise (which is continuous), and  $K$  is compact. However,  $f_n$  does not converge to  $f$  uniformly. In this case, condition (c) of the above Theorem fails;  $f_n$  is not monotonic in  $n$ .



Figure 44: Plot of  $f_n$  on  $K = [0, 1]$  from the above example.

### Proof

Let  $g_n = f_n - f$ . We can then see that:

- (a)  $g_n$  is continuous (the difference of two continuous functions is continuous by Theorem 4.9)
- (b)  $g_n \rightarrow 0$  pointwise for all  $x \in K$
- (c)  $g_n \geq g_{n+1} \geq 0$  for all  $x \in K$ .

The goal will be to show that  $g_n \rightarrow 0$  uniformly on  $K$ . We will use the finite intersection property of compact sets to show this. Let  $\epsilon > 0$ . We will show that there exists  $N$  such that  $0 \leq g_n(x) < \epsilon$  for all  $n \geq N$  and for all  $x \in K$ . Note that it suffices to show that  $g_N(x) < \epsilon$  for some  $N$ , as  $g$  is monotone decreasing in  $n$ . Define  $K_n = g_n^{-1}([\epsilon, \infty))$  (i.e. the set of “bad  $x$ ”). We are done if we are able to show that there exists a  $N$  with  $K_N = \emptyset$ . Since  $g_n$  is continuous,  $K_n$  is closed as  $[\epsilon, \infty)$  is closed. Since  $K_n \subset K$ ,  $K$  is therefore compact as a closed subset of a compact set (Theorem 2.35). Additionally, we have that  $K_{n+1} \subset K_n$ , as  $g_{n+1} \geq \epsilon$  implies that  $g_n \geq \epsilon$ . Since  $g_n \rightarrow 0$  pointwise, given  $x \in K$ , there exists  $N_x$  such that  $x \notin K_n$  for all  $n \geq N_x$  (as  $g_n(x) < \epsilon$  for large enough  $n$ ). We therefore have that  $x \notin \bigcap_n K_n$  for all  $x \in K$ . Then, applying the corollary to Theorem 2.36, we obtain that  $K_N$  is empty. This means that for this  $N$ ,  $g_N^{-1}([\epsilon, \infty)) = \emptyset$ , and hence  $g_N^{-1}([0, \epsilon]) = K$ , which is to say that  $0 \leq g_n(x) < \epsilon$  for all  $x \in K$ .  $\square$

### Definition 7.14: $\mathcal{C}(X)$ and the Supremum Norm

For a metric space  $X$ , define:

$$\mathcal{C}(X) = \{f : X \mapsto \mathbb{C} \text{ such that } f \text{ is bounded and continuous.}\}$$

The **supremum norm** of  $f \in \mathcal{C}(X)$  is then defined as  $\|f\| = \sup_{x \in X} |f(x)|$ . We claim that  $\|f - g\|$  defines a metric on  $\mathcal{C}(X)$ , and we prove this assertion below. Thus, we have that:

$$\begin{aligned} f_n \rightarrow f \text{ uniformly} &\iff \forall \epsilon > 0, \exists N \text{ such that } |f_n(x) - f(x)| < \epsilon \forall n \geq N \text{ and } \forall x \in X \\ &\iff \forall \epsilon > 0, \exists N \text{ such that } \|f_n(x) - f(x)\| < \epsilon \forall n \geq N \\ &\iff f_n \rightarrow f \text{ in the metric space } \mathcal{C}(X) \end{aligned}$$

We have hence “metrized” uniform convergence.

### Theorem

$\|f - g\|$  defines a metric on  $\mathcal{C}(X)$ .

### Proof

We recall the three properties of a metric as per Definition 2.15:

- (a)  $d(f, g) = 0 \iff f = g$
- (b)  $d(f, g) = d(g, f)$
- (c)  $d(f, g) \leq d(f, h) + d(h, g)$

We now show that  $\|f - g\|$  satisfies these three properties.

- (a)  $\|f - g\| = 0$  means that  $0 = \sup_{x \in X} |f(x) - g(x)| \implies |f(x) - g(x)| = 0$  for all  $x$ , hence  $f(x) = g(x)$ .
- (b)  $\|f - g\| = \sup_{x \in X} |f(x) - g(x)| = \sup_{x \in X} |g(x) - f(x)| = \|g - f\|$
- (c) We have that  $|f(x) - g(x)| \leq |f(x) - h(x)| + |h(x) - g(x)|$  for all  $x \in X$ , so  $\|f - g\| \leq \|f - h\| + \|h - g\|$ .  $\square$

Note that sometimes  $\|f\|$  is written as  $\|f\|_\infty$  as it is the  $n \rightarrow \infty$  limit of the  $L_p$  norm. See HW3Q3 for the proof that the supremum norm is the limit of the  $L_p$  norm.

### Theorem 7.15

$\mathcal{C}(X)$  is a complete metric space (every Cauchy sequence in  $\mathcal{C}(X)$  has a limit in  $\mathcal{C}(X)$ ).

### Proof

Let  $\{f_n\}$  be a Cauchy sequence in  $\mathcal{C}(X)$ . Then, given  $\epsilon > 0$ ,  $\exists N$  such that  $m, n \geq N$  implies  $\|f_m - f_n\| = \sup_{x \in X} |f_m(x) - f_n(x)| < \epsilon$ . By the Cauchy criterion (Theorem 7.8),  $f_n \rightarrow f$  for some  $f$ . What is left to show is that  $f \in \mathcal{C}(X)$ .  $f$  is continuous as it is the uniform limit of continuous functions (Theorem 7.12). Additionally,  $f$  is bounded as there exists  $N_0$  such that  $|f(x) - f_{N_0}(x)| < 1$  for all  $x$ , and hence  $|f(x)| \leq |f_{N_0}(x)| + |f(x) - f_{N_0}(x)| \leq M_0 + 1$  for all  $x$  where  $M_0$  is the bound on  $f_{N_0}(x)$  that exists as  $f_{N_0} \in \mathcal{C}(X)$ . As  $f$  is continuous and bounded, we conclude that  $f \in \mathcal{C}(X)$ .  $\square$

## 7.3 Uniform Convergence and Integration

### Theorem 7.16

Suppose  $f_n \in \mathcal{R}_a[a, b]$  for all  $n \in \mathbb{N}$  and that  $f_n \rightarrow f$  uniformly on  $[a, b]$ . Then,  $f \in \mathcal{R}_a[a, b]$ , and:

$$\lim_{n \rightarrow \infty} \int_a^b f_n d\alpha = \int_a^b f d\alpha$$

In other words, the above Theorem tells us that we can interchange the integral with the limit if the sequence is uniformly convergent. Compare this to our earlier example with pointwise convergence, where such an interchange was not possible (as it yielded different values).

#### Proof

First, we show that  $f \in \mathcal{R}_\alpha[a, b]$ . Let  $\epsilon > 0$ . Since  $f_n \rightarrow f$  uniformly, there exists  $N$  such that  $|f_n(x) - f(x)| < \epsilon$  if  $n \geq N$  for all  $x \in [a, b]$ . So,  $f_n(x) - \epsilon < f(x) < f_n(x) + \epsilon$ . Hence,

$$\int_a^b (f_n - \epsilon) d\alpha \leq \int_a^b f d\alpha \leq \int_a^b (f_n + \epsilon) d\alpha$$

Since  $f_n \pm \epsilon \in \mathcal{R}_\alpha[a, b]$ , we have that:

$$\int_a^b (f_n - \epsilon) d\alpha \leq \int_a^b f d\alpha \leq \int_a^b (f_n + \epsilon) d\alpha$$

Therefore:

$$0 \leq \int_a^b f d\alpha - \int_a^b f_n d\alpha \leq \int_a^b 2\epsilon d\alpha \implies \int_a^b f d\alpha - \int_a^b f_n d\alpha \leq 2\epsilon(\alpha(b) - \alpha(a))$$

Since  $\epsilon$  is arbitrary, we have that  $\int_a^b f d\alpha = \lim_{n \rightarrow \infty} \int_a^b f_n d\alpha$  and hence  $f \in \mathcal{R}_\alpha[a, b]$ .

Next, we show that  $\lim_{n \rightarrow \infty} \int_a^b f_n d\alpha = \int_a^b f d\alpha$ . To do this, we show that  $\left| \int_a^b f d\alpha - \int_a^b f_n d\alpha \right|$  goes to 0 as  $n \rightarrow \infty$ . We have that:

$$\left| \int_a^b f d\alpha - \int_a^b f_n d\alpha \right| = \left| \int_a^b (f - f_n) d\alpha \right| \leq \int_a^b |f - f_n| d\alpha \leq \int_a^b \epsilon d\alpha = \epsilon(\alpha(b) - \alpha(a))$$

Where in the first equality we use Linearity (Theorem 6.12), the first inequality we apply Theorem 6.13, and in the second inequality, we use that for any  $\epsilon > 0$ , there exists  $N$  such that  $|f - f_n| < \epsilon$  for  $n \geq N$ . Since  $\epsilon$  is arbitrary, we conclude that  $\lim_{n \rightarrow \infty} \int_a^b f_n d\alpha = \int_a^b f d\alpha$ .  $\square$

#### Corollary

If  $f_n \in \mathcal{R}_\alpha[a, b]$  and  $f(x) = \sum_{n=1}^{\infty} f_n(x)$  converges uniformly on  $[a, b]$ , then  $\int_a^b f d\alpha = \sum_{n=1}^{\infty} \int_a^b f_n d\alpha$ . That is to say, the infinite series and the integral can be interchanged.

#### Proof

Let  $S_n(x) = \sum_{i=1}^n f_i(x)$ . Then,  $S_n(x) \rightarrow f(x)$  uniformly by assumption, so:

$$\int_a^b f d\alpha = \lim_{n \rightarrow \infty} \int_a^b S_n d\alpha = \lim_{n \rightarrow \infty} \sum_{i=1}^n \int_a^b f_i d\alpha = \sum_{i=1}^{\infty} \int_a^b f_i d\alpha$$

Where the first equality follows from the previous theorem, and the second equality follows from the fact that a finite sum and integral can be interchanged by Linearity.  $\square$

## 7.4 Uniform Convergence and Differentiation

Recall Example 7.5, where we looked at the sequence of functions  $f_n(x) = \frac{\sin nx}{\sqrt{n}}$ . We showed that  $f_n \rightarrow 0$  uniformly on  $\mathbb{R}$ , but we found in the example that  $f'_n(x)$  does *not* converge. We are therefore motivated to find a condition that if a function converges and is differentiable, then  $f'_n$  converges.

As a point of notation, note that for  $a < b$  we denote  $\int_b^a f d\alpha = -\int_a^b f d\alpha$ .

### Theorem 7.17

Suppose:

- (a)  $f_n$  is differentiable on  $[a, b]$ ;
- (b)  $\exists x_0 \in [a, b]$  such that  $f_n(x_0)$  converges as  $n \rightarrow \infty$ ;
- (c)  $f'_n$  converges uniformly on  $[a, b]$ .

Then, there exists  $f$  such that  $f_n \rightarrow f$  uniformly on  $[a, b]$ , and:

$$\lim_{n \rightarrow \infty} f'_n(x) = f'(x) \quad \forall x \in [a, b]$$

A couple remarks before we move to the proof. First, we note that hypothesis (b) seems strange; why would we require convergence  $f_n$  at a single point? This has to do with the fact that in differentiating, we lost our constants. For example, let  $f_n(x) = n$  as the simplest example. In this case, we have that  $f_n$  is differentiable everywhere (with derivative zero everywhere on  $[a, b]$ ) and that  $gf_n'$  uniformly converges (it is just the sequence of the zero function). However,  $f_n$  does not even converge!

Note that we can and will assume that  $f_n(x_0) \rightarrow 0$  at the specified  $x_0$ ; if this is not true, we can simply replace  $f_n(x)$  by  $f_n(x) - f_n(x_0)$ .

### Proof

The proof of the above theorem is not so trivial. We will therefore prove a weaker theorem. Namely, we add a fourth hypothesis (d) that  $f'_n$  is continuous on  $[a, b]$ . The proof of the stronger/original theorem can be found in Rudin.

First, by (c) there exists a  $g$  such that  $f'_n \rightarrow g$  uniformly on  $[a, b]$  (and also on any subinterval of  $[a, b]$ ). Furthermore, by (d) and Theorem 7.12,  $g$  is continuous.

Next, applying Theorem 7.16 (to either  $[x_0, x]$  or  $[x, x_0]$ ) we have that:

$$\int_{x_0}^x f'_n(t) dt \rightarrow \int_{x_0}^x g(t) dt = f(x)$$

Then applying the Fundamental theorem of calculus (Theorem 6.21), we have  $f_n(x) - f_n(x_0) \rightarrow f(x)$  and  $f'(x) = g(x)$ . But we also assume that  $f_n(x_0) \rightarrow 0$ , so  $f_n(x) \rightarrow f(x)$  and  $f'_n(x) \rightarrow g(x) = f'(x)$ . So, we have shown pointwise convergence of  $f_n$  to  $f$ ! We have obtained that  $\lim_{n \rightarrow \infty} f'_n(x) = f'(x)$  for all  $x \in [a, b]$ . Finally, we show  $f_n \rightarrow f$  uniformly on  $[a, b]$ . We have that:

$$\begin{aligned} |f(x) - f_n(x)| &= \left| \int_{x_0}^x g(t) dt - \int_{x_0}^x f'_n(t) dt + f_n(x_0) \right| \leq \int_{x_0}^x |g(t) - f'_n(t)| dt + |f_n(x_0)| \\ &< \int_{x_0}^x \frac{\epsilon}{2(b-a)} dt + \frac{\epsilon}{2} \leq \frac{\epsilon}{2(b-a)}(b-a) + \frac{\epsilon}{2} = \epsilon \end{aligned}$$

Where we apply Theorem 6.13 for the first inequality, the fact that  $f'_n(t) \rightarrow g$  and  $f_n(x_0) \rightarrow 0$  in the second last inequality, and Theorem 6.12(d) in the last inequality.  $\square$

### Theorem 7.18

There exists a continuous function  $f : \mathbb{R} \mapsto \mathbb{R}$  such that  $f'(x)$  does not exist for any  $x \in \mathbb{R}$ .

The proof of the above theorem will follow by the construction of an “infinitely spiky” real function. Though this might seem like a very pathological counterexample, there are actually many examples of non-differentiable phenomena in mathematics. Looking at the field of probability, we find that brownian motion, brownian maps, and discrete exploration processes (to name a few) all have this property. A visualization of the brownian map, as well as other beautiful probability pictures can be found here [https://secure.math.ubc.ca/Links/Probability/pages/pic\\_gallery.html](https://secure.math.ubc.ca/Links/Probability/pages/pic_gallery.html).




Figure 45: Plot of the  $\phi$  function defined in the proof of Theorem 7.18.




Figure 46: Desmos visualization of the nowhere-differentiable  $f$  constructed in the proof of Theorem 7.18. Note that a partial sum ( $N = 10$ ) of the series that  $f$  is defined to be is shown, as the infinite series is impossible to plot. Readers can play around the function with themselves at <https://www.desmos.com/calculator/onhkmb1go6>. There is a notion that  $f$  is “infinitely spiky”, no matter how much one is to zoom into the above graph.

### Proof

Define  $\phi : \mathbb{R} \mapsto \mathbb{R}$  by  $\phi(x) = |x|$  for  $-1 \leq x \leq 1$  and  $\phi(x+2) = \phi(x)$  for all  $x \in \mathbb{R}$ . (See figure 45 above). Then,  $\phi$  is continuous; moreover, it is Lipschitz continuous, with  $|\phi(s) - \phi(t)| \leq |s - t|$  for all  $s, t \in \mathbb{R}$  (with equality where there are no integers between  $s, t$ ). Define  $f(x) = \sum_{n=0}^{\infty} \left(\frac{3}{4}\right)^n \phi(4^n x)$ . The series converges uniformly on  $\mathbb{R}$  by Theorem 7.10, since  $0 \leq \left(\frac{3}{4}\right)^n \phi(4^n x) \leq \left(\frac{3}{4}\right)^n$  and  $\sum_n \left(\frac{3}{4}\right)^n$  converges (it is a geometric series with  $r < 1$ ). Hence,  $f$  is continuous as it is a uniform limit of a continuous function (Theorem 7.12). We now prove that  $f'(x)$  does not exist for any  $x \in \mathbb{R}$ ; let us then fix  $x$ . It suffices to find  $\delta_m \rightarrow 0$  such that:

$$\left| \frac{f(x + \delta_m) - f(x)}{\delta_m} \right| \rightarrow \infty \text{ as } m \rightarrow \infty.$$

We then choose  $\delta_m = \pm \frac{1}{2} \frac{1}{4^m}$ . We choose the sign of  $\delta_m$  depending on the choice of  $x$  as follows. At most one of  $(4^m x - \frac{1}{2}, 4^m x)$  and  $(4^m x, 4^m x + \frac{1}{2})$  contains an integer. We choose the sign such that no integer lies between  $4^m x$  and  $4^m(x + \delta_m)$ . Note that we may choose a different sign for each  $m$ . Next, we make the observation that  $|\phi(4^m(x + \delta_m)) - \phi(4^m x)| = 4^m x$ ; this holds as for the difference between two  $\phi(x)$  values at two points without an integer between them is just the difference between the  $x$  values. Looking back at our definition of  $\delta_m$ , we then see that  $|\phi(4^m(x + \delta_m)) - \phi(4^m x)| = \frac{1}{2}$ .

Furthermore, we see that if  $n > m$ , we have that  $\phi(4^n(x + \delta_m)) - \phi(4^n x \pm \frac{1}{2} 4^{n-m}) = \phi(4^n x)$  as  $\frac{1}{2} 4^{n-m}$  is an even integer and  $\phi$  is 2-periodic. This leads us to conclude that  $\phi(4^n(x + \delta_m)) - \phi(4^n x) = 0$  if  $n > m$ . Given  $m$ , then define:

$$\gamma_n = \frac{\phi(4^n(x + \delta_m)) - \phi(4^n x)}{\delta_m}$$

Then,  $\gamma_n = 0$  if  $n > m$ ,  $|\gamma_m| = \left| \frac{4^m \delta_m}{\delta_m} \right| = |4^m| = 4^m$ , and if  $0 \leq n < m$ ,  $|\gamma_n| \leq \frac{1}{\delta_m} |4^n(x + \delta_m) - 4^n x| = \frac{1}{|\delta_m|} |4^n \delta_m| = 4^n$ . Finally, we have that:

$$\begin{aligned} \left| \frac{f(x + \delta_m) - f(x)}{\delta_m} \right| &= \left| \sum_{n=0}^{\infty} \left(\frac{3}{4}\right)^n \gamma_n \right| = \left| \sum_{n=0}^m \left(\frac{3}{4}\right)^n \gamma_n \right| \geq \left(\frac{3}{4}\right)^m |\gamma_m| - \sum_{n=0}^{m-1} \left(\frac{3}{4}\right)^n |\gamma_n| \\ &\geq \left(\frac{3}{4}\right)^m 4^m - \sum_{n=0}^{m-1} \left(\frac{3}{4}\right)^n 4^n \\ &= 3^m - \sum_{n=1}^{m-1} 3^n \\ &= 3^m - \frac{3^m - 1}{3 - 1} \\ &= \frac{1}{2}(3^m + 1) \end{aligned}$$

Where the second equality follows as all terms  $n > m$  are zero, the first inequality follows by the reverse triangle inequality, the second inequality follows by the bounds on  $|\gamma_n|$ , and the third-to-last equality is the geometric sum formula. As  $m \rightarrow \infty$ , the difference quotient goes to infinity, and we therefore conclude that  $f$  is differentiable nowhere.  $\square$

## 7.5 Equicontinuous Families of Functions

### Definition 7.22: Equicontinuity

A family  $\mathcal{F}$  of functions on  $E$  (that is, a possibly finite, countable, or uncountable set of functions on  $E$ ) is **equicontinuous** on  $E$  if for every  $\epsilon > 0$ , there exists  $\delta > 0$  such that if  $f \in \mathcal{F}$  and  $x, y \in E$  with  $d(x, y) < \delta$ , then  $|f(x) - f(y)| < \epsilon$ . Note that the functions  $f \in \mathcal{F}$  are either real or complex valued.

The above definition of equicontinuity is essentially an even stronger version of uniform continuity; the  $\delta$  works not just for all  $x$  and  $y$  for a single  $f$ , but for all  $x$  and  $y$  for all of the  $f$ s in  $\mathcal{F}$ . If  $\mathcal{F} = \{f\}$ , then this is just uniform continuity.

### Theorem

- (a) If a family  $\mathcal{F}$  of functions on  $E$  is equicontinuous, then every  $f \in \mathcal{F}$  is uniformly continuous on  $E$ .
- (b) Any finite family  $\mathcal{F} = \{f_1, \dots, f_n\}$  of uniformly continuous functions on  $E$  is equicontinuous.

### Proof

- (a) The claim follows immediately from the definition.
- (b) Let  $\epsilon > 0$ . Then, by the uniform continuity of each  $f_i \in \mathcal{F}$ , there exists  $\delta_i$  such that if  $d(x, y) < \delta_i$ , then  $|f_i(x) - f_i(y)| < \epsilon$ . Taking  $\delta = \min \delta_1, \dots, \delta_n$ , we have that for any  $f \in \mathcal{F}$ , if  $d(x, y) < \delta$  then  $|f(x) - f(y)| < \epsilon$ . Hence  $\mathcal{F}$  is equicontinuous.  $\square$

### Example

Let  $\mathcal{F} = \{f_1, f_2, \dots\}$  with  $f_n(x) = \frac{\sin(nx)}{\sqrt{n}}$  for  $x \in [0, 1] \in E$ . Then,  $\mathcal{F}$  is equicontinuous.

### Proof

We have that  $|f_n(x) - f_n(y)| = \frac{1}{\sqrt{n}} |\sin(nx) - \sin(ny)| \leq \frac{2}{\sqrt{n}}$  for all  $x, y \in E$ . Let  $\epsilon > 0$ . Choose  $N$  such that  $\frac{2}{\sqrt{n}} < \epsilon$  if  $n > N$ . Since the remaining  $f_n$  (i.e.  $\{f_1, \dots, f_N\}$ ) are a finite collection of uniformly continuous functions (they are uniformly continuous by Theorem 4.19, as they are continuous functions on a closed and bounded interval), by the above theorem,  $\{f_1, \dots, f_N\}$  is equicontinuous. So, there exists a  $\delta$  such that for  $n \geq N$  and  $|x - y| < \delta$ ,  $|f_n(x) - f_n(y)| < \epsilon$ . We then have that for any  $n \in \mathbb{N}$  and for any  $x, y \in E$  with  $|x - y| < \delta$ , then  $|f_n(x) - f_n(y)| < \epsilon$ . We conclude that  $\mathcal{F}$  is equicontinuous.  $\square$

### Theorem (Problem 7.16)

Let  $\{f_n\}$  be an equicontinuous sequence of functions such that  $f_n : K \mapsto \mathbb{C}$  with  $K$  compact. Suppose there is a pointwise limit  $f(x) = \lim_{n \rightarrow \infty} f_n(x)$  that exists for all  $x \in K$ . Then,  $f_n \rightarrow f$  uniformly on  $K$ .



### Proof

We use an “ $\frac{\epsilon}{3}$  argument”. Let  $\epsilon > 0$ . Then, choose  $\delta > 0$  such that  $|f_n(x) - f_n(y)| < \frac{\epsilon}{3}$  for all  $n$  and for all  $x, y$  such that  $d(x, y) < \delta$  (such a choice is possible by the equicontinuity of the sequence). Take an open cover of  $K$  by considering the set of neighbourhoods of radius  $\delta$  around every point  $x \in K$ . Since  $K$  is compact, the open cover  $\{N_\delta(x) : x \in K\}$  has a finite subcover  $\{N_\delta(x_1), \dots, N_\delta(x_k)\}$ . Thus, given  $x \in K$ , there exists  $x_j$  such that  $x \in N_\delta(x_j)$  and hence  $d(x_j, x) < \delta$ . Therefore by the triangle inequality:

$$\begin{aligned} |f_n(x) - f_m(x)| &\leq |f_n(x) - f_n(x_j)| + |f_n(x_j) - f_m(x_j)| + |f_m(x_j) - f_m(x)| \\ &< \frac{\epsilon}{3} + |f_n(x_j) - f_m(x_j)| + \frac{\epsilon}{3} \end{aligned}$$

where the last inequality follows from the equicontinuity. For each  $i \in \{1, \dots, k\}$ , we know that  $\{f_n(x_i)\}$  is a convergent sequence as  $f_n$  converges pointwise by assumption. Hence, it is a Cauchy sequence. Therefore, there exists a  $N_i$  such that  $m, n \geq N_i$  implies  $|f_n(x_i) - f_m(x_i)| < \frac{\epsilon}{3}$ . Take  $N = \max N_1, \dots, N_k$ . Then, we have that  $m, n \geq N$  implies:

$$|f_n(x) - f_m(x)| < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon.$$

So,  $f_n$  satisfies the Cauchy criterion for uniform convergence, and hence  $\{f_n\}$  converges uniformly on  $K$ .  $\square$

### Theorem 7.24

If  $f_n : K \mapsto \mathbb{C}$  is continuous,  $K$  is compact, and  $f_n \mapsto f$  uniformly on  $K$ , then  $\{f_n\}$  is equicontinuous.

### Proof

We again use an “ $\frac{\epsilon}{3}$  argument”. Let  $\epsilon > 0$ . Since  $f_n \rightarrow f$  uniformly, we have that there exists  $N$  such that  $m, n \geq N$  implies  $|f_n(x) - f_m(x)| < \frac{\epsilon}{3}$  for all  $x \in K$ . Also, since  $K$  is compact, each  $f_i$  is uniformly continuous, so  $\{f_1, \dots, f_N\}$  is equicontinuous for any  $N \in \mathbb{N}$  (as it is a finite set of uniformly continuous functions). Hence, there exists  $\delta > 0$  such that  $|f_i(x) - f_i(y)| < \frac{\epsilon}{3}$  if  $d(x, y) < \delta$  and  $i \leq N$ . Finally, for  $n > N$ , we have that:

$$|f_n(x) - f_n(y)| \leq |f_n(x) - f_N(x)| + |f_N(x) - f_N(y)| + |f_N(y) - f_n(y)| < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon$$

where the first/third  $\frac{\epsilon}{3}$ s come from uniform convergence and the second from the equicontinuity. We conclude that  $\{f_n\}$  is equicontinuous.  $\square$

### Example

We here discuss a set of functions which is not equicontinuous, by returning to a prior example. Let  $K = [0, 1]$ , and define:

$$f_n(x) = \begin{cases} 2nx & 0 \leq x < \frac{1}{2n} \\ 2 - 2nx & \frac{1}{n} \leq x \leq \frac{1}{n} \\ 0 & \frac{1}{n} < x \leq 1 \end{cases}.$$

See Figure 44 for a visualization. We have that  $\{f_n\}$  obeys  $|f_n(x)| \leq 1$  for all  $x \in [0, 1]$ , but that  $\{f_n\}$  is not equicontinuous, as  $|f_n(\frac{1}{2n}) - f_n(0)| = 1 - 0 = 1$  for all  $n$ . We can get as close as we like to 0, but the difference will remain large. Also, there is no subsequence of  $\{f_n\}$  that can be uniformly convergent on  $[0, 1]$ , as  $f_n(x) \rightarrow 0$  for all  $x \in [0, 1]$  pointwise but  $f_n(\frac{1}{2n}) = 1$  for all  $n$ .  $f_n(x)$  “stays far” from the limit. The takeaway message is that a sequence that converges pointwise but is not equicontinuous is not guaranteed to have a uniformly convergent subsequence. This is motivation for the later Theorem 7.25, which gives criteria for a sequence of functions having a uniformly convergent subsequence.

### Definition 7.19: Pointwise/Uniform Bounded Functions

$\{f_n\}$  is **pointwise bounded** on  $E$  if there exists a  $\phi : E \mapsto \mathbb{R}$  such that  $|f_n(x)| < \phi(x)$  for all  $x \in E$  and for all  $n \in \mathbb{N}$ .  $\{f_n\}$  is **uniformly bounded** on  $E$  if there exists  $M$  such that  $|f_n(x)| \leq M$  for all  $x \in E$  and all  $n \in \mathbb{N}$ .

### Example

Let  $f_n(x) = \frac{1}{x} + \frac{1}{n}$ . Then,  $\{f_n\}$  is pointwise bounded, by (for example)  $\phi(x) = \frac{1}{x} + 2$ . But, it is not uniformly bounded, as  $\frac{1}{x}$  grows arbitrarily large as  $x \rightarrow 0$ .

### Theorem 7.23: Selection Theorem

Suppose  $f_n : E \mapsto \mathbb{C}$  is pointwise bounded on a countable set  $E$ . Then, some subsequence  $\{f_{n_k}\}$  of  $\{f_n\}$  is pointwise convergent on  $E$ ; that is to say,  $\lim_{k \rightarrow \infty} f_{n_k}(x)$  exists for all  $x \in E$ .

Note that the above theorem plays a large role in probability!

### Proof

We invoke a “diagonal argument”. Let  $E = \{x_1, x_2, \dots\}$ . Consider the sequence  $\{f_n(x_1)\}$ . We have that it is pointwise bounded by hypothesis, so there exists a subsequence  $\{f_{1_k}\}$  such that  $\lim_{n \rightarrow \infty} f_{1_n}(x_1)$  converges. (Theorem 2.42). We can apply the same logic for  $x_2, x_3, \dots$  in term, such that the proceeding sequence is a subsequence of the former. In other words, we form the array:

$$\begin{array}{cccc} S_1 : & f_{1_1} & f_{1_2} & f_{1_3} & \dots \\ S_2 : & f_{2_1} & f_{2_2} & f_{2_3} & \dots \\ S_3 : & f_{3_1} & f_{3_2} & f_{3_3} & \dots \\ & \vdots & & & \end{array}$$

In doing so, we have that  $S_1$  converges on  $x_1$ ,  $S_2$  is a subsequence of  $S_1$  that converges on  $x_1$  and  $x_2$ ,  $S_3$  is a subsequence of  $S_2$  that converges on  $x_1$  and  $x_2$  and  $x_3$  and so on. Then, we consider the sequence formed by the diagonal of the above array, with  $S : f_{1_1}, f_{2_2}, f_{3_3}, \dots$ . This is a subsequence of our original sequence  $f_n$ . If we fix some  $N$ , then this subsequence is a subsequence of  $S_n$  for  $n \geq N$  (as  $S$  is eventually a subsequence of each  $S_n$ ), so it converges on the same points that  $S_n$  does, namely,  $x_1, \dots, x_n$ . But this is true for every  $n \in \mathbb{N}$ , so the subsequence  $S$  converges for  $x_i \in E$  for every  $i \in \mathbb{N}$ .  $\square$

### Lemma (Problem 2.25)

If  $K$  is compact, then  $K$  has a countable dense subset  $E \subset K$  (i.e.  $\bar{E} = E \cup E' = K$ ). Alternatively, for all  $x \in K$ , there exists  $r > 0$  such that there exists  $p \in E$  such that  $d(p, x) < r$ . In other words,  $K$  is separable.

### Proof

For  $n \in \mathbb{N}$ ,  $\{N_{1/n}(p)\}_{p \in K}$  is an open cover. So, it has a finite subcover  $\{N_{1/n}(p)\}_{p \in E_n}$  where  $E_n \subset K$  is finite. Let  $E = \bigcup_{n=1}^{\infty} E_n$ , then  $E$  is at most countable (Theorem 2.12). To see that it is dense, let  $x \in K$ , and  $r > 0$ . Then, choose  $n_0$  such that  $\frac{1}{n_0} < r$ . We can then find  $p_0 \in E_{n_0} \subset E$  such that  $x \in N_{1/n_0}(p_0)$  as  $E_{n_0}$  is an open cover of  $K$ . Then,  $d(x, p_0) < \frac{1}{n_0} < r$ . Hence,  $E$  is dense.  $\square$

### Theorem 7.25: Arzela-Ascoli

Suppose  $K$  is compact, and that  $\mathcal{F} = \{f_n\} \subset \mathcal{C}(K)$  is equicontinuous and pointwise bounded. Then,

- (a)  $\{f_n\}$  is uniformly bounded.
- (b)  $\{f_n\}$  has a uniformly convergent subsequence (i.e. a subsequence that converges in  $\mathcal{C}(K)$ ).

## Proof

- (a) The goal is to find  $M$  such that  $|f_n(x)| \leq M$  for all  $n \in \mathbb{N}$  and for all  $x \in K$ . Let  $\epsilon > 0$  (though we can take  $\epsilon = 0$  for this proof of part (a)). Since  $\mathcal{F}$  is equicontinuous, we have that there exists  $\delta > 0$  such that  $d(x, y) < \delta$  implies  $|f_n(x) - f_n(y)| < \epsilon$  for all  $n$ .  $K$  is compact, we can cover  $K$  with balls of radius  $\delta$  around each point in  $K$  and then take a finite subcover; i.e. there exists a finite set  $\{p_1, \dots, p_r\} \in K$  such that  $\{N_\delta(p_i)\}_{i=1, \dots, r}$  covers  $K$ . For each  $i$ ,  $\{f_n(p_i)\}_n$  is bounded, that is,  $|f_n(p_i)| \leq M_i$  for all  $n$ . Let  $M_0 = \max M_1, \dots, M_r$ . Given  $x \in K$ , choose  $p_i$  such that  $x \in N_\delta(p_i)$ . Then:

$$|f_n(x)| \leq |f_n(p_i)| + |f_n(x) - f_n(p_i)| < M_i + \epsilon \leq M_0 + \epsilon$$

letting  $M = M_0 + \epsilon$ , we see that  $\{f_n\}$  is a uniformly bounded.

- (b) Our goal is to construct a uniformly convergent subsequence. We do this in three steps. First, we construct a subsequence (we will show it is uniformly convergent afterwards!). By the above Lemma,  $K$  has a countable dense subset  $E$ . By Theorem 7.23, there exists a subsequence  $\{f_{n_i}\}$  such that  $\lim_{i \rightarrow \infty} f_{n_i}(x)$  exists for all  $x \in E$ . Write  $g_i = f_{n_i}$ .

Secondly, we set up the argument to show uniform convergence of the subsequence constructed in the first step. Let  $\epsilon > 0$ . By the equicontinuity assumption, there exists  $\delta > 0$  such that  $d(x, y) < \delta$  implies  $|g_i(x) - g_i(y)| < \frac{\epsilon}{3}$  for all  $i$ . Consider  $\{N_\delta(p)\}_{p \in E}$ , which covers  $K$  since  $E$  is dense. By the compactness of  $K$ , there exists a finite subset  $\{N_\delta(x_1), \dots, N_\delta(x_m)\}$  with  $x_i \in E$ . Hence, given  $x \in K$ , there exists  $x_s$  such that  $d(x, x_s) < \delta$ .

For the third step, we complete the proof with an " $\frac{\epsilon}{3}$  argument". Using the triangle inequality, we have that:

$$\begin{aligned} |g_i(x) - g_j(x)| &\leq |g_i(x) - g_i(x_s)| + |g_i(x_s) - g_j(x_s)| + |g_j(x_s) - g_j(x)| \\ &< \frac{\epsilon}{3} + |g_i(x_s) - g_j(x_s)| + \frac{\epsilon}{3} \end{aligned}$$

where the last inequality follows from the arguments in step 2. For the second term, we consider that for  $s = 1, \dots, m$ , we can choose  $N_s$  such that  $|g_j(x_s) - g_i(x_s)| < \frac{\epsilon}{3}$  for  $i, j \geq N_s$  (this choice of  $N_s$  is possible as  $\{g_n(x_s)\}$  converges). There are finitely many  $N_s$ s, so let  $N = \max N_1, \dots, N_m$ . Then, we have that:

$$|g_i(x) - g_j(x)| < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon$$

for  $i, j \geq N$  and for all  $x \in K$ . Hence,  $\{g_i\}$  converges uniformly on  $K$ . □

## 7.6 The Stone-Weierstrass Theorem

### Theorem 7.26: Weierstrass

Let  $f : [a, b] \mapsto \mathbb{R}$  be continuous. Then, there exists polynomials  $P_n$  such that  $P_n \rightarrow f$  uniformly on  $[a, b]$ .

Note that it will suffice to consider the case where  $[a, b] = [0, 1]$ ; we can get to arbitrary  $[a, b]$  to  $[0, 1]$  by a change of variable, and the composition of a polynomial with a change of variable is still a polynomial.

Note that our proof will take a different angle from Rudin's proof of the theorem; we shall be exploring the proof by Bernstein. However, before we begin the proof, we will need to establish some basic background in probability.

#### Definition: Bernoulli Trials

A **Bernoulli trial** is a random experiment with a success outcome of probability  $p$  and a failure outcome with probability  $1 - p$  (here,  $p \in [0, 1]$  is fixed). Consider  $n$  independent Bernoulli trials (where each experiment does not affect any of the others) and let  $S_n$  be the number of successes. Then, we have that:

$$p_m = P(S_n = m) = \binom{n}{m} p^m (1 - p)^{n-m} \quad (m = 0, 1, \dots, n).$$

Also note that:

$$\sum_{m=0}^n p_m = [p + (1 - p)]^n = 1^n = 1$$

#### Definition: Random Variables

A **random variable** is a function  $X : \{0, 1, \dots, n\} \mapsto \mathbb{R}$ .

For example,  $S_n$  is the identity function, with  $X(m) = S_n(m) = m$ .

#### Definition: Expectation

The **expectation** of a random variable  $X$ , denoted  $EX$ , is defined as:

$$EX = \sum_{m=0}^n X(m) p_m$$

We can interpret the expectation of  $X$  as the sum over the values of  $X$ , weighted by the likelihoods. As an example, we have that:

$$ES_n = \sum_{m=0}^n S_n(m) p_m = \sum_{m=0}^n m p_m = \sum_{m=0}^n m \binom{n}{m} p^m (1 - p)^{n-m} = \dots = np$$

#### Definition: Variance & Standard Deviation

The **variance** of a random variable  $X$ , denoted  $\text{Var}(X)$ , is defined as:

$$\text{Var}(X) = E[(X - EX)^2] = E(X^2) - (EX)^2.$$

The **standard deviation** of  $X$ , denoted by  $\sigma_X$ , is then defined as  $\sigma_X = \sqrt{\text{Var}(X)}$ .

For example,  $\text{Var}(S_n) = E(S_n^2) - (ES_n)^2 = np(1 - p)$ , and  $\sigma_{S_n} = \sqrt{np(1 - p)}$ .

**Definition: Proportion of Successes**

Define  $X_n$  to be the proportion of successes in  $n$  independent Bernoulli trials, with  $X_n = \frac{1}{n}S_n$ . Then, we have that  $EX_n = \frac{1}{n}np = p$ ,  $\text{Var}(X_n) = \text{Var}(\frac{1}{n}S_n) = \frac{1}{n^2}\text{Var}(S_n) = \frac{1}{n}p(1-p)$ . We then have that  $\sigma_{X_n} = \sqrt{\frac{p(1-p)}{n}}$ .

Analyzing the standard deviation  $\sigma_{X_n}$ , we see that as we do more trials ( $n$  increases), the fluctuation of the proportion of successes gets smaller. This phenomena is known as the *Law of large numbers*, which states that the proportion of successes should converge to the probability of success in a single trial.  $\sigma_{X_n} \rightarrow 0$  as  $n \rightarrow \infty$  tells us this fact.



Figure 47: Visualization of how  $P(X_n)$ . Since  $\sigma_{X_n}$  (the width of the distribution) scales as  $\frac{1}{\sqrt{n}}$ , as  $n$  grows, the distribution becomes more sharply peaked around  $p$ .

**Theorem: Chebychev's Inequality**

$P(|X_n - p| > \delta) \leq \frac{1}{\sigma^2}p(1-p)\frac{1}{n}$ . Note that this inequality can be generalized to random variables in general, but here it suffices to consider the inequality just for the case of  $X_n$ .

**Proof**

We have that:

$$\begin{aligned} P(|X_n - p| > \delta) &= \sum_{m: |\frac{m}{n} - p| > \sigma} p_m \leq \sum_{m=0}^n \left| \frac{\frac{m}{n} - p}{\sigma} \right|^2 p_m = \frac{1}{\sigma^2} \sum_{m=0}^n \left( \frac{m}{n} - p \right)^2 p_m = \frac{1}{\sigma^2} \text{Var}(X_n) \\ &= \frac{1}{\sigma^2} p(1-p) \frac{1}{n}. \end{aligned}$$

where in the first inequality we use the fact that  $\left| \frac{\frac{m}{n} - p}{\sigma} \right|^2 \geq 1$  and we hence add non-negative terms to the sum, and in the second to last equality we invoke the definition of the variance.  $\square$

With the machinery of basic probability established, we move to the proof of the Weierstrass theorem.

### Proof

Take  $p = x \in [0, 1]$ . We then have that  $p_m = \binom{n}{m} x^m (1-x)^{n-m}$ . Let  $P_n(x) = E f(x_n) = \sum_{m=0}^n f\left(\frac{m}{n}\right) p_m$  (why? as we take  $n$  large, we have that  $x_n \rightarrow x$ , so  $f(x_n) \rightarrow f(x)$ , showing that  $P_n(x)$  approximates  $f(x)$  well). We note that  $\sum_{m=0}^n f\left(\frac{m}{n}\right) p_m$  is a polynomial in  $x$  of degree  $n$ . This is our candidate for a uniformly convergent polynomial. We then have that:

$$f(x) - P_n(x) = \sum_{m=0}^n \left( f(x) - f\left(\frac{m}{n}\right) \right) p_m$$

We will show that this is small by dividing it into two parts. For  $\sigma > 0$ , we have that:

$$\begin{aligned} |f(x) - P_n(x)| &\leq \sum_{m: \left| \frac{m}{n} - x \right| \leq \sigma} \left| f(x) - f\left(\frac{m}{n}\right) \right| p_m + \sum_{m: \left| \frac{m}{n} - x \right| > \sigma} \left| f(x) - f\left(\frac{m}{n}\right) \right| p_m \\ &\leq \sum_{m: \left| \frac{m}{n} - x \right| \leq \sigma} \left| f(x) - f\left(\frac{m}{n}\right) \right| p_m + \sum_{m: \left| \frac{m}{n} - x \right| > \sigma} 2M p_m \end{aligned}$$

where  $M = \sup \{ f(x) : x \in [0, 1] \}$ . Let  $\epsilon > 0$ . we choose  $\delta > 0$  such that  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{2}$ . This choice is possible by the uniform continuity of  $f$  (it is a continuous (polynomial) function on a compact set  $([0, 1])$ ). Then, for the first term above we have that:

$$\sum_{m: \left| \frac{m}{n} - x \right| \leq \sigma} \left| f(x) - f\left(\frac{m}{n}\right) \right| p_m \leq \frac{\epsilon}{2} \sum_{m=0}^n p_m = \frac{\epsilon}{2}.$$

For the second term, we apply Chebychev's inequality to get:

$$\sum_{m: \left| \frac{m}{n} - x \right| > \sigma} 2M p_m \leq 2M \frac{1}{\delta^2} \frac{x(1-x)}{n}.$$

Since  $x(1-x) \leq \frac{1}{4}$  for  $x \in [0, 1]$  we have:

$$\sum_{m: \left| \frac{m}{n} - x \right| > \sigma} 2M p_m \leq 2M \frac{1}{\delta^2} \frac{x(1-x)}{n} \leq \frac{M}{2\delta^2} \frac{1}{n}.$$

Now, choose  $n$  such that  $n > N \geq \frac{4\delta^2}{M\epsilon}$ . We then have that:

$$\frac{M}{2\delta^2} \frac{1}{n} < \frac{\epsilon}{2}$$

Then, we have that:

$$|f(x) - P_n(x)| \leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

which proves the claim. □

Our conclusion is that  $P_n(x)$  is very close to  $f(x)$ . In the above proof, we split up the sum into two parts, and used different methods to obtain nice estimates/bounds on each. The next topic we will look at is generalizing this theorem; we will be building up to Rudin 7.32 (Stone-Weierstrass).

## **8 Some Special Functions**

### **8.1 Power Series, Revisited**

### **8.2 The Exponential Function**

### **8.3 The Logarithm**

### **8.4 Sine and Cosine**

### **8.5 Fourier Series**

## **9 Functions of Several Variables**

### **9.1 Banach Fixed Point Theorem**

### **9.2 Differentiation of Functions of Several Variables**

### **9.3 The Inverse Function Theorem**