

MODELING BIOLOGICAL SYSTEMS

INSTRUCTIONS TP - TD6

Project 2 : a continuous model

Let us consider a cell density ρ of low grade glioma LGG (grade I/II brain tumors).
General diffusion equations, as heat equation, follow:

$$\frac{\partial \rho}{\partial t} = D \Delta \rho$$

where D is a diffusion rate.

We want to apply this equation to glioma cell density, assuming the brain is homogeneous. For our biological application, D might depends on cell positions in the brain (diffusivity should be heterogeneous), but it will be assumed constant here.

$\Delta \rho$ here denotes the Laplacian of the ρ function .

- A. To introduce a proliferation term, it is generally accepted to add a proliferation rate κ of this cell density ρ . To limit an infinite number of cells due to an infinite proliferation, we will coupled it with a saturation term $1 - \rho$ to obtain finally a logistic function of ρ . Write the new equation for ρ .
- B. What is the general shape of a low grade tumor? What spatial coordinates are you likely to use?
- C. What symmetrical considerations could you make on ρ ? We assume ρ depending only on its radial distance r , (i.e. $\rho = \rho(r)$).

Show that ρ follows the Diffusion Proliferation Saturation DPS equation :

$$\frac{\partial \rho}{\partial t} = D \left(\frac{\partial^2 \rho}{\partial r^2} + \frac{2}{r} \frac{\partial \rho}{\partial r} \right) + \kappa \rho (1 - \rho) \quad (E)$$

- D. Set $u = \rho \cdot r$ and find a new equation (E') depending on the variable u . We set $u(r = 0, t) = 0$ and $\frac{\partial u}{\partial r}(r = 0, t) = \rho(r = 0, t)$.
- E. This equation (E') is not analytically solvable (because of the nonlinear proliferation saturation term). We will use a numerical approach to solve it. Instead of using a classical but unstable Euler's method, we will use a Crank-Nicolson method* that was first invented to solve the heat equation.

To avoid chaotic oscillations, we will assume that $(u_j^n)^2 \approx u_j^n u_j^{n+1}$ for $n > 1$.

Starting from your equation (E'), show that the numerical scheme of the function u can be written as the following :

$$u_j^{n+1} = u_j^n \left(1 - 2\alpha + \kappa \Delta t - \kappa \frac{\Delta t}{r_j} u_j^{n+1} \right) + \alpha (u_{j+1}^n - u_{j-1}^n)$$

where $\alpha = \frac{D\Delta t}{(\Delta r^2)}$, $r_j = j\Delta r$ and $t_n = n\Delta t$, with $\rho_j^n = \frac{u_j^n}{r_j}$. Note that $u_j^n = u_{-j}^n$.

This expression can be written only depending on u_j^{n+1} as the following :

$$u_j^{n+1} = \frac{1}{\left(1 + \frac{\kappa \Delta t u_j^n}{r_j} \right)} \left[u_j^n (1 - 2\alpha + \kappa \Delta t) + \alpha (u_{j+1}^n + u_{j-1}^n) \right]$$

- F. Use your numerical scheme to compute your solution in your favorite programming language. We want ρ with respect to t (in years).

This link could help you especially if you use Python :

<https://levelup.gitconnected.com/solving-2d-heat-equation-numerically-using-python-3334004aa01a>

For this part, you could use these informations :

- $\Delta r = 0.02$ mm, as the average size of a LGG cell ($\approx 20 \mu\text{m}$) and will watch the evolution of ρ up to 100 mm;
- $\Delta t = 0.001$ year, up to a many years evolution;
- $\rho(r_0, t_0) = 1$;
- final tumor expected : between 10^3 and 10^4 cells for a small tumor;
- initial condition : one single mutated cell;
- in the case of a compact tumor, $\kappa = 10\text{year}^{-1}$ and $D = 0.01\text{mm}^2/\text{year}$;
- in the case of a diffuse tumor, $\kappa = 1.0\text{year}^{-1}$ and $D = 0.7\text{mm}^2/\text{year}$;
- edge growth velocity approximated : $v = 2\sqrt{\kappa D}$

Extra questions:

We define the radius of the tumor R :

$$R(\rho^*, t) = \sqrt{4D(t + t_0) \cdot \left[\kappa t - \frac{3}{2} \ln\left(\rho^* \frac{t + t_0}{t_0}\right) \right]}$$

where $\rho(r = R, t) = \rho^*$, the density at the edge of the tumor, and $t_0 = r_0^2/(4D)$

G. The detection threshold is known to be around $\rho_{\text{detected}} = 2\%$. Can you find the duration $t_{\text{detection}}$ before the glioma cells get detected ?

Appendix :

$$u(r, t) \leftrightarrow u_j(r_j, t_n)$$

$$\frac{\partial u}{\partial t} \leftrightarrow \frac{u_j^{n+1} - u_j^n}{\Delta t}$$

$$\frac{\partial^2 u}{\partial r^2} \leftrightarrow \frac{1}{(\Delta r)^2} [u_{j+1}^n - 2u_j^n + u_{j-1}^n]$$

Butcher's tableau of Crank-Nicolson scheme:

0	0	0
1	1/2	1/2
<hr/>		
	1/2	1/2