# ■ KisanMitra AI Crop Report

Automated Farm Health Advisory Multi-Agent System

Date: 05-Jul-2026
Report Ref: KM-DX-031202
Status: Complete / Verified

## I. Farmer & Land Profile

| | | | |
|---|---|---|---|
| **Farmer Name:** | Test Farmer AP | **Location:** | Amaravati, Guntur, Andhra Pradesh |
| **Land Size:** | 3.5 Acres | **Crop Type:** | Cotton (Primary) |
| **Soil Type:** | Black Cotton Soil | **System Config:** | Simulated AI Agent Execution Mode |

## II. Health Diagnosis Summary

| | |
|---|---|
| **Target Plant Crop:** | Cotton |
| **Diagnosed Disease:** | **Cotton Leaf Curl Virus (CLCuV)** |
| **Classification Severity:** | **Moderate** |

## III. Prescribed Treatment Advisory

### ■ Organic / Biological Control Plan

Biological/Organic Control: Spray neem seed kernel extract (5%) to repel whiteflies. Install yellow sticky traps (1015 per acre) to monitor vector population.

### ■ Chemical Control Schedule

Chemical Control: Control whiteflies using Diafenthiuron 50% WP at 240 g/acre or Imidacloprid 17.8% SL at 100 ml/acre.

### ■ Agronomic Preventive Recommendations

Agronomic Preventive Advice: Keep fields weedfree as weeds act as alternative hosts. Remove and destroy infected plants during early stages.

## IV. Full Agent Advisory Remarks

*DIAGNOSIS REPORT (Simulator Mode):*
*- Target Crop: Cotton*
*- Diagnosed Disease: Cotton Leaf Curl Virus (CLCuV)*

*- Confidence Score: 88.0%*
*- Severity: Moderate*
*- Symptoms Observed: Upward or downward curling of leaf margins, thickening of veins, and formation of leaf-like enations on the underside of leaves.*
*- Cause: Viral pathogen transmitted by the Silverleaf Whitefly (Bemisia tabaci).*
*- Biological/Organic Control: Spray neem seed kernel extract (5%) to repel whiteflies. Install yellow sticky traps (10-15 per acre) to monitor vector population.*
*- Chemical Control: Control whiteflies using Diafenthiuron 50% WP at 240 g/acre or Imidacloprid 17.8% SL at 100 ml/acre.*
*- Agronomic Preventive Advice: Keep fields weed-free as weeds act as alternative hosts. Remove and destroy infected plants during early stages.*

---