

KisanMitra AI Crop Report

Automated Farm Health Advisory Multi-Agent System

Date: 05-Jul-2026
Report Ref: KM-DX-F74D68
Status: Complete / Verified

I. Farmer & Land Profile

Farmer Name:	Banavathu Prasanth Kumar	Location:	Chevuluru, Guntur, Andhra Pradesh
Land Size:	3.5 Acres	Crop Type:	Cotton (Primary)
Soil Type:	Black Cotton Soil	System Config:	Simulated AI Agent Execution Mode

II. Health Diagnosis Summary

Target Plant Crop:

Cotton

Diagnosed Disease:

Cotton Leaf Curl Virus (CLCuV)

Classification

Severity: Moderate

CROP DISEASES

Common Diseases in Crops and Their Symptoms

RICE

Blast



Spindle-shaped lesions on leaves and neck.

WHEAT

Leaf Rust



Orange-brown pustules on leaves.

MAIZE

Northern Corn Leaf Blight



Long, gray-green lesions on leaves.

POTATO

Late Blight



Dark, water-soaked lesions on leaves and stems.

TOMATO

Early Blight



Dark spots with concentric rings on leaves.

CHILLI

Anthracnose



Dark, sunken lesions on fruits and stems.

GRAPES

Powdery Mildew



White powdery growth on leaves and berries.

COTTON

Bacterial Blight



Irregular water-soaked lesions on leaves.

SOYBEAN

Rust



Reddish-brown pustules on leaves.

CITRUS

Canker



Raised, corky lesions on leaves and fruits.

III. Prescribed Treatment Advisory

Organic / Biological Control Plan

Biological/Organic Control: Spray neem seed kernel extract (5%) to repel whiteflies. Install yellow sticky traps (1015 per acre) to monitor vector population.

Chemical Control Schedule

Chemical Control: Control whiteflies using Diafenthiuron 50% WP at 240 g/acre or Imidacloprid 17.8% SL at 100 ml/acre.

Agronomic Preventive Recommendations

Agronomic Preventive Advice: Keep fields weedfree as weeds act as alternative hosts. Remove and destroy infected plants during early stages.

IV. Full Agent Advisory Remarks

DIAGNOSIS REPORT (Simulator Mode):

- Target Crop: Cotton

- Diagnosed Disease: Cotton Leaf Curl Virus (CLCuV)

- Confidence Score: 88.0%
- Severity: Moderate
- Symptoms Observed: Upward or downward curling of leaf margins, thickening of veins, and formation of leaf-like enations on the underside of leaves.
- Cause: Viral pathogen transmitted by the Silverleaf Whitefly (*Bemisia tabaci*).
- Biological/Organic Control: Spray neem seed kernel extract (5%) to repel whiteflies. Install yellow sticky traps (10-15 per acre) to monitor vector population.
- Chemical Control: Control whiteflies using Diafenthiuron 50% WP at 240 g/acre or Imidacloprid 17.8% SL at 100 ml/acre.
- Agronomic Preventive Advice: Keep fields weed-free as weeds act as alternative hosts. Remove and destroy infected plants during early stages.

Disclaimer: This report is automatically generated by KisanMitra multi-agent advisory system. Recommendations are based on visual classification models and retrieved scheme databases. Consult local Krishi Vigyan Kendra (KVK) for final chemical application approvals.