

Workshop 3

Sources of Errors

FIT 3139

Computational Modelling and Simulation



COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

WARNING


This material has been reproduced and communicated to you by or on behalf of Monash University pursuant to Part VB of the Copyright Act 1968 (the Act). The material in this communication may be subject to copyright under the Act. Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act.

Do not remove this notice.




What's the surface area of the earth?


Earth / Radius

6,371 km



People also search for

 Mars 3.39K km	 Moon 1.737K km	 Sun 695.508K ...
--	---	---



https://www.youtube.com/watch?v=6EzQEdBX_30&t=306s

$\approx 510064472 \text{ km}^2$

- **Modelling error:**
the earth is not a sphere.
- **Data errors:**
Measuring earth radius is error prone.
- **Computational errors:**
the number π has to be truncated in a computer

Sources of errors...

- **Modelling error:**

Arise from simplifying assumptions inherent to models.

- **Data errors:**

Arising from:

- **Empirical measurements**, which have physical limitations.
- **Previous computations**, which “propagate” an error

- **Computational Errors:**

Arising from:

- **Truncation errors**, e.g, approximate an infinite sum with a finite sum.
- **Rounding error**, because computers can only cope with finite precision.

Measuring error...



Measure

9999 cm

7 cm

True value

10000 cm

10 cm

Error

1 cm

3 cm

Errors often need to be put in **context**, which we do by normalising.

Measuring error...



Measure

9999 cm

7 cm

True value

10000 cm

10 cm

**Absolute
Error**

1 cm

3 cm

**Relative
Error**

$(10000 - 9999) / 10000$
0.01%

$(10 - 7) / 10$
30%

Absolute Error

Absolute error = | Approximate - True value |

Relative Error

Relative error = (Absolute error / True value) x 100%



Data error vs Computational Error

$$\sin\left(\frac{\pi}{8}\right)$$

$$f(x) = \sin(x)$$

$$f(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$$

$$\tilde{f}(x) \approx x \quad \pi \approx 3$$



$$\sin\left(\frac{\pi}{8}\right) \xrightarrow{\text{red}} \sin\left(\frac{3}{8}\right) \xrightarrow{\text{orange}} \frac{3}{8} = 0.37500$$

Data error: $f(\tilde{x}) - f(x) = \sin\left(\frac{3}{8}\right) - \sin\left(\frac{\pi}{8}\right) = -0.0164$

Computation Error: $\tilde{f}(\tilde{x}) - f(\tilde{x}) = \frac{3}{8} - \sin\left(\frac{3}{8}\right) = 0.0087$

Total Error: $\tilde{f}(\tilde{x}) - f(x) = \frac{3}{8} - \sin\left(\frac{\pi}{8}\right)$
 $= 0.37500 - 0.38267 = -0.0077$

Total error = Data error + Computational Error

- Let f be a true function, and x a true value.
- Let \tilde{f} be an approximation of f and \tilde{x} be an inexact measurement of x

$$\begin{aligned}\text{Total error} &= \tilde{f}(\tilde{x}) - f(x) \\ &= \underbrace{\tilde{f}(\tilde{x}) - f(\tilde{x})}_{\substack{\downarrow \\ \text{Computational} \\ \text{Error}}} + \underbrace{f(\tilde{x}) - f(x)}_{\substack{\downarrow \\ \text{Data} \\ \text{Error}}}\end{aligned}$$

Forward and Backward Error

Forward error: Difference between **computed** and **true value**

$$f(x) = \sqrt{x} \qquad \tilde{y} = 1.4 \qquad y = \sqrt{2} = 1.41421\dots$$

$$|\Delta y| = |\tilde{y} - y| = |1.4 - 1.41421\dots| \approx 0.0142$$

Backward error:

Discrepancy in the input that would lead to the observed discrepancy in output

$$|\Delta x| = |\tilde{x} - x| = |1.96 - 2| = 0.04$$

$$\tilde{x} = f^{-1}(\tilde{y}) \leftarrow \boxed{\text{Input } \tilde{x} \text{ for which } f \text{ produces } \tilde{y}}$$

$$\sqrt{\tilde{x}} = 1.4$$


$$\tilde{x} = 1.4^2 = 1.96$$

Forward and Backward Error

Forward error: Difference between **computed** and **true value**

Backward error:

Discrepancy in the input that would lead to the observed discrepancy in output




Some functions deal better
than others...

with propagation of error

Sensitivity: A qualitative statement of propagated data error.

Conditioning: Quantitive measure of sensitivity.



A problem is **insensitive** or **well-conditioned** if a given relative change in the input data causes a reasonably commensurate **change in the solution**.

$$\text{Condition number} = \frac{\left| \frac{\Delta y}{y} \right|}{\left| \frac{\Delta x}{x} \right|} = \frac{\text{Relative forward error}}{\text{Relative backward error}}$$

For an ill-conditioned problem, condition number $\gg 1$

Approximating the condition number

$$f'(x) \approx \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

$$\left| \frac{\Delta y}{y} \right| = \left| \frac{\tilde{y} - y}{y} \right| \approx \frac{f'(x)\Delta x}{f(x)} \longrightarrow \text{relative forward error}$$

$$\left| \frac{\Delta x}{x} \right| \longrightarrow \text{relative backward error}$$

$$\frac{\left| \frac{f'(x)\Delta x}{f(x)} \right|}{\left| \frac{\Delta x}{x} \right|} \approx \left| \frac{xf'(x)\Delta x}{f(x)\Delta x} \right| = \left| \frac{xf'(x)}{f(x)} \right|$$

Stability is to an algorithm
what **conditioning** is to a problem.....