

Workshop 13

Runge Kutta

FIT 3139
Computational Science



COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

WARNING


This material has been reproduced and communicated to you by or on behalf of Monash University pursuant to Part VB of the Copyright Act 1968 (the Act). The material in this communication may be subject to copyright under the Act. Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act.

Do not remove this notice.

$$y_{i+1} = y_i + h\phi$$

Error in Euler method

$$y_{i+1} = y_i + hf(x_i, y_i)$$



Predictor: Slope 1

$$f(x_i, y_i)$$


Corrector: Slope 2

$$f(x_{i+1}, y_{i+1})$$

$$f(x_i + h, \underline{\quad})$$


$$f(x_i + h, y_i + hf(x_i, y_i))$$


$$y_{i+1} = y_i + \frac{h}{2} \left(f(x_i, y_i) + f(x_i + h, y_i + hf(x_i, y_i)) \right)$$




Heun's method

$$y_{i+1} = y_i + \frac{h}{2} \left(f(x_i, y_i) + f(x_i + h, y_i + hf(x_i, y_i)) \right)$$
$$x_{i+1} = x_i + h$$






$$y' = y$$



$$y' = y$$

Midpoint method




$$f(x_i, y_i)$$

$$y_{i+1} = y_i + hf(x_i, y_i)$$

$$f\left(x_i + \frac{h}{2}, \right)$$

$$f\left(x_i + \frac{h}{2}, y_i + \frac{h}{2}f(x_i, y_i)\right)$$

Midpoint method



$$f\left(x_i + \frac{h}{2}, y_i + \frac{h}{2} f(x_i, y_i)\right)$$

$$y_{i+1} = y_i + h\phi$$

$$y_{i+1} = y_i + hf\left(x_i + \frac{h}{2}, y_i + \frac{h}{2} f(x_i, y_i)\right)$$

$$x_{i+1} = x_i + h$$

So far...

Method	
Euler	RK1
Heun's	RK2
Midpoint	RK2

$$y_{i+1} = y_i + \underbrace{hy'(x_i)}_{RK1} + \underbrace{\frac{h^2}{2}y''(x_i) + \dots}_{RK2}$$

$$y_{i+1} \approx y_i + hy'(x_i) + \frac{h^2}{2}y''(x_i)$$

Runge-Kutta 2

General formula

$$\frac{dy}{dx} = f(x, y)$$

$$y_{i+1} \approx y_i + hy'(x_i) + \frac{h^2}{2}y''(x_i)$$

$$y_{i+1} \approx y_i + h\left(y'(x_i) + \frac{h}{2}y''(x_i)\right)$$

$$y_{i+1} \approx y_i + h\left(f(x_i, y_i) + \frac{h}{2}\boxed{f'(x_i, y_i)}\right) \longrightarrow \textbf{unknown}$$

Can be approximated as follows...

$$y_{i+1} = y_i + h(ak_1 + bk_2)$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + \alpha h, y_i + \beta k_1 h)$$

$$a + b = 1$$

$$b\alpha = \frac{1}{2}$$

$$b\beta = \frac{1}{2}$$

Heun's method

General RK2

$$y_{i+1} = y_i + h(ak_1 + bk_2)$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + \alpha h, y_i + \beta k_1 h)$$

$$a + b = 1$$

$$b\alpha = \frac{1}{2} \quad b\beta = \frac{1}{2}$$

$$y_{i+1} = y_i + \frac{h}{2} \left(f(x_i, y_i) + f(x_i + h, y_i + hf(x_i, y_i)) \right)$$

$$a = \frac{1}{2} \quad b = \frac{1}{2} \quad \alpha = 1 \quad \beta = 1$$

$$y_{i+1} = y_i + h \left(\frac{1}{2}k_1 + \frac{1}{2}k_2 \right)$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + h, y_i + k_1 h)$$

Midpoint method

General RK2

$$y_{i+1} = y_i + h(ak_1 + bk_2)$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + \alpha h, y_i + \beta k_1 h)$$

$$a + b = 1$$

$$b\alpha = \frac{1}{2} \quad b\beta = \frac{1}{2}$$

$$y_{i+1} = y_i + hf\left(x_i + \frac{h}{2}, y_i + \frac{h}{2}f(x_i, y_i)\right)$$

$$a = 0 \quad b = 1 \quad \alpha = \frac{1}{2} \quad \beta = \frac{1}{2}$$

$$y_{i+1} = y_i + hk_2$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f\left(x_i + \frac{h}{2}, y_i + k_1 \frac{h}{2}\right)$$

Ralston's method

General RK2

$$y_{i+1} = y_i + h(ak_1 + bk_2)$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + \alpha h, y_i + \beta k_1 h)$$

$$a + b = 1$$

$$b\alpha = \frac{1}{2} \quad b\beta = \frac{1}{2}$$

$$a = \frac{1}{3} \quad b = \frac{2}{3} \quad \alpha = \frac{3}{4} \quad \beta = \frac{3}{4}$$

$$y_{i+1} = y_i + h\left(\frac{1}{3}k_1 + \frac{2}{3}k_2\right)$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f\left(x_i + \frac{3h}{4}, y_i + k_1 \frac{3h}{4}\right)$$

Homework: graphic interpretation

Runge-Kutta 2

$$y_{i+1} = y_i + h(ak_1 + bk_2)$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + \alpha h, y_i + \beta k_1 h)$$

$$a + b = 1$$

$$b\alpha = \frac{1}{2} \quad b\beta = \frac{1}{2}$$

Please refer to lecture recording for a full derivation.

Named methods

Method	
Heun's	RK2
Midpoint	RK2
Ralston's	RK2

You can choose any b.....

Not examinable

Runge Kutta 4

$$y_{i+1} = y_i + hy'(x_i) + \frac{h^2}{2}y''(x_i) + \frac{h^3}{3!}y'''(x_i) + \frac{h^4}{4!}y''''(x_i)\dots$$

Not examinable

Runge Kutta 4

$$y_{i+1} = y_i + \frac{1}{6} (k_1 + 2k_2 + 2k_3 +$$

$$k_4 = f(x_i, y_i)$$

$$k_2 = f\left(x_i + \frac{1}{2}h, y_i + \frac{1}{2}k_1\right)$$

$$k_3 = f\left(x_i + \frac{1}{2}h, y_i + \frac{1}{2}k_2\right)$$

$$k_4 = f(x_i + h, y_i + k_3)$$

Not examinable

Runge Kutta 4

$$y_{i+1} = y_i + \frac{1}{8} (k_1 + 3k_2 + 3k_3 +$$

$$k_4 = f(x_i, y_i)$$

$$k_2 = f\left(x_i + \frac{1}{3}h, y_i + \frac{1}{3}k_1h\right)$$

$$k_3 = f\left(x_i + \frac{2}{3}h, y_i - \frac{1}{3}k_1h + \frac{2}{3}k_2h\right)$$

$$k_4 = f\left(x_i + h, y_i + k_1h - k_2h + k_3h\right)$$

Not examinable

Runge Kutta 4

$$y_{i+1} = y_i + \frac{1}{6}(k_1 + 2k_2 + 2k_3 + k_4)h$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f\left(x_i + \frac{1}{2}h, y_i + \frac{1}{2}k_1h\right)$$

$$k_3 = f\left(x_i + \frac{1}{2}h, y_i + \frac{1}{2}k_2h\right)$$

$$k_4 = f(x_i + h, y_i + k_3h)$$

Notes...

- The RK methods can be generalised for systems of several ODE.
- State of the art algorithms common in computational science libraries: RK4, RK5 and variations.