

Prepared by Julian García, based on material by  
David Albrecht and María García de la Banda

# Workshop 13

## Intro to Stochastic Models: Montecarlo

# FIT 3139 Computational Modelling and Simulation



**MONASH** University  
Information Technology

COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

WARNING


This material has been reproduced and communicated to you by or on behalf of Monash University pursuant to Part VB of the Copyright Act 1968 (the Act). The material in this communication may be subject to copyright under the Act. Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act.  
Do not remove this notice.

# Outline

- Basic elements of probability
- Buffon needle problem:
  - Axiomatic
  - Montecarlo
- (Pseudo) random numbers


# Deterministic models

- No stochastic (i.e., **random**) components.
- Every state of the variables is *uniquely* determined by previous states of the variables.
- A set of input **always** leads to same outputs.
- Same initial conditions, same output



# Stochastic models

- Inherent **randomness**
- State of variables are described by **probability distributions**, not unique values.
- Same initial conditions lead to different outcomes.
- **Stochastic Simulations**



An **experiment**



Let's think of something that has a  
“random” outcome...

(and think of some useful definitions on the way)

7

## An **experiment**



$S = \{ \text{ all possible orderings of } (1,2,3,4,5,6,7) \}$



**sample  
space**

(4,1,2,5,3,7,6)

→ **outcome**

of the random experiment

---

$A = \{ \text{all outcomes starting with 4 } \}$  → **event**

Any subset of the sample space is an event.

---

For any two events  $A$  and  $B$ , the event  $A \cup B$  consists of all events in either  $A$  or  $B$ . It's called the **Union**

$\bigcup_{i=1}^n A_i$  it's the union of all outcomes in any of  $A_1, A_2, \dots, A_n$

---

Similarly for **intersections** of events.

# Axioms of probability

Consider an **experiment** with **sample space**  $S$ , for each event  $A$ , there is a number  $P(A)$ , called **the probability** of the event  $A$ , which is in accord with the following axioms:

$$0 \leq P(A) \leq 1$$

the probability that the outcome  
of the experiment lies in  $A$  is some number  
between 0 and 1

$$P(S) = 1$$

with probability 1 the outcome of the  
experiment is in the sample space

$$P\left(\bigcup_{i=1}^n A_i\right) = \sum_{i=1}^n P(A_i)$$

$$i = 1, 2, \dots, \infty$$

for any sequence of **mutually exclusive** events  
 $\{A_i\}$  the probability that at least one of these  
events occurs is equal to the sum of their  
respective probabilities

These axioms, can be used to prove a variety of results....

Example:

$$S = A \cup A^c$$

$$1 = P(S) = P(A \cup A^c) = P(A) + P(A^c)$$

axiom 2

axiom 3

$$P(A^c) = 1 - P(A)$$

“The probability that an event does not occur is 1 minus the probability that it does”

# Empirical probability

If a **random experiment** is repeated  $n$  times and  $n_a$  is the number of times an event  $E$  occurs, then the relative frequency of occurrence of event  $E$  is  $\frac{n_a}{n}$ .

This relative frequency converges to the probability of  $E$  as  $n \rightarrow \infty$ :

$$Pr(E) = \lim_{n \rightarrow \infty} \frac{n_a}{n}$$

**This is the basis of the Montecarlo technique, to be discussed..**



## Experiment

Throw two dice.

$$S = \{(d_1, d_2), 0 \leq d_i \leq 6, d_i \in \mathbb{Z}, i = 1, 2\}$$

Say I am interested, in the sum of the dice  $X$

$X$  is a random variable

# Random variable

A **quantity** of interest determined by the result of an experiment.

We say a random variable is **discrete**, if it can take a finite (or at most countable) number of values.

$p(x) = P\{X = x\}$  ————— Probability mass function

[for a discrete RV]

We say a random variable is **continuous**, if the set of possible values is an interval.


$P\{X \in C\} = \int_C f(x)dx$        $f(x)$  ————— Probability density function

[for a continuous RV]



# Monte Carlo method

- An (approximate) method of simulation based on **random sampling**.
- Typically used to attack some **intractable models**, integrals, or sums.
- Initially used in 1930's by Enrico Fermi (*neutron diffusion*), popularised by Stanislaw Ulam in the 1940's at Los Alamos Nat Lab.
- **Computers** make this method extremely relevant.
- **Used extensively** in mathematical, computational, statistical modelling




$$A = \pi r^2$$

Use Montecarlo to estimate  $\pi$

# Use Montecarlo to estimate $\pi$

$$A = \pi r^2$$

$$A = \frac{\pi}{4}$$



throw *many* random points  
ratio of inside/total samples  
must equal  $\pi/4$

## Montecarlo method:

1. Define a domain for the samples.
2. Generate samples (using a Random Number Generator).
3. Apply a deterministic computation on the samples.
4. Aggregate the results.







Georges-Louis Leclerc, Comte de Buffon

# Buffon's needle




An infinite family of long vertical lines are spaced one unit apart in the  $(x,y)$  plane. If a needle of length  $r > 0$  (and negligible width) is dropped at random onto the plane, what is the probability that it will land crossing at least one line?





$$u + r \cos \theta < 1$$


$$u = 1 - r \cos \theta$$




$$u + r \cos \theta > 1$$

# Sample space


$$u = 1 - r \cos \theta$$




# Montecarlo simulation



# Montecarlo simulation



- Randomly sample:  $(\theta_1, u_1), (\theta_2, u_2), \dots, (\theta_n, u_n)$
- If some  $(\theta_i, u_i)$  falls above the curve (i.e.,  $u_i > 1 - r \cos \theta_i$ ) call it a **hit**, otherwise a **miss**.
- Probability of the needle crossing can be approximated as number of **hits** over number of samples ( $n$ )



$$\text{Probability of not crossing} = \frac{\text{Area of sector } \theta}{\text{Area of circle}} = \frac{\theta}{\pi}$$

$$\text{Probability of crossing} = \frac{\text{Area of sector } u - \text{Area of sector } \theta}{\text{Area of circle}} = \frac{u - \theta}{\pi} = \frac{2r}{\pi}$$

$$\square = \pi$$

$$\text{Area of sector } \theta = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} 1 - r \cos \theta d\theta$$

$$\text{Area of sector } u = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \theta - r \sin \theta d\theta$$

$$\text{Area of circle} = \pi - 2r$$

# How do we generate random numbers?

<https://www.random.org/>