

Workshop 11

Solving ODEs

- FIT3139: Computational Modelling and Simulation



COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

WARNING

This material has been reproduced and communicated to you by or on behalf of Monash University pursuant to Part VB of the Copyright Act 1968 (the Act). The material in this communication may be subject to copyright under the Act. Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act.


Do not remove this notice.

Odeint: Numerical integration

$$\frac{dx}{dt} = f(x, t)$$

$$x(t) = \int f(x)$$

$$x(0) = x_0$$



Terminology

The diagram shows the differential equation $2x^2 \left(\frac{dy}{dx} \right)^3 - 3y = 0$. A red box highlights the dy in the derivative, with an arrow pointing to the text "order 1". A blue box highlights the superscript 3, with an arrow pointing to the text "degree 3".


$$2x^2 \left(\frac{dy}{dx} \right)^3 - 3y = 0$$

The **order** of an ordinary differential equation is determined by the order of the highest appearing derivative

The **degree** of an ODE is the power to which the highest-order derivative is raised.

This **ODE**, where the O is for **ordinary**; i.e., no partial derivatives.

Terminology

order 4 *degree 1*


$$5x \frac{d^4 y}{dx^4} + 2x^2 \left(\frac{dy}{dx} \right)^3 - 3y =$$

notational shorthand $y' = \frac{dy}{dx}$ $y'' = \frac{d^2 y}{dx^2}$ $y^{(n)} = \frac{d^n y}{dx^n}$

$$5xy'''' + 2x^2(y')^3 - 3y =$$

Linear differential equation

$$a_0(x)y + a_1(x)y' + a_2(x)y'' + a_3(x)y''' + \cdots + a_n y^{(n)} + b(x) = 0$$

- $a_1(x)$, $a_2(x)$... $a_n(x)$, and $b(x)$ are arbitrary differentiable functions that do not need to be linear.
- Note that derivatives are degree 1.
- A solution, is a function $y(x)$ that satisfies the equation.
- Anything else is **non-linear**.

An **initial value problem** involves an ODE together with a specified value, called the **initial condition**, of an unknown function at a given point in the domain of the solution.

We'll focus on **first order** initial value problems.

$$y' = \frac{dy}{dx} = f(x, y)$$

$$(x_0, y_0) \quad \text{i.e.,} \quad y_0 = y(x_0)$$

$y(x)$ is unknown.




Euler's method

known...

$$y' = f(x, y), (x_0, y_0)$$

$y(x)$ is unknown.



$$\tan(\theta) = \frac{r}{h}$$


$$f(x_0, y_0) = \frac{r}{h}$$

$$hf(x_0, y_0) = r$$

$$x_1 = x_0 + h$$

$$y_1 = y_0 + r$$

$$y_1 = y_0 + hf(x_0, y_0)$$



$$x_1 = x_0 + h$$

$$y_1 = y_0 + hf(x_0, y_0)$$

$$x_2 = x_1 + h$$

$$y_2 = y_1 + hf(x_1, y_1)$$

$$x_3 = x_2 + h$$


$$y_3 = y_2 + hf(x_2, y_2)$$

⋮

$$x_n = x_{n-1} + h$$

$$y_n = y_{n-1} + hf(x_{n-1}, y_{n-1})$$

$(x_0, y_0), (x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ approximates $y(x)$



Another view...

Remember Taylor...

$$f(x + h) = f(x) + f'(x)h + f''(x)\frac{h^2}{2!} + \dots$$

known...

$$y' = f(x, y)$$

$y(x)$ is unknown.

$$f(x + h) = f(x) + f'(x)h + f''(x)\frac{h^2}{2!} + \dots$$

$$y_{i+1} = y(x_{i+1}) = y(x_i + h)$$

Taylor

$$= y(x_i + h) = y(x_i) + hy'(x_i) + \boxed{\frac{h^2}{2}y''(x_i) + \dots}$$

small h

$$y_{i+1} \approx y(x_i) + hy'(x_i)$$

$$y_{i+1} \approx y_i + hf'(x_i, y_i)$$

$$y_{i+1} \approx y_i + hf(x_i, y_i)$$

$$x_{i+1} \approx x_i + h$$

Start at initial
value and iterate

known...

$$y' = f(x, y)$$

$$(x_0, y_0)$$

$y(x)$ is unknown.

$$y_{i+1} \approx y_i + hf(x_i, y_i)$$

$$x_{i+1} \approx x_i + h$$

Start at initial
value and iterate

Example

$$\frac{dP}{dt} = r \left(1 - \frac{P(t)}{K} \right) P(t)$$

Carl **Runge**



Martin **Kutta**



Runge-Kutta methods circa 1901

Runge-Kutta methods circa 1901

$$y_{i+1} = y(x_{i+1}) = y(x_i + h) = y(x_i) + hy'(x_i) + \frac{h^2}{2}y''(x_i) + \dots$$

$$\approx y(x_i) + h \left(f(x_i, y_i) + \frac{h}{2}y''(x_i) \right)$$



how is this computed

Runge-Kutta methods circa 1901

$$y_{i+1} = y_i + \underbrace{\phi}_{\text{slope}} h$$

Methods of this family
will differ in **how the slope is computed**.

$$\phi = \frac{dy}{dx} = f(x, y) \quad \text{RK1}$$

$$y_{i+1} = y_i + f(x_i, y_i)h$$

$$y_i = f(x_i, y_i)(x_{i+1} - x_i)$$

Euler

$$\frac{dy}{dx} = f(x, y)$$

$$\int dy = \int f(x, y) dx$$


$$\int_{y_i}^{y_{i+1}} dy = \int_{x_i}^{x_{i+1}} f(x, y) dx$$

$$y_{i+1} - y_i = \int_{x_i}^{x_{i+1}} f(x, y) dx$$

$$y_{i+1} = y_i + \int_{x_i}^{x_{i+1}} f(x, y) dx$$

Error in Euler method

$$y' = y$$



How to improve our prediction?

Predictor: **Slope 1**

$$f(x_i, y_i)$$


Corrector: **Slope 2**

$$f(x_{i+1}, y_{i+1})$$

$$f(x_i + h, \underline{\quad})$$

$$f(x_i + h, y_i + hf(x_i, y_i))$$


$$y_{i+1} = y_i + \frac{h}{2} \left(f(x_i, y_i) + f(x_i + h, y_i + hf(x_i, y_i)) \right)$$



Heun's method

$$y_{i+1} = y_i + \frac{h}{2} (f(x_i, y_i) + f(x_i + h, y_i + hf(x_i, y_i)))$$

$$x_{i+1} = x_i + h$$



What about Dynamical Systems?

$$\begin{aligned}\frac{dS}{dt} &= -\beta SI \\ \frac{dI}{dt} &= +\beta SI - \alpha I \\ \frac{dR}{dt} &= +\alpha I\end{aligned}$$

We know the starting values
 S_0 , I_0 , and R_0 ...

Can we use them to find
 $S(t_1)$, $I(t_1)$, and $R(t_1)$?

Recall

$$y_{i+1} \approx y_i + hf(x_i, y_i)$$
$$x_{i+1} \approx x_i + h$$

Let's apply this schema to each
Differential equation

$$\frac{dS}{dt} = -\beta SI$$
$$\frac{dI}{dt} = +\beta SI - \alpha I$$
$$\frac{dR}{dt} = +\alpha I$$



$$S_{i+1} = S_i - h\beta S_i I_i$$
$$I_{i+1} = I_i + h(\beta S_i I_i - \alpha I_i)$$
$$R_{i+1} = R_i + h\alpha I_i$$
$$t_{i+1} = t_i + h$$

Try it out....