

Workshop 4

Linear Systems

FIT 3139

Computational Modelling and Simulation



COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

WARNING

This material has been reproduced and communicated to you by or on behalf of Monash University pursuant to Part VB of the Copyright Act 1968 (the Act). The material in this communication may be subject to copyright under the Act. Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act.

Do not remove this notice.

Linear system

$$A\vec{x} = \vec{b} \quad ?$$

General elimination matrix

$$\mathbf{M}_k \mathbf{a} = \begin{bmatrix} 1 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 & \cdots & 0 \\ 0 & \cdots & -m_{k+1} & 1 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & -m_n & 0 & \cdots & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_k \\ a_{k+1} \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} a_1 \\ \vdots \\ a_k \\ 0 \\ \vdots \\ 0 \end{bmatrix},$$

$$a_k \neq 0$$

$$m_i = \frac{a_i}{a_k} \forall k < i \leq n$$

a_k is know as the common **pivot**

Elementary Elimination Matrix

$$\underbrace{\begin{pmatrix} 1 & 0 & 0 \\ -\frac{a_2}{a_1} & 1 & 0 \\ -\frac{a_3}{a_1} & 0 & 1 \end{pmatrix}}_{\text{Elimination matrix}} \underbrace{\begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}}_{\text{Column vector}} = \underbrace{\begin{pmatrix} a_1 \\ 0 \\ 0 \end{pmatrix}}_{\text{column vector with eliminated terms}}$$

$$A\vec{x} = \vec{b}$$

GOAL

$$U\vec{x} = \vec{z}$$

$$M_1 A \vec{x} = M_1 \vec{b}$$

Build M_i using
elements in first column of A

Eliminate terms
below **first** diagonal element

$$M_2 M_1 A \vec{x} = M_2 M_1 \vec{b}$$

Eliminate terms
below **second** diagonal element

\vdots

$$M_{n-1} \dots M_2 M_1 A \vec{x} = M_{n-1} \dots M_2 M_1 \vec{b}$$

$$U \vec{x} = \vec{z}$$



backward
substitution

Properties of Elementary Elimination Matrices

M_k is **non-singular** (the diagonal is made of 1 and it is lower triangular).

M_k^{-1} is the same as M_k except the **signs of the multipliers are reversed**.

$M_k M_j$ is “the union” of M_k and M_j , provided $j > k$.

This is amazing!

$$\overbrace{\begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix}}^A \overbrace{\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix}}^x = \overbrace{\begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix}}^b$$

$$\overbrace{\begin{pmatrix} u_{11} & u_{12} & u_{13} & u_{14} \\ 0 & u_{22} & u_{23} & u_{24} \\ 0 & 0 & u_{33} & u_{34} \\ 0 & 0 & 0 & u_{44} \end{pmatrix}}^U \overbrace{\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix}}^x = \overbrace{\begin{pmatrix} z_1 \\ z_2 \\ z_3 \\ z_4 \end{pmatrix}}^z$$

$$u_{44}x_4 = z_4 \Rightarrow x_4 = \frac{z_4}{u_{44}}$$

$$u_{33}x_3 + u_{34}x_4 = z_3 \Rightarrow x_3 = \frac{z_3 - u_{34}x_4}{u_{33}}$$

$$u_{22}x_2 + u_{23}x_3 + u_{24}x_4 = z_2 \Rightarrow x_2 = \frac{z_2 - u_{23}x_3 - u_{24}x_4}{u_{22}}$$

$$u_{11}x_1 + u_{12}x_2 + u_{13}x_3 + u_{14}x_4 = z_1 \Rightarrow x_1 = \frac{z_1 - u_{12}x_2 - u_{13}x_3 - u_{14}x_4}{u_{11}}$$

For a **lower triangular** matrix we have a similar **forward substitution** procedure.

Algorithm 1 Back substitution for upper triangular system

BACK-SUBSTITUTION(U, z)

- 1 **for** j in $\{n, \dots, 1\}$: ▷ Loop backwards over columns
 - 2 **do**
 - 3 **if** $u_{jj} = 0$ ▷ Stop if matrix is singular
 then Stop
 - 4 $x_j = \frac{z_j}{u_{jj}}$ ▷ Compute solution component
 - 5 **for** i in $\{1, \dots, j-1\}$:
 - 6 **do**
 - 7 $z_i = z_i - u_{ij}x_j$ ▷ Update right-hand side
-

Let's go **back to the task**
(transforming the problem into an easy one)

Example

$$\begin{aligned}x_1 + 2x_2 + 2x_3 &= 3 \\4x_1 + 4x_2 + 2x_3 &= 6 \\4x_1 + 6x_2 + 4x_3 &= 10\end{aligned}$$

$$Ax = b$$

$$\begin{pmatrix} 1 & 2 & 2 \\ 4 & 4 & 2 \\ 4 & 6 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 \\ 6 \\ 10 \end{pmatrix}$$

$$M_1 A = \begin{pmatrix} 1 & 0 & 0 \\ -4 & 1 & 0 \\ -4 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 & 2 \\ 4 & 4 & 2 \\ 4 & 6 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 2 \\ 0 & -4 & -6 \\ 0 & -2 & -4 \end{pmatrix}$$

$$M_1 b = \begin{pmatrix} 1 & 0 & 0 \\ -4 & 1 & 0 \\ -4 & 0 & 1 \end{pmatrix} \begin{pmatrix} 3 \\ 6 \\ 10 \end{pmatrix} = \begin{pmatrix} 3 \\ -6 \\ -2 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 2 & 2 \\ 0 & -4 & -6 \\ 0 & -2 & -4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 \\ -6 \\ -2 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 2 & 2 \\ 0 & -4 & -6 \\ 0 & -2 & -4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 \\ -6 \\ -2 \end{pmatrix}$$

Gaussian elimination
= Forward elimination + Back substitution

$$M_2 M_1 A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -0.5 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 & 2 \\ 0 & -4 & -6 \\ 0 & -2 & -4 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 2 \\ 0 & -4 & -6 \\ 0 & 0 & -1 \end{pmatrix}$$

$$M_2 M_1 b = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -0.5 & 1 \end{pmatrix} \begin{pmatrix} 3 \\ -6 \\ -2 \end{pmatrix} = \begin{pmatrix} 3 \\ -6 \\ 1 \end{pmatrix}$$

$$\left(\begin{array}{c} \text{orange triangle} \end{array} \right) \leftarrow \begin{pmatrix} 1 & 2 & 2 \\ 0 & -4 & -6 \\ 0 & 0 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 \\ -6 \\ 1 \end{pmatrix}$$

backward
substitution

$$U \vec{x} = \vec{z}$$

LU factorisation

$$Ax = b$$

$$\left(\begin{array}{c|c} \text{blue triangle} & \end{array} \right) M_{n-1} \dots M_1 Ax = M_{n-1} \dots M_1 b \left(\begin{array}{c|c} & \text{green triangle} \end{array} \right)$$

Upper triangular **Lower triangular**



$$(M_{n-1} \dots M_1)^{-1} (M_{n-1} \dots M_1) Ax = b$$

$$(M_1^{-1} M_2^{-1} \dots M_{n-1}^{-1}) (M_{n-1} \dots M_1) Ax = b$$

$$(L_1 L_2 \dots L_{n-1}) (M_{n-1} \dots M_1) Ax = b$$

Lower triangular **Upper triangular**

$$L U x = b$$

A turns into **L** and **U**

$$Ax = b$$

$$\textcolor{green}{L} \textcolor{blue}{U} x = b$$

$$\begin{array}{ccc} \textcolor{green}{L} & y = & b \\ \downarrow & & \downarrow \\ \text{known} & & \text{known} \\ & \downarrow & \\ & \underline{\text{unknown}} & \\ & \text{known!} & \end{array}$$

$$Ux = y$$

$n \times 1$ unknown

$$\left(\begin{array}{c|c} \textcolor{green}{\triangle} & \end{array} \right) \quad \begin{array}{l} \text{forward} \\ \text{substitution} \end{array}$$

$$\left(\begin{array}{c|c} & \textcolor{blue}{\triangle} \end{array} \right) \quad \begin{array}{l} \text{backward} \\ \text{substitution} \end{array}$$

DONE!


Summary...

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix}$$

$$\begin{pmatrix} u_{11} & u_{12} & u_{13} & u_{14} \\ 0 & u_{22} & u_{23} & u_{24} \\ 0 & 0 & u_{33} & u_{34} \\ 0 & 0 & 0 & u_{44} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{pmatrix} = M_3 M_2 M_1 \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix}$$

$$M_1^{-1} M_2^{-1} M_3^{-1} \begin{pmatrix} u_{11} & u_{12} & u_{13} & u_{14} \\ 0 & u_{22} & u_{23} & u_{24} \\ 0 & 0 & u_{33} & u_{34} \\ 0 & 0 & 0 & u_{44} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix}$$

Forward
elimination



Summary...

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix}$$

$$\begin{pmatrix} u_{11} & u_{12} & u_{13} & u_{14} \\ 0 & u_{22} & u_{23} & u_{24} \\ 0 & 0 & u_{33} & u_{34} \\ 0 & 0 & 0 & u_{44} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{pmatrix} = M_3 M_2 M_1 \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix}$$

Forward
elimination



$$\begin{pmatrix} l_{11} & 0 & 0 & 0 \\ l_{21} & l_{22} & 0 & 0 \\ l_{31} & l_{23} & l_{33} & 0 \\ l_{41} & l_{42} & l_{43} & l_{44} \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} & u_{13} & u_{14} \\ 0 & u_{22} & u_{23} & u_{24} \\ 0 & 0 & u_{33} & u_{34} \\ 0 & 0 & 0 & u_{44} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix}$$

Summary...

$$\begin{pmatrix} l_{11} & 0 & 0 & 0 \\ l_{21} & l_{22} & 0 & 0 \\ l_{31} & l_{23} & l_{33} & 0 \\ l_{41} & l_{42} & l_{43} & l_{44} \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} & u_{13} & u_{14} \\ 0 & u_{22} & u_{23} & u_{24} \\ 0 & 0 & u_{33} & u_{34} \\ 0 & 0 & 0 & u_{44} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix}$$

$$\begin{pmatrix} l_{11} & 0 & 0 & 0 \\ l_{21} & l_{22} & 0 & 0 \\ l_{31} & l_{23} & l_{33} & 0 \\ l_{41} & l_{42} & l_{43} & l_{44} \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix}$$

Forward substitution
to find **y**

$$\begin{pmatrix} u_{11} & u_{12} & u_{13} & u_{14} \\ 0 & u_{22} & u_{23} & u_{24} \\ 0 & 0 & u_{33} & u_{34} \\ 0 & 0 & 0 & u_{44} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{pmatrix}$$

Backward substitution
to find **x**

Which is better?

HOMEWORK

Potential issues

- Division by zero
- Round-off error

$$\begin{pmatrix} 10 & -7 & 0 \\ -3 & 2.099 & 6 \\ 5 & -1 & 5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 7 \\ 3.901 \\ 6 \end{pmatrix}$$

6 significant digits

with chopping

$$\begin{pmatrix} 10 & -7 & 0 \\ 0 & -0.001 & 6 \\ 0 & 0 & 15005 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 7 \\ 6.001 \\ 15005 \end{pmatrix}$$

$$x_3 = 1$$

$$x_2 = -1$$

$$x_1 = 0$$

5 significant digits


with chopping

$$\begin{pmatrix} 10 & -7 & 0 \\ 0 & -0.001 & 6 \\ 0 & 0 & 15004 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 7 \\ 6.001 \\ 15005 \end{pmatrix}$$

$$x_3 = 0.99993$$

$$x_2 = -1.5$$

$$x_1 = -0.35$$



In a column, if you **use the largest entry on or below the diagonal as the pivot**... then multipliers are bounded by 1

Use a permutation matrix to swap rows...

Partial pivoting

$$Ax = \begin{pmatrix} 1 & 2 & 2 \\ 4 & 4 & 2 \\ 4 & 6 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 \\ 6 \\ 10 \end{pmatrix} = b$$

$$P_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$P_1Ax = \begin{pmatrix} 4 & 4 & 2 \\ 1 & 2 & 2 \\ 4 & 6 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 6 \\ 3 \\ 10 \end{pmatrix} = P_1b$$

$$M_1 = \begin{pmatrix} 1 & 0 & 0 \\ -0.25 & 1 & 0 \\ -1 & 0 & 1 \end{pmatrix}$$

$$M_1P_1Ax = \begin{pmatrix} 4 & 4 & 2 \\ 0 & 1 & 1.5 \\ 0 & 2 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 6 \\ 1.5 \\ 4 \end{pmatrix} = M_1P_1b$$

$$M_1 P_1 A x = \begin{pmatrix} 4 & 4 & 2 \\ 0 & 1 & 1.5 \\ 0 & 2 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 6 \\ 1.5 \\ 4 \end{pmatrix} = M_1 P_1 b$$

$$P_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

$$P_2 M_1 P_1 A x = \begin{pmatrix} 4 & 4 & 2 \\ 0 & 2 & 2 \\ 0 & 1 & 1.5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 6 \\ 4 \\ 1.5 \end{pmatrix} = P_2 M_1 P_1 b$$

$$M_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -0.5 & 1 \end{pmatrix}$$

$$M_2 P_2 M_1 P_1 A x = \begin{pmatrix} 4 & 4 & 2 \\ 0 & 2 & 2 \\ 0 & 0 & 0.5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 6 \\ 4 \\ -0.5 \end{pmatrix} = M_2 P_2 M_1 P_1 b$$

$$M_2 P_2 M_1 P_1 A x = \begin{pmatrix} 4 & 4 & 2 \\ 0 & 2 & 2 \\ 0 & 0 & 0.5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 6 \\ 4 \\ -0.5 \end{pmatrix} = M_2 P_2 M_1 P_1 b$$

For explicit **LU** decomposition


$$L = M^{-1} = (M_2 P_2 M_1 P_1)^{-1} = P_1^T L_1 P_2^T L_2 = \begin{pmatrix} 0.25 & 0.5 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 0 \end{pmatrix}$$

$$A = \begin{pmatrix} 1 & 2 & 2 \\ 4 & 4 & 2 \\ 4 & 6 & 4 \end{pmatrix} = \begin{pmatrix} 0.25 & 0.5 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 0 \end{pmatrix} \begin{pmatrix} 4 & 4 & 2 \\ 0 & 2 & 2 \\ 0 & 0 & 0.5 \end{pmatrix} = LU$$

But **L is not lower triangular!**



$$P = P_2 P_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$$



$$L = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0.25 & 0.5 & 1 \end{pmatrix}$$

$$PA = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 2 & 2 \\ 4 & 4 & 2 \\ 4 & 6 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0.25 & 0.5 & 1 \end{pmatrix} \begin{pmatrix} 4 & 4 & 2 \\ 0 & 2 & 2 \\ 0 & 0 & 0.5 \end{pmatrix} = LU$$

$$PA = LU$$

Partial pivoting... extra (thin) layer of complexity, with permutations....

scipy.linalg.lu

scipy.linalg.lu(*a*, *permute_l=False*, *overwrite_a=False*, *check_finite=True*)

[\[source\]](#)

Compute pivoted LU decomposition of a matrix.

The decomposition is:

$$A = P L U$$

where P is a permutation matrix, L lower triangular with unit diagonal elements, and U upper triangular.

Parameters: *a* : (M, N) array_like

Array to decompose

permute_l : bool

Perform the multiplication P*L (Default: do not permute)

overwrite_a : bool

Whether to overwrite data in a (may improve performance)

check_finite : boolean, optional

Whether to check that the input matrix contains only finite numbers. Disabling may give a performance gain, but may result in problems (crashes, non-termination) if the inputs do contain infinities or NaNs.

Returns:

(If **permute_l == False**)

p : (M, M) ndarray

Permutation matrix

l : (M, K) ndarray

Lower triangular or trapezoidal matrix with unit diagonal. K = min(M, N)

u : (K, N) ndarray

Upper triangular or trapezoidal matrix

(If **permute_l == True**)

pl : (M, K) ndarray

Permuted L matrix. K = min(M, N)

u : (K, N) ndarray

Upper triangular or trapezoidal matrix

Previous topic

[scipy.linalg.eigvals_banded](#)

Next topic

[scipy.linalg.lu_factor](#)