

# Geometry: Maths Olympiad

G V V Sharma\*

**Abstract**—In this letter, an algorithm for evaluating the exact analytical bit error rate (BER) for the piecewise linear (PL) combiner for multiple relays is presented. Previous results were available only for upto three relays. The algorithm is unique in the sense that the actual mathematical expressions, that are prohibitively large, need not be explicitly obtained. The diversity gain due to multiple relays is shown through plots of the analytical BER, well supported by simulations.

1. Consider triangle  $P_1P_2P_3$  and a point P within the triangle. Lines  $P_1P$ ,  $P_2P$ ,  $P_3P$  intersect the opposite sides in points  $Q_1$ ,  $Q_2$ ,  $Q_3$  respectively. Prove that, of the numbers

$$\frac{P_1P}{PQ_1}, \frac{P_2P}{PQ_2}, \frac{P_3P}{PQ_3}$$

at least one is  $\leq 2$  and at least one is  $\geq 2$ .

2. Construct triangle ABC if  $AC = b$ ,  $AB = c$  and  $\angle AMB = \omega$ , where M is the midpoint of segment BC and  $\omega < 90^\circ$ . Prove that a solution exists if and only if

$$b \tan \frac{\omega}{2} \leq c < b.$$

In what case does the equality hold?

3. Consider a plane  $\varepsilon$  and three non-collinear points A, B, C on the same side of  $\varepsilon$ ; suppose the plane determined by these three points is not parallel to  $\varepsilon$ . In plane  $\varepsilon$  take three arbitrary points  $A_0, B_0, C_0$ . Let L, M, N be the midpoints of segments  $AA_0, BB_0, CC_0$ ; let G be the centroid of triangle LMN. (We will not consider positions of the points  $A', B', C'$  such that the points L, M, N do not form a triangle.) What is the locus of point G as  $A', B', C'$  range independently over the plane  $\varepsilon$ ?

4. Given a triangle ABC, let I be the center of its inscribed circle. The internal bisectors of the angles A, B, C meet the opposite sides in  $A', B', C'$  respectively. Prove that

$$\frac{1}{4} < \frac{AI \cdot BI \cdot CI}{AA' \cdot BB' \cdot CC'} \leq \frac{8}{27}.$$

5. Let ABC be a triangle and P an interior point of ABC. Show that at least one of the angles  $\angle PAB$ ,  $\angle PBC$ ,  $\angle PCA$  is less than or equal to  $30^\circ$ .
6. Consider nine points in space, no four of which are coplanar. Each pair of points is joined by an edge (that is, a line segment) and each edge is either colored blue or red or left uncolored. Find the smallest value of n such that whenever exactly n edges are colored, the set of colored edges necessarily contains a triangle all of whose edges have the same color.
7. In the plane let C be a circle, L a line tangent to the circle C, and M a point on L. Find the locus of all points P with the following property: there exists two points Q, R on L such that M is the midpoint of QR and C is the inscribed circle of triangle PQR.
8. Let D be a point inside acute triangle ABC such that  $\angle ADB = \angle ACB + \frac{\pi}{2}$  and  $AC \cdot BD = AD \cdot BC$ .
  - a) Calculate the ratio  $(AB \cdot CD)/(AC \cdot BD)$ .
  - b) Prove that the tangents at C to the circumcircles of  $\triangle ACD$  and  $\triangle BCD$  are perpendicular.
9. ABC is an isosceles triangle with  $AB = AC$ . Suppose that
  - a) M is the midpoint of BC and O is the point on the line AM such that OB is perpendicular to AB;
  - b) Q is an arbitrary point on the segment BC different from B and C;
  - c) E lies on the line AB and F lies on the line AC such that E, Q, F are distinct and collinear.
10. Let A, B, C, D be four distinct points on a line, in that order. The circles with diameters

\*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in. All content in this manual is released under GNU GPL. Free and open source.

AC and BD intersect at X and Y . The line XY meets BC at Z. Let P be a point on the line XY other than Z. The line CP intersects the circle with diameter AC at C and M, and the line BP intersects the circle with diameter BD at B and N. Prove that the lines AM, DN, XY are concurrent.

11. Let ABCDEF be a convex hexagon with  $AB = BC = CD$  and  $DE = EF = FA$ , such that  $\angle BCD = \angle EFA = \frac{\pi}{3}$ . Suppose G and H are points in the interior of the hexagon such that  $\angle AGB = \angle DHE = \frac{2\pi}{3}$ . Prove that  $AG + GB + GH + DH + HE \geq CF$ .
12. We are given a positive integer r and a rectangular board ABCD with dimensions  $|AB| = 20$ ,  $|BC| = 12$ . The rectangle is divided into a grid of  $20 \times 12$  unit squares. The following moves are permitted on the board: one can move from one square to another only if the distance between the centers of the two squares is  $\sqrt{r}$ . The task is to find a sequence of moves leading from the square with A as a vertex to the square with B as a vertex.
  - a) Show that the task cannot be done if r is divisible by 2 or 3.
  - b) Prove that the task is possible when  $r = 73$ .
  - c) Can the task be done when  $r = 97$ ?
13. Let P be a point inside triangle ABC such that

$$\angle APB - \angle ACB = \angle APC - \angle ABC.$$

Let D, E be the incenters of triangles APB, APC, respectively. Show that AP, BD, CE meet at a point.

14. Let ABCDEF be a convex hexagon such that AB is parallel to DE, BC is parallel to EF, and CD is parallel to FA. Let  $R_A$ ,  $R_C$ ,  $R_E$  denote the circumradii of triangles FAB, BCD, DEF, respectively, and let P denote the perimeter of the hexagon. Prove that
 
$$R_A + R_C + R_E \geq \frac{P}{2}.$$
15. In the plane the points with integer coordinates are the vertices of unit squares. The squares are colored alternately black and white (as on a chessboard). For any pair of positive integers m and n, consider a right-angled triangle

whose vertices have integer coordinates and whose legs, of lengths m and n, lie along edges of the squares.

Let  $S_1$  be the total area of the black part of the triangle and  $S_2$  be the total area of the white part. Let

$$f(m, n) = |S_1 - S_2|.$$

- a) Calculate  $f(m, n)$  for all positive integers m and n which are either both even or both odd.
  - b) Prove that  $f(m, n) \leq \frac{1}{2} \max m, n$  for all m and n.
  - c) Show that there is no constant C such that  $f(m, n) < C$  for all m and n.
16. The angle at A is the smallest angle of triangle ABC. The points B and C divide the circumcircle of the triangle into two arcs. Let U be an interior point of the arc between B and C which does not contain A. The perpendicular bisectors of AB and AC meet the line AU at V and W, respectively. The lines BV and CW meet at T. Show that
 
$$AU = TB + TC.$$
  17. In the convex quadrilateral ABCD, the diagonals AC and BD are perpendicular and the opposite sides AB and DC are not parallel. Suppose that the point P, where the perpendicular bisectors of AB and DC meet, is inside ABCD. Prove that ABCD is a cyclic quadrilateral if and only if the triangles ABP and CDP have equal areas.
  18. Let I be the incenter of triangle ABC. Let the incircle of ABC touch the sides BC, CA, and AB at K, L, and M, respectively. The line through B parallel to MK meets the lines LM and LK at R and S, respectively. Prove that angle RIS is acute.
  19. Two circles  $G_1$  and  $G_2$  are contained inside the circle G, and are tangent to G at the distinct points M and N, respectively.  $G_1$  passes through the center of  $G_2$ . The line passing through the two points of intersection of  $G_1$  and  $G_2$  meets G at A and B. The lines MA and MB meet  $G_1$  at C and D, respectively.
  20. AB is tangent to the circles CAMN and NMBD. M lies between C and D on the line

CD, and CD is parallel to AB. The chords NA and CM meet at P; the chords NB and MD meet at Q. The rays CA and DB meet at E. Prove that  $PE = QE$ .

side BC. The bisectors of the angles  $\angle BAC$  and  $\angle MON$  intersect at R. Prove that the circumcircles of the triangles BMR and CNR have a common point lying on the side BC.

21.  $A_1A_2A_3$  is an acute-angled triangle. The foot of the altitude from  $A_i$  is  $K_i$  and the incircle touches the side opposite  $A_i$  at  $L_i$ . The line  $K_1K_2$  is reflected in the line  $L_1L_2$ . Similarly, the line  $K_2K_3$  is reflected in  $L_2L_3$  and  $K_3K_1$  is reflected in  $L_3L_1$ . Show that the three new lines form a triangle with vertices on the incircle.

22. Let ABC be an acute-angled triangle with circumcentre O. Let P on BC be the foot of the altitude from A.

Suppose that  $\angle BCA \geq \angle ABC + 30^\circ$ .

Prove that  $\angle CAB + \angle COP < 90^\circ$ .

23. In a triangle ABC, let AP bisect  $\angle BAC$ , with P on BC, and let BQ bisect  $\angle ABC$ , with Q on CA. It is known that  $\angle BAC = 60^\circ$  and that  $AB + BP = AQ + QB$ . What are the possible angles of triangle ABC?

24. BC is a diameter of a circle center O. A is any point on the circle with  $\angle AOC > 60^\circ$ . EF is the chord which is the perpendicular bisector of AO. D is the midpoint of the minor arc AB. The line through O parallel to AD meets AC at J. Show that J is the incenter of triangle CEF.

25.  $n > 2$  circles of radius 1 are drawn in the plane so that no line meets more than two of the circles. Their centers are  $O_1, O_2, \dots, O_n$ . Show that  $\sum_{i < j} \frac{1}{O_i O_j} \leq (n-1)\frac{\pi}{4}$ .

26. A convex hexagon has the property that for any pair of opposite sides the distance between their midpoints is  $\sqrt{3}/2$  times the sum of their lengths. Show that all the hexagon's angles are equal.

27. ABCD is cyclic. The feet of the perpendicular from D to the lines AB, BC, CA are P, Q, R respectively. Show that the angle bisectors of ABC and CDA meet on the line AC if  $RP = RQ$ .

28. Let ABC be an acute-angled triangle with  $AB \neq AC$ . The circle with diameter BC intersects the sides AB and AC at M and N respectively. Denote by O the midpoint of the

29. In a convex quadrilateral ABCD the diagonal BD does not bisect the angles ABC and CDA. The point P lies inside ABCD and satisfies

$$\angle PBC = \angle DBA \text{ and } \angle PDC = \angle BDA.$$

Prove that ABCD is a cyclic quadrilateral if and only if  $AP = CP$ .