# Assignment 6 Papoullis Textbook Chapter 5 Example 33

Rishi Manoj - CS21BTECH11045

May 31, 2022



#### Outline

- Question
- Solution
- Probability P
- 4 Probability  $p_n(k)$
- Solving
- Moment Generating Function
- **7** Solving for  $p_n(0)$
- Probability Pr

### Question

A person writes n letters and addresses n envelopes. Then one letter is randomly placed into each envelope. What is the probability that at least one letter will reach its correct destination?



#### Solution

Let  $X_k$  represents the event that there are exactly k coincidences among the n envelopes. All the events corresponding to the values of k starting from 0 to n are mutually exclusive and exhaustive. So, by theorem of total probability,

$$p_n(0) + p_n(1) + p_n(2) + \cdots + p_n(n) = 1$$
 (1)

$$p_n(k) \triangleq P(X_k) \tag{2}$$

Let P be the probability of generating k coincidences with the chosen k letters and Pr be the required probability that atleast one letter reaches its correct destination.



## Probability P

Number of ways of drawing k letters from a group of n is  $\binom{n}{k}$ . Probability of generating k coincidences with the chosen k letters is,

$$P = \left(\frac{1}{n}\right) \left(\frac{1}{n-1}\right) \cdots \left(\frac{1}{n-k+1}\right) \tag{3}$$



# Probability $p_n(k)$

Probability of no coincidences with remaining n-k letters is given by  $p_{n-k}(0)$ . So,

$$p_n(k) = \binom{n}{k} \left( \frac{1}{n(n-1)\cdots(n-k+1)} \right) p_{n-k}(0) \tag{4}$$

$$=\frac{p_{n-k}(0)}{k}\tag{5}$$

# Solving

We know that.

$$p_n(n) = \frac{1}{n!} \tag{6}$$

Substituiting eq.(5) in eq.(1) along with the above one gives,

$$p_n(0) + \frac{p_{n-1}(0)}{1!} + \dots + \frac{p_1(0)}{(n-1)!} + \frac{1}{n!} = 1$$
 (7)

From this we can get the following,

$$p_1(0) = 0 (8)$$

$$p_2(0) = \frac{1}{2}$$
 (9)  
$$p_3(0) = \frac{1}{6}$$
 (10)

$$p_3(0) = \frac{1}{6} \tag{10}$$



## Moment Generating Function

We define the moment generating function in the following way to get an explicit expression for  $p_n(0)$ ,

$$\phi(z) = \sum_{n=0}^{\infty} \rho_n(0) z^n \tag{11}$$

$$e^{z}\phi(z) = \left(\sum_{k=0}^{\infty} \frac{z^{k}}{k!}\right) \left(\sum_{n=0}^{\infty} p_{n}(0)z^{n}\right)$$
 (12)

$$=1+z+z^2+\cdots+z^n+\cdots \tag{13}$$

$$=\frac{1}{1-z}\tag{14}$$

# Solving for $p_n(0)$

Using the above equations, we get,

$$\phi(z) = \frac{e^{-z}}{1-z} \tag{15}$$

$$= \sum_{n=0}^{\infty} \left( \sum_{k=0}^{n} \frac{(-1)^k}{k!} \right) z^n \tag{16}$$

$$p_n(0) = \sum_{k=0}^n \frac{(-1)^k}{k!} \to \frac{1}{e}$$
 (17)

$$= 0.378$$
 (18)

Using eq.(5) we get,

$$p_n(k) = \frac{1}{k!} \sum_{m=0}^{n-k} \frac{(-1)^m}{m!}$$
 (19)



## Probability Pr

The required probability is given by,

$$Pr = 1 - p_n(0) \tag{20}$$

$$=1-\sum_{k=0}^{n}\frac{(-1)^{k}}{k!}\to 0.632 \tag{21}$$

