


Scanning Physical Interaction Behavior of 3D Objects

Dinesh K. Pai, Kees van den Doel, Doug L. James, Jochen Lang, John E. Lloyd, Joshua L. Richmond, Som H. Yau


Presented by
Rishit Dagli




some content taken
from the paper

what did we do up until now?

- start with physical laws
- develop some numerical methods to solve these equations
- make sure you have good accuracy and stability in your simulation
- **predict behavior** from the first laws




[BZCGZ] 2023]




[S 1999]

let us introduce a new problem

- start with physical laws
- develop some numerical methods to solve these equations
- make sure you have good accuracy and stability in your simulation
- predict behavior from the first laws




[BZCGZJ 2023]



[S 1999]


- observe the behavior of a real object
- reproduce the observed phenomena
- make sure you can well capture or measure the phenomena




let us introduce a new problem

- start with physical laws

- no longer need models of complex phenomena
- can easily capture certain behaviors which are hard to simulate
- often very efficient for real-time (producing behaviors in real-time)




[BZCGZJ 2023]



[S 1999]

- observe the behavior of a real object
- reproduce the observed phenomena
- make sure you can well capture or measure the phenomena



what will we talk about today?

To appear in SIGGRAPH 2001 Conference Proceedings

Scanning Physical Interaction Behavior of 3D Objects

Dinesh K. Pai, Kees van den Doel, Doug L. James, Jochen Lang, John E. Lloyd, Joshua L. Richmond, Som H. Yau

Department of Computer Science, University of British Columbia, Vancouver, Canada
{pai,kvdoel,djames,jlang,lloyd,jlrichmo,shyau}@cs.ubc.ca



(a) Real toy tiger. By design, it is soft to touch and exhibits significant deformation behavior.



(b) Deformable model of tiger scanned by our system, with haptic interaction.



(c) Real clay pot, with glazed regions. The pot exhibits a variety of contact textures and sounds.



(d) Virtual interaction with scanned model of pot; includes contact texture and contact sounds.

Figure 1: Examples of behavior models scanned by our system

from 2001 

what will we talk about today?

To appear in SIGGRAPH 2001 Conference Proceedings

Scanning **Physical Interaction Behavior** of 3D Objects

Dinesh K. Pai, Kees van den Doel, Doug L. James, Jochen Lang, John E. Lloyd, Joshua L. Richmond, Som H. Yau

Department of Computer Science, University of British Columbia, Vancouver, Canada
{pai,kvdoel,djames,jlang,lloyd,jlrichmo,shyan}@cs.ubc.ca

- deformation
- contact texture
- sound

deformation behavior. tion. tact textures and sounds. ture and contact sounds.

Figure 1: Examples of behavior models scanned by our system

from 2001 

what will we talk about today?

focus on
measurement

To appear in SIGGRAPH 2001 Conference Proceedings

Scanning Physical Interaction Behavior of 3D Objects

Dinesh K. Pai, Kees van den Doel, Doug L. James, Jochen Lang, John E. Lloyd, Joshua L. Richmond, Som H. Yau

Department of Computer Science, University of British Columbia, Vancouver, Canada
{pai,kvdoel,djames,jlang,llloyd,jlrichmo,shyau}@cs.ubc.ca

- deformation
- contact texture
- sound


deformation behavior. tion. tact textures and sounds. ture and contact sounds.

Figure 1: Examples of behavior models scanned by our system


from 2001 

motivation

- there were some deformation models back then that kind of worked
- there did exist haptic simulations
- and acoustic modelling techniques or more accurately spatialization techniques



[FCEPSW 1999]




[RKK 1997]

motivation


- there were some deformation models back then that kind of worked
- there did exist haptic simulations
- and acoustic modelling techniques or more accurately spatialization techniques



often require completely different frameworks



[FCEPSW 1999]




[RKK 1997]


motivation

- there were some deformation models back then that kind of worked
- there did exist haptic simulations
- and acoustic modelling techniques or more accurately spatialization techniques

very far from simulating
all the relevant information



[FCEPSW 1999]




[RKK 1997]


motivation

- there were some deformation models back then that kind of worked
- there did exist haptic simulations
- and acoustic modelling techniques or more accurately spatialization techniques

very far from simulating
all the relevant information



[FCEPSW 1999]




[RKK 1997]

very hard to make this
work to mimic a real object

motivation

make models which have such parameters such that they can all be efficiently (automatically) captured just through a real object



[RKK 1997]



very hard to make this work to mimic a real object




what to measure and how to measure?

measurement system

[PLLW 1999]




field
measurement system




force/position
measurement system

measurement system




measurement system

3 DOF placement




measurement system

field measurement system
(stereo vision + rgb camera + mic)
on a 5 DOF positioning robot




light field



sound field



[MID 2002]



[PLLW 1999]

measurement system



[MID 2002]


stereo range data is highly noisy
we need some kind of filtering to
decide what stereo range data we
need

some classical techniques:

- variable mask size voting
- estimate (local) surface normal
- decide some handcrafted ways of choosing what to keep



measurement system

force/position measurement system
(force/torque sensor)
on a linear stage



measurement system

time-of-flight laser range finder




measurement system

field measurement system
(stereo vision + rgb camera + mic)
on a 5 DOF positioning robot

3 DOF placement


force/position measurement system
(force/torque sensor)
on a linear stage

time-of-flight laser range finder




mesh reconstruction


multiresolution mesh




range data




subdivision with displaced subdivision + tricks





multiresolution mesh



range data



subdivision with displaced subdivision + tricks



[RW 1997]

multiresolution textured mesh



multiview color images

To appear in SIGGRAPH 2001 Conference Proceedings

Scanning **Physical Interaction Behavior** of 3D Objects

Dinesh K. Pai, Kees van den Doel, Doug L. James, Jochen Lang, John E. Lloyd, Joshua L. Richmond, Som H. Yau

Department of Computer Science, University of British Columbia, Vancouver, Canada

let us focus on this now



(a) Real toy tiger. By design, it is soft to touch and exhibits significant deformation behavior.



(b) Deformable model of tiger scanned by our system, with haptic interaction.




(c) Real clay pot, with glazed regions. The pot exhibits a variety of contact textures and sounds.




(d) Virtual interaction with scanned model of pot; includes contact texture and contact sounds.

Figure 1: Examples of behavior models scanned by our system



deformation

what model do we use?



boundaries

linear elasticity operator

body forces

$$G \sum_{k=1}^3 \left(\frac{\partial^2 u_i}{\partial x_k^2} + \frac{1}{1-2\nu} \frac{\partial^2 u_k}{\partial x_k \partial x_i} \right) + b_i = 0,$$

$$(\mathbf{N} \mathbf{u})(\mathbf{x}) + \mathbf{b}(\mathbf{x}) = 0, \quad \mathbf{x} \in \Omega.$$

$$\underbrace{\mathbf{G}\nabla^2\mathbf{u}}_{\text{Pure shear}} + \underbrace{\frac{G}{1-2\nu}\nabla(\nabla\cdot\mathbf{u})}_{\text{Volume change}} + \underbrace{\mathbf{b}}_{\text{Body force}} = \mathbf{0}$$

$$\begin{aligned}\mathbf{u} &= \bar{\mathbf{u}} && \text{on } \Gamma_u \\ \mathbf{p} &= \bar{\mathbf{p}} && \text{on } \Gamma_p\end{aligned}$$

$$\overbrace{G\nabla^2 \mathbf{u}}^{\text{Pure shear}} + \overbrace{\frac{G}{1-2\nu} \nabla(\nabla \cdot \mathbf{u})}^{\text{Volume change}} + \overbrace{\mathbf{b}}^{\text{Body force}} = \mathbf{0}$$

but body force is
a point force

$$G\nabla^2 \mathbf{u}^* + \frac{G}{1-2\nu} \nabla(\nabla \cdot \mathbf{u}^*) = -\delta(\mathbf{x} - \mathbf{y})\mathbf{I}$$

aside: make greens
functions

Pure shear $\underbrace{G\nabla^2 \mathbf{u}}$ + $\underbrace{\frac{G}{1-2\nu} \nabla(\nabla \cdot \mathbf{u})}_{\text{Volume change}}$ + $\underbrace{\mathbf{b}}_{\text{Body force}} = \mathbf{0}$

aside: make greens functions

but body force is a point force

$$G\nabla^2 \mathbf{u}^* + \frac{G}{1-2\nu} \nabla(\nabla \cdot \mathbf{u}^*) = -\delta(\mathbf{x} - \mathbf{y}) \mathbf{I}$$

exploit linearity

$$G\nabla^2 u_{ij}^* + \frac{G}{1-2\nu} \frac{\partial}{\partial x_i} \left(\underbrace{\sum_{k=1}^3 \frac{\partial u_{kj}^*}{\partial x_k}}_{\text{divergence}} \right) = - \overbrace{\delta(\mathbf{x} - \mathbf{y}) \delta_{ij}}^{\text{point force in direction j}}$$

aside: make greens functions


$$\underbrace{G\nabla^2 \mathbf{u}}_{\text{Pure shear}} + \underbrace{\frac{G}{1-2\nu} \nabla(\nabla \cdot \mathbf{u})}_{\text{Volume change}} + \underbrace{\mathbf{b}}_{\text{Body force}} = \mathbf{0}$$

but body force is
a point force

$$G\nabla^2 \mathbf{u}^* + \frac{G}{1-2\nu} \nabla(\nabla \cdot \mathbf{u}^*) = -\delta(\mathbf{x} - \mathbf{y}) \mathbf{I}$$

exploit linearity

$$G\nabla^2 u_{ij}^* + \frac{G}{1-2\nu} \frac{\partial}{\partial x_i} \left(\underbrace{\sum_{k=1}^3 \frac{\partial u_{kj}^*}{\partial x_k}}_{\text{divergence}} \right) = - \underbrace{\delta(\mathbf{x} - \mathbf{y}) \delta_{ij}}_{\text{point force in direction j}}$$



fourier transform

$$G(-|\xi|^2) \hat{u}_{ij}^* + \frac{G}{1-2\nu} (-\xi_i \xi_j \hat{u}_{ij}^*) = -e^{-i\xi \cdot \mathbf{y}} \delta_{ij}$$

$$G(-|\xi|^2)\hat{u}_{ij}^* + \frac{G}{1-2\nu}(-\xi_i\xi_j\hat{u}_{ij}^*) = -e^{-i\xi\cdot y}\delta_{ij}$$

aside: make greens
functions

$$G(-|\xi|^2)\hat{u}_{ij}^* + \frac{G}{1-2\nu}(-\xi_i\xi_j\hat{u}_{ij}^*) = -e^{-i\xi\cdot y}\delta_{ij}$$

aside: make greens
functions


solve this in fourier
space

$$u_{ij}^*(x, y) = \frac{1}{16\pi G(1-\nu)} \left[\overbrace{\frac{(3-4\nu)\delta_{ij}}{r}}^{\text{isotropic expansion}} + \underbrace{\frac{r_i r_j}{r^3}}_{\text{directional deformation}} \right]$$

$$\underbrace{G\nabla^2 \mathbf{u}}_{\text{Pure shear}} + \underbrace{\frac{G}{1-2\nu} \nabla(\nabla \cdot \mathbf{u})}_{\text{Volume change}} + \underbrace{\mathbf{b}}_{\text{Body force}} = \mathbf{0}$$

$$\mathbf{u} = \bar{\mathbf{u}} \quad \text{on } \partial\Omega$$

$$\mathbf{p} = \bar{\mathbf{p}} \quad \text{on } \partial\Omega$$



Elastic operator

$$\underbrace{\mathcal{N} \mathbf{u}^*(x, y)}_{\text{green function}} + \underbrace{\delta(x - y) \mathbf{I}}_{\text{Point force}} = \mathbf{0}$$

green function

$$u_{ij}^*(x, y) = \frac{1}{16\pi G(1-\nu)}$$

$$\left[\underbrace{\frac{(3-4\nu)\delta_{ij}}{r}}_{\text{Isotropic expansion}} + \underbrace{\frac{r_i r_j}{r^3}}_{\text{Directional deformation}} \right]$$

$$p_{ij}^*(x, y) = -\frac{1}{8\pi(1-\nu)r^2} \left[\underbrace{\frac{\partial r}{\partial n} \left((1-2\nu)\delta_{ij} + 3\frac{r_i r_j}{r^2} \right)}_{\text{Normal traction}} + \underbrace{(1-2\nu)(n_i r_j - n_j r_i)}_{\text{Shear traction}} \right]$$

weighted
residual


$$\overbrace{\int_{\Omega} \mathbf{u}^* \cdot \mathcal{N} \mathbf{u} d\Omega}^{\text{Volume work}} = \underbrace{\oint_{\Gamma} \mathbf{u}^* \cdot \mathbf{p} d\Gamma}_{\text{Boundary work}} - \overbrace{\int_{\Omega} \nabla \mathbf{u}^* : \boldsymbol{\sigma}(\mathbf{u}) d\Omega}^{\text{Internal strain energy}}$$

reciprocal theorem
and by parts

Difference of virtual works in volume

$$\overbrace{\int_{\Omega} (\mathbf{u}^* \cdot \mathcal{N} \mathbf{u} - \mathbf{u} \cdot \mathcal{N} \mathbf{u}^*) d\Omega}^{\text{Difference of virtual works in volume}} = \underbrace{\oint_{\Gamma} (\mathbf{u}^* \cdot \mathbf{p} - \mathbf{u} \cdot \mathbf{p}^*)}_{\text{Difference of virtual works on boundary}} d\Gamma$$

Difference of virtual works on boundary



$$\mathcal{Q} = \underbrace{\oint_{\Gamma} \mathbf{u}^* \cdot \mathbf{p} d\Gamma}_{\text{Boundary work}} - \underbrace{\int_{\Omega} \nabla \mathbf{u}^* : \boldsymbol{\sigma}(\mathbf{u}) d\Omega}_{\text{Internal strain energy}}$$

reciprocal theorem
and by parts

Difference of virtual works in volume

$$\underbrace{\int_{\Omega} (\mathbf{u}^* \cdot \mathcal{N} \mathbf{u} - \mathbf{u} \cdot \mathcal{N} \mathbf{u}^*) d\Omega}_{\text{Difference of virtual works in volume}} = \underbrace{\oint_{\Gamma} (\mathbf{u}^* \cdot \mathbf{p} - \mathbf{u} \cdot \mathbf{p}^*) d\Gamma}_{\text{Difference of virtual works on boundary}}$$

discretize the boundary into non-overlapping pieces
do field approximations for \mathbf{u} and \mathbf{p}

$$\underbrace{\overbrace{c_{ij}(x^m) \mathbf{u}_j(x^m)}^{\text{Point displacement}} + \sum_{e=1}^N \underbrace{\int_{\Gamma_e} \mathbf{p}_{ij}^*(x^m, y) \phi_k(y) d\Gamma(y) \mathbf{u}_j^k}_{\text{Influence of all displacements}}}_{\text{Influence of all displacements}} = \underbrace{\sum_{e=1}^N \int_{\Gamma_e} \mathbf{u}_{ij}^*(x^m, y) \phi_k(y) d\Gamma(y) \mathbf{p}_j^k}_{\text{Influence of all tractions}}$$


now let us bring it all together

$$\begin{aligned}
 \mathbf{H}_{ij} &= \underbrace{c_{ij}\delta_{ij}}_{\text{Jump term}} + \underbrace{\sum_{e=1}^N \int_{\Gamma_e} \mathbf{p}_{ij}^*(x, y) \phi_k(y) d\Gamma(y)}_{\text{Double layer influence coefficients}} \\
 \mathbf{G}_{ij} &= \underbrace{\sum_{e=1}^N \int_{\Gamma_e} \mathbf{u}_{ij}^*(x, y) \phi_k(y) d\Gamma(y)}_{\text{Single layer influence coefficients}}
 \end{aligned}$$

$$\underbrace{\mathbf{H}}_{\text{System matrix}} = \underbrace{\mathbf{G}}_{\text{Influence matrix}} \underbrace{\mathbf{u}}_{\text{Displacements}}$$

$\underbrace{\mathbf{p}}_{\text{Tractions}}$

now let us bring it all together




now let us bring it all together

$$\begin{aligned}
 \mathbf{H}_{ij} &= \underbrace{c_{ij}\delta_{ij}}_{\text{Jump term}} + \underbrace{\sum_{e=1}^N \int_{\Gamma_e} \mathbf{p}_{ij}^*(x, y) \phi_k(y) d\Gamma(y)}_{\text{Double layer influence coefficients}} \\
 \mathbf{G}_{ij} &= \underbrace{\sum_{e=1}^N \int_{\Gamma_e} \mathbf{u}_{ij}^*(x, y) \phi_k(y) d\Gamma(y)}_{\text{Single layer influence coefficients}}
 \end{aligned}$$

$$\underbrace{\mathbf{H}}_{\text{System matrix}} \underbrace{\mathbf{u}}_{\text{Displacements}} = \underbrace{\mathbf{G}}_{\text{Influence matrix}} \underbrace{\mathbf{p}}_{\text{Tractions}}$$

does this look familiar?


getting back to capturing



- apply a controlled force
- records 6D force/torque data at 1000Hz
- trinocular vision system captures the surface deformation at 30Hz
- repeat

use vision system:

- track visual features from optical flow
- reconstructing 3D motion field
- map the deformation onto the mesh vertices



get model parameters

- we already have displacement field and force for vertices of mesh

$$[\mathbf{p}_j^1 \mathbf{p}_j^2 \cdots \mathbf{p}_j^M]^T \mathbf{U}_{ij}^T = [\mathbf{u}_i^1 \mathbf{u}_i^2 \cdots \mathbf{u}_i^M]^T$$

does this look familiar?
tsvd + least trimmed squares

get model parameters

- we already have displacement field and force for vertices of mesh

$$[\mathbf{p}_j^1 \mathbf{p}_j^2 \cdots \mathbf{p}_j^M]^T \mathbf{U}_{ij}^T = [\mathbf{u}_i^1 \mathbf{u}_i^2 \cdots \mathbf{u}_i^M]^T$$

does this look familiar?

tsvd + least trimmed squares
also interpolate missing values
also interpolate across resolutions

System matrix

\mathbf{H}

\mathbf{u}

Displacements

Influence matrix


\mathbf{G}

\mathbf{p}

Tractions


=

make this system



contact texture


in general: model, measurement, rendering



$$\mathbf{f}_f = -\mu \|\mathbf{f}_n\| \mathbf{u}_m$$

$$\mu_e(x) = \mu + \tilde{\mu}(x) \quad \text{non-typical}$$

model



friction

surface roughness

$$\mathbf{f}_f = -\mu \|\mathbf{f}_n\| \mathbf{u}_m$$

$$\mu_e(x) = \mu + \tilde{\mu}(x) \quad \text{non-typical}$$

model all these variations as an auto-regressive process

$$\tilde{\mu}(x) = \tilde{\mu}(k\Delta) \equiv \tilde{\mu}(k) = \sum_{i=1}^p a_i \tilde{\mu}(k-i) + \sigma \epsilon(k)$$

index

how many times
do you do this

ar coefficients

std dev of noise

zero mean and
std. dev 1 noise

model

friction

$$\mathbf{f}_f = -\mu \|\mathbf{f}_n\| \mathbf{u}_m$$

surface roughness


$$\mu_e(x) = \mu + \tilde{\mu}(x) \quad \text{non-typical}$$

model all these variations as an auto-regressive process

$$\begin{bmatrix} \mu(x) \\ \mu(x - \Delta) \end{bmatrix} = \begin{bmatrix} a_1 & a_2 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \mu(x - \Delta) \\ \mu(x - 2\Delta) \end{bmatrix} + \begin{bmatrix} \sigma \\ 0 \end{bmatrix} \epsilon(x)$$

index how many times do you do this p a_i σ zero mean and std. dev 1 noise

measurement



use multi resolution mesh

$$\mu = \tan(\theta/2)$$

measurement

$$\mathbf{n}(x) = \frac{\mathbf{f}_f^+ + \mathbf{f}_f^-}{\|\mathbf{f}_f^+ + \mathbf{f}_f^-\|}$$

measurement

$$\mathbf{n}(x) = \frac{\mathbf{f}_f^+ + \mathbf{f}_f^-}{\|\mathbf{f}_f^+ + \mathbf{f}_f^-\|}$$

$$\mathbf{f} = \mathbf{f}_n [\mathbf{n}(x) - \mu_e \mathbf{u}_m(x)]$$




sensor

from our
estimate



direction of motion
from a sensor

rendering




barycentric interpolation

ar process
discretize by Δ



find effective friction



sound

model

$$M = \{ \overbrace{\mathbf{f}}^{\text{frequencies}}, \overbrace{\mathbf{d}}^{\text{decay rates}}, \underbrace{\mathbf{A}}_{\text{mode gains}} \}$$

$$y_k(t) = \sum_{n=1}^N \overbrace{a_{nk}}^{\text{mode amplitude}} \underbrace{e^{-d_n t}}_{\text{decay}} \overbrace{\sin(2\pi f_n t)}^{\text{oscillation}}$$

model

$$M = \{ \overbrace{\mathbf{f}}^{\text{frequencies}}, \overbrace{\mathbf{d}}^{\text{decay rates}}, \underbrace{\mathbf{A}}_{\text{mode gains}} \}$$


$$y_k(t) = \sum_{n=1}^N \overbrace{a_{nk}}^{\text{mode amplitude}} \underbrace{e^{-d_n t}}_{\text{decay}} \overbrace{\sin(2\pi f_n t)}^{\text{oscillation}}$$

$$F_{audio}(t) = \overbrace{F_N}^{\text{normal force}} \cdot \underbrace{\mu_e(v \cdot t)}_{\text{friction variation}} \cdot \overbrace{v(t)}^{\text{sliding velocity}}$$

measurement


measure impulse response

$$h(t) = \sum_{n=1}^N \underbrace{A_n e^{-d_n t}}_{\text{amplitude envelope}} \cdot \underbrace{\cos(2\pi f_n t + \phi_n)}_{\text{oscillation}}$$




rendering

$$\overbrace{\ddot{x}_n + 2d_n\dot{x}_n + \omega_n^2 x_n}^{\text{resonator dynamics}} = \underbrace{F_{audio}(t)}_{\text{input force}}$$



rendering

$$\overbrace{\ddot{x}_n + 2d_n\dot{x}_n + \omega_n^2 x_n}^{\text{resonator dynamics}} = \underbrace{F_{audio}(t)}_{\text{input force}}$$



$$\begin{bmatrix} x_n(t + \Delta t) \\ \dot{x}_n(t + \Delta t) \end{bmatrix} = \overbrace{e^{A\Delta t}}^{\text{state transition}} \begin{bmatrix} x_n(t) \\ \dot{x}_n(t) \end{bmatrix} + \underbrace{\int_t^{t+\Delta t} e^{A(t+\Delta t-\tau)} b F_{audio}(\tau) d\tau}_{\text{force response}}$$

$A = \begin{bmatrix} 0 & 1 \\ -\omega_n^2 & -2d_n \end{bmatrix}, \quad b = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$

summary





Synthesized or Real?

2

references

- [PDJLLRY 2001] D. K. Pai, K. van den Doel, D. L. James, J. Lang, J. E. Lloyd, J. L. Richmond, S. H. Yau, "Scanning Physical Interaction Behavior of 3D Objects," in *Computer Graphics (ACM SIGGRAPH 2001 Conference Proceedings)*, August 2001.
- [LPS 2003] Lang, J., Pai, D. K., & Seidel, H-P. Scanning Large-Scale Articulated Deformations. In *Proceedings of Graphics Interface*, Halifax, June 2003.
- [LP 2001] Lloyd, J. E., & Pai, D. K. Robotic Mapping of Friction and Roughness for Reality-based Modeling. In *Proceedings of the 2001 IEEE International Conference on Robotics and Automation*, 1884-1890.
- [RP 2000] Richmond, J. L., & Pai, D. K. Active Measurement and Modeling of Contact Sounds. In *Proceedings of the 2000 IEEE International Conference on Robotics and Automation*, San Francisco, April 2000, 2146-2152.
- [LP 1999] Lang, J., & Pai, D. K. Bayesian Estimation of Distance and Surface Normal with a Time-of-Flight Laser Rangefinder. In *Proceedings of the Second International Conference on 3-D Digital Imaging and Modeling (IEEE)*, Ottawa, October 1999, 109-117.
- [PLLW 1999] Pai, D. K., Lang, J., Lloyd, J. E., & Woodham, R. J. ACME, A Telerobotic Active Measurement Facility. In *Proceedings of the Sixth International Symposium on Experimental Robotics*, Sydney, Australia, March 1999. Also in *Experimental Robotics VI*. Springer London, 2000, 391-400.
- [P 1999] Pai, D. K. Robotics in Reality-based Modeling. In *Proceedings of the International Symposium on Robotics Research*, Snowbird, UT, October 8-12, 1999. Also in *Robotics Research: the Ninth International Symposium*, Springer-Verlag, 2000, 353-358.
- [S 1999] Stam, Jos. "Stable Fluids." (1999).
- [BZCGZJ 2023] Benchekroun, Otman, Jiayi Eris Zhang, Siddhartha Chaudhuri, Eitan Grinspun, Yi Zhou, and Alec Jacobson. "Fast Complementary Dynamics via Skinning Eigenmodes." *ACM Transactions on Graphics (TOG)* 42, no. 4 (2023): 1-21.
- [RKK 1997] Ruspini, Diego C., Krasimir Kolarov, and Oussama Khatib. "The haptic display of complex graphical environments." In *Proceedings of the 24th annual conference on Computer graphics and interactive techniques*, pp. 345-352. 1997.
- [FCEPSW 1999] Funkhouser, Thomas, Ingrid Carlbom, Gary Elko, Gopal Pingali, Mohan Sondhi, and Jim West. "Interactive acoustic modeling of complex environments." *J. Acoust. Soc. Am* 105, no. 2 (1999): 1357.
- [MID 2002] Mulligan, Jane, Volkan Isler, and Kostas Daniilidis. "Trinocular stereo: A real-time algorithm and its evaluation." *International Journal of Computer Vision* 47 (2002): 51-61.
- [RW 1997] Roth, Gerhard, and Eko Wibowoo. "An Efficient Volumetric Method for Building Closed Triangular Meshes from 3-D Image and Point Data." In the *proceedings of Graphics Interface*, pp. 173-180. 1997.
- [LMH 2000] Lee, Aaron, Henry Moreton, and Hugues Hoppe. "Displaced subdivision surfaces." In *Proceedings of the 27th annual conference on Computer graphics and interactive techniques*, pp. 85-94. 2000.
- [MMC 2016] Macklin, Miles, Matthias Müller, and Nuttapong Chentanez. "XPBD: position-based simulation of compliant constrained dynamics." In *Proceedings of the 9th International Conference on Motion in Games*, pp. 49-54. 2016.
- [JP 1999] James, Doug L., and Dinesh K. Pai. "Artdefo: accurate real time deformable objects." In *Proceedings of the 26th annual conference on Computer graphics and interactive techniques*, pp. 65-72. 1999.
- [DPWK 2024] Dagli, Rishit, Shivesh Prakash, Robert Wu, and Houman Khosravani. "SEE-2-SOUND: Zero-Shot Spatial Environment-to-Spatial Sound." *ICML Workshop on Foundation Models in the Wild* (2024).
- [DKP 2001] Van Den Doel, Kees, Paul G. Kry, and Dinesh K. Pai. "FoleyAutomatic: physically-based sound effects for interactive simulation and animation." In *Proceedings of the 28th annual conference on Computer graphics and interactive techniques*, pp. 537-544. 2001.
- [CWLWCZZL 2024] Chen, Yiwen, Yikai Wang, Yihao Luo, Zhengyi Wang, Zilong Chen, Jun Zhu, Chi Zhang, and Guosheng Lin. "Meshanything v2: Artist-created mesh generation with adjacent mesh tokenization." *arXiv preprint arXiv:2408.02555* (2024).

Scanning Physical Interaction Behavior of 3D Objects

