

Healthcare delivery gaps in pain management within the first 3 months after discharge from inpatient noncardiac surgeries: a scoping review

Janny X. C. Ke^{1,2,3,*†}, Maya de Vos^{2†}, Katarina Kojic^{1,2,†}, Mark Hwang^{4,†}, Jason Park^{5,6}, Heather Stuart^{5,6}, Jill Osborn^{1,2}, Alana Flexman^{1,2}, Lindsay Blake⁷ and Daniel I. McIsaac⁸

¹Department of Anesthesia, Providence Health Care, Vancouver, BC, Canada, ²Department of Anesthesiology, Pharmacology and Therapeutics, Faculty of Medicine, University of British Columbia, Vancouver, BC, Canada, ³Department of Anesthesia, Pain Management & Perioperative Medicine, Dalhousie University, Halifax, NS, Canada, ⁴Undergraduate Medical Education Program, Faculty of Medicine, University of British Columbia, Vancouver, BC, Canada, ⁵Department of Surgery, Faculty of Medicine, University of British Columbia, Vancouver, BC, Canada, ⁶Department of Surgery, Vancouver General Hospital, Vancouver, BC, Canada, ⁷University of Arkansas for Medical Sciences Library, Little Rock, AK, USA and ⁸Department of Anesthesiology and Pain Medicine, University of Ottawa, Ottawa, ON, Canada

*Corresponding author. E-mail: janny.ke@ubc.ca, [@jannyke](https://twitter.com/jannyke)

†Co-first authors.

‡Co-second authors.

Abstract

Background: Poor pain control during the postoperative period has negative implications for recovery, and is a critical risk factor for development of persistent postsurgical pain. The aim of this scoping review is to identify gaps in healthcare delivery that patients undergoing inpatient noncardiac surgeries experience in pain management while recovering at home.

Methods: Searches were conducted by a medical librarian in PubMed, MEDLINE, EMBASE, EBSCO CINAHL, Web of Science, and Cochrane Database of Systematic Reviews for articles published between 2016 and 2022. Inclusion criteria were adults (≥ 18 yr), English language, inpatient noncardiac surgery, and included at least one gap in care for acute and/or persistent pain management after surgery within the first 3 months of recovery at home. Two reviewers independently screened articles for inclusion and extracted data. Quotations from each article related to gaps in care were synthesised using thematic analysis.

Results: There were 4794 results from databases and grey literature, of which 38 articles met inclusion criteria. From these, 23 gaps were extracted, encompassing all six domains of healthcare delivery (capacity, organisational structure, finances, patients, care processes and infrastructure, and culture). Identified gaps were synthesised into five overarching themes: education (22 studies), provision of continuity of care (21 studies), individualised management (10 studies), support for specific populations (11 studies), and research and knowledge translation (10 studies).

Conclusions: This scoping review identified health delivery gaps during a critical period in postoperative pain management. These gaps represent potential targets for quality improvement and future research to improve perioperative care and longer-term patient-centred outcomes.

Scoping review protocol: Open Science Framework (<https://osf.io/cq5m6/>).

Keywords: gaps in care; health delivery; pain management; persistent postsurgical pain; post-discharge; postoperative pain; transitional pain service

Editor's key points

- Poorly controlled postsurgical pain has significant implications, including persistent postsurgical pain.
- In this scoping review, the authors examined delivery gaps in pain management during the 3 months after discharge following inpatient noncardiac surgery. The gaps included common themes of education, continuity of care, individualisation, support for patient populations, and research and knowledge translation.
- Anaesthetists can work with stakeholders to help better understand and manage pain during this critical period of acute pain control and chronic pain development.

Globally, more than 312 million surgeries were performed in 2012, a rate that is expected to increase.¹ After surgery, more than 80% of patients experience pain, with 80% of these patients rating their pain as moderate, severe, or extreme.² Poorly controlled pain after surgery can have significant implications on recovery and quality of life, including morbidity, disability, readmission, and prolonged opioid use.^{3–8} Many patients with poorly controlled acute surgical pain also develop persistent postsurgical pain (also known as chronic postsurgical pain), which is defined as ‘pain that develops or increases in intensity after a surgical procedure and persists beyond the healing process’, typically greater than 3 months.^{3,9,10} Studies show that moderate to severe persistent postsurgical pain is common after inpatient surgeries, including hip replacement (7–23%), hernia repair (5–35%), Cesarean section (10–21%), hysterectomy (5–32%), mastectomy (20–50%), thoracotomy (61–70%), limb amputation (30–81%), and cholecystectomy (3–56%).^{3,11} In fact, a year after surgery, 12% of patients still report moderate to severe persistent postsurgical pain, with more severe pain correlating with greater functional impairment.⁸ Persistent postsurgical pain also has societal implications as a result of lost work and increased healthcare resource utilisation.^{2,12,13} For example, patients with persistent postsurgical pain after lumbar spine surgery experienced £5000, £10,000, and £14,000 higher healthcare costs at 2, 5, and 10 yr postoperatively, respectively, than patients without persistent pain.¹⁴

Experiencing more severe acute pain is associated with a greater risk of persistent postsurgical pain; for every 10% increase in duration of severe pain on postoperative day 1, the incidence of persistent postsurgical pain increases by 30% at 1 yr.⁸ Although anaesthetists, surgeons, and allied health professionals are involved in the management of acute and chronic pain, a substantial gap exists in research and clinical care for pain management during the transition from acute hospitalisation to home and before the diagnosis of chronic pain can be made.¹⁵ Currently, we lack evidence-based pharmacological therapies to prevent persistent postsurgical pain.¹⁶ Furthermore, after discharge, patients are often expected to manage their own pain with little professional support.¹⁵ Transitional Pain Services are recent innovations seeking to address gaps in care between discharge and the development of chronic pain, and are associated with improved patient knowledge, function, and pain scores while reducing long-term opioid prescribing.¹⁵

However, Transitional Pain Services are not widely accessible outside of tertiary centres. These issues are exacerbated by variability in postoperative analgesic prescribing¹⁷ as well as the push towards shorter hospital stays,¹⁸ where patients spend a greater proportion of their recovery without pain management support.

Although most pain management research and care focus on the individual patient level, a better understanding of pain management needs from a health services perspective may reveal opportunities to mitigate the transition from acute pain to persistent postsurgical pain. According to the Agency for Healthcare Research and Quality's (AHRQ) Effective Health Care Stakeholders Group Delivery Systems Committee, the delivery of healthcare can be conceptualised using a framework of six domains: capacity, organisational structure, finances, patients, care processes and infrastructure, and culture.¹⁹ By identifying gaps, which are discontinuities in care,²⁰ we can unveil areas of vulnerability for research and quality improvement. The objective of this scoping review is to identify gaps in healthcare delivery that patients undergoing inpatient noncardiac surgeries experience in pain management within the first 3 months of their recovery at home.

Methods

A scoping review was conducted according to Joanna Briggs Institute methodology for scoping reviews²¹ and reported using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for scoping reviews (PRISMA-ScR).²² The protocol was registered on May 26, 2022, in the Open Science Framework (<https://osf.io/cq5m6/>).

Search strategy

The search strategy was codeveloped with an experienced medical librarian (LB) and included articles published in PubMed MEDLINE, EMBASE, EBSCO CINAHL, Web of Science, and Cochrane Database of Systematic Reviews (Supplementary material 1). The database search was performed on June 2, 2022 and updated to December 31, 2022. The grey literature search was done according to published methods²³ involving a search of pre-print and grey literature databases, a Google search, a search within websites of Canadian foundations for surgical patients, and a review of included article references for further articles (Supplementary material 1). The grey literature search was performed between June 1 and 4, 2022.

Definition of gaps

We defined gaps in care based on the existing patient safety literature, which defines gaps as discontinuities in care that ‘may appear as losses of information or momentum or interruptions in delivery of care’ and may occur ‘with organisational and institutional boundaries that mark changes in responsibility or authority, different roles of professionals, or formal divisions of labour’ or as a side-effect of systems change.²⁰ We approached the definition as inclusively as possible, and included any problems, issues, challenges, and/or barriers in healthcare delivery. A gap was operationalised in the search strategy using both broad Medical Subject Headings (“Continuity of Patient Care,” “Root Cause Analysis,” “Professional Practice Gaps,” or “Preventive Health Services”) as well

as keyword search terms (gap, barrier, problem, issue, or challenge) ([Supplementary material 1](#)).

Inclusion and exclusion criteria

We included studies involving adults (≥ 18 yr), English language, inpatient noncardiac surgery, and published within the past 6 yr to obtain contemporary perspectives (2016–22). Articles must have included at least one gap in care for acute and/or persistent pain management after surgery within the first 3 months of recovery at home. We chose 3 months post-operatively since postsurgical pain beyond 3 months in duration would be considered persistent, which opens a new dimension of logistical and disease considerations for health delivery that would be beyond the scope of this study. Also, focusing on the period prior to 3 months allows for in-depth exploration of gaps that contribute to the transition from acute to persistent pain. To obtain a broad perspective and acknowledging that experimental research relevant to the research question would be limited, all types of published full-text articles (including experimental and observational study designs, reviews, commentaries, editorials, and study protocols) were included. Cardiac surgeries and surgeries without overnight hospital stay were excluded as these have unique considerations in terms of health system structures and processes.

Article selection

After removal of duplicates, articles were uploaded into Covidence (Veritas Health Information, Melbourne, VIC, Australia). Titles and abstracts were screened by two independent reviewers based on the study inclusion and exclusion criteria. Articles selected for potential inclusion were retrieved in full-text and assessed by two independent reviewers against the inclusion/exclusion criteria to select studies for data extraction. Disagreements between reviewers for both the title/abstract and full-text screening were resolved through discussion. If not resolved, additional reviewers were involved in the discussion until a consensus was achieved.

Data extraction

Two independent reviewers extracted data from each article using a standardised Microsoft Excel (Microsoft Corporation, Redmond, WA, USA) form developed in accordance with the Joanna Briggs Institute methodology for scoping reviews.²¹ The data extraction form ([Supplementary material 2](#)) included details about the participants (population characteristics), methodology of the study, and all text that discusses gaps in postsurgical pain management within the first 3 months after discharge from inpatient surgery. If conflict arose between the reviewers, they were resolved through discussion and if necessary, additional reviewers were involved to help reach a consensus.

Scoping review data analysis and presentation

Text from each article related to gaps in care underwent thematic analysis using a six-step process of familiarisation, initial codes, search for themes, review themes, define and name themes, and report.²⁴ During inductive analysis, the text data were reviewed to generate a preliminary coding structure. Similar codes were grouped together by consensus

amongst the study team to create a summarised list of gaps. Then, common themes amongst similar gaps were generated as potential targets for further research and intervention. We subsequently applied a deductive framework of six domains as defined by the AHRQ. These were capacity, 'the physical assets and their ownership, personnel, and organizational characteristics of a delivery system that determine the number of individuals and breadth of conditions for which the system can provide care'; organizational structure, 'the components of an organisation, both formal and informal, that describe functional operations in terms of hierarchy of authority and the flow of information, patients, and resources'; finances, 'mechanisms by which a healthcare delivery system is paid for its services and the financial arrangements and practices of the system and organizations within the system to allocate those funds, as well as the system's financial status'; patients, 'demographic characteristics, as well as wants, needs, and preferences of individuals and families of individuals who receive healthcare services from a healthcare delivery system'; care processes and infrastructure, 'the methods by which a healthcare delivery system provides healthcare services to its customers and patients as well as the degree of coordination of those methods'; and culture, 'shared values, beliefs, and assumptions that influence behaviour, attitudes, and meaning in an organization'.¹⁹

Results


Study characteristics

The search, screening, inclusion, and exclusion process is reported in [Figure 1](#). After removal of duplicates, 3740 results were screened for potential inclusion, with 36 included studies (20 from databases, 16 from grey literature search). A search of the references of these included articles revealed an additional two articles that met criteria, for a total of 38 studies. The results from the grey literature search are detailed in [Supplementary material 3](#).

The characteristics of the included studies are detailed in [Table 1](#). Sixteen countries were represented, though representation from low-income countries was limited: USA (17 studies), Canada (five studies), Brazil (two studies), Germany (two studies), and single reports from Belgium, China, Columbia, Denmark, Finland, France, Japan, Netherlands, Norway, Pakistan, Spain, Sweden, and UK. There were three systematic reviews, 13 narrative reviews, eight cohort studies (five prospective and three retrospective), five interview and/or focus group studies for patients, two conference summaries, and one each of cross-sectional survey, interview study for physicians, book chapter, editorial, expert interview, quality standards, and study protocol. Seventeen studies focused on specific subtypes of surgery, predominantly hip and/or knee arthroplasty ($n=8$), total joint arthroplasty ($n=2$), with single studies in colorectal cancer, general and plastic, general only, major upper abdominal, major oncological, orthopaedic trauma, spine, and thoracic surgeries. Further patient and surgical characteristics are detailed in [Supplementary material 4](#).

Gaps and themes

[Supplementary material 5](#) summarises the quotations regarding gaps in health delivery identified in the care of postoperative pain management for adult patients undergoing



noncardiac surgery. From these, 23 gaps were extracted (Table 2), which encompassed all six AHRQ domains of healthcare delivery: care processes and infrastructure (19 gaps), patients (seven gaps), organisational structure (three gaps), capacity (two gaps), finances (two gaps), and culture (four gaps). Five overarching themes emerged (Table 2): education (22 studies), provision of continuity of care (21 studies), individualised management (10 studies), support for specific populations (11 studies), and research and knowledge translation (10 studies).

Education

The most frequently mentioned gaps were related to education. Included studies reported that there was insufficient patient education on self-management of opioid and non-opioid pain medications, including purpose, adherence, perception, administration, weaning, disposal, and side-effects in an evidence-based, understandable format.^{15,27,30,32,33,35,38,40–44,46,49,50,53–55} Lack of patient comprehension was prevalent, with one study³² reporting that 70% (114 out of 147) 'could not accurately reproduce their pain medication regimen, regardless of age, employment status, income level, education level, and time between surgery and follow-up'. In a cross-sectional survey of 155 patients, Nasir and Ahmed⁴⁶ found that 'Excellent and good knowledge [about postoperative pain and its management] were observed in 11.61% and 21.94% [of] patients, respectively, whereas fair and poor knowledge were seen in 42.58% and 23.87%, respectively. Inadequate knowledge was more marked regarding analgesic side-effects and addiction risk'. Similarly, Lemay and colleagues⁴² found that '44% of 1609

patients reported that they did not receive or received unhelpful information regarding postoperative pain self-management options for surgical knee or hip pain'.

Patients also appeared to have limited understanding about expected levels of postoperative pain and risk of persistent postsurgical pain.^{15,27,38–40,50,53,55} Patients and providers reported negative perception of pain medications, including fear of opioids.^{43,44,46,49,50,57,58} Communication gaps existed amongst providers and patients, including the reporting of pain and what to do when there are pain issues.^{27,33,43,44,55,57} The issues in patient education were compounded by provider knowledge gaps including facilitating behaviour change, sharing tools for pain management, care coordination and treatment for patients with or at high risk of opioid dependence, appropriate perioperative opioid therapy and tapering, multimodal management options, and cultural competency.^{30,33,35,39,43,45,57,58}

Provision of continuity of care

According to Lopes and colleagues,⁴³ 'it is acknowledged that there is a period of subacute pain during which its management has been neglected' and 'this pain can last for several weeks, have a negative psychological impact, contribute to the process of central sensitisation, and is associated with hospital readmission and delayed postoperative recovery'. Through interviews of eight patients within the first 12 weeks after hip and knee surgery, Specht and colleagues⁵⁴ reported that patients had difficulty managing their pain at home. Patients expected 'more interaction with staff once they had gone home, to ask questions about their pain and pain medication

Table 1 Characteristics of included studies. N/A, not applicable.

Author and year	Country	Specific surgical group	Study design	Study aim	Sample size
Andrés and colleagues, 2017 ²⁵	Netherlands	N/A	Conference session summary and case report	To summarise the quality of acute pain management systems in Europe.	N/A
Beloeil and Sulpice 2016 ²⁶	France	N/A	Narrative review	To discuss the risk factors, management, and consequences of perioperative pain.	N/A
Bleicher and colleagues, 2021 ²⁷	USA	General surgery	Patient interviews	To understand patients' perspectives on postoperative pain management education.	30
Cata and colleagues, 2019 ²⁸	USA	Major oncological procedures	Narrative review	To summarise challenges in acute postoperative pain management in patients with cancer.	N/A
European Federation of National Association of Orthopaedics and Traumatology 2019 ²⁹	Belgium	Hip and knee arthroplasty	Expert interview	To improve postoperative pain management and improve postoperative outcomes relating to orthopaedic surgical practice.	1
Esses and colleagues, 2020 ³⁰	USA	N/A	Narrative review	To discuss prevention and management of postoperative pain in patients aged >65 yr living with frailty.	N/A
Finnegan and colleagues, 2017 ³¹	USA	Hip and knee arthroplasty	Retrospective cohort	To characterise emergency department visits in the acute postoperative period after total knee and hip arthroplasty.	152 782
Gan 2017 ²	USA	N/A	Narrative review	To provide an update on the issues surrounding poorly controlled postoperative pain.	N/A
Gangavalli and colleagues, 2017 ³²	USA	Orthopaedic trauma surgery	Prospective cohort	To identify patients who are at risk of non-compliance and misunderstanding in regards to their postoperative pain management.	147
Garcia and colleagues, 2017 ³³	Brazil	N/A	Narrative review	To discuss issues in acute postoperative pain management and identify potential solutions.	N/A
Hadlandsmayth 2021 ³⁴	USA	Total joint replacement	Study protocol	An RCT to study the efficacy of a perioperative pain self-management program to prevent the development of chronic postsurgical pain and prolonged opioid use.	N/A
Health Quality Ontario 2018 ³⁵	Canada	N/A	Quality standards	To summarise recommendations for improving opioid prescribing for acute pain, including postsurgical pain.	N/A
Jackman and colleagues, 2019 ³⁶	USA	N/A	Narrative review	To discuss challenges in postoperative pain management in patients with chronic pain and opioid dependence.	N/A
Joseph and colleagues, 2022 ³⁷	USA	N/A	Systematic review	To evaluate the quality and availability of postoperative pain management measures.	19 pain quality measures; 206 publications
Kennedy and colleagues, 2017 ³⁸	Canada	Hip and knee arthroplasty	Patient focus groups and interviews	To explore patients' experiences and areas for improvement following hip and knee arthroplasty.	32 (6 focus groups and 7 phone semi-structured interviews)
Klueh and colleagues, 2019 ³⁹	USA	N/A	Physician interviews	To identify opportunities for improvement in transitions of care in postoperative opioid prescribing.	10 surgeons and 10 primary care physicians

Continued

Table 1 Continued

Author and year	Country	Specific surgical group	Study design	Study aim	Sample size
Kojic and Clarke 2021 ¹⁵	Canada	N/A	Editorial	To discuss key considerations in the transition from acute to persistent postsurgical pain.	N/A
Kuusniemi and Pöyhiäk 2016 ⁴⁰	Finland	N/A	Conference summary	To discuss challenges and propose solutions to improve postoperative pain management.	N/A
Lemay and colleagues, 2019 ⁴¹	USA	Total joint replacement	Patient interviews	To summarise postoperative pain management education needs after elective joint replacement.	27
Lemay and colleagues, 2017 ⁴²	USA	Hip and knee arthroplasty	Prospective cohort	To evaluate the patient perspective of postoperative pain management education after elective joint replacement.	1609
Lopes and colleagues, 2021 ⁴³	Brazil	N/A	Narrative review	To review postoperative pain prevention, diagnosis, and treatment.	N/A
Makimoto and colleagues, 2020 ⁴⁴	Japan	Hip and knee arthroplasty	Systematic review	To explore patients' experiences in the first 6 weeks after total hip or knee arthroplasty.	253 (16 qualitative studies)
Meissner and Zaslansky 2019 ⁴⁵	Germany	N/A	Narrative review	To review postoperative pain management, and identify areas for improvement.	N/A
Nasir and Ahmed 2020 ⁴⁶	Pakistan	Major upper abdominal surgery	Cross-sectional survey	To evaluate patients' knowledge of postoperative pain and its management.	155
Nguyen and colleagues, 2022 ⁴⁷	USA	Total knee arthroplasty	Retrospective cohort	To investigate the relationship between limited English proficiency and requests for postoperative opioid prescription refills up to 90 days after hospital discharge for total knee arthroplasty.	2148
Pogatzki-Zahn and Lavandhomme 2021 ⁴⁸	Germany, Belgium	N/A	Narrative review	To review the prevention of chronic postsurgical pain.	N/A
Rahman and colleagues, 2021 ⁴⁹	USA	Spine surgery	Prospective cohort	To assess patients' knowledge and barriers to appropriate use of opioids postoperatively.	174
Reaza-Alarcón and Rodríguez-Martín 2019 ⁵⁰	Spain, Ireland	N/A	Systematic review	To evaluate the benefits of nurse education in managing postoperative pain.	12 studies
Simpson and Jackson 2017 ⁵¹	UK	N/A	Narrative review	To review methods for managing postoperative pain in opioid-tolerant patients.	N/A
Sjøveian and Leegaard 2017 ⁵²	Norway	Hip and knee arthroplasty	Patient interviews	To assess patients' experiences with postoperative pain management after hip and knee arthroplasty.	12
Smith and colleagues, 2018 ⁵³	Sweden	Colorectal cancer surgery	Prospective cohort	To explore stressful events and management strategies in the 4–6 weeks after colorectal cancer surgery.	105
Specht and colleagues, 2018 ⁵⁴	Denmark	Hip and knee arthroplasty	Patient interviews	To explore patients' experiences in the first 12 weeks after hip and knee arthroplasty.	8
Sugar and colleagues, 2018 ⁵⁵	Canada	N/A	Retrospective cohort	To describe emergency department visits within 6 weeks of surgery and explore issues of continuity of care.	248
Tawfic and colleagues, 2017 ⁵⁶	Canada	N/A	Narrative review	To discuss prevention of chronic postsurgical pain, focusing on early identification of at-risk patients.	N/A
Thakur 2018 ⁵⁷	USA	General and plastic surgery	Book chapter	To discuss the barriers to management of acute postoperative pain.	N/A

Continued

Table 1 Continued

Author and year	Country	Specific surgical group	Study design	Study aim	Sample size
Wang and colleagues, 2017 ⁵⁸	China	Thoracic surgery	Prospective cohort	To describe postoperative pain follow thoracotomy and video-assisted thoracic surgery.	262
Yelton and Jildeh 2023 ⁵⁹	USA	N/A	Narrative review	To highlight some of the disparities [of pain management and rehabilitation] while emphasising the importance of cultural competency in pain control and rehabilitation, particularly within the orthopaedic field.	N/A
Zaccagnino and colleagues, 2017 ⁶⁰	USA	N/A	Narrative review	To describe the role of a comprehensive perioperative pain service to reduce costs and improve outcomes for surgical patients.	N/A

and other concerns'.⁵⁴ We delineated five gaps in continuity of care. First, there is a need for multidisciplinary patient follow-up after hospital discharge, including assessment, adequate treatments, and referral.^{26,29,31,35,39,43,44,52,54–56,58} Second, there was a gap in communication and care coordination amongst healthcare providers.^{25,39,44,55,56} There needs to be a better understanding of the division in responsibility of care in pain management.^{25,39} Funding and operational resources are required to develop effective specialised pain services that are accessible.^{15,29,39,45,56,60} Another gap lies in equity in access especially for racial and ethnic minorities, including transportation, availability of primary care, and affordability.^{31,33,35,47,57,59} Meissner and Zaslansky⁴⁵ remarked 'there remain gaps between outcomes obtained in [randomised controlled trials] as opposed to the clinical routine. Questions remain such as how to set up effective specialized pain services in many different settings, worldwide'.⁴⁵ Lastly, there was a lack of standardised practice protocols to provide consistent pain management.^{40,58}

Individualised pain management

The need for individualised pain management was clearly identified, including difficulty in accurate patient stratification of the expected degree and trajectory of postsurgical pain.^{25,45,48,56} Next, pain management strategies and education are often not tailored to patient and surgical risks of postsurgical pain, including increased follow-up for high-risk patients.^{15,43,48,55,56} There was also a gap in individualised discharge planning and education on pain management, rehabilitation, and contact.^{44,52,54–56} As Specht and colleagues⁵⁴ summarised, 'there is a need to develop in partnership with each individual patient a post-discharge plan of care and rehabilitation to meet their individual needs, preferences, and mode of motivation'.

Support for specific patient populations

There were challenges unique to certain populations that need to be addressed. Patients with opioid tolerance^{28,36,43,51,56} 'have higher rest and dynamic pain scores, greater opioid use, and require frequent prescription alterations compared with opioid-naïve patients'.⁵¹ Similarly, patients with chronic pain may have 'tolerance to opioid medications, coexisting depression and anxiety, and higher levels of postoperative pain [which] all add to the difficulty of adequately treating [postoperative] pain'.³⁶ In particular, patients with chronic pain related to cancer may struggle with 'preoperative pain (i.e. rapidly growing tumours and painful neuropathies), opioid tolerance, and contraindications to non-opioid analgesics or regional anesthesia', with a rate of persistent post-surgical pain ranging from 5% to 59%.²⁸ The increased risk of persistent postsurgical pain (40%) was also present in older surgical patients, which may be an underestimate as a result of underreporting of pain in older individuals.³⁰ The physiology, comorbidity, cognition, and polypharmacy considerations in older individuals necessitated the 'need for a careful and nuanced approach to treating both acute pain and chronic postsurgical pain in this population'.³⁰ Furthermore, there were disparities in pain management and experience among underrepresented demographic subgroups^{31,35,47,57,59}: 'there are fractures along the lines of racial, ethnic, and gender divisions', and 'the groups that are at most risk for inadequate pain relief are racial and ethnic minorities'.⁵⁷

Table 2 Themes, gaps, and health delivery domains. Details of each gap and example quotations are listed in [Supplementary material 5](#).

Themes	Gap and number of studies	Healthcare delivery domain ¹⁹
Education (n=22)	Patient education on self-management of pain medications (n=18) ^{15,27,30,32,33,35,38,40–44,46,49,50,53–55}	Care processes and infrastructure
	Setting patient expectation on postoperative pain (n=8) ^{15,27,38–40,50,53,55}	Care processes and infrastructure
	Negative perception of pain medication by patients and providers (n=7) ^{43,44,46,49,50,57,58}	Culture
	Communication amongst providers and patients (n=6) ^{27,33,43,44,55,57}	Care processes and infrastructure
	Provider knowledge on pain management and cultural competence (n=8) ^{30,33,35,39,43,45,57,58}	Organisational structure
Provision of continuity of care (n=21)	Multidisciplinary patient follow-up after hospital discharge (n=12) ^{26,29,31,35,39,43,44,52,54–56,58}	Care processes and infrastructure
	Communication and care coordination amongst healthcare providers (n=5) ^{25,39,44,55,56}	Care processes and infrastructure
	Clear division of responsibility of care with regards to postoperative pain management (n=2) ^{25,39}	Care processes and infrastructure
	Resources to facilitate access to effective specialised pain services (n=6) ^{15,29,39,45,56,60}	Finances, capacity, and organisational structure
	Equity in access to quality and efficient healthcare (n=6) ^{31,33,35,47,57,59}	Patients, capacity, finances, care processes and infrastructure, culture
Individualised pain management (n=10)	Standardised practice protocols (n=2) ^{40,58}	Care processes and infrastructure
	Accurate patient stratification of the degree and trajectory of postsurgical pain (n=4) ^{25,45,48,56}	Care processes and infrastructure, patients
	Pain management strategies and education tailored to patient and surgical risks of postsurgical pain (n=5) ^{15,43,48,55,56}	Care processes and infrastructure, patients
Support for specific patient populations (n=11)	Individualised discharge planning and education (n=5) ^{44,52,54–56}	Care processes and infrastructure
	Patients with opioid tolerance (n=5) ^{28,36,43,51,56}	Care processes and infrastructure
	Patients with chronic pain (n=2) ^{36,56}	Patients, care processes and infrastructure
	Patients with chronic cancer pain (n=1) ²⁸	Patients, care processes and infrastructure
	Older surgical patients (n=1) ³⁰	Patients, care processes and infrastructure
Research and knowledge translation (n=10)	Underrepresented demographic subgroups (racial and ethnic minorities, non-English speakers, gender, insurance status) (n=5) ^{31,35,47,57,59}	Patients, care processes and infrastructure, culture
	Research and development for effective pharmacologic and non-pharmacological strategies to prevent or treat persistent postsurgical pain (n=6) ^{2,30,34,35,48,56}	Organisational structure
	Translation of evidence-based findings into clinical practice and patient education (n=3) ^{35,40,45}	Care processes and infrastructure
	Evidence-based quality measures to guide and promote postoperative pain management (n=1) ³⁷	Care processes and infrastructure
	Research involving underrepresented demographic subgroups (racial and ethnic minorities, gender, insurance status) (n=1) ⁵⁹	Care processes and infrastructure, culture

Research and knowledge translation

Included studies identified a lack of evidence-based pharmacologic and non-pharmacologic strategies to prevent or treat persistent postsurgical pain.^{2,30,34,35,48,56} Moreover, gaps in knowledge translation were apparent, as differences exist between research findings and what is delivered via clinical practice and patient education.^{35,40,45} As Tawfic and colleagues⁵⁶ remarked, ‘not only do we have a limited grasp on when to treat, but we are also uncertain about what is both effective and safe’, and ‘well-designed clinical trials are required to identify the role of the current pain pharmacological agents, in addition to focusing on internal factors (such as psychological factors) in preventing and treating [chronic postsurgical pain]’. Moreover, we lack quality indicators to assess the delivery of pain management care. A systematic review of quality measures found only three on postoperative

pain management: multimodal analgesia, limiting opioid use in opioid-naïve patients, and monitoring opioid refills.³⁷ Importantly, more studies need to be performed in under-represented groups.⁵⁹

Discussion

Acute and chronic pain after surgery are prevalent and expensive problems, with inadequate management contributing to negative recovery trajectories and the development of persistent postsurgical pain.³ This scoping review of pain management in the first 3 months after discharge following inpatient noncardiac surgeries identified and synthesised gaps that spanned all domains of health delivery. Despite substantial variations across healthcare systems, we identified five common themes: education, provision of continuity of care, individualised pain management, support for patient

populations, and research and knowledge translation. With limited pharmacologic options to prevent the development of persistent postsurgical pain,¹⁶ these health delivery gaps represent potential targets for future process design and research aimed at mitigating the transition from acute pain to persistent postsurgical pain. Anaesthetists have an opportunity to play leadership roles in multidisciplinary perioperative pain management beyond the hospital setting to help better understand and manage pain during a critical and vulnerable period of acute pain control and chronic pain development.

This scoping review highlights several areas that deserve further attention from quality improvement and research. Firstly, anaesthetists and perioperative care providers need to more frequently engage with patients, caregivers, and health providers in pain education, using verbal and written approaches with checks for understanding and reviews at multiple time points.^{41,61} Tailored patient education, counselling, and shared decision-making can reduce postoperative opioid use and pain scores, anxiety, length of hospital stay, and emergency room visits, and improve patient satisfaction and physical function.^{42,55,62–66} For example, a systematic review found that psychoeducation about surgery and pain reduced the duration of hospitalisation by 1–3 days.⁶⁴ Involvement of patients and caregivers in the design of perioperative patient education may help increase engagement and feasibility. Educating healthcare providers is also important to optimise pain control and avoid delays in treatment. In addition, health providers need to be conscientious of potential biases and disparities in pain management, including race, ethnicity, and gender.^{57,67–69} To address this, more education about pain management and cultural competence must be incorporated into the curriculums of health science faculties and continuing education.^{59,70}

Secondly, health systems need to create standardised, patient-centred care pathways to provide continuity of care in pain management. Access to transitional pain services should be expanded. Postoperative pain medication management follow-up should be in place for patients who continue to have pain beyond 7 days,³⁵ with coordinated multidisciplinary management of pain medication tapering and timely referral to chronic pain specialists.³⁹ Electronic health records (EHRs) order sets optimised for each surgery may help reduce unnecessary discharge opioid prescriptions.⁷¹

Thirdly, we need to better understand and provide individualised postoperative pain management. This needs to start with developing accessible and accurate tools to assess pain and function,⁷² particularly in outpatient settings. The increasing availability of data from EHRs and consumer devices may help bridge the data gap of pain experiences in the outpatient setting, and shed light on the transition between acute and chronic pain. Moreover, a population cohort involving genome-wide analysis may reveal genetic factors associated with acute and chronic postoperative pain.⁷³

Fourthly, patients with chronic opioid medications, cancer pain, and older individuals deserve further clinical and research attention in post-discharge pain management. Health systems need to be adaptable to facilitate equitable, tailored strategies for vulnerable clinical and demographic populations including race, ethnicity, and gender.^{57,67–69}

Finally, we identified important gaps in research and knowledge translation. Healthcare providers have limited data and understanding of the best approaches to pain management to mitigate the transition from acute to

persistent postsurgical pain in and out of hospital. The knowledge gap is intricately linked to many of the above themes. For example, the difficulty in delivering individualised care is closely related to the lack of evidence-based risk stratification tools and the dearth of studies in underrepresented groups. As health system resources are finite, we need data-driven strategies to optimise resource allocation and provide consistent pain management at both population and individual levels.

Although similar gaps exist in diverse health systems, health providers and administrators must review the gaps and design tailored strategies that would be feasible for their respective milieu, and increase knowledge sharing to help integrate best practices in low-income countries. Cost analysis research demonstrated that preoperative patient education is associated with decreased overall healthcare expenditures in elective spine and hip arthroplasty surgeries.^{74,75} Similarly, a national cohort study showed that hip fracture care adhering to a bundle of performance measures that included systematic pain assessment translated into cost savings within 1 yr after surgery.⁷⁶ Thus, the costs of intervening on the gaps must be viewed in the context of potential cost savings and increased efficiency of the overall health system. Importantly, our review showed that the potential solutions to many of the gaps (e.g. education and provider communication) need not be resource-intensive, but would require working with stakeholders to redesign post-discharge pain management to align with patients' journey and priorities of recovery.

Strength and limitations

There are several strengths of this review. The search was comprehensive, providing a contemporary synthesis of the literature. Both inductive and deductive analyses were performed to delineate gaps and themes to provide tangible targets for intervention. There was diversity in geographic location of the included papers, representing perspectives from around the world. Several limitations also exist. The operationalisation of the term 'gap' can be subjective, although we were as inclusive as possible during screening and data extraction. We applied an inductive and deductive framework to the categorisation of the gaps, but there may be alternative schemes of organisation. Half of the studies were on specific surgical populations, which may limit the generalisability of our findings in other surgeries. This review may not have captured gaps from all types of health systems globally because of potential publication bias, limited research in low-income countries, and the inclusion of only English-language articles. Moreover, because of the variations in health delivery around the world, it may be difficult to ascertain the root cause of the gaps (e.g. lack of implementation of best practices and provision of existing services, vs insufficient resources and pathways). Nevertheless, our findings highlight the multifactorial, interlinked nature of health delivery gaps and provide a starting point to guide further research and quality improvement. More qualitative and quantitative studies across diverse healthcare systems, surgery types, and demographics would be helpful in elucidating health delivery needs. Importantly, although there is an assumption that the identified gaps are important to address, further research is needed to conclude whether intervention of many of the identified gaps would improve or worsen patient outcomes.

Conclusions

This scoping review revealed gaps during a critical period in postoperative pain management spanning all domains of health delivery. These gaps represent targets for care improvement and research to improve perioperative care and longer-term patient-centred outcomes. Anaesthetists have an opportunity to play a leadership role to facilitate individualised, streamlined pain management during the first few months after surgery.

Authors' contributions

Study design: all authors

Data extraction: JK, MD, KK, MH

Data analysis: JK, MD, KK, MH, DM

Search strategy and article management: LB

Interpretation: JK, MD, KK, MH, DM, JP, HS, JO, AF

Manuscript first draft: JK, MD, KK, MH, DM

Manuscript revisions: all authors

Declarations of interest

JK received salary support as the Clinical Data Lead, St. Paul's Hospital, Vancouver, BC, Canada, for the Project 'Reducing Opioid Use for Pain Management' from the Canadian Digital Technology Supercluster and Consortium (Careteam Technologies Inc, Thrive Health Inc, Excelar Technologies (Connected Displays Inc), Providence Health Care Ventures Inc, and Xerus Inc). JK provided paid consulting for Careflow Technologies (Connected Displays Inc), funded via Providence Health Care Ventures, Vancouver, BC, Canada. Starting September 2023, JK will receive research and salary support for Project 'Continuous Connected Patient Care' in patients with high perioperative risk, funded by the Canadian Digital Supercluster and Consortium: Medtronic Canada ULC, Cloud Diagnostics Canada ULC, Excelar Technologies (Connected Displays Inc), Providence Health Care Ventures Inc, 3D Bridge Solutions Inc, and FluidAI (NERv Technology Inc) in Vancouver, BC, Canada. DM receives salary support from The Ottawa Hospital Anesthesia Alternate Funds Association, A Clinical Research Chair from the University of Ottawa Faculty of Medicine, and the Physician Services Incorporated Mid-Career Knowledge Translation Fellowship Award (Ottawa, ON, Canada). The remaining authors do not have any conflicts of interest to disclose. JK and AF gratefully acknowledge research support from the Providence Health Care Department of Anesthesia, Vancouver, BC, Canada.

Acknowledgements

We are thankful for the time and advice from two patient partners (Kathryn Gamble and anonymous), who highlighted the challenges of post-discharge pain management and inspired this review.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.bja.2023.08.006>.

References

- Clarke H, Soneji N, Ko DT, Yun L, Wijeyesundera DN. Rates and risk factors for prolonged opioid use after major

- surgery: population based cohort study. *BMJ* 2014; **348**: g1251
- Gan TJ. Poorly controlled postoperative pain: prevalence, consequences, and prevention. *J Pain Res* 2017; **10**: 2287–98
- Richebé P, Capdevila X, Rivat C. Persistent postsurgical pain: pathophysiology and preventative pharmacologic considerations. *Anesthesiology* 2018; **129**: 590–607
- Jivraj NK, Raghavji F, Bethell J, et al. Persistent postoperative opioid use: a systematic literature search of definitions and population-based cohort study. *Anesthesiology* 2020; **132**: 1528–39
- Pagé MG, Kudrina I, Zomahoun HTV, et al. A systematic review of the relative frequency and risk factors for prolonged opioid prescription following surgery and trauma among adults. *Ann Surg LWV* 2020; **271**: 845–54
- Lawal OD, Gold J, Murthy A, et al. Rate and risk factors associated with prolonged opioid use after surgery: a systematic review and meta-analysis. *JAMA Netw Open Am Med Assoc* 2020; **3**, e207367
- Rosenberger DC, Pogatzki-Zahn EM. Chronic post-surgical pain—update on incidence, risk factors and preventive treatment options. *BJA Educ* 2022; **22**: 190–6
- Fletcher D, Stamer UM, Pogatzki-Zahn E, et al. Chronic postsurgical pain in Europe: an observational study. *Eur J Anaesthesiol* 2015; **32**: 725–34
- World Health Organization. ICD-11 for Mortality and morbidity Statistics. Available from: <https://icd.who.int/browse11/l-m/en#/http://id.who.int/icd/entity/985186256?data=%7B%22dataType%22%3A%22pc%22%2C%22postCoordinationCodeSet%22%3A%7B%22stemId%22%3A%22http%3A%2F%2Fid.who.int%2Ficd%2Fentity%2F985186256%22%2C%22axisToValueIds%22%3A%7B%22hasAlternativeSeverity%22%3A%5B%7B%22stemId%22%3A%22http%3A%2F%2Fid.who.int%2Ficd%2Fentity%2F171227995%22%7D%5D%7D%7D> (accessed 7 June 2023).
- Treede RD, Rief W, Barke A, et al. A classification of chronic pain for ICD-11. *Pain* 2015; **156**: 1003–7
- Weibel S, Neubert K, Jeltz Y, et al. Incidence and severity of chronic pain after caesarean section: a systematic review with meta-analysis. *Eur J Anaesthesiol* 2016; **33**: 853–65
- Wise B, Wilson LZ, Taber DJ, Pilch NA, Rohan V, Fleming JN. The impact of pretransplant opioid exposure on healthcare utilization and costs in kidney transplant. *Pharmacother J Hum Pharmacol Drug Ther* 2021; **41**: 6–13
- Hopman WM, Belliveau P, Gilron I. Chronic pain, healthcare utilization, and quality of life following gastrointestinal surgery. *Can J Anesth* 2012; **59**: 670
- Weir S, Samnaliev M, Kuo TC, et al. The incidence and healthcare costs of persistent postoperative pain following lumbar spine surgery in the UK: a cohort study using the Clinical Practice Research Datalink (CPRD) and Hospital Episode Statistics (HES). *BMJ Open* 2017; **7**, e017585
- Kojic K, Clarke H. Important considerations with respect to reducing the transition from acute to persistent postoperative pain. *Expert Opin Pharmacother* 2021; **22**: 779–82
- Carley ME, Chaparro LE, Choinière M, et al. Pharmacotherapy for the prevention of chronic pain after surgery in adults: an updated systematic review and meta-analysis. *Anesthesiology* 2021; **135**: 304–25

17. Blay Jr E, Nooromid MJ, Bilimoria KY, et al. Variation in post-discharge opioid prescriptions among members of a surgical team. *Am J Surg* 2018; **216**: 25–30
18. Regenbogen SE, Cain-Nielsen AH, Norton EC, Chen LM, Birkmeyer JD, Skinner JS. Costs and consequences of early hospital discharge after major inpatient surgery in older adults. *JAMA Surg* 2017; **152**, e170123
19. Piña IL, Cohen PD, Larson DB, et al. A framework for describing health care delivery organizations and systems. *Am J Public Health* 2015; **105**: 670–9
20. Cook RI, Render M, Woods DD. Gaps in the continuity of care and progress on patient safety. *BMJ* 2000; **320**: 791–4
21. Joanna Briggs Institute. Appendix 5.1: Critical Appraisal Instrument for Studies Reporting Prevalence Data - JBI Manual for Evidence Synthesis - JBI Global Wiki [Internet]. JBI Man. Evid. Synth. Available from: <https://jbi-global-wiki.refined.site/space/MANUAL/468835/Appendix+5.1%3A+Critical+Appraisal+Instrument+for+Studies+Reporting+Prevalence+Data> (accessed 7 June 2023).
22. Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018; **169**: 467–73
23. Godin K, Stapleton J, Kirkpatrick SI, Hanning RM, Leatherdale ST. Applying systematic review search methods to the grey literature: a case study examining guidelines for school-based breakfast programs in Canada. *Syst Rev* 2015; **4**: 1–10
24. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006; **3**: 77–101
25. Andrés JD, Pogatzki-Zahn E, Huygen F, Neugebauer E, Fawcett W. Controlling acute pain to improve the quality of postoperative pain management: an update from the European Society of Regional Anesthesia meeting held in Maastricht (September 2016). *Pain Manag* 2017; **7**: 513–22
26. Beloeil H, Sulpice L. Peri-operative pain and its consequences. *J Visc Surg* 2016; **153**: S15–8
27. Bleicher J, Esplin J, Blumling AN, et al. Expectation-setting and patient education about pain control in the perioperative setting: a qualitative study. *J Opioid Manag* 2021; **17**: 455–64
28. Cata JP, Corrales G, Speer B, Owusu-Agyemang P. Post-operative acute pain challenges in patients with cancer. *Best Pract Res Clin Anaesthesiol* 2019; **33**: 361–71
29. Morlion B. Expert interview: pain management treatment gap after discharge. 2019. Available from: <https://www.efort.org/popm08/> (accessed 7 June 2023).
30. Esses G, Deiner S, Ko F, Khelemsky Y. Chronic post-surgical pain in the frail older adult. *Drugs Aging* 2020; **37**: 321–9
31. Finnegan MA, Shaffer R, Remington A, Kwong J, Curtin C, Hernandez-Boussard T. Emergency department visits following elective total hip and knee replacement surgery: identifying gaps in continuity of care. *J Bone Jt Surg Am* 2017; **99**: 1005–12
32. Gangavalli AK, Malige A, Rehman S, Nwachuku CO. Patient comprehension and compliance survey to assess postoperative pain regimens in the orthopaedic trauma population. *J Orthop Trauma* 2017; **31**: e190–4
33. Garcia JBS, Bonilla P, Kraychette DC, Flores FC, de Valtolina EDP, Guerrero C. Optimizing post-operative pain management in Latin America. *Rev Bras Anesthesiol* 2017; **67**: 395–403
34. Hadlandsmayth K. Preventing chronic post-surgical pain and prolonged opioid use: The Perioperative Pain Self-Management Program. *NIH RePORTER* 2021
35. Health Quality Ontario. Recommendations for adoption: opioid prescribing for acute pain. Available from <http://www.hqontario.ca/Portals/0/documents/evidence/quality-standards/qs-opioid-acute-pain-recommendations-for-adoption-en.pdf>.
36. Jackman C. Perioperative pain management for the chronic pain patient with long-term opioid use. *Orthop Nurs* 2019; **38**: 159–63
37. Joseph JM, Gori D, Curtin C, et al. Gaps in standardized postoperative pain management quality measures: a systematic review. *Surgery* 2022; **171**: 453–8
38. Kennedy D, Wainwright A, Pereira L, et al. A qualitative study of patient education needs for hip and knee replacement. *BMC Musculoskelet Disord* 2017; **18**: 413
39. Klueh MP, Sloss KR, Dossett LA, et al. Postoperative opioid prescribing is not my job: a qualitative analysis of care transitions. *Surgery* 2019; **166**: 744–51
40. Kuusniemi K, Pöyhä R. Present-day challenges and future solutions in postoperative pain management: results from PainForum 2014. *J Pain Res* 2016; **9**: 25–36
41. Lemay CA, Saag KG, Franklin PD. A qualitative study of the postoperative pain management educational needs of total joint replacement patients. *Pain Manag Nurs* 2019; **20**: 345–51
42. Lemay CA, Lewis CG, Singh JA, Franklin PD. Receipt of pain management information preoperatively is associated with improved functional gain after elective total joint arthroplasty. *J Arthroplasty* 2017; **32**: 1763–8
43. Lopes A, Seligman Menezes M, Antonio Moreira de Barros G. Chronic postoperative pain: ubiquitous and scarcely appraised: narrative review. *Braz J Anesthesiol Engl Ed* 2021; **71**: 649–55
44. Makimoto K, Fujita K, Konno R. Review and synthesis of the experience of patients following total hip or knee arthroplasty in the era of rapidly decreasing hospital length of stay. *Jpn J Nurs Sci* 2020; **17**, e12361
45. Meissner W, Zaslansky R. A survey of postoperative pain treatments and unmet needs. *Best Pract Res Clin Anaesthesiol* 2019; **33**: 269–86
46. Nasir M, Ahmed A. Knowledge about postoperative pain and its management in surgical patients. *Cureus* 2020; **12**: e6685
47. Nguyen KH, Rambachan A, Ward DT, Manuel SP. Language barriers and postoperative opioid prescription use after total knee arthroplasty. *Explor Res Clin Soc Pharm* 2022; **7**, 100171
48. Pogatzki-Zahn E, Lavandhomme P. Prevention of chronic post-surgical pain. Int Assoc study pain 2020 prevention of pain. Published 9 July 2021. Available from: <https://www.iasp-pain.org/resources/fact-sheets/prevention-of-chronic-post-surgical-pain/> (accessed 7 June 2023).
49. Rahman R, Wallam S, Zhang B, et al. Appropriate opioid use after spine surgery: psychobehavioral barriers and patient knowledge. *World Neurosurg* 2021; **150**: e600–12
50. Reaza-Alarcón A, Rodríguez-Martín B. Effectiveness of nursing educational interventions in managing post-surgical pain. Systematic review. *Invest Educ Enferm* 2019; **37**: e10
51. Simpson GK, Jackson M. Perioperative management of opioid-tolerant patients. *BJA Educ Elsevier*; 2017. p. 124–8
52. Sjøveian AKH, Leegaard M. Hip and knee arthroplasty - patient's experiences of pain and rehabilitation after discharge from hospital. *Int J Orthop Trauma Nurs* 2017; **27**: 28–35

53. Smith F, Öhlén J, Persson LO, Carlsson E. Daily assessment of stressful events and coping in early post-operative recovery after colorectal cancer surgery. *Eur J Cancer Care* 2018; **27**, e12829
54. Specht K, Agerskov H, Kjaersgaard-Andersen P, Jester R, Pedersen BD. Patients' experiences during the first 12 weeks after discharge in fast-track hip and knee arthroplasty - a qualitative study. *Int J Orthop Trauma Nurs* 2018; **31**: 13–9
55. Sugar R, Jansen A, McDonald S. The red dot initiative: an analysis of postoperative visits to the emergency department. *Br Columbia Med J* 2018; **60**: 398–402
56. Tawfic Q, Kumar K, Pirani Z, Armstrong K. Prevention of chronic post-surgical pain: the importance of early identification of risk factors. *J Anesth* 2017; **31**: 424–31
57. Thakur AC. Barriers to optimal pain management in the general surgery population. In: Narayan D, Kaye AD, Vadivelu N, editors. *Perioperative pain management for general and plastic surgery*. Oxford: Oxford University Press; 2018
58. Wang H, Li S, Liang N, Liu W, Liu H, Liu H. Postoperative pain experiences in Chinese adult patients after thoracotomy and video-assisted thoracic surgery. *J Clin Nurs John Wiley Sons Inc*; 2017. p. 2744–54
59. Yelton MJ, Jildeh TR. Cultural competence and the post-operative experience: pain control and rehabilitation. *Arthrosc Sports Med Rehabil* 2023; **5**, 100733
60. Zaccagnino MP, Bader AM, Sang CN, Correll DJ. The perioperative surgical home: a new role for the acute pain service. *Anesth Analg* 2017; **125**: 1394–402
61. Clarke HA, Manoo V, Pearsall EA, et al. Consensus statement for the prescription of pain medication at discharge after elective adult surgery. *Can J Pain* 2020; **4**: 67–85
62. Chou R, Gordon DB, de Leon-Casasola OA, et al. Management of postoperative pain: a clinical practice guideline from the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists' committee on regional anesthesia, executive committee, and administrative council. *J Pain* 2016; **17**: 131–57
63. Lee CH, Liu JT, Lin SC, Hsu TY, Lin CY, Lin LY. Effects of educational intervention on state anxiety and pain in people undergoing spinal surgery: a randomized controlled trial. *Pain Manag Nurs* 2018; **19**: 163–71
64. Horn A, Kaneshiro K, Tsui BCH. Preemptive and preventive pain psychoeducation and its potential application as a multimodal perioperative pain control option: a systematic review. *Anesth Analg* 2020; **130**: 559–73
65. Uhrbrand P, Rasmussen MM, Haroutounian S, Nikolajsen L. Shared decision-making approach to taper postoperative opioids in spine surgery patients with pre-operative opioid use: a randomized controlled trial. *Pain* 2022; **163**: e634
66. Ghaddaf AA, Alsharef JF, Alhindi AK, et al. Influence of perioperative opioid-related patient education: a systematic review and meta-analysis. *Patient Educ Couns* 2022; **105**: 2824–40
67. Salamonson Y, Everett B. Demographic disparities in the prescription of patient-controlled analgesia for post-operative pain. *Acute Pain* 2005; **7**: 21–6
68. Lee P, Le Saux M, Siegel R, et al. Racial and ethnic disparities in the management of acute pain in US emergency departments: meta-analysis and systematic review. *Am J Emerg Med* 2019; **37**: 1770–7
69. Safdar B, Heins A, Homel P, et al. Impact of physician and patient gender on pain management in the emergency department—a multicenter study. *Pain Med* 2009; **10**: 364–72
70. Watt-Watson J, McGillion M, Hunter J, et al. A survey of prelicensure pain curricula in health science faculties in Canadian universities. *Pain Res Manag Hindawi* 2009; **14**: 439–44
71. Mou Z, Chau H, Kalavacherla S, et al. Tailored order set in the electronic health record decreases postoperative opioid prescriptions. *Surgery* 2022; **172**: 677–82
72. Baamer RM, Iqbal A, Lobo DN, Knaggs RD, Levy NA, Toh LS. Utility of unidimensional and functional pain assessment tools in adult postoperative patients: a systematic review. *Br J Anaesth* 2022; **128**: 874–88
73. Li S, van Boekel RLM, van den Heuvel SAS, Coenen MJH, Visser KCP. Pain predict genetics: protocol for a prospective observational study of clinical and genetic factors to predict the development of postoperative pain. *BMJ Open* 2022; **12**, e066134
74. Burgess LC, Arundel J, Wainwright TW. The effect of preoperative education on psychological, clinical and economic outcomes in elective spinal surgery: a systematic review. *Healthcare* 2019; **7**: 48
75. Moulton LS, Evans PA, Starks I, Smith T. Pre-operative education prior to elective hip arthroplasty surgery improves postoperative outcome. *Int Orthop* 2015; **39**: 1483–6
76. Kristensen PK, Søgaard R, Thillemann TM, Søballe K, Johnsen SP. High quality of care did not imply increased hospital spending—nationwide cohort study among hip fracture patients. *Int J Qual Health Care* 2019; **31**: 22–9. Oxford University Press

Handling Editor: Jonathan Hardman