Comprehensive Clinical Strategy for Hypertension Management: Nutritional Pathophysiology and Evidence-Based Dietary Protocols

The clinical management of hypertension, defined by the updated 2025 American College of Cardiology (ACC) and American Heart Association (AHA) guidelines as a persistent blood pressure (BP) of \ge 130/80 mmHg, has entered a new era of precision nutrition and risk-stratified intervention. Hypertension remains the primary modifiable risk factor for cardiovascular disease (CVD), stroke, and chronic kidney disease (CKD), affecting approximately 50% of the adult population in the United States and a growing proportion globally. The evolution of clinical standards reflects a deeper understanding of the linear relationship between blood pressure and vascular damage; specifically, for every 10 mmHg reduction in systolic blood pressure (SBP), there is a corresponding 17% reduction in coronary heart disease, a 27% reduction in stroke, and a 28% reduction in heart failure. Consequently, the 2025 guidelines have shifted toward a more intensive management paradigm, encouraging targets of <120/80 mmHg for most adults to mitigate not only macrovascular events but also the risk of cognitive impairment and dementia.

Overview of Hypertension and the Pathophysiology of Blood Pressure Control

Hypertension is a multifaceted hemodynamic disorder characterized by the dysregulation of systemic vascular resistance and cardiac output. The basic physiological equation BP = CO \times SVR, where CO represents cardiac output and SVR represents systemic vascular resistance, underscores the mechanisms by which dietary habits influence blood pressure. Excessive sodium intake expands extracellular fluid volume, thereby increasing CO, while inadequate intake of vasodilatory minerals like potassium and magnesium increases SVR by promoting vasoconstriction and endothelial dysfunction.

The diagnostic landscape has been refined in 2025 to emphasize the use of the PREVENT equation, which estimates 10-year and 30-year cardiovascular risk. This tool replaces the older Pooled Cohort Equations and integrates broader data sets, including heart failure risk and markers of chronic kidney disease. For individuals in the "Stage 1" category (130–139/80–89 mmHg) with a PREVENT 10-year risk of <7.5\%, the guidelines advocate for a three-to-six-month window of intensive lifestyle modification before initiating pharmacotherapy. However, if target thresholds are not met through diet and exercise alone, pharmacological intervention is now strongly recommended even in these lower-risk cohorts to prevent long-term target-organ damage.

Current Clinical Categories and Diagnostic Thresholds

The 2025 framework maintains the core categories established in 2017 but provides nuanced instructions for management based on risk stratification. The term "Hypertensive Urgency" has been officially retired and replaced with "Severe Hypertension," which refers to elevations of >180/120 mmHg without acute target-organ damage.

BP Category	Systolic BP (mmHg)	Relationship	Diastolic BP (mmHg)	2025 Clinical Intervention Threshold
Normal	< 120	and	< 80	Annual screening; health promotion
Elevated	120–129	and	< 80	Non-pharmacological (lifestyle) therapy
Stage 1 Hypertension	130–139	or	80–89	Lifestyle + Rx if risk \ge 7.5\% or comorbidities
Stage 2 Hypertension	\ge 140	or	\ge 90	Immediate pharmacological + lifestyle
Severe Hypertension	> 180	and/or	> 120	Urgent evaluation; outpatient or inpatient

The role of diet in this continuum is foundational. Nutritional therapy serves as the first-line defense in the "Elevated" and "Stage 1" categories and acts as a potent synergistic adjunct to medication in "Stage 2" and "Severe" hypertension. In fact, the Dietary Approaches to Stop Hypertension (DASH) eating plan has demonstrated blood pressure-lowering effects comparable to single-agent drug therapy in many clinical trials.

Core Dietary Principles: The Electrolyte and Macro-Nutrient Synergy

The efficacy of nutritional intervention in hypertension management is not derived from a single nutrient but from a synergistic pattern of mineral intake and fat quality. The primary therapeutic targets of this approach include the modulation of the sodium-potassium pump, the reduction of oxidative stress in the endothelium, and the inhibition of the renin-angiotensin-aldosterone system (RAAS).

Sodium Reduction and the Natriuretic Response

Sodium chloride is the most pervasive dietary contributor to elevated blood pressure. Excessive sodium intake leads to water retention, increasing the intravascular volume and stretching the vascular walls. For individuals categorized as "salt-sensitive," even minor increases in sodium can trigger significant rises in blood pressure. The 2025 guidelines and the DASH protocol set a standard ceiling of 2,300 mg of sodium per day, with the more therapeutic "low-sodium" DASH variant aiming for 1,500 mg per day.

Clinical evidence suggests that reducing sodium by approximately 1,000 mg per day can lower SBP by 5–6 mmHg in hypertensive individuals. This reduction is mediated by the decreased

stimulation of the sympathetic nervous system and the downregulation of RAAS activity. To achieve these targets, clinicians must advise patients to transition away from ultra-processed "convenience" foods, which account for the vast majority of sodium in the modern diet.

Potassium, Magnesium, and Calcium: The Vasodilatory Trio

While sodium causes volume expansion, potassium serves as its physiological antagonist. High potassium intake stimulates the kidneys to excrete more sodium through a process called natriuresis and promotes the relaxation of vascular smooth muscle cells. The typical goal for potassium intake in the DASH diet is 4,700 mg per day, though this must be monitored in patients with CKD.

Magnesium and calcium are equally critical for blood pressure regulation. Magnesium helps modulate the transport of other electrolytes across cell membranes and acts as a natural calcium channel blocker, preventing the excessive constriction of blood vessels. Calcium, meanwhile, is essential for the stabilization of the cellular membrane and the proper functioning of the vascular endothelium.

Mineral	Recommended Daily Intake (2,000 kcal)	Biochemical Mechanism	Clinical Evidence
Sodium	1,500–2,300 mg	Volume expansion; RAAS activation	High-sodium intake linked to 1.9 million deaths annually
Potassium	4,700 mg	Natriuresis; vascular wall relaxation	50 mmol reduction in Na paired with K lowers SBP
Magnesium	310–420 mg	Endothelial nitric oxide support	Involved in 300+ biological responses
Calcium	1,000–1,200 mg	Vascular tone modulation	Essential for BP reduction in DASH-Sodium trials

DASH Diet Overview and Fat Quality

The Dietary Approaches to Stop Hypertension (DASH) plan is a comprehensive nutritional framework rich in fiber, protein, and minerals, while being low in saturated fats and refined sugars. Unlike many restrictive diets, DASH emphasizes the *addition* of nutrient-dense foods rather than just the *subtraction* of harmful ones. A major component of the DASH diet is the reduction of saturated fat and the elimination of trans fats, which are replaced by heart-healthy monounsaturated (MUFA) and polyunsaturated fats (PUFA).

Saturated fats, found in fatty meats and full-fat dairy, contribute to vascular inflammation and the development of atherosclerosis, which stiffens the arteries and increases blood pressure. In contrast, unsaturated fats from sources like olive oil, nuts, and fish help maintain endothelial flexibility and reduce circulating LDL cholesterol, a common comorbidity in hypertensive patients.

Recommended Foods: The Nutrient-Dense Portfolio

The clinical application of the DASH diet requires a precise understanding of food categories and serving sizes. For a standard 2,000-calorie plan, the following serving guidelines serve as

the blueprint for nutritional therapy.

Fruits and Vegetables: The Foundation of Nitric Oxide Support

Vegetables and fruits provide the highest concentration of potassium and fiber per calorie. They are also rich in dietary nitrates, which are converted in the body to nitric oxide (NO), a potent vasodilator that improves arterial elasticity.

- **Serving Goal:** 4–5 servings of vegetables and 4–5 servings of fruits daily.
- **Standard Servings:** 1 cup of raw leafy greens, 1/2 cup of cooked vegetables, or 1 medium fruit.
- **Strategic Choices:** Leafy greens (spinach, kale), cruciferous vegetables (broccoli, cauliflower), berries (anthocyanin-rich), and citrus (vitamin C supports NO bioavailability).

Whole Grains: Fiber and Metabolic Stability

Whole grains retain the bran and germ, providing significantly higher levels of magnesium and fiber than refined counterparts. Fiber slows down the absorption of carbohydrates, preventing the insulin spikes that can contribute to sodium retention in the kidneys.

- **Serving Goal:** 6–8 servings daily.
- **Standard Servings:** 1 slice of whole-wheat bread, 1/2 cup of cooked rice/pasta, or 1 ounce of dry whole-grain cereal.
- **Strategic Choices:** Oats, brown rice, barley, quinoa, and whole-wheat roti (in the Indian context).

Low-Fat Dairy: The Calcium-Peptide Connection

Low-fat and fat-free dairy products are emphasized in the DASH diet because they provide the necessary calcium and potassium without the atherosclerotic risk of saturated fats. Some dairy proteins may also have mild ACE-inhibiting properties.

- **Serving Goal:** 2–3 servings daily.
- **Standard Servings:** 1 cup of milk or yogurt, or 1.5 ounces of cheese. * **Strategic Choices:** Greek yogurt (low sugar), skim milk, and low-fat cottage cheese.

Lean Proteins: Plant-Based and Omega-3 Sources

The DASH diet shifts the protein focus from red meats to poultry, fish, and legumes. Legumes (beans, peas, lentils) are particularly beneficial as they provide plant-based protein alongside significant amounts of fiber and magnesium.

- **Serving Goal:** 6 servings or fewer of lean meat/fish per day; 4–5 servings of nuts/seeds/legumes *per week*.
- **Standard Servings:** 1 ounce of cooked meat/fish or 1 egg; 1/2 cup cooked beans or 1/3 cup nuts.
- **Strategic Choices:** Fatty fish (salmon, mackerel for omega-3s), skinless chicken breast, chickpeas, and lentils.

Foods to Avoid: Identifying Sodium and Metabolic

Triggers

Success in hypertension management is often determined by the ability to exclude high-sodium and inflammatory foods that override the benefits of nutrient-dense choices.

The "Salty Six" and Processed Food Hazards

The primary threat to sodium management is the prevalence of hidden salt in everyday foods. The "Salty Six," as identified by heart associations, include bread, cold cuts/cured meats, pizza, poultry (often injected with sodium), soup, and sandwiches.

- **Processed Meats:** Bacon, sausages, ham, and deli turkey are often cured with sodium nitrate and contain excessive salt.
- **Canned and Prepared Foods:** Standard canned vegetables and soups can contain over 800 mg of sodium per serving. Clinicians should recommend "low-sodium" versions and advise rinsing canned beans/vegetables to remove up to 40% of the salt.

Sugar-Sweetened Beverages and Refined Sugars

Recent research indicates that high sugar intake, especially high-fructose corn syrup, may be as detrimental to blood pressure as excessive salt. Sugar increases insulin levels, which stimulates the sympathetic nervous system and causes the kidneys to retain sodium.

- **Impact:** Consumption of sugary drinks is linked to an increased risk of obesity and Type 2 diabetes, both of which exacerbate hypertension.
- **Avoid:** Regular soda, sweetened coffees, energy drinks, and desserts.

Saturated Fats and Tropical Oils

While fats are essential, the DASH diet strictly limits saturated fats to less than 6% of total calories. This involves avoiding fatty cuts of red meat (lamb, pork, beef) and tropical oils like coconut and palm oil, which are high in saturated fats and can worsen lipid profiles.

- **Substitutes:** Use liquid vegetable oils like olive, canola, or mustard oil instead of butter, lard, or ghee.

Sample Daily Meal Plan: Integrating Western and Indian Options

A successful meal plan must be culturally adaptable to ensure long-term adherence. The following menus focus on a 2,000-calorie target with less than 1,500 mg of sodium.

Day 1: Western DASH Focus

- **Breakfast:** 1 cup of oatmeal (prepared without salt), topped with 1/4 cup raisins and 1 medium banana; 1 cup fat-free milk; coffee or tea.
- **Lunch:** 3 oz grilled skinless chicken breast on a whole-wheat pita with lettuce, tomato, and 1 tbsp low-fat mayonnaise; served with a medium apple and a side of raw carrot sticks.

- **Dinner:** 4 oz roasted salmon with a lemon-herb glaze; 1 cup cooked brown rice or wild rice; 1 cup steamed broccoli and a spinach salad with cucumber and unsalted vinaigrette.
- **Snack:** 1/3 cup unsalted almonds or a cup of low-fat yogurt with fresh peaches.

Day 2: Indian DASH Focus

- **Morning Drink:** Warm water with lemon or Jeera (cumin) water.
- **Breakfast:** Oats upma with mixed peas, carrots, and beans, seasoned with mustard seeds and curry leaves (no salt added); herbal tea.
- **Mid-Morning:** 1 cup papaya or watermelon cubes.
- **Lunch:** 1 cup Brown rice; 1 bowl Arhar dal (seasoned with garlic and roasted cumin); 1 bowl stir-fried Bhindi (ladyfinger) using minimal mustard oil; 1 bowl unsalted curd.
- **Evening Snack:** Roasted unsalted Makhana (fox nuts) or sprouted moong chaat with lemon juice and coriander.
- **Dinner:** 2 small Phulkas (whole wheat, no salt); 1 bowl Lauki (bottle gourd) and tomato curry; a side of cucumber and beet salad.

Strategic Nutritional Hacks for Indian Households

In Indian cooking, salt is often deeply integrated into traditional sides. Clinical advice must emphasize the following replacements to maintain flavor without the sodium burden :

Traditional Item	High-Sodium Ingredient	Low-Sodium Alternative
Pickle (Achar)	Salt, Preservatives	Fresh lemon squeeze or raw mango pieces
Papad	Sodium Bicarbonate, Salt	Roasted homemade grain crackers (no salt)
Chutney	Added Table Salt	Mint-Coriander with green chili and lemon
Table Salt	Sodium Chloride	Potassium-based salt substitute (if no CKD)
Rice	White Polished Rice	Brown rice, Ragi, Jowar, or Bajra millets

Lifestyle Considerations: Exercise, Weight, and Stress

Nutritional therapy for hypertension does not exist in a vacuum. The 2025 guidelines emphasize a "team-based" approach that integrates diet with physical activity and psychosocial support.

The Bioenergetics of Exercise

Aerobic exercise induces a physiological "shear stress" on the vascular walls, which stimulates the production of nitric oxide and improves the efficiency of the sodium-potassium pump. Regular physical activity can lead to an average reduction of 4–9 mmHg in systolic blood pressure.

- **Recommendation:** 150 minutes of moderate-intensity aerobic exercise (e.g., brisk walking, cycling) per week, supplemented by resistance training twice weekly.
- **The "Weight Loss" Multiplier:** When the DASH diet is combined with exercise and weight loss, BP reductions can be as high as 16/10 mmHg, compared to 11/7 mmHg with

DASH alone.

Adipose Tissue and Vascular Tension

Obesity is a pro-inflammatory state that directly contributes to hypertension. Adipose tissue secretes cytokines and adipokines that stimulate the sympathetic nervous system and cause vascular stiffening.

- **Clinical Impact:** A modest weight loss of 5–10\% of total body weight has been shown to improve glycemia, lower blood pressure, and in some cases, induce remission of Type 2 diabetes.
- **Mechanism:** Weight loss reduces the physical compression of the kidneys and lowers the metabolic demand for cardiac output.

Stress-Induced Hypertension and Sleep Hygiene

Psychosocial stressors and sleep disturbances (such as obstructive sleep apnea) are major contributors to "resistant hypertension." Chronic stress elevates cortisol levels, which increases the sensitivity of blood vessels to vasoconstrictors like norepinephrine.

- **Intervention:** Stress management through meditation, yoga, or deep breathing can lower SBP.
- **Sleep:** Ensuring 7–9 hours of restful sleep is vital for the nocturnal "dipping" of blood pressure, a necessary physiological process for heart health.

Special Cases: Diabetes and Obesity Comorbidities

Management becomes more complex when hypertension is part of a metabolic cluster involving Type 2 diabetes (T2DM) and obesity.

Hypertension and Type 2 Diabetes (T2DM)

Over 50% of adults with diabetes also have hypertension. Both conditions cause microvascular damage, leading to a high risk of diabetic nephropathy (kidney disease) and retinopathy (vision loss).

- **Dietary Adjustment:** The standard DASH diet can be high in carbohydrates for some T2DM patients. It is recommended to prioritize complex, high-fiber carbohydrates and pair them with lean protein and healthy fats to blunt postprandial glucose spikes.
- **Pharmacological Note:** In the presence of albuminuria (protein in the urine), ACE inhibitors or ARBs are the first-line treatment to protect the kidneys while managing BP.
- **Goal:** A target BP of <130/80 mmHg is mandatory for this group to prevent the progression of heart failure and kidney disease.

Obesity-Related Hypertension

In cases of morbid obesity where lifestyle interventions fail, the 2025 guidelines and 2026 Standards of Care for Diabetes suggest considering metabolic (bariatric) surgery. This is particularly effective for patients with a BMI \ge 30 kg/m² (\ge 27.5 for Asian Americans) who have uncontrolled hypertension and diabetes.

- **Benefits:** Surgery often leads to a rapid, sustained reduction in blood pressure and the discontinuation of multiple antihypertensive medications.

Safety Guidelines: From Sodium Limits to Medical Emergencies

Nutritional planning must include clear safety boundaries, particularly regarding the risk of electrolyte imbalances and the recognition of cardiovascular crises.

Sodium and Potassium Safety

While reducing sodium and increasing potassium is generally beneficial, certain populations face significant risks.

- **Sodium Floor:** While 1,500 mg is a common target, reducing sodium too drastically in certain heart failure patients may lead to compensatory activation of the RAAS. Consultation with a specialist is required.
- **Potassium and CKD:** Patients with Stage 3 or higher Chronic Kidney Disease (eGFR < 60 mL/min) cannot excrete potassium efficiently. In these individuals, high-potassium diets or salt substitutes can lead to hyperkalemia (dangerously high blood potassium), which can cause cardiac arrest.
- **Drug-Nutrient Interactions:** Medications such as Spironolactone, Lisinopril, and Losartan cause the body to "hold onto" potassium. Patients on these drugs should avoid potassium-based salt substitutes and "muscle-building" supplements without medical supervision.

When to Seek Immediate Medical Help

The 2025 terminology shift to "Severe Hypertension" (>180/120 mmHg) simplifies the triage process. Patients should be taught to monitor for signs of target-organ damage.

Red Flag Symptom	Potential Underlying Complication	Action Required
Severe Headache / Blurred Vision	Encephalopathy / Intracranial Hemorrhage	Call 911 Immediately
Chest Pain / Shortness of Breath	Myocardial Infarction / Pulmonary Edema	Call 911 Immediately
Sudden Numbness or Weakness	Ischemic Stroke	Call 911 Immediately
Significant Leg Swelling (Edema)	Heart Failure	Urgent Medical Visit
Blood Pressure > 180/120 (No Symptoms)	Asymptomatic Severe Hypertension	Contact Provider Today

Summary and RAG System Data Integration

For an AI-driven diet planning system (RAG), the following structured data summarizes the essential parameters for hypertension management.

Structured Nutritional Data (JSON)

```
{
  "recommended_foods": {
    "fruits_vegetables":,
    "whole_grains":,
    "low_fat_dairy":,
    "lean_proteins":,
    "flavorings":
  },
  "foods_to_avoid": {
    "high_sodium":,
    "condiments":,
    "sugary_items":,
    "saturated_fats":
  },
  "meal_plan_logic": {
    "base_calories": 2000,
    "sodium_targets": {"standard": 2300, "therapeutic": 1500},
    "potassium_target": 4700,
    "magnesium_target": 420,
    "calcium_target": 1200,
    "plate_method": {
      "vegetables_fruit": "50%",
      "whole_grains": "25%",
      "lean_protein": "25%"
    }
  },
  "key_guidelines":
}
```

Conclusion

The nutritional management of hypertension has transitioned from a general "low-salt" recommendation to a sophisticated, mineral-rich, and risk-stratified therapeutic approach. The DASH diet, particularly when culturally adapted and combined with modern pharmacological guidelines, remains the most effective non-invasive strategy for lowering blood pressure and preventing cardiovascular mortality. By focusing on the synergy of sodium reduction, increased intake of vasodilatory minerals (potassium, magnesium, calcium), and healthy weight maintenance, patients can achieve blood pressure levels that not only protect the heart and kidneys but also preserve long-term cognitive function. For healthcare professionals and RAG AI systems, the integration of these evidence-based serving sizes, safety parameters, and cultural adaptations is essential for delivering high-quality, actionable nutrition planning that meets the intensive goals of the 2025 hypertension standards.

Works cited

1. Implementing the 2025 Guideline for the Prevention, Detection ..., https://www.ahajournals.org/doi/10.1161/HYPERTENSIONAHA.125.25418 2. New in Clinical Guidance | High Blood Pressure Focus of New ACC/AHA Guideline, https://www.acc.org/latest-in-cardiology/articles/2025/10/01/01/new-in-clinical-guidance-hbp 3. AHA Clinical Slide Series: 2025 Guideline for the Prevention, Detection, Evaluation and Management of High Blood Pressure in Adults - Professional Heart Daily, https://professional.heart.org/-/media/PHD-Files-2/Science-News/Clinical-Slide-Series/2025-High-Blood-Pressure-Guideline-Clinical-Slides.pptx?sc_lang=en 4. Hypertension - World Health Organization (WHO), https://www.who.int/news-room/fact-sheets/detail/hypertension 5. Updates in the 2025 AHA/ACC Hypertension Guideline - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC12995957/ 6. Hypertensive Emergency - StatPearls - NCBI Bookshelf - NIH, https://www.ncbi.nlm.nih.gov/books/NBK470371/ 7. DASH eating plan - Healthy eating to lower your blood pressure - MC6872 - Mayo Clinic, https://mcforms.mayo.edu/mc6800-mc6899/mc6872.pdf 8. The DASH diet in diabetes related complications or comorbidities ..., https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1689467/full 9. Overview of the 2025 Hypertension Guidelines - Drug Information Group, https://dig.pharmacy.uic.edu/faqs/2025-2/december-2025-faqs/overview-of-the-2025-hypertension-guidelines/ 10. 2025 AHA/ACC/AANP/AAPA/ABC/ACCP/ACPM/AGS/AMA/ASPC/NMA/PCNA/SGIM Guideline for the Prevention, Detection, Evaluation and Management of High Blood Pressure in Adults - American Heart Association Journals, https://www.ahajournals.org/doi/10.1161/CIR.0000000000001356 11. Hypertensive Crisis: Types, Causes & Symptoms - Cleveland Clinic, https://my.clevelandclinic.org/health/diseases/24470-hypertensive-crisis 12. Management of Hypertension and Diabetes in Obesity: Non-Pharmacological Measures, https://pmc.ncbi.nlm.nih.gov/articles/PMC3095960/ 13. Weight-Loss Strategies for Prevention and Treatment of Hypertension: A Scientific Statement From the American Heart Association, https://www.ahajournals.org/doi/10.1161/HYP.0000000000000202 14. The Dash Diet and Diabetes | Diabetes Care | Newsroom - Abbott, https://www.abbott.com/en-us/corpnewsroom/diabetes-care/the-dash-diet-and-diabetes 15. DASH Diet: Care Instructions | Cardiovascular Institute of Central Florida, https://cvicfl.com/dash-diet-care-instructions/ 16. Can You Really Do DASH With Diabetes? - The Well by Northwell, https://thewell.northwell.edu/healthy-living-fitness/dash-diet-diabetes 17. DASH diet: Guide to recommended servings - Mayo Clinic, https://www.mayoclinic.org/healthy-lifestyle/nutrition-and-healthy-eating/in-depth/dash-diet/art-20050989 18. DASH Diet Indian Plan for High BP | Heart-Healthy Eating – RiteBite ..., https://maxprotein.in/blogs/health-nutrition/dash-diet-indian-diet-plan-for-high-blood-pressure 19. DASH diet: Healthy eating to lower your blood pressure - Mayo Clinic, https://www.mayoclinic.org/healthy-lifestyle/nutrition-and-healthy-eating/in-depth/dash-diet/art-20048456 20. DASH Eating Plan | NHLBI, NIH, https://www.nhlbi.nih.gov/health/dash-eating-plan 21. ACE Inhibitors and ARBs - National Kidney Foundation, https://www.kidney.org/kidney-topics/ace-inhibitors-and-arbs 22. DASH Eating Plan | Kaiser Permanente, https://mydoctor.kaiserpermanente.org/ncal/Images/DASH%20Eating%20Plan_tcm75-2794478.pdf 23. Launch of the WHO guideline on the use of lower-sodium salt substitutes,

https://www.who.int/news-room/events/detail/2025/01/27/default-calendar/launch-of-the-who-guideline-on-the-use-of-lower-sodium-salt-substitutes 24. Heart Healthy Diet: Simple Indian Foods Guide, https://www.nivabupa.com/health-wellness-articles/heart-healthy-diet-indian-foods.html 25. Heart-Healthy Indian Diet | Pune Cardiology Clinic, https://punecardiologyclinic.com/heart-healthy-indian-diet-what-to-eat-and-what-not/ 26. DASH Eating Plan: An Eating Pattern for Diabetes Management - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC5439361/ 27. Indian Vegetarian Foods That Can Lower Blood Pressure - Everyday Health, https://www.everydayhealth.com/hypertension/indian-vegetarian-foods-that-can-lower-blood-pressure/ 28. Low-Sodium Indian Meal Plan for a Week- Your Ultimate Blood Pressure Diet, https://www.lupindiagnostics.com/blog/low-sodium-indian-meal-plan-for-a-week-your-ultimate-blood-pressure-diet 29. DASH Eating Plan Sample Menu - Cigna Healthcare, https://www.cigna.com/knowledge-center/hw/dash-eating-plan-sample-menu-zp3302 30. DASH diet: Sample menus - Mayo Clinic, https://www.mayoclinic.org/healthy-lifestyle/nutrition-and-healthy-eating/in-depth/dash-diet/art-20047110 31. A Week With the DASH Eating Plan - nhlbi - NIH, https://www.nhlbi.nih.gov/sites/default/files/publications/WeekOnDASH.pdf 32. 8. Obesity and Weight Management for the Prevention and ..., https://diabetesjournals.org/care/article/49/Supplement_1/S166/163915/8-Obesity-and-Weight-Management-for-the-Prevention 33. YOUR KIDNEYS AND HIGH POTASSIUM (HYPERKALEMIA), https://www.kidney.org/sites/default/files/01-10-7269_ABG_PatBro_Hyperkalemiap7.pdf 34. Hyperkalemia: Medicines That Can Raise Potassium - WebMD, https://www.webmd.com/a-to-z-guides/medication-affect-potassium-levels 35. 11 Medications That Can Cause High Potassium Levels, And How to Prevent Them - GoodRx, https://www.goodrx.com/conditions/hyperkalemia/what-drugs-can-cause-high-potassium-levels 36. What medications can affect potassium levels? - Drugs.com, https://www.drugs.com/medical-answers/medications-affect-potassium-levels-3566412/ 37. Drugs That Cause Hyperkalemia - Can You Name These Seven? - Med Ed 101, https://www.meded101.com/drugs-that-cause-hyperkalemia-high-potassium-ace-arbs/ 38. Top 10 Things to Know About the AHA/ACC High Blood Pressure Guideline, https://www.heart.org/en/health-topics/high-blood-pressure/the-facts-about-high-blood-pressure/high-bp-top-10