

## Properties of Eigen values:

① Sum of eigen values = trace of matrix  $A$   $\lambda_1 + \lambda_2 + \lambda_3 =$

$$\lambda_1 + \lambda_2 + \lambda_3 = \text{trace}(A)$$

$$\begin{bmatrix} \lambda_1 & & \\ & \lambda_2 & \\ & & \lambda_3 \end{bmatrix}$$

Product of eigen values = determinant of matrix

$$\lambda_1 \cdot \lambda_2 \cdot \lambda_3 = |A|$$

## Properties of Eigen values

16:15

① Sum of eigen values = trace of matrix  $A$   $\lambda_1 + \lambda_2 + \lambda_3 =$   
 $\lambda_1 + \lambda_2 + \lambda_3 = \text{trace}(A)$

$$\begin{bmatrix} \lambda_1 & & \\ & \lambda_2 & \\ & & \lambda_3 \end{bmatrix}$$

Product of eigen values = determinant of matrix

$$\lambda_1 \cdot \lambda_2 \cdot \lambda_3 = |A| \quad \lambda_1 \cdot \lambda_2 \cdot \lambda_3 = |A|$$

Find the sum & product of the eigen values of  
matrix  $A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$

Let  $\lambda_1, \lambda_2, \lambda_3$  be the eigen values of A

$$\lambda_1 + \lambda_2 + \lambda_3 = \text{trace of } A$$

$$\lambda_1 + \lambda_2 + \lambda_3 = 8 + 7 + 3$$

02211051... Note Snowflake Downloads SP

MAXHUB Note5 Brazil-Bi

This PC Microsoft Office 2010 Telegram

Recycle Bin Snowflake Installati... New docum

Control Panel WhatsApp Image 2022... Zoom

Snowflake Installati... supremecon... 2022021

Google Chrome Acrobat Reader 2022030

Letter Writing Tips

Introduction to Enginee...

Topics 1 to (1)

SP

Window Tools Help

CASIO CLASSWIZ

Det(MatA) 0

7 8 9 DEL AC

4 5 6 X ÷

1 2 3 + -

0 .  $\times 10^x$  Ans

02211051...



Pro  $\rightarrow \lambda_1 + \lambda_2 + \lambda_3 = \text{trace of } A$

$$\lambda_1 + \lambda_2 + \lambda_3 = 8 + 7 + 3$$

$$\underline{\lambda_1 + \lambda_2 + \lambda_3 = 18}$$

Pro  $\Rightarrow \lambda_1 \cdot \lambda_2 \cdot \lambda_3 = |A|$

$$\underline{\underline{\lambda_1 \cdot \lambda_2 \cdot \lambda_3 = 0}}$$

16:23



If the product of 2 eigen values of  $\begin{bmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{bmatrix}$  is 6, Find the third eigen value

Let  $\lambda_1, \lambda_2, \lambda_3$  to be the eigen value

$$\lambda_1 \cdot \lambda_2 = 6$$

Pr

is 16, Find the third eigen value

$$\begin{bmatrix} 2 & 0 & 1 \\ 2 & -1 & 3 \end{bmatrix}$$

16:26

Let  $\lambda_1, \lambda_2, \lambda_3$  to be the eigen value

$$\lambda_1 \cdot \lambda_2 = 16$$

$$\Rightarrow \lambda_1 \lambda_2 \lambda_3 = |A|$$

$$16 \lambda_3 = 32$$

$$\lambda_3 = 2$$

Try:

If sum of the eigen values of  $3 \times 3$  matrix is 6 & product of the eigen value is also 6. If one of the eigen value is 1. Find the other eigen values

$\lambda_1, \lambda_2, \lambda_3$

Answer:

$\lambda_2, \lambda_3 = 1, 2, 3$  or  $1, 3, 2$



Try:

If sum of the eigen values of  $3 \times 3$  matrix is 6 & product of the eigen value is also 6. If one of the eigen value is 1. Find the other eigen values

Answer:

$$\begin{aligned} \lambda_1 \lambda_2 \lambda_3 &= 6 \\ \lambda_1 + \lambda_2 + \lambda_3 &= 6 \\ \lambda_2 \lambda_3 &= 6 \\ \lambda_2 &= 6/\lambda_3 \\ \lambda_3 &= 6/\lambda_2 \end{aligned}$$

$$\lambda_3 = 1, 2, 3 \text{ or } 1, 3, 2$$

$$1 + \lambda_2 + \lambda_3 = 6$$

## Properties of eigen values:

\* Eigen value of  $A$  &  $A'$  (transpose) are same

\* If  $\lambda_1, \lambda_2, \lambda_3$  are eigen values of  $A$  then eigen values of

①  $A^n \Rightarrow \lambda_1^n, \lambda_2^n, \lambda_3^n$

②  $KA \Rightarrow K\lambda_1, K\lambda_2, K\lambda_3$

③  $A^{-1} \Rightarrow \frac{1}{\lambda_1}, \frac{1}{\lambda_2}, \frac{1}{\lambda_3}$

④  $\text{adj}(A) \Rightarrow \frac{|A|}{\lambda_1}, \frac{|A|}{\lambda_2}, \frac{|A|}{\lambda_3}$

\* If  $\lambda_1, \lambda_2, \lambda_3$  are eigen values of  $A$  then eigen value

①  $A^n$   $\Rightarrow \lambda_1^n, \lambda_2^n, \lambda_3^n$

②  $kA$   $\Rightarrow k\lambda_1, k\lambda_2, k\lambda_3$

③  $A^{-1}$   $\Rightarrow \frac{1}{\lambda_1}, \frac{1}{\lambda_2}, \frac{1}{\lambda_3}$

$$A \quad 1 \quad 2 \quad 3$$

$$A^{-1} \quad \frac{1}{1}, \frac{1}{2}, \frac{1}{3}$$

④  $\text{adj}(A)$   $\Rightarrow \frac{|A|}{\lambda_1}, \frac{|A|}{\lambda_2}, \frac{|A|}{\lambda_3}$

Eigen value of identity matrix is 1, 1, 1.

$$(4) \underline{\text{adj}(A)} \Rightarrow \frac{|A|}{\lambda_1}, \frac{|A|}{\lambda_2}, \frac{|A|}{\lambda_3}$$

16:33

\* Eigen value of identity matrix is 1, 1, 1.

\* Eigen value of upper / Lower triangular matrix are diagonal elements.

17  $A = \begin{bmatrix} -1 & 2 & 3 \\ 0 & 3 & 5 \\ 0 & 0 & -2 \end{bmatrix}$  find eigen values of  $A^3 + 5A + 8I$

Let  $\lambda_1, \lambda_2, \lambda_3$

$$\lambda_1 = -1$$

$$\lambda_2 = 3$$

$$\lambda_3 = -2$$



$$\lambda_1 = -1$$

$$\lambda_2 = 3$$

$$\lambda_3 = -2$$

$$A^n$$

$$kA$$

$$8(I)$$

$$\Rightarrow A^n = \lambda_1^n$$

$$A^3 + 5A + 8I$$

Eigen values	$A^3$	$5A$	$8I$	
-1	$(-1)^3$	$5(-1)$	$8(1)$	
3	$(3)^3$	$5(3)$	$8(1)$	
-2	$(-2)^3$	$5(-2)$	$8(1)$	

$$\lambda_3 = -2$$

$$kA$$
  

$$8(1)$$

$$A^n \lambda_1^n$$

16:40

$$A^3 + 5A + 8I$$

Eigen values	$A^3$	$5A$	$8I$
-1	$(-1)^3 = -1$	$5(-1) = -5$	$8(1) = 8$
2	$(2)^3 = 8$	$5(2) = 10$	$8(1) = 8$
-2	$(-2)^3 = -8$	$5(-2) = -10$	$8(1) = 8$

$$\begin{aligned} & -6 + 8 \\ & \rightarrow -1 - 5 + 8 \end{aligned}$$

$$\rightarrow 8 + 10 + 8$$

$$\rightarrow -8 - 10 + 8$$

Eigen values of  $A^3 + 5A + 8I =$   
 $\lambda_1 =$

3	$(3)^3_{27}$	$5(3)_{15}$	$8(1)_8$	50	$\rightarrow 27+15+8$
-2	$(-2)^3_{-8}$	$5(-2)_{-10}$	$8(1)_8$	-10	$\rightarrow \cancel{-8} - 10 + \cancel{8}$

16:41

The eigen values of  $A^3 + 5A + 8I =$

$$\lambda_1' = 2$$

$$\lambda_2' = 50$$

$$\lambda_3' = -10$$

Try:

If  $A = \begin{bmatrix} 1 & 0 \\ 2 & 4 \end{bmatrix}$  then find the eigen values of  
 $4A^{-1} + 3A + 2I$

Answer: 9 & 15.