WARNINGS:

- Hematologic: This tablets should not be administered if the absolute neutrophil count is less than 500 cells/JJL, the platelet count is less than 25,000/JJL, or the hemoglobin is less than 8 g/dl.
- Impairment of fertility, teratogenis and mutagenesis and acute renal failure.

While receiving valganciclovir:


- Tell your doctor if you are taking any other medicines including herbal medicines and OTC medicines since sume medicines may interfere with Valganciclovir.
- Tell your doctor if you become pregnant while taking this medicine.
- Laboratory Testing: Due to the frequency of neutropenia, anemia and thrombocytopenia in patients receiving this tablets, it is recommended that CBC test be performed frequently.

Product Availability:

- Tablets: 450 mg
- Oral Solution: 50 mg/ml

STORAGE:

- Store the medicine at room temperature 25'C (77'F).
- Should be protected from moisture and kept away from direct sunlight.
- Do not leave it in the car or on window sills.



rela Hospital

DEPARTMENT OF CLINICAL PHARMACY

Rela Institute & Medical Centre

7, CLC Works Road, Chromepet, Chennai-600044

Contact

+91 44 6666 7788

www.relainstitute.com

Disclaimer: Not medical advice. For educational use only.

rela Hospital

DEPARTMENT OF CLINICAL PHARMACY


DR. RELA INSTITUTE & MEDICAL CENTRE

An International Medical Facility



VALGANCICLOVIR TABLET & SOLUTION

- This leaflet answers some common questions about VALGANCICLOVIR tablets/solution.
- It does not contain all the available information.
- It does not take the place of talking to your doctor or pharmacist.
- All medicines have risks and benefits, if you have any concerns about taking this medicine, ask your doctor or pharmacist.



What is Valganciclovir?

- It is an antiviral medicine which acts against a virus called cytomegalovirus or CMV by preventing it's growth and multification in the body.
- Valganciclovir is a pro drug which after oral administration, rapidly converted to Ganciclovir.

What is Valganciclovir used for?

Valganciclovir is used in the treatment and prevention of CMV Infections In high-risk individuals, those with poor immune function. It is used in,

- CMV retinitis (eye Infections)
- AIDS patients
- Prevention of CMV Infection in patients following organ transplantation.

When should I take the medication?

- This medicine should be administered only with a doctor's prescription.
- It is usually given 1 to 2 times daily and advisable to take the tablets at the same time each day.

How should it be taken?

- Valganciclovir should be taken with food or as soon.
- As after your food to improve absorption.
- If you are taking the oral suspension, shake well before use.
- Take valganciclovlr with plenty of water every day.
- Unless you are told to drink less liquid by your doctor.

What should I do if I miss a dose?

- If you forget a dose, take it as soon as you remember.
- If it ls nearer to the time of your next dose. skip the missed dose.
- Do not take two doses at the same time.
- Do not repeat dose If vomiting occurs after dose has been administered, wait until the next scheduled dose.
- Do not repeat dose If vomiting occurs after dose has been administered, wait until the next scheduled dose.

What are the possible side effects?

- Diarrhoea, Nausea, Vomiting, Abdominal pain, Neutropenia, Anaemia, Headache, Decreased appetite, Cough, Dyspnoea, fever and Fatigue,
- Tell your doctor if you notice anything else that is making you feel unwell or any unusual changes.

SPECIAL INSTRUCTIONS:

- It is contraindicated in patients with hypersensitivity to valganciclovir or Ganciclovir.
- Pregnant women - should avoid this medicine as it may harm the fetus. • Nursing mothers - should not breastfeed If they are receiving this medicine.
- Renal Impairment - Since ganciclovir is excreted by the kidneys, normal clearance depends on adequate renal function. Hence dosage adjustment is required according to creatinine clearance.
- It should not be taken by patients receiving hemodialysis.