What are Oral Hypoglycemic Agents (OHAs)?

They are group of medications which are given orally which to help reduce the blood sugar level primarily used in patients with type II Diabetes Mellitus.



What is Diabetes?

Diabetes is a chronic condition that occurs due to the body's inability to produce enough insulin or when the body loses sensitivity to the insulin produced.This in turn causes increased blood Sugar levels.

rela Hospital

DEPARTMENT OF CLINICAL PHARMACY

Rela Institute & Medical Centre

7, CLC Works Road, Chromepet, Chennai-600044

Contact

+91 44 6666 7788

www.relainstitute.com

Disclaimer: Not medical advice. For educational use only.

rela Hospital

DEPARTMENT OF CLINICAL PHARMACY

DR. RELA INSTITUTE & MEDICAL CENTRE

An International Medical Facility

A STEP TOWARDS A BETTER LIFE



Knowing more about Diabetes and Oral Hypoglycemic agents you consume everyday.

Recommended Schedule and Frequency

DRUGS	DIRECTION OF USE	HOW DO THEY ACT?	EFFECT ON WEIGHT	SIDE EFFECTS
Biguanides Metformin	To be taken after meals	Decreases glucose output from the liver	Weight loss	Abodominal discomfort, Gastric irritation
Sulphonylureas Gliclazide, Gliclazide Glimepiride	To be taken 30 minutes before food	Increases insulin secretion from the pancreas.	Weight gain	Hypoglycemia Hyperpigmentation of Skin
DPPA Inhibitors Vidagliptin , Teneligliptin	Can be taken before or after food.	Increases insulin secretion and decreases glucose secretion	Weight neutrat	Headache muscle and joint pain
SGLT2 Inhibitors Dapagkiflozin, Canaglifozin	Except Canagliflozin others can be taken before or after food	Increases urinary glucose creation	Weight loss	Increased urination urinary tract infection
Thaizolidnedione pioglitazone , Rosiglitazone	Can be taken before or after food.	Increases insulin sensitivity	Weight gain	Swelling of arms and feet Anemia
a Glucosidase inibitors Acarbose, Voglibose	To be taken before food	Delays glucose absorption	Small weight loss	Bloating Diarrhea Abdominal discomfort
Meglitinides Repaglinide Nateglinide	To be taken 30mins before food.	Increases inslin secretion from the	Weight gain	Hypogecemia