



Machine Learning


Linear Algebra
review (optional)

Matrices and
vectors

Matrix: Rectangular array of numbers:



$$\rightarrow \boxed{\mathbb{R}^{4 \times 2}}$$



$$\boxed{\mathbb{R}^{2 \times 3}}$$

Dimension of matrix: number of rows x number of columns

Matrix Elements (entries of matrix)

$$A = \begin{bmatrix} 1402 & 191 \\ 1371 & 821 \\ 949 & 1437 \\ 147 & 1448 \end{bmatrix}$$

A_{ij} = " i, j entry" in the i^{th} row, j^{th} column.

$$A_{11} = 1402$$

$$A_{12} = 191$$

$$A_{32} = 1437$$

$$A_{41} = 147$$

~~A_{43}~~ = Undefined (error)

Vector: An $n \times 1$ matrix.

$$\textcircled{y} = \begin{bmatrix} \textcircled{460} \\ \textcircled{232} \\ \textcircled{315} \\ 178 \end{bmatrix}$$

$n = 4$

← 4-dimensional vector.

~~$\mathbb{R}^{3 \times 2}$~~

\mathbb{R}^4

$y_i = i^{th}$ element

$$y_1 = 460$$

$$y_2 = 232$$

$$y_3 = 315$$

→ A, B, C, X

a, b, x, y

1-indexed vs 0-indexed:

$y[1]$

$$y = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix}$$

1-indexed

$y[0]$

$$y = \begin{bmatrix} y_0 \\ y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

0-indexed



Machine Learning

Linear Algebra review (optional)

Addition and scalar multiplication

Matrix Addition

$$\begin{array}{c} \downarrow \quad \downarrow \\ \rightarrow \begin{bmatrix} \textcircled{1} & 0 \\ \textcircled{2} & 5 \\ \textcircled{3} & 1 \end{bmatrix} + \begin{bmatrix} \textcircled{4} & 0.5 \\ \textcircled{2} & 5 \\ \textcircled{0} & 1 \end{bmatrix} = \begin{bmatrix} 5 & 0.5 \\ 4 & 10 \\ 3 & 2 \end{bmatrix} \\ \text{3} \times \text{2} \quad \text{3} \times \text{2} \quad \text{3} \times \text{2} \end{array}$$

$$\begin{array}{c} \rightarrow \begin{bmatrix} 1 & 0 \\ 2 & 5 \\ 3 & 1 \end{bmatrix} + \begin{bmatrix} 4 & 0.5 \\ 2 & 5 \end{bmatrix} = \\ \text{3} \times \text{2} \quad \text{2} \times \text{2} \end{array}$$

error

Scalar Multiplication

← real number

$$3 \times \begin{bmatrix} 1 & 0 \\ 2 & 5 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 6 & 15 \\ 9 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 2 & 5 \\ 3 & 1 \end{bmatrix} \times 3$$

3x2 3x2

$$\begin{bmatrix} 4 & 0 \\ 6 & 3 \end{bmatrix} / 4 = \frac{1}{4} \begin{bmatrix} 4 & 0 \\ 6 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ \frac{3}{2} & \frac{3}{4} \end{bmatrix}$$

Combination of Operands

$$\begin{aligned}
 & \text{Scalar multiplication} \rightarrow 3 \times \begin{bmatrix} 1 \\ 4 \\ 2 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 5 \end{bmatrix} - \begin{bmatrix} 3 \\ 0 \\ 2 \end{bmatrix} \text{ / 3} \quad \text{Scalar division} \\
 & = \begin{bmatrix} 3 \\ 12 \\ 6 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 5 \end{bmatrix} - \begin{bmatrix} 1 \\ 0 \\ \frac{2}{3} \end{bmatrix} \quad \begin{array}{l} \text{matrix subtraction /} \\ \text{vector subtraction} \end{array} \\
 & = \begin{bmatrix} 2 \\ 12 \\ 10 \frac{1}{3} \end{bmatrix} \quad \begin{array}{l} \text{matrix addition /} \\ \text{vector addition} \end{array} \\
 & \quad \begin{array}{l} 3 \times 1 \text{ matrix} \\ 3\text{-dimensional vector} \end{array}
 \end{aligned}$$



Machine Learning

Linear Algebra review (optional)

Matrix-vector multiplication

Example

$$\begin{bmatrix} 1 & 3 \\ 4 & 0 \\ 2 & 1 \end{bmatrix}_{3 \times 2} \begin{bmatrix} 1 \\ 5 \end{bmatrix}_{2 \times 1} = \begin{bmatrix} 16 \\ 4 \\ 7 \end{bmatrix}_{3 \times 1} \text{ matrix}$$

$$1 \times 1 + 3 \times 5 = 16$$

$$4 \times 1 + 0 \times 5 = 4$$

$$2 \times 1 + 1 \times 5 = 7$$

Details:

$$\underline{A} \times \underline{x} = \underline{y}$$

\underline{A} is an $m \times n$ matrix (m rows, n columns).
 \underline{x} is an $n \times 1$ matrix (n-dimensional vector).
 \underline{y} is an m -dimensional vector.

→ To get y_i , multiply \underline{A} 's i^{th} row with elements of vector \underline{x} , and add them up.

Example

$$\begin{bmatrix} 1 & 2 & 1 & 5 \\ 0 & 3 & 0 & 4 \\ -1 & -2 & 0 & 0 \end{bmatrix}_{3 \times 4} \begin{matrix} \downarrow \\ \begin{bmatrix} 1 \\ 3 \\ 2 \\ 1 \end{bmatrix}_{4 \times 1} \end{matrix} = \begin{bmatrix} 14 \\ 13 \\ -7 \end{bmatrix}_{3 \times 1} = \begin{bmatrix} 14 \\ 13 \\ -7 \end{bmatrix}$$

$$\left. \begin{array}{l} 1 \times 1 + 2 \times 3 + 1 \times 2 + 5 \times 1 = 14 \\ 0 \times 1 + 3 \times 3 + 0 \times 2 + 4 \times 1 = 13 \\ -1 \times 1 + (-2) \times 3 + 0 \times 2 + 0 \times 1 = -7 \end{array} \right\}$$

House sizes:

→ 2104

→ 1416

→ 1534

→ 852

Matrix

4x2

1	2104
1	1416
1	1534
1	852

$$h_{\theta}(x) = -40 + 0.25x$$

$h_{\theta}(x)$

2x1

Vector

X

$$\begin{bmatrix} -40 \\ 0.25 \end{bmatrix}$$

=

4x1 matrix

$-40 \times 1 + 0.25 \times 2104$
$-40 \times 1 + 0.25 \times 1416$

$h_{\theta}(1416)$

Prediction = Data Matrix * Parameters

4x1

for $i = 1:1000$,
prediction(i) = ...



Machine Learning

Linear Algebra review (optional)

Matrix-matrix multiplication

Example

$$\begin{array}{l} \begin{bmatrix} 1 & 3 & 2 \\ 4 & 0 & 1 \end{bmatrix} \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 0 & 1 \\ \hline 5 & 2 \\ \hline \end{array} = \begin{bmatrix} 11 & 10 \\ 9 & 14 \end{bmatrix} \\ \textcircled{2 \times 3} \quad \textcircled{3 \times 2} \\ \begin{bmatrix} 1 & 3 & 2 \\ 4 & 0 & 1 \end{bmatrix} \times \begin{array}{|c|} \hline 1 \\ \hline 0 \\ \hline 5 \\ \hline \end{array} = \begin{bmatrix} 11 \\ 9 \end{bmatrix} \\ \begin{bmatrix} 1 & 3 & 2 \\ 4 & 0 & 1 \end{bmatrix} \times \begin{array}{|c|} \hline 3 \\ \hline 1 \\ \hline 2 \\ \hline \end{array} = \begin{bmatrix} 10 \\ 14 \end{bmatrix} \end{array}$$

Handwritten green annotations: The first row of the first matrix is underlined. In the second equation, the first matrix is underlined. In the third equation, the first matrix is underlined. In the second equation, the second matrix is underlined. In the third equation, the second matrix is underlined. Arrows point from the 2x2 matrix to the 2x1 and 2x1 matrices, indicating that the 2x2 matrix is the result of the 2x3 matrix multiplied by the 3x2 matrix.

Details:

$$\underline{A} \times \underline{B} = \underline{C}$$

$m \times n$ matrix
(m rows,
 n columns)

$n \times o$ matrix
(n rows,
 o columns)

$m \times o$
matrix

The i^{th} column of the matrix C is obtained by multiplying A with the i^{th} column of B . (for $i = 1, 2, \dots, o$)

Example

$$\overset{2 \times 2}{\begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix}} \overset{2 \times 2}{\begin{bmatrix} 0 & 1 \\ 3 & 2 \end{bmatrix}} =$$

$$\overset{2 \times 2}{\begin{bmatrix} 9 & 7 \\ 15 & 12 \end{bmatrix}}$$

$$\begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} 0 \\ 3 \end{bmatrix} =$$

$$= \begin{bmatrix} 1 \times 0 + 3 \times 3 \\ 2 \times 0 + 5 \times 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 15 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} =$$

$$= \begin{bmatrix} 1 \times 1 + 3 \times 2 \\ 2 \times 1 + 5 \times 2 \end{bmatrix} = \begin{bmatrix} 7 \\ 12 \end{bmatrix}$$

House sizes:

$$\begin{Bmatrix} 2104 \\ 1416 \\ 1534 \\ 852 \end{Bmatrix}$$

Matrix

$$\begin{bmatrix} 1 & 2104 \\ 1 & 1416 \\ 1 & 1534 \\ 1 & 852 \end{bmatrix} \times$$

Matrix

$$\begin{bmatrix} -40 \\ 0.25 \end{bmatrix} \begin{bmatrix} 200 \\ 0.1 \end{bmatrix} \begin{bmatrix} -150 \\ 0.4 \end{bmatrix} =$$

Have 3 competing hypotheses:

$$\begin{aligned} 1. & h_{\theta}(x) = -40 + 0.25x \\ 2. & h_{\theta}(x) = 200 + 0.1x \\ 3. & h_{\theta}(x) = -150 + 0.4x \end{aligned}$$

$$\begin{bmatrix} 486 \\ 314 \\ 344 \\ 173 \end{bmatrix} \begin{bmatrix} 410 \\ 342 \\ 353 \\ 285 \end{bmatrix} \begin{bmatrix} 692 \\ 416 \\ 464 \\ 191 \end{bmatrix}$$

Prediction
of first
 h_{θ}

Predictions
of 2nd
 h_{θ}



Machine Learning

Linear Algebra review (optional)

Matrix multiplication properties

$$3 \times 5 = 5 \times 3$$


"Commutative"

Let A and B be matrices. Then in general,
 $A \times B \neq B \times A$. (not commutative.)

E.g.


$$\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 2 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix}$$


$$\begin{bmatrix} 0 & 0 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 2 & 2 \end{bmatrix}$$


$A \times B$
 $m \times n \quad n \times m$

$A \times B$ is $m \times m$

$B \times A$ is $n \times n$



$$\underline{3 \times 5 \times 2}$$

$$3 \times 10 = 30 = 15 \times 2$$

$$3 \times (5 \times 2) = (3 \times 5) \times 2$$

"Associative"

$$\begin{array}{l} A \times (B \times C) \leftarrow \\ \underline{(A \times B)} \times C \leftarrow \end{array}$$

$$A \times B \times C.$$

Let $D = B \times C$. Compute $A \times D$.

Let $E = A \times B$. Compute $E \times C$.

$A \times (B \times C)$
 $(A \times B) \times C$
 Some
 answer.

1 is identity

$$1 \times z = z \times 1 = z$$

for any z

Identity Matrix

Denoted I (or $I_{n \times n}$).

Examples of identity matrices:

$[1]$
 1×1

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

2×2

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

3×3

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

4×4

Informally:

$$\begin{bmatrix} 1 & & & \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{bmatrix}$$

For any matrix A ,

$$A \cdot I = I \cdot A = A$$

$m \times n$ $n \times n$ $m \times m$ $m \times n$ $m \times n$

$I_{n \times n}$

Note:

$AB \neq BA$ in general

$$AI = IA \checkmark$$



Machine Learning

Linear Algebra review (optional)

Inverse and transpose

$$\underline{1 = \text{"identity"}}$$

$$3 \underbrace{(3^{-1})}_{\frac{1}{3}} = 1$$

$$12 \times \underbrace{(12^{-1})}_{\frac{1}{12}} = 1$$

$$0 \underbrace{(0^{-1})}_{\text{undefined}}$$

Not all numbers have an inverse.

Matrix inverse: \swarrow square matrix
(#rows = #columns) A^{-1}

If A is an $m \times m$ matrix, and if it has an inverse,

$$\rightarrow \underline{A(A^{-1})} = \underline{A^{-1}A} = \underline{I}.$$

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \swarrow$$

e.g. $\underbrace{\begin{bmatrix} 3 & 4 \\ 2 & 16 \end{bmatrix}}_A \underbrace{\begin{bmatrix} 0.4 & -0.1 \\ -0.05 & 0.075 \end{bmatrix}}_{A^{-1}} = \underbrace{\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}}_{A^{-1}A} = I_{2 \times 2}$

Matrices that don't have an inverse are "singular" or "degenerate"

Matrix Transpose

Example:

$$\underline{A} = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 5 & 9 \end{bmatrix}_{2 \times 3}$$
$$\underline{B} = \underline{A}^T = \begin{bmatrix} 1 & 3 \\ 2 & 5 \\ 0 & 9 \end{bmatrix}_{3 \times 2}$$

Let A be an $m \times n$ matrix, and let $B = A^T$.

Then B is an $n \times m$ matrix, and

$$\underline{B}_{ij} = \underline{A}_{ji}.$$

$$B_{12} = A_{21} = 2$$

$$B_{32} = 9$$

$$A_{23} = 9.$$