

Rust in Pulses

Uredo cajani Thurunoi / பரம்பா வகைஞ்சுரா

	Prevention	Monitoring	Direct Control	Direct Control	Restrictions
 Rust pustules on leaves (Dongxin Feng, ecoport.org)	- Plough the soil twice before sowing the crop to kill the inoculum. - Remove the stubble after harvesting and destroy by burning to reduce the primary and secondary inoculums. - Sow/plant the seeds with 60x60cm spacing. - Avoid excess usage of nitrogenous fertilizers (recommended dosage of nitrogen fertilizer is 12.25 kg). - Maintain soil moisture with a proper irrigation schedule. - Irrigate immediately after sowing and light irrigation on the third day. Irrigate at intervals of 7 to 10 days depending upon soil and climatic conditions.	- Disease infestation occurs at the vegetative stage. - Observe leaves for red/brown raised pustules under a microscope or hand lens. - Disease incidence is severe during Jan 15th to Feb 15th due to heavy wind. - Red gram and Blackgram: Dark brown rusty pustules (uredia) appear on the lower surface of the leaves. - Bengal gram and Cowpea: Dark brown rusty pustules (uredia) appear on the upper surface of the leaves. - In severe conditions observe for desiccated leaves and leaf drop at seed setting stage or pod formation stage.	Treat the seeds with <i>Pseudomonas fluorescence</i> at 10 gm/kg of seeds and shade dry for 15 minutes before sowing.	- Spray of wettable Sulphur at 1500 g/Ha at 10 day intervals. - Spray the crop with 0.2% Mancozeb 75 WP followed by two more sprays at 10 day intervals.	- WHO Class III (Slightly Hazardous). REI: 24 hrs. - WHO class U (Unlikely to present acute hazard in normal use).
 Severe stage of rust in pulses (iNaturalist.org)					

India

CREATED/UPDATED: Dec. 2017

AUTHOR(S): Kanchana, Rajasekar, T. Sunil Kumar, D. Adhikari. NAF Phone no. 9790707020, Email: sriramamills@gmail.com

EDITED BY: Plantwise