

Theoretical Computer Science

Tutorial Week 5

Prof. Andrey Frolov



- Myhill-Nerode criteria (Negative examples)
- Pumping Lemma
- Pushdown automata

Myhill-Nerode criteria

For a language L over an alphabet A ,

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \ \& \ s_2 t \in L) \vee (s_1 t \in L \ \& \ s_2 t \notin L)]$$

Myhill-Nerode theorem

A language L is regular iff \equiv_L has a finite number of equivalent classes.

Myhill-Nerode method. Examples

Negative Example

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

$$L_{01} = \{\epsilon, 01, 0011, 000111, \dots\}$$

Proof

For $m \neq k$,

$$0^m \not\equiv_{L_{01}} 0^k,$$

since $0^m 1^k \notin L_{01}$, $0^k 1^k \in L_{01}$ (a distinguishing ext. is 1^k).

Therefore, there are **infinity many equivalence classes**!

So, L_{01} is not regular.

Agenda

- Myhill-Nerode criteria (Negative examples)
- **Pumping Lemma**
- Pushdown automata

Pumping Lemma

Negative Example

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

$$L_{01} = \{\epsilon, 01, 0011, 000111, \dots\}$$

Suppose that a FSA with k states recognize L_{01} .

$$\underbrace{\epsilon, 0, 00, \dots, 0^k}_{k+1}$$

$$\underbrace{q_0, q_{i_1}, q_{i_2}, \dots, q_{i_k}}_{k+1}$$

Pumping Lemma

Suppose that a FSA with k states recognizes L_{01} .

$$\underbrace{\epsilon, 0, 00, \dots, 0^k}_{k+1}$$

$$\underbrace{q_0, q_{i_1}, q_{i_2}, \dots, q_{i_k}}_{k+1}$$

$$q_{i_s} \xrightarrow{0} q_{i_{s+1}} \xrightarrow{0} q_{i_{s+2}} \xrightarrow{0} \dots \xrightarrow{0} q_{i_s}$$

Pumping Lemma

Suppose that a FSA with k states recognizes L_{01} .


$$\underbrace{\epsilon, 0, 00, \dots, 0^k}_{k+1}$$

$$\underbrace{q_0, q_{i_1}, q_{i_2}, \dots, q_{i_k}}_{k+1}$$

$$q_{i_s} \xrightarrow{0} q_{i_{s+1}} \xrightarrow{0} q_{i_{s+2}} \xrightarrow{0} \dots \xrightarrow{0} q_{i_s}$$

Therefore, if the FSA recognizes $0^n 1^n$ then it recognizes also $0^{n+t} 1^n$ for some $t > 0$. It's a contradiction.

Pumping lemma



Pumping lemma

If $L \subseteq \Sigma^*$ is a regular language then there exists $m \geq 1$ such that any $w \in L$ with $|w| \geq m$ can be represented as $w = xyz$ such that

- $y \neq \epsilon$,
- $|xy| \leq m$,
- $xy^iz \in L$ for any $i \geq 0$.

How Pumping lemma is useful?

- Can we use this theorem to prove that a set is regular?

How Pumping lemma is useful?

- Can we use this theorem to prove that a set is regular?
No, because it gives only a necessary condition for a language to be regular (and not a sufficient condition).

How Pumping lemma is useful?

- Can we use this theorem to prove that a set is regular?
No, because it gives only a necessary condition for a language to be regular (and not a sufficient condition).
- We can use it to prove that a language is not regular.
How?

Pumping lemma

Pumping lemma

If $L \subseteq \Sigma^*$ is a regular language then there exists $m \geq 1$ such that any $w \in L$ with $|w| \geq m$ can be represented as $w = xyz$ such that

- $y \neq \epsilon$ and $|xy| \leq m$,
- $xy^iz \in L$ for any $i \geq 0$.

Corollary (Contrapositive)

If for any $m \geq 1$ there is $w \in L$ such that $|w| \geq m$ and for any representation $w = xyz$ with $y \neq \epsilon$ and $|xy| \leq m$

$$xy^iz \notin L \text{ for some } i \geq 0.$$

Then L is **not** a regular language.

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

For $0^n 1^n = xyz \in L_{01}$ with $y \neq \epsilon$, we have 3 cases:

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

For $0^n 1^n = xyz \in L_{01}$ with $y \neq \epsilon$, we have 3 cases:

1) $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

For $0^n 1^n = xyz \in L_{01}$ with $y \neq \epsilon$, we have 3 cases:

- 1) $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$
- 2) $\underbrace{0^n 1^{p_1}}_x \underbrace{(1^{p_2})}_y \underbrace{1^{n-p_1-p_2}}_z$ and $p_2 \neq 0$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

For $0^n 1^n = xyz \in L_{01}$ with $y \neq \epsilon$, we have 3 cases:

- 1) $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$
- 2) $\underbrace{0^n 1^{p_1}}_x \underbrace{(1^{p_2})}_y \underbrace{1^{n-p_1-p_2}}_z$ and $p_2 \neq 0$
- 3) $\underbrace{0^{n-p_1}}_x \underbrace{(0^{p_1} 1^{p_2})}_y \underbrace{1^{n-p_2}}_z$ and $p_1, p_2 \neq 0$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

1) for $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$, we have
 $xy^2z =$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

1) for $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$, we have

$$xy^2z = 0^{n-p_1-p_2}(0^{p_1})^2 0^{p_2} 1^n$$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

1) for $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$, we have

$$xy^2z = 0^{n-p_1-p_2}(0^{p_1})^2 0^{p_2} 1^n = 0^{n+p_1} 1^n$$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

1) for $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$, we have

$$xy^2z = 0^{n-p_1-p_2}(0^{p_1})^2 0^{p_2} 1^n = 0^{n+p_1} 1^n \notin L,$$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

1) for $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$, we have

$$xy^2z = 0^{n-p_1-p_2}(0^{p_1})^2 0^{p_2} 1^n = 0^{n+p_1} 1^n \notin L,$$

2) for $\underbrace{0^n 1^{p_1}}_x \underbrace{(1^{p_2})}_y \underbrace{1^{n-p_1-p_2}}_z$ and $p_2 \neq 0$, we have

$$xy^2z =$$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

1) for $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$, we have

$$xy^2z = 0^{n-p_1-p_2}(0^{p_1})^2 0^{p_2} 1^n = 0^{n+p_1} 1^n \notin L,$$

2) for $\underbrace{0^n 1^{p_1}}_x \underbrace{(1^{p_2})}_y \underbrace{1^{n-p_1-p_2}}_z$ and $p_2 \neq 0$, we have

$$xy^2z = 0^n 1^{n+p_2} \notin L,$$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

1) for $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$, we have

$$xy^2z = 0^{n-p_1-p_2}(0^{p_1})^2 0^{p_2} 1^n = 0^{n+p_1} 1^n \notin L,$$

2) for $\underbrace{0^n 1^{p_1}}_x \underbrace{(1^{p_2})}_y \underbrace{1^{n-p_1-p_2}}_z$ and $p_2 \neq 0$, we have

$$xy^2z = 0^n 1^{n+p_2} \notin L,$$

3) for $\underbrace{0^{n-p_1}}_x \underbrace{(0^{p_1} 1^{p_2})}_y \underbrace{1^{n-p_2}}_z$ and $p_1, p_2 \neq 0$, we have $xy^2z =$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

1) for $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$, we have

$$xy^2z = 0^{n-p_1-p_2}(0^{p_1})^2 0^{p_2} 1^n = 0^{n+p_1} 1^n \notin L,$$

2) for $\underbrace{0^n 1^{p_1}}_x \underbrace{(1^{p_2})}_y \underbrace{1^{n-p_1-p_2}}_z$ and $p_2 \neq 0$, we have

$$xy^2z = 0^n 1^{n+p_2} \notin L,$$

3) for $\underbrace{0^{n-p_1}}_x \underbrace{(0^{p_1} 1^{p_2})}_y \underbrace{1^{n-p_2}}_z$ and $p_1, p_2 \neq 0$, we have $xy^2z =$
 $0^{n-p_1} (0^{p_1} 1^{p_2})^2 1^{n-p_2} =$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

1) for $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$, we have

$$xy^2z = 0^{n-p_1-p_2}(0^{p_1})^2 0^{p_2} 1^n = 0^{n+p_1} 1^n \notin L,$$

2) for $\underbrace{0^n 1^{p_1}}_x \underbrace{(1^{p_2})}_y \underbrace{1^{n-p_1-p_2}}_z$ and $p_2 \neq 0$, we have

$$xy^2z = 0^n 1^{n+p_2} \notin L,$$

3) for $\underbrace{0^{n-p_1}}_x \underbrace{(0^{p_1} 1^{p_2})}_y \underbrace{1^{n-p_2}}_z$ and $p_1, p_2 \neq 0$, we have $xy^2z =$

$$0^{n-p_1} (0^{p_1} 1^{p_2})^2 1^{n-p_2} = 0^{n-p_1} 0^{p_1} 1^{p_2} 0^{p_1} 1^{p_2} 1^{n-p_2} =$$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

1) for $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$, we have

$$xy^2z = 0^{n-p_1-p_2}(0^{p_1})^2 0^{p_2} 1^n = 0^{n+p_1} 1^n \notin L,$$

2) for $\underbrace{0^n 1^{p_1}}_x \underbrace{(1^{p_2})}_y \underbrace{1^{n-p_1-p_2}}_z$ and $p_2 \neq 0$, we have

$$xy^2z = 0^n 1^{n+p_2} \notin L,$$

3) for $\underbrace{0^{n-p_1}}_x \underbrace{(0^{p_1} 1^{p_2})}_y \underbrace{1^{n-p_2}}_z$ and $p_1, p_2 \neq 0$, we have $xy^2z =$

$$0^{n-p_1} (0^{p_1} 1^{p_2})^2 1^{n-p_2} = 0^{n-p_1} 0^{p_1} 1^{p_2} 0^{p_1} 1^{p_2} 1^{n-p_2} = 0^n 1^{p_2} 0^{p_1} 1^n$$

Pumping lemma. Examples

Example 1

$L_{01} = \{0^n 1^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

1) for $\underbrace{0^{n-p_1-p_2}}_x \underbrace{(0^{p_1})}_y \underbrace{0^{p_2} 1^n}_z$ and $p_1 \neq 0$, we have

$$xy^2z = 0^{n-p_1-p_2}(0^{p_1})^2 0^{p_2} 1^n = 0^{n+p_1} 1^n \notin L,$$

2) for $\underbrace{0^n 1^{p_1}}_x \underbrace{(1^{p_2})}_y \underbrace{1^{n-p_1-p_2}}_z$ and $p_2 \neq 0$, we have

$$xy^2z = 0^n 1^{n+p_2} \notin L,$$

3) for $\underbrace{0^{n-p_1}}_x \underbrace{(0^{p_1} 1^{p_2})}_y \underbrace{1^{n-p_2}}_z$ and $p_1, p_2 \neq 0$, we have $xy^2z =$

$$0^{n-p_1} (0^{p_1} 1^{p_2})^2 1^{n-p_2} = 0^{n-p_1} 0^{p_1} 1^{p_2} 0^{p_1} 1^{p_2} 1^{n-p_2} = 0^n 1^{p_2} 0^{p_1} 1^n \notin L$$

Pumping lemma

Pumping lemma

If $L \subseteq \Sigma^*$ is a regular language then there exists $m \geq 1$ such that any $w \in L$ with $|w| \geq m$ can be represented as $w = xyz$ such that

- $y \neq \epsilon$ and $|xy| \leq m$,
- $xy^iz \in L$ for any $i \geq 0$.

Corollary (Contrapositive)

If for any $m \geq 1$ there is $w \in L$ such that $|w| \geq m$ and for any representation $w = xyz$ with $y \neq \epsilon$ and $|xy| \leq m$

$$xy^iz \notin L \text{ for some } i \geq 0.$$

Then L is **not** a regular language.

Pumping lemma. Examples

Example 2

$L_2 = \{a^n ba^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

Let $w = xyz \in L_2$ and $y \neq \epsilon$

Pumping lemma. Examples

Example 2

$L_2 = \{a^n b a^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

Let $w = xyz \in L_2$ and $y \neq \epsilon$

1) if y contains b (that is $w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1} b a^{p_2}}_y \underbrace{a^{n-p_2}}_z$)

Pumping lemma. Examples

Example 2

$L_2 = \{a^n b a^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

Let $w = xyz \in L_2$ and $y \neq \epsilon$

1) if y contains b (that is $w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1} b a^{p_2}}_y \underbrace{a^{n-p_2}}_z$)

then xy^2z contains b twice and hence $xy^2z \notin L_2$

Pumping lemma. Examples

Example 2

$L_2 = \{a^n b a^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

Let $w = xyz \in L_2$ and $y \neq \epsilon$

1) if y contains b (that is $w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1} b a^{p_2}}_y \underbrace{a^{n-p_2}}_z$)

then xy^2z contains b twice and hence $xy^2z \notin L_2$

2) if y does not contain b , that is

$$w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1}}_y \underbrace{b a^n}_z, \text{ or } w = \underbrace{a^n}_x \underbrace{b}_{y_1} \underbrace{a^{p_2}}_y \underbrace{a^{n-p_2}}_z$$

Pumping lemma. Examples

Example 2

$L_2 = \{a^n ba^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

Let $w = xyz \in L_2$ and $y \neq \epsilon$

1) if y contains b (that is $w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1} ba^{p_2}}_y \underbrace{a^{n-p_2}}_z$)

then xy^2z contains b twice and hence $xy^2z \notin L_2$

2) if y does not contain b , that is

$$w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1}}_y \underbrace{ba^n}_z, \text{ or } w = \underbrace{a^n}_x \underbrace{b}_y \underbrace{a^{p_2}}_y \underbrace{a^{n-p_2}}_z$$

then $xy^2z = a^k ba^m$ where $k \neq m$ and hence $xy^2z \notin L_2$

Pumping lemma. For practice

Pumping lemma

If $L \subseteq \Sigma^*$ is a regular language then there exists $m \geq 1$ such that any $w \in L$ with $|w| \geq m$ can be represented as $w = xyz$ such that

- $y \neq \epsilon$,
- $|xy| \leq m$,
- $xy^iz \in L$ for any $i \geq 0$.

Corollary

If for any $m \geq 1$ there is $w \in L$ such that $|w| \geq m$ and for any representation $w = xyz$ with $y \neq \epsilon$ and $|xy| \leq m$

$$xy^iz \notin L \text{ for some } i \geq 0.$$

Then L is **not** a regular language.

Pumping lemma. For practice

Corollary

If for any $m \geq 1$ there is $w \in L$ such that $|w| \geq m$ and for any representation $w = xyz$ with $y \neq \epsilon$ and $|xy| \leq m$

$$xy^iz \notin L \text{ for some } i \geq 0.$$

Then L is **not** a regular language.

Negative Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular.

Proof

For each m , let $w = a^m b^m$. If $w = xyz$ and $|xy| \leq m$, then $w = \underbrace{a^{m-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^m}_z$ and hence $xy^2z \notin L_1$

Pumping lemma. For practice

Corollary

If for any $m \geq 1$ there is $w \in L$ such that $|w| \geq m$ and for any representation $w = xyz$ with $y \neq \epsilon$ and $|xy| \leq m$

$$xy^iz \notin L \text{ for some } i \geq 0.$$

Then L is **not** a regular language.

Negative Example 2

$L_2 = \{a^n ba^n \mid n \in \mathbb{N}\}$ is not regular.


Proof

For each m , let $w = a^m ba^m$. If $w = xyz$ and $|xy| \leq m$, then $w = \underbrace{a^{m-p_1}}_x \underbrace{a^{p_1}}_y \underbrace{ba^m}_z$ and hence $xy^2z \notin L_2$

Agenda

- Myhill-Nerode criteria (Negative examples)
- Pumping Lemma
- **Pushdown automata**


FSA




Pushdown Automata (PDA)




Finite State Automata



Pushdown Automata




Pushdown automata




$\Gamma = \{Z_0, A\}$, Z_0 is the initial stack symbol.

Pushdown automata




*a*abb

Pushdown automata




$aabb$

Pushdown automata




$aa**b**b$

Pushdown automata



*aab***b**

Pushdown automata




$aabb \in L_1$

PDA. Examples


Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular, but is **recognized by a PDA**.



Example 2

$L_2 = \{a^n b a^n \mid n \in \mathbb{N}\}$ is not regular, but is **recognized by a PDA**.



Thank you for your attention!