

داخلي

1- Which of the following muscle has dual nerve supply:

- ✓ Pectineus
- Sartorius
- Adductor longus
- Rectus femoris

2- Which of the following does not pass through the greater sciatic foramen?

- ✓ the nerve to obturator externus
- the pudendal nerve and vessels
- the posterior cutaneous nerve of the thigh
- the sciatic nerve

3- All the following are branches of the femoral nerve except:

- saphenous nerve
- nerve to vastus medialis
- ✓ lateral femoral cutaneous nerve
- medial femoral cutaneous nerve

4- Which one of the following muscle is inserted into adductor tubercle of femur:

- Adductor longus
- Adductor brevis
- ✓ Adductor magnus
- Biceps femoris

5- The following muscle is NOT responsible for hip flexion:

- iliopsoas
- . Sartorius
- ✓ . vastus lateralis
- . rectus femoris

6- The 1st toe cleft is supplied by

- Saphenous n
- Sural n
- Superficial peroneal n
- ✓ Deep peroneal n

7- What type of joint is hip joint:

- ✓ Ball & socket joint
- Pivot joint
- Ellipsoid joint
- Hinge joint

8- Which part of quadriceps femoris has attachment to the hip bone?

vastus medialis

vastus lateralis

✓ . rectus femoris

vastus intermedius

9- A 45-year-old man has trouble walking. At his physician's clinic, when he is asked to stand on his right foot, his left hip drops. Which of the following nerves is most likely damaged, causing his problem:

✓ Right superior gluteal

Left superior gluteal

Right inferior gluteal

Left inferior gluteal

10-

The lateral sural nerve is a branch of which of the following nerves?

tibial

femoral

✓ common fibular

obturator

11- Contents of the femoral sheath include:

Superficial inguinal lymph nodes

Femoral n

✓ Femoral a

Profunda femoris a

12- Profunda femoral artery is normally the vessel of supply:

To the anterior muscles of thigh

To the lateral muscles of thigh

To the medial muscles of thigh

✓ To the posterior muscles of thigh

13- At the subtalar/talocalcaneal joint the primary movements are

✓ Inversion and eversion

Inversion and adduction

Flexion and extension

Eversion and abduction

14- Which is not a component of the 2nd layer of the foot

Tendon of FHL

The lumbricals

✓ Adductor hallucis

Flexor accessorius

15- Obturator nerve is divided into anterior & posterior divisions by:

- . adductor brevis
- . obturator internus
- . medial circumflex femoral artery
- ✓ . obturator externus

16- Which of the following muscles is the strongest dorsiflexor and invertor of the foot?

- ✓ tibialis anterior
- extensor digitorum longus
- extensor hallucis longus
- fibularis tertius

17- The most medial structure passing under the inferior extensor retinaculum of the foot is:

- extensor hallucis longus
- deep peroneal nerve
- anterior tibial artery
- ✓ tibialis anterior

18- Which of the following is not a border for TIBIA:

- Lateral
- Anterior
- Medial
- ✓ Posterior

19- A 20-year-old college student receives a severe blow on the inferolateral side of the left knee joint while playing football. Radiographic examination reveals a fracture of the head and neck of the fibula. Which of the following nerves is damaged:

- Tibial
- Deep peroneal
- Superficial peroneal
- ✓ Common peroneal

20- Medial plantar nerve does NOT supply:

- . flexor hallucis brevis
- ✓ . adductor hallucis
- . abductor hallucis
- . flexor digitorum brevis

21- All of the following attach to the greater trochanter except

- Gluteus medius
- Piriformis
- ✓ Iliacus
- Obturator internus

22- Which structure is intrasynovial at the knee joint:

- anterior cruciate ligament
- ✓ tendon of popliteus
- medial and lateral menisci
- oblique popliteal ligament

23- Floor of the femoral triangle EXCLUDES:

- ✓ adductor brevis
- . psoas major
- iliacus
- pectineus

24- Difficulty extending the knee can result from damage to the

- inferior gluteal nerve
- tibial nerve
- common peroneal nerve
- ✓ femoral nerve

25- Which of the following is NOT a content of the adductor canal?

- ☐ femoral artery
- ☒ femoral nerve
- ☐ femoral vein
- ☐ nerve to vastus medialis

26- The muscle that unlocks the knee joint is

- ☒ Popliteus
- ☐ Gastrocnemius
- ☐ Plantaris
- ☐ Quadriceps femoris

27- Which of the following is found superficially immediately anterior to the medial malleolus?

- ☐ Small saphenous vein
- ☐ Arcuate artery
- ☒ Greater saphenous vein
- ☐ Dorsalis pedis artery

28- Iliotibial tract is inserted on:

- ☒ Lateral condyl of tibia
- ☐ Upper smooth part of tibial tuberosity
- ☐ Medial condyl of tibia
- ☐ Lower roughened part of tibial tuberosity

29- The femoral nerve is formed by the posterior division of the anterior rami of:

L4, L5

L1, L2, L3

L2, L3, L4

✓ L3, L4, L5

30- Posterior tibial artery

Divides superior to the flexor retinaculum

Branches form 2 plantar arches in the foot

✓ Divides into the medial and lateral plantar branches

Lateral palmar branch supplies the big toe