

1. Regarding the clavicle, all are correct EXCEPT

- ☐ It is the first bone to ossify in the fetal life.
- ☐ It is the commonly fractured bone
- ☐ It is an example of the short bones.
- ☐ It ossifies in membrane.

3. It performs its function in the unlocking of the knee joint:

- ☐ Popliteus
- ☐ Soleus
- ☐ Plantaris
- ☐ Gastrocnemius

6. Iliotibial tract receives the insertion of:

- ☐ Gluteus med. and gluteus min
- ☐ Gluteus max. and gluteus min.
- ☐ Gluteus max. and tensor fascia
- ☐ Gluteus max. and gluteus med.

5. The medial boundary of the femoral triangle is formed by:

- ☐ Adductor longus
- ☐ Sartorius
- ☐ Adductor magnus
- ☐ Iliopsoas

2. It is situated between superior gemelus and inferior gemelus:

- ☐ Quadratus femoris
- ☐ Obturator externus
- ☐ Obturator internus
- ☐ Piriformis

4. The tubercle separating the tendons of peroneus longus and peroneus brevis is:

- ☐ Posterior tubercle
- ☐ Anterior tubercle
- ☐ Peroneal tubercle
- ☐ Medial tubercle

9. The medial femoral circumflex artery is the branch of:

- ☐ Internal iliac artery
- ☐ Femoral artery
- ☐ Profunda femoris artery
- ☐ External iliac artery

8. Structure crossing superficial to the flexor retinaculum:

- ☐ Median nerve.
- ☐ Flexor pollicis longus
- ☐ Ulnar nerve.
- ☐ Flexor digitorum superficialis.

7. The most common site of fracture of the clavicle is:

- ☐ Lateral end.
- ☐ Junction of the medial two-thirds and the lateral third
- ☐ Medial end.
- ☐ Midpoint of the clavicle.

12. The quadrangular space is bounded by all of the following EXCEPT:

- ☐ Teres major.
- ☐ Teres minor.
- ☐ Long head of biceps.
- ☐ Long head of triceps

11. The lateral condyle of tibia possesses articular facet for the head of the fibula on its:

- ☐ Medial aspect
- ☐ Lateral aspect
- ☐ Posterior aspect
- ☐ Anterior aspect

15. ?-Which muscle passes through the greater sciatic foramen

- ☐ Obturator internus
- ☐ Obturator externus
- ☐ Piriformis
- ☐ Superior gemelus

14. Femoral vein receives blood from:

- ☐ Great saphenous vein
- ☐ External iliac vein
- ☐ Internal thoracic vein
- ☐ Internal iliac vein

13. The largest nerve of the body is:

- ☐ Femoral nerve
- ☐ Radial nerve
- ☐ Sural nerve
- ☐ Sciatic nerve

18. : Paralysis of the serratus anterior muscle causes

- ☐ Ape hand.
- ☐ Wrist drop
- ☐ Claw hand.
- ☐ Winging of the scapula.

17. The subsartorial (adductor canal) is located in:

- ☐ Middle third of thigh
- ☐ Upper third of thigh
- ☐ Lower third of thigh
- ☐ Leg

16. Rectus femoris takes origin from:

- ☐ Posterior inferior iliac spine
- ☐ Anterior superior iliac spine
- ☐ Posterior superior iliac spine
- ☐ Anterior inferior iliac spine

21. Which of the following structure does not take part in the formation of the knee joint?

- ☐ Head of fibula
- ☐ Condyle of tibia
- ☐ Lateral femoral condyle
- ☐ Medial femoral condyle

20. The surgical neck of the humerus is related to the:

- ☐ Ulnar nerve.
- ☐ Median nerve.
- ☐ Axillary nerve.
- ☐ Radial nerve.

19. Which structure does not enter the femoral sheath:

- ☐ Lymph nodes
- ☐ Femoral vein
- ☐ Femoral nerve
- ☐ Femoral artery

24. It is the largest and longest bone of the body:

- ☐ Hip bone
- ☐ Vertebra
- ☐ Tibia
- ☐ Femur

23. Superficial peroneal nerve supplies:

- ☐ Anterior compartment of leg
- ☐ Posterior compartment of leg
- ☐ Anteromedial surface of leg
- ☐ Anteriolateral surface of leg

22. Sustentaculum tali is located on_____ of calcaneum:

- ☐ Lateral surface
- ☐ Medial surface
- ☐ Superior surface
- ☐ Anterior surface

27. One of the following is not attached to the medial border of the scapula:

- ☐ Teres minor.
- ☐ Levator scapulae.
- ☐ Serratus anterior.
- ☐ Rhomboides minor

26. Muscles taking origin from the radius:

- ☐ Pronator quadratus.
- ☐ Flexor pollicis longus
- ☐ Supinator.
- ☐ Biceps.

25. The symphysis pubis is:

- ☐ Fibrous joint
- ☐ Secondary cartilaginous joint
- ☐ Synovial joint
- ☐ Primary cartilaginous joint

31. In Erb's paralysis, the deformity is called:

- ☐ Ape hand.
- ☐ Claw hand.
- ☐ Winging of the scapula.
- ☐ Policeman's tip position

29. Carpal tunnel syndrome causes:

- ☐ Ape-like hand
- ☐ Wrist drop.
- ☐ Claw hand.
- ☐ Paralysis of all muscles of the hand.

28. The sesmoid bone of the lower limb is:

- ☐ Calcaneum
- ☐ Ilium
- ☐ Pubis
- ☐ Patela

33. One of the following is a branch of the second part of the axillary artery:

- ☐ Superior thoracic artery
- ☐ Lateral thoracic artery.
- ☐ Posterior circumflex humeral artery.
- ☐ Anterior circumflex humeral artery.

30. Which of the following muscles crosses two joints?

- ☐ Vastus medialis
- ☐ Vastus intermedius
- ☐ Rectus femoris
- ☐ Vastus lateralis

32. The following muscles are innervated by the median nerve EXCEPT:

- ☐ Adductor pollicis
- ☐ Pronator quadratus.
- ☐ Flexor pollicis longus.
- ☐ Pronator teres.

10. The glenoid cavity articulates with the head of the humerus by

- ☐ Cartilaginous articulation.
- ☐ a. Fibrous articulation.
- ☐ Ball and socket synovial articulation
- ☐ Plane synovial articulation.