

UPPER LIMB**1- What is the action of the brachioradialis?**

extension of wrist

extension of elbow

**Flexion of elbow**

flexion of wrist

2- In the cubital fossa, which of the following structures located most medially?

Brachial art.



Median nerve.

Radial art.

Tendon of biceps

3- One of the following muscles rotates arm laterally?

teres major

latissimus dorsi



infraspinatus

subscapularis

4- The muscle that forms the anterior wall of axillary fossa:

latissimus dorsi

subscapularis

teres minor

☒ pectoralis major**5- All of the following muscles innervated by Median nerve Except:**☒ Add. Pollicis

Opponens Pollicis

Abd Pollicis Brevis

Palmaris longus

6- The anterior compartment of the arm innervated by:

Median Nerve

☒ Musculocutaneous Nerve

Ulnar Nerve

Radial Nerve

7- One of the below muscle supply by two nerves.

brachioradialis

☒ . Brachialis

biceps brachii

triceps brachii

8- One of the below choice is in the clavicle.

- styloid process
- coronoid fossa
- ✓ conoid tubercle
- coracoid process

9- All of the following muscles innervated by ulnar nerve except:

- 4th lumbrical
- dorsal interosseous
- ✓ c.1st lumbrical
- palmar interosseous

10- structures that found in the deltopectoral groove:

- brachial vein and acromial art
- ✓ cephalic vein and deltoid art
- basilic vein and deltoid art
- radial vein and acromial art

11- One of the following muscle is innervated by both the ulnar and median nerves.

- palmaris longus
- ✓ flexor digitorum profundus
- flexor digitorum superficialis
- pronator quadratus

12- . This is the strong supinator:

Brachialis



Biceps Brachii

Triceps Brachii

Coraco Brachialis

13-

One of the below muscles abduct the arm to 90 degree.

Supra spinatous

trapezius



middle fibers of deltoid

anterior fibers of deltoid

14-

This muscle attached to the intertubercular groove:

teres major

subscapularis



latissimus dorsi

infrapinatus

15- skin of the superior lateral of the arm innervated by :

ulnar nerve



axillary nerve

median nerve

radial nerve

16- One of the following is not branches from the radial artery.

deep artery arch



common palmar digital

radialis indicis

superficial branch

17- Axillary nerve innervated one of the below muscles.



terse minor

pronator terse

terse major

brachioradialis

18- . Lateral border of the cubital fossa is:

palmaris longus



pronator teres

extensor carpi radialis

brachioradialis

19- The long head of the biceps brachii muscle originated from the:

supraglenoid tubercle

Coracoid process

✓ infraglenoid tubercle

acromion process

20- Dorsal tubercle located at one of the following bone.

Scapula

Ulnar

Humerus

✓ Radius

21- . Profunda brachii artery is along with one of the nerve below

. median nerve

✓ radial nerve

. axillary nerve

ulnar nerve

22- It is action of dorsal interosseous muscle.

flexion



abduction

adduction

extension

23- All of the choice is branches from axillary artery except:

superior thoracic

lateral thoracic



. internal thoracic

thoraco acromial

24- The superficial palmar arterial arch is formed primarily by the:

Radial artery

Posterior interosseous artery

Anterior interosseous artery



Ulnar artery

25- All of the bellow choice is in the anterior view of the humerus except :

capitulum

coronoid fossa



. olecranon fossa

radial fossa

26- This nerve located between brachialis and brachioradialis.

- ✓ radial nerve
- axillary nerve
- median nerve
- ulnar nerve

27- All of the below nerves derived from posterior cord of the brachial plexus except:

- lower subscapular nerve
- thoracodorsal
- upper subscapular nerve
- ✓ long thoracic

28- .All of these muscles belong to the anterior compartment of forearm except:

- pronator quadratus
- pronator teres
- ✓ supinator
- palmaris longus

29- One of the below structures is located in the depth of the snuffbox.

- ulnar artery
- radial nerve
- cephalic vein
- ✓ radial artery

30- .This joint is pivot

elbow

radiocarpal

carpometacarpal

✓ radioulnar

31- . This muscle attached to the pisiform bone.

extensor carpi radialis

✓ flexor carpi ulnaris

flexor carpi radialis

extensor carpi ulnaris

32- . Insertion of the triceps brachii muscle is:

✓ olecranon process

Coronoid process

ulnar tuberosity

styloid process

33- .Recurrent radial artery anastomosis with:

- inferior ulnar collateral artery
- middle collateral artery
- ✓ radial collateral artery
- superior ulnar collateral artery

34- This tendon located medially of the snuff box.

- abductor pollicis longus
- ✓ extensor pollicis longus
- extensor pollicis brevis
- abductor pollicis brevis

35- .One of the below structures passes posteriorly to the flexor retinaculum at the wrist?

- tendon of the palmaris longus
- ✓ median nerve
- ulnar nerve
- ulnar artery

36- . All of the following muscles attach to the medial border of the scapula, except:

- ✓ . teres major
- rhomboid minor
- . levator scapulae
- rhomboid major

37- .These structures are located in the quadrate space .

axillary nerve and anterior circumflex humeral artery

axillary nerve and sub scapular artery .



axillary nerve and circumflex scapular artery .

. axillary nerve and posterior circumflex humeral artery .

38- This is not branches from the median nerve.



posterior inter osseous

anterior inter osseous

recurrent

palmar cutaneous

39- . The extensor expansion of the fingers is formed by the tendon from the:

extensor pollicis longus

extensor pollicis brevis



extensor digitorum

. extensor carpi radialis longus

مدت آزمون: 34 بر اساس دفترچه کاربر UPPER LIMB: آزمون

40- One of the below muscles is an initiator abduction of Gleno-humeral joint:

Deltoid

Infraspinatus

Coraco Brachialis



Supraspinatus