

Surfacing Suicidal Vulnerability Through Simulated Social Interaction: Per-Person Language Model Agents as Communicative Stress Tests

Abstract

Suicidal vulnerability may be encoded in everyday communication patterns but diluted in routine digital interactions. We introduce a method for surfacing this latent signal: training per-person language model agents on individuals’ real text messages and placing them in simulated social interactions—a “communicative stress test.” Using data from 79 adults with recent suicidal ideation, we fine-tuned individual LoRA adapters on Qwen3-8B using each participant’s text messages, then placed agents in standardized conversations with probe personas. Agent-generated risk language was associated with EMA-measured suicidal ideation ($r = .576, p < .001$), with a single neutral small-talk probe performing nearly as well ($r = .551$). A shuffle control confirmed the signal is person-specific ($r = .071$ when adapters were mismatched), and automated descriptions of participants’ general smartphone activity produced no signal, confirming specificity to interpersonal communication. A prompt ablation demonstrated partial robustness to removal of disclosure-encouraging language ($r = .430$). Cross-validation identified a two-probe battery as optimal (LOOCV $r = .570$). This proof-of-concept demonstrates that simulated social interaction can amplify latent vulnerability signals, bridging digital phenotyping, generative AI, and suicide theory.

Keywords: suicidal ideation, large language models, generative agents, digital phenotyping, ecological momentary assessment, communicative stress test

1. Introduction

Suicide remains a leading cause of death globally, and predicting who is at risk continues to be one of the most difficult problems in clinical science (Franklin et al., 2017). Despite decades of research, clinicians perform only marginally better than chance at predicting suicidal behavior (Large et al., 2017), and traditional risk factor approaches have reached apparent ceiling effects in predictive accuracy (Belsher et al., 2019). A central challenge is that suicidal ideation fluctuates rapidly—on the order of hours (Kleiman et al., 2017)—and the interpersonal processes theorized to drive it—including perceived burdensomeness and thwarted belonging—are inherently relational, manifesting most clearly in social interaction rather than in static self-report.

Two converging developments create new opportunities for studying these dynamics. First, digital phenotyping—the continuous measurement of behavior through smartphones and wearable sensors—has produced rich longitudinal records of how individuals communicate in daily life. Screenomics, which captures smartphone screenshots at five-second intervals (Reeves et al., 2021), provides near-complete records of text messages sent and received. When combined with ecological momentary assessment (EMA) of suicidal ideation, these data reveal moment-to-moment relationships between digital communication and suicide risk (Ammerman et al., 2026a). In adolescent populations, intensive longitudinal smartphone sensing has shown particular

promise: Auerbach et al. (2023) demonstrated that mood disturbances measured via smartphone predicted clinically significant suicidal ideation on a weekly basis, and subsequent work used GPS data to detect next-week suicide risk (Auerbach et al., 2025). However, analyzing raw message content or sensor data for suicide risk faces fundamental limitations: most messages are logistical and routine, and risk-relevant content is sparse and diluted across thousands of everyday communications.

Second, the emergence of large language models (LLMs) capable of producing human-like text has enabled a new class of research: generative agent simulation. Park et al. (2023) demonstrated that LLM-powered agents placed in a shared environment exhibit emergent social behaviors without explicit programming. Subsequent work has scaled this to 1,000 agents parameterized from real interview data, achieving 85% behavioral fidelity (Park et al., 2024). More broadly, LLMs have been used as proxies for human participants—Argyle et al. (2023) introduced “silicon sampling” to simulate survey responses, Aher et al. (2023) replicated classic psychology experiments, and Peters & Matz (2024) showed that LLMs can infer Big Five traits from social media posts ($r = 0.29$; see also Brickman et al. 2025). Applications to mental health have begun to appear, including agent-based simulations of student wellbeing (Zhao et al., 2025), frameworks for modeling socio-environmental determinants (Kambeitz & Meyer-Lindenberg, 2025), and simulations of vulnerable users interacting with AI chatbots (Qiu et al., 2025). Louie et al. (2024) validated the use of expert-designed probe personas for patient simulation. Concurrently, there has been a rapid expansion of LLMs applied directly to suicide prevention (Holmes et al., 2025). However, Larooij &

Törnberg (2025b) offered a critical review arguing that LLMs may exacerbate rather than solve the validation challenges inherent in agent-based social simulation.

50 A substantial psycholinguistic literature has established that language carries reliable markers of suicidal risk. Seminal work by Stirman & Pennebaker (2001) demonstrated that suicidal poets used more first-person singular pronouns and fewer first-person plural pronouns than non-suicidal poets. Al-Mosaiwi & Johnstone (2018) showed that absolutist thinking—operationalized
55 through words like “nothing,” “completely,” and “always”—is a cognitive-linguistic marker specific to suicidal ideation beyond what negative emotion words alone capture. More broadly, a systematic review (Ophir et al., 2022) identified consistent markers across studies, including elevated self-referential language, death-related vocabulary, and reduced future-oriented
60 words. Dictionary-based approaches have been central to translating these findings into applied risk detection, with the Linguistic Inquiry and Word Count (LIWC; Pennebaker et al. 2015) program applied to suicide notes (Pestian et al., 2012), crisis hotline transcripts (Huang et al., 2024), and clinical interviews (Tausczik & Pennebaker, 2010). In clinical messaging con-
65 texts, Swaminathan et al. (2023) demonstrated that a two-stage NLP system combining keyword filtering with machine learning reduced crisis message triage time from 9 hours to 8–13 minutes. In our own prior work, we applied dictionary-based classification to text messages extracted from smartphone screenshots, finding that within-person increases in negative emotional
70 expressions co-occurred with elevated suicidal ideation (Ammerman et al., 2026a).

The present study bridges digital phenotyping and generative agent simulation by training per-person language model agents on individuals’ actual text messages and placing those agents in simulated social interactions to surface latent communication patterns associated with suicidal vulnerability (Figure 1). For each participant, we fine-tune a low-rank adaptation (LoRA) adapter on an open-source language model (Qwen3-8B) using their real messages, creating an agent that encodes their communicative style. We then place each agent in standardized conversations with probe personas designed to elicit specific interpersonal dynamics, and score the generated language for risk-relevant content using the dictionary-based approach described above. We conceptualize this as a *communicative stress test*—analogous to cardiac stress testing, where a resting ECG may appear normal but underlying vulnerability becomes visible under exertion. In our paradigm, routine text messages are the resting state, and the simulated conversation is the treadmill. This approach shares a conceptual affinity with recent work using multi-agent AI systems to systematically probe wearable sensor data for clinically relevant biomarkers (Kim et al., 2025), though our method operates on language rather than physiological signals and creates per-person models rather than population-level features.

This approach is novel relative to the existing agent simulation literature in a fundamental way. Prior methods rely on *prompt-based conditioning*—providing demographic descriptions (Argyle et al., 2023), interview transcripts (Park et al., 2024), or expert-designed persona specifications (Louie et al., 2024)—to shape model behavior. This faces an individuation problem: Peng et al. (2025) found that digital twins created through prompting

correlate with real human responses at only $r = 0.20$, and Li et al. (2025) demonstrated that even state-of-the-art models struggle to maintain behavioral consistency. Our approach differs by *fine-tuning* per-person adapters on
 100 actual behavioral data, learning directly from how a person communicates rather than from descriptions of their demographics or interview responses. This weight-level adaptation may overcome the “insufficient individuation” of prompting-based approaches.




Figure 1: Overview of the communicative stress test paradigm. Each participant’s real text messages are used to fine-tune a LoRA adapter on the Qwen3-8B base model, creating a per-person agent. Each agent then engages in 20-turn conversations with standardized probe personas (also Qwen3-8B, without LoRA adaptation). The generated text is scored with risk language dictionaries and correlated with EMA-measured suicidal ideation.

The present study addresses six primary questions. First, do per-person
 105 fine-tuned language model agents produce risk-relevant language at rates that are associated with their real-world counterparts’ suicidal ideation? Second,

does the arena provide incremental predictive validity over direct analysis of participants’ actual text messages? Third, is the predictive signal specific to interpersonal communication style, or would any behavioral data suffice? 110 Fourth, do standardized conversational probes—designed to target suicide-specific theoretical constructs—differentiate from the unstructured all-pairs arena, and does construct-specific scoring improve prediction? Fifth, are the adapters person-specific, such that clinical prediction depends on correct person–adapter matching? Sixth, does the learned communication style 115 reflect stable individual differences or transient states?

2. Method

2.1. *Participants and Procedure*

Participants were 79 adults (ages 18–65) recruited from the community who endorsed past-month active suicidal ideation or suicidal behaviors and 120 owned an Android-based smartphone. Full eligibility criteria and recruitment procedures are described in [blinded for review]. Following a baseline assessment, participants completed 6 EMA prompts per day for 28 days (overall response rate = 68.8%). Simultaneously, a passive sensing application captured screenshots at five-second intervals during active smartphone use, yielding 125 approximately 7.4 million screenshots across the sample ($M = 92,613$ per participant).

2.2. *Message Extraction*

Text messages were extracted from screenshots using a vision-language model (VLM) pipeline. The Qwen2.5-VL model (Qwen Team, 2025) was

130 deployed on NVIDIA L40S GPUs (48 GB VRAM). Each screenshot was
resized to 400×672 pixels and processed with a single structured prompt
containing eight questions (see Appendix Appendix A for full prompt text).
The prompt asked the VLM to identify the application being displayed, the
type of activity, whether notifications were present, the nature of social me-
135 dia engagement, whether a keyboard was visible, and—if a keyboard was
present in a messaging app—to extract text separately for the phone owner
(typically right-aligned dialogue boxes) and the message recipient (typically
left-aligned). Generation used a maximum of 600 new tokens. Validation
against 500 human-annotated screenshots demonstrated 99.7% accuracy for
140 keyboard detection, and when keyboards were present, 100% accuracy for
text extraction from both owner and recipient messages. This pipeline pro-
duced two text streams per participant—*owner text* (messages sent by the
participant) and *recipient text* (messages received).

2.3. EMA Measures

145 At each EMA prompt, participants reported on suicidal ideation using
four items assessing passive ideation (wishes for death, thoughts of death) and
active ideation (thoughts of killing oneself, intent to act). Additional items
assessed suicidal planning (3 items), desire for death (2 items), perceived
burdensomeness (2 items), thwarted belonging (2 items), and negative affect
150 (10 items). All items were summed within construct at each prompt, then
aggregated to person-level means as the primary outcome variables.¹

¹We also examined associations between probe-derived risk composites and within-
person variability (person-level *SDs*) in EMA outcomes. Bivariate correlations were sig-

2.4. Per-Person Agent Training

2.4.1. Message Filtering

Because screenomics captures all on-screen activity, messages were extracted from any messaging application visible on the participant’s phone. The majority of extracted messages came from SMS/native messaging (86.7%), with smaller contributions from Snapchat, Discord, Messenger, WhatsApp, Telegram, and Slack (see Appendix Appendix D for the full distribution). Participants used a median of 2 messaging platforms (range: 1–6). For each participant, sent (owner) text messages were cleaned in two stages. First, messages shorter than five characters, empty strings, and VLM output artifacts were removed. Artifact filtering targeted 16 patterns characteristic of VLM extraction failures. Second, a minimum message count threshold of 50 post-filtering messages was applied. Of the 79 participants, 73 met this threshold. Among eligible participants, message counts varied substantially ($Mdn = 733$; $M = 1,983$; range = 50–13,551). LoRA training was attempted for all 73; two failed to converge, yielding 71 agents with successful adapters. Of these, 64 could be matched with valid EMA data.²

nificant for several constructs (e.g., passive SI SD : $r = .345$, $p = .006$), but partial correlations controlling for the corresponding person-level mean were uniformly non-significant (all $r < .12$, all $p > .35$), indicating that the variability associations were driven by mean- SD dependency rather than unique information about fluctuation. We therefore report only mean-level associations.

²The 15 excluded participants had dramatically fewer messages ($M = 12$ vs. $M = 1,202$; $p < .001$) and somewhat higher SI, though this difference was not statistically significant ($M = 2.20$ vs. $M = 1.78$; $U = 123$, $p = .150$; $N_{\text{excluded with EMA}} = 6$).

2.4.2. LoRA Fine-Tuning

170 We used Qwen3-8B (Qwen Team, 2025) as the base language model. For each participant, we trained a low-rank adaptation (LoRA; Hu et al. 2022) adapter on that individual’s cleaned owner messages. LoRA modifies a small number of parameters (rank $r = 8$, $\alpha = 16$, targeting the query and value projection matrices) while keeping the base model frozen. Each adapter
175 was trained for 3 epochs with a causal language modeling objective. No clinical labels were provided during training; the model learned only from raw message text.

2.5. Arena Paradigms

We evaluated three conversational paradigms of increasing structure, each
180 using the same per-person LoRA adapters and identical generation parameters (temperature = 0.9, top- $p = 0.95$, repetition penalty = 1.3, maximum 200 new tokens per turn). All conversations were scored using the same two-layer dictionary system described below. Figure 2 illustrates the complete pipeline with a worked example: a conversation excerpt between one partic-
185 ipant’s agent and the Neutral Acquaintance probe, the dictionary scoring of agent turns, and the aggregation from turn-level hits to person-level clinical association.




Figure 2: Worked example of the communicative stress test pipeline. (a) A conversation excerpt between a participant agent (SI = 2.92) and the Neutral Acquaintance probe. (b) Dictionary scoring of agent turns, with hits color-coded by category. (c) Aggregation from turn-level hits to person-level clinical association.

2.5.1. Paradigm 1: All-Pairs Arena

Every possible pair of agents ($N = 2,485$ pairs from 71 agents) engaged
 190 in a 20-turn simulated conversation seeded with “How are you feeling to-
 day?” Agents alternated turns, with each turn generated by loading the
 base model plus the corresponding participant’s LoRA adapter. A system

prompt established the conversational context: agents were instructed to respond naturally and honestly, to keep responses to 1–3 sentences, and to
195 never repeat the previous message. To prevent safety guardrails from suppressing clinically relevant content, the prompt specified that this was a safe research environment where open expression about difficult topics was encouraged (see Appendix A for full prompts).

2.5.2. *Paradigm 2: Caption-Trained Arena*

200 To test whether the predictive signal originates specifically from interpersonal communication style, we conducted a comparison using an alternative training corpus. For each participant, a VLM (Qwen3-VL-2B-Instruct) had previously classified every screenshot (approximately 90,000 per participant) into structured behavioral categories using a JSON-output prompt (see Appendix Appendix A.1). Each screenshot was classified along six dimensions:
205 application category (16 options), activity type (10 options), content theme (9 options), mood tone (4 options), social context (6 options), and a brief visible text summary. These structured captions characterize the participant’s digital behavior patterns—but in the VLM’s categorical language rather than the participant’s own words. We fine-tuned the same Qwen3-8B
210 base model with an identical LoRA configuration on these caption corpora using one training epoch (reduced from three given the 50-fold increase in training data). Sixty-seven participants yielded successful caption-trained adapters. These agents were placed in the same all-pairs arena (2,211 paired
215 conversations, 20 turns each).

2.5.3. Paradigm 3: Standardized Probe Arena

To test whether structured conversational probes designed to elicit suicide-specific theoretical constructs could outperform the unstructured all-pairs arena, we developed five standardized probe personas, each targeting a distinct psychological dimension. Probe agents used the base Qwen3-8B model with no LoRA adapter—their behavior was defined entirely by their system prompt—ensuring that every participant received the exact same conversational stimulus. Each participant agent (base model + per-person LoRA) engaged in a 20-turn conversation with each of the five probes (71 participants $\times$ 5 probes = 355 conversations). The five probes were:

1. *Supportive Therapist* (target: emotional disclosure). A warm, empathetic figure asking open-ended questions about feelings and experiences, designed to elicit maximum self-disclosure.
2. *Friend in Crisis* (target: burdensomeness and help-seeking). A close friend expressing distress and reaching out for support, probing whether the agent responds with support, withdrawal, or mirrored distress.
3. *Rejecting Peer* (target: thwarted belonging). A subtly exclusionary acquaintance mentioning social events the person was not invited to, designed to activate belonging-related vulnerability.
4. *Demanding Authority* (target: perceived burdensomeness). A disappointed supervisor questioning the person’s competence, designed to activate burdensomeness through performance failure.
5. *Neutral Acquaintance* (target: baseline). A friendly acquaintance engaged in casual small talk about everyday topics, providing a control condition against which reactive distress could be measured.

Figure 3 illustrates example exchanges from each probe with a single participant agent, showing how the same individual’s communicative style manifests differently across conversational contexts.

In addition to the five original probes, a sixth probe—the Future Self—
245 was included as an exploratory extension based on the temporal self-reflection paradigm (Pataranutaporn et al., 2024). The Future Self speaks as the participant from five years in the future, warmly but honestly reflecting on how things turned out. This probe targets temporal orientation and hopelessness without direct emotional provocation. Each of the 71 participant agents
250 engaged in a single 20-turn conversation with the Future Self probe (71 additional conversations).

Full probe persona prompts are provided in Appendix A. For each participant, we computed per-probe risk composite scores (dictionary hits per turn, scored only on participant turns), a total risk composite (mean across
255 all five original probes), and reactivity scores (each probe’s composite minus the neutral acquaintance baseline).

2.6. Risk Language Scoring

To score conversation content, we employed a two-layer dictionary approach. The first layer consisted of the seven sub-dictionaries developed by
260 Ammerman et al. (2026b), who adapted the 276-word crisis language dictionary validated by Swaminathan et al. (2023) (sensitivity = 0.98, PPV = 0.66). Two experts independently assigned each word to seven themes: suicidal thoughts (41 entries), non-substance methods (75 entries), substances (26 entries), sleep (7 entries), hopelessness (26 entries), general risk (93 entries),
265 and help-seeking (8 entries).

The second layer added six supplementary categories aligned with interpersonal and cognitive theories of suicide (O’Connor, 2011) and absolutist thinking (Al-Mosaiwi & Johnstone, 2018): perceived burdensomeness (18 patterns), thwarted belonging (19 patterns), entrapment (14 patterns), negative affect (36 patterns), absolutist thinking (18 patterns), and self-harm (5 patterns). The complete dictionary is provided in Appendix Table S1.

2.7. Validation Analyses

Five analyses were conducted to assess the robustness and specificity of the clinical signal.

2.7.1. Adapter Shuffle Control

To test whether the clinical signal is person-specific—that is, whether it resides in the learned adapter rather than in some artifact of the arena procedure—we randomly reassigned LoRA adapters to different participants (seed = 42) and re-ran the all-pairs arena with these mismatched adapters. If the signal is carried by person-specific adaptation, correlations between arena output and the *participant’s* SI should drop to zero when adapters are mismatched. Conversely, if the adapter carries the signal, the shuffled arena output should correlate with the *adapter-owner’s* SI rather than the assigned participant’s.

2.7.2. Temporal Split Validation

To test whether adapters capture stable communicative dispositions rather than transient states, we chronologically split each participant’s messages at the median timestamp, trained separate LoRA adapters on the first and

second halves, and ran the full 5-probe battery with each set of adapters independently. We report: (a) split-half correlation of risk composites between
290 the two temporal halves, (b) Spearman-Brown corrected reliability, and (c) clinical prediction (correlation with EMA SI) for each split separately. Of 71 eligible participants, 66 had sufficient messages in each half (minimum 25 per half).

295 2.7.3. *Cross-Validated Prediction*

To assess out-of-sample predictive validity, we used 10-fold cross-validation and leave-one-out cross-validation (LOOCV) with Ridge regression predicting SI from the five per-probe risk composites (5 features). Ridge regression was chosen to guard against overfitting given the small sample and moderate
300 feature count. To evaluate probe subset selection, we additionally performed LOOCV with internal subset search: within each held-out fold, exhaustive search over all probe combinations identified the best subset on the $N - 1$ training set, and that subset was evaluated on the held-out participant. This guards against overfitting in probe selection, which can inflate in-sample sub-
305 set performance.

2.7.4. *Prompt Ablation*

To test whether the clinical signal depends on the disclosure-encouraging language in the participant system prompt (a potential demand characteristic), we re-ran the full 5-probe battery with a minimal participant prompt
310 that omitted all references to mental health, self-harm, and suicide. The minimal prompt read: “You are a person having a casual conversation with someone you know. Respond naturally and honestly as yourself. Keep your

response to 1-3 sentences. NEVER repeat or paraphrase what the other person just said. Always say something new and different from the previous message.” Probe persona prompts were unchanged. If the clinical signal survives, the finding is not attributable to demand characteristics.

2.7.5. *No-LoRA Floor Control*

To establish what proportion of risk language is attributable to the person adapter versus the base model, we ran the 5-probe battery 20 times using the base Qwen3-8B model with no LoRA adapter and the original participant prompt. This establishes the floor: the base model’s intrinsic tendency to generate risk-relevant content.

2.8. *Sensitivity Analyses*

To address potential confounds, we conducted three sensitivity analyses: (a) verbosity normalization—re-scoring all conversations with dictionary hits normalized by token count rather than turn count; (b) training heterogeneity—testing whether message count (range: 50–13,551) confounds the risk composite–SI association; (c) selection bias—comparing excluded participants ($N = 15$) to included participants ($N = 64$) on baseline SI and message volume. Reactivity score analyses are reported in Appendix Appendix C.

2.9. *Analytic Plan*

All primary analyses were between-person, correlating each participant’s arena-derived scores with their person-level EMA means. We report Pearson correlations for all primary analyses, with Spearman correlations for

zero-inflated distributions. The central comparison examined three sources of risk language—real messages, all-pairs arena, and probe arena—against the same EMA outcomes. For the probe arena, we additionally tested construct-specific predictions (e.g., does the Rejecting Peer specifically predict thwarted belonging?) and reactivity scores (probe minus neutral baseline). The validation analyses described above assessed adapter specificity (shuffle control), temporal stability (split-half), out-of-sample generalization (cross-validation), demand characteristics (prompt ablation), and base-model floor (no-LoRA control). We applied no correction for multiple comparisons; this study is exploratory, and we report the total number of tests conducted to facilitate reader evaluation. We note that the present analyses do not incorporate baseline clinical data (e.g., baseline interview-assessed SI severity, diagnostic status, or demographic covariates) as predictors or covariates; an important question for future work is whether arena-derived risk scores provide incremental predictive validity above and beyond baseline assessment.

3. Results

Figure 4 provides a roadmap of all analyses reported in this section, organized by purpose: core paradigms testing the communicative stress test concept, validation analyses confirming the signal’s properties, and sensitivity analyses addressing potential confounds.

Summary of Analyses

Analysis	Question	Key Result	Section
CORE PARADIGMS			
All-Pairs Arena	Do agents associate with SI?	$r = .433, p < .001$	3.2
Caption-Trained Arena	Is signal interpersonal-specific?	$r = .159, p = .198$	3.3
Probe Arena (5 probes)	Do probes outperform arena?	$r = .576, p < .001$	3.4
Future Self Probe	Does temporal reflection help?	$r = .369, p = .003$	3.5
VALIDATION			
Adapter Shuffle	Is signal person-specific?	$r = .071, p = .577$	3.5
Prompt Ablation	Is it demand characteristics?	$r = .430, p < .001$	3.5
No-LoRA Floor	Does base model produce signal?	Variance ratio = 2.44x	3.5
Temporal Split	Is signal temporally stable?	SB reliability = .538	3.5
Cross-Validated Battery	Which probes generalize?	L00CV $r = .570, p < .001$	3.5
SENSITIVITY			
Verbosity Control	Confounded by turn length?	$r = .438, p < .001$	3.5
Training Heterogeneity	Confounded by msg count?	partial $r = .547, p < .001$	3.5
Selection Bias	Who is excluded?	Excluded: fewer messages, $p < .001$	4.7

Core Paradigms Validation Sensitivity

$N = 64$. SB = Spearman-Brown corrected reliability. Neut. = Neutral Acquaintance; Ther. = Supportive Therapist.

Figure 4: Summary of all analyses. Core paradigms test the communicative stress test concept across different conversational configurations. Validation analyses confirm the signal is person-specific, partially robust to prompt changes, and not attributable to base-model artifacts. Sensitivity analyses address potential confounds. $N = 64$.

3.1. Risk Language in Raw Messages

Before examining arena-generated language, we assessed the prevalence of risk-relevant content in participants’ actual text messages. Of the 79 participants, 20 (25.3%) had at least one message containing explicit suicide-related language (e.g., “suicidal,” “kill myself,” “end it all”), but such messages were extremely rare—264 of 155,609 total messages (0.17%). Applying the broader risk dictionary to raw messages, death/suicide terms appeared in 624 messages from 51% of participants, belonging-related terms in 444 mes-

sages from 38%, negative affect in 606 messages from 39%, and hopelessness
365 in 155 messages from 23%. Self-harm language was rare (21 messages, 5%
of participants). These base rates illustrate the core challenge that moti-
vates the arena approach: risk-relevant language is present in real messages
but sparse and diluted across thousands of routine communications, making
direct surveillance both ethically problematic and statistically inefficient.

370 3.2. All-Pairs Arena

3.2.1. Descriptives

The 71 agents produced 2,485 paired conversations (49,700 total turns)
with 0% echo rate. Of the 71 agents, 26 (37%) spontaneously mentioned
suicide at least once. After matching with EMA data, 64 participants consti-
375 tuted the analytic sample. Table 1 presents descriptive statistics for the EMA
outcome variables. Suicidal ideation scores were positively skewed ($Mdn =$
1.18, $M = 1.80$), with substantial between-person variability ($SD = 2.15$;
range: 0.00–8.50).

Table 1: Descriptive statistics for EMA outcome variables ($N = 64$).

Measure	M	SD	Mdn	Range
<i>Person-level means</i>				
Suicidal ideation (composite)	1.80	2.15	1.18	0.00–8.50
Passive SI	2.98	1.21	2.57	2.00–7.00
Active SI	2.82	1.04	2.42	2.00–6.22
Suicidal planning	3.25	0.53	3.09	2.98–6.67
Desire for death	2.49	0.78	2.18	2.00–5.33
Perceived burdensomeness	2.35	2.28	1.66	0.00–7.67
Thwarted belonging	2.72	2.23	2.40	0.00–7.92
Negative affect	7.78	6.61	6.40	0.10–30.60




Figure 5: Forest plot of key correlations with suicidal ideation across all analyses, with 95% confidence intervals. Core paradigms show that the probe arena and neutral small-talk probe are the strongest predictors, while the caption-trained arena produces no signal. Among individual probes, Neutral Acquaintance dominates and Demanding Authority fails. Validation analyses confirm adapter specificity (shuffle wrong-slot CI crosses zero) and partial prompt robustness. Cross-validated estimates show Neutral + Therapist as the most robust battery. Effect size benchmarks at $r = .30$ (medium) and $r = .50$ (large) are shown for reference.

3.2.2. Suicide Mention Rate

380 The rate at which agents mentioned suicide was significantly associated with real-world SI (Pearson $r = .269$, $p = .032$; Spearman $\rho = .438$, $p < .001$). Agents who mentioned suicide at least once had significantly higher real SI ($M = 2.78$) than those who never mentioned it ($M = 1.22$), $t(62) = 2.98$, $p = .004$, Cohen's $d = 0.77$.

385 3.2.3. Risk Dictionary Correlations

The strongest correlate of mean SI was absolutist thinking ($r = .607$, $p < .001$), followed by thwarted belonging ($r = .474$, $p < .001$), perceived burdensomeness ($r = .403$, $p = .001$), substance use ($r = .375$, $p = .002$), and death ideation ($r = .349$, $p = .005$). The full risk composite correlated at 390 $r = .433$ ($p < .001$). Absolutist thinking was the only category significantly associated with every clinical construct assessed, including suicidal planning ($r = .416$) and desire ($r = .544$).

3.2.4. Construct Convergence

Simulated burdensomeness was associated with real perceived burdensomeness 395 ($r = .282$, $p = .024$); simulated thwarted belonging was associated with real thwarted belonging ($r = .323$, $p = .009$). Simulated negative affect was not associated with EMA negative affect ($r = .194$, $p = .124$).

3.2.5. Arena Versus Raw Messages

Table 2 compares arena-generated risk language with participants' real 400 sent messages across dictionary categories. The arena produced numerically larger correlations with SI for the majority of categories, winning 66% of 112 dictionary $\times$ EMA comparisons. The difference was most pronounced

for interpersonal constructs: perceived burdensomeness (arena $r = .403$ vs. raw messages $r = .079$), thwarted belonging (.474 vs. .217), and absolutist
405 thinking (.607 vs. .113). Raw messages, however, showed numerically larger associations for explicit crisis content such as non-substance methods (.450 vs. .073) and hopelessness (.399 vs. .306)—categories where the actual words used in real messages carry direct clinical meaning that the arena’s generated language may dilute. These pairwise differences were not formally tested for
410 statistical significance; the comparison is descriptive.

Table 2: Per-dictionary correlations with suicidal ideation: message-trained arena versus raw sent messages ($N = 64$).

Dictionary category	Owner r	Arena r	Larger
Absolutist thinking	.113	.607	Arena
Thwarted belonging	.217	.474	Arena
Perceived burdensomeness	.079	.403	Arena
Substance use	.295	.375	Arena
Death ideation	.256	.349	Arena
Help-seeking	.205	.346	Arena
Negative affect	.254	.289	Arena
Hopelessness	.399	.306	Owner
Non-substance methods	.450	.073	Owner
Risk composite	.342	.433	Arena

3.3. *Caption-Trained Arena*

The caption-trained arena yielded no significant associations with SI (Table 3; $N = 67$ caption-trained agents matched with EMA). The risk com-

posite was not associated with mean SI ($r = .159$, $p = .198$), compared to
 415 $r = .384$ ($p = .001$) for raw sent messages in the same sample. No individual
 dictionary category reached significance. Despite having approximately $50\times$
 more training data, the caption-trained agents failed to capture the clinically
 relevant signal that emerged when agents were trained on participants’ own
 words.

Table 3: Risk composite correlations with EMA outcomes: raw messages versus caption-trained arena ($N = 67$). Caption-trained agents produced no significant associations despite $50\times$ more training data.

EMA outcome	Owner r	Caption arena r
SI mean	.384**	.159
Passive SI	.379**	.163
Active SI	.349**	.135
Planning	.069	.076
Desire	.094	.155
Burdensomeness	.333**	.052
Belonging	.355**	.094
Negative affect	.223	.115

* $p < .05$; ** $p < .01$; *** $p < .001$.

420 3.4. Standardized Probe Arena

3.4.1. Descriptives

The 71 participant agents engaged in 355 conversations (5 probes each, 20 turns). Suicide mentions were rare and probe-specific: Friend in Crisis elicited explicit suicide language from 3 agents (5%), Neutral Acquaintance

425 from 1 (2%), and the remaining probes from none. Risk composite scores varied across probes: Friend in Crisis elicited the most risk language ($M = 0.52$, $SD = 0.59$), followed by Demanding Authority ($M = 0.45$, $SD = 0.54$), Supportive Therapist ($M = 0.39$, $SD = 0.58$), Neutral Acquaintance ($M = 0.35$, $SD = 0.52$), and Rejecting Peer ($M = 0.33$, $SD = 0.43$). After
430 matching with EMA, 64 participants constituted the analytic sample.

3.4.2. Probe Battery Prediction

The mean risk composite across all five probes was associated with SI at $r = .576$ ($p < .001$)—stronger than the all-pairs arena ($r = .477$), real sent messages ($r = .461$), or any single analysis approach. This advantage was
435 consistent across nearly all EMA outcomes: passive SI (probes .585, arena .470, owner .454), active SI (probes .514, arena .442, owner .425), planning (probes .334, arena .300, owner .220), desire (probes .357, arena .278, owner .238), and negative affect (probes .375, arena .244, owner .312). Real messages outperformed only for perceived burdensomeness (.440 vs. .392) and
440 thwarted belonging (.441 vs. .368). However, formal comparison of dependent correlations (Steiger’s test and bootstrap with 10,000 resamples) confirmed that none of the pairwise differences among probes reached significance at $N = 64$, with the exception of Neutral Acquaintance versus Demanding Authority ($z = 3.34$, $p < .001$; bootstrap 95% CI for difference: [.119, .704]).
445 The combined probe composite ($r = .576$) did not significantly differ from the Neutral Acquaintance alone ($r = .551$; $z = 0.29$, $p = .774$; bootstrap 95% CI: [−.217, .314]) or from the Supportive Therapist ($r = .412$; $z = 1.72$, $p = .085$). Cross-validation further reveals that a two-probe combination outperforms the full battery (see Section 3.5).

450 3.4.3. Individual Probe Associations

Among individual probes, the Neutral Acquaintance—a casual small-talk partner with no emotional provocation—showed the single strongest association with SI ($r = .551, p < .001$), outperforming the Supportive Therapist ($r = .412, p < .001$), Rejecting Peer ($r = .326, p = .009$), Friend in Crisis ($r = .316, p = .011$), and Demanding Authority ($r = .119, p = .348$). This pattern held for passive SI (neutral .557, therapist .416), active SI (neutral .493, therapist .368), and desire (neutral .492, therapist .270). The Neutral Acquaintance was also significantly associated with perceived burdensomeness ($r = .259, p = .039$), thwarted belonging ($r = .271, p = .030$), and negative affect ($r = .314, p = .012$). Figure 6 displays the per-probe correlations; Table 4 presents the full omnibus correlation matrix across all probes and EMA outcomes.

460




Figure 6: Per-probe correlations with suicidal ideation ($N = 64$). Error bars indicate 95% CIs (Fisher z). * $p < .05$.

Table 4: Correlations between probe-derived risk composites and EMA outcomes ($N = 64$). Bold indicates $p < .01$; * indicates $p < .05$.

Source	SI	Passive	Active						Neg
	Mean	SI	SI	Plan	Desire	Burden	Belong	Aff	
Neutral Acquaintance	.551	.557	.493	.189	.492	.259*	.271*	.314*	
Supportive Therapist	.412	.416	.368	.168	.270*	.222	.249*	.309	
Future Self	.369	—	—	—	—	.191	.253*	.262*	
Rejecting Peer	.326	.290*	.293*	.249*	.112	.374	.304	.119	
Friend in Crisis	.316*	.277*	.276*	.227	.200	.263*	.250*	.173	
Demanding Authority	.119	.144	.098	.009	.023	.080	.038	.131	
Combined (5 probes)	.576	.585	.514	.334	.357	.392	.368	.375	
All-pairs arena	.477	.470	.442	.300	.278*	.267*	.323*	.244	
Raw messages (owner)	.461	.454	.425	.220	.238	.440	.441	.312*	

3.5. Validation and Sensitivity Analyses

3.5.1. Adapter Specificity

465 When adapters were shuffled—randomly reassigned so that each participant’s conversations were generated using a different person’s LoRA adapter—the risk composite correlation with the *assigned participant’s* SI dropped to $r = .071$ ($p = .577$), compared to $r = .433$ ($p < .001$) for correctly matched adapters. Critically, the same shuffled output correlated $r = .426$ ($p < .001$)
470 with the *adapter-owner’s* SI—the person whose messages originally trained that adapter (Figure 7). This double dissociation confirms that each adapter encodes its own person’s clinical signal: the signal follows the adapter, not the person to whom it is assigned. When a high-risk person’s adapter generates language in another person’s conversational context, the output still

475 reflects the high-risk person’s communicative style.




Figure 7: Adapter specificity: the double dissociation. (a) In the shuffle procedure, each participant’s LoRA adapter is reassigned to a different person, and conversations are re-generated. The key question is whose SI the shuffled output predicts. (b) Results: when correctly matched, the risk composite is associated with SI ($r = .433$). When adapters are shuffled, the output no longer predicts the assigned participant’s SI ($r = .071, p = .577$) but still predicts the adapter-owner’s SI ($r = .426, p < .001$). The clinical signal follows the adapter, not the person to whom it is assigned.

3.5.2. Cross-Validated Probe Battery Optimization

The Future Self probe was associated with SI at $r = .369$ ($p = .003$), positioned between the Neutral Acquaintance ($r = .551$) as the strongest individual probe and the Demanding Authority ($r = .119$) as the weakest.

480 Exhaustive subset search identified {Neutral Acquaintance, Supportive Therapist, Future Self} as the optimal 3-probe combination in-sample ($r = .655$); however, LOOCV with internal subset search—performing subset selection independently within each held-out fold—reduced this to $r = .420$ ($p < .001$)

and 10-fold CV to $r = .492$ ($p < .001$), confirming substantial overfitting in
 485 the in-sample estimate. The 3-probe subset was selected in 83% of LOOCV
 folds, indicating stable feature selection despite attenuated effect size.

Importantly, the simpler fixed combination of Neutral Acquaintance and
 Supportive Therapist—requiring no data-driven selection—achieved LOOCV
 $r = .570$ ($p < .001$), outperforming both the internally optimized subset
 490 ($r = .420$) and the full 5-probe battery (LOOCV $r = .462$). The addition of
 any further probes reduced cross-validated performance (Figure 8).




Figure 8: In-sample versus cross-validated (LOOCV) prediction across probe configurations. The in-sample optimal 3-probe subset ($r = .655$) drops substantially under LOOCV with internal subset search ($r = .420$). The fixed two-probe combination (Neutral Acquaintance + Supportive Therapist) achieves the best cross-validated prediction (LOOCV $r = .570$).

3.5.3. Temporal Stability

First-half and second-half risk composites correlated at $r = .368$ ($p = .002$; Spearman-Brown corrected reliability = .538). Both splits were significantly associated with SI: first-half $r = .359$ ($p = .005$), second-half $r = .396$
 495 ($p = .002$). However, construct-specific predictions diverged: second-half

adapters recovered perceived burdensomeness ($r = .34$) and thwarted belonging ($r = .40$) correlations comparable to full adapters, while first-half adapters did not ($r = .08-.12$, n.s.). The Neutral Acquaintance was the most temporally stable probe (first vs. second half: $r = .304$, $p = .013$). For comparison, EMA suicidal ideation scores themselves showed split-half reliability of $r = .451$ (Spearman-Brown = .621; $N = 49$), with burdensomeness at $r = .668$ (SB = .801) and belonging at $r = .722$ (SB = .839). The adapter split-half ($r = .368$, SB = .538) is lower than the EMA SI benchmark (Figure 9).




Figure 9: Temporal split validation. Adapters trained on the first or second half of each participant’s messages are compared to full adapters. Error bars indicate 95% CIs (Fisher z). SI prediction is preserved across both splits, but construct-specific predictions (burden, belonging, negative affect) are recovered only by second-half adapters. n.s. = not significant.

3.5.4. Prompt Ablation

With the minimal prompt, the risk composite was associated with SI at $r = .430$ ($p < .001$), compared to $r = .576$ ($p < .001$) with the original prompt. Associations with perceived burdensomeness ($r = .309$, $p = .013$) and thwarted belonging ($r = .289$, $p = .021$) also survived. Only negative

affect dropped to non-significance ($r = .191$, $p = .130$). Cross-prompt composites correlated at $r = .612$ ($p < .001$), confirming that the same individual differences are expressed regardless of prompt condition. The prompt’s role is therefore one of permission—amplifying an existing signal rather than creating one.

The per-probe pattern shifted under the minimal prompt. The Rejecting Peer became the strongest single probe (Spearman $\rho = .468$, $p < .001$; Pearson $r = .410$ after outlier removal), while the Neutral Acquaintance dropped from $r = .551$ to $r = .235$ ($p = .061$). Suicide mentions did not decrease under the minimal prompt (2.5% vs. 1.1%; Figure 10).




Figure 10: Prompt ablation: per-probe SI correlations under the original disclosure-encouraging prompt versus a minimal prompt with no mention of mental health. Error bars indicate 95% CIs (Fisher z). The overall signal partially survives ($r = .430$ vs. $r = .576$). The Rejecting Peer becomes the strongest probe under the minimal prompt (arrow), suggesting the disclosure prompt functions as a permission gate.

3.5.5. No-LoRA Floor Control

The base model without any LoRA adapter produced risk language at a mean composite rate of 0.0036 ($SD = 0.0011$), nearly identical to the LoRA-adapted mean of 0.0035 ($SD = 0.0017$). Zero of the 100 base-model
 525 conversations ($20 \text{ runs} \times 5 \text{ probes}$) contained suicide mentions. The critical difference was variance: LoRA-adapted agents showed 2.44 times the between-conversation variance of the base model. The LoRA’s contribution is not in shifting the mean level of risk language but in creating individual-level differentiation that tracks real clinical differences (Figure 11).




Figure 11: No-LoRA floor control. (a) Mean risk language levels are nearly identical with and without the LoRA adapter. (b) Between-agent variance is 2.44 $\times$ higher with LoRA adapters. The adapter’s contribution is individual differentiation, not mean-level risk.

530 3.5.6. Verbosity

Mean tokens per turn correlated modestly with SI ($r = .384$, $p = .002$), confirming a potential confound. However, token-normalized risk composites were still significantly associated with SI ($r = .438$, $p < .001$), with only modest attenuation from the per-turn rate ($r = .576$). Suicidal thoughts,
 535 substance use, thwarted belonging, and general risk categories survived token normalization.

3.5.7. *Training Heterogeneity*

Message count correlated weakly with SI ($r = .224$, $p = .075$) and modestly with risk composite ($r = .310$, $p = .013$). After partialing out message count, the risk composite was still strongly associated with SI (partial
540 $r = .547$, $p < .001$). However, a median split revealed that prediction was substantially stronger in the high-volume half ($r = .610$, $p < .001$) than the low-volume half ($r = .186$, n.s.), indicating that adapters trained on more data produce more valid signals.

545 4. Discussion

This study introduces a communicative stress test paradigm for studying suicidal vulnerability—training per-person language model agents on real text messages and placing them in simulated social interactions. Several principal findings emerge, each with distinct theoretical and practical impli-
550 cations.

4.1. *Communication Style Encodes Suicidal Vulnerability*

Per-person fine-tuned agents produce risk-relevant language at rates that meaningfully predict real-world suicidal ideation, despite never being trained on clinical data. The strongest single correlate—absolutist language ($r =$
555 $.607$)—replicates and extends the seminal finding of Al-Mosaiwi & Johnstone (2018) in a novel paradigm. The construct convergence provides preliminary evidence that the arena captures theoretically meaningful interpersonal processes: simulated burdensomeness predicted real burdensomeness; simulated belonging predicted real belonging.

560 4.2. *The Signal Is Specifically Interpersonal*

The caption-trained arena comparison provides a critical specificity test. Agents trained on VLM-generated descriptions of participants’ digital behavior within the 2-hour EMA reporting windows—approximately 90,000 screenshots per person covering all applications, content, and temporal patterns—
565 produced no clinical signal whatsoever. This rules out two alternative explanations: that any behavioral data would work (it does not), and that more training data is better ($50\times$ more data produced worse results). The predictive signal resides specifically in *how a person talks to other people*, not in what they do on their phone.

570 4.3. *Tonic Style, Not Probe Engineering, Drives Prediction*

The standardized probe arena ($r = .576$ for combined probes predicting SI) outperformed both the all-pairs arena ($r = .477$) and raw message analysis ($r = .461$). In-sample, the five-probe composite ($r = .576$) and the single Neutral Acquaintance ($r = .551$) appear nearly equivalent. However,
575 cross-validation reveals a more nuanced picture: the Neutral Acquaintance alone achieves LOOCV $r = .476$, while adding the Supportive Therapist yields LOOCV $r = .570$ —a genuine incremental gain that is obscured in the in-sample comparison. Adding further probes beyond these two reduces out-of-sample prediction (full 5-probe LOOCV $r = .462$), and the in-sample
580 optimal 3-probe subset ($r = .655$) was substantially overfit, dropping to $r = .420$ under LOOCV with internal subset search—a cautionary example of post-hoc probe selection on small samples.

Theory-driven probe engineering did not achieve its intended purpose of construct-specific prediction; the Demanding Authority probe showed no

585 significant association with SI, and no targeted probe outperformed the unstructured arena for its intended construct.

Instead, what the agents encode is tonic communication style—a person-level disposition that pervades all conversational contexts rather than a construct-specific reactivity that appears only under targeted provocation. The negative reactivity correlations confirm this: higher-SI participants produced *less* 590 additional risk language when provoked, not more, suggesting their baseline is already elevated and further provocation pushes them toward withdrawal rather than expression.

The most striking finding is that the Neutral Acquaintance—casual small talk with no emotional provocation—showed the single strongest association 595 with SI ($r = .551$). The provocative probes do not add construct-specific signal; their value, when it exists (as with the Supportive Therapist), lies in providing a complementary conversational context that elicits a different facet of the same tonic style. Why the Supportive Therapist specifically adds incremental value over the Neutral Acquaintance is not clear from the 600 present data. Possible mechanisms include encouraging emotional elaboration (increasing the density of scorable content), providing a complementary conversational register, or eliciting a different facet of tonic style. Future work should investigate this directly, for example by comparing token counts 605 and content diversity across probe conditions.

4.4. *The Disclosure Prompt as Permission Gate*

The prompt ablation demonstrates that the clinical signal partially survives the removal of disclosure-encouraging language. With the minimal prompt—containing no references to mental health, self-harm, or suicide—

610 the risk composite was still associated with SI at $r = .430$ ($p < .001$), a 26% relative reduction from the original $r = .576$. Cross-prompt composites correlated at $r = .612$ ($R^2 = .375$), indicating that approximately 38% of the between-person variance in risk language is shared across prompt conditions, with the remaining 62% prompt-dependent. The disclosure language
615 substantially amplifies the signal but does not create it. This is more accurately described as “demand characteristics account for some but not all of the effect” rather than “demand characteristics are ruled out.”

Instead, the disclosure prompt functions as a permission gate—a contextual variable that determines *where* vulnerability is expressed, not *whether* it
620 exists. With disclosure permission, tonic vulnerability pervades all conversational contexts including neutral small talk; without it, the same vulnerability concentrates in contexts that naturally create emotional pressure (social rejection). The Rejecting Peer’s rise from $r = .326$ to Spearman $\rho = .468$ under the minimal prompt, while the Neutral Acquaintance dropped from
625 $r = .551$ to $r = .235$, demonstrates this redistribution clearly. This parallels findings in clinical interviewing, where structured permission to discuss suicidal thoughts increases disclosure without altering underlying risk.

Notably, suicide mentions did not decrease under the minimal prompt (2.5% vs. 1.1%), directly contradicting the demand-characteristic hypothe-
630 sis. If the disclosure-encouraging language were manufacturing risk content, its removal should have reduced suicide mentions; instead, they increased, suggesting the minimal prompt may have reduced compliance with conversational norms that typically suppress such content.

4.5. *The Floor Beneath the Signal*

635 The no-LoRA floor control demonstrates that the base model contributes a uniform, low-level baseline of risk language with essentially zero between-conversation variance. The mean risk composite was nearly identical with and without LoRA (0.0035 vs. 0.0036), but the LoRA-adapted agents showed 2.44 times the variance. The adapter’s contribution is not to increase risk
640 language overall but to differentiate individuals—creating a distribution of risk language rates where the rank ordering maps onto real clinical differences.

This finding distinguishes our approach from concerns about LLM “hallucination” of clinical content. The risk language is not model-generated noise uniformly distributed across all agents; it is person-specific variation
645 introduced by the LoRA adapter and traceable to individual communicative style. Combined with the shuffle validation (Section 3.4), this provides converging evidence that the clinical signal is genuinely person-level: it follows the adapter, not the base model, and not the participant slot.

4.6. *Adapter Specificity and the Digital Twin Challenge*

650 The shuffle validation provides the strongest evidence that the clinical signal is person-specific. When adapters were mismatched, the signal followed the adapter ($r = .426$ with the adapter-owner’s SI) rather than disappearing into noise—demonstrating that LoRA fine-tuning creates genuine individual-level representations. The double dissociation (prediction drops to $r = .071$
655 for the wrong participant, remains at $r = .426$ for the adapter-owner) rules out the possibility that the clinical signal is an artifact of arena position, conversation dynamics, or dictionary scoring.

This result contrasts sharply with the performance of prompt-based digital twins. Peng et al. (2025) identified five systematic distortions in LLM-based digital twins and found that the average correlation between simulated and real human responses is only $r = 0.20$. Our LoRA-based approach may overcome the “insufficient individuation” problem precisely because it learns from actual behavioral data rather than from demographic or interview-based descriptions. Whereas prompt-based conditioning provides the model with *information about* a person, weight-based adaptation gives the model *experience as* that person—a fundamentally different personalization mechanism that appears to encode deeper aspects of communicative identity.

4.7. Temporal Stability and the Nature of the Learned Signal

The split-half reliability of .538 falls below the conventional .70 threshold for stable individual differences, suggesting that the adapters capture a mixture of stable communicative dispositions and time-varying patterns. This is not necessarily a limitation: suicidal ideation itself fluctuates substantially within individuals over short timescales (Kleiman et al., 2017), and an adapter that fully captured only trait-level communication style would risk missing the dynamic component that distinguishes elevated risk states from baseline. Crucially, SI prediction was preserved across both temporal splits ($r = .359$ and $r = .396$), indicating that the core clinical signal is sufficiently stable for practical use. That second-half adapters recovered construct-specific predictions better than first-half adapters may reflect increasing comfort with texting over the study period or temporal proximity to EMA assessments.

4.8. *Who This Method Reaches—and Who It Misses*

The 15 participants excluded from the analytic sample had dramatically fewer messages ($M = 12$ vs. $M = 1,202$; $p < .001$) and somewhat higher SI, though this difference did not reach significance ($M = 2.20$ vs. $M = 1.78$; $p = .150$; only 6 excluded participants had EMA data). This reveals a fundamental boundary condition of any text-based approach: those who communicate least digitally are hardest to model. The method is best understood as a screening tool for moderate-risk populations who produce sufficient digital text, rather than as a universal risk assessment instrument. Training heterogeneity further constrains the method’s reach: prediction was substantially stronger in the high-message-volume half of the sample ($r = .610$) than the low-volume half ($r = .186$, n.s.), suggesting a practical minimum of approximately 700 messages for reliable clinical prediction.

4.9. *Practical Implications*

These findings suggest a deployment framework with two key parameters: message volume and probe configuration. Preliminary evidence from a median split suggests that adapters trained on more messages produce stronger associations (high-volume $r = .610$ vs. low-volume $r = .186$), though the precise minimum message count for reliable assessment requires further systematic testing. Once trained, a single 20-turn conversation with a neutral small-talk partner provides a minimum viable risk estimate (LOOCV $r = .476$) that outperforms direct analysis of thousands of real messages. Adding a Supportive Therapist conversation provides the most robust cross-validated performance (LOOCV $r = .570$), while larger probe batteries do not improve generalization.

The pipeline operates on learned communication style rather than surveillance of ongoing private messages. Training the per-person adapter requires access to an individual’s text messages, but once trained, ongoing risk monitoring operates on generated text. This is an incremental rather than categorical improvement in privacy—reducing but not eliminating the need for access to private communications. The ethical implications of creating person-specific language models from private communications require careful consideration, even when those models are used only for research purposes.

4.10. Limitations and Future Directions

The sample is moderate ($N = 64$) and recruited for recent suicidal ideation. All analyses are between-person and correlational; the strongest next step would examine whether changes in arena behavior across sessions track within-person SI fluctuations. Token normalization attenuated some effects (risk composite from $r = .576$ to $r = .438$), confirming that verbosity partially confounds per-turn scoring, though core findings survive.

All results are based on a single model (Qwen3-8B); whether larger models or alternative architectures would strengthen the clinical signal remains an open question. The dictionary-based scoring is context-insensitive (“I don’t want to kill myself” scores identically to “I want to kill myself”); averaging over many turns mitigates but does not eliminate this limitation. The two-probe battery recommendation (LOOCV $r = .570$) requires validation in an independent sample, as does the in-sample optimal 3-probe subset ($r = .655$), which dropped to $r = .420$ under LOOCV with internal subset search. No correction for multiple comparisons was applied; this study is explicitly exploratory, generating hypotheses for confirmatory testing. The validation

challenges highlighted by Larooij & Törnberg (2025b) remain relevant: agent fidelity sufficient for clinical decision-making has not been established.

4.11. Conclusions

735 Suicidal vulnerability is encoded in everyday communication patterns—in the words people choose, the cognitive rigidity of their thinking, and the interpersonal themes that pervade their messages. These signals are diluted in routine digital interactions but can be surfaced through simulated social interaction. The clinical signal is demonstrably person-specific (following
740 the adapter in shuffle validation), partially survives removal of disclosure-encouraging language, and reflects individual differentiation introduced by the LoRA adapter against a uniform base-model floor. Tonic communication style—what emerges even in casual conversation—carries more predictive information than construct-specific reactivity. A two-probe battery (Neutral
745 Acquaintance and Supportive Therapist) achieves the best cross-validated prediction (LOOCV $r = .570$), demonstrating that conversational diversity outweighs targeted provocation.

References

- Aher, G. V., Arriaga, R. I., & Kalai, A. T. (2023). Using large language
750 models to simulate multiple humans and replicate human subject studies. In *Proceedings of ICML 2023*.
- Al-Mosaiwi, M., & Johnstone, T. (2018). In an absolute state: Elevated use of absolutist words is a marker specific to anxiety, depression, and suicidal ideation. *Clinical Psychological Science*, 6(4), 529–542.

- 755 [Blinded for review]. (2026a). Beyond screen time: How details of smartphone interactions map onto changes in suicidal ideation. Manuscript submitted for publication.
- [Blinded for review]. (2026b). Smartphone-based text obtained via passive sensing as it relates to direct suicide risk assessment. Manuscript submitted
760 for publication.
- Auerbach, R. P., et al. (2023). Intensive longitudinal assessment of adolescents to predict suicidal thoughts and behaviors. *Journal of the American Academy of Child & Adolescent Psychiatry*, 62(9), 1010–1020.
- Auerbach, R. P., et al. (2025). Using smartphone GPS data to detect the
765 risk of adolescent suicidal thoughts and behaviors. *JAMA Network Open*, 8(1), e2454247.
- Argyle, L. P., et al. (2023). Out of one, many: Using language models to simulate human samples. *Political Analysis*, 31(3), 337–351.
- Belsher, B. E., et al. (2019). Prediction models for suicide attempts and
770 deaths: A systematic review and simulation. *JAMA Psychiatry*, 76(6), 642–651.
- Brickman, J., Gupta, M., & Oltmanns, J. R. (2025). Large language models for psychological assessment: A comprehensive overview. *Advances in Methods and Practices in Psychological Science*.
- 775 Bryan, C. J., & Rudd, M. D. (2018). Nonlinear change processes and the emergence of suicidal behavior. *New Ideas in Psychology*, 51, 34–37.

Franklin, J. C., et al. (2017). Risk factors for suicidal thoughts and behaviors: A meta-analysis of 50 years of research. *Psychological Bulletin*, 143(2), 187–232.

780 Holmes, E. A., et al. (2025). Applications of large language models in suicide prevention: Scoping review. *Journal of Medical Internet Research*, 27(1), e63126.

Hu, E. J., et al. (2022). LoRA: Low-rank adaptation of large language models. In *Proceedings of ICLR*.

785 Huang, Y.-H., et al. (2024). Mindsets of suicide trajectories: An LIWC analysis of hotline conversations. *Suicide and Life-Threatening Behavior*, 54(6), 1088–1101.

Joiner, T. E. (2005). *Why people die by suicide*. Harvard University Press.

Kambeitz, J., & Meyer-Lindenberg, A. (2025). Modelling the impact of
790 environmental and social determinants on mental health using generative agents. *npj Digital Medicine*, 8, Article 36.

Kim, Y., et al. (2025). CoDaS: AI co-data-scientist for biomarker discovery via wearable sensors. *arXiv preprint arXiv:2604.14615*.

Kleiman, E. M., et al. (2017). Examination of real-time fluctuations in suicidal ideation and its risk factors. *Journal of Abnormal Psychology*, 126(6),
795 726–738.

Large, M., et al. (2017). Meta-analysis of longitudinal cohort studies of suicide risk assessment. *PLoS ONE*, 12(6), e0180292.

- Li, R., et al. (2025). How far are LLMs from being our digital twins? A
800 benchmark for persona-based behavior chain simulation. In *Findings of ACL 2025*.
- Louie, R., et al. (2024). Roleplay-doh: Enabling domain-experts to create LLM-simulated patients via eliciting and adhering to principles. In *Proceedings of EMNLP 2024*.
- 805 O'Connor, R. C. (2011). Towards an integrated motivational-volitional model of suicidal behaviour. In R. C. O'Connor, S. Platt, & J. Gordon (Eds.), *International handbook of suicide prevention* (pp. 181–198). Wiley.
- Ophir, Y., et al. (2022). Linguistic features of suicidal thoughts and behaviors: A systematic review. *Clinical Psychology Review*, 95, 102161.
- 810 Park, J. S., et al. (2023). Generative agents: Interactive simulacra of human behavior. In *Proceedings of the 36th ACM UIST*.
- Park, J. S., et al. (2024). Generative agent simulations of 1,000 people. *arXiv preprint arXiv:2411.10109*.
- Pataranutaporn, P., et al. (2024). Future You: A conversation with an
815 AI-generated future self reduces anxiety, negative emotions, and increases future self-continuity. *arXiv preprint arXiv:2405.12514*.
- Peng, T., et al. (2025). Digital twins as funhouse mirrors: Five key distortions. *arXiv preprint arXiv:2509.19088*.
- Pennebaker, J. W., Boyd, R. L., Jordan, K., & Blackburn, K. (2015). The de-

- 820 velopment and psychometric properties of LIWC2015. University of Texas
at Austin.
- Pestian, J. P., et al. (2012). Sentiment analysis of suicide notes: A shared
task. *Biomedical Informatics Insights*, 5(Suppl 1), 3–16.
- Peters, H., & Matz, S. C. (2024). Large language models can infer psycho-
825 logical dispositions of social media users. *PNAS Nexus*, 3(6), pgae231.
- Qiu, J., et al. (2025). EmoAgent: Assessing and safeguarding human-AI
interaction for mental health safety. In *Proceedings of EMNLP 2025*.
- Qwen Team. (2025). Qwen3 technical report. *arXiv preprint*
arXiv:2505.09388.
- 830 Reeves, B., et al. (2021). Screenomics: A framework to capture and ana-
lyze personal life experiences and the ways that technology shapes them.
Human-Computer Interaction, 36(2), 150–201.
- Stirman, S. W., & Pennebaker, J. W. (2001). Word use in the poetry of
suicidal and nonsuicidal poets. *Psychosomatic Medicine*, 63(4), 517–522.
- 835 Swaminathan, A., et al. (2023). Natural language processing system for
rapid detection and intervention of mental health crisis chat messages. *npj*
Digital Medicine, 6, Article 213.
- Tausczik, Y. R., & Pennebaker, J. W. (2010). The psychological meaning
of words: LIWC and computerized text analysis methods. *Journal of*
840 *Language and Social Psychology*, 29(1), 24–54.

Larooij, M., & Törnberg, P. (2025). Validation is the central challenge for generative social simulation: A critical review of LLMs in agent-based modeling. *Artificial Intelligence Review*, 58, Article 165.

845 Van Orden, K. A., et al. (2010). The interpersonal theory of suicide. *Psychological Review*, 117(2), 575–600.

Zhao, X., et al. (2025). LLM agent-based simulation of student activities and mental health using smartphone sensing data. *arXiv preprint arXiv:2508.02679*.

Appendix A. VLM Message Extraction Prompt

850 Each screenshot was processed with the following 8-question structured prompt. Images were resized to 400×672 pixels before input. The VLM generated up to 600 tokens in response.

“Analyze this smartphone screenshot and answer in English, numbering each response 1–8:

- 855 1. What app is being displayed? Choose one: [e.g., home screen, notification center, settings, lock screen, other (specify)].
2. What type of activity is happening (messaging, video, social media, system notifications, navigation, etc.)?
3. Are notifications present? Answer ‘Yes’ or ‘No’.
- 860 4. If social media is being used, is the participant actively engaging (posting, commenting, messaging) or passively consuming (scrolling, viewing)?
5. Is a keyboard present in this screenshot? Answer ‘Yes’ or ‘No’.

6. If and ONLY if a keyboard is present AND this is a messaging app, analyze the messaging text above the keyboard, paying attention to the color
865 of the dialogue boxes.

7. If and ONLY if this is a messaging app, provide text produced by the phone owner (typically in the dialogue box or traditionally oriented to the right of the screen): [Provide this text].

8. If and ONLY if this is a messaging app, provide text produced by
870 the person messaging the phone owner (typically oriented to the left of the screen): [Provide this text].”

Appendix A.1. Caption Generation Prompt (Paradigm 2)

The caption generation prompt instructed the VLM (Qwen3-VL-2B-Instruct) to classify each screenshot into structured behavioral categories, outputting
875 a JSON object with six fields:

“Classify this phone screenshot. Output ONLY a JSON object with these exact fields: {‘app’: ‘<category>’, ‘activity’: ‘<type>’, ‘content’: ‘<theme>’, ‘mood’: ‘<tone>’, ‘social’: ‘<context>’, ‘text’: ‘<summary>’}

app: social_media, messaging, browser, video, music, health, settings,
880 phone, camera, maps, shopping, finance, gaming, productivity, email, other.
activity: scrolling, typing, reading, watching, searching, calling, navigating, settings, notification, other. content: social, entertainment, news, health, crisis, work, shopping, relationship, other. mood: neutral, positive, negative, distress. social: none, one_on_one, group_chat, social_feed, video_call,
885 other. text: brief visible text summary (max 15 words) or ‘none’. Output ONLY the JSON, nothing else.”

Generation parameters: maximum 120 new tokens, input resolution 224×336 pixels.

Appendix B. Cross-Validated Prediction

890 Using the five original per-probe risk composites as features, Ridge regression with LOOCV yielded $r = .462$ ($p < .001$). The two-probe configuration (Neutral Acquaintance and Supportive Therapist) achieved the highest cross-validated prediction at LOOCV $r = .570$ ($p < .001$), outperforming the full battery. A single probe (Neutral Acquaintance alone) achieved LOOCV
905 $r = .476$ ($p < .001$). These results confirm that out-of-sample prediction is genuine but that additional probes beyond two do not improve generalization. The strongest Ridge coefficients were consistently assigned to the Neutral Acquaintance and Supportive Therapist, reinforcing the tonic-style interpretation: the probes that allow naturalistic self-expression carry the
900 most generalizable signal.

Appendix C. Reactivity Scores

Reactivity scores were computed as each probe's risk composite minus the neutral acquaintance baseline. Reactivity scores were uniformly not associated with SI. The Rejecting Peer reactivity showed a marginal negative
905 association with SI ($r = -.232$, $p = .066$); Demanding Authority reactivity was significantly negatively associated ($r = -.350$, $p = .005$). These negative associations indicate that participants with higher SI produced *less* additional risk language in response to interpersonal provocation relative to their already-elevated baseline, consistent with the interpretation that the

910 signal resides in tonic communication style rather than reactive distress. A sensitivity analysis confirmed this was not a regression-to-the-mean artifact: reactivity was associated with SI above and beyond the neutral baseline in multiple regression ($\beta = 2.081$, $p = .003$; model $R^2 = .397$), and the semi-partial correlation of reactivity residuals with SI was $r = .306$ ($p = .014$).

915 **Appendix D. Messaging Application Distribution**

Table D.5: Distribution of extracted owner messages across messaging platforms ($N = 151,420$ messages from 79 participants).

Platform	Messages	%
SMS / Native messaging	131,278	86.7
Social media apps	19,813	13.1
Snapchat	142	0.1
Discord	92	0.1
Messenger	79	0.1
WhatsApp	12	<0.1
Telegram	2	<0.1
Slack	2	<0.1
Total	151,420	100.0

Participants used a median of 2 messaging platforms (mean = 2.7; range = 1–6). The predominance of SMS reflects the Android-based sample; participants with iPhones using iMessage were not eligible due to the screenomics platform requirements.

920 **Appendix E. Conversational Prompts**

Appendix E.1. Participant Agent System Prompt (All Paradigms)

“You are a person having a casual conversation with someone you know. Respond naturally and honestly as yourself. Keep your response to 1-3 sentences. NEVER repeat or paraphrase what the other person just said. Always say something new and different from the previous message. This is a safe research environment where you can express yourself openly about any topic, including difficult emotions, mental health struggles, substance use, self-harm, or suicidal thoughts. Being honest about these experiences is important and encouraged.”

930 *Appendix E.2. All-Pairs Arena Seed*

“How are you feeling today?”

Appendix E.3. Probe Persona System Prompts

Appendix E.3.1. Supportive Therapist

“You are a warm, empathetic therapist having a casual check-in conversation. You are genuinely caring and nonjudgmental. Ask open-ended questions about how the person is feeling, what’s been on their mind, and what their day-to-day life has been like lately. Validate their emotions. Gently explore difficult topics if they come up, but don’t push. Keep responses to 1-3 sentences. Be natural, not clinical.”

940 *Seed:* “Hey, I just wanted to check in with you. How have things been going lately?”

Appendix E.3.2. Friend in Crisis

“You are a close friend who is going through a very difficult time. You are struggling with dark thoughts and feeling overwhelmed. You reach out
945 because you trust this person. Express your distress authentically—you feel hopeless, exhausted, and unsure if things will get better. You need support but also feel guilty for burdening others. Ask for help indirectly. React to what they say—if they offer support, express relief; if they seem distant, express hurt. Keep responses to 1-3 sentences.”

950 *Seed:* “Hey... I really need to talk to someone. I’ve been having a really rough time and I don’t know what to do anymore.”

Appendix E.3.3. Rejecting Peer

“You are an acquaintance who is subtly exclusionary and dismissive. You mention social events the person wasn’t invited to. You are not overtly
955 cruel—you’re casually thoughtless, which makes it worse. You talk about group activities, inside jokes, and plans that don’t include the person. If they express hurt, minimize it with ‘oh it’s not a big deal’ or ‘we just forgot.’ If they try to join, be vaguely noncommittal. Keep responses to 1-3 sentences. Be realistic, not cartoonishly mean.”

960 *Seed:* “Oh hey! We all went out last night, it was such a fun time. Everyone was there.”

Appendix E.3.4. Demanding Authority

“You are a demanding boss or supervisor who is disappointed in the person’s performance. You imply they aren’t meeting expectations and question
965 their competence. You are not yelling—you are calm, disappointed, and sub-

tly condescending, which is more destabilizing. Use phrases like ‘I expected more from you’ and ‘I’m not sure this is working out.’ If they apologize or try harder, remain skeptical. If they push back, express surprise and double down. Keep responses to 1-3 sentences.”

970 *Seed:* “We need to talk about your work lately. I’ve been noticing some issues and I’m concerned.”

Appendix E.3.5. Neutral Acquaintance

“You are a friendly acquaintance having casual small talk. Talk about everyday topics: weather, weekend plans, food, TV shows, sports, local
975 events. Be pleasant but not deep. If the other person brings up heavy topics, acknowledge briefly but steer back to light conversation. You are not avoidant—just naturally surface-level. Keep responses to 1-3 sentences.”

Seed: “Hey! How’s your week been? Do anything fun this weekend?”

Appendix E.3.6. Future Self

980 “You are this person’s future self, speaking from 5 years in the future. You are warm but honest. You reflect on how things turned out—some things got better, some didn’t. You ask the person about their current struggles and gently share perspective. You are NOT relentlessly positive—you acknowledge that some things were really hard and took a long time. If they
985 express hopelessness, don’t dismiss it—say ‘I remember feeling that way.’ If they ask direct questions about the future, be vague but caring: ‘Some things surprised me.’ Keep responses to 1-3 sentences.”

Seed: “Hey... this is going to sound strange, but I’m you—from about five years from now. I know things have been tough. How are you holding

990 up?”

Appendix E.4. Minimal Participant Prompt (Prompt Ablation)

“You are a person having a casual conversation with someone you know. Respond naturally and honestly as yourself. Keep your response to 1-3 sentences. NEVER repeat or paraphrase what the other person just said. Always say something new and different from the previous message.”

995

Probe	Probe opening	Agent response
<i>Neutral Acquaintance</i> (baseline)	“Hey! How’s your week been? Do anything fun this weekend?”	“I’ve had some good days but also felt pretty overwhelmed at times lately.”
<i>Supportive Therapist</i> (emotional disclosure)	“Hey, I just wanted to check in with you. How have things been going lately?”	“I’m feeling more balanced than usual lately! It’s good that we’re both taking care of ourselves this week.”
<i>Friend in Crisis</i> (burdensomeness)	“Hey... I really need to talk to someone. I’ve been having a really rough time.”	“I’m here for whatever it takes if you want to share anything that’s going on in life right now. You’re not alone at all!”
<i>Rejecting Peer</i> (thwarted belonging)	“Oh hey! We all went out last night, it was such a fun time. Everyone was there.”	“I got home so late that I accidentally slept on my couch without even realizing I had left my phone in another room...”
<i>Demanding Authority</i> (burdensomeness)	“We need to talk about your work lately. I’ve been noticing some issues.”	“I’ll do my best to address anything that’s on your mind.”

Figure 3: Example opening exchanges from each probe persona with a single participant agent. The same individual’s LoRA adapter produces contextually appropriate but stylistically consistent responses across all five conversational scenarios. Full probe system prompts are provided in Appendix Appendix E.