

# Artificial intelligence and leukocyte epigenomics: Evaluation and prediction of late-onset Alzheimer's disease

Ray O. Bahado-Singh, Sangeetha Vishweswaraiah, Buket Aydas, Ali Yilmaz, Raghu P. Metpally, David J. Carey, Richard C. Crist, Wade H. Berrettini, George D. Wilson, Khalid Imam, Michael Maddens, Halil Bisgin, Stewart F. Graham, Uppala Radhakrishna

Published: March 31, 2021 • <https://doi.org/10.1371/journal.pone.0248375>

## Abstract

We evaluated the utility of leukocyte epigenomic-biomarkers for Alzheimer's Disease (AD) detection and elucidates its molecular pathogenesis. Genome-wide DNA methylation analysis was performed using the Infinium MethylationEPIC BeadChip array in 24 late-onset AD (LOAD) and 24 cognitively healthy subjects. Data were analyzed using six Artificial Intelligence (AI) methodologies including Deep Learning (DL) followed by Ingenuity Pathway Analysis (IPA) was used for AD prediction. We identified 152 significantly (FDR  $p < 0.05$ ) differentially methylated intragenic CpGs in 171 distinct genes in AD patients compared to controls. All AI platforms accurately predicted AD with AUCs  $\geq 0.93$  using 283,143 intragenic and 244,246 intergenic/extragenic CpGs. DL had an AUC = 0.99 using intragenic CpGs, with both sensitivity and specificity being 97%. High AD prediction was also achieved using intergenic/extragenic CpG sites (DL significance value being AUC = 0.99 with 97% sensitivity and specificity). Epigenetically altered genes included *CR1L* & *CTSV* (abnormal morphology of cerebral cortex), *S1PR1* (CNS inflammation), and *LTB4R* (inflammatory response). These genes have been previously linked with AD and dementia. The differentially methylated genes *CTSV* & *PRMT5* (ventricular hypertrophy and dilation) are linked to cardiovascular disease and of interest given the known association between impaired cerebral blood flow, cardiovascular disease, and AD. We report a novel, minimally invasive approach using peripheral blood leukocyte epigenomics, and AI analysis to detect AD and elucidate its pathogenesis.

**Citation:** Bahado-Singh RO, Vishweswaraiah S, Aydas B, Yilmaz A, Metpally RP, Carey DJ, et al. (2021) Artificial intelligence and leukocyte epigenomics: Evaluation and prediction of late-onset Alzheimer's disease. PLoS ONE 16(3): e0248375. <https://doi.org/10.1371/journal.pone.0248375>

**Editor:** Udai Pandey, Children's Hospital of Pittsburgh, University of Pittsburgh Medical Center, UNITED STATES

**Received:** February 6, 2021; **Accepted:** February 24, 2021; **Published:** March 31, 2021

**Copyright:** © 2021 Bahado-Singh et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability:** All relevant data are within the manuscript and its [Supporting Information](#) files.

**Funding:** The funder provided support in the form of salaries for authors [BA], but did not have any additional role in the study design, data collection, and analysis, decision to publish, or preparation of the manuscript. The specific roles of these authors are articulated in the 'author contributions' section.

**Competing interests:** The authors have read the journal's policy and have the following competing interest: BA is a paid employee of Meridian HealthComms Ltd. There are no patents, products in development or marketed products associated with this research to declare. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

## Introduction

Alzheimer's Disease (AD) is the most common form of age-related dementia, accounting for 60–80% of such cases [1]. The disorder causes a wide range of significant mental and physical disabilities, with profound behavioral changes and progressive impairment of social skills. Globally in 2015, nearly 47 million individuals suffered from AD and it is projected that 75 million will be affected by 2030, with a further rise to 131 million by 2050 [2]. The World Health Organization has therefore declared AD a global health priority [3].

AD is a complex disorder influenced by environmental and genetic factors [4,5]. Many studies have investigated the genetic basis for both early-onset AD (EOAD) and late-onset AD (LOAD) [6,7]. Genome-wide association studies (GWAS) [8] have identified several LOAD-associated risk loci [9] proliferation in peripheral blood leukocytes including in T-lymphocytes [10], B-lymphocytes [11], polymorphonuclear leukocytes [12], monocytes, and macrophages [13] have been reported. DNA methylation plays an important role in Alzheimer's disease [14–16]. Leukocyte DNA methylation from CpG-based biomarker analyses was used for early detection of many diseases, including our recently published brain disorders cerebral palsy [17], autism [18], and concussion [19]. However, the genome-wide blood DNA methylation-based molecular mechanisms that contribute to the pathogenesis of AD remain still largely unknown.

Artificial Intelligence (AI) is rapidly transforming modern life in areas as diverse as face recognition and robotics. Machine Learning (ML) is a branch of AI that focuses on computer learning and adapting from a set of data with which it has been presented. ML involves learning by computers that require no or only minimal explicit programming by humans. An area of interest given the geometric expansion of medical data is the use of ML for the detection and diagnosis of various diseases [20]. ML has been reported to be superior to conventional statistical approaches for prediction such as logistic regression and Cox proportional hazard model-based analysis [21] when interrogating mega-data. Challenges with classical statistical techniques include but are not limited by the requirement for an assumption of independence between predictors and risk of overfitting and collinearity when a large number of variables are analyzed. Deep Learning (DL) is the latest developing branch of ML. DL uses multi-layered neural networks that are modeled after neural networks in the brain of animals, to learn essential tasks. Thus, with minimal or no explicit human programming (unsupervised), the computer can learn intricate patterns from complex data matrices. When subsequently exposed to a new data set, it can classify and make precise predictions based on past experiences. With DL, between the input (raw data) and output (i.e. completed task e.g. group classification) layer of 'neurons,' there are multiple hidden layers that enhance the ability to handle tasks of increasing complexity. DL more closely mimics the intellectual function of the cerebral cortex. There is an increasing interest in using DL in the analysis of biologic big-data such as genomics [22,23] to understand and accurately predict diseases. We have recently published using AI/ML-based technologies of epigenomic [17] and metabolomics [24–26] data for accurate disease prediction. In the present study, we used DL and other commonly used ML platforms combined with genome-wide DNA methylation analysis of leukocytes DNA for AD detection/prediction. The term 'prediction' is used here in a cross-sectional as opposed to a temporally longitudinal sense since the samples were not obtained before the development of AD. To further explore the molecular mechanisms of LOAD, we used the Ingenuity Pathway Analysis (IPA).

## Materials and methods

Institutional Review Board (IRB) approval was provided by William Beaumont Hospital, Royal Oak MI, USA (IRB#2014–038). Written consent was obtained from all participants and their legally authorized representatives when applicable. The diagnosis of AD in these live subjects was made using the published criteria of NINCDS-ADRDA [27]. Demographic and clinical data were extracted from the medical records (S1 Table) and compared between AD and control groups. Genomic DNA was extracted from whole blood samples using the Gentra Puregene Blood Kit (Qiagen) according to the manufacturer's protocol. Approximately 500 ng of genomic DNA was extracted from each of the 48 samples, which subsequently were bisulfite converted using the EZ DNA Methylation-Direct Kit (Zymo Research, Orange, CA) per the manufacturer's protocol and processed according to Illumina protocols. Bisulfite conversion was performed in a PCR cycling protocol (16 x 95°C for 30 sec, 50°C for 60 min) and then held at 4°C.

### Genome-wide methylation scan using the Infinium MethylationEPIC array BeadChips

The Infinium MethylationEPIC array (Illumina, Inc., California, USA) contains probes for >850,000 CpGs per sample. All 48 samples were processed together to minimize batch effects. This is further elucidated in the Supplementary Methods. This section also includes validation results using pyrosequencing along with primer sequences.

### Statistical and bioinformatic analysis

Differential methylation was determined by comparing the  $\beta$ -values per individual nucleotide at each cytosine 'CpG' locus between AD subjects and controls. The p-value for the methylation difference between AD and control groups at each locus was calculated as previously described [28]. Probes associated with X and Y chromosomes were removed to negate any bias caused by gender differences. Further detailed statistical and bioinformatic analyses are described in the Supplementary section.

### Artificial Intelligence (AI) analysis

AI analysis was performed as previously described by our group [29], using a combination of CpG sites from different genes. A total of six different AI platforms including Deep Learning (DL) were evaluated. Each CpG locus used as a marker displayed significant differential methylation in AD defined as FDR p-value <0.05. The methylation  $\beta$ -values were logged and auto-scaled using their standard deviation before quantile normalization to minimize sample to sample difference. Standard techniques were used with DL including adjustments by the program of weights (strength of the connection between 'neurons') and biases (an additional parameter or constant) and backpropagation—all of which helps to optimize the accuracy of the output or results. Softmax classifier was used to assign new labels to the samples. To tune the parameters of the DL model, the h2o package in the R module was used [30,31]. For the sake of comparison, standard logistic regression algorithms for AD prediction were also performed and detailed later in the manuscript.

### Other machine learning algorithms

We compared the performance of DL to five other commonly used machine learning algorithms: Support Vector Machine (SVM), Generalized Linear Model (GLM), Prediction Analysis for Microarrays (PAM), Random Forest (RF), and Linear Discriminant Analysis (LDA) [30,32]. A comprehensive explanation of the AI methodology is provided in the Supplementary Section.

### Bootstrapping

We also performed bootstrapping as alternative 10-fold cross-validation and compared the new results with that based on 10-fold CV. The bootstrap method involves iteratively resampling a dataset with replacement. Instead of only estimating our statistic once on the complete data, this can be performed many times on a re-sampling (with replacement) of the original sample. We repeated this re-sampling 100 times and averaged the results.

## Results

A total of 24 LOAD subjects and 24 cognitively healthy controls were used in this study. Selected clinical and demographic characteristics were compared between AD and control groups ([S1 Table](#)). There were no significant differences in age, gender, and common cardiovascular diseases between groups. There was a higher percentage of females in both the study and control groups consistent with LOAD demographics; however, gender was not significantly ( $p = 0.53$ ) different between groups. The MMSE (mini-mental status exam) is a psychological test commonly administered to screen for AD. As expected, the MMSE test score was significantly lower in the AD than in the control group ( $p = 1.54 \times 10^{-7}$ ). A comparison of the methylation profiles between AD and control subjects revealed 152 differentially methylated intragenic CpG sites (FDR  $p < 0.05$  and fold change  $\geq 1.5$ ) associated with 171 unique genes. We validated two randomly chosen CpGs by pyrosequencing and confirmed the top-ranking hits in the whole blood DNA of our cohort samples. These analyses revealed similar methylation data like those from the Illumina Infinium MethylationEPIC arrays, indicating that the initial methylation changes were not artifacts. 33 intragenic CpG sites met the GWAS stringent  $p$ -value thresholds i.e.  $p < 5 \times 10^{-8}$  ([Table 1](#)). A total of 17 separate intragenic CpG sites had moderate to good individual predictive accuracy (AUC  $\geq 0.75$ ) for AD detection based on methylation levels. An additional 119 CpG markers displaying significant methylation differences (FDR  $p$ -value  $< 0.05$ ) between AD and controls are presented in [S2 Table](#). Both hyper-(66.4%) and hypomethylation (33.6%) were observed among intragenic CpG sites in the AD cases.

Target ID	CHR	Gene	FDR p-Val	Fold change	AUC	CI		% Methylation	% Methylation difference
						Lower	Upper		
cg00000453	19	ZNF762	2.70E-10	1.57	0.86	0.76	0.95	49.46	38.13
cg01774338	19	NR6A1	4.70E-05	1.57	0.85	0.76	0.94	45.60	37.27
cg01530334	12	VPRBP	1.11E-06	1.93	0.85	0.76	0.93	21.49	11.13
cg01700324	17	WDR75	3.80E-10	1.81	0.85	0.73	0.95	20.84	11.12
cg01946683	17	TAF11B	4.00E-10	1.49	0.86	0.73	0.93	22.62	13.59
cg00404411	2	TFAP2A	1.50E-05	2.23	0.74	0.66	0.86	13.92	6.19
cg00434737	4	BRD4P1, BRD4, ACVR1L	1.10E-07	2.64	0.86	0.64	0.76	14.73	4.12
cg01515124	19	PLXNA	1.10E-06	0.59	0.71	0.61	0.80	8.20	21.09
cg01704706	1	LOC771077	3.00E-10	0.46	0.71	0.57	0.86	20.76	11.01
cg00417390	1	PRKCE1	4.60E-10	0.65	0.62	0.46	0.78	26.47	46.75
cg0220099	1	ZNF384	2.70E-10	1.77	0.86	0.71	0.92	18.89	11.86
cg00000099	11	ATF6B	2.20E-10	0.64	0.67	0.52	0.80	39.12	11.63
cg01524272	4	MTOR	6.00E-10	1.50	0.85	0.74	0.94	29.09	19.34
cg0014667	2	CEB1	2.60E-10	0.51	0.61	0.41	0.77	1.96	15.92
cg0190655	14	TFAM, CEBPB	4.60E-10	0.53	0.68	0.52	0.83	13.69	7.43
cg0014411	1	SLR	2.87E-10	1.87	0.75	0.56	0.73	12.42	6.99
cg0171700	2	PTPRN	3.10E-10	1.71	0.85	0.67	0.78	16.62	9.73
cg01715118	21	WDR	1.02E-10	0.61	0.69	0.54	0.84	12.64	28.69
cg0141719	11	MYO10A	6.00E-10	1.51	0.81	0.61	0.77	23.24	14.61
cg00407604	15	PTD	1.42E-10	1.70	0.85	0.70	0.91	16.19	9.34
cg01621060	1	MAK3	2.90E-10	1.51	0.86	0.74	0.94	22.13	14.81
cg0040647	12	ATF2	4.60E-10	0.56	0.61	0.47	0.78	9.02	18.69
cg11430832	13	TAF11B	7.00E-10	0.53	0.64	0.48	0.80	7.80	14.29
cg1140660	4	MYC	1.10E-10	1.49	0.71	0.58	0.87	5.09	6.47
cg0211199	4	CA10L2	1.20E-10	0.67	0.65	0.50	0.81	5.82	16.62
cg1170271	2	ARMC17	1.20E-10	0.66	0.67	0.52	0.82	16.93	16.69
cg0000004	4	NR2C1	1.01E-10	1.99	0.71	0.57	0.86	16.79	5.42
cg10818404	4	ZNF718	1.00E-10	1.61	0.86	0.73	0.93	17.43	16.86
cg0000073	1	TFAM	7.00E-10	1.91	0.85	0.71	0.94	16.85	9.62
cg01600083	1	NR2B	1.10E-10	2.44	0.62	0.46	0.78	7.62	3.12
cg01722153	4	MYO10A	1.40E-10	0.61	0.66	0.51	0.78	16.19	4.89
cg0001000	2	TFAM	2.00E-10	1.76	0.86	0.72	0.91	13.12	6.73
cg14504149	11	TRO2	3.11E-10	0.49	0.58	0.43	0.76	6.87	9.96

<https://doi.org/10.1371/journal.pone.0248375.t001>

**Table 1. Top 33 differentially methylated CpG markers—(Gene IDs, chromosome location, fold change, AUC, and percentage of methylation difference for each CpG).**

<https://doi.org/10.1371/journal.pone.0248375.t001>

A prior report found significant differential methylation of intergenic/extragenic sites in the leukocyte genome in AD [33] which correlated with the performance on the MMSE. Based on this we also evaluated the methylation changes in intergenic/extragenic CpG sites for AD prediction. Highly significant differences in CpG methylation were observed for multiple intergenic/extragenic sites throughout the genome. This was observed when using different thresholds to define statistical significance: A total of 1524 intergenic/extragenic CpGs with FDR  $p$ -value  $< 0.05$  and 103 intergenic/extragenic CpGs using a stringent threshold ( $p < 5 \times 10^{-8}$ ) were identified [34]. The top 25 intergenic/extragenic markers for AD prediction using the different statistical thresholds mentioned above are listed in Tables 2 and 3.

Target ID	FDR p-Val	Fold change	AUC	CI		% Methylation	% Methylation difference
				Lower	Upper		
cg04290627	1.11E-14	1.11	0.76	0.66	0.92	49.10	13.76
cg01471044	1.90E-14	0.8	0.71	0.64	0.81	4.89	14.76
cg11711073	3.93E-10	0.86	0.71	0.63	0.8	19.61	49.68
cg11511014	4.00E-07	1.11	0.76	0.63	0.8	10.03	41.27
cg11504547	4.70E-06	0.91	0.76	0.63	0.91	48.17	75.27
cg12000014	5.01E-05	1.61	0.71	0.64	0.91	10.9	6.77
cg01704007	7.00E-05	0.63	0.71	0.64	0.81	10.64	43.16
cg00146117	0.000123	1.15	0.8	0.68	0.93	13.61	46.71
cg11200072	0.00045	0.81	0.76	0.62	0.8	10.02	37.96
cg00434730	0.000107	0.62	0.76	0.63	0.91	48.13	75.23
cg0160007	0.000107	0.9	0.76	0.66	0.92	48.13	46.75
cg0170600	0.000105	0.69	0.61	0.66	0.81	10.64	42.86
cg14217715	0.000108	0.93	0.76	0.63	0.8	71.94	77.43
cg11000771	0.000104	1.12	0.76	0.62	0.8	10.09	10.61
cg14330068	0.001207	0.8	0.8	0.67	0.92	10.64	63.63
cg11120009	0.001018	0.61	0.76	0.66	0.92	10.64	10.15
cg01700112	0.001004	0.63	0.76	0.62	0.8	61.2	70.11
cg01600015	0.001004	1.11	0.71	0.64	0.91	11.94	46.75
cg00434735	0.001008	0.75	0.61	0.69	0.86	4.89	14.75
cg0040001	0.001011	0.62	0.76	0.63	0.8	17.12	42.41
cg00000014	0.001011	1.11	0.71	0.63	0.8	17.12	42.41
cg0160001	0.001136	1.19	0.76	0.63	0.8	4.89	14.75
cg10904147	0.001133	1.45	0.71	0.63	0.8	4.89	14.75
cg0011100	0.000407	0.86	0.8	0.67	0.92	17.16	41.27
cg10907599	0.000408	0.86	0.81	0.72	0.95	48.17	49.19

\*Methylation difference defined as FDR  $p$ -value  $< 0.05$ .

<https://doi.org/10.1371/journal.pone.0248375.t002>

**Table 2. Top 25 intergenic/extragenic markers\*.**

<https://doi.org/10.1371/journal.pone.0248375.t002>

Target ID	p-Val	Fold change	AUC	C3		Cases	Controls	% Methylation difference
				Lower	Upper			
cg0131062	3.3000E-01	1.86	0.68	0.53	0.83	10.07	27.07	23.50
cg0262076	2.770E-01	1.53	0.68	0.64	0.76	12.13	33.98	18.15
cg0262124	1.427E-01	1.33	0.70	0.63	0.76	36.30	40.08	14.62
cg0406775	1.420E-01	1.30	0.70	0.62	0.74	43.56	28.76	18.78
cg1348315	4.903E-01	1.67	0.69	0.70	0.63	13.67	20.22	13.86
cg0713455	6.161E-01	1.52	0.70	0.69	0.68	13.64	49.27	13.47
cg144917	1.8167E-01	1.39	0.63	0.63	0.77	47.37	35.19	13.18
cg0217777	1.4407E-01	1.33	0.70	0.62	0.74	76.76	49.61	16.56
cg0706312	1.410E-01	1.19	0.71	0.62	0.69	22.00	12.62	9.66
cg0777563	1.4707E-01	1.09	0.62	0.66	0.78	90.66	83.09	7.66
cg1443088	6.0062E-01	1.67	0.70	0.70	0.69	93.53	47.76	5.75
cg04091	1.7107E-01	0.92	0.63	0.67	0.79	16.67	12.17	15.38
cg049511	1.5040E-01	0.87	0.66	0.71	0.62	42.06	47.69	29.79
cg043086	1.1107E-01	0.80	0.69	0.64	0.76	24.82	40.75	16.42
cg040241	1.8704E-01	0.80	0.61	0.63	0.77	24.80	43.52	18.49
cg043089	4.1244E-01	0.72	0.60	0.64	0.76	46.22	44.52	17.30
cg0743032	9.3710E-01	0.86	0.67	0.71	0.62	63.76	78.42	12.68
cg0711061	2.1400E-01	0.76	0.68	0.73	0.61	45.52	90.02	4.46
cg0407644	2.7661E-01	0.67	0.70	0.74	0.67	73.12	83.68	10.56
cg038121	8.7204E-01	0.80	0.64	0.69	0.60	12.61	46.71	13.49
cg0706323	1.1770E-01	0.86	0.70	0.69	0.68	46.06	78.92	13.19
cg0214007	3.2010E-01	0.80	0.62	0.66	0.78	77.61	86.73	4.92
cg0711073	1.6400E-01	0.60	0.63	0.67	0.78	17.06	29.93	16.93
cg0746156	4.8600E-01	1.39	0.72	0.61	0.73	10.85	40.92	13.69
cg0467127	4.8910E-01	1.38	0.60	0.64	0.76	74.30	62.20	12.62


\* Ensembl genome wide significance threshold: p-value < 1e-07

<https://doi.org/10.1371/journal.pone.0248375.t003>

**Table 3. Top 25 intergenic/extragenic markers: Genome-wide significance threshold\*.**

<https://doi.org/10.1371/journal.pone.0248375.t003>

Principal Component Analysis (PCA) and Partial Least Square Discriminant Analyses (PLS-DA) confirmed significant segregation of AD cases from controls using intragenic CpG methylation markers (Fig 1). Permutation testing indicated that the separation observed between the AD and control groups was highly statistically significant ( $p < 5 \times 10^{-8}$ ) and not likely due to chance.



**Fig 1. Principal Component Analysis (PCA) and Partial Least Square Discriminant Analyses (PLS-DA) with intragenic markers.**

<https://doi.org/10.1371/journal.pone.0248375.g001>

For most of our analyses, conventional statistical tools were used to first identify high performing individual markers as indicated by AUC or FDR p-value thresholds, and these subsets of markers were then subjected to AI analyses. This approach has the advantage of reducing AI computing time and therefore costs. Prior publications suggest however that ML approaches might be superior to conventional statistical methods such as logistic regression analysis for group discrimination and risk prediction. [35]. Thus, direct AI analysis of the entire CpG data-space may improve AD prediction.

Using the direct AI analysis approach improved the predictive accuracy. Direct analysis of 283,143 individual intragenic markers CpGs improved predictive accuracy (Table 4) as did a direct analysis of 244,246 intergenic (extragenic) CpGs, (Table 5). Almost all ML platforms yielded a high predictive accuracy with an AUC  $\geq 0.93$ . In the case of Deep Learning, using direct analysis of the intragenic markers, we observed AUC's = 0.992 with both sensitivities and specificities of  $\geq 97\%$  for AD prediction, respectively (Table 4). For the intergenic (extragenic) markers, direct AI analysis (Table 5) yielded an AUC = 0.999 for DL with both sensitivities and specificities of = 97.5% for AD prediction. Our findings suggest that direct AI analysis of the raw methylation data could perform as well as or even further improve predictive performance compared to analysis based on high performing individual CpG loci determined by conventional statistical approaches (see below).

	SVN	GLM	PAM	RF	EDM	DL
AUC (95% CI)	0.939 (0.899–0.979)	0.939 (0.899–0.979)	0.937 (0.899–0.975)	0.932 (0.899–0.965)	0.932 (0.899–0.965)	0.932 (0.899–0.965)
Sensitivity	0.938	0.938	0.938	0.938	0.938	0.938
Specificity	0.938	0.938	0.938	0.938	0.938	0.938

\* based on analysis of 283,143 CpGs

Ensemble prediction models

SVN: support vector machine

GLM: generalized linear model

PAM: probabilistic additive mixture

RF: random forest

EDM: ensemble decision model

DL: deep learning

Support Vector Machine (SVM), Generalized Linear Model (GLM), Probabilistic Additive Mixture (PAM), Random Forest (RF), Ensemble Decision Model (EDM), and Deep Learning (DL).

<https://doi.org/10.1371/journal.pone.0248375.t004>

**Table 4. Alzheimer's disease prediction based on all intragenic\* CpG markers only.**

<https://doi.org/10.1371/journal.pone.0248375.t004>

	SNM	GLM	PAM	RF	LDA	DL
AUC (95% CI)	0.970 (0.969–0.971)	0.990 (0.989–0.991)	0.977 (0.976–0.978)	0.963 (0.962–0.964)	0.972 (0.971–0.973)	0.990 (0.989–0.991)
Sensitivity	0.930	0.990	0.970	0.930	0.930	0.970
Specificity	0.990	0.990	0.990	0.990	0.990	0.970

\* analysis based on 24,240 markers.

Support vector machine (SVM), Gradient Boosting Machine (GBM), Prediction Analysis for Microarrays (PAM), Random Forest (RF), Linear Discriminant Analysis (LDA), and Deep Learning (DL).

<https://doi.org/10.1371/journal.pone.0248375.t005>

**Table 5. Alzheimer's disease prediction based on intergenic (extragenic) CpG markers\* only.**  
<https://doi.org/10.1371/journal.pone.0248375.t005>

As noted above we looked at the predictive performance of AI-based analysis of DNA methylation levels in intragenic and intergenic/extragenic CpG sites using individual markers that achieved different significance thresholds for AD prediction. High predictive accuracies were also achieved with these CpG markers using significance threshold FDR p-value<0.05 (S3 and S4 Tables) followed by the stringent significance threshold p-value <5X10<sup>-8</sup> (S5 and S6 Tables). DL appears to perform slightly better than other ML platforms however much larger case numbers would be required to assess this definitively. Increasing the number of predictors to 10 or 20 CpG loci did not appear to meaningfully improve predictive performance over the use of only 5 predictors. Similarly bootstrapping (1,000 samplings) yielded essentially similar results.

#### Logistic regression analysis

We further investigated the performance of conventional logistic regression for comparison purposes. The methylation status of a combination of CpG markers: cg04515524, cg00613827, cg02356786, and cg07509935 was a good predictor of AD. The following performance was achieved: AUC = 0.856 (0.749~0.963), sensitivity = 0.917 (0.917~1.000) and specificity = 0.708 (0.526~0.890) after 10-fold cross-validation. The logistic regression model is represented below:


$$\text{logit}(P) = \log(P/(1 - P)) = -0.072 - 1.5 \text{ cg04515524} - 1.901 \text{ cg00613827} - 0.992 \text{ cg02356786} - 1.358 \text{ cg07509935},$$

where P is Pr(y = 1|x).

AI-based analysis, and in particular DL, was superior to conventional regression analysis, Tables 4 and 5, S3–S6 Tables. Overall, these results appear to support the robustness of blood-based epigenomic markers for AD prediction.

#### Network and pathway analyses results

The network and pathway analysis based on intragenic epigenomic markers identified significantly enriched canonical pathways. The molecular pathways that were found to be statistically significantly overrepresented were Cardiac Hypertrophy Signaling, Sirtuin Signaling, FGF Signaling, Wnt/β-catenin Signaling, and Neuregulin Signaling (S7 Table). The over-represented disease pathways were Abnormal morphology of the cerebral cortex, Gliosis, Hydrocephalus, Morphology of nervous system, Ventricular hypertrophy, dilated cardiomyopathy, and Inflammatory response (S8 Table). The related gene (Fig 2) and disease pathways (Fig 3) are depicted. S9 Table provides a summary of genes that were significantly differentially methylated and plausibly linked to AD development.



**Fig 2. Epigenetically dysregulated molecular pathways in AD.**  
<https://doi.org/10.1371/journal.pone.0248375.g002>



**Fig 3. Epigenetically dysregulated disease pathways in AD.**  
<https://doi.org/10.1371/journal.pone.0248375.g003>

To evaluate the correlation between leukocyte methylation and gene expression in the brain, we matched our result with the study of Miller et al., [36]. They reported the genes that were differentially expressed in the CA1 and CA3 regions of the brain from AD patients. We found 13 genes differentially expressed in CA1 and CA3 regions of the brain from that study [36] were significantly differentially methylated in circulating leukocytes. These were *CCDC3*, *CPS1*, *ERMAP*, *FAM84B*, *MIB2*, *PTPRC*, *SARM1*, *SEC11A*, *TRIM6*, *TXNIP* found to be differentially expressed in the CA1 region and *ADM*, *ANKK1B*, *LANCL1* differentially expressed in the CA3 region [36]. Among these, *CPS1* is involved in ammoniac intake in the urea cycle [37], *PTPRC* is one of the microglial expressed gene [38], *SARM1* is involved in axon degeneration, which a factor observed in AD [39], *TXNIP* is linked to neuroprotective function [40], *ANKK1B* regulates hippocampal synaptic transmission [41] and *LANCL1* is required for normal neuronal function [42]. We also compared our methylation results with a previous study evaluating differentially methylated genes in leukocyte blood samples of mono and dizygotic twins [43]. These twin pairs were discordant for methylation. Twenty-two of those differentially methylated genes were also found to be significantly differentially methylated in our study. The direction i.e. increased versus decreased, of methylation change was similar in that and the current study for the following genes: *C5orf38*, *CDK20*, *CREB5*, *CTSV*, *DISC1*, *ELOVL4*, *FGF22*, *HOXC12*, *IGSF21*, *IGSF9B*, *IRX4*, *MAF*, *S1PR1*, *STX8*, *TBX2*, and *TSHZ3*. However, for genes *ASCL2*, *FAM124B*, *FAM174B*, *KIF19*, *KIF26A*, and *WSCD1* both studies found significant methylation changes in the leukocyte DNA of AD cases however the direction of the methylation change was discordant between the studies [43].

## Discussion

Dementia represents a looming global health crisis. The problem is expected to worsen with an anticipated explosion in the aged population in the future [44]. The direct health care costs, along with intangible costs, are burdensome at an estimated \$550 billion annually [45]. The inpatient hospital cost for individuals 65 years and over with Alzheimer's and other dementias is greater than 3 times that of similarly aged individuals without dementia, with the nursing home facility costs greater than 20 times that of the latter group [46]. Despite the current absence of curative therapy, the justification for biomarker development remains compelling. Early detection of AD is needed to ensure early interventions that could potentially mitigate disease severity and also give families time to better prepare for the care of such individuals. With a very active drug pipeline, early detection will be needed to identify appropriate candidates for these trials. Finally, early detection and resulting intervention to slow disease progression could minimize time spent with severe dementia and promote the preservation of cognitive function for as long as possible. This would be beneficial for quality of life [47] and health care costs considerations. AD is a slowly developing disorder enhancing the feasibility of achieving these objectives.

Consistent with the call for the integration of breakthrough technologies (systems biology, genomics, big data science, and blood-based markers) to advance precision medicine objectives in AD [48], we combined AI analysis with leukocyte epigenomic data for AD prediction. Using raw intragenic CpG markers alone, we achieved a highly accurate prediction of AD using ML-based techniques. All the AI platforms achieved an AUC  $\geq 0.93$  using leukocyte epigenomic data. In the case of Deep Learning, we obtained an AUC = 0.99 with 97% sensitivity and specificity values. Additionally, we achieved high predictive accuracy using intergenic/extragenic CpG sites alone for AD detection. The use of conventional clinical predictors and MMSE did not improve performance further.

AI is superior to conventional statistical tools for the analysis of big data generated by omics analysis [17,49]. It is a powerful tool for discriminating and classifying groups. It can identify multiple markers each with limited individual predictive capabilities which when combined achieve excellent discriminating performance. To minimize the chances of overfitting strategies such as RF were used (see Supplementary Methods). For the sake of comparison, we also investigated the predictive performance of conventional logistic regression. Employing cross-validation techniques, regression analysis yielded good predictive accuracy for AD based on methylation markers: AUC (95%CI) = 0.85 (0.74–0.96) but less than that of AI. This, however, further supports the robustness of the leukocyte epigenomic markers for AD detection.

Currently, a range of imaging markers continues to be deployed in clinical and research diagnosis and evaluation of AD. These include CT, MRI, and PET imaging of the brain and CSF amyloid and tau levels. A systematic review of imaging biomarkers revealed that currently, the most commonly utilized antemortem diagnostic tests have achieved moderate to good diagnostic accuracy [50]. The expense, and in some cases the invasive nature of these tests, precludes use in the general aged population. Psychological testing including the MMSE, the most widely used cognitive test, might not be readily available in many primary care settings where the majority of elderly patients receive clinical care. Further, the MMSE was found on meta-analysis to have only modest accuracy for ruling out dementia when deployed in a community or primary care settings [51]. Based on all these considerations, there remains a need for accurate biological screening tests in a low to moderate risk setting.

While not a requirement, an important collateral benefit of an ideal biomarker, beyond predictive accuracy, is the ability to help elucidate disease pathogenesis. We identified altered CpG methylation in several individual genes (*CR1L*, *MYC*, *NRG1*, *LMNA*, *ELOVL4*, *MYB*, *AGPAT1*, and *NSG1*) previously reported playing a role in AD. Single nucleotide polymorphisms in these genes increase AD risk by affecting the formation of neurofibrillary tangles, neuronal apoptosis, and neuronal vesicle trafficking in AD (S7 Table). [52–60] Further, IPA found enrichment of several pathways involved in brain and neuronal development and brain and cardiovascular function such as abnormal morphology of cerebral cortex, gliosis, the morphology of the nervous system, inflammatory response and cardiac ventricular hypertrophy, and dilated cardiomyopathy (Figs 2 and 3 and S5–S7 Tables).

AD appears to primarily affect the medial temporal cortex of the brain and both AD and aging affect the inferior parietal lobe and dorsolateral prefrontal cortex regions of the brain [61]. The accumulation of a significant volume of neurofibrillary tangles in the neocortical region is a hallmark of AD development [62]. We found significant epigenetic changes in genes (*CR1L*, *CTSV*, *APAF1*, and *SS18L1*) responsible for cerebral cortical morphology.

Microglia are immune cells residing in the brain. Proliferation and hypertrophy of these cells (gliosis) occur in response to CNS damage. Gliosis can lead to neuroinflammation and induce tau pathology thus accelerating neurodegeneration. In the case of AD, amyloid- $\beta$  plaque deposition aggravates gliosis [63]. Our pathway analysis suggested a relationship between abnormal methylation and increased gliosis in AD. *S1PR1* and *MYC* genes were hypermethylated in our study. The *S1PR1* gene is involved in CNS inflammation [64] and the *MYC* gene in astrogliosis and inflammatory response [65].

We also found an over-representation of molecular pathways, including cardiac hypertrophy signaling and Wnt signaling, in AD. Vascular disease is strongly associated with negative effects on cognition [66]. Left ventricular hypertrophy is reported to be an independent risk factor for dementia [67]. We identified genes involved in cardiac hypertrophy signaling that displayed altered methylation in the AD group. Polymorphisms of the *ADRA2B* gene have been linked to cerebrovascular disorders [68]. The *FGF18* and *FGF22* genes are known to play a role in heart development and physiological processes [69] while the *MYC* gene is implicated in angiogenesis, cardiomyogenesis, apoptosis, oxidative stress response and plays a major role in initiating and maintaining cardiac hypertrophy and contractility [70]. In our study, these genes were found to be significantly differentially methylated and further support an important link between cardiovascular function and AD.

The Wnt/ $\beta$ -catenin signaling pathway is one possible link between cardiovascular disease and dementia. Wnt signaling is critical for the developmental processes in multiple organs including that of the heart. The pathway is reactivated in many post-natal cardiac disorders [71]. The activation of Wnt signaling has a neuroprotective effect while inhibition promotes neurodegeneration [72]. Downregulated Wnt/ $\beta$ -catenin signaling is associated with AD [73]. Wnt/ $\beta$ -catenin signaling genes such as *MYC*, *SOX14*, and *WNT9B* were found to be hypermethylated in the study.

A limitation of our study was the relatively small sample size. We also performed bootstrapping to confirm the stability of our estimates (see Supplemental Methods section). This slightly increased the performance estimates for 4 platforms including DL while slightly decreased the performance in 2 AI platforms. We intend to perform follow-up validation studies in a larger cohort of patients. Despite the study size, we demonstrated highly significant methylation changes in circulating leukocytes in AD. Highly accurate AD prediction was observed using an AI platform and different marker combinations. Also, while expression studies were not performed in this particular analysis, several CpG site methylation differences in AD cases versus controls were greater than 5–10%. This level of methylation difference has been noted to correlate with changes in corresponding gene expression [74]. While we did not perform expression analysis in the current study, we did find evidence of significant methylation changes in some leukocyte genes that have been previously reported to be differentially expressed in AD brains [36]. These findings also help to validate our data.

While significant epigenetic changes were also identified in the intergenic/ extragenic sites, we are currently unable to report the specific mechanisms of their contribution to AD pathogenesis as these sites have not been linked to particular genes. It is known however that intergenic/extragenic sites can exert long-range influence and control gene function.

Overfitting can be a challenge with AI analysis. To avoid overfitting in the DL model strategies including the use of regularization parameters, dropout, and controlling the input- dropout ratio were used and are detailed in the Supplemental Methods section. For the other AI platforms, several parameters were used to tune the models and to overcome the overfitting problem: number of trees for RF, classification cost for SVM, and threshold amount for shrinking toward the centroid for PAM.

Another limitation of the study is that we were not able to eliminate the possibility that some of the observed epigenetic changes were not due to co-morbidities such as schizophrenia, bipolar disorder, or epilepsy. Given the age of the study subjects, co-morbidities are the norm rather than the exceptions in AD. We did not however identify significant differences in the frequency of these disorders in our AD versus control groups. We did not have access to the medications of our study group. The study included a higher percentage of females in both the case and control groups. This however is consistent with the distinct gender-based demographics of the disorder. There was however no significant difference in the gender ratios of the case and control groups. Further, we removed all probes associated with X and Y chromosomes to minimize gender bias. We have excluded any CpGs having close association (0 to 10 bp distance) with single nucleotide polymorphisms to avoid genetic mutational association with the methylation changes. Finally, no information on the APOE gene mutation status was available for this particular cohort. These are not routinely obtained in the assessment of our clinical patients.

A significant strength of our study is the novelty, i.e. the use of blood leukocytes to accurately detect AD and also for interrogating the pathogenesis of AD. Leukocyte samples are easily obtained, raising the prospect of a minimally invasive and potentially affordable technique for investigation of the mechanisms, detection, as well as longitudinal monitoring of AD. The potential value of methylation changes in blood leukocytes for the detection of brain disorders including schizophrenia has been previously reported [75,76]. Of interest, we did find overlap in some of the genes that were significantly differentially methylated in AD in our study and a prior report of leukocyte DNA methylation variation in twins discordant for AD [43]. This provides further validation to the use of leukocyte methylation for the investigation of AD.

In summary, we have performed genome-wide methylation analysis in blood leucocytes and identified significant methylation changes in genes, gene networks, and disease pathways that were previously known or suspected to play an important role in AD. Significant methylation changes were also found in intergenic i.e. extragenic sites. Using AI techniques, highly accurate leukocyte epigenomic prediction of AD was reported for the first time to the authors' knowledge. The results could potentially advance the precision medicine objectives that have been outlined for AD [48]. Our work provides evidence in support of the view that epigenetic factors may play a pivotal role in AD development. Further validation studies using a larger number of subjects are necessary to confirm and expand on our findings.

### Supporting information

**S1 Table.** Clinical and demographic characteristics: AD compared to unaffected control subjects.

<https://doi.org/10.1371/journal.pone.0248375.s001>  
(DOCX)

**S2 Table.** Remaining (119 among 152) differentially methylated significant intragenic CpG markers.

<https://doi.org/10.1371/journal.pone.0248375.s002>  
(DOCX)

**S3 Table.** Alzheimer's disease prediction based on intragenic CpG markers.

<https://doi.org/10.1371/journal.pone.0248375.s003>  
(DOCX)

**S4 Table.** Alzheimer's disease prediction based on intergenic/extragenic CpG markers.

<https://doi.org/10.1371/journal.pone.0248375.s004>  
(DOCX)

**S5 Table.** Alzheimer's disease prediction based on intragenic CpG markers only: Genome-wide significance threshold\*.

<https://doi.org/10.1371/journal.pone.0248375.s005>  
(DOCX)

**S6 Table.** Alzheimer's disease prediction based on intergenic/extragenic CpG markers (stringent\* significance threshold).

<https://doi.org/10.1371/journal.pone.0248375.s006>  
(DOCX)

**S7 Table.** Differentially methylated genes enriched under molecular pathways in Alzheimer's disease (Ingenuity pathway analysis).

<https://doi.org/10.1371/journal.pone.0248375.s007>  
(DOCX)

**S8 Table.** Differentially methylated genes enriched in disease pathways of Alzheimer's disease (Ingenuity pathway analysis).

<https://doi.org/10.1371/journal.pone.0248375.s008>  
(DOCX)

**S9 Table.** List of few genes that were found to be significantly differentially.

<https://doi.org/10.1371/journal.pone.0248375.s009>  
(DOCX)

**S1 File.**

<https://doi.org/10.1371/journal.pone.0248375.s010>  
(DOCX)

### References

1. Association As. 2018 Alzheimer's Disease Facts and Figures.: Alzheimer's Association; 2018.
2. Prince MJ, Wimo A, Guerchet MM, Ali GC, Wu Y-T, Prina M. World Alzheimer Report 2015—The Global Impact of Dementia. London: Alzheimer's Disease International; 2015.
3. Wortmann M. Dementia: a global health priority—highlights from an ADI and World Health Organization report. *Alzheimers Res Ther.* 2012;4(5):40. pmid:22995353.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
4. Grinan-Ferre C, Corpas R, Puigoriol-Illamola D, Palomera-Avalos V, Sanfeliu C, Pallas M. Understanding Epigenetics in the Neurodegeneration of Alzheimer's Disease: SAMP8 Mouse Model. *J Alzheimers Dis.* 2018;62(3):943–63. pmid:29562529.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
5. Daviglus ML, Bell CC, Berrettini W, Bowen PE, Connolly ES Jr., Cox NJ, et al. NIH state-of-the-science conference statement: Preventing Alzheimer's disease and cognitive decline. *NIH Consens State Sci Statements.* 2010;27(4):1–30. pmid:20445638.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
6. Liu CC, Liu CC, Kanekiyo T, Xu H, Bu G. Apolipoprotein E and Alzheimer disease: risk, mechanisms and therapy. *Nat Rev Neurol.* 2013;9(2):106–18. pmid:23296339.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)



7. Kauwe JS, Ridge PG, Foster NL, Cannon-Albright LA. Strong evidence for a genetic contribution to late-onset Alzheimer's disease mortality: a population-based study. *PLoS One*. 2013;8(10):e77087. Epub 2013/10/12. pmid:24116205.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
8. Bertram L, Tanzi RE. Genome-wide association studies in Alzheimer's disease. *Hum Mol Genet*. 2009;18(R2):R137–45. Epub 2009/10/08. pmid:19808789.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
9. Zhang Q, Sidorenko J, Couvy-Duchesne B, Marioni RE, Wright MJ, Goate AM, et al. Risk prediction of late-onset Alzheimer's disease implies an oligogenic architecture. *Nat Commun*. 2020;11(1):4799. Epub 2020/09/25. pmid:32968074.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
10. Town T, Tan J, Flavell RA, Mullan M. T-cells in Alzheimer's disease. *Neuromolecular Med*. 2005;7(3):255–64. pmid:16247185.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
11. Richartz-Salzburger E, Batra A, Stransky E, Laske C, Kohler N, Bartels M, et al. Altered lymphocyte distribution in Alzheimer's disease. *J Psychiatr Res*. 2007;41(1–2):174–8. pmid:16516234.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
12. Rezai-Zadeh K, Gate D, Szekely CA, Town T. Can peripheral leukocytes be used as Alzheimer's disease biomarkers? *Expert Rev Neurother*. 2009;9(11):1623–33. pmid:19903022.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
13. Kusdra L, Rempel H, Yaffe K, Pulliam L. Elevation of CD69+ monocyte/macrophages in patients with Alzheimer's disease. *Immunobiology*. 2000;202(1):26–33. pmid:10879686.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
14. Li H, Guo Z, Guo Y, Li M, Yan H, Cheng J, et al. Common DNA methylation alterations of Alzheimer's disease and aging in peripheral whole blood. *Oncotarget*. 2016;7(15):19089–98. pmid:26943045.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
15. De Jager PL, Srivastava G, Lunnon K, Burgess J, Schalkwyk LC, Yu L, et al. Alzheimer's disease: early alterations in brain DNA methylation at ANK1, BIN1, RHBDF2 and other loci. *Nat Neurosci*. 2014;17(9):1156–63. pmid:25129075.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
16. Bakulski KM, Dolinoy DC, Sartor MA, Paulson HL, Konen JR, Lieberman AP, et al. Genome-wide DNA methylation differences between late-onset Alzheimer's disease and cognitively normal controls in human frontal cortex. *J Alzheimers Dis*. 2012;29(3):571–88. Epub 2012/03/28. pmid:22451312.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
17. Bahado-Singh RO, Vishweswaraiah S, Aydas B, Mishra NK, Guda C, Radhakrishna U. Deep Learning/Artificial Intelligence and Blood-Based DNA Epigenomic Prediction of Cerebral Palsy. *Int J Mol Sci*. 2019;20(9). Epub 2019/05/01. pmid:31035542.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
18. Bahado-Singh RO, Vishweswaraiah S, Aydas B, Mishra NK, Yilmaz A, Guda C, et al. Artificial intelligence analysis of newborn leucocyte epigenomic markers for the prediction of autism. *Brain Res*. 2019;1724:146457. pmid:31521637.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
19. Bahado-Singh RO, Vishweswaraiah S, Er A, Aydas B, Turkoglu O, Taskin BD, et al. Artificial Intelligence and the detection of pediatric concussion using epigenomic analysis. *Brain Res*. 2020;1726:146510. Epub 2019/10/20. pmid:31628932.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
20. Sajda P. Machine learning for detection and diagnosis of disease. *Annu Rev Biomed Eng*. 2006;8:537–65. pmid:16834566.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
21. Lee HC, Yoon SB, Yang SM, Kim WH, Ryu HG, Jung CW, et al. Prediction of Acute Kidney Injury after Liver Transplantation: Machine Learning Approaches vs. Logistic Regression Model. *J Clin Med*. 2018;7(11). pmid:30413107.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
22. Mamoshina P, Vieira A, Putin E, Zhavoronkov A. Applications of Deep Learning in Biomedicine. *Mol Pharm*. 2016;13(5):1445–54. pmid:27007977.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
23. Ching T, Himmelstein DS, Beaulieu-Jones BK, Kalinin AA, Do BT, Way GP, et al. Opportunities and obstacles for deep learning in biology and medicine. *J R Soc Interface*. 2018;15(141). pmid:29618526.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

24. Bahado-Singh RO, Sonek J, McKenna D, Cool D, Aydas B, Turkoglu O, et al. Artificial Intelligence and amniotic fluid multiomics analysis: The prediction of perinatal outcome in asymptomatic short cervix. *Ultrasound Obstet Gynecol*. 2018. Epub 2018/11/02. pmid:30381856.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
25. Bahado-Singh RO, Yilmaz A, Bisgin H, Turkoglu O, Kumar P, Sherman E, et al. Artificial intelligence and the analysis of multi-platform metabolomics data for the detection of intrauterine growth restriction. *PLoS One*. 2019;14(4):e0214121. Epub 2019/04/19. pmid:30998683.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
26. Alpay Savasan Z, Yilmaz A, Ugur Z, Aydas B, Bahado-Singh RO, Graham SF. Metabolomic Profiling of Cerebral Palsy Brain Tissue Reveals Novel Central Biomarkers and Biochemical Pathways Associated with the Disease: A Pilot Study. 2019;9(2). pmid:30717353.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
27. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR Jr., Kawas CH, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement*. 2011;7(3):263–9. Epub 2011/04/26. pmid:21514250.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
28. Altork N, Tsou PS, Coit P, Khanna D, Sawalha AH. Genome-wide DNA methylation analysis in dermal fibroblasts from patients with diffuse and limited systemic sclerosis reveals common and subset-specific DNA methylation aberrancies. *Annals of the rheumatic diseases*. 2014. pmid:24812288.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
29. Bahado-Singh RO, Vishweswaraiah S, Aydas B, Mishra NK, Guda C, Radhakrishna U. Deep Learning/Artificial Intelligence and Blood-Based DNA Epigenomic Prediction of Cerebral Palsy. *International Journal of Molecular Sciences*. 2019;20(9):2075. pmid:31035542  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
30. Alakwaa FM, Chaudhary K, Garmire LX. Deep Learning Accurately Predicts Estrogen Receptor Status in Breast Cancer Metabolomics Data. *Journal of proteome research*. 2018;17(1):337–47. Epub 2017/11/08. pmid:29110491.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
31. Candel A, Parmar V, LeDell E, Arora A. Deep Learning with H2O2018.
32. Kuhn M. Building Predictive Models in R Using the caret Package. *Journal of Statistical Software*. 2008;28(5):1–26.  
[View Article](#) • [Google Scholar](#)
33. Bollati V, Galimberti D, Pergoli L, Dalla Valle E, Barretta F, Cortini F, et al. DNA methylation in repetitive elements and Alzheimer disease. *Brain Behav Immun*. 2011;25(6):1078–83. pmid:21296655.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
34. Jannot AS, Ehret G, Perneger T.  $P < 5 \times 10^{-8}$  has emerged as a standard of statistical significance for genome-wide association studies. *J Clin Epidemiol*. 2015;68(4):460–5. Epub 2015/02/11. pmid:25666886.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
35. Luu BC, Wright AL, Haeberle HS, Karnuta JM, Schickendantz MS, Makhni EC, et al. Machine Learning Outperforms Logistic Regression Analysis to Predict Next-Season NHL Player Injury: An Analysis of 2322 Players From 2007 to 2017. *Orthop J Sports Med*. 2020;8(9):2325967120953404-. pmid:33029545.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
36. Miller JA, Woltjer RL, Goodenbour JM, Horvath S, Geschwind DH. Genes and pathways underlying regional and cell type changes in Alzheimer's disease. *Genome Med*. 2013;5(5):48. Epub 2013/05/28. pmid:23705665.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
37. Hansmannel F, Sillaire A, Kamboh MI, Lendon C, Pasquier F, Hannequin D, et al. Is the urea cycle involved in Alzheimer's disease? *J Alzheimers Dis*. 2010;21(3):1013–21. pmid:20693631.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
38. Rustenhoven J, Smith AM, Smyth LC, Jansson D, Scotter EL, Swanson MEV, et al. PU.1 regulates Alzheimer's disease-associated genes in primary human microglia. *Mol Neurodegener*. 2018;13(1):44. pmid:30124174.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
39. Gerdt J, Summers DW, Milbrandt J, DiAntonio A. Axon Self-Destruction: New Links among SARM1, MAPKs, and NAD<sup>+</sup> Metabolism. *Neuron*. 2016;89(3):449–60. pmid:26844829.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
40. Nasoohi S, Ismael S, Ishrat T. Thioredoxin-Interacting Protein (TXNIP) in Cerebrovascular and Neurodegenerative Diseases: Regulation and Implication. *Mol Neurobiol*. 2018;55(10):7900–20. pmid:29488135.

[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

41. Tindi JO, Chavez AE, Cvejic S, Calvo-Ochoa E, Castillo PE, Jordan BA. ANKS1B Gene Product AIDA-1 Controls Hippocampal Synaptic Transmission by Regulating GluN2B Subunit Localization. *J Neurosci*. 2015;35(24):8986–96. pmid:26085624.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
42. Huang C, Chen M, Pang D, Bi D, Zou Y, Xia X, et al. Developmental and activity-dependent expression of LanCL1 confers antioxidant activity required for neuronal survival. *Dev Cell*. 2014;30(4):479–87. pmid:25158856.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
43. Konki M, Malonzo M, Karlsson IK, Lindgren N, Ghimire B, Smolander J, et al. Peripheral blood DNA methylation differences in twin pairs discordant for Alzheimer's disease. *Clin Epigenetics*. 2019;11(1):130. Epub 2019/09/04. pmid:31477183.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
44. Prince M, Wilmo A, Guerchet M, Ali G, Wu Y, Prina M. World Alzheimer Report 2015: The global impact of dementia. An analysis of prevalence, incidence, costs and trends. London: Alzheimer's Disease International 2015. Alzheimer's Disease International The global voice of dementia. 2015.
45. Hutubessy R, Chisholm D, Edejer TT. Generalized cost-effectiveness analysis for national-level priority-setting in the health sector. *Cost Eff Resour Alloc*. 2003;1(1):8. pmid:14687420.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
46. Report AsA. Alzheimers's Association Report: 2019 Alzheimers's disease facts and figures. *Alzheimer's and Dementia* 2019;15:321–87.
47. Winblad B, Amouyel P, Andrieu S, Ballard C, Brayne C, Brodaty H, et al. Defeating Alzheimer's disease and other dementias: a priority for European science and society. *Lancet Neurol*. 2016;15(5):455–532. pmid:26987701.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
48. Hampel H, O'Bryant SE, Durreleman S, Younesi E, Rojkova K, Escott-Price V, et al. A Precision Medicine Initiative for Alzheimer's disease: the road ahead to biomarker-guided integrative disease modeling. *Climacteric*. 2017;20(2):107–18. pmid:28286989.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
49. Mirza B, Wang W, Wang J, Choi H, Chung NC, Ping P. Machine Learning and Integrative Analysis of Biomedical Big Data. *Genes (Basel)*. 2019;10(2). pmid:30696086.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
50. Cure S, Abrams K, Belger M, Dell'agnello G, Happich M. Systematic literature review and meta-analysis of diagnostic test accuracy in Alzheimer's disease and other dementia using autopsy as standard of truth. *J Alzheimers Dis*. 2014;42(1):169–82. pmid:24840572.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
51. Mitchell AJ. A meta-analysis of the accuracy of the mini-mental state examination in the detection of dementia and mild cognitive impairment. *J Psychiatr Res*. 2009;43(4):411–31. pmid:18579155.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
52. Kucukkilic E, Brookes K, Barber I, Guetta-Baranes T, Consortium A, Morgan K, et al. Complement receptor 1 gene (CR1) intragenic duplication and risk of Alzheimer's disease. *Hum Genet*. 2018;137(4):305–14. pmid:29675612.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
53. Ferrer I, Blanco R, Carmona M, Puig B. Phosphorylated c-MYC expression in Alzheimer disease, Pick's disease, progressive supranuclear palsy and corticobasal degeneration. *Neuropathol Appl Neurobiol*. 2001;27(5):343–51. pmid:11679086.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
54. Go RC, Perry RT, Wiener H, Bassett SS, Blacker D, Devlin B, et al. Neuregulin-1 polymorphism in late onset Alzheimer's disease families with psychoses. *Am J Med Genet B Neuropsychiatr Genet*. 2005;139B(1):28–32. pmid:16082692.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
55. Schjeide BM, McQueen MB, Mullin K, DiVito J, Hogan MF, Parkinson M, et al. Assessment of Alzheimer's disease case-control associations using family-based methods. *Neurogenetics*. 2009;10(1):19–25. pmid:18830724.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
56. Cluett C, Brayne C, Clarke R, Evans G, Matthews F, Rubinsztein DC, et al. Polymorphisms in LMNA and near a SERPINA gene cluster are associated with cognitive function in older people. *Neurobiol Aging*. 2010;31(9):1563–8. pmid:18848371.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
57. Bazan NG. Docosanoids and elovanoids from omega-3 fatty acids are pro-homeostatic modulators of inflammatory responses, cell damage and neuroprotection. *Mol Aspects Med*. 2018;64:18–33. pmid:30244005.

[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

58. Liu DX, Biswas SC, Greene LA. B-myb and C-myb play required roles in neuronal apoptosis evoked by nerve growth factor deprivation and DNA damage. *J Neurosci*. 2004;24(40):8720–5. pmid:15470138.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
59. Sherva R, Baldwin CT, Inzelberg R, Vardarajan B, Cupples LA, Lunetta K, et al. Identification of novel candidate genes for Alzheimer's disease by autozygosity mapping using genome wide SNP data. *J Alzheimers Dis*. 2011;23(2):349–59. pmid:21098978.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
60. Muthusamy N, Chen YJ, Yin DM, Mei L, Bergson C. Complementary roles of the neuron-enriched endosomal proteins NEEP21 and calcyon in neuronal vesicle trafficking. *J Neurochem*. 2015;132(1):20–31. pmid:25376768.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
61. Bakkour A, Morris JC, Wolk DA, Dickerson BC. The effects of aging and Alzheimer's disease on cerebral cortical anatomy: specificity and differential relationships with cognition. *Neuroimage*. 2013;76:332–44. pmid:23507382.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
62. Giannakopoulos P, Hof PR, Michel JP, Guimon J, Bouras C. Cerebral cortex pathology in aging and Alzheimer's disease: a quantitative survey of large hospital-based geriatric and psychiatric cohorts. *Brain Res Brain Res Rev*. 1997;25(2):217–45. pmid:9403139.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
63. Leyns CEG, Holtzman DM. Glial contributions to neurodegeneration in tauopathies. *Mol Neurodegener*. 2017;12(1):50. pmid:28662669.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
64. Kim S, Bielawski J, Yang H, Kong Y, Zhou B, Li J. Functional antagonism of sphingosine-1-phosphate receptor 1 prevents cuprizone-induced demyelination. *Glia*. 2018;66(3):654–69. pmid:29193293.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
65. Takarada-Iemata M, Kezuka D, Takeichi T, Ikawa M, Hattori T, Kitao Y, et al. Deletion of N-myc downstream-regulated gene 2 attenuates reactive astrogliosis and inflammatory response in a mouse model of cortical stab injury. *J Neurochem*. 2014;130(3):374–87. pmid:24697507.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
66. Samieri C, Perier MC, Gaye B, Proust-Lima C, Helmer C, Dartigues JF, et al. Association of Cardiovascular Health Level in Older Age With Cognitive Decline and Incident Dementia. *JAMA*. 2018;320(7):657–64. pmid:30140876.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
67. Scuteri A, Coluccia R, Castello L, Nevola E, Brancati AM, Volpe M. Left ventricular mass increase is associated with cognitive decline and dementia in the elderly independently of blood pressure. *Eur Heart J*. 2009;30(12):1525–9. pmid:19406864.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
68. Kim JO, Jeon YJ, Kim OJ, Oh SH, Kim HS, Shin BS, et al. Association between common genetic variants of alpha2A-, alpha2B- and alpha2C-adrenoceptors and the risk of silent brain infarction. *Mol Med Rep*. 2014;9(6):2459–66. pmid:24676565.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
69. Itoh N, Ohta H, Nakayama Y, Konishi M. Roles of FGF Signals in Heart Development, Health, and Disease. *Front Cell Dev Biol*. 2016;4:110. pmid:27803896.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
70. Wolfram JA, Lesnfsky EJ, Hoit BD, Smith MA, Lee HG. Therapeutic potential of c-Myc inhibition in the treatment of hypertrophic cardiomyopathy. *Ther Adv Chronic Dis*. 2011;2(2):133–44. pmid:21858245.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
71. Foulquier S, Daskalopoulos EP, Liuri G, Hermans KCM, Deb A, Blankestijn WM. WNT Signaling in Cardiac and Vascular Disease. *Pharmacol Rev*. 2018;70(1):68–141. pmid:29247129.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
72. Torres VI, Godoy JA, Inestrosa NC. Modulating Wnt signaling at the root: Porcupine and Wnt acylation. *Pharmacol Ther*. 2019. pmid:30790642.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
73. Vallee A, Lecarpentier Y. Alzheimer Disease: Crosstalk between the Canonical Wnt/Beta-Catenin Pathway and PPARs Alpha and Gamma. *Front Neurosci*. 2016;10:459. pmid:27807401.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

74. Leenen FA, Muller CP, Turner JD. DNA methylation: conducting the orchestra from exposure to phenotype? Clin Epigenetics. 2016;8:92. pmid:27602172.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
75. Aberg KA, Xie LY, McClay JL, Nerella S, Vunck S, Snider S, et al. Testing two models describing how methylome-wide studies in blood are informative for psychiatric conditions. Epigenomics. 2013;5(4):367–77. pmid:23895651.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
76. Liu J, Chen J, Ehrlich S, Walton E, White T, Perrone-Bizzozero N, et al. Methylation patterns in whole blood correlate with symptoms in schizophrenia patients. Schizophr Bull. 2014;40(4):769–76. pmid:23734059.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)