

---

## Figures

1.1	Main metabolic pathways	2
1.2	Metabolic pathways of glycolysis	4
1.3	Structure of ATP	5
1.4	Structure of NADH and NADPH	5
1.5	(a) Pentose phosphate pathway, and (b) the change in carbon numbers	11
1.6	Entner Doudoroff pathway	15
1.7	PDH and TCA cycle	16
1.8	Acetate producing pathways	21
1.9	Anaplerotic pathways	22
1.10	Gluconeogenetic pathways	24
1.11	Modularity of respiratory chains	26
1.12	Proton translocating values per electron in the respiratory chain	26
1.13	Schematic illustration for the electron transfer and proton translocation	28
1.14	Chemical structure of quinones	28
1.15	Pathway of UQ biosynthesis in <i>E. coli</i>	30
1.16	Anaerobic pathways	31
1.17	Nitrate respiration	33
1.18	Calvin-Benson cycle	35
1.19	Amino acid synthesis from their precursors	37
1.20	Alanine synthesis from PYR	38
1.21	Valine, leusine, and isoleusine biosynthesis	39
1.22	Glutamate and glutamine synthesis as well as aspartate and asparagine synthesis	39
1.23	Proline biosynthesis from glutamate	40
1.24	Arginine, ornithine, and citrulline synthesis pathway	40
1.25	Lysine, threonine, and methionine biosynthesis	41
1.26	Aromatic amino acid synthesis pathways	43
1.27	Several control schemes for aromatic amino acid biosynthesis	44

1.28 Serine, glycine, and cystein synthesis pathways	44
1.29 Histisine synthesis pathways	45
1.30 Nucleic acids synthesis pathways	46
1.31 $\beta$ oxidation and biosynthesis of fatty acid	47
1.32 Phosphotransferase system (PTS)	49
1.33 Fructose metabolism	50
1.34 Xylose metabolism	51
1.35 ATP balance for aerobic and anaerobic conditions	52
2.1 Batch cultivations of <i>E. coli</i> K12 using different DO levels and different carbon sources	58
2.2 2-DE gel maps of the total lysate of <i>E. coli</i> cells	63
2.3 The relative expression levels of <i>E. coli</i> K-12 proteins of central metabolic pathways under different conditions based on 2DE results	66
2.4 Comparison of the logarithmic protein expression ratios based on 2DE and the corresponding enzyme activity ratios	77
2.5 Growth curves of <i>E. coli</i> BW25113, which was grown in minimal media containing 10 g glucose/l as the sole carbon source	78
2.6 Relative gene expression of different global regulators and the metabolic pathway genes known to be regulated by those global regulators during different phases of growth	80
2.7 Concentration of different intracellular metabolites	83
2.8 Total and relative expression of iso-genes of <i>E. coli</i> metabolic pathways	84
2.9 Acetate metabolism	86
3.1 Overall metabolic regulation scheme	96
3.2 Outer and inner membrane and periplasm	98
3.3 Inducer exclusion and the activation of adenylate cyclase in the glucose-lactose system	101
3.4 The multiple regulations by Mlc and cAMP-Crp	103
3.5 Batch cultivation of (a) <i>E. coli</i> BW25113 and (b) its <i>cra</i> mutant	105
3.6 Comparison of enzyme activities of <i>cra</i> mutant as compared to the wild type (BW25113)	110
3.7 The effect of dilution rate on the gene transcript levels	113
3.8 Comparison of gene transcript levels of the wild type, <i>crp</i> knockout mutant, and <i>crp</i> <sup>+</sup> mutant	116

3.9	Glucose PTS and fructose PTS	118
3.10	Central metabolic pathways and $\text{NH}_3$ -assimilation pathways	121
3.11	Ammonia assimilation under different $\text{NH}_4^+$ concentration	122
3.12	Effect of C/N ratio on the fermentation characteristics for the continuous culture at the dilution rate of $0.2 \text{ h}^{-1}$	123
3.13	Schematic illustration of the interaction among several metabolic regulations. Comparison of the transcriptional mRNA levels of the wild type <i>E.coli</i> genes cultivated at 100% (C/N = 1.68), 40% (C/N = 4.21), 20% (C/N = 8.42) and 10% (C/N = 1.68) $\text{N}^-$ concentration: (a) global regulatory, (b) $\text{N}^-$ regulatory, (c) metabolic pathway, (d) respiratory chain	124
3.14	The interaction between nitrogen regulation and catabolite regulation	126
3.15	Overall mechanism of nitrogen assimilation in <i>E. coli</i> under C-limited (N-rich) and N-limited conditions	129
3.16	Molecular mechanism of phosphate regulation	131
3.17	Comparison of the transcript levels of the wild type <i>E.coli</i> cultivated with different P concentrations of the feed (100%, 55%, 10%)	134
3.18	Schematic illustration of the interaction among several metabolic regulation mechanisms	136
3.19	Comparison of some gene expressions for parent <i>E. coli</i> BW25113 and <i>arcB</i> mutant at 4 h of batch cultivation along with the gel picture	141
3.20	Comparison of specific enzyme activities of <i>E. coli</i> BW25113, its <i>arcB</i> and <i>arcA</i> mutant at 4 h of batch cultivation	141
3.21	Comparison of the transcript levels between wild type and <i>fnr</i> mutant under micro-aerobic continuous culture conditions	144
3.22	Comparison of enzymes activities during micro-aerobic batch culture	145
3.23	Metabolic flux distributions of wild type and <i>fnr</i> mutant under micro-aerobic conditions	147
3.24	Comparison of gene expressions	151
3.25	Specific enzyme activities in cell extracts	152
3.26	The role of glutamate decarboxylase for acid resistance	154
3.27	Acid resistance mechanism under acidic conditions	157

3.28	Effect of temperature up-shift on gene expressions in <i>E. coli</i> BW25113 under aerobic continuous culture at the dilution rate of $0.2 \text{ h}^{-1}$	160
3.29	Effect of heat shock on gene and protein expressions and the fermentation characteristics	163
3.30	Metabolic pathways showing levels of enzymes (or proteins) and intracellular metabolite concentrations in the <i>fadR</i> mutant <i>E. coli</i> relative to those in the parent at the exponential phase grown in glucose minimal medium under aerobic conditions	171
3.31	Different kinds of sigma factor in <i>E. coli</i>	176
3.32	Schematic diagram on the function of sigma factor as a transcription factor	176
3.33	Various levels of $\sigma^s$ regulation are differentially affected by various stress conditions	177
3.34	Growth curves of: (a) <i>E. coli</i> BW25113 (parent strain); and (b) <i>E. coli</i> JWK 2711 ( <i>rpoS</i> mutant)	179
3.35	Intracellular metabolite concentrations of <i>E. coli</i> BW25113 (parent strain) and <i>E. coli</i> JWK 2711 ( <i>rpoS</i> mutant)	184
4.1	Central reaction network for microalgae metabolism	221
4.2	Cultivation results of <i>C. pyrenoidosa</i> under autotrophic conditions	228
4.3	Cultivation results of <i>C. pyrenoidosa</i> under mixotrophic conditions	229
4.4	Cultivation results of <i>C. pyrenoidosa</i> under cyclic light-autotrophic/dark-heterotrophic conditions	230
4.5	Metabolic flux distribution of <i>Chlorella</i> cells	233
4.6	Energy conversion efficiency between the energy supplied to the culture, the energy absorbed by the cells, and the high free energy stored in ATP	238
4.7	Theoretical thermodynamic efficiency of ATP formation from the absorbed energy ( $Y_{\text{ATP}}/AE_{\text{max}}$ ) as a linear function of the fraction of the absorbed light energy of the total	239
4.8	Conversion efficiency of: (a) total energy; and (b) light energy during the various growth phases of the mixotrophic cultivation	241
4.9	Energy conversion efficiency between the three energy forms during the first light/dark cycle of the cyclic autotrophic/heterotrophic cultivation	242

4.10	General metabolic pathway of <i>E. coli</i> under oxygen-limited conditions	248
4.11	Factors influencing lactate producing flux	252
4.12	Factors influencing flux through Pta–Ack pathway	255
4.13	The effect of Pyk activity on: (a) Pyk flux; and (b) glycolytic flux	257
5.1	Simple examples of flux calculation	267
5.2	IDV and MAV	268
5.3	Schematic illustration of the data NMR and MS, compared to isotopomer distribution	270
5.4	The fate of labeled carbons for when using: (a) glucose; and (b) acetate or pyruvate, as a carbon source	276
5.5	Transformation of IDVs into MDVs and FMDVs for a C <sub>3</sub> molecule	285
5.6	Schematic illustration of net flux and exchange flux	286
5.7	The relationship between the precursor and amino acid	290
5.8	Derivatization to generate <i>M</i> -57 <sup>+</sup> and <i>M</i> -159 <sup>+</sup>	291
5.9	Net flux distribution in acetate metabolism of <i>E. coli</i> K12 in chemostat cultures at D of 0.11 and 0.22 h <sup>-1</sup>	295
5.10	Net flux distribution in glucose metabolism of <i>E. coli</i> K12 in chemostat cultures at D of 0.11 and 0.22 h <sup>-1</sup>	296
5.11	<sup>13</sup> C isotopomer patterns and NMR multiple spectral patterns	300
5.12	The fate of carbons of precursor metabolites to amino acids	301
5.13	Carbon position and <i>f</i> values	302
5.14	<sup>13</sup> C– <sup>13</sup> C scalar coupling	309
5.15	Metabolic flux distribution of the wild type and <i>pckA</i> mutant	311
5.16	Predicted dependencies of the flux through Pck	316
5.17	Bioreaction network of <i>E. coli</i> central carbon metabolism	320
5.18	<sup>13</sup> C– <sup>13</sup> C scalar coupling multiplets observed for aspartate from glucose-limited chemostat cultures of <i>E. coli</i> W3110 and the <i>pgi</i> mutant	323
5.19	<sup>13</sup> C– <sup>13</sup> C scalar coupling multiplets observed for C-4 of glucose from ammonia-limited chemostat cultures of <i>E. coli</i> W3110 and the <i>pgi</i> mutant	326
5.20	<sup>13</sup> C– <sup>13</sup> C scalar coupling multiplets observed for Asp-α from ammonia-limited chemostat cultures of <i>E. coli</i> W3110 and the <i>zwf</i> mutant	330

5.21 Metabolic flux distribution in chemostat cultures of <i>E. coli</i> W3110 under glucose-limited conditions	331
5.22 Specific rates of NADPH production and consumption in glucose (C)- and ammonia (N)-limited chemostat cultures of <i>E. coli</i> W3110, the <i>pgi</i> mutant, and the <i>zwf</i> mutant	333
5.23 <i>In vivo</i> flux distributions in the central metabolism of <i>Synechocystis</i> sp. PCC6803 cultivated (a) heterotrophically and (b) mixotrophically	343
5.24 Synthesis of the precursor for histidine (R5P or X5P) via the reactions of Calvin cycle	346
6.1 Metabolic flux distribution in the chemostat culture of the wild type <i>E. coli</i> (a) and <i>ppc</i> mutant (b)	365
6.2 Metabolic flux distributions in chemostat culture of glucose-grown <i>E. coli</i> parent strain ( <i>upper values</i> ), <i>gnd</i> ( <i>middle values</i> ), and <i>zwf</i> ( <i>lower values</i> ) mutants at $D = 0.2 \text{ h}^{-1}$	372
6.3 Metabolic flux distribution in chemostat culture of pyruvate-grown <i>E. coli</i> parent strain ( <i>upper values</i> ), <i>gnd</i> ( <i>middle values</i> ), and <i>zwf</i> ( <i>lower values</i> ) mutants at $D = 0.2 \text{ h}^{-1}$	373
6.4 Metabolic flux distribution of wild type (upper value) and <i>pykF</i> mutant (lower value) at dilution rate ( $D$ ) of $0.1 \text{ h}^{-1}$	380
6.5 Enzyme activities for the <i>pyk F</i> mutant at two dilution rates ( $D$ values)	381
6.6 Comparison of concentrations of intracellular metabolites in the <i>pykF</i> mutant at two $D$ values	381
6.7 Aerobic batch cultivation of (a) <i>E. coli</i> BW25113 and (b) <i>E. coli lpdA</i> mutant using glucose as carbon source in LB medium	383
6.8 Enzyme activities of <i>E. coli</i> BW25113 and <i>E. coli lpdA</i> mutant under aerobic conditions in LB medium	384
6.9 Comparison of the intracellular metabolite concentrations of <i>E. coli</i> BW25113 and <i>E. coli lpdA</i> mutant in the batch cultivation	385
6.10 Metabolic flux distributions of wild type (upper values) at dilution rate of $0.2 \text{ h}^{-1}$ and <i>lpdA</i> mutant (lower values, underlined) at dilution rate of $0.22 \text{ h}^{-1}$	387

6.11 Comparison of enzyme activities of <i>E. coli</i> BW25113 and <i>E. coli</i> <i>poxB</i> mutant under aerobic conditions using synthetic medium	388
6.12 Aerobic batch cultivation of: (a) <i>E. coli</i> BW25113; (b) <i>E. coli</i> <i>sucA</i> mutant; and (c) <i>E. coli</i> <i>sucC</i> mutant using glucose as carbon source	391
6.13 Comparison of enzyme activities at a dilution rate of $0.2 \text{ h}^{-1}$ in a continuous culture	393
6.14 Intracellular metabolite concentrations of <i>E. coli</i> BW25113, <i>sucA</i> and <i>sucC</i> gene knockout mutants at $0.2 \text{ h}^{-1}$ specific growth rate in chemostat cultures	394
6.15 Comparison of some gene expression at a dilution rate of $0.2 \text{ h}^{-1}$ in a continuous culture	395
6.16 Metabolic flux distributions of wild-type (upper values), <i>sucA</i> mutant (lower values) at dilution rate of $0.2 \text{ h}^{-1}$	396
6.17 Batch cultivation results of: (a) parent <i>E. coli</i> ; and (b) <i>icd</i> mutant, grown on glucose under aerobic conditions	399
6.18 Batch cultivation results of: (a) parent <i>E. coli</i> ; and (b) <i>icd</i> mutant, grown on glucose under microaerobic conditions	400
6.19 Batch cultivation results of: (a) parent <i>E. coli</i> ; and (b) <i>icd</i> mutant, grown on acetate under aerobic conditions	401
6.20 Relative protein and enzyme levels of <i>icd</i> mutant JW1122 from exponential growth phase in comparison to the parent <i>E. coli</i> BW25113 grown on glucose under aerobic conditions	407
6.21 The cultivation results of: (a) <i>E. coli</i> BW25113, and (b) the <i>pflB</i> <sup>-</sup> mutant, grown on glucose under the aerobic conditions	413
6.22 The cultivation results of: (a) <i>E. coli</i> BW25113; and (b) the <i>pflB</i> <sup>-</sup> mutant, grown on glucose under the microaerobic conditions	414
6.23 The specific glucose uptake rates and the specific product formation rates for: (a) the aerobic conditions; and (b) the microaerobic conditions	415
6.24 Biomass and metabolite yields on glucose for: (a) the aerobic conditions; and (b) the microaerobic conditions	416
6.25 The pyruvate formate lyase activities of <i>E. coli</i> <i>pfl</i> mutants and the parent strain <i>E. coli</i> BW25113	417

6.26 Enzyme activities for strains grown on glucose in the microaerobic conditions	418
6.27 Metabolic regulation mechanisms of lactate production in <i>E. coli pflA</i> and <i>pflB</i> mutants	423
6.28 Fermentative pathways of <i>E. coli</i> grown on glucose	427
6.29 Batch cultivation results of: (a) parent <i>E. coli</i> ; and (b) <i>ldhA</i> mutant <i>E. coli</i> , grown on glucose under anaerobic conditions	429
6.30 Intracellular metabolite concentrations in the central metabolic pathways at the exponential growth phase for both <i>ldhA</i> mutant and parent <i>E. coli</i>	432
6.31 Comparison of gene expressions for parent and <i>ldhA</i> mutant <i>E. coli</i>	434
6.32 Anaerobic metabolism of glucose in <i>E. coli</i>	436
6.33 NADH balances in both <i>ldhA</i> mutant and parent <i>E. coli</i> calculated based on the metabolic fluxes	438