

Notes on VAR Analysis for Doctorate

Rob Hayward

June 27, 2013

1 Preparation of data

The preparation of data is straight forward. The figures are downloaded and adjustments are made. It is possible to chose from a number of options for the data to be analysed in the VAR.

- The full dataset includes CNB, CNE, CNFDI, RTWI, SPREAD1 and SPREAD2 as well as S1 and S2.
- There are three dummy variables
- The data can be normalised to deal with the issue of hetorscedasticity that is evident in some of the series
- The current account can be added.
- It is possible to consider breaks in the data to deal with the issue of parameter instability. This can take three forms:
 - Select a break period and check the stability of parameters before and after the break
 - Impose an intervention dummy after the break and check statistical significance.
 - Look at rolling estimates of the parameters to assess stability

These can all be tested for unit roots to begin and as cointegrated vectors as a second move. The labtop has a loop to calcualte the Augmented Dickey-Fuller tests of the key variables.

2 Unit roots

The τ_3 test statistic is the test of the null hypothesis that the coefficient on the difference of the lagged dependent variable is equal to zero and that there is a *unit root* as ρ is equal to one.

The critical value for a sample size of 100 comes from (?).

An F-test of the null hypothesis that the coefficients on the lagged change in the dependend variable and the coefficient on the time trend are jointly equal to zero is also supplied (ϕ_3). The critical values come from Table VI (?) testing the null $(\alpha, \beta, \rho) = (\alpha, 0, 1)$. It seems that unit root and lack of time trend cannot be rejected. A joint test of the null that the coefficients on the drift, time trend and lagged difference of the dependent variable is suppoed in (ϕ_2). The critical values come from Table V (?) testing the null $(\alpha, \beta, \rho) = (0, 0, 1)$.

3 Cointegration tests with capital flow data

Testing for cointegration of the series. For the F-Test, that the coefficient on the lagged difference of the dependent variable and the coefficient on the It appears that the null of a unit root cannot be rejected for CNB, RTWI, Spread2. It is rejected at the 5% level for CNE, CFDI, COT and S1. Therefore, it appears that the former should be introduced in first difference.

According to the results from the "Test type: maximal eigenvalue statistic (lambda max), with linear trend in cointegration", there are 3 or 4 cointegrating realtionships.

The next step would be to use the cajorls function to estimate an error correction model. The results of this activity are shown in the following table.

test	10pct	5pct	1pct	test stat
r j= 6	1.91	6.50	8.18	11.65
r j= 5	13.07	15.66	17.95	23.52
r j= 4	31.44	28.71	31.52	37.22
r j= 3	59.99	45.23	48.28	55.43
r j= 2	106.41	66.49	70.60	78.87
r j= 1	168.18	85.18	90.39	104.20
r = 0	251.31	118.99	124.25	136.06

This suggests that there are three or four conintegrating variables. The results are the same no matter what the combination that is adopted.