Binary Choice Analysis

Robert Hickman

Data shown for:
date

[1] "10-Jan-2018"

monkey

[1] "Ulysses"


#plot p1
p1

Monkey Bid Positions on Binary Choice Task

10-Jan-2018; Ulysses; 1 blocks block 1 juice chosen **=** 0 0.75 NA trial number fractal position 100 50 right bid left bid juice offered 1.0 1.5 0.25 2.0 2.5 3.0 0.00 0.5 -0.5 0.0 bid position


Graph of choices for each block. Circles indicate bid selecting the bundle, squares are bid selecting the budget. A fractal bid position of 1 means that the bundle is on the left hand side of the screen. Bids range from -1 (all the way to the left) to 1 (all the way to the right)






Graph showing all choices and how far away they are from the edge of the screen on the bundle side. 0 indicates full movement to the bundle side of the screen and 1 represent full movement away. Count is over all blocks for all values of the fractal (in ml of juice).





10-Jan-2018 ; Ulysses ; 1 blocks



Graphs of various factors against the distance from the bundle side of the screen the monkey bids.


A bundle position of 1 indicates that the bundle is on the left hand side of the screen. A bundle water percentage of 1 indicates that the bundle contains no water [CHECK THIS- PRETTY SURE ITS CORRECT], whereas zero means it contains the full 1.2ml. Offer values of 1, 2, and 3 represent 0.15ml, 0.45ml, and 0.75mls of apple and mango juice (150ml in 950ml of water).

Fit lines use LOESS method.

```
#generate a model of likelihood to bid for the fractal dependent on it's position,
#value and associated water
model <- glm(data = task_data,</pre>
            fractal_bid ~ bundle_position + bundle_water_perc + offer_value + trial,
            family = "binomial")
#summarise the parameters
summary(model)
##
## Call:
## glm(formula = fractal_bid ~ bundle_position + bundle_water_perc +
      offer_value + trial, family = "binomial", data = task_data)
##
## Deviance Residuals:
##
       Min
                        Median
                  1Q
                                       3Q
                                                Max
                        0.03084
## -2.74230 -0.26287
                                  0.25652
                                            2.27362
##
## Coefficients:
##
                     Estimate Std. Error z value Pr(>|z|)
## (Intercept)
                     -2.739776
                                1.035542 -2.646 0.00815 **
## bundle position
                                0.687298 -1.344 0.17883
                    -0.923974
## bundle_water_perc -9.100556
                                1.988569 -4.576 4.73e-06 ***
## offer value
                     4.761118
                                0.920081
                                           5.175 2.28e-07 ***
## trial
                     -0.013728
                                0.008382 -1.638 0.10148
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
## (Dispersion parameter for binomial family taken to be 1)
##
##
       Null deviance: 180.871 on 134 degrees of freedom
## Residual deviance: 62.858 on 130 degrees of freedom
     (16 observations deleted due to missingness)
## AIC: 72.858
##
```


Number of Fisher Scoring iterations: 7






Graph showing the number of trials the monkey carried out for each bundle combination. Does not include failed trials.








Graph showing the proportion of bids for the bundle that a monkey makes, separated by the values of the juice offered in the bundles. Fits using a binomial glm model.

р6



Same graph as above but with 95% confidence intervals. Uses the default method of calculating this for the tidyverse libraries in R which I'm not convinced are the best way. Looking into calculating and plotting it myself.