

Binary Choice Analysis

Robert Hickman


Data shown for: date ## [1] "24-Jan-2018" monkey ## [1] "Ulysses" #plot p1 p1


Monkey Bid Positions on Binary Choice Task






Graph of choices for each block. Circles indicate bid selecting the bundle, squares are bid selecting the budget. A fractal bid position of 1 means that the bundle is on the left hand side of the screen. Bids range from -1 (all the way to the left) to 1 (all the way to the right)






Graph showing all choices and how far away they are from the edge of the screen on the bundle side. 0 indicates full movement to the bundle side of the screen and 1 represent full movement away. Count is over all blocks for all values of the fractal (in ml of juice).





24-Jan-2018 ; Ulysses ; 1 blocks




Graphs of various factors against the distance from the bundle side of the screen the monkey bids.

A bundle position of 1 indicates that the bundle is on the left hand side of the screen. A bundle water percentage of 1 indicates that the bundle contains no water [CHECK THIS- PRETTY SURE ITS CORRECT], whereas zero means it contains the full 1.2ml. Offer values of 1, 2, and 3 represent 0.15ml, 0.45ml, and 0.75mls of apple and mango juice (150ml in 950ml of water).

Fit lines use LOESS method.

```
#generate a model of likelihood to bid for the fractal dependent on it's position,
#value and associated water
model <- glm(data = task_data,</pre>
             fractal_bid ~ bundle_position + bundle_water_perc + offer_value + trial,
             family = "binomial")
#summarise the parameters
summary(model)
##
## Call:
## glm(formula = fractal_bid ~ bundle_position + bundle_water_perc +
      offer_value + trial, family = "binomial", data = task_data)
##
## Deviance Residuals:
##
      Min
                     Median
                 1Q
                                   3Q
                                           Max
## -2.1863 -0.4351 -0.0853
                               0.3726
                                        2.4724
##
## Coefficients:
##
                      Estimate Std. Error z value Pr(>|z|)
## (Intercept)
                     -5.866488
                                 0.932481 -6.291 3.15e-10 ***
## bundle position
                                            0.704
                      0.305285
                                 0.433517
                                                     0.481
## bundle_water_perc -4.380752
                                 0.913580 -4.795 1.63e-06 ***
## offer value
                      3.591311
                                 0.449576
                                           7.988 1.37e-15 ***
## trial
                      0.001053
                                 0.001808
                                            0.583
                                                     0.560
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
## (Dispersion parameter for binomial family taken to be 1)
##
##
       Null deviance: 327.33 on 236 degrees of freedom
## Residual deviance: 140.23 on 232 degrees of freedom
     (230 observations deleted due to missingness)
## AIC: 150.23
##
```

Number of Fisher Scoring iterations: 6



Graph showing the number of trials the monkey carried out for each bundle combination. Does not include failed trials.

0.75

bundle juice offered (ml)


0.25

0.0

0.00








Graph showing the proportion of bids for the bundle that a monkey makes, separated by the values of the juice offered in the bundles. Fits using a binomial glm model.

р6



Same graph as above but with 95% confidence intervals. Uses the default method of calculating this for the tidyverse libraries in R which I'm not convinced are the best way. Looking into calculating and plotting it myself.