Binary Choice Analysis

Robert Hickman

Data shown for:
dates

[1] "22-Feb-2018"

monkey

[1] "Ulysses"

precise_dates


[1] "22-Feb-2018 "

#plot p1


p1


Monkey Bid Positions on Binary Choice Task
Ulysses; 1 blocks

block 1



Graph of choices for each block. Circles indicate bid selecting the bundle, squares are bid selecting the budget. A fractal bid position of 1 means that the bundle is on the left hand side of the screen. Bids range from -1 (all the way to the left) to 1 (all the way to the right)






Graph showing all choices and how far away they are from the edge of the screen on the bundle side. 0 indicates full movement to the bundle side of the screen and 1 represent full movement away. Count is over all blocks for all values of the fractal (in ml of juice).



Monkey Bid Distance From Bundle on Binary Choice Task

Ulysses; 1 blocks




Graphs of various factors against the distance from the bundle side of the screen the monkey bids.

A bundle position of 1 indicates that the bundle is on the left hand side of the screen. A bundle water percentage of 1 indicates that the bundle contains no water [CHECK THIS- PRETTY SURE ITS CORRECT], whereas zero means it contains the full 1.2ml. Offer values of 1, 2, and 3 represent 0.15ml, 0.45ml, and 0.75mls of apple and mango juice (150ml in 950ml of water).

Fit lines use LOESS method.


```
#generate a model of likelihood to bid for the fractal dependent on it's position,
#value and associated water
model <- glm(data = task_data,</pre>
             fractal_bid ~ bundle_position + bundle_water_perc + offer_value + trial,
             family = "binomial")
#summarise the parameters
summary(model)
##
## Call:
## glm(formula = fractal_bid ~ bundle_position + bundle_water_perc +
      offer_value + trial, family = "binomial", data = task_data)
##
## Deviance Residuals:
##
       Min
                         Median
                   1Q
                                       3Q
                                                Max
## -2.56508 -0.39901
                        0.07749
                                  0.44109
                                            2.44801
##
## Coefficients:
##
                      Estimate Std. Error z value Pr(>|z|)
## (Intercept)
                     -1.602041
                                 0.636219 -2.518
                                                    0.0118 *
## bundle position
                                 0.414688 -2.036
                     -0.844273
                                                    0.0418 *
## bundle_water_perc -7.559530
                                 1.104194 -6.846 7.58e-12 ***
## offer value
                      2.840509
                                 0.392227
                                           7.242 4.42e-13 ***
## trial
                      0.003427
                                 0.002290
                                            1.496
                                                    0.1345
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
## (Dispersion parameter for binomial family taken to be 1)
##
       Null deviance: 327.29 on 239 degrees of freedom
##
## Residual deviance: 154.90 on 235 degrees of freedom
     (89 observations deleted due to missingness)
## AIC: 164.9
##
```

Number of Fisher Scoring iterations: 6



Graph showing the number of trials the monkey carried out for each bundle combination. Does not include failed trials.






Graph showing the proportion of bids for the bundle that a monkey makes, separated by the values of the juice offered in the bundles. Fits using a binomial glm model.

water offered (ml)

р6



Same graph as above but with 95% confidence intervals. Uses the default method of calculating this for the tidyverse libraries in R which I'm not convinced are the best way. Looking into calculating and plotting it myself.