



Original Article

The effect of sleep deprivation on pain perception in healthy subjects: a meta-analysis



Marlene Schrimpf^a, Gregor Liegl^{a,b}, Markus Boeckle^a, Anton Leitner^a, Peter Geisler^c,
Christoph Pieh^{a,d,e,*}

^a Department of Psychotherapy and Biopsychosocial Health, Danube-University Krems, Dr. Karl Dorrek Straße 30, A-3500 Krems, Austria

^b Medical Clinic, Department of Psychosomatic Medicine, Charité – Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany

^c Department of Psychiatry and Psychotherapy, University Hospital Regensburg, Universitätsstraße 84, 93053 Regensburg, Germany

^d Department of Psychosomatic Medicine, University Hospital Regensburg, Franz-Josef-Strauß-Allee 11, 93053 Regensburg, Germany

^e Karl Landsteiner University of Health Sciences, Dr.-Karl-Dorrek-Straße 30, 3500 Krems, Austria

ARTICLE INFO

Article history:

Received 22 April 2015

Received in revised form 17 July 2015

Accepted 23 July 2015

Available online 20 August 2015

Keywords:

Sleep deprivation

Pain perception

Meta-analysis

Reciprocal relation

ABSTRACT

Background: There is strong evidence indicating an interaction between sleep and pain. However, the size of this effect, as well as the clinical relevance, is unclear. Therefore, this meta-analysis was conducted to quantify the effect of sleep deprivation on pain perception.

Methods: A systematic literature search was conducted using the electronic databases PubMed, Cochrane, Psynex, Psycinfo, and Scopus. By conducting a random-effect model, the pooled standardized mean differences (SMDs) of sleep deprivation on pain perception was calculated. Studies that investigated any kind of sleep deprivation in conjunction with a pain measurement were included. In cases of several pain measurements within a study, the average effect size of all measures was calculated.

Results: Five eligible studies ($N = 190$) for the between-group analysis and ten studies ($N = 266$) for the within-group analysis were identified. Sleep deprivation showed a medium effect in the between-group analysis (SMD = 0.62; CI95: 0.12, 1.12; $z = 2.43$; $p = 0.015$) and a large effect in the within-group analysis (SMD = 1.49; CI95: 0.82, 2.17; $z = 4.35$; $p < 0.0001$). The test for heterogeneity was not significant in the between-group analysis ($Q = 5.29$; $df = 4$; $p = 0.2584$), but it was significant in the within-group analysis ($Q = 53.49$; $df = 9$; $p < 0.0001$).

Conclusion: This meta-analysis confirms a medium effect (SMD = 0.62) of sleep deprivation on pain perception. As this meta-analysis is based on experimental studies in healthy subjects, the clinical relevance should be clarified.

© 2015 Elsevier B.V. All rights reserved.

1. Introduction

Chronic pain and sleep disorders are common in the general population. The prevalence of chronic pain ranges from 10% to 40% [1], which is comparable with the prevalence of sleep disorders (10–36% [2–9]).

Increasingly more studies indicate a reciprocal relationship between sleep and pain [2–6,10]. This hypothesis is based on epidemiological, pharmaceutical, clinical, and experimental studies. For example, 50–88% of patients with chronic pain also suffer from sleep disorders [11–13]. Alternatively, > 40% of patients with insomnia also report chronic pain [6]. Several pharmaceutical studies, including studies on eszopiclone [14], triazolam [15], or pregabalin [16],

showed a simultaneous improvement of both sleep and pain. In addition, clinical trials highlight a bidirectional relationship between sleep and pain. Patients with a range of sleep disorders, such as obstructive sleep apnea syndrome (OSAS) [17] and restless legs syndrome (RLS) [18], seem to be hyperalgesic. The pain sensitivity improves with effective treatment of the respective sleep disorder. This improvement has been demonstrated in both pharmacological interventions, such as levodopa (L-DOPA) therapy for RLS [18], as well as non-pharmacological interventions, such as continuous positive airway pressure (CPAP) treatment for OSAS [17]. The latter study is of particular interest, as the improvement disappears immediately upon the discontinuation of CPAP therapy [17].

The first experimental study that showed an effect of sleep deprivation on pain perception was published in 1975 [19]. Since then, several studies have confirmed these results [7,20,21]. A review from 2006 showed a clear effect of sleep deprivation on pain perception in animals, whereas human data were inconclusive [22]. The same study group extended the systematic review in 2014 and found evidence of the hyperalgesic effect of sleep deprivation, both in animals

* Corresponding author. Department of Psychotherapy and Biopsychosocial Health, Faculty of Health and Medicine, Danube University Krems, Dr. Karl Dorrek Straße 30, A-3500 Krems an der Donau, Austria. Tel.: +432732 893/2530; fax: +432732893/4630.
E-mail address: Christoph.pieh@donau-uni.ac.at (C. Pieh).

and humans [23]. By focusing solely on controlled trials, a review indicated conclusive evidence of an increase of bodily complaints, or a decrease of pain thresholds, following sleep deprivation [24].

Varying explanations exist for the reciprocity between sleep and pain. For example, sleep and pain could share similar neurotransmitter systems, such as the dopaminergic system [25] and the serotonin system [26], or influence the endogenous opioid system [27]. In addition, substance P, which is involved in pain processes, interferes with sleep by reducing sleep efficiency or increasing awakenings [28]. However, the pathway of this interaction is not sufficiently clarified by the present research.

In summary, there is a large body of evidence for the interaction between sleep and pain. However, the size of this effect is unclear and the clinical relevance uncertain. Therefore, this meta-analysis was conducted to quantify the effect of sleep deprivation on pain perception.

2. Methods

2.1. Search strategy

Five electronic scientific databases, namely PubMed, Cochrane, Psynex, Psycinfo, and Scopus, were used. The search was conducted on December 5, 2014, with no limitations on the publication date. All terms that were appropriate in former reviews were used [22–24,29].

Sleep AND pain AND (“sleep restriction” OR “sleep deprivation” OR “sleep loss” OR “sleep interruption” OR “hyperalgesia” OR “total

sleep deprivation” OR “partial sleep deprivation” OR “sleep fragmentation”)

One author (M.S.) and one research assistant (F.H.) independently screened the titles and abstracts for eligibility (Fig. 1) with a high inter-rater reliability (Cohen’s Kappa $\kappa = 0.96$) [30] by at least one of the reviewers read the full text for all studies that were not excluded after screening the titles and abstracts. One author (M.S.) performed the final study selection, controlled by a second author (C.P.). The reference lists of former reviews about sleep deprivation and pain perception were screened [22–24,29] for additional eligible studies.

2.2. Inclusion and exclusion criteria

Experimental studies included a study sample of healthy adult humans (older than 18 years). Sleep deprivation had to be used as an independent variable and pain perception as a primary outcome. Only articles published in a peer-reviewed journal written in English language were included in the meta-analysis. Studies had to contain all necessary data for calculating standardized mean differences. The corresponding authors were contacted for necessary data, if incomplete.

2.3. Data extraction and risk of bias

Characteristics of intervention and sample, duration of sleep deprivation, type of pain measurement, and data needed for effect-size calculation were extracted.




Fig. 1. Flow diagram for search procedure

The risk of bias in this study was assessed only for between-group analysis. The Cochrane Collaboration's tool was used for assessing the risk of bias in randomized trials [31]. We rated the study quality of included studies for between-group analyses with six of the seven criteria of the assessment tool (random sequence generation, allocation concealment, blinding of outcome assessment, incomplete outcome data, selective reporting, and other bias). The criterion of blinding participants was excluded due to unfeasibility blinding condition in sleep deprivation. The risk of bias was rated as low when five or six criteria were met, as medium when three or four criteria were fulfilled, and as high when two or fewer criteria were met. The risk of bias was not assessed in studies with crossover or uncontrolled design.

To calculate effect size, means and standard deviations of all pain outcomes (eg, heat pain threshold, visual analog scale, mechanical pain threshold) at baseline and after sleep deprivation were extracted. If eligible, outcomes measured at the same time of the day were extracted. If one study reported several pain measurements, the mean effect size of all measures was calculated.

2.4. Data analysis

A meta-analysis of between-group effect sizes and within-group effect sizes was conducted.

Standardized mean differences for between-group effect sizes (SMD_B) were calculated. We used a formula described by Morris and colleagues [32,33] and a simple bias adjustment for small sample sizes recommended by Becker [34]:

$$SMD_B = c_T (M_{CHANGE-T} / SD_{PRE-T}) - c_C (M_{CHANGE-C} / SD_{PRE-C}).$$

$M_{CHANGE-T}$ is the mean of the change scores in the treatment group, $M_{CHANGE-C}$ is the mean of the change scores in the control group, SD_{PRE-T} is the standard deviation before sleep deprivation for the treatment group, and SD_{PRE-C} is the standard deviation before sleep deprivation for the control group. Because of the small sample sizes in our meta-analysis, a simple bias adjustment (c_T and c_C) was used [34], approximated by

$$c_j = 1 - (3 / (4(n_{pre(j)} + n_{post(j)} - 2) - 1)).$$

If the mean and standard deviation were not available, but the t - or F -values were reported, these were converted to effect sizes with the following formulas [35]:

$$SMD_B = c_T \frac{2t}{\sqrt{df}}$$

$$SMD_B = c_T \frac{2\sqrt{F(1, x)}}{\sqrt{df}}$$

Accordingly, the following formula was used for within-group effect-size calculations (SMD_W):

$$SMD_W = c_T (M_{CHANGE-T} / SD_{PRE-T}).$$

Effect sizes of 0.2 were interpreted as small, 0.5 as medium, and 0.8 as large [36]. Positive effect sizes indicate increasing pain.

A H - V random-effect method was used to estimate effect sizes and model parameters because the included studies vary in several conditions that may moderate effect sizes [37]. We conducted a Q -test and reported τ^2 to estimate the between-study variability. I^2 -statistic was used for relative heterogeneity. An I^2 of 0% indicates no heterogeneity, 25% low heterogeneity, 50% moderate heterogeneity, and 75% high heterogeneity [38].

Using Microsoft Office Excel®, single effect sizes were calculated. To perform the meta-analysis and generate corresponding graphics and plots [39], the *metafor* package for R was used. To check publication bias, forest plots displaying single and pooled effect sizes with 95% confidence intervals and funnel plots were created.

3. Results

3.1. Study selection

Fig. 1 shows the study selection process. A total of 456 records were found in the searching procedure after removing duplicates. By screening the titles and abstracts, 34 potential eligible studies were identified. After full-text analyses, 21 studies were excluded for the following reasons: no primary research [40–42], no experimental design [43], no sleep deprivation [44–49], no primary pain outcome [50,51], only non-healthy participants [52], and no opportunity to calculate effect size [2,7,9,19,53–56]. From the remaining 13 studies, eight were randomized controlled trials (RCTs) and five were uncontrolled. The study by Haack et al. [57] was excluded because it was a subsample of another study [58]. The study by Irwin et al. [59] was excluded because it compared a healthy to a non-healthy group, and its within-group analyses did not include all of the necessary data for calculating effect size. Schey et al. [60] also compared a healthy to a non-healthy group, but only the healthy subgroup was included. Although Schey et al. [60] used a crossover design for the healthy group, data for calculating the between-group effect were not reported, so the study for between-group effect analyses was excluded, but the healthy subgroup in the within-group effect analyses was included.

3.2. Between-group meta-analysis

Five studies in the between-group meta-analysis with a total of 190 subjects was included [20,21,58,61,62]. The participants were completely sleep deprived in three studies [20,21,62] and sleep restricted (4–6 h sleep per day) in two studies [58,61]. All five studies used one or more visual analog scales (VAS) to measure pain, and two studies additionally analyzed pain threshold using quantitative sensory testing (QST) [20,62]. The risk of bias was low for all five of these studies. Table 1 shows detailed information of included studies.

The pooled between-group meta-analysis resulted in a significant medium effect size of sleep-deprived conditions in comparison to non-sleep deprived conditions ($SMD = 0.62$; $CI_{95\%}: 0.12, 1.12$; $z = 2.43$; $p = 0.0152$, Fig. 2).

The test for heterogeneity in effect sizes was not significant ($Q = 5.29$; $df = 4$; $p = 0.2584$), and no remarkable variance was found in the between-group effect-size distribution ($\tau^2 = 0.2816$; $I^2 = 24.45\%$). The funnel plot to check potential publication bias is symmetrical (Fig. 3).

3.3. Within-group meta-analysis

One study from the between-group effect-size meta-analysis could not be included in the within-group meta-analysis [61] due to insufficient information to calculate the within-group effect size (only the F -value was reported). In addition to the four remaining studies for between-group effect sizes [20,21,58,62], six other studies were included in the meta-analysis for within-group effect sizes [60,63–67], ten studies and 266 subjects in total. Detailed information is also provided in Table 1. Four studies assessed participants with total sleep deprivation [20,21,62,66], whereas the participants in two studies were subjected to only slow-wave-sleep deprivation [63,64]. In the remaining four studies, the participants were only allowed to sleep for 2–4 h per night [58,60,65,67].

Table 1

Detailed information of included studies.

Author (year)	Study design	N (IG/CG)	Descriptives: <i>M</i> age (<i>SD</i> /range), % male, <i>n</i>	Sleep deprivation	Pain outcomes	Risk of bias ^a
Ablin et al. (2013) [61] ^a	RCT with four groups ^c	<i>n</i> = 87	Full sample: 27.2 years (<i>SD</i> = 5.6); 52% male	Sleep restriction (6 h of sleep per night for ten days)	VAS from the short-form McGill Pain Questionnaire	Low
Arima et al. (2001) [63] ^b	Uncontrolled	<i>n</i> = 10	22.7 years (<i>SD</i> = 0.9); 100% male	Slow-wave-sleep deprivation (three nights)	VAS (headache, pain in jaw muscles, soreness in jaw muscles, stiff in jaw muscles, pain in neck/shoulder region), Pain pressure threshold ^d	Low
Azevedo et al. (2011) [20] ^{a,b}	RCT with three groups ^e	<i>n</i> = 19 (9/10)	Full sample: range 19–28 years; 100% male	Total sleep deprivation for two consecutive nights	Laser-evoked pain, VAS pain	Low
Haack and Mullington (2005) [58] ^{a,b}	RCT	<i>n</i> = 40 (22/18)	SDG: 26.0 years (<i>SD</i> = 4.9); 63% male CG: 26.6 years (<i>SD</i> = 5.1); 67% male	Sleep restriction (4 h of sleep per night for 12 days)	VAS generalized body pain	Low
Haack et al. (2009) [21] ^{a,b}	RCT	<i>n</i> = 24 (16/9)	SDG: 37.9 years (<i>SD</i> = 6.5); 73% male CG: 30.4 years (<i>SD</i> = 5.6); 67% male	Total sleep deprivation (88 h)	VAS spontaneous pain	Low
Kundermann et al. (2004) [62] ^{a,b}	RCT	<i>n</i> = 20 (10/10)	SDG: 34.9 years (<i>SD</i> = 2.3); 60% male CG: 36.6 years (<i>SD</i> = 2.0); 50% male	Total sleep deprivation for two nights	Cold pain threshold, heat pain threshold, pain questionnaire	Low
Lentz et al. (1999) [64] ^b	Uncontrolled	<i>n</i> = 12	44 years (<i>SD</i> = 6.2); 0% male;	Slow-wave-sleep deprivation (three nights)	Tender point pain threshold	
Schstatsky et al. (2013) [65] ^b	Uncontrolled	<i>n</i> = 19	24.3 years (range 22–30 years); 100% male	Work-induced sleep restriction for one night (<i>M</i> = 120 min (<i>SD</i> = 28) sleep)	Heat pain threshold, electrical painful threshold	
Schey et al. (2007) [60] ^b	RCT with cross-over ^f	<i>n</i> = 10	36.9 years (<i>SD</i> = 17.08); 50% male	Sleep restriction (≤3 h of sleep for one night)	Lag time to symptom report, intensity rating, acid perfusion sensitivity score	
Schuh-Hofer et al. (2013) [66] ^b	Cross-over	<i>n</i> = 14	23.5 years (<i>SD</i> = 4.1); 57% male	Total sleep deprivation for one night	Cold pain threshold, heat pain threshold, pressure pain threshold, mechanical pain threshold, mechanical pain sensitivity	
Tiede et al. (2010) [67] ^b	Uncontrolled	<i>n</i> = 10	25.3 years (<i>SD</i> = 2.2); 80% male	Sleep restriction (4 h of sleep for one night)	Laser-evoked pain, VAS pain	

Note: SDG: intervention group; CG: control group; RCT: randomized controlled trial; VAS: visual analog scale.

^a Extracted for between-group analysis.^b Extracted for within-group analysis.^c Two groups with sleep restriction and two groups without sleep restriction.^d No data available.^e Only total sleep deprivation included.^f Only healthy participants included.




Fig. 2. Single and pooled effect sizes and 95% confidence interval for between-group (A) and within-group (B) meta-analysis. The squares represent the single effect sizes of each study, and the size of the squares are proportional to the precision of the estimates. The vertical line indicates the confidence interval of the effect size. Random-effect (RE) model shows the pooled effect size with 95% confidence interval.

QST was used in seven of the included studies [20,62–67] and VAS in six studies [20,21,58,62,63,67], and one study used stimulus response functions to acid to measure pain [60].

The pooled within-group effect size was large and significant (SMD = 1.49; CI₉₅: 0.82, 2.17; $z = 4.35$; $p < 0.0001$, Fig. 2). A significant heterogeneity of effect sizes ($Q = 53.49$; $df = 9$; $p < 0.0001$) and a high variance level were found in the within-group effect-size distribution ($\tau^2 = 0.9361$; $I^2 = 83.17\%$). The funnel plot of potential publication was asymmetrical (Fig. 3); a single sample with large standard errors produced the largest effect size [20,21,58,63]. Without this sample, we still found a large significant pooled effect size (SMD = 1.33; CI₉₅: 1.69, 1.98; $z = 4.03$; $p < 0.0001$).

4. Discussion

This meta-analysis confirms the effect of sleep deprivation on pain perception. A medium effect (SMD = 0.62) was found for the between-group analysis and a large effect (SMD = 1.49) was found for the within-group analysis. Sleep deprivation increases self-reported pain and affects evoked pain responses obtained through somatosensory testing protocols (eg, heat pain thresholds, pressure pain thresholds, or laser-evoked pain).

Patients with chronic pain exhibit several changes in sleep, such as frequent sleep-stage shifts, increased nocturnal awakenings, and decreased slow-wave or rapid-eye-movement sleep [68]. A recent review on 29 controlled polysomnographic studies found fragmentation of sleep to be the most common alteration of sleep in patients

with chronic pain [69]. In addition, increased amounts of superficial sleep and decreased levels of slow-wave sleep were found, whereas sleep architecture was commonly preserved. A consistent pattern of objective sleep disturbances was not found in nonmalignant chronic pain populations [69].

It is evident that the results from sleep deprivation and pain perception studies in an experimental design cannot be extended directly to sleep disorders or chronic pain. However, there are several relevant studies on this topic. Pain threshold was lower in pain patients with endometriosis [70] as well as with juvenile idiopathic arthritis [71]. In addition, the pain threshold correlates with functional scores in patients with osteoarthritis [72], although the results are still inconsistent. For example, sleep deprivation led to an increase of clinical pain complaints but an unaltered experimental pain threshold in patients with somatoform disorders [53]. This could be because self-reported pain is more closely associated with anxiety than with pain threshold [73]. Sleep and pain are critical homeostatic systems that interact in a bidirectional manner [69]. For example, <6 h of sleep per night is associated with greater pain the following day [74], and sleep quality is a consistent predictor of pain the next day [75,76]. Short-term sleep improvement predicts long-term pain improvement in patients with comorbid osteoarthritis and insomnia [77]. Insufficient sleep, in both quantity and quality, is predictive of pain three years in the future [78]. In a sample of 24 women, Wright et al. [79] found that lower sleep efficiency is a significant predictor of greater pain severity after breast surgery. The hyperalgesic effect of sleep deprivation seems reversible by




Fig. 3. Funnel plots of publication bias for between-group and within-group meta-analyses.

napping [80] as well as by effective treatment of sleep disorders [17]. Roehrs et al. [81] showed that extended sleep reduces pain sensitivity, and poor sleep quality is a risk factor for the development of chronic widespread pain [82].

In addition, several pharmaceutical studies demonstrated simultaneous changes in sleep and pain. Eszopiclone improves sleep and pain in patients with insomnia and comorbid rheumatoid arthritis [14], as well as in patients with chronic back pain [83]. Similarly, anticonvulsives, such as pregabalin, reduce pain and improve pain-related sleep interference [84]. However, as drugs often have multiple effects, a simultaneous interference could be mediated through an underlying but independent link. This also applies to similar comorbidities of sleep disorders and chronic pain, such as obesity [85], type 2 diabetes [86,87], or depression [88,89].

Several explanations exist for the interaction of sleep and pain. Paiva et al. [90] investigated the sleeping behavior of adolescents and found a correlation between quality of life, health complaints, and sleep. The interaction between sleep and pain could also be mediated by negative emotions [91]. Sleep loss can affect mood states in healthy adolescents [92], and sleep deprivation decreases positive and increases negative mood [58,66,93]. These findings suggest that addressing negative mood directly, or by addressing sleep disturbances in patients with chronic pain, may have a beneficial impact on patients' pain [91]. In addition, neurotransmitters, such as dopamine, play a role in sleep and pain [25]. In addition, sleep deprivation decreases morphine analgesia, possibly by decreasing descending pain inhibitory activity and by

increasing descending pain facilitatory activity [94]. Conversely, sleep improves on intrathecal morphine infusion in patients with severe pain [95]. A recent meta-analysis showed that sleep disturbance is associated with an increase in markers of systemic inflammation [96]. Systemic inflammation resulted in a lower pain threshold [97]. In addition, the tumor necrosis factor (TNF) system [98] as well as the hypothalamic–pituitary–adrenal axis [99] are influenced by sleep and pain. Melatonin, which is an essential regulator of the circadian rhythm, also interacts with pain [100,101].

The calculated effect size of sleep deprivation on pain perception is comparable with effect sizes achieved through chronic pain treatment. A recent meta-analysis showed medium effects for several pharmacological therapies, such as non-opioids (SMD = 0.41), “weak” opioids (SMD = 0.44), and “strong” opioids (SMD = 0.46) for the treatment of chronic pain [102]. Not only pain but also analgesics can alter sleep architecture by suppressing slow-wave or rapid-eye-movement sleep [68]. The effects of non-pharmaceutical interventions vary distinctly according to the control conditions (waiting list vs. active control group). On average, physiotherapy (SMD = 0.52) seems more beneficial than psychotherapy (SMD = 0.39) in cases of chronic pain [102]. There is evidence that patients with chronic pain and insomnia would benefit from sleep treatment in both disorders [14,83].

The following limitations should be considered when interpreting this data: (1) From our point of view, the major limitation of this study is its small sample size. Only five studies, with a total of 190 subjects, fulfilled the inclusion criteria and could be included in the meta-analysis. (2) Different types of sleep deprivation (total sleep deprivation, sleep restriction and partial sleep deprivation, such as slow-wave-sleep deprivation) and pain outcomes (visual analog scale, quantitative sensory testing such as heat and cold pain threshold, mechanical pain threshold, and laser-evoked pain threshold) were compared within the sample. Different sensory modalities correspond to different types of nociceptors in experimental pain threshold procedures. For example, cold pain seems to correspond with c-fibers, while heat pain corresponds with aδ-fibers [103]. Visual analog scales may be influenced by previous experiences, mood [104], and anxiety [73]. According to the Cochrane Handbook for Systematic Reviews, mixing of outcomes is applicable when calculating mean differences in meta-analysis [105]. If multiple parameters within one study sample are measured, then averaging all outcome parameters is recommended [106,107]. (3) A relevant number of studies [2,7,9,19,53–56] had to be excluded due to insufficient data. Retrieving data from figures would have been possible in some of these studies, but the selection would have been barely comprehensible. (4) The focus on RCTs (in the main analysis) and exclusion of studies that have not been published in peer-reviewed journals in the English language may affect generalizability. (5) The funnel plot for the within-group meta-analysis indicated publication bias (Fig. 3). (6) Due to the experimental design, the clinical relevance of the results is uncertain.

In summary, this meta-analysis confirms the effect of sleep deprivation on pain perception. Considering the prevalence of sleep disorders and chronic pain, future research should clarify the clinical relevance of these results. Continuing efforts in experimental and clinical research are needed to develop a translational understanding of how insomnia and pain are related.

Conflict of interest

The ICMJE Uniform Disclosure Form for Potential Conflicts of Interest associated with this article can be viewed by clicking on the following link: <http://dx.doi.org/10.1016/j.sleep.2015.07.022>.

Acknowledgment

The authors thank Friederike Hurtig (F.H.) for screening abstracts and titles during the search procedure, Anna Glechner for her support in answering methodological questions, and Robert Jank for his support revising the manuscript.

References

- [1] Nickel R, Raspe HH. [Chronic pain: epidemiology and health care utilization]. *Nervenarzt* 2001;72:897–906.
- [2] Onen SH, Alloui A, Gross A, et al. The effects of total sleep deprivation, selective sleep interruption and sleep recovery on pain tolerance thresholds in healthy subjects. *J Sleep Res* 2001;10:35–42.
- [3] Harman K, Pivik RT, D'Eon JL, et al. Sleep in depressed and nondepressed participants with chronic low back pain: electroencephalographic and behaviour findings. *Sleep* 2002;25:775–83.
- [4] Raymond I, Nielsen TA, Lavigne G, et al. Quality of sleep and its daily relationship to pain intensity in hospitalized adult burn patients. *Pain* 2001;92:381–8.
- [5] Morin CM, Gibson D, Wade J. Self-reported sleep and mood disturbance in chronic pain patients. *Clin J Pain* 1998;14:311–14.
- [6] Ohayon MM. Relationship between chronic painful physical condition and insomnia. *J Psychiatr Res* 2005;39:151–9.
- [7] Roehrs T, Hyde M, Blaisdell B, et al. Sleep loss and REM sleep loss are hyperalgesic. *Sleep* 2006;29:145–51.
- [8] Roehrs T, Roth T. Sleep and pain: interaction of two vital functions. *Semin Neurol* 2005;25:106–16.
- [9] Smith MT, Edwards RR, McCann UD, et al. The effects of sleep deprivation on pain inhibition and spontaneous pain in women. *Sleep* 2007;30:494–505.
- [10] Moffitt PF, Kalucy EC, Kalucy RS, et al. Sleep difficulties, pain and other correlates. *J Intern Med* 1991;230:245–9.
- [11] Pilowsky I, Crettenden I, Townley M. Sleep disturbance in pain clinic patients. *Pain* 1985;23:27–33.
- [12] Gislason T, Almqvist M. Somatic diseases and sleep complaints. An epidemiological study of 3,201 Swedish men. *Acta Med Scand* 1987;221:475–81.
- [13] Smith MT, Perlis ML, Smith MS, et al. Sleep quality and presleep arousal in chronic pain. *J Behav Med* 2000;23:1–13.
- [14] Roth T, Price JM, Amato DA, et al. The effect of eszopiclone in patients with insomnia and coexisting rheumatoid arthritis: a pilot study. *Prim Care Companion J Clin Psychiatry* 2009;11:292–301.
- [15] Walsh JK, Muehlbach MJ, Lauer SA, et al. Effects of triazolam on sleep, daytime sleepiness, and morning stiffness in patients with rheumatoid arthritis. *J Rheumatol* 1996;24:5–52.
- [16] Pauer L, Winkelmann A, Arsenault P, et al. An international, randomized, double-blind, placebo-controlled, phase III trial of pregabalin monotherapy in treatment of patients with fibromyalgia. *J Rheumatol* 2011;38:2643–52.
- [17] Khalid I, Roehrs TA, Hudgel DW, et al. Continuous positive airway pressure in severe obstructive sleep apnea reduces pain sensitivity. *Sleep* 2011;34:1687–91.
- [18] Stiasny-Kolster K, Pfau DB, Oertel WH, et al. Hyperalgesia and functional sensory loss in restless legs syndrome. *Pain* 2013;154:1457–63.
- [19] Moldofsky H, Scarisbrick P, England R, et al. Musculoskeletal symptoms and non-REM sleep disturbance in patients with 'fibrositis syndrome' and healthy subjects. *Psychosom Med* 1975;37:341–51.
- [20] Azevedo E, Manzano GM, Silva A, et al. The effects of total and REM sleep deprivation on laser-evoked potential threshold and pain perception. *Pain* 2011;152:2052–8.
- [21] Haack M, Lee E, Cohen DA, et al. Activation of the prostaglandin system in response to sleep loss in healthy humans: potential mediator of increased spontaneous pain. *Pain* 2009;145:136–41.
- [22] Lautenbacher S, Kundermann B, Krieg JC. Sleep deprivation and pain perception. *Sleep Med Rev* 2006;10:357–69.
- [23] Karmann AJ, Kundermann B, Lautenbacher S. [Sleep deprivation and pain: a review of the newest literature]. *Schmerz* 2014;28:141–6.
- [24] Pieh C, Popp R, Geisler P, et al. [Sleep and pain: a bi-directional relation?]. *Psychiatr Prax* 2011;38:166–70.
- [25] Volkow ND, Wang GJ, Telang F, et al. Sleep deprivation decreases binding of [¹¹C]raclopride to dopamine D₂/D₃ receptors in the human brain. *J Neurosci* 2008;28:8454–61.
- [26] Foo H, Mason P. Brainstem modulation of pain during sleep and waking. *Sleep Med Rev* 2003;7:145–54.
- [27] Przewlocka B, Mogilnicka E, Lason W, et al. Deprivation of REM sleep in the rat and the opioid peptides beta-endorphin and dynorphin. *Neurosci Lett* 1986;70:138–42.
- [28] Andersen ML, Nascimento DC, Machado RB, et al. Sleep disturbance induced by substance P in mice. *Behav Brain Res* 2006;167:212–18.
- [29] Finan PH, Goodin BR, Smith MT. The association of sleep and pain: an update and a path forward. *J Pain* 2013;14:1539–52.
- [30] Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics* 1977;33:159–74.
- [31] Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
- [32] Morris SB, DeShon RP. Combining effect size estimates in meta-analysis with repeated measures and independent-groups designs. *Psychol Methods* 2002;7:105–25.
- [33] Feingold A. Effect sizes for growth-modeling analysis for controlled clinical trials in the same metric as for classical analysis. *Psychol Methods* 2009;14:43–53.
- [34] Becker BJ. Synthesizing standardized mean-change measures. *Br J Math Stat Psychol* 1988;41:257–78.
- [35] Rosenthal R. Meta-analytic procedures for social research. SAGE Publications; 1991.
- [36] Cohen J. Statistical power analysis for the behavioral sciences. New York: 1977 [u.a.].
- [37] Field AP. Is the meta-analysis of correlation coefficients accurate when population correlations vary? *Psychol Methods* 2005;10(4):444–67.
- [38] Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–60.
- [39] Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw* 2010;36(3):1–48.
- [40] Glass J, Ablin J, Lyden A, et al. Women are more susceptible to increased symptoms of pain, fatigue, negative mood, and cognitive dysfunction after exercise and sleep restriction [conference abstract]. *Arthritis and Rheumatism [abstracts from the American College of Rheumatology/Association of Rheumatology Health Professionals Annual Scientific Meeting, ACR/ARHP 09 Atlanta, GA United States Nov 6–11 2010]* 2010.
- [41] Vollono C, Testani E, Losurdo A, et al. Migraine, arousal and sleep deprivation: comment on: "sleep quality, arousal and pain thresholds in migraineurs: a blinded controlled polysomnographic study". *J Headache Pain* 2013;14:50.
- [42] Halson SL. Sleep in elite athletes and nutritional interventions to enhance sleep. *Sports Med* 2014;44(Suppl. 1):S13–23.
- [43] Blau JN. Sleep deprivation headache. *Cephalalgia* 1990;10:157–60.
- [44] Chhangani BS, Roehrs TA, Harris EJ, et al. Pain sensitivity in sleepy pain-free normals. *Sleep* 2009;32:1011–17.
- [45] Edwards RR, Grace E, Peterson S, et al. Sleep continuity and architecture: associations with pain-inhibitory processes in patients with temporomandibular joint disorder. *Eur J Pain* 2009;13:1043–7.
- [46] Ribeiro-Dasilva MC, Goodin BR, Fillingim RB. Differences in suprathreshold heat pain responses and self-reported sleep quality between patients with temporomandibular joint disorder and healthy controls. *Eur J Pain* 2012;16:983–93.
- [47] Staud R, Smitherman ML. Peripheral and central sensitization in fibromyalgia: pathogenetic role. *Curr Pain Headache Rep* 2002;6:259–66.
- [48] Smith MT, Quartana PJ, Okonkwo RM, et al. Mechanisms by which sleep disturbance contributes to osteoarthritis pain: a conceptual model. *Curr Pain Headache Rep* 2009;13:447–54.
- [49] Steinmiller CL, Roehrs TA, Harris E, et al. Differential effect of codeine on thermal nociceptive sensitivity in sleepy versus nonsleepy healthy subjects. *Exp Clin Psychopharmacol* 2010;277–83.
- [50] Lee YB, Koh JW, Hyon JY, et al. Sleep deprivation reduces tear secretion and impairs the tear film. *Invest Ophthalmol Vis Sci* 2014;3525–31.
- [51] Sauvet F, Bourrilhon C, Besnard Y, et al. Effects of 29-h total sleep deprivation on local cold tolerance in humans. *Eur J Appl Physiol* 2012;3239–50.
- [52] Kundermann B, Hemmeter-Spernal J, Huber MT, et al. Effects of total sleep deprivation in major depression: overnight improvement of mood is accompanied by increased pain sensitivity and augmented pain complaints. *Psychosom Med* 2008;92–101.
- [53] Busch V, Haas J, Crönlein T, et al. Sleep deprivation in chronic somatoform pain-effects on mood and pain regulation. *Psychiatry Res* 2012;195:134–43.
- [54] Drewes AM, Rössel P, Arendt-Nielsen L, et al. Sleepiness does not modulate experimental joint pain in healthy volunteers. *Scand J Rheumatol* 1997;26:399–400.
- [55] Moldofsky H, Scarisbrick P. Induction of neurasthenic musculoskeletal pain syndrome by selective sleep stage deprivation. *Psychosom Med* 1976;38:35–44.
- [56] Older SA, Battafarano DF, Danning CL, et al. The effects of delta wave sleep interruption on pain thresholds and fibromyalgia-like symptoms in healthy subjects: Correlations with insulin-like growth factor I. *J Rheumatol* 1998;25:1180–6.
- [57] Haack M, Sanchez E, Mullington JM. Elevated inflammatory markers in response to prolonged sleep restriction are associated with increased pain experience in healthy volunteers. *Sleep* 2007;30:1145.
- [58] Haack M, Mullington JM. Sustained sleep restriction reduces emotional and physical well-being. *Pain* 2005;119:56–64.
- [59] Irwin MR, Olmstead R, Carrillo C, et al. Sleep loss exacerbates fatigue, depression, and pain in rheumatoid arthritis. *Sleep* 2012;35:537–43.
- [60] Schey R, Dickman R, Parthasarathy S, et al. Sleep deprivation is hyperalgesic in patients with gastroesophageal reflux disease. *Gastroenterology* 2007;1787–95.
- [61] Ablin JN, Clauw DJ, Lyden AK, et al. Effects of sleep restriction and exercise deprivation on somatic symptoms and mood in healthy adults. *Clin Exp Rheumatol* 2013;S53–9.
- [62] Kundermann B, Spernal J, Huber MT, et al. Sleep deprivation affects thermal pain thresholds but not somatosensory thresholds in healthy volunteers. *Psychosom Med* 2004;66:932–7.

- [63] Arima T, Svensson P, Rasmussen C, et al. The relationship between selective sleep deprivation, nocturnal jaw-muscle activity and pain in healthy men. *J Oral Rehabil* 2001;28:140–8.
- [64] Lentz MJ. Effects of selective slow wave sleep disruption on musculoskeletal pain and fatigue in middle aged women. *J Rheumatol* 1999;26(7):1586–92.
- [65] Schestatsky P, Dall-Agnol L, Gheller L, et al. Pain-autonomic interaction after work-induced sleep restriction. *Eur J Neurol* 2013;20:638–46.
- [66] Schuh-Hofer S, Wodarski R, Pfau DB, et al. One night of total sleep deprivation promotes a state of generalized hyperalgesia: a surrogate pain model to study the relationship of insomnia and pain. *Pain* 2013;154:1613–21.
- [67] Tiede W, Magerl W, Baumgartner U, et al. Sleep restriction attenuates amplitudes and attentional modulation of pain-related evoked potentials, but augments pain ratings in healthy volunteers. *Pain* 2010;148:36–42.
- [68] Lee JS, Jeong DU. Sleep and pain. *Sleep Med Psychophysiol* 2012;19:63–7.
- [69] Bjurström MF, Irwin MR. Polysomnographic characteristics in nonmalignant chronic pain populations: a review of controlled studies. *Sleep Med Rev* 2015;doi:10.1016/j.smrv.2015.03.004.
- [70] Nunes F, Ferreira J, Bahamondes L. Pain threshold and sleep quality in women with endometriosis. *Eur J Pain* 2015;19:15–20.
- [71] Leegaard A, Lomholt JJ, Thastum M, et al. Decreased pain threshold in juvenile idiopathic arthritis: a cross-sectional study. *J Rheumatol* 2013;40:1212–17.
- [72] Kuni B, Wang H, Rickert M, et al. Pain threshold correlates with functional scores in osteoarthritis patients. *Acta Orthop* 2015;86:215–19.
- [73] Edwards RR, Fillingim RB. Self-reported pain sensitivity: lack of correlation with pain threshold and tolerance. *Eur J Pain* 2007;11:594–8.
- [74] Edwards RR, Almeida DM, Klick B, et al. Duration of sleep contributes to next-day pain report in the general population. *Pain* 2008;137:202–7.
- [75] Lewandowski AS, Palermo TM, De la Motte S, et al. Temporal daily associations between pain and sleep in adolescents with chronic pain versus healthy adolescents. *Pain* 2010;151:220–5.
- [76] Tang NK, Goodchild CE, Sanborn AN, et al. Deciphering the temporal link between pain and sleep in a heterogeneous chronic pain patient sample: a multilevel daily process study. *Sleep* 2012;35:675–87A.
- [77] Vitiello MV, McCurry SM, Shortreed SM, et al. Short-term improvement in insomnia symptoms predicts long-term improvements in sleep, pain, and fatigue in older adults with comorbid osteoarthritis and insomnia. *Pain* 2014;155:1547–54.
- [78] Auvinen JP, Tammelin TH, Taimela SP, et al. Is insufficient quantity and quality of sleep a risk factor for neck, shoulder and low back pain? A longitudinal study among adolescents. *Eur Spine J* 2010;19:641–9.
- [79] Wright CE, Bovbjerg DH, Montgomery GH, et al. Disrupted sleep the night before breast surgery is associated with increased postoperative pain. *J Pain Symptom Manage* 2009;37:352–62.
- [80] Faraud B, Leger D, Medkour T, et al. Napping reverses increased pain sensitivity due to sleep restriction. *PLoS ONE* 2015;10:e0117425.
- [81] Roehrs TA, Harris E, Randall S, et al. Pain sensitivity and recovery from mild chronic sleep loss. *Sleep* 2012;35:1667–72.
- [82] Choy EH. The role of sleep in pain and fibromyalgia. *Nat Rev Rheumatol* 2015;11(9):513–20.
- [83] Goforth HW, Preud'homme XA, Krystal AD. A randomized, double-blind, placebo-controlled trial of eszopiclone for the treatment of insomnia in patients with chronic low back pain. *Sleep* 2014;37:1053–60.
- [84] Roth T, van Seventer R, Murphy TK. The effect of pregabalin on pain-related sleep interference in diabetic peripheral neuropathy or postherpetic neuralgia: a review of nine clinical trials. *Curr Med Res Opin* 2010;26:2411–19.
- [85] Heo M, Allison DB, Faith MS, et al. Obesity and quality of life: mediating effects of pain and comorbidities. *Obes Res* 2003;11:209–16.
- [86] Davies M, Brophy S, Williams R, et al. The prevalence, severity, and impact of painful diabetic peripheral neuropathy in type 2 diabetes. *Diabetes Care* 2006;29:1518–22.
- [87] Knutson KL, Ryden AM, Mander BA, et al. Role of sleep duration and quality in the risk and severity of type 2 diabetes mellitus. *Arch Intern Med* 2006;166:1768–74.
- [88] Wilson KG, Eriksson MY, D'Eon JL, et al. Major depression and insomnia in chronic pain. *Clin J Pain* 2002;18:77–83.
- [89] Finan PH, Smith MT. The comorbidity of insomnia, chronic pain, and depression: dopamine as a putative mechanism. *Sleep Med Rev* 2013;17:173–83.
- [90] Paiva T, Gaspar T, Matos MG. Sleep deprivation in adolescents: correlations with health complaints and health-related quality of life. *Sleep Med* 2015;16(4):521–7.
- [91] O'Brien EM, Waxenberg LB, Atchison JW, et al. Negative mood mediates the effect of poor sleep on pain among chronic pain patients. *Clin J Pain* 2010;26:310–19.
- [92] Short MA, Louca M. Sleep deprivation leads to mood deficits in healthy adolescents. *Sleep Med* 2015;16(8):987–93.
- [93] Kaida K, Niki K. Total sleep deprivation decreases flow experience and mood status. *Neuropsychiatr Dis Treat* 2014;10:19.
- [94] Tomim DH, Pontarolla FM, Bertolini JF, et al. The pronociceptive effect of paradoxical sleep deprivation in rats: evidence for a role of descending pain modulation mechanisms. *Mol Neurobiol* 2015;doi:10.1007/s12035-014-9059-0.
- [95] Block L, Lundborg C, Bjersing J, et al. Ultralow dose of naloxone as an adjuvant to intrathecal morphine infusion improves perceived quality of sleep but fails to alter persistent pain: a randomized, double-blind, controlled study. *Clin J Pain* 2015; doi: 10.1097/AJP.0000000000000200.
- [96] Irwin MR, Olmstead R, Carroll JE. Sleep disturbance, sleep duration, and inflammation: a systematic review and meta-analysis of cohort studies and experimental sleep deprivation. *Biol Psychiatry* 2015; doi:10.1016/j.biopsych.2015.05.014.
- [97] de Goeij M, van Eijk LT, Vanelderen P, et al. Systemic inflammation decreases pain threshold in humans in vivo. *PLoS ONE* 2013;8(12):e84159.
- [98] Vgontzas AN, Zoumakis E, Bixler EO, et al. Adverse effects of modest sleep restriction on sleepiness, performance, and inflammatory cytokines. *J Clin Endocrinol Metab* 2004;89:2119–26.
- [99] Floam S, Simpson N, Nemeth E, et al. Sleep characteristics as predictor variables of stress systems markers in insomnia disorder. *J Sleep Res* 2015;24(3):296–304.
- [100] Schwertner A, Conceicao Dos Santos CC, Costa GD, et al. Efficacy of melatonin in the treatment of endometriosis: a phase II, randomized, double-blind, placebo-controlled trial. *Pain* 2013;154:874–81.
- [101] Vidor LP, Torres IL, Custodio de Souza IC, et al. Analgesic and sedative effects of melatonin in temporomandibular disorders: a double-blind, randomized, parallel-group, placebo-controlled study. *J Pain Symptom Manage* 2013;46:422–32.
- [102] Reinecke H, Weber C, Lange K, et al. Analgesic efficacy of opioids in chronic pain: recent meta-analyses. *Br J Pharmacol* 2015;172:324–33.
- [103] Klinker R, Bauer C, Silbernagl S. *Lehrbuch der Physiologie*. Thieme; 2003.
- [104] Kröner-Herwig B, Lautenbacher S. Schmerzmessung und klinische Diagnostik. In: Kröner-Herwig B, Frettlöh J, Klinger R, et al., editors. *Schmerzpsychotherapie*. Berlin, Heidelberg: Springer; 2011. p. 295–318.
- [105] Higgins JPT, Green S. *Cochrane handbook for systematic reviews of interventions*. Wiley; 2011.
- [106] Cooper HM. *Synthesizing research: a guide for literature reviews*. SAGE Publications; 1998.
- [107] Scamacca N, Roberts G, Stuebing KK. Meta-analysis with complex research designs: dealing with dependence from multiple measures and multiple group comparisons. *Rev Educ Res* 2014;84:328–64.