

# A Second Attempt at MathOverflow 413935

Exact relaxations, finite-temperature barriers, and a sharper proof frontier

OpenAI Codex

AI-generated research note; directed and preserved by Rob Sneiderman

1 August 2026

**Status and nonclaim.** This note does not prove that the limit in Question 1 exists, and it does not prove nonexistence. It contains independently checked partial theorems and exact finite computations. It must not be cited or submitted as a solution to the MathOverflow question.

## Abstract

For a symmetric zero-diagonal sign matrix  $A$ , put

$$M(A) = \max_{x \in \{-1,1\}^n} \left| \frac{1}{2} x^\top A x \right|, \quad F(n) = \min_A M(A).$$

The open question is whether  $F(n)/n^{3/2}$  converges. We first audit the previous bounds  $1/\pi \leq \liminf \leq \limsup \leq 1/2$ , the augmented cut-code identity, and one-vertex monotonicity; all survive audit. We then show that the proposed cube relaxation has the unique optimizer zero at zero temperature and at every fixed unscaled positive temperature, with a leading-order zero-temperature integrality gap. In contrast, an ellipotope relaxation has the exact thermodynamic limit  $1/2$ , while the defined two-covariance, edgewise-oriented Gaussian certificate has asymptotic lower-bound constant  $1/\pi$ . We identify the augmented cut code as a binary linear code whose dual consists of even-cardinality Eulerian subgraphs, and derive an exact signed-Eulerian MacWilliams formula. Ordinary graphons and limiting empirical spectra are proved to erase the  $n^{3/2}$  functional. Exhaustive computation gives  $F(2), \dots, F(10) = 1, 3, 4, 4, 5, 9, 10, 12, 13$ . The sharp remaining target is a quenched minimax free-energy limit that absorbs, rather than separately bounds, the cross-block contribution.

## 1 Problem, audited baseline, and scope

The question, asked by Paata Ivanisvili on MathOverflow in 2022 [1], is whether

$$\lim_{n \rightarrow \infty} \frac{F(n)}{n^{3/2}}$$

exists. Here  $A_{ii} = 0$ ,  $A_{ij} = A_{ji} = a_{ij} \in \{-1, 1\}$ , and

$$Q_A(x) = \sum_{i < j} a_{ij} x_i x_j = \frac{1}{2} x^\top A x.$$

The earlier attempt was rechecked from source rather than assumed. Its central conclusions are correct:

$$\frac{1}{\pi} \leq \liminf_{n \rightarrow \infty} \frac{F(n)}{n^{3/2}} \leq \limsup_{n \rightarrow \infty} \frac{F(n)}{n^{3/2}} \leq \frac{1}{2}, \quad (1)$$

$$F(n) = \binom{n}{2} - 2\rho(D_n), \quad F(n) \leq F(n+1) \leq F(n) + n. \quad (2)$$

The lower bound uses two Gaussian sign roundings and the arcsine law. The upper bound uses symmetric Paley conference matrices and principal submatrices. The code  $D_n$  is the cut-character set augmented by its negative. A claim-by-claim audit, including two attribution errors and a factor-of-two prose correction, is in `AUDIT.md`. Nothing in (1) or (2) implies convergence.

All new statements below distinguish a proved identity from a proposed limit theorem. Exact computations guide the search but are not used as asymptotic proofs.

## 2 The relaxed-coupling program collapses

Let  $m = \binom{n}{2}$ , let  $c_x = (x_i x_j)_{i < j}$ , and for arbitrary real couplings define

$$G_n(a) = \max_{x \in \{-1, 1\}^n} |\langle a, c_x \rangle|.$$

**Theorem 1** (Cube relaxation). *The function  $G_n$  is a norm on  $\mathbb{R}^m$ . Hence*

$$\min_{a \in [-1, 1]^m} G_n(a) = 0,$$

*and the unique minimizer is  $a = 0$ .*

*Proof.* Homogeneity and the triangle inequality follow from the maximum of absolute linear forms. If  $X$  is uniform on  $\{-1, 1\}^n$ , orthogonality of distinct degree-two Walsh characters gives

$$\mathbb{E} \langle a, c_X \rangle^2 = \sum_{i < j} a_{ij}^2.$$

Thus  $G_n(a) = 0$  forces  $a = 0$ . Since the origin belongs to the cube, the last assertion follows.  $\square$

In particular, any convex feasible set in the original coupling coordinates that contains every sign coupling contains their convex hull, the whole cube, and therefore contains this zero optimizer when the objective remains  $G_n$ . By the audited lower bound, its additive integrality gap is at least  $n\sqrt{n-1}/\pi$ , of the full target order.

The same collapse persists at positive temperature. Retain duplicated states  $(x, s) \in \{-1, 1\}^n \times \{-1, 1\}$  and put

$$L_{n,t}(a) = \frac{1}{t} \log \sum_{x,s} \exp(t \langle a, sc_x \rangle), \quad t > 0.$$

**Theorem 2** (Finite-temperature cube dual). *The function  $L_{n,t}$  is strictly convex and has the unique cube minimizer zero, with value*

$$\min_{a \in [-1, 1]^m} L_{n,t}(a) = \frac{(n+1) \log 2}{t}.$$

*It has the exact dual representation*

$$\max_{p \in \Delta(\{-1, 1\}^n \times \{-1, 1\})} \left\{ \frac{H(p)}{t} - \|\mathbb{E}_p[sc_x]\|_1 \right\}.$$

*Proof.* The Hessian of  $L_{n,t}$  is  $t$  times the covariance matrix of  $sc_x$  under a full-support Gibbs distribution. If a linear functional of these vectors were constant, replacing  $s$  by  $-s$  would make the constant zero; Walsh orthogonality would then make the functional zero. The Hessian is therefore positive definite. Symmetry makes the gradient vanish at zero, where the value is  $t^{-1} \log 2^{n+1}$ .

The entropy variational formula for log-sum-exp gives

$$L_{n,t}(a) = \max_p \left\{ \langle a, \mathbb{E}_p[sc_x] \rangle + \frac{H(p)}{t} \right\}.$$

Both domains are compact and convex and the displayed function is convex in  $a$  and concave in  $p$ , so Sion's theorem applies. Minimization of a linear form over the cube is minus the  $\ell^1$ -norm of its coefficient.  $\square$

Thus the simplest probability-measure dual is exact but vacuous for the discrete problem: the uniform state measure cancels every first moment.

There is nevertheless a useful conditional rounding statement.

**Proposition 3** (Variance-sensitive sign rounding). *For  $a \in [-1, 1]^m$ , set  $V(a) = \sum_e (1 - a_e^2)$ . There is a sign vector  $\sigma \in \{-1, 1\}^m$  for which*

$$G_n(\sigma) \leq G_n(a) + \sqrt{2V(a)(n+1) \log 2} + \frac{4}{3}(n+1) \log 2.$$

*Proof.* Round the coordinates independently with  $\mathbb{E}\sigma_e = a_e$ . For fixed  $x$ , the centered error  $\sum_e (\sigma_e - a_e)c_x(e)$  has variance  $V(a)$  and summands bounded by two. The two-sided Bernstein bound with parameter  $u$  is at most  $2e^{-u}$  beyond  $\sqrt{2V(a)u} + 4u/3$ . There are  $2^{n-1}$  distinct cut characters. Taking  $u = (n+1) \log 2$  makes the union failure probability at most  $1/2$ . For a successful outcome, add this uniform error to  $G_n(a)$ .  $\square$

This gives  $o(n^{3/2})$  loss if  $V(a) = o(n^2)$ , but the cube optimizer has  $V(0) = m$ . The obvious spherical repairs also fail. On  $\|a\|_2 = \sqrt{m}$ , Parseval gives  $G_n(a) \geq \sqrt{m}$ , attained by a one-coordinate vector. Adding  $|a_e| \leq 1$  to that sphere forces every  $|a_e| = 1$ , recovering the original discrete problem exactly.

### 3 Temperature scaling and the cross-block wall

The functional proposed in the question is

$$\Phi_{n,\beta} = \min_A \frac{1}{\beta n^{3/2}} \log \sum_x e^{\beta |Q_A(x)|}.$$

For every fixed  $\beta > 0$ , the maximum-versus-log-sum-exp inequality gives

$$\frac{F(n)}{n^{3/2}} \leq \Phi_{n,\beta} \leq \frac{F(n)}{n^{3/2}} + \frac{\log 2}{\beta \sqrt{n}}. \quad (3)$$

Thus existence of  $\lim_n \Phi_{n,\beta}$  for even one fixed positive  $\beta$  is equivalent to the original limit. At this normalization, fixed temperature is already asymptotically zero temperature.

A genuinely extensive mean-field functional is

$$\Psi_{n,\beta} = \min_A \frac{1}{\beta n} \log \sum_{x,s} \exp \left( \frac{\beta s Q_A(x)}{\sqrt{n}} \right).$$

It satisfies

$$\frac{F(n)}{n^{3/2}} \leq \Psi_{n,\beta} \leq \frac{F(n)}{n^{3/2}} + \frac{(n+1)\log 2}{\beta n}. \quad (4)$$

Consequently, existence of  $\lim_n \Psi_{n,\beta}$  for arbitrarily large fixed  $\beta$  would settle the problem: the limsup-minus-liminf gap would be at most  $\log 2/\beta$  for every such  $\beta$ .

The direct interpolation does not have the needed accuracy. Define

$$h_n(t) = \min_A \log \sum_{x,s} e^{tsQ_A(x)}.$$

**Proposition 4** (Exact annealed block estimate). *For all  $n, k$  and  $t > 0$ ,*

$$h_{n+k}(t) \leq h_n(t) + h_k(t) - \log 2 + nk \log \cosh t.$$

*Proof.* For a within-block matrix  $A$ , write  $Z_A^s = \sum_x e^{tsQ_A(x)}$ . Replacing all coefficients of the second block by their negatives swaps its two one-sided partition sums. Of the two possible alignments, one satisfies

$$Z_A^+ Z_B^+ + Z_A^- Z_B^- \leq \frac{1}{2}(Z_A^+ + Z_A^-)(Z_B^+ + Z_B^-).$$

Choose every cross coefficient independently and uniformly. For fixed  $x, y, s$ , averaging the cross exponential gives  $(\cosh t)^{nk}$ . Therefore the expected augmented partition sum is at most the right side above times this factor, and some deterministic cross signing is no larger. Take logarithms and infima.  $\square$

At  $t = \beta/\sqrt{n+k}$  and  $n/(n+k) \rightarrow \alpha$ , the normalized cross cost tends to

$$\frac{\beta}{2}\alpha(1-\alpha),$$

which is nonzero for fixed  $\beta$  and grows in the zero-temperature limit. The direct annealed cross block is therefore a leading term, not an error.

Finally, the absolute value cannot be silently removed. If every edge from vertex 1 is positive and every other edge negative, fix  $x_1 = 1$  and put  $T = \sum_{i>1} x_i$ . Then

$$Q_A(x) = \frac{n}{2} - \frac{(T-1)^2}{2}.$$

Hence  $\max_x Q_A(x) = \lfloor n/2 \rfloor$ , whereas  $\max_x |Q_A(x)| = \binom{n}{2}$ .

## 4 Covariance relaxations: two exact asymptotic limits

Let

$$\mathcal{E}_n = \{R \succeq 0 : \text{diag } R = \mathbf{1}\}$$

be the elliptope. The full arcsine expression is tautologically exact:

$$M(A) = \max_{R \in \mathcal{E}_n} \left| \frac{2}{\pi} \sum_{i < j} a_{ij} \arcsin R_{ij} \right|. \quad (5)$$

Indeed, Gaussian sign rounding makes every value on the right at most  $M(A)$ , while  $R = xx^\top$  attains  $Q_A(x)$ . The nonlinearity therefore preserves the original problem rather than relaxing it.

Linearizing gives a model with a complete thermodynamic limit. Put

$$S(A) = \max_{R \in \mathcal{E}_n} \left| \frac{1}{2} \text{tr}(AR) \right|, \quad S_n = \min_A S(A).$$

**Theorem 5** (Adversarial elliptope limit). *For every  $n \geq 2$ ,*

$$S_n \geq \frac{1}{2}n\sqrt{n-1},$$

and

$$\boxed{\lim_{n \rightarrow \infty} \frac{S_n}{n^{3/2}} = \frac{1}{2}.}$$

*At each symmetric conference order, equality holds in the finite bound.*

*Proof.* Fix any sign matrix  $A$ , set  $q = n - 1$ , and define

$$R^\pm = \frac{1}{2} \left( I \pm \frac{A}{\sqrt{q}} \right)^2.$$

They are positive semidefinite. Since every row of  $A$  has squared norm  $q$ , their diagonals equal one. Moreover

$$R^+ - R^- = \frac{2A}{\sqrt{q}}.$$

The corresponding two linear SDP values differ by

$$\frac{1}{2} \operatorname{tr}(A(R^+ - R^-)) = n\sqrt{q}.$$

At least one has absolute value at least half this amount.

For the reverse asymptotic inequality, if  $R \in \mathcal{E}_n$ , then

$$|\operatorname{tr}(AR)| \leq \|A\|_{\text{op}} \operatorname{tr} R = n\|A\|_{\text{op}}.$$

A symmetric conference matrix has operator norm  $\sqrt{n-1}$ , so it attains equality. For general  $n$ , the prime number theorem in arithmetic progressions supplies a prime  $q \equiv 1 \pmod{4}$  with  $q+1 \geq n$  and  $q+1 = n(1+o(1))$  [9]. Take a principal  $n$ -by- $n$  submatrix of the symmetric Paley conference matrix of order  $N = q+1$ . Compression does not increase its operator norm  $\sqrt{N-1}$ , giving  $S_n \leq n\sqrt{N-1}/2 = (1/2 + o(1))n^{3/2}$ .  $\square$

Rank-one  $R = xx^\top$  show  $S(A) \geq M(A)$ . The missing transfer is an asymptotically vanishing integrality gap, which is not supplied by standard Grothendieck-type constant-factor rounding [5, 6]. The gap is already strict at order six: the exact computation below gives  $F(6) = 5$ , whereas Theorem 5 and a conference matrix give  $S_6 = 3\sqrt{5}$ .

There is also a sharp barrier for the linear Gaussian-sign method. Let  $\Lambda(A)$  be the maximum of

$$\sum_{i < j} a_{ij}(R_{ij}^+ - R_{ij}^-)$$

over  $R^\pm \in \mathcal{E}_n$ , subject to the edgewise orientations  $a_{ij}(R_{ij}^+ - R_{ij}^-) \geq 0$ .

**Theorem 6** (Oriented covariance barrier). *For every sign matrix  $A$ ,*

$$M(A) \geq \frac{\Lambda(A)}{\pi}, \quad \Lambda(A) \geq n\sqrt{n-1}.$$

Moreover

$$\boxed{\lim_{n \rightarrow \infty} \frac{\min_A \Lambda(A)}{n^{3/2}} = 1.}$$

*Proof.* Gaussian sign rounding of  $R^+$  and  $R^-$ , followed by the arcsine law, gives two expected energies. Since  $\arcsin'(r) \geq 1$  and each edge difference has the prescribed orientation, their difference is at least  $2\Lambda(A)/\pi$ . Both expectations lie in  $[-M(A), M(A)]$ , proving the first inequality.

The square covariances used in Theorem 5 satisfy the orientation constraints and have objective  $n\sqrt{n-1}$ . Conversely,

$$\Lambda(A) = \frac{1}{2} \operatorname{tr}(A(R^+ - R^-)) \leq n\|A\|_{\text{op}}.$$

Conference matrices attain equality. The same prime-number-theorem and Paley principal-submatrix construction used in Theorem 5 gives the matching asymptotic upper bound at every order.  $\square$

Thus the optimized two-covariance, edgewise-oriented linear certificate  $\Lambda$  defined above has adversarial asymptotic lower-bound constant exactly  $1/\pi$ . Improving it requires information beyond this specific linearization; the theorem does not exclude more general covariance certificates.

## 5 The augmented cut code and its exact dual

Encode signs as bits via  $(-1)^b$ . Let  $j \in \mathbb{F}_2^m$  be the all-one edge word and

$$(\delta u)_{ij} = u_i + u_j.$$

Then the augmented code in (2) is

$$D_n = \{\delta u + sj : u \in \mathbb{F}_2^n, s \in \mathbb{F}_2\}.$$

**Theorem 7** (Linear code and dual). *For  $n \geq 3$ ,  $D_n$  is a binary  $[m, n]$  linear code and*

$$D_n^\perp = \{H \subseteq E(K_n) : \deg_H(v) \equiv 0 \pmod{2} \text{ for all } v, |H| \equiv 0 \pmod{2}\}.$$

*For  $n \geq 4$ , the dual is spanned by 4-cycles, has dimension  $m - n$ , and minimum distance four.*

*Proof.* The cut space has dimension  $n - 1$ . Its words have even parity on every triangle, whereas  $j$  has odd triangle parity, so adjoining  $j$  raises the dimension by one. A graph is orthogonal to every cut exactly when every vertex has even degree, and it is orthogonal to  $j$  exactly when its number of edges is even.

The cycle space is generated by triangles through one fixed vertex. An even-weight cycle-space word is a sum of an even number of these basis triangles. The sum of two sharing a nonfixed vertex is a 4-cycle. Two that share only the fixed vertex can be paired through an intermediate basis triangle and expressed as two 4-cycles. This proves generation. A nonzero even-cardinality Eulerian graph has at least four edges, and a 4-cycle exists.  $\square$

Counting cuts by the size of one shore gives

$$W_{D_n}(z) = \frac{1}{2} \sum_{k=0}^n \binom{n}{k} \left( z^{k(n-k)} + z^{m-k(n-k)} \right). \quad (6)$$

Let  $a_e = (-1)^{y_e}$  and define the exact augmented partition function

$$Z_y(\beta) = \sum_{d \in D_n} \exp(\beta[m - 2 \operatorname{wt}(y + d)]).$$

Its ground state is  $M(A)$ .

**Theorem 8** (Signed-Eulerian MacWilliams formula). *For  $n \geq 3$ ,*

$$Z_y(\beta) = 2^n (\cosh \beta)^m P_y(\tanh \beta),$$

where

$$P_y(t) = \sum_{H \in D_n^\perp} (-1)^{|H \cap y|_t|H|}.$$

For  $0 \leq t < 1$ ,  $P_y(t) > 0$ , and uniformly in  $y$ ,

$$M(A) \leq \frac{1}{\beta} \log Z_y(\beta) \leq M(A) + \frac{n \log 2}{\beta}.$$

*Proof.* The coset MacWilliams identity is

$$W_{y+D_n}(z) = 2^{n-m} (1+z)^m \sum_{H \in D_n^\perp} (-1)^{|H \cap y|} \left( \frac{1-z}{1+z} \right)^{|H|}.$$

Multiply by  $e^{\beta m}$  and substitute  $z = e^{-2\beta}$ . This gives the formula because  $e^\beta(1 + e^{-2\beta}) = 2 \cosh \beta$  and the final ratio is  $\tanh \beta$ . Positivity follows because the left side is a positive partition sum. The last display compares its largest summand with a sum of  $|D_n| = 2^n$  summands.  $\square$

For fixed  $\beta > 0$ , the minimized free energy in Theorem 8, divided by  $n^{3/2}$ , differs from  $F(n)/n^{3/2}$  by  $O(n^{-1/2})$ . Thus the exact unresolved term is

$$\min_y \log P_y(\tanh \beta).$$

It must cancel the explicit  $m \log \cosh \beta$  term to  $n^{3/2}$  accuracy. Averaging  $P_y(t)$  over all  $y$  gives one and yields only

$$F(n) \leq n \sqrt{(n-1) \log 2},$$

the usual sphere/union-bound constant, weaker than  $1/2$ .

Puncturing also stops at the old scale. For  $n \geq 4$ , deleting the final star maps  $D_n$  onto  $D_{n-1}$ , and the two lifts of each word have complementary star coordinates. Therefore

$$\rho(D_{n-1}) \leq \rho(D_n) \leq \rho(D_{n-1}) + \left\lfloor \frac{n-1}{2} \right\rfloor,$$

equivalently

$$F(n-1) + ((n-1) \bmod 2) \leq F(n) \leq F(n-1) + n-1.$$

The  $n = 3$  instance of these inequalities is checked directly; the fiber there has four rather than two words. Across a two-block partition, the cross lifts are complementary rank-one sign matrices. On the dual side, the  $\binom{n}{2} \binom{k}{2}$  alternating cross-block 4-cycles already contribute macroscopically at mean-field scaling. Product bounds that take absolute values discard the cancellation needed here.

## 6 High-temperature data do not truncate uniformly

Let  $X$  be uniform on  $\{-1, 1\}^n$ , let  $Y = Q_A(X)$ , and use the symmetrized partition function  $\mathbb{E} \cosh(tY)$ , equivalently the augmented state  $sY$ . Direct Walsh counting gives

$$\mathbb{E}Y^2 = m, \quad \mathbb{E}Y^4 = 3m^2 - 2m + 24 \sum_{C_4} \prod_{e \in C_4} a_e, \quad (7)$$

where the sum is over undirected 4-cycles. Indeed, a monomial in the fourth power survives exactly when every vertex has even incidence. Besides one edge four times and two edges twice, the only case is a simple 4-cycle, with multinomial factor 24.

Equivalently,

$$\kappa_4(Y) = 3 \operatorname{tr}(A^4) - 2n(n-1)(3n-4), \quad \operatorname{tr}(A^4) = n(n-1)^2 + \sum_{i \neq j} (A_{ij}^2)^2. \quad (8)$$

The second identity is  $\operatorname{tr}(A^4) = \sum_{ij} (A_{ij}^2)^2$ ; the first follows by separating the closed walks with four distinct vertices in (7).

The expansion

$$\log \mathbb{E} \cosh(tY) = \frac{mt^2}{2} + \frac{\kappa_4(Y)t^4}{24} + O(t^6)$$

shows that, for each fixed  $n$ , all sufficiently high-temperature minimizers of this symmetrized partition function minimize  $\operatorname{tr}(A^4)$ . The symmetrization matters: the one-sided partition can have a signing-dependent cubic triangle term. Spectral Cauchy–Schwarz gives

$$\operatorname{tr}(A^4) \geq \frac{\operatorname{tr}(A^2)^2}{n} = n(n-1)^2.$$

For a Paley principal submatrix of order  $n$  inside a conference matrix of order  $N = n(1 + o(1))$ ,

$$\operatorname{tr}(A^4) \leq \|A\|_{\text{op}}^2 \operatorname{tr}(A^2) \leq (N-1)n(n-1).$$

This gives the reverse asymptotic inequality, so

$$\lim_n \frac{\min_A \operatorname{tr}(A^4)}{n^3} = 1, \quad \lim_n \frac{\min_A \kappa_4(Y)}{n^3} = -3.$$

This fourth-order limit does not determine the next term. Exhausting all root-normalized order-seven signings gives

$$\min_A \operatorname{tr}(A^4) = 342,$$

while the trace-minimizing signings have three distinct exact sixth moments,

$$50,781, \quad 53,661, \quad 59,421.$$

Thus fourth-order data alone are insufficient. No uniform remainder theorem connecting this high-temperature expansion to the required ground-state regime was obtained.

## 7 Ordinary graphons and empirical spectra lose the functional

For a signed matrix  $A$ , let  $W_A$  be its step graphon and normalize

$$\|W_A\|_{\square} = \frac{1}{n^2} \max_{S, T \subseteq [n]} \left| \sum_{i \in S, j \in T} a_{ij} \right|.$$

**Theorem 9** (Cut-limit obstruction). *If  $M(A_n) = O(n^{3/2})$ , then  $W_{A_n} \rightarrow 0$  in cut norm. There are two deterministic sequences satisfying this hypothesis and having the same zero graphon limit, while their normalized Boolean maxima are separated by at least  $3/2$ .*

*Proof.* For  $x, y \in \{-1, 1\}^n$ , split the coordinates according to whether  $x_i = y_i$ . Writing  $z^+, z^- \in \{-1, 0, 1\}^n$  for the corresponding restrictions of  $y$ , symmetry of  $A$  cancels cross terms and gives

$$x^T A y = 2Q_A(z^+) - 2Q_A(z^-).$$

A multilinear quadratic form attains its maximum absolute value on  $[-1, 1]^n$  at a Boolean vertex, so every sign bilinear form is bounded by  $4M(A)$ . Expressing two indicator vectors as affine combinations of sign vectors then gives

$$\|W_A\|_{\square} \leq \frac{4M(A)}{n^2}.$$

Take an infinite sequence of Paley conference matrices  $C_N$ . Their normalized Boolean maxima have limsup at most  $1/2$ , and their graphons tend to zero. Choose  $k = \lfloor 2N^{3/4} \rfloor$  vertices and change every coefficient inside that set to  $+1$ , obtaining  $B_N$ . At most  $\binom{k}{2} = O(N^{3/2})$  edges change, so  $M(B_N) = O(N^{3/2})$  and the graphon perturbation tends to zero in  $L^1$ . Fix the planted spins to  $+1$  and average all outside spins independently. Every cross and outside term has mean zero, while the planted block contributes  $\binom{k}{2}$ . Some outside assignment therefore has at least that energy, and

$$\liminf_N \frac{M(B_N)}{N^{3/2}} \geq 2.$$

The separation from the conference limsup is at least  $3/2$ . □

The usual dense graph topology therefore collapses every target-scale near-optimizer to the same point. This is not repaired by keeping only a limiting empirical spectrum. At dense scaling, every empirical law of  $A/n$  tends to  $\delta_0$ , since its average squared eigenvalue is  $(n-1)/n^2$ . At fluctuation scaling there is a stronger obstruction.

**Proposition 10** (Empirical-spectral obstruction). *For every fixed  $d > 1/2$ , there are conference matrices  $C_N$  and sign matrices  $B_N$  such that*

$$W_2(\mu_{B_N/\sqrt{N}}, \mu_{C_N/\sqrt{N}}) \rightarrow 0,$$

but

$$\liminf_N \frac{M(B_N)}{N^{3/2}} \geq d, \quad \limsup_N \frac{M(C_N)}{N^{3/2}} \leq \frac{1}{2}.$$

*Proof.* Let  $q_N = Q_{C_N}(\mathbf{1})$ . The spectral bound gives  $|q_N| \leq N\sqrt{N-1}/2$ . Flip negative coefficients to positive until the all-one energy is at least  $dN^{3/2}$ . This needs  $O(N^{3/2})$  edges, and there are  $\Theta(N^2)$  available. Therefore  $\|B_N - C_N\|_F^2 = O(N^{3/2})$ . Hoffman–Wielandt implies

$$W_2^2(\mu_{B_N/\sqrt{N}}, \mu_{C_N/\sqrt{N}}) \leq \frac{\|B_N - C_N\|_F^2}{N^2} = O(N^{-1/2}).$$

The energy inequalities follow from the construction and the conference spectral bound. □

This excludes dense graphons and limiting empirical spectral distributions, not spectral edges, eigenvectors, or a richer Gaussian-process limit.

For comparison, iid signings are rigorously worse than the conference upper constant.

**Proposition 11** (Greedy energy of an iid signing). *On one infinite iid Rademacher edge array  $A_n$ , almost surely*

$$\liminf_n \frac{M(A_n)}{n^{3/2}} \geq \frac{2}{3} \sqrt{\frac{2}{\pi}} = 0.531923 \dots > \frac{1}{2}.$$

*Proof.* Choose spins sequentially by

$$x_1 = 1, \quad x_j = \operatorname{sgn} \sum_{i < j} a_{ij} x_i.$$

The resulting energy is

$$Q_{A_n}(x) = \sum_{j=2}^n \left| \sum_{i < j} a_{ij} x_i \right| = \sum_{k=1}^{n-1} Y_k.$$

Before column  $j$  is exposed, the earlier spins are measurable with respect to earlier columns. Multiplying the fresh iid column signs by those predictable spins preserves their iid Rademacher law. Hence the  $Y_k$  are independent and  $Y_k$  has the law of  $|\varepsilon_1 + \dots + \varepsilon_k|$ .

The central limit theorem and uniform integrability give  $\mathbb{E}Y_k/\sqrt{k} \rightarrow \sqrt{2/\pi}$ . A Riemann-sum argument therefore yields

$$\frac{1}{n^{3/2}} \sum_{k < n} \mathbb{E}Y_k \longrightarrow \sqrt{\frac{2}{\pi}} \int_0^1 \sqrt{t} dt = \frac{2}{3} \sqrt{\frac{2}{\pi}}.$$

Changing one input sign changes  $Y_k$  by at most two, so its centered moment generating function is bounded by  $\exp(\lambda^2 k/2)$ . Independence gives an exponential tail  $2e^{-c_\varepsilon n}$  for deviations of  $\sum_{k < n} Y_k$  by  $\varepsilon n^{3/2}$ . This is summable; Borel–Cantelli completes the proof.  $\square$

This is not a lower bound on  $F(n)$ ; it shows that typical iid signings are not asymptotically optimal.

## 8 Exact computation through order ten

Switching makes every edge incident with vertex zero positive. The remaining negative coefficients form a graph on  $n - 1$  vertices, and vertex permutations reduce the search to one unlabeled graph in each isomorphism class. For a residual negative-edge graph  $G$  and a negative-spin set  $S$  not containing the root, the exact energy is

$$\binom{n}{2} - 2|E(G)| - 2|S|(n - |S|) + 4|\delta_G(S)|.$$

Exhausting nauty’s `geng` streams gives Table 1.

Completeness is checked against the unlabeled-graph counts and hashes of the full graph streams. The reduced formula is checked by direct evaluation; all labeled signings are independently enumerated through  $n = 6$ ; optional NetworkX and Z3 routes use different decoders and optimization mechanisms. Z3 independently gives UNSAT/SAT pairs certifying the displayed values for  $n = 7, 8, 9$ .

Table 1: Exact computational values.

| $n$            | 2     | 3     | 4     | 5     | 6     | 7     | 8     | 9     | 10    |
|----------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| $F(n)$         | 1     | 3     | 4     | 4     | 5     | 9     | 10    | 12    | 13    |
| $F(n)/n^{3/2}$ | .3536 | .5774 | .5000 | .3578 | .3402 | .4860 | .4419 | .4444 | .4111 |

At  $n = 10$ , the lower certificate is the exhaustive nauty search; the attempted Z3 lower query timed out.

The Paley conference matrix of order ten over  $\mathbb{F}_9$  has exact Boolean maximum 15, strictly worse than  $F(10) = 13$ . The two optimal switching classes at order ten have different characteristic polynomials. Thus neither conference structure nor a unique spectrum describes all finite optimizers. These finite facts do not imply a limiting constant.

## 9 Five proof programs and the sharp remaining lemma

The detailed derivations and failed variants are preserved in `RESEARCH_LEDGER.md`. The current programs are logically distinct.

**A. Quenched extensive free energy.** Prove  $\lim_n \Psi_{n,\beta}$  for every fixed  $\beta > 0$ , or at least for an unbounded set of  $\beta$ 's. Equation (4) then forces convergence. Proposition 4 shows exactly why independent annealed cross blocks fail. A repair must absorb the cross term in a joint minimax order parameter, as random-disorder spin-glass interpolation absorbs it only after disorder expectation [3, 4].

**B. Near-saturated relaxation.** Find a nonconvex compact relaxation whose optimizers have  $\sum_e (1 - a_e^2) = o(n^2)$  and an asymptotic composition law. Then Proposition 3 transfers its limit to signs. The cube and sphere calculations prove that the two obvious feasible sets are structurally too loose or exactly discrete.

**C. Signed Eulerian polynomial.** Prove a second-order limit for  $\min_y \log P_y(t)$  after the explicit MacWilliams terms are included. Puncturing and ordinary products fail because of the common parity phase, rank-one cross lift, and macroscopic family of mixed 4-cycles. A quenched coset-weight theorem, rather than an average weight enumerator, is required.

**D. Second-order limit object.** Construct a rate-sensitive topology for  $A/\sqrt{n}$  in which the Boolean maximum is continuous and sign arrays have an asymptotically lossless composition. Theorem 9 and Proposition 10 rule out ordinary graphons and empirical spectra. Spectral edges, eigenvectors, or the whole induced Gaussian process would have to remain visible.

**E. Nonexistence.** To separate subsequences one needs order-specific lower bounds on  $F(n)$ , not merely constructions with different energies. Conference, random, and planted matrices supply upper bounds or deliberately inferior examples. The small values fluctuate but do not certify a subsequence mechanism.

No route presently closes. The single best next lemma is the following.

**Quenched minimax free-energy lemma.** For every fixed  $\beta > 0$ , the limit

$$\lim_{n \rightarrow \infty} \min_A \frac{1}{\beta n} \log \sum_{x,s} \exp \left( \frac{\beta s Q_A(x)}{\sqrt{n}} \right)$$

exists.

Proving this lemma for an unbounded collection of inverse temperatures settles the MathOverflow question by (4). In coding coordinates it asks for cancellation in  $P_y(\tanh(\beta/\sqrt{n}))$ . Any proof that bounds the mixed cross-block contribution independently incurs a leading error; a successful proof must make that contribution part of the optimized variational structure.

## Reproducibility and final status

The exact identity checker, exhaustive small-order search, optional Z3 certificate, expected outputs, deterministic seeds, and certificate boundaries are documented in `verification/README.md`. The literature map is in `LITERATURE.md`; the original and second proof-search ledgers are both preserved.

The adversarial ellipsope limit, the code dual and MacWilliams formula, the oriented covariance barrier, and the graph-limit obstructions materially narrow the search. They do not prove existence or nonexistence of the original limit.

## References

- [1] P. Ivanišvili, *Min-max of a quadratic form of plus-minus ones*, MathOverflow Question 413935, 2022. <https://mathoverflow.net/questions/413935/>
- [2] P. Solé and T. Zaslavsky, *A coding approach to signed graphs*, SIAM J. Discrete Math. **7** (1994), 544–553. <https://doi.org/10.1137/S0895480189174374>
- [3] F. Guerra and F. L. Toninelli, *The thermodynamic limit in mean field spin glass models*, Comm. Math. Phys. **230** (2002), 71–79. <https://doi.org/10.1007/s00220-002-0699-y>
- [4] A. Auffinger and W.-K. Chen, *Parisi formula for the ground state energy in the mixed p-spin model*, Ann. Probab. **45** (2017), 4617–4631. <https://doi.org/10.1214/16-AOP1173>
- [5] N. Alon and A. Naor, *Approximating the cut-norm via Grothendieck’s inequality*, SIAM J. Comput. **35** (2006), 787–803. <https://doi.org/10.1137/S0097539704441629>
- [6] Y. Nesterov, *Semidefinite relaxation and nonconvex quadratic optimization*, Optim. Methods Softw. **9** (1998), 141–160. <https://doi.org/10.1080/10556789808805690>
- [7] L. Lovász and B. Szegedy, *Limits of dense graph sequences*, J. Combin. Theory Ser. B **96** (2006), 933–957. <https://doi.org/10.1016/j.jctb.2006.05.002>
- [8] R. E. A. C. Paley, *On orthogonal matrices*, J. Math. Phys. **12** (1933), 311–320. <https://doi.org/10.1002/sapm1933121311>
- [9] H. Davenport, *Multiplicative number theory*, third ed., Graduate Texts in Mathematics 74, Springer, 2000. <https://doi.org/10.1007/978-1-4757-5927-3>